Structural basis of EHEP-mediated offense against phlorotannin-induced

defense from brown algae to protect *aku*BGL activity

Xiaomei Sun[1,¶], Yuxin Ye[1,#a,¶], Naofumi Sakurai[1], Hang Wang[1], Koji Kato[1,#b],

Jian Yu[1], Keizo Yuasa[2,#c], Akihiko Tsuji[2], Min Yao[1,*]

[1] Faculty of Advanced Life Science, Hokkaido University, Sapporo, Japan

[2] Graduate School of Bioscience and Bioindustry, Tokushima University, Tokushima,

Japan

[#a] Current address: Pingshan translational medicine center, Shenzhen Bay Laboratory,

Shenzhen, China

[#b] Current address: Structural Biology Division, Japan Synchrotron Radiation Research

Institute (JASRI), Hyogo, Japan

[#c] Current address: Faculty of Science and Engineering, Setsunan University, Neyagawa,

Japan

*Corresponding author

E-mail: yao@castor.sci.hokudai.ac.jp (MY)

¶ These authors contributed equally to this work.

Abstract

The defensive-offensive associations between algae and herbivores determine marine ecology. Brown algae utilize phlorotannin as their chemical defense against the predator *Aplysia kurodai*, which uses β-glucosidase (*aku*BGL) to digest the laminarin in algae to glucose. Moreover, *A. kurodai* employs *Eisenia* hydrolysis-enhancing protein (EHEP) as an offense to protect *aku*BGL activity from phlorotannin inhibition by precipitating phlorotannin. To underpin the molecular mechanism of this digestive-defensive-offensive system, we determined the structures of apo and tannic-acid (TNA, a phlorotannin-analog) bound form of EHEP, as well as *aku*BGL. EHEP consisted of three peritrophin-A domains formed in a triangle and bound TNA in the center without significant conformational changes. Structural comparison between EHEP and EHEP–TNA led us to find that EHEP can be resolubilized from phlorotannin-precipitation at an alkaline pH, which reflects a requirement in the digestive tract. *aku*BGL contained two GH1 domains, only one of which conserved the active site. Combining docking analysis, we propose the mechanisms by which phlorotannin inhibits *aku*BGL by occupying the substrate-binding pocket, and EHEP protects *aku*BGL against the inhibition by binding with phlorotannin to free the *aku*BGL pocket.

Introduction

Over millions of years of evolution, predators have successfully coevolved with their prey to maintain an ecological balance [1]. In marine habitats, interactions between algae and marine herbivores dominate marine ecosystems. Most algae are consumed by marine herbivores [2]. They produce secondary metabolites as a chemical defense to protect themselves against predators. The sea hare *Aplysia kurodai*, a marine gastropod, preferentially feeds on the laminarin-abundant brown algae *Eisenia bicyclis* (laminarin constitutes 20%–30% of the dry weight of *E. bicyclis* and acts as a major storage carbohydrate), releasing large amounts of glucose through hydrolysis mediated by 110 and 210 kDa β-glucosidases (*aku*BGLs). Interestingly, such a feeding strategy has attracted attention for producing glucose as a renewable biofuel source [3]. However, to protect themselves against predators, brown algae produce phlorotannin, a secondary metabolite, thereby reducing the digestion of *A. kurodai* by inhibiting the hydrolytic activity of *aku*BGLs. This inhibition has a negative impact on the application of brown algae for producing renewable biofuel sources. As the 110 kDa *aku*BGL is more sensitive to phlorotannin than the 210 kDa BGL [4], we focused on the 110 kDa *aku*BGL in this study (hereafter, *aku*BGL refers to 110 kDa *aku*BGL).

To counteract the antipredator adaptations of algae, herbivores use diverse approaches, such as detoxification, neutralization, defense suppression, and physiological adaptations [5]. *A. kurodai* inhibits the phlorotannin-defense of brown algae through *Eisenia* hydrolysis-enhancing protein (EHEP), a protein from their digestive system that protects *aku*BGL activity from phlorotannin inhibition [4]. Previous studies have shown that incubating *E. bicyclis* with *aku*BGL in the presence of EHEP results in increased glucose production because EHEP binds to phlorotannin and forms an insoluble complex [4].

The *aku*BGL–phlorotannin/laminarin–EHEP system exemplifies the digestion process of *A. kurodai* as well as the defense and antidefense strategies between *E.*

bicyclis and *A. kurodai*. Although the defense/antidefense strategy has been established,

the detailed molecular mechanism of this interplay remains unknown. Further,

phlorotannin inhibition hinders the potential application of brown algae as feedstocks

for enzymatically producing biofuel from laminarin. Thus, understanding the

underlying molecular mechanisms will be beneficial for the application of this system

in the biofuel industry.

Despite the potential use of laminarin hydrolytic enzymes in the biofuel industry,

only a few BGLs of glycoside hydrolases belonging to the GH3 and GH1 family are

known to hydrolyze laminarin (e.g., *Talaromyces amestolkiae* BGL[6], *Ustilago*

esculenta BGL [7], and V*ibrio campbellii* BGL [8] from the GH3 family and

Saccharophagus degradans 2-40[T] BGL [9] from the GH1 family). GH3 is a multidomain

enzyme family characterized by N-terminal $(\beta/\alpha)_8$ (NTD) and C-terminal $(\beta/\alpha)_6$ (CTD)

domains, with or without auxiliary domains [10]; the nucleophile aspartate and the

acid/base glutamate residues exist in the NTD and CTD, respectively. In contrast, the

members of the GH1 family generally share a single $(\beta/\alpha)_8$-fold domain (hereafter

referred to as GH1 domain [GH1D]), and the two glutamic acid catalytic residues are

located in the carboxyl termini of β-strands 4 and 7. Therefore, the two families may

use different substrate-recognition and catalytic mechanisms for laminarin. Intriguingly,

although *aku*BGL possesses laminarin hydrolytic activity and belongs to the GH1

family, its molecular weight is considerably greater than that of other GH1 members.

Sequence analysis has indicated that *aku*BGL consists of ≥2 GH1Ds. Because no

structural information of BGL active on polysaccharides is available, the catalytic

mechanism toward laminarin is unclear.

There is limited information on EHEP, a novel cysteine-rich protein (8.2% of the

amino acid content), because no structural or functional homologous protein exists in

other organisms. EHEP was predicted to consist of three peritrophin-A domains (PADs)

with a cysteine-spacing pattern of $CX_{15}CX_5CX_9CX_{12}CX_{5-9}C$. The PADs consist of

peritrophic matrix proteins, which have been proposed to play an important role in

detoxifying ingested xenobiotics [11]. For instance, *Aedes aegypti* intestinal mucin 1

(*Ae*IMUC1) consists of a signal peptide followed by three PADs with an intervening

mucin-like domain; its expression is induced by blood feeding. *Ae*IMUC1-mediated

blood detoxification during digestion is completed by binding to toxic heme molecules

[12]. Despite the similar domain organization of EHEP and *Ae*IMUC1, their function and

binding partner are completely different, implying their different characteristics.

However, the characteristics of the EHEP-phlorotannin insoluble complex remain

unknown; moreover, it remains unclear why and how EHEP protects *aku*BGL from

phlorotannin inhibition.

In this study, we determined the structures of apo and tannic acid (TNA,

phlorotannin-analog) bound-form of EHEP, as well as *aku*BGL; all isolated from *A.*

kurodai. The structure of EHEP consists of three PADs arranged in a triangle shape,

with TNA bound at the surface of the triangle center. A structural comparison of EHEP

and EHEP–TNA revealed no significant changes in conformation upon TNA binding,

implying that EHEP maintains its structure when precipitated with TNA. Then, we

found the conditions to resolubilize EHEP–TNA precipitate for EHEP recycling. The

obtained *aku*BGL structure suggests that only one GH1D (GH1D2) possesses laminarin

hydrolytic activity; subsequently, ligand-docking experiments demonstrated that

TNA/phlorotannin has a higher binding affinity than laminarin. Our results revealed the

mechanisms by which EHEP protects *aku*BGL from phlorotannin inhibition and

phlorotannin inhibits the hydrolytic activity of *aku*BGL, providing structural support

for the potential application of brown algae for biofuel production.

Results

Effects of TNA on akuBGL activity with or without EHEP

As a chemical defense metabolite of brown algae, phlorotannins are a type of

tannins. It is difficult to isolate a compound from phlorotannins because they are a

group of polyphenolic compounds with different sizes and varying numbers of

phloroglucinol units [13]. Previous studies have reported that the phlorotannin-analog

TNA has a comparable inhibition effect on *aku*BGL to that of phlorotannin [4]. Hence,

we used TNA instead of a phlorotannin to explore phlorotannin binding with EHEP and

*aku*BGL. First, we confirmed TNA inhibition of *aku*BGL activity and clarified the

protective effects of EHEP from TNA inhibition. The inhibition experiments showed

that the galactoside hydrolytic activity of *aku*BGL decreased with increasing TNA

concentration, indicating that TNA inhibits *aku*BGL activity in a dose-dependent

manner (Fig. 1A). Approximately 70% *aku*BGL activity was inhibited at a TNA

concentration of 40 μM. Moreover, protection ability analysis revealed that EHEP

protects *aku*BGL activity from TNA inhibition in a dose-dependent manner, as

indicated by the recovery of the inhibited *aku*BGL activity with increasing EHEP

concentration (Fig. 1B). Further, approximately 80% of *aku*BGL activity was recovered

at an EHEP concentration of 3.36 μM.

Overall structure of EHEP

Considering the lack of known homologous proteins of EHEP, we determined the

structure of natural EHEP using the native-SAD method at a resolution of 1.15 Å, with

a R_{work} and R_{free} of 0.18 and 0.19, respectively (Table 1). The residues A21–V227 of

A21–K229 in purified EHEP (1–20 aa were cleaved during maturation) were

appropriately visualized, whereas two C-terminal residues were disordered. The

structure of EHEP consists of three PADs: PAD1 (N24–C79), PAD2 (I92–C146), and

PAD3 (F164–C221), which are linked by two long loops, LL1 (Q80–N91) and LL2

(R147– G163), and arranged in a triangle shape (Fig. 2A). These three PADs share a

similar structure, with a root-mean-squared difference (RMSD) of 1.065 Å over 46 Cα

atoms and only ~20.3% sequence identity (Figs. 2B and 2C). The three PADs share a

canonical CBM14 fold consisting of two β-sheets containing three N-terminal and two

C-terminal antiparallel β-strands. Additionally, two small α-helices were appended to

the N- and C-terminus in PAD1 and PAD3 but not in PAD2 (Fig. 2B).

Although the Dali server [14] did not provide similar structures using the overall structure

of EHEP as the search model, six structures showed similarities with a single PAD of

EHEP. These structures were the members of the PAD family, including the chitin-

binding domain of chitotriosidase (PDB ID 5HBF) [15], avirulence protein 4 from

Pseudocercospora fuligena [*Pf*Avr4 (PDB ID 4Z4A)] [16] and *Cladosporium fulvum*

[*Cf*Avr4 (PDBID 6BN0)] [17], allergen Der p 23 (PDB ID 4ZCE) [18], tachytitin (PDB ID

1DQC) [19], and allergen Blot 12 (PDB ID 2MFK), with Z-scores of 4.7–8.4, RMSD

values of 1.2–2.8 Å, and sequence identity of 19%–37%. The highest sequence

disparity was detected in PAD2, whereas the greatest structural differences were noted

in PAD3. The $C^{No1}X_{15}C^{No2}X_5C^{No3}X_9C^{No4}X_{12}C^{No5}X_{5-9}C^{No6}$ motif (superscripts and

subscripts indicate the cysteine number and number of residues between adjacent

cysteines, respectively) in each PAD of EHEP formed three disulfide bonds between

the following pairs: C^{No1}–C^{No3}, C^{No2}–C^{No6}, and C^{No4}–C^{No5} (Fig. 2B). Such rich disulfide

bonds may play a folding role in the structural formation of EHEP, with >70% loop

conformation. A similar motif with disulfide bonds was observed in tachycitin [19],

*Pf*Avr4 [16], *Cf*Avr4 [17], and the chitin-binding domain of chitinase [15]. Although these

proteins share a highly conserved core structure, they have different biochemical

characteristics. For example, the chitin-binding domain of human chitotriosidase Avr4

and tachycitin possess chitin-binding activity, but the critical residues for chitin-binding

are not conserved [15, 17, 20], indicating that they employ different binding mechanisms. In

contrast, EHEP and allergen Der p 23 do not possess chitin-binding activity [4, 18]. Thus,

the PAD family may participate in several biochemical functions.

Modification of EHEP

Among our structures, the apo structure2 (1.4 Å resolution) clearly showed that the

cleaved N-terminus of Ala21 underwent acetylation (Fig. 3A), consistent with the

molecular weight results obtained using MALDI–TOF MS [21]. N-terminal acetylation is

a common modification in eukaryotic proteins. Such acetylation is associated with

various biological functions, such as protein half-life regulation, protein secretion,

protein–protein interaction, protein–lipid interaction [22], metabolism, and apoptosis [23].

Further, N-terminal acetylation may stabilize proteins [24]. To explore whether

acetylation affects the protective effects of EHEP on *aku*BGL, we measured the TNA-

precipitating assay of recombEHEP (A21–K229) without acetylation. The results

revealed that recombEHEP precipitated after incubation with TNA at a comparable

level to that of natural EHEP (Fig. 3B), indicating that acetylation is not indispensable

for the phlorotannin binding activity and stabilization of EHEP. Future studies are

warranted to verify the exact role of N-terminal acetylation of EHEP in *A.kurodai*.

TNA binding to EHEP

To understand the mechanism by which TNA binds to EHEP, we determined the

structure of EHEP complexed with TNA (EHEP–TNA) using the soaking method. In

the obtained structure, both $2F_o–F_c$ and $F_o–F_c$ maps showed an electron density blob of

1,2,3,4,6-pentagalloylglucose, a core part of TNA missing the five external gallic acids

(Fig. 4A, Fig. S1A). Previous studies have shown that acid catalytic hydrolysis of TNA

requires a high temperature of 130°C [25]; even with a polystyrene-hollow sphere catalyst,

a temperature of 80°C is required [26]. Therefore, the five gallic acids could not be

visualized in the EHEP–TNA structure most likely due to the structural flexibility of

TNA.

The apo EHEP and EHEP–TNA structures were extremely similar, with an RMSD

value of 0.283 Å for 207 Cα atoms (Fig. S1B). However, the superposition of the two

structures showed a decrement of the loop part in EHEP–TNA, indicating that EHEP is

more stable when bound to TNA. TNA binding caused a slight increase in the α-helix

and β-sheet contents of PAD2 and PAD3 (Fig. S1B). In the EHEP–TNA structure, the

residues C93–Y96 of PAD2 folded into an α-helix and each β-sheet of the first β-strand

in PAD3 elongated by incorporating one residue in the first (G^{176}) and second β-sheets

(S^{186}) and three residues in the third β-sheet ($H^{197}MP^{199}$).

The EHFP–TNA structure revealed that TNA binds to the center of the triangle formed

by the three PADs, a positively charged surface (Fig. 4A and Fig. S1C). The binding

pocket on EHEP surface was formed by the C-terminal α-helix of PAD1, the N-terminal

α-helix of PAD2, and the middle part (loop) of PAD3 assisted by two long linker loops

(LL1 and LL2). TNA was primarily bound to EHEP via hydrogen bonds and

hydrophobic interactions (Fig. 4B). Gallic acid1, 4, and 6 interacted with EHEP via

hydrogen bonds and additional hydrophobic contacts, whereas gallic acid2 and 3 only

hydrophobically interacted with EHEP. The 3-hydroxyl groups of gallic acid1 and 6

individually formed a hydrogen bond with the main chain of G74 and the side chain of

N75 in PAD1. The backbone carbonyl of Y96 and P199 in PAD2 and PAD3,

respectively, formed a hydrogen bond with the 3,5-hydroxyl groups of gallic acid4.

Additionally, some hydrogen bonds were formed between TNA and water molecules.

TNA binding was also stabilized by hydrophobic interactions between the benzene

rings of gallic acid and EHEP. For instance, gallic acid4 and 6 showed alkyl-π

interaction with P77 and P201, respectively; moreover, gallic acid3 and 4 formed

amide-π stackings with P199.

The EHEP–TNA structure clearly showed that TNA binds to EHEP without covalent

bonds and the binding does not induce significant structural changes; thus, we

attempted to recover EHEP from EHEP–TNA precipitates by adjusting the pH. As

hypothesized, re-solubilization of the EHEP–phlorotannin precipitate is pH-dependent

(Fig. 4C). The EHEP–TNA precipitate did not resolubilize at pH 7.0; however, after

incubating for >1 h at pH 7.5, the precipitate started resolubilizing. Most of the

precipitate rapidly resolubilized at an alkaline pH (≥8.0) after incubation for 15 min.

Further, the resolubilized EHEP had the same elution profile as that of the natural EHEP

(Fig. 4D) in SEC, suggesting that resolubilized EHEP maintained the native structure

and its phlorotannin–precipitate activity (Fig. S1E).

Two domains of *aku*BGL

To reveal the structural basis of *aku*BGL recognition of laminarin and inhibition by TNA, we attempted to determine its structure. We soaked crystals in TNA as well as various substrate solutions but finally obtained the optimal resolution using crystal soaking in TNA. There was no blob of TNA in the electron density map of the obtained structure; thus, we considered this structure as the apo form of *aku*BGL.

Two *aku*BGL molecules were observed in an asymmetrical unit (MolA and MolB), without the N-terminal 25 residues (M1–D25), as confirmed using N-terminal sequencing analysis of purified natural *aku*BGL. This N-terminal fragment was predicted to be a signal peptide using the web server SignalP-5.0. The residues L26–P978 of L26–N994 were constructed in both MolA and MolB with glycosylation, whereas the remaining C-terminal residues (A979–M994) could not be visualized as they were disordered. The electron density map of Fo–Fc revealed N-glycosylation at three residues, i.e., N113, N212, and N645 (Figs. S2A, B). N-glycosylation of GH enzymes prevents proteolysis and increases thermal stability [27, 28]. Additionally, a study on β-glucosidase *Aspergillus terreus* BGL demonstrated that N-glycosylation of N224 affected the folding stability, even when it is located close to a catalytic residue [29]. In *aku*BGL, all N-glycosylation sites were present on the surface, far away from the catalytic site. Therefore, we speculate that *aku*BGL glycosylation does not affect its activity. Except for the difference in visualized glycans resulting from glycosylation, MolA and MolB were similar, with a RMSD value of 0.182 for 899 Ca atoms; therefore, we used MolA for further descriptions and calculations.

The structure of *aku*BGL consisted of two independent GH1 domains, GH1D1 (L26–T494) and GH1D2 (D513–P978), linked by a long loop (D495–Y512) (Fig. 5A). There was little interaction between GH1D1 and GH1D2, only in a buried surface area comprising 2% of the total surface (708.9 $Å^2$) (Fig. S2C). GH1D1 and GH1D2 have a sequence identity of 40.47% and high structural similarity with an RMSD value of 0.59

Å for 371 Cα atoms (Fig. S3A up).

Glucosidases of the GH1 family utilize the retaining mechanism with two glutamic

acids for catalyzing glucoside hydrolysis. In general, the distance between the two

catalytic oxygen atoms of the side chain of two glutamic acids is approximately 5 Å [30].

Sequence and structure alignment of GH1D1 and GH1D2 of *aku*BGL with other

members of the GH1 family revealed that the second glutamate is conserved (E404),

but the first glutamate is replaced by D192 in GH1D1. The oxygen atoms of the side

chains between D192 and E404 of GHD1 were 8.4 Å apart. In contrast, GH1D2

conserved two glutamic acids (E675 and E885) at the carboxyl termini of β-strands 4

and 7; the distance between oxygen atoms of E675 and E885 side chains was 5.1 Å

(Fig. S3A down), similar to that of *Neotermes koshunensis* BGL (*Nk*BGL) [31],

Nannochloropsis oceanica BGL (*No*BGL) [32], and *Spodoptera frugiperda* BGL (3.9–

4.9Å) [33]. Furthermore, regarding the two other conserved essential regions for β-

glucosidase activity, namely, glycone-binding site (GBS) and catalysis-related residues

(CR), GH1D1 conserved neither GBS nor CR, whereas GH1D2 conserved both (Fig.

5B). Altogether, we suggest that GH1D1 does not possess catalytic activity. We

expressed and purified the recombinant GH1D1, which did not show any hydrolytic

activity toward O-PNG (Figs. S2D, E), although we could not rule out the effect of N-

glycosylation.

A multi-GH1D assembly has been reported in β-glucosidase *Cj*CEL1A of *Corbicula*

japonica and glycosidase *Lp*MDGH1 of the shipworm *Lyrodus pedicellatus*.

*Lp*MDGH1 has both exo- and endo-glucanase activity and is possibly implicated in

cellulose and hemicellulose digestion. *Cj*CEL1A has two tandem GH1Ds with a

sequence identity of 43.41% [34]. Two catalytic glutamic acids and the residues related to

substrate binding are conserved in the second GH1D, whereas the first GH1 domain

lacks these conserved residues and may play a role in folding the catalytic domain.

*Lp*MDGH1 consists of six GH1Ds, among which GH1D2, 4, 5, and 6 contain the

conserved residues for activity, whereas others do not contain these residues and might

be involved in protein folding or substrate interactions [35].

Structural comparison of GH1D2 with other BGLs, including *Nk*BGL, rice (*Oryza*

sativa L.) BGL (*Os*BGL), and microalgae *No*BGL, revealed the characteristics of each

active pocket (Fig. S3B). *Os*BGL and *No*BGL have a deep, narrow, and straight pocket,

whereas GH1D2 and *Nk*BGL have a broad and crooked pocket. Such active pocket

shapes reflect the substrate preferences of *Os*BGL and *No*BGL; they hydrolyze

laminaribiose with no detectable activity toward laminaritetraose [32, 36]. Furthermore,

the difference in the features of large active pockets between *Nk*BGL and GH1D2,

wherein GH1D2 often possesses an auxiliary site with several aromatic residues bound

to the carbohydrate via CH-π interactions [37], may explain their substrate specificity.

*Nk*BGL efficiently hydrolyzes laminaribiose and cellobiose but has weak hydrolytic

activity toward laminarin [38]. In contrast, the GH1D2 of *aku*BGL has similar activity

levels toward cellobiose and laminarin [39]. Therefore, the GH1D2 of *aku*BGL may

recognize larger substrates than that of other BGLs. Laminarin typically has a curved

conformation; accordingly, narrow- and straight-shaped pockets are incompatible for

binding. Furthermore, we docked GH1D2 with laminaritetraose, wherein the four

glucose units formed extensive contacts with GH1D2. Hydrogen bonds involved the

catalytic residues E675 and E885. In addition, several aromatic residues, such as F677,

F689, Y819, W857, and W935, formed π-π stacking (Fig. S4). Some interacting

residues belonged to GBS and CR sites, such as E675, Y819, E885, and W935.

Additionally, the docking structure revealed that the +4 glucose of laminaritetraose is

located at the auxiliary binding site and that atom O1 of the +4 glucose is positioned

outside the pocket (Fig. S4), implying that the auxiliary binding site with several

aromatic residues (F677, W681, and F689) of GH1D2 facilitates laminarin binding.

Inhibitor binding of *aku*BGL

As we could not obtain the complex structure of *aku*BGL with TNA, we performed

docking calculations of *aku*BGL GH1D2 with TNA to explore the inhibition

mechanism. The docking model of *aku*BGL–TNA showed that seven gallic acid rings

of TNA formed an extensive hydrogen bond network with *aku*BGL in the binding

pocket (Fig. 6). The hydroxyl groups of TNA formed hydrogen bonds with the residues

N552, E675, D735, K739, K759, Q840, T844, D852, and K859 of GH1D2. Moreover,

benzene rings showed hydrophobic interactions with several hydrophobic residues. In

particular, stable π-π stacking was observed between TNA and residues F547, W631,

F689, Y846, W857, and W935. Among these residues, the conserved E675 was the

catalytic residue, and W631, W935, and E934 contributed to GBS and CR sites.

In addition, we performed a docking calculation of GH1D2 with the characteristic

inhibitors eckol and phloroglucinol [40]. The binding mechanisms of eckol and

phloroglucinol were similar to those of TNA but with different contact residues (Fig.

S5). For eckol, the six hydroxyl groups formed hydrogen bonds with residues E675,

D735, E737, K759, E885, and E934. Additionally, residues W631, F677, F689, Y819,

W857, W935, F943, and W927 formed π-π stacking with eckol. For phloroglucinol, the

three hydroxyl groups formed hydrogen bonds with E675, E885, and E934, whereas

residues W631, F689, Y819, W857, W935, and W927 formed π-π stacking with the

benzene ring.

In summary, the three inhibitors inhibited *aku*BGL activity through similar binding

mechanisms to occupy the substrate-binding site, suggesting a reversible competitive

inhibition mechanism. The docking scores of the inhibitors TNA, eckol, and

phloroglucinol were −8.8, −7.3, and −5.7 kcal/mol, respectively, whereas the substrate

laminaritetraose had a docking score of −6.6 kcal/mol. TNA binding and phloroglucinol

had the highest and lowest negative docking score, respectively, indicating that TNA

has a higher binding affinity to *aku*BGL. This finding is consistent with that of a

previous study showing that phloroglucinol binding has a weaker inhibitory activity

than TNA [4].

Discussion

In marine habitats, the ecological interactions between brown algae and herbivores dominate marine ecosystems [41]. The *aku*BGL–phlorotannin/laminarin–EHEP system represents the feeding defense-offense associations between *A. kurodai* and brown algae. We focused on this system to understand the molecular mechanism at the atomic level. In contrast to most GH1 BGLs containing one catalytic GH1 domain, *aku*BGL consists of noncatalytic GH1D1 and catalytic GH1D2. The noncatalytic GH1D1 may act as a chaperone of GH1D2, as we successfully overexpressed GH1D1 but failed to do the same for GH1D2. A similar function has been suggested in *Cj*CEL1A [34] and *LpM*DGH1 [35].

BGLs have different substrate preferences in the degree of polymerization and type of glycosidic bond. In general, BGLs prefer to react with mono-oligo sugars over polysaccharides. For instance, *Os*BGL, *No*BGL, and *Nk*BGL hydrolyze disaccharides (cellobiose and laminaribiose) but display no or weak activity toward polysaccharides (cellulose and laminarin) [32, 36, 38]. The structure of GH1D2 explained the substrate preference for the polysaccharide laminarin. GH1D2 contains an additional auxiliary site composed of aromatic residues (Fig. S3B) in the substrate entrance pocket, which enables it to accommodate a long substrate, contributing to *aku*BGL activity toward laminarin, as supported by docking calculations (Fig. S4). In addition, docking analysis of *aku*BGL GH1D2 with inhibitors (TNA, phlorotannin, eckol, and phloroglucinol) revealed that these inhibitors bound to the substrate-binding site via hydrogen bonds and hydrophobic interactions similar to laminarin. Such binding mechanisms suggest the presence of competitive inhibition to occupy the binding site, consistent with previous research [4].

EHEP, expressed in the midgut of *A. kurodai*, was identified as an antidefense protein, protecting the hydrolysis activity of *aku*BGL from phlorotannin inhibition [4]. Such an ecological balance also exists between plants and their predator mammals and

insects. Similar to brown algae, plants use the toxic secondary metabolite tannins as their defense mechanism against predators, which constitute 5%–10% of dry weight of leaves. In vertebrate herbivores, tannins reduce protein digestion. In phytophagous insects, tannins may be oxidized at an alkaline pH of insect midgut and cause damage to cells. The evolution of plant-herbivore survival competition has led to the development of remarkably unique adaptation strategies. Mammals feeding on plants that contain tannin may overcome this defense by producing tannin-binding proteins, proline-rich proteins, and histatins [42]. Proline constitutes at least 20% of the total amino acid content in proline-rich proteins; for some species, the proportion of proline reaches 40%. Histidine constitutes 25% of total amino acid content in histatins. Both proline-rich proteins and histatins are nonstructural proteins in solution. In caterpillars, the oxidation damage of tannin is reduced by the low oxygen level. Some insects use the peritrophic membrane to transport tannins into the hemolymph, where they are excreted [43]. Additionally, the peritrophic envelop protects insects from tannins by forming an impermeable barrier to tannins [44]. *A. kurodai* uses a similar strategy with mammals by secreting the tannin-binding protein EHEP. Although EHEP has a completely different amino acid composition with proline-rich proteins and histatins, EHEP also binds to phlorotannin. Therefore, EHEP may be a specific counteradaptation that allows *A.kurodai* to feed on brown algae, as there are no homologous proteins in other organisms.

The three PADs of EHEP are arranged in a triangle shape, forming a large cavity on the surface at the triangle center to provide a ligand-binding site. Interestingly, EHEP–TNA crystal packing revealed that each TNA simultaneously binds to three EHEP molecules and crosslinks them together (Fig. S1D); this may be responsible for EHEP precipitation by TNA. EHEP has a positively charged surface at a pH of <6.0, whereas the surface becomes negatively charged at a pH of >7.0 (Fig. S1C). Meanwhile, TNA has a pKa of 4.9–8 [45-47], showing minor negative charges at an acidic pH and the highest negative charge at a pH of >7.0 [48]. Therefore, TNA binds to EHEP at a pH of <6.0 (pH

of crystallization = 4.5), but it shows charge repulsion with EHEP at a pH of >8.0.

Altogether, TNA is protonated and behaves as a hydrogen bond donor when the pH is

below its pKa, whereas when the pH is above its pKa, TNA is deprotonated and the

hydrogen bonding cannot be maintained. As losing hydrogen bonds and increasing

repulsive forces at a pH >8.0, the precipitated EHEP–TNA could not dissolve in the

buffer of pH > 8.0. This pH-induced reversible interaction also occurred in other

proteins, such as BSA, pepsin, and cytochrome C [49]. The phlorotannin members share

a similar structure with TNA; thus, we speculate that the EHEP–phlorotannin complex

also exhibits a pH-induced reversible interaction. *In vivo*, the pH of the digestive fluid

of *A. kurodai* is approximately 5.5, which favors the binding of EHEP to phlorotannin.

In the alkaline hindgut [50], the EHEP–phlorotannin disassociates (Fig. 7), and the

phlorotannin is subsequently excreted from the anus.

Based on the EHEP–TNA structure and docking models of *aku*BGL-

inhibitor/substrate, we proposed a mechanism of phlorotannin inhibition on *aku*BGL

activity and EHEP protection from phlorotannin inhibition (Fig. 7). Because laminarin

lacks the benzene rings essentially to forming π-stacking interactions with EHEP, the

EHEP can be considered no binding with the laminarin. In the absence of EHEP,

phlorotannin occupies the substrate-binding site of *aku*BGL, inhibiting the substrate

from entering the activity pocket and resulting in no glucose production. In the presence

of EHEP, it competitively binds to phlorotannin, freeing the *aku*BGL pocket. Then, the

substrate can enter the active pocket of *aku*BGL and glucose can be produced normally.

The digestive fluid of *A. kurodai* contains EHEP at a high concentration (>4.4 μM) [4],

which is slightly higher than the concentration of EHEP (3.36 μM) that protects

*aku*BGL activity (Fig. 1B). The high concentration of EHEP allows *A. kurodai* feeding

of phlorotannin-rich brown algae. The balance between phlorotannin inhibition and

protection is controlled by the concentrations of phlorotannin and EHEP *in vivo*.

The *aku*BGL–phlorotannin/laminarin–EHEP system is the digestive-defensive-

offensive associations between algae and herbivores. Our study presented the molecular

mechanism of this system at the atomic level, providing a molecular explanation for

how the sea hare *A. kurodai* utilizes EHEP to protect *aku*BGL activity from

phlorotannin inhibition. Further, such a feeding strategy has attracted attention for

producing glucose as a renewable biofuel source, so our studies provide a molecular

basis for the biofuel industry applications of brown algae.

Materials and Methods

EHEP and *aku*BGL preparation

Natural EHEP (22.5 kDa) and *aku*BGL (110 kDa) were purified from *A. kurodai*

digestive fluid as described previously [4]. For crystallization, we further added one step

purification of EHEP using size exclusion chromatography (HiLoad 16/60 Superdex

75, GE Healthcare), for which the column was equilibrated with 20 mM MES–NaOH

buffer (pH 6.5). Obtained EHEP was then concentrated to 15–25 mg/mL using

Vivaspin-4 10K columns (Sartorius, Göttingen, Germany). About *aku*BGL, we

exchanged buffer from 20 mM Tris-HCl pH 7.0 to 20 mM Bis-tris pH 6.0 and

concentrated it to 11 mg/mL using Amicon with a cutoff of 50 kDa.

To verify whether chemical modifications which was indicated by previous study (13)

affect the function of EHEP, we prepared recombinant EHEP (recombEHEP) without

the N-terminal signal peptide (1–20 aa) and chemical modifications [21]. EHEP cDNA

was obtained via reverse transcription–polymerase chain reaction (RT–PCR) using the

total RNA of *A. kurodai* as a template. The reamplified fragment was digested and

ligated to a plasmid derived from pET28a (Novagene, Darmstadt, Germany). We

transformed the plasmid containing recombEHEP into *E. coli* B834(DE3) pARE2 and

expressed it with a C-terminal hexahistidine-tag. The cells were cultured in lysogeny

broth (LB) medium with the antibiotics kanamycin (25 mg/L) and chloramphenicol (34

mg/L) until the optical density at 600 nm (OD_{600}) reached 0.6. Subsequently,

overexpression was induced by adding 0.5 mM isopropyl b-D-L-thiogalactopyranoside

for 20 h at 20 °C. After harvesting by centrifugation, the cells were resuspended in a

buffer containing 50 mM Tris–HCl pH 7.4, 300 mM NaCl, DNase, and lysozyme and

were disrupted via sonication. The insoluble part was removed by centrifugation for 30

min at 40000 ×*g* at 4 °C. We loaded the supernatant onto a 5-mL Histrap HP column

and the recombEHEP was eluted using increasing concentrations of imidazole (0–500

mM). The purified proteins were dialyzed against a solution containing 50 mM Tris–

HCl pH 7.4 and 50 mM NaCl and subsequently loaded onto a Hitrap Q column and

eluted by a linear gradient of a solution containing 50 mM Tris–HCl and 1 M NaCl.

Fractions containing recombEHEP were concentrated and further purified using a gel

filtration column (Hiload 16/60 superdex 75 pg) equilibrated with 20 mM sodium

acetate pH 6.0 and 100 mM NaCl. We collected the fractions containing recombEHEP

and concentrated them to 2.1 mg/mL using Amicon (Merck, American).

TNA binding assay for recombEHEP

456   We measured the binding activity of recombEHEP using precipitation analysis method,

457 as described previously [4]. Briefly, recombEHEP was incubated with TNA at 25 °C for

458 90 min and centrifuged for 10 min at 12000 ×g at 4 °C. Then, we washed the precipitates

459 twice and resuspended them in an SDS–PAGE loading buffer for binding analysis.

Effects of TNA on *aku*BGL activity with or without EHEP

Ortho-nitrophenyl-β-galactoside (ONPG) was used as a substrate to measure *aku*BGL

activity. The reaction system (100μL) included 2.5 mM ONPG, 49 nM *aku*BGL, and

different TNA concentrations in a reaction buffer (50 mM CH_3COONa pH 5.5, 100 mM

NaCl, and 10 mM $CaCl_2$). After incubation for 10 min at 37 °C, 100 μL of methanol

was added to each sample to terminate the reaction. Then, the mixture was centrifuged

for 10 min at 15000 ×g at 4 °C and the supernatant was used for analyzing *aku*BGL

activity via HPLC. To measure the protective effect of EHEP on *aku*BGL, we added

different amounts of EHEP to the reaction system.

Resolubilization of the EHEP–eckol precipitate

470 A mixture of 2 mg of EHEP and 0.4 mg of eckol was incubated at 37 °C for 1 h,

471 followed by centrifugation at 12000 ×g for 10 min, and the supernatant was removed.

472 The sediment was dissolved in 50 mM Tris–HCl buffer at different pH (7.0–9.0) and

473 the absorbance at 560 nm was measured over time.

Crystallization and data collection

The crystallization, data collection, and initial phase determination of EHEP were

described previously [21]. As EHEP precipitates when bound to TNA, we could not

cocrystallize EHEP with TNA. Therefore, we used the soaking method to obtain the

EHEP–TNA complex. Owing to the poor reproducibility of EHEP crystallization, we

used a co-cage-1 nucleant [51] to prepare EHEP crystals for forming the complex with

TNA. Finally, we obtained high-quality EHEP crystals under the reservoir solution

containing 1.0 M sodium acetate, 0.1 M imidazole (pH 6.5) with co-cage-1 nucleant.

Subsequently, we soaked the EHEP crystals in a reservoir solution containing 10 mM

TNA at 37 ℃ for 2 days; further, they were maintained at 20 ℃ for 2 weeks. Next, we

soaked the EHEP crystals in a reservoir solution containing 10 mM phloroglucinol. For

data collection, the crystal was soaked in a cryoprotectant solution containing 20% (v/v)

glycerol along with the reservoir solution. Diffraction data were collected under a cold

nitrogen gas stream at 100 K using Photon Factory BL-17 (Tsukuba, Japan) or Spring

8 BL-41XU (Hyogo, Japan).

For *aku*BGL crystallization, the initial crystallization screening was performed using

the sitting-drop vapor-diffusion method with Screen classics and Classics II

crystallization kits (Qiagen, Hilden, Germany) and PACT kits (Molecular Dimensions,

Anatrace, Inc.) at 20 ℃. Crystallization drops were set up by mixing 0.5 μL of protein

solution with an equal volume of the reservoir solution. The initial crystals were

obtained under condition no. 41 (0.1 M sodium acetate pH 4.5 and 25% polyethylene

glycol [PEG] 3350) of Classics II, no. 13 (0.1 M MIB buffer [25 mM sodium malonate

dibasic monohydrate, 37.5 mM imidazole, and 37.5 mM boric acid], with pH 4.0 and

25% PEG 1500), and no. 37 (0.1 M MMT buffer [20 mM DL-malic acid, 40 mM MES

monohydrate, and 40 mM Tris], with pH 4.0 and 25% PEG 1500) of PACT. After

optimization by varying the buffer pH and precipitant concentration and adding co-

cage1 nucleant, the optimal crystals were obtained using 0.1 M sodium acetate pH 4.5,

and 20% PEG 3350 with a co-cage1 nucleant at a protein concentration of 5.4 mg/mL.

Diffraction data were collected under a cold nitrogen gas stream at 100 K using Photon

Factory BL-1A (Tsukuba, Japan) after cryoprotection by adding glycerol to a 20% final

concentration into the reservoir solution. The optimal resolution of diffraction data was

obtained by soaking a crystal with 5 mM TNA in the reservoir buffer at 37 °C for 4 h.

All datasets were indexed, integrated, scaled, and merged using *XDS/XSCALE* program

[52]. Statistical data collection and process are summarized in Table 1.

Structure determination and refinement

For EHEP structure determination, after initial phasing via the native-SAD method[21,

53], the model was obtained and refined with *auto-building* using *Phenix.autobuil* of

Phenix software suite [54]. The obtained native-SAD structure was used as a model for

rigid body refinement using *phenix.refine* [55] of *Phenix* software suite with a native data

at high resolution of 1.15 Å. The structure of EHEP was automatically rebuilt using

Phenix.autobuil of the *Phenix* software suite again [54]. Several rounds of refinement

were performed using *Phenix.refine* of the *Phenix* software suite [54], alternating with

manual fitting and rebuilding using *COOT* program [56]. The final refinement statistics

and geometry are shown in Table 1.

The structure of the EHEP–TNA complex was determined via the molecular

replacement (MR) method using the EHEP structure as a search model with *Phaser*

of *Pheni*x software suite [57]. The electron density block of TNA was clearly shown in

both $2F_o$–F_c and F_o–F_c maps. Subsequently, TNA structure was manually constructed,

followed by several rounds of refinement using *Phenix.refine* [54], with manual fitting

and rebuilding using *COOT* [56]. We also determined the structure of phloroglucinol-

soaked crystals at a resolution of 1.4 Å via the MR method using the refined EHEP

structure as a search model with *Phaser*, but no electron density block of phloroglucinol

was obtained. Therefore, we referred to this structure as the apo form (apo structure2).

The final refinement statistics and geometry are shown in Table 1.

We determined the structure of *aku*BGL via the MR method using *Phaser* of *Phenix*

software suite [57]. We used one GH domain (86–505 aa) of β-klotho (PDB entry: 5VAN)

[58] as the search model. This GH domain of β-klotho shares 30% sequence identity with

*aku*BGL. Four GH domains of two molecules in an asymmetric unit were found and

subsequently rebuilt with *Phenix_autobuild* of *Phenix* software suite [54]. Finally,

refinement of *aku*BGL structure was performed as described for EHEP.

Docking studies of *aku*BGL with phlorotanins and laminarins

535 We used Schrodinger Maestro program for performing docking studies [59]. First, we

536 superimposed the structure of *Os*BGL mutant complexed with cellotetraose (PDB ID

4QLK) to that of *aku*BGL GH1D2 to define the ligand position in the ligand-binding

cavity. Then, we modified the structure of the *aku*BGL GH1D2 using wizard module

to remove water molecules and add hydrogen atoms for docking. The 2D structures of

the inhibition ligands, including TNA, phloroglucinol, and eckol, were downloaded

from Pubchem [60] and further converted to 3D structures using the LigPrep module of

Schrodinger Maestro program. The structure of the substate laminaritetraose was

extracted from the *Zobellia galactanivorans* β-glucanase–laminaritetraose complex

structure (PDB ID: 4BOW) [61]. Then, a receptor grid was constructed in the center of

the ligand-binding cavity. We performed docking using the Glide standard precision

mode without any constraints. The optimal binding pose was determined using the

lowest Glide score, and docked structures were analyzed using PyMol.

Data Availability

The atomic coordinates were deposited in the PDB with the accession codes as follows:
EHEP with 1.15 Å resolution (8IN3), EHEP with 1.4 Å resolution (8IN4), EHEP
complexed with tannic acid (8IN6), *aku*BGL(8IN1).

Acknowledgments

This work was supported in part by Grant-in-Aid for Scientific Research (B)
(Grant Number 21H01754 To M. Y) and Platform Project for Supporting Drug
Discovery and Life Science Research (Basis for Supporting Innovative Drug Discovery
and Life Science Research (BINDS)) from Japan Agency for Medical Research and

558 Development (AMED) under Grant Number JP18am0101071 and JP19am0101083.

559 We are grateful to the Photon Factor and SPring-8 (No. 2017B2545, 2017A2551,

560 2018B2538) for beam time and the beamline staff for their assistance for data collection.

References

1. Becklin, K.M. A Coevolutionary Arms Race: Understanding Plant-Herbivore

Interactions. *The American Biology Teacher* **70**, 288-292 (2008).

2. Jormalainen, V. & Honkanen, T. in Algal Chemical Ecology 57-89 (2008).

3. Enquist-Newman, M. et al. Efficient ethanol production from brown

macroalgae sugars by a synthetic yeast platform. *Nature* **505**, 239-243 (2014).

4. Tsuji, A., Kuwamura, S., Shirai, A. & Yuasa, K. Identification and

Characterization of a 25 kDa Protein That Is Indispensable for the Efficient

Saccharification of Eisenia bicyclis in the Digestive Fluid of Aplysia kurodai.

PLoS One **12**, e0170669 (2017).

5. Erb, M. & Reymond, P. Molecular Interactions Between Plants and Insect

Herbivores. *Annu Rev Plant Biol* **70**, 527-557 (2019).

6. Mendez-Liter, J.A., de Eugenio, L.I., Prieto, A. & Martinez, M.J. The beta-

glucosidase secreted by Talaromyces amestolkiae under carbon starvation: a

versatile catalyst for biofuel production from plant and algal biomass.

Biotechnol Biofuels **11**, 123 (2018).

7. Nakajima, M., Yamashita, T., Takahashi, M., Nakano, Y. & Takeda, T.

Identification, cloning, and characterization of beta-glucosidase from Ustilago

esculenta. *Appl Microbiol Biotechnol* **93**, 1989-1998 (2012).

8.   Wang, Z. et al. A novel Vibrio beta-glucosidase (LamN) that hydrolyzes the

algal storage polysaccharide laminarin. *FEMS Microbiol Ecol* **91**, fiv087 (2015).

9.   Kim, D.H., Kim, D.H., Lee, S.H. & Kim, K.H. A novel beta-glucosidase from

Saccharophagus degradans 2-40(T) for the efficient hydrolysis of laminarin

from brown macroalgae. *Biotechnol Biofuels* **11**, 64 (2018).

10.   Mohsin, I., Poudel, N., Li, D.C. & Papageorgiou, A.C. Crystal Structure of a

GH3 beta-Glucosidase from the Thermophilic Fungus Chaetomium

thermophilum. *Int J Mol Sci* **20** (2019).

11.   Hegedus, D., Erlandson, M., Gillott, C. & Toprak, U. New Insights into

Peritrophic Matrix Synthesis, Architecture, and Function. *Annual Review of*

Entomology **54**, 285-302 (2009).

12.   Devenport, M. et al. Identification of the Aedes aegypti peritrophic matrix

protein AeIMUCI as a heme-binding protein. *Biochemistry* **45**, 9540-9549

(2006).

13.   Cassani, L. et al. Seaweed-based natural ingredients: Stability of phlorotannins

during extraction, storage, passage through the gastrointestinal tract and

potential incorporation into functional foods. *Food Res Int* **137**, 109676 (2020).

14.   Holm, L. & Rosenstrom, P. Dali server: conservation mapping in 3D. *Nucleic*

Acids Res **38**, W545-549 (2010).

15.   Fadel, F. et al. X-Ray Crystal Structure of the Full Length Human

Chitotriosidase (CHIT1) Reveals Features of Its Chitin Binding Domain. *PLoS*

One **11**, e0154190 (2016).

16.   Kohler, A.C. et al. Structural Analysis of an Avr4 Effector Ortholog Offers

Insight into Chitin Binding and Recognition by the Cf-4 Receptor. *Plant Cell*

28, 1945-1965 (2016).

17.   Hurlburt, N.K., Chen, L.H., Stergiopoulos, I. & Fisher, A.J. Structure of the

Cladosporium fulvum Avr4 effector in complex with (GlcNAc)6 reveals the

ligand-binding mechanism and uncouples its intrinsic function from recognition

by the Cf-4 resistance protein. *PLoS Pathog* **14**, e1007263 (2018).

18.   Mueller, G.A. et al. Serological, genomic and structural analyses of the major

mite allergen Der p 23. *Clin Exp Allergy* **46**, 365-376 (2016).

19.   Suetake, T. et al. Chitin-binding proteins in invertebrates and plants comprise a

common chitin-binding structural motif. *J Biol Chem* **275**, 17929-17932 (2000).

20.   Madland, E., Crasson, O., Vandevenne, M., Sorlie, M. & Aachmann, F.L. NMR

and Fluorescence Spectroscopies Reveal the Preorganized Binding Site in

Family 14 Carbohydrate-Binding Module from Human Chitotriosidase. *ACS*

Omega **4**, 21975-21984 (2019).

21.   Sun, X. et al. Crystallographic analysis of Eisenia hydrolysis-enhancing protein

using a long wavelength for native-SAD phasing. *Acta Crystallographica*

Section F-Structural Biology Communications **76**, 20-24 (2020).

22.   Silva, R.D. & Martinho, R.G. Developmental roles of protein N-terminal

acetylation. *Proteomics* **15**, 2402-2409 (2015).

23.   Hollebeke, J., Van Damme, P. & Gevaert, K. N-terminal acetylation and other

functions of Nalpha-acetyltransferases. *Biol Chem* **393**, 291-298 (2012).

24.   Lange, P.F. & Overall, C.M. TopFIND, a knowledgebase linking protein termini

with function. *Nat Methods* **8**, 703-704 (2011).

25.   Jie Fu, J.x., Zhang, Y. & Lu, X. A Greener Process for Gallic Acid Production

from Tannic Acid Hydrolysis with Hydrochloric Acid. *Asian Journal of*

Chemistry **27**, 3328-3332 (2015).

26.   Luo, Q. et al. A novel green process for tannic acid hydrolysis using an

internally sulfonated hollow polystyrene sphere as catalyst. *RSC Advances* **8**,

17151-17158 (2018).

27.   Amore, A. et al. Distinct roles of N- and O-glycans in cellulase activity and

stability. *Proc Natl Acad Sci U S A* **114**, 13667-13672 (2017).

28.   Han, C. et al. Improvement of the catalytic activity and thermostability of a

hyperthermostable endoglucanase by optimizing N-glycosylation sites.

Biotechnol Biofuels **13**, 30 (2020).

29.   Wei, W. et al. N-glycosylation affects the proper folding, enzymatic

characteristics and production of a fungal beta-glucosidas. *Biotechnol Bioeng*

110, 3075-3084 (2013).

30.   Hayashi, Y. et al. Klotho-related protein is a novel cytosolic neutral beta-

glycosylceramidase. *J Biol Chem* **282**, 30889-30900 (2007).

31. Jeng, W.Y. et al. High-resolution structures of Neotermes koshunensis beta-

glucosidase mutants provide insights into the catalytic mechanism and the

synthesis of glucoconjugates. *Acta Crystallogr D Biol Crystallogr* **68**, 829-838

(2012).

32. Dong, S. et al. Structural insight into a GH1 beta-glucosidase from the

oleaginous microalga, Nannochloropsis oceanica. *Int J Biol Macromol* **170**,

196-206 (2021).

33. Tamaki, F.K. et al. Using the Amino Acid Network to Modulate the Hydrolytic

Activity of beta-Glycosidases. *PLoS One* **11**, e0167978 (2016).

34. Sakamoto, K., Uji, S., Kurokawa, T. & Toyohara, H. Molecular cloning of

endogenous beta-glucosidase from common Japanese brackish water clam

Corbicula japonica. *Gene* **435**, 72-79 (2009).

35. Sabbadin, F. et al. Uncovering the molecular mechanisms of lignocellulose

digestion in shipworms. *Biotechnol Biofuels* **11**, 59 (2018).

36. Opassiri, R. et al. Characterization of a rice β-glucosidase highly expressed in

flower and germinating shoot. *Plant Science* **165**, 627-638 (2003).

37. Hudson, K.L. et al. Carbohydrate-Aromatic Interactions in Proteins. *J Am Chem*

Soc **137**, 15152-15160 (2015).

38. Ni, J., Tokuda, G., Takehara, M. & Watanabe, H. Heterologous expression and

enzymatic characterization of .BETA.-glucosidase from the drywood-eating

termite, Neotermes koshunensis. *Applied Entomology and Zoology* **42**, 457-463

(2007).

39.   Tsuji, A., Tominaga, K., Nishiyama, N. & Yuasa, K. Comprehensive enzymatic

analysis of the cellulolytic system in digestive fluid of the Sea Hare Aplysia

kurodai. Efficient glucose release from sea lettuce by synergistic action of 45

kDa endoglucanase and 210 kDa ss-glucosidase. *PLoS One* **8**, e65418 (2013).

40.   Jung, H.A., Oh, S.H. & Choi, J.S. Molecular docking studies of phlorotannins

from Eisenia bicyclis with BACE1 inhibitory activity. *Bioorg Med Chem Lett*

20, 3211-3215 (2010).

41.   Amsler, C.D. & Fairhead, V.A. in Advances in Botanical Research, Vol 43:

Incorporating Advances in Plant Pathology, Vol. 43. (ed. J.A. Callow) 1-91

(2006).

42.   Shimada, T. Salivary proteins as a defense against dietary tannins. *J Chem Ecol*

32, 1149-1163 (2006).

43.   War, A.R. et al. in Co-Evolution of Secondary Metabolites 795-822 (2020).

44.   Barbehenn, R.V. & Peter Constabel, C. Tannins in plant-herbivore interactions.

Phytochemistry **72**, 1551-1565 (2011).

45.   Lin, D. et al. The effect of ionic strength and pH on the stability of tannic acid-

facilitated carbon nanotube suspensions. *Carbon* **47**, 2875-2882 (2009).

46.   Ge, D., Yuan, H., Shen, Y., Zhang, W. & Zhu, N. Improved sludge

dewaterability by tannic acid conditioning: Temperature, thermodynamics and

684  mechanism studies. *Chemosphere* **230**, 14-23 (2019).

685 47. Yi, Z. et al. Green, effective chemical route for the synthesis of silver nanoplates

686  in tannic acid aqueous solution. *Colloids and Surfaces A: Physicochemical and*

687  *Engineering Aspects* **392**, 131-136 (2011).

688 48. Dultz, S., Mikutta, R., Kara, S.N.M., Woche, S.K. & Guggenberger, G. Effects

689  of solution chemistry on conformation of self-aggregated tannic acid revealed

690  by laser light scattering. *Sci Total Environ* **754**, 142119 (2021).

691 49. Han, Y. et al. Polyphenol-Mediated Assembly of Proteins for Engineering

692  Functional Materials. *Angew Chem Int Ed Engl* **59**, 15618-15625 (2020).

693 50. Lemke, T., Stingl, U., Egert, M., Friedrich, M.W. & Brune, A. Physicochemical

694  conditions and microbial activities in the highly alkaline gut of the humus-

695  feeding larva of Pachnoda ephippiata (Coleoptera: Scarabaeidae). *Appl Environ*

696  *Microbiol* **69**, 6650-6658 (2003).

697 51. Yao, M. & Li, L. (NATIONAL UNIVERSITY CORPORATION

698  HOKKAIDO UNIVERSITY 2022).

699 52. Kabsch, W. Xds. *Acta Crystallogr D Biol Crystallogr* **66**, 125-132 (2010).

700 53. Yu, J., Shinoda, A., Kato, K., Tanaka, I. & Yao, M. A solution-free crystal-

701  mounting platform for native SAD. *Acta Crystallogr D Struct Biol* **76**, 938-945

702  (2020).

703 54. Adams, P.D. et al. PHENIX: a comprehensive Python-based system for

704  macromolecular structure solution. *Acta Crystallogr D Biol Crystallogr* **66**,

705  213-221 (2010).

706 55. Afonine, P.V. et al. Towards automated crystallographic structure refinement

707  with phenix.refine. *Acta Crystallogr D Biol Crystallogr* **68**, 352-367 (2012).

708 56. Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics.

709  *Acta Crystallogr D Biol Crystallogr* **60**, 2126-2132 (2004).

710 57. McCoy, A.J. et al. Phaser crystallographic software. *J Appl Crystallogr* **40**, 658-

711  674 (2007).

712 58. Lee, S. et al. Structures of beta-klotho reveal a 'zip code'-like mechanism for

713  endocrine FGF signalling. *Nature* **553**, 501-505 (2018).

714 59. Sastry, G.M., Adzhigirey, M., Day, T., Annabhimoju, R. & Sherman, W. Protein

715  and ligand preparation: parameters, protocols, and influence on virtual

716  screening enrichments. *Journal of Computer-Aided Molecular Design* **27**, 221-

717  234 (2013).

718 60. Wang, Y.L. et al. PubChem: a public information system for analyzing

719  bioactivities of small molecules. *Nucleic Acids Research* **37**, W623-W633

720  (2009).

721 61. Labourel, A. et al. The beta-glucanase ZgLamA from Zobellia galactanivorans

722  evolved a bent active site adapted for efficient degradation of algal laminarin. *J*

723  *Biol Chem* **289**, 2027-2042 (2014).

725 Fig 1

Figure 1. Galactoside hydrolytic activity of *aku*BGL toward ortho-Nitrophenyl-β-

galactoside. (A) The hydrolytic activity of akuBGL with TNA at different

concentrations. (B) The hydrolytic activity of akuBGL (0.049 μM) with 40 μM TNA

and EHEP at different concentrations. The average and standard deviation of the

relative activity were estimated from three independent replicates (N = 3).

Fig 2

Figure 2. EHEP structure. (A) Cartoon representation of EHEP. The three PAD

domains are colored green, light blue, and pink, respectively. Linker long loop1 (LL1)

and loop2 (LL2) are colored yellow and blue. (B) Structural superposition of the three

PAD domains of EHEP. The three domains are colored as in (A). The disulfide bonds

are shown as yellow sticks. (C) Sequence alignment of three PAD domains. Alignment

was performed by CLUSTALW and displayed with ESPript3.

Fig 3

Figure. 3. (A) Acetylation modification on the N-terminal residue A21. The structure

is shown as sticks with an omitted map of A21-ace at a 3.3 σ level (blue-white). (B)

TNA binding activity of recombEHEP. SDS–PAGE was run using a mixture of

recombEHEP and TNA.

Figure 4. Structure of EHEP-TNA. (a) The overall structure of EHEP-TNA. (A) the

overall structure of EHEP-TNA. EHEP and TNA are shown by the cartoon and stick

model, respectively. EHEP is colored as in Fig. 1. The C and O atoms of TNA are

colored lemon and red, respectively. (B) Interaction of TNA (ball-stick in same color

as (A)) with EHEP (cartoon in same color as (A)) in EHEP-TNA structure. The residues

of EHEP in contact are labeled and shown by a ball-stick with N, O, and S atoms in

blue, red, and brown, respectively. The C and O atoms of TNA are colored the same as

(A), lemon and red, respectively. Dashed lines show hydrogen bonds. The water

molecules stabilizing TNA was shown as light orange spheres. (C) Effect of pH on

758   resolubilization of an EHEP–eckol precipitate. Buffers with pH 9.0, 8.0, 7.5, and 7.0

759   are presented as hollow square, solid circle, hollow circle, and solid square, respectively.

760   (D) The EHEP–eckol precipitate was dissoloved in 50 mM Tris–HCl (pH 8.0) and

761   analyzed by Sephacryl S-100.

763   Fig 5

Figure 5. Structure of *aku*BGL. (A) Overall structure. The GH1D1 (light blue) and

GH1D2 (cyan) domains are linked by a long loop (linker-loop) colored in pink. The N-

linked glycans were shown in the orange stick. (B) Residues superposition of the GBS

and CR sites of the domains GH1D1 (cyan), GH1D2 (light blue) with β-glucosidase

structures from termite *Neotermes koshunensis* (*Nk*BGL, grey), β-glucosidase from rice

(*Os*BGL, gray), β-glucosidase from *Bacillus circulans* sp. *Alkalophilus* (grey). Only the

residues numbers of GH1D1 (cyan), GH1D2 (light blue), and *Nk*BGL (grey) are shown

for clarity.

774　Fig 6

Figure 6. Docking model of *aku*BGL with TNA. (A) Detailed interaction between

*aku*BGL and TNA in the docking model. TNA is shown in the green stick model. The

778   hydrogen bonds are shown as dashed lines. (B) A 2D diagram of the interaction between

779   *aku*BGL and TNA shown in (A). The hydrogen bonds are shown as dash lines, and the

780   hydrophobic contacts as circular arcs.

781   Fig 7

Figure 7. Proposed molecular mechanisms. Proposed molecular mechanisms of TNA

inhibition of *aku*BGL activity and EHEP's protective effects of *aku*BGL in *aku*BGL–

phlorotannin/laminarin–EHEP system (light purple triangle). The digestive tract of

A.kurodai consists of foregut (blue), midgut (purple), and hindgut (blue) (top). The bar

chart above depicts the pH of the digestive tract, with pink denoting acid and blue

denoting alkalinity.

790 **Table 1** X-ray data collection and structure-refinement statistics.

	Native-SAD of EHEP	EHEP1	EHEP2	EHEP-TNA	*aku*BGL
Data Collection					
Beamline	PF BL17A	PF BL17A	Spring 8 BL-41XU	PF BL17A	PF BL1A
Wavelength (Å)	2.1000	0.9800	1.0000	0.9800	1.0000
Resolution range (Å)	46.61-2.48 (2.54-2.48)	46.56–1.15 (1.20–1.15)	47.02- 1.4 (1.45-1.4)	46.84–1.9 (1.97–1.9)	49.65-2.7 (2.80-2.7)
Space group	$P2_12_12_1$	$P2_12_12_1$	$P2_12_12_1$	$P2_12_12_1$	$P6_2$
Unit-cell parameters a, b, c (Å)	42.2, 65.3, 66.5	42.2, 65.3, 66.5	40.6, 65.6, 67.5	42.5, 65.4, 67.2	191.7, 191.7, 112.6
Completeness (%)	97.7 (87.5)	93.4 (75.8)	99.30 (97.98)	99.9 (99.7)	99.9 (99.1)
Redundancy	99.7 (38.9)	6.6 (6.1)	6.4 (5.8)	6.4 (6.4)	10.7 (10.9)
Average $I/\sigma(I)$	105.1 (74.4)	19.28 (1.97)	14.34 (3.41)	15.39 (1.83)	10.69 (2.57)
R_{meas}(%)[a]	6.9 (14.2)	7.3 (90.5)	8.6 73.7(55.3)	8.9 (89.4)	19.4 (83.0)
$CC^{1/2}$ (%)	100 (99.8)	99.9（70.4）	99.8 (86.2)	99.9（73.7）	99.5 (84.2)
Ano/Sig	3.1 (1.7)				
Molecules/ asymmetric unit	1	1	1	1	2
Refinement					
R_{work}[b]/R_{free}[c] (%)		18.19/18.91	16.57/18.39	19.87/23.54	18.39/21.98
No. of atoms		1955	1861	1732	15595
No. of residues		1600	1573	1580	15256
No. of water molecules		343	273	87	96
No. of Ligands		12	15	67	243
RMSD from ideality					
bond length (Å)		0.005	0.006	0.008	0.004
bond angle (°)		0.84	0.84	0.91	0.65
Ramachandran plot (%)					

Favoured	99.02	98.54	98.52	96.16
Allowed	0.98	1.46	1.48	3.74
Outliers	0.00	0.00	0.00	0.11
PDB accession code	8IN3	8IN4	8IN6	8IN1

The highest resolution shell is shown in parentheses.

$^{a}R_{meas} = \Sigma_{hkl}\{N(hkl) / [N(hkl) -1] \}^{1/2} \Sigma_i|I_i(hkl)-< I(hkl)> | /\Sigma_{hkl} \Sigma_i|I_i(hkl)$, where where

$I_i(hkl)$ is the ith observation of the intensity of reflection *hkl* and $\langle I(hkl)\rangle$ is the mean

over n observations.

$^{b}R_{work} = \Sigma_{hkl}||F_{obs}(hkl)| - |F_{calc}(hkl)||/\Sigma_{hkl}|F_{obs}(hkl)|$.

$^{c}R_{free}$ was calculated with an approximate 5% fraction of randomly selected reflections

evaluated from refinement.

Supporting information

 Fig.S1

Figure S1. EHEP–TNA structure and structural comparisons. (A) TNA structure (stick

model) in EHEP–TNA with an omitted map countered at the 2.0 σ level. The O and C

atoms are colored red and lemon, respectively. (B) Structure superposition of apo EHEP

(orange) with EHEP–TNA (green). Two arrows mark the conformational changes in a

β-sheet of PAD1 and an α-helix of PAD2, respectively. (C) Electrostatic potential of the

EHEP surface in the EHEP–TNA complex at different pH values. (D) Diagram of

EHEP–TNA packing in the crystal structure. The triangle represents the EHEP, with the

three colors denoting the three PAD domains, as in Fig. 3a. TNA is shown in a lemon

pentagram. (E) Activity of the resolubilized EHEP. The EHEP-eckol precipitate was

resolubilized in Tris–HCl (pH 8.0) and then the supernatant was purified by Sephacrys

S-100 column to measure eckol-binding activity.

815 Fig.S2

Figure S2. *aku*BGL structure. (A) Two *aku*BGL molecules in the asymmetry unit,

colored green and grey, respectively. The glycosylation sites are shown as orange sticks,

with O and N atoms in red and blue, respectively. (B) The glycosylation chains with

omitted density maps are countered at the 2.0 σ level. (C) The interface between the

GH1D1 (cyan) and GH1D2 (blue). The key interactions between the two domains are

shown as black dashed lines. (D) Size exclusion chromatogram of the purified GH1D1

domain. The inset shows the SDS–PAGE analysis of the GH1D1 domain. (E)

Galactoside hydrolytic activity of the GH1D1 toward ortho-Nitrophenyl-β-galactoside.

The average and standard deviation of the relative activity were estimated from three

independent replicates (N = 3).

Figure S3. Structural comparison of BGLs. (A) Structure superposition of

GH1D1(cyan) and GH1D2 (blue). The enlarged picture shows the distance of

conceivable catalytic residues in GH1D1 and GH1D2. (B) Surface representations of

GH1D2, *Nk*BGL, *Os*BGL, and *No*BGL with the aromatic and catalytic residues colored

green and red, respectively. The red arrow indicates the location of the auxiliary site of

GH1D2. The active pockets are highlighted by a black circle on each surface.

Fig. S4

Figure S4. The model of GH1D2 docking with the substrate laminaritetraose. GH1D2

is shown as a grey surface representation and laminaritetraose as marine sticks. The

aromatic and catalytic residues of GH1D2 are colored green and red, respectively.

Fig. S5

eckol

phloroglucinol

Figure S5. The models of GH1D2 docking (A) With eckol; (B) With phloroglucinol.

845   The left panel shows the 3D structures, and the right panel shows the 2D diagrams. The

846   C, N, and O atoms of residues are colored light blue, dark blue, and red, respectively.

847   The C atoms of eckol and phloroglucinol are shown in orange and yellow, respectively.

848