Clustering temporal step-counting patterns for 24

hours with machine learning revealed potential

heterogeneity in the categorization by a traditional

tertile procedure.

Saida Salima Nawrin[1,¶], Hitoshi Inada[1¶*,#a] , Haruki Momma[2], and Ryoichi

nagatomi[1*, 2]

[1] Laboratory of Health and Sports Sciences, Tohoku University Graduate
School of Biomedical Engineering, Sendai, Miyagi, Japan

[2] Department of Medicine and Science in Sports and Exercise, Tohoku
University Graduate School of Medicine, Sendai, Miyagi, Japan

[#a] Current address: Department of Biochemistry & Cellular Biology,
National Center of Neurology and Psychiatry, Kodaira, Tokyo, Japan

*Corresponding author
E-mail: hinada@med.tohoku.ac.jp (HI)
E-mail: nagatomi@med.tohoku.ac.jp (RN)

[¶]These authors contributed equally to this work.

Abstract

Objectives

Physical activity is a crucial aspect of health benefits in the public society. Although studies on the temporal physical activity patterns might lead to the protocol for efficient intervention/program, a standardized procedure to determine and analyze the temporal physical activity patterns remains to be developed. Here, we attempted to develop a procedure to cluster 24-hour patterns of physical activity as step counts measured with an accelerometer-based wearable sensor.

Methods

Data was collected from 42 healthy participants, comprising 35 males and 7 females, at the Sendai Oroshisho center in 2008. Using unsupervised machine learning, specifically the kernel k-means algorithm with the global alignment kernel was applied on a total of 815 days from 42 participants, and 6 activity patterns were identified. Further, the probability of each 24-hour step-counting pattern was calculated for every participant., and was used in MATLAB to apply spectral clustering, and 5 activity behaviors were identified.

Results

We could identify six 24-hour step-counting patterns and five daily step-behavioral clusters. When the amount of physical activity was categorized into tertile groups reflecting highly active, moderately active, and low active, each tertile group consisted of different proportions of six 24-hour step-counting patterns.

Conclusions

Our study introduces a novel approach using an unsupervised machine learning method to

categorize daily hourly activity, revealing six distinct step counting patterns and five

clusters representing daily step behaviors. Our procedure would be reliable for finding and

clustering physical activity patterns/behaviors and reveal heterogeneity in the

categorization by a traditional tertile procedure using total step amount.

Keywords: Physical activity, step-counting patterns, daily step behaviors,

unsupervised   machine   learning.

Introduction

55   Physical activity is a crucial aspect of maintaining good health and well-being.

World Health Organization (WHO) defines "physical activity as any bodily movement

produced by skeletal muscles that requires energy expenditure" and recommends at least

75-150/150-300 minutes of vigorous/moderate-intensity aerobic physical activity

throughout the week (Organization WHO. Physical Activity 2022). WHO also mentioned

that physical inactivity is one of the leading risk factors for global mortality, causing an

estimated 3.2 million deaths per year (Organization WHO Physical Inactivity). Many

observational and intervention studies suggested that regular physical activity has been

linked to numerous physical and mental health benefits, including reducing the risk of

chronic diseases, improving cardiovascular health, maintaining a healthy weight,

improving mental health, and enhancing the overall quality of life (Chekroud et al., 2018;

Kraus et al., 2019; Madigan et al., 2021; Pearce et al., 2022; Piercy et al., 2018).

However, a recent systematic review and meta-analysis of randomized controlled trials

indicated that interventions to maintain physical activity behavior was effective in short

periods (at least three months) but provided a small impact on the long-term maintenance

of physical activity (Madigan et al., 2021). Therefore, it is critical to figure out efficient

ways integrated into daily life to promote and maintain physical activity for the

improvement of the health and well-being of the public.

A historical topic in public health is what kind, how much, and how intense physical

activity is required for health benefits (Powell, Paluch, & Blair, 2011). One of the

simplest ways to increase and maintain physical activity is to walk/move more and longer

(Morris & Hardman, 1997; National Heart, Lung, and Blood Institute. Tips for Getting

Active; Ogilvie et al., 2007) as shown in the 10,000-steps campaign (Eyler, Brownson,

Bacak, & Housemann, 2003; Hatano, Y. 1993 ; Tudor-Locke & Bassett, 2004). A recent

meta-analysis reported that taking more steps per day was associated with lower mortality

risk, where the risk plateaued at approximately 6,000–8,000 steps or 8,000–10,000 steps

per day for older (≥60 years) or younger adults (<60 years), respectively (Paluch et al.,

2022). Most of the previous research on physical activity has concentrated on how the

total amount of activity based on its intensity and duration is linked to health (Amagasa et

al., 2018; Brailey et al., 2022; Lu et al., 2022). However, analyzing only the total amount

of physical activity might overlook a complete understanding of the activity behavior

since the metrics of the total amount of physical activity tends to exclude or

underestimate the effect of temporal variations of physical activity.

Recently, several studies focused on the 24-hour temporal physical activity

patterns using a narrower criterion for physical activity, such as steps or body movements

objectively measured with an accelerometer (Aqeel et al., 2021; Niemelä et al., 2019;

Smagula et al., 2015; Smagula, Krafty, Thayer, Buysse, & Hall, 2018; Smagula et al.,

2022). An earlier study that analyzed the relationship between activity rhythm

disturbances and depression identified eight sub-clusters in older adults (≥65 years)

(Smagula et al., 2015). Late or combined early/compressed/dampened activity rhythms

may independently contribute to depression symptom development (Smagula et al., 2015).

Another recent report using activity counts for seven days in US adults indicated that,

although the lowest physical activity counts cluster showed a higher obesity index

including body mass index and waist circumference than the other clusters showing

higher activity counts, the clusters also expressed different temporal patterns of physical

activity (Aqeel et al., 2021). Moreover, an increase in cardiovascular disease risk was

observed in the inactive cluster than the other clusters (evening active, moderately active,

and very active) (Niemelä et al., 2019) from the data of The Northern Finland Birth

Cohort 1966 study (Northern Finland Birth Cohorts | University of Oulu). The latest

report combining activity timing/period and pattern robustness identified sub-groups with

progressive depression symptoms and impaired cognitive performance in aging (Smagula

et al., 2022). Therefore, understanding temporal physical activity patterns would help

extract informative health outcomes.

In the previous studies above, participants were first clustered by several selected

features indicating physical activity patterns, and the physical activity patterns in 24

hours were analyzed. However, those procedures have difficulty analyzing temporal

physical activity patterns directly because of their methodological restrictions, in which

participants were first clustered by extracted individual features of the physical activity

patterns and then analyzed. In this study, we propose a novel procedure for the direct

clustering of physical activity based on the temporal step-counting patterns with

unsupervised machine learning irrespective of the total volume of physical activity.

Subsequently, we could identify five step-behavioral clusters of participants, based on the

proportion of each step-counting pattern. Furthermore, we analyzed the proportion of the

temporal step-counting patterns and step-behavioral clusters of groups categorized by a

traditional tertile procedure The results revealed potential heterogeneity in categorizing

by the tertile procedure using total step amount. Our approach could provide a useful

procedure to analyze temporal patterns of physical activity in a direct manner.

Materials and Methods

Participants and data collection

We used the anonymized activity counting data obtained in the previous study (Guo

et al., 2012), in which participatns were recruited in August 2008 throughout an

annual health examination at the Sendai Oroshisho center, provided and agreed with

informed consent for their data to be analyzed (Guo et al., 2012). The data collected

from forty-two participants between August and September 2008 in Sendai Oroshisho

center was used for a series of analyses. The Institutional Review Board of the Tohoku

University Graduate School of Medicine have approved the current study protocol

(Approval number: 2019-1-394).

Step count data was continuously collected by a hip-worn triaxial accelerometer

(Fig S1, Nipro Welsupport, model no wat3663, Japan) for up to 25 days. Data from

participants with less than three days was excluded for statistical reasons (Hart, Swartz,

Cashin, & Strath, 2011). The demography of the forty-two participants (35 males and 7

females). Each participant has 19 ± 4 days of data. The data sampling rate was 1 Hz.

Eight types of activities are recorded as below: 0, Resting; 1, Moving while standing; 2,

Going down using a vehicle; 3, Going up using vehicle; 4, Going down using stairs; 5,

Walking on the ground level; 6, Going up using stairs; 7, Running; 8, Lying down.

Numbers of indices 5-7 were counted as steps. The other indices were counted as zero.

143   Software for data processing and visualization

144   MATLAB (2021a, MathWorks, MA) and Python (3.11.0) were used for data

145   processing and visualization.

147   Twenty-four-hour step-counting pattern

The activity counting data at 1 Hz sampling rate was first converted hourly for each 24-

hour. A total of 815 days of hourly data was used to cluster the 24-hour step-counting

patterns by unsupervised machine learning using tslearn, a Python package for time-

series analysis (Tavenard et al., 2020). The kernel k-means algorithm was used for

clustering with the global alignment kernel (GAK). The cluster number was determined

by elbow plot using distortion values. Six clusters of 24-hour physical activity patterns

were obtained.

156   Dominant daily step behavior

157   First, the probability of each 24-hour step-counting pattern is calculated for each

158   participant based on the formula below.

$$Probability\ of\ each\ 24-hour\ step-counting\ pattern$$

$$= \frac{Number\ of\ a\ paticular\ pattern}{Total\ number\ of\ days}$$

161   Vectors of the probability of each 24-hour step-counting pattern (with six values)

162   were used to cluster the daily step behaviors by unsupervised machine learning using a

163   spectral clustering algorithm with MATLAB. Five clusters of daily step-behavior were

164   obtained.

Tertile clustering of daily physical activity based on the daily

step amount

168   The daily step amount for 815 days was divided into the tertile group (low,

169   medium, and high). Each group is defined as:

170   Low: Daily step amount <= 4127

171   Medium: 4127 < Daily step amount < 6946

172   High: Daily step amount >= 6946

173 **Statistical analysis**

174   Pearson chi-squared test was performed with (JMP pro 16.0.0) for all statistical

175   analyses where P<.05 is considered significant.

Results

The data set was collected from forty-two healthy participants (35 males and 7 females) from August to September 2008 in Miyagi prefecture, Japan. The demography of the participants is shown in Table 1.

Twenty-four-hour step-counting pattern

Step count data obtained with a triaxial accelerometer was first converted hourly for each 24-hour and then applied for the clustering with unsupervised machine learning (See Methods). Fig 1 shows six 24-hour step-counting patterns identified: all-day (AD), bi-phasic (BP), morning (M), evening (E), irregular morning (IM), and irregular night (IN). AD showed a continuous high step count from 9:00 to 19:00. BP showed increased step counts from 8:00 to 21:00 but had two peak periods at 9:00 and 19:00. Step counts between 10:00-16:00 was slightly low in BP. M started actively around 6:00, kept on high step counts until 14:00, and gradually decreased in activity after 15:00. By contrast, E started active afternoon and showed peak step counts between 16:00-18:00. IM showed high step counts mainly in the morning from 5:00 to 13:00 and decreases in activity after 15:00. IN showed high step-counts from 19:00 to midnight. Table 2 summarizes the proportion of each step-counting pattern. Among 815 days, AD was the most dominant pattern (330/815, 40.5 %), and the BP pattern followed (146/815, 17.9%). M and E patterns were 15.8 % and 13.0%, respectively. IM and IN were minor patterns, less than 10% (7.9% and 4.9 %, respectively). The proportion of each pattern showed significant differences except for a difference between IM and IN (Table S1). Each step-counting

pattern showed a different step-count amount (Table 3). The AD showed the highest step-

count amount per day (7020.0 ± 3271.4 counts). The M and BP patterns were similar, and

the E pattern followed. The IN and IM patterns showed lower step count amounts. AD,

BP, and M patterns differed significantly from E, IM, and IN (Table S2). Among E, IM,

and IN patterns, a significant difference was observed between E and IM patterns.

Regarding duration, the BP pattern showed the longest duration of activity per day (14.54

± 2.61 hours). The AD and M followed. Generally, the AD and BP patterns shared 58.4

% of total days and showed higher/longer step count amounts per day. The M pattern was

more active than the E pattern. The IM and IN shared a minor proportion in days (12.8

%) and showed the lowest step-count amount. All the pairs show a difference ($p < 0.05$)

except BP-AD (Table S3).

211   Clustering dominant daily step-behaviors

According to the six 24-hour step-counting patterns, we further clustered

participants based on the probability of each pattern. We found five daily step-behavioral

clusters; AD dominant, AD + BP, BP dominant, AD + E, and M dominant (Fig 2). Half

of the participants (21) belong to the AD-dominant (S4 Table), consistent with the result

that AD was the most frequent step-counting pattern. The AD+BP was the following

frequent behavior. The other daily step behaviors shared minor proportions. There were

no IM dominant or IN dominant step behaviors due to their minor proportions in the 24-

hour step-counting patterns. Interestingly, mean values of step count amounts for each

step behavior were similar (Table S4).

Table 4 represents the probabilities of six step-counting patterns in each daily

step-behavioral cluster. The AD dominant step-behavior accounts for most of the AD

pattern (0.5515 ± 0.1492). AD dominant step behavior also consists of a significant

portion with the M pattern (0.1227 ± 0.0888). Other step-counting patterns are less.

AD+BP step-behavior is characterized by a significant share of AD and BP step-counting

patterns (0.4031 ± 0.1021 and 0.2964 ± 0.0586, respectively). AD+BP step-behavior also

contains a significant amount with the M step-counting pattern (0.1587±0.0748), while

other step-counting patterns are less frequent. BP dominant step-behavior shares a

significant proportion of the BP step-counting pattern (0.3984 ± 0.1167). This step

behavior also contains a minor amount with the IN and E step-counting patterns (0.1482

± 0.1694 and 0.2045 ± 0.1356, respectively). AD+E step-behavior consists of a similar

amount with the AD and E step-counting patterns (0.4195 ± 0.0168 and 0.3815 ± 0.0771,

respectively), while other step-counting patterns are less. M dominant behavior is

characterized by most of the M step-counting pattern (0.6235 ± 0.1509) and also consists

of less frequency of the AD and IM step-counting patterns (0.1267 ± 0.1378 and 0.1105 ±

0.0301, respectively). Though IM and IN step-counting patterns share the least number of

days and do not show any dominant step-behavioral clusters, these step-counting patterns

show some prevalence

in all step-behavioral clusters, especially in BP dominant and M dominant step-

behavioral clusters.

Scale effect evaluation by simulations

243  We then investigated the scale effects of day and participant numbers on cluster

244 identification (Fig 3 and Figs S2-5). First, the number of days was randomly selected

245 from 3 to 18 days per participant, and our procedure was applied. When 3 days of data

246 per participant were used, only 4 step-counting patterns (Fig 3A and Fig S2A) and 3 step

247 behaviors (Fig 3B and Fig S3A) were identified. By increasing the number of days, the

248 number of step-counting patterns and step behaviors increased and reached a plateau over

249 9 days of data per participant (Fig 3A and 3B). The step-counting patterns identified

250 from 3-18 days of data were consistent with those from the full data set (Fig 1 and Fig

251 S2), although step behaviors were slightly different from those of the full data set (Fig 2

252 and Fig S3). When 15 days per participant (540 days from 36 participants) were selected,

253 step behaviors became consistent with those from the full data set were detected set (Fig

254 S3B).

255  We then examined the scale effect of participant number on the step behavior

256 clustering by our procedure. When 10 participants were randomly selected, 5 step-

257 counting patterns were identified (Fig 3C and Fig S4), while only 2 step behaviors were

258 detected (Fig 3D and Fig S5). When 20 participants, step-counting patterns increased to 6

259 but the number of step behaviors was four (Fig 3C and 3D). Analysis using 30 participant

260 data set (572 days) showed a similar result to the full data set (Fig 2 and Fig S5). These

261 results suggest that at least 15 days per participant (42 participants) or data set from 30

262 participants (572 days) are required to obtain robust results.

264 **Categorization by a traditional tertile procedure shows**

265 **heterogeneity in step-counting patterns.**

We identified six 24-hour step-counting patterns and five daily step behavior

clusters using unsupervised machine-learning approaches. Then, we compare the step

amounts from these step-counting patterns to those from traditional grouping approaches

such as the tertile procedure.

Table 5 summarized the composition of step-counting patterns in each tertile

group. AD is the most dominant step-counting pattern in the high and mid groups (52.7%

and 46.6%, respectively). However, the composition of BP, M, and E step-counting

patterns differed between the high and mid groups. On the other hand, the low tertile

group consists of a relatively similar proportion for each step-counting pattern. The

proportion of tertile groups in each step-counting pattern was consistent with the result

above (Table 6). In the AD pattern, the high tertile group is dominant (44.2%) and

followed by the mid (37.9%) and low groups (17.9%). In of BP pattern, mid tertile group

is the most dominant (50.7%) and followed by the high (26.0%) and low (23.3%) tertile

groups. The low tertile and high tertile groups were similar (39.5%) and (38.0%) of days

in the M pattern. In the E pattern, the low tertile group is the dominant (42.2%), followed

by mid (31.1%) and high (21.7%). IM and IN patterns consist of low tertile groups

(76.6% and 67.5%, respectively. Each group showed significantly different compositions

(Table S5 – S8).

Discussion

In this study, we attempted to develop a procedure to cluster 24-hour step-counting

patterns, as physical activity patterns, using unsupervised machine learning. We identified

six step-counting patterns and five daily step behavior clusters. Comparing traditional

tertile procedure (high, medium, and low) using daily step amounts, a different proportion

of six 24-hour step-counting patterns were observed in each tertile group, suggesting

heterogeneity in the categorization by the traditional procedure.

The "temporal patterns of physical activity" in the general term have been paid more

attention in recent decades (Aqeel et al., 2021; Creasy et al., 2021; De Baere, Lefevre, De

Martelaer, Philippaerts, & Seghers, 2015; Hallman, Mathiassen, Gupta, Korshøj, &

Holtermann, 2015) . However, the situation is confusing since this general term includes

different "activity" aspects, such as simple body movements or steps (Aqeel et al., 2021),

active/sedentary behaviors (Creasy et al., 2021), social behaviors (De Baere et al., 2015), or

those combinations (Hallman et al., 2015). Only several studies focused on the 24-hour

temporal physical activity patterns, such as body movements or steps in a narrower

criterion (Aqeel et al., 2021; Niemelä et al., 2019; Smagula et al., 2015, 2018, 2022). For

clustering, some previous studies extracted and used several features, such as rhythm

height, timing and robustness of physical activity (Smagula et al., 2015), several parameters

affecting phase shape and length of rest-activity rhythm (Smagula et al., 2018, 2022), or

timing, intensity, and duration of activity (Aqeel et al., 2021). Another study used 7-day-

continuous data of physical activity counts for clustering and then analyzed intensity and

temporal patterns of physical activity for 24 hours (Niemelä et al., 2019). These studies

adopted a two-step strategy, that is, 1) clustering by extracted features and 2) analyzing 24-

hour patterns of physical activity. By contrast, our procedure first directly identified the 24-

hour patterns of physical activity from bulk data of daily step-counting records using time-

series clustering analysis (Fig 1) and further clustered participants depending on the

probability of each pattern (Fig 2). Therefore, our strategy appears more straightforward,

direct, and intuitive than previous studies.

The scale effect analyses by simulation using different numbers of randomly selected

data sets indicated the limitation of our procedure when applying for a smaller number of

data (Fig 3 and Figs S2-5). The analyses indicated that at least 15 days per participant (540

days from 36 participants) or data set from 30 participants (572 days) are required to obtain

robust results, suggesting that over 500 days of data set could be essential. The relationship

between the number of days per participant and participants remains to be investigated

using a larger data set.

Conventional quantile procedure has been employed on the total amount of activity

to categorize data into some clusters because of its simple application. Recently, however,

quantile procedure has been pointed for a few potential problems; (1) reduction of detection

power, (2) multiple comparison testing, (3) assuming the homogeneity of risk within the

group, and (4) difficulty in comparing results within studies (Bennette & Vickers, 2012;

Greenland, 1995). Unfortunately, these problems remain to be unsolved to date as

discussed in a recent review (Jones & Ekelund, 2019). Our procedure described in this

study might provide a complementary strategy to compensate for the heterogeneity in the

quantile-based clustering. Our study showed an association between step-counting patterns

and step amounts and heterogeneity in step-counting patterns among tertile groups based

on the step amount, producing a typical example of the second problem (Tables 6 and 7).

Our approach may be helpful for understanding and handling the heterogeneity derived

from different physical activity patterns between/within the groups, considering it as a

confounding factor.

A significant limitation of the current study is that our dataset was small and

collected from office workers in the Miyagi area in the Tohoku region, Japan. As daily

activity depends on job type and the geographical area where participants live, it is still

challenging to mention how our procedure can be generalized for different populations. A

more extensive and diverse dataset needs to be analyzed to confirm our procedure's

reliability, robustness, and reproducibility. Our approach using unsupervised machine

learning to daily activity behavior and patterns may help to understand people's

prominent behaviors considering heterogeneity. Although the relationship between our

results and health outcome is beyond the goal of this study, our procedure might be

reliable for identifying specific patterns of physical activity and activity behavior

associated with positive health outcomes. It could help the development of targeted

interventions aimed at promoting healthy behavior.

347 Conclusion

348   In our study, we introduced a new approach using unsupervised machine learning

349   to group 24-hour step-counting patterns. This approach revealed six distinct patterns and

350   five clusters of daily step behaviors. We found significant heterogeneity in daily step

351   behavior categorization when applying a traditional tertile procedure based on total step

352   amount, emphasizing the importance of considering temporal variations in activity.

354 References

Amagasa, S., Machida, M., Fukushima, N., Kikuchi, H., Takamiya, T., Odagiri,

Y., & Inoue, S. (2018, July 9). Is objectively measured light-intensity

physical activity associated with health outcomes after adjustment for

moderate-to-vigorous physical activity in adults? A systematic review.

International Journal of Behavioral Nutrition and Physical Activity, Vol. 15.

Int J Behav Nutr Phys Act. https://doi.org/10.1186/s12966-018-0695-z

Aqeel, M., Guo, J., Lin, L., Gelfand, S., Delp, E., Bhadra, A., … Eicher-Miller, H.

A. (2021). Temporal Physical Activity Patterns are Associated with Obesity

in U.S. Adults. *Preventive Medicine*, *148*, 106538.

https://doi.org/10.1016/J.YPMED.2021.106538

Bennette, C., & Vickers, A. (2012). Against quantiles: Categorization of

continuous variables in epidemiologic research, and its discontents. *BMC*

Medical Research Methodology, *12*. https://doi.org/10.1186/1471-2288-12-21

Brailey, G., Metcalf, B., Lear, R., Price, L., Cumming, S., & Stiles, V. (2022, June

1). A comparison of the associations between bone health and three different

intensities of accelerometer-derived habitual physical activity in children and

adolescents: a systematic review. *Osteoporosis International*, Vol. 33, pp.

1191–1222. Osteoporos Int. https://doi.org/10.1007/s00198-021-06218-5

Chekroud, S. R., Gueorguieva, R., Zheutlin, A. B., Paulus, M., Krumholz, H. M.,

Krystal, J. H., & Chekroud, A. M. (2018). Association between physical

exercise and mental health in 1·2 million individuals in the USA between

2011 and 2015: a cross-sectional study. *The Lancet Psychiatry*, *5*(9), 739–

746. https://doi.org/10.1016/S2215-0366(18)30227-X

Creasy, S. A., Hibbing, P. R., Cotton, E., Lyden, K., Ostendorf, D. M., Willis, E.

A., … Catenacci, V. A. (2021). Temporal patterns of physical activity in

successful weight loss maintainers. *International Journal of Obesity (2005)*,

45(9), 2074–2082. https://doi.org/10.1038/S41366-021-00877-4

De Baere, S., Lefevre, J., De Martelaer, K., Philippaerts, R., & Seghers, J. (2015).

Temporal patterns of physical activity and sedentary behavior in 10-14 year-

old children on weekdays. *BMC Public Health*, *15*(1).

https://doi.org/10.1186/S12889-015-2093-7

Eyler, A. A., Brownson, R. C., Bacak, S. J., & Housemann, R. A. (2003). The

epidemiology of walking for physical activity in the United States. *Medicine*

and Science in Sports and Exercise, *35*(9), 1529–1536.

https://doi.org/10.1249/01.MSS.0000084622.39122.0C

Greenland, S. (1995). Avoiding power loss associated with categorization and

ordinal scores in dose-response and trend analysis. *Epidemiology*

(Cambridge, Mass.), *6*(4), 450. https://doi.org/10.1097/00001648-199507000-

00025

Guo, H., Niu, K., Monma, H., Kobayashi, Y., Guan, L., Sato, M., … Nagatomi, R.

(2012). Association of Japanese dietary pattern with serum adiponectin

concentration in Japanese adult men. *Nutrition, Metabolism, and*

Cardiovascular Diseases : NMCD, *22*(3), 277–284.

https://doi.org/10.1016/J.NUMECD.2010.06.006

Hallman, D. M., Mathiassen, S. E., Gupta, N., Korshøj, M., & Holtermann, A.

(2015). Differences between work and leisure in temporal patterns of

objectively measured physical activity among blue-collar workers. *BMC*

Public Health, *15*(1). https://doi.org/10.1186/S12889-015-2339-4

Hart, T. L., Swartz, A. M., Cashin, S. E., & Strath, S. J. (2011). How many days of

monitoring predict physical activity and sedentary behaviour in older adults?

International Journal of Behavioral Nutrition and Physical Activity, Vol. 8.

https://doi.org/10.1186/1479-5868-8-62

Hatano, Y. (1993) Use of the pedometer for promoting daily walking exercise.

Journal of the International Com- mittee on Health, Physical Education and

Recreation, 29, 4-8. - References - Scientific Research Publishing. (n.d.).

Retrieved June 1, 2023, from

https://www.scirp.org/(S(i43dyn45teexjx455qlt3d2q))/reference/ReferencesP

apers.aspx?ReferenceID=454558

Jones, P. R., & Ekelund, U. (2019). Physical Activity in the Prevention of Weight

Gain: the Impact of Measurement and Interpretation of Associations. *Current*

Obesity Reports, *8*(2), 66–76. https://doi.org/10.1007/S13679-019-00337-1

Kraus, W. E., Powell, K. E., Haskell, W. L., Janz, K. F., Campbell, W. W., Jakicic,

J. M., … Piercy, K. L. (2019). Physical Activity, All-Cause and

Cardiovascular Mortality, and Cardiovascular Disease. *Medicine and Science*

in Sports and Exercise, *51*(6), 1270–1281.

https://doi.org/10.1249/MSS.0000000000001939

Lu, Y., Wiltshire, H. D., Baker, J. S., Wang, Q., Ying, S., Li, J., & Lu, Y. (2022).

Objectively determined physical activity and adiposity measures in adult

women: A systematic review and meta-analysis. *Frontiers in Physiology*, *13*. https://doi.org/10.3389/FPHYS.2022.935892

Madigan, C. D., Fong, M., Howick, J., Kettle, V., Rouse, P., Hamilton, L., … Daley, A. J. (2021). Effectiveness of interventions to maintain physical activity behavior (device-measured): Systematic review and meta-analysis of randomized controlled trials. *Obesity Reviews*, *22*(10). https://doi.org/10.1111/obr.13304

Morris, J. N., & Hardman, A. E. (1997). Walking to health. *Sports Medicine*, Vol. 23, pp. 306–332. Sports Med. https://doi.org/10.2165/00007256-199723050-00004

National Heart, Lung, and Blood Institute. Tips for getting active. (n.d.). Retrieved from https://www.nhlbi.nih.gov/health/educational/wecan/get-active/getting-active.htm

Niemelä, M., Kangas, M., Farrahi, V., Kiviniemi, A., Leinonen, A.-M., Ahola, R., … Jämsä, T. (2019). *Intensity and temporal patterns of physical activity and cardiovascular disease risk in midlife*. https://doi.org/10.1016/j.ypmed.2019.04.023

Northern Finland Birth Cohorts | University of Oulu. (n.d.). Retrieved from https://www.oulu.fi/en/university/faculties-and-units/faculty-medicine/northern-finland-birth-cohorts-and-arctic-biobank

Ogilvie, D., Foster, C. E., Rothnie, H., Cavill, N., Hamilton, V., Fitzsimons, C. F., & Mutrie, N. (2007). Interventions to promote walking: systematic review.

BMJ (Clinical Research Ed.), *334*(7605), 1204–1207.

https://doi.org/10.1136/BMJ.39198.722720.BE

Organization WHO. Physical activity 2022. (n.d.). Retrieved from

https://www.who.int/news-room/fact-sheets/detail/physical-activity

Organization WHO Physical inactivity. (n.d.). Retrieved from

https://www.who.int/data/gho/indicator-metadata-registry/imr-details/3416

Paluch, A. E., Bajpai, S., Bassett, D. R., Carnethon, M. R., Ekelund, U., Evenson,

K. R., … Fulton, J. E. (2022). Daily steps and all-cause mortality: a meta-

analysis of 15 international cohorts. *The Lancet. Public Health*, *7*(3), e219–

e228. https://doi.org/10.1016/S2468-2667(21)00302-9

Pearce, M., Garcia, L., Abbas, A., Strain, T., Schuch, F. B., Golubic, R., …

Woodcock, J. (2022). Association Between Physical Activity and Risk of

Depression: A Systematic Review and Meta-analysis. *JAMA Psychiatry*,

79(6), 550–559. https://doi.org/10.1001/JAMAPSYCHIATRY.2022.0609

Piercy, K. L., Troiano, R. P., Ballard, R. M., Carlson, S. A., Fulton, J. E., Galuska,

D. A., … Olson, R. D. (2018). The Physical Activity Guidelines for

Americans. *JAMA*, *320*(19), 2020–2028.

https://doi.org/10.1001/JAMA.2018.14854

Powell, K. E., Paluch, A. E., & Blair, S. N. (2011). Physical activity for health:

What kind? How much? How intense? On top of what? *Annual Review of*

Public Health, *32*, 349–365. https://doi.org/10.1146/ANNUREV-

PUBLHEALTH-031210-101151

Smagula, S. F., Boudreau, R. M., Stone, K., Reynolds, C. F., Bromberger, J. T.,

Ancoli-Israel, S., … Cauley, J. A. (2015). Latent activity rhythm disturbance

sub-groups and longitudinal change in depression symptoms among older

men. *Chronobiology International*, *32*(10), 1427–1437.

https://doi.org/10.3109/07420528.2015.1102925

Smagula, S. F., Krafty, R. T., Thayer, J. F., Buysse, D. J., & Hall, M. H. (2018).

Rest-activity rhythm profiles associated with manic-hypomanic and

depressive symptoms. *Journal of Psychiatric Research*, *102*, 238–244.

https://doi.org/10.1016/J.JPSYCHIRES.2018.04.015

Smagula, S. F., Zhang, G., Gujral, S., Covassin, N., Li, J., Taylor, W. D., …

Krafty, R. T. (2022). Association of 24-Hour Activity Pattern Phenotypes

With Depression Symptoms and Cognitive Performance in Aging. *JAMA*

Psychiatry, *79*(10), 1023–1031.

https://doi.org/10.1001/JAMAPSYCHIATRY.2022.2573

Tavenard, R., Faouzi, J., Vandewiele, G., Divo, F., Androz, G., Holtz, C., …

Rußwurm, M. (2020). Tslearn, A Machine Learning Toolkit for Time Series

Data. *Journal of Machine Learning Research*, *21*, 1–6. Retrieved from

https://github.com/tslearn-team/tslearn.

Tudor-Locke, C., & Bassett, D. R. (2004). How Many Steps/Day Are Enough?

Preliminary Pedometer Indices for Public Health. *Sports Medicine*, Vol. 34,

pp. 1–8. Sports Med. https://doi.org/10.2165/00007256-200434010-00001

Author contribution

SSN and HI for conceptualization, methodology, data analysis, interpretation, and original draft writing; HM for writing, review, and editing; RN for supervision. All authors read and approved the final manuscript.

Data availability

Data is available on request from the authors.

Declaration of interests

The authors have no competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

We would like to thank all the participants and staff who have contributed to the experiment. A pioneering research support grant from Tohoku University partially supported this project. The funding body has no role in the study design, data analysis, or manuscript submission.

Figure legends

Fig 1. Six step-counting patterns for 24 hours. Temporal step-counting pattern for 24 hours identified using unsupervised machine learning. (A) All traces of step-counting activity in each pattern. (B) Averaged traces (mean ± SD) of step-counting activity in each pattern. AD, all-day; BP, bi-phasic; M, morning; E, evening; IM, irregular morning; IN, irregular night.

Fig 2. Five step behaviors. Step behavior clusters were identified using unsupervised machine learning. (A) All traces of proportion of step-counting patterns in each behavior. (B) Averaged traces (mean) of the proportion of step-counting patterns in each behavior. AD, all-day; BP, bi-phasic; M, morning; E, evening; IM, irregular morning; IN, irregular night.

Fig 3. Scale effect evaluation by simulations. The effects of day and participant numbers on cluster identification were investigated. The effects of different numbers of days on identifying the numbers of step-counting patterns (A) and step behaviors (B). The effects of different numbers of participants on identifying the numbers of step-counting patterns (C) and step behaviors (D).

Fig 1. Six step-counting patterns for 24 hours. Temporal step-counting pattern for 24 hours identified using unsupervised machine learning. (A) All traces of step-counting activity in each pattern. (B) Averaged traces (mean ± SD) of step-counting activity in each pattern. AD, all-day; BP, bi-phasic; M, morning; E, evening; IM, irregular morning; IN, irregular night.

Fig 2. Five step behaviors. Step behavior clusters were identified using unsupervised machine learning. (A) All traces of proportion of step-counting patterns in each behavior. (B) Averaged traces (mean) of the proportion of step-counting patterns in each behavior. AD, all-day; BP, bi-phasic; M, morning; E, evening; IM, irregular morning; IN, irregular night.

Fig 3. Scale effect evaluation by simulations. The effects of day and participant numbers on cluster identification were investigated. The effects of different numbers of days on identifying the numbers of step-counting patterns (A) and step behaviors (B). The effects of different numbers of participants on identifying the numbers of step-counting patterns (C) and step behaviors (D).

Figure 3

Table 1. Demographical information

Variables	Mean ± SD
Age (Years)	45.59 ± 10.96
Weight (kg)	55.83 ± 10.69
Body fat (%)	29.37 ± 6.29
BMI	22.10 ± 3.42
Waist (cm)	78.04 ± 9.72

Note: Demographic information for 42 participants

Table 2. Proportion of each step-counting pattern

Pattern	Days	% of days
All-Day (AD)	330	40.5
Bi-phasic (BP)	146	17.9
Morning (M)	129	15.8
Evening (E)	106	13.0
Irregular morning (IM)	64	7.9
Irregular night (IN)	40	4.9
Total	815	100

Note: In total 815 days, the AD pattern was the most common, followed by BP, M, and E. IM and IN patterns were less frequent.

Table 3. Step count amount and duration for each pattern

Pattern	Step count amount mean ± SD	Activity duration (hours) mean ± SD
All-Day (AD)	7020.0 ± 3271.4	14.29 ± 2.28
Bi-phasic (BP)	6178.2 ± 3102.1	14.54 ± 2.61
Morning (M)	6473.7 ± 4220.1	12.49 ± 3.97
Evening (E)	4871.0 ± 3091.9	10.91 ± 4.30
Irregular morning (IM)	2597.6 ± 3481.4	5.67 ± 5.39
Irregular night (IN)	3977.2 ± 5540.0	7.82 ± 6.48

Note: Each step-counting pattern had different daily step counts. AD had the highest, followed by M and BP, then E. IN and IM had the lowest counts. In terms of duration, the BP pattern exhibited the longest activity duration per day, followed by AD, M and E patterns. IM and IN has the lowest duration of activity.

Table 4. Probability of 24-hour step-counting patterns in each daily step-behavior

Behavior	Mean (SD)					
	AD	BP	M	E	IM	IN
AD dominant	0.5515 (0.1492)	0.0887 (0.0606)	0.1227 (0.0888)	0.0998 (0.0702)	0.0886 (0.1190)	0.0486 (0.0610)
AD+BP	0.4031 (0.1021)	0.2964 (0.0586)	0.1587 (0.0748)	0.0850 (0.0800)	0.0398 (0.0362)	0.0117 (0.0237)
BP dominant	0.0740 (0.0468)	0.3984 (0.1167)	0.0833 (0.0701)	0.2045 (0.1356)	0.0915 (0.0752)	0.1482 (0.1694)
AD+E	0.4195 (0.0168)	0.0916 (0.0422)	0.0770 (0.0938)	0.3815 (0.0771)	0.0156 (0.0313)	0.0147 (0.0294)
M dominant	0.1267 (0.1378)	0.0903 (0.0604)	0.6235 (0.1509)	0.0175 (0.0304)	0.1105 (0.0301)	0.0314 (0.0278)

Note: This table displays probabilities of six step-counting patterns in daily step-behavioral clusters. AD dominant behavior is mostly AD pattern, with some M pattern. AD+BP is mainly AD pattern and BP pattern, with some M pattern. BP dominant behavior is primarily BP pattern, with minor IN pattern and E pattern. AD+E behavior is evenly split between AD pattern and E pattern. M dominant behavior is mostly M pattern, with some AD pattern and IM pattern. IM and IN are less common but present in all clusters, especially in M dominant and BP dominant behavior respectively.

Table 5. Proportion and number of days of 24-hour step-counting pattern in each tertile group

Activity	AD	BP	M	E	IM	IN	Total
High	52.7% 146	13.7% 38	17.7% 49	8.3% 23	4.0% 11	3.6% 10	277
Mid	46.6% 125	27.6% 74	10.8% 29	12.3% 33	1.5% 4	1.1% 3	268
Low	21.9% 59	12.6% 34	18.9% 51	18.5% 50	18.1% 49	10.0% 27	270

Note: This table outlines step-counting pattern compositions across tertile groups. AD predominates in high and mid groups, while BP, M, and E patterns vary between high and mid groups. Conversely, the low tertile group shows a more balanced distribution across patterns.

Table 6. Proportion and number of days of tertile groups for each 24-hour step-counting pattern

Pattern	High	Mid	Low	Total
AD	44.2 %	37.9 %	17.9 %	
	146	125	59	330
BP	26.0 %	50.7 %	23.3 %	
	38	74	34	146
M	38.0 %	22.5 %	39.5 %	
	49	29	51	129
E	21.7 %	31.1 %	42.2 %	
	23	33	50	106
IM	17.2 %	6.3 %	76.6 %	
	11	4	49	64
IN	25.0 %	7.5 %	67.5 %	
	10	3	27	40

Note: Tertile group proportions align with previous findings as in Table 5. In the AD pattern, high tertile dominates, followed by mid and low. BP is mostly in the mid tertile, followed by high and low. M pattern is similar in low and high tertile. E pattern is dominated by the low tertile, followed by mid and high. IM and IN patterns are mainly in the low tertile groups.