Running title: *Solanum* trichome metabolite diversity

Trading acyls and swapping sugars: metabolic innovations in *Solanum* trichomes

Paul D. Fiesel[1], Rachel E. Kerwin[1], A. Daniel Jones[1], and Robert L. Last[1,2]

[1]Department of Biochemistry and Molecular Biology, Michigan State University, East Lansing,

MI 48823 USA

[2]Department of Plant Biology, Michigan State University, East Lansing, MI 48823 USA

P.D.F and R.E.K contributed equally

Corresponding Author: Robert L. Last, Michigan State University, Department of Biochemistry

and Molecular Biology, 603 Wilson Road, East Lansing, MI 48824; Phone: 517-432-3278;

email: lastr@msu.edu

ORCID:

Paul D. Fiesel 0000-0002-2038-5417

Rachel E. Kerwin 0000-0001-7395-5973

A. Daniel Jones 0000-0002-7408-6690

Robert L. Last 0000-0001-6974-9587

Abstract

Solanaceae (nightshade family) species synthesize a remarkable array of clade- and tissue-specific specialized metabolites. Protective acylsugars, one such class of structurally diverse metabolites, are produced by acylsugar acyltransferases from sugars and acyl-Coenzyme A esters. We characterized trichome acylsugars of the Clade II species *Solanum melongena* (brinjal eggplant) using liquid chromatography-mass spectrometry (LC-MS), gas chromatography-MS and nuclear magnetic resonance (NMR) spectroscopy. This led to the identification of eight unusual structures with inositol or inositol glycoside cores, and hydroxyacyl chains. LC-MS analysis of 31 species in the megadiverse *Solanum* genus revealed striking acylsugar diversity with some traits restricted to specific clades and species. Acylinositols were found throughout each clade while acylglucoses were restricted to DulMo and VANAns species. Medium length hydroxyacyl chains were identified in many species. Analysis of tissue-specific transcriptomes and interspecific acylsugar acetylation differences led to the unexpected identification of the *S. melongena* AcylSugar AcylTransferase 3-Like 1 (SmASAT3-L1; SMEL4.1_12g015780) enzyme. This enzyme is distinct from previously characterized acylsugar acetyltransferases, which are in the ASAT4 clade, and appears to be a functionally divergent ASAT3. This study provides a foundation for investigating the evolution and function of diverse *Solanum* acylsugar structures and harnessing this diversity in breeding and synthetic biology.

Introduction

Plants are remarkable synthetic chemists, producing a multitude of structurally complex specialized metabolites that differ from the products of general, or primary, metabolism in their lineage-specific distribution and tissue or cell type-specific biosynthesis. In contrast to the negative selection against changes in primary metabolism (for example, amino acids, energy metabolism intermediates and vitamin cofactors), less constrained evolution of specialized metabolism led to accumulation of hundreds of thousands of taxonomically restricted metabolites in broad classes. Specialized metabolites play many critical roles such as in abiotic and biotic stress adaptation (Agati and Tattini, 2010; De Moraes et al., 2001; Landry et al., 1995), pollinator attraction (Kretschmar and Baumann, 1999) and mediation of interactions with beneficial and pathogenic microbes (Yu et al., 2021). These diverse and bioactive small molecules have historical and modern uses in human medicine, including the chemotherapeutics vinblastine and paclitaxel, antimalarial artemisinin, and painkillers such as morphine. Their distinct phylogenetic distributions and relatively rapid diversification make specialized metabolic enzymes excellent models to study evolution.

Acylsugars are specialized metabolites produced across the Solanaceae (nightshade) family, aiding in defense against herbivores, fungi, and bacteria (Goffreda et al., 1989; Leckie et al., 2016; Luu et al., 2017; Weinhold and Baldwin, 2011). In Types I- and IV-glandular trichomes, BAHD-type AcylSugar AcylTransferase (ASAT) enzymes assemble acylsugars from the basic building blocks of sugars, often sucrose, and short- to medium-length acyl chains derived from acyl-CoenzymeAs (acyl-CoAs) (Fan et al., 2015; Lou et al., 2021; Moghe et al., 2017; Schilmiller et al., 2015; Schilmiller et al., 2012). Despite their simple components, acylsugars exhibit remarkable chemical diversity arising from variations in sugar core

composition and acyl chain length, branching pattern, position, and number (Fan et al., 2019;

Fiesel et al., 2022; Ghosh et al., 2014; Hurney, 2018; Lou et al., 2021; Moghe et al., 2017;

Schenck et al., 2022). For example, acylsucroses, composed of a sucrose disaccharide core,

accumulate in cultivated tomato (*Solanum lycopersicum*) trichomes, while acylsucroses and

acylglucoses have been observed in the trichomes of wild tomato species, *Solanum pennellii*.

Acylsugar structural variation also impacts biological activity; for example, differential

oviposition deterrence was demonstrated from naturally derived acylsugar mixtures (Leckie et

al., 2016).

Solanaceae acylsugars have become an exemplary model to study evolution of a diverse,

biologically relevant trait in a plant family with genomic and phylogenetic resources. Utilization

of this model revealed gene duplication, neofunctionalization, co-option, and loss involved in

acylsugar evolution (Fan et al., 2020, 2017; Leong et al., 2019; Moghe et al., 2017). For

example, neofunctionalization of an invertase-like enzyme, AcylSucrose FructoFuranosidase 1

(ASFF1) and functional divergence of core ASAT enzymes is responsible for differences in

sugar core type and acyl chain positions between cultivated tomato *S. lycopersicum* and wild

tomato *S. pennellii* acylsugars (Fan et al., 2017; Leong et al., 2019). Identifying mechanisms of

acylsugar evolution across the broader *Solanum* requires a detailed understanding of acylsugar

diversity and biosynthesis, which is lacking for many species.

Nearly half of the Solanaceae falls into the large (>1200 species) and phenotypically

diverse *Solanum* genus. This genus is split into several major clades, including Potato,

Regmandra, DulMo, VANAns, and Clade II; the Potato clade contains cultivated tomato and

potato and their wild relatives while Clade II contains the cultivated brinjal eggplant, *Solanum*

melongena, and other 'spiny *Solanums*' (Bohs, 2004; Bohs and Olmstead, 1997; Gagnon et al.,

2022; Levin et al., 2006; PBI Solanum Project, 2022; Särkinen et al., 2013; Stern et al., 2011;

Tepe et al., 2016; Weese and Bohs, 2007). To date, documentation of Solanum acylsugar

diversity largely focused on a handful of species within the Potato clade, including cultivated

tomato and its close relatives. These efforts identified at least 38 nuclear magnetic resonance

(NMR) spectroscopy-resolved acylsugar structures and liquid chromatography mass

spectrometry (LC-MS) supported annotations of many more (Fan et al., 2017; Ghosh et al., 2014;

Lybrand et al., 2020; Schilmiller et al., 2016). While limited acylsugar screening outside of the

Potato clade was reported, novel structural variants not observed among cultivated tomato

relatives were identified. For example, acylsugars with *myo*-inositol sugar cores (i.e.,

acylinositols) were characterized in three species: *S. nigrum*, from the DulMo clade*,* and *S.*

lanceolatum and *S. quitoense,* from Clade II (Herrera-Salgado et al., 2005; Hurney, 2018; Leong

et al., 2020; Lou et al., 2021). These discoveries highlight the benefits of a comprehensive

description of *Solanum* acylsugars within the well-developed phylogenetic framework.

Here we report analysis of *Solanum* acylsugar chemical diversity in species from the

relatively unexplored *Solanum* clades DulMo, VANAns, and Clade II, which together comprise

~1000 *Solanum* species. We first established the Clade II brinjal eggplant, *S. melongena*, as a

reference species. Eggplant is a major worldwide fruit crop with extensive genomic,

transcriptomic, and germplasm resources (Barchi et al., 2021; Li et al., 2021; Mennella et al.,

2010). We characterized eggplant trichome acylinositols, acylinositol glycosides, and acylsugars

with unusual hydroxylated acyl chains using electrospray ionization LC-quadrupole time of

flight-MS (ESI LC-QToF-MS) and NMR. These atypical structures likely reflect participation of

enzymes that are distinct from the cultivated tomato acylsucrose pathway. Moving beyond this

model organism framework, LC-MS phylogenetic screening of 31 Clade II, DulMo, and

VANAns species, including *S. melongena*, led to the identification of remarkable acylsugar

structural variation. LC-MS features with characteristics of acylinositols were found in 25 of the

26 acylsugar-producing species, suggesting one or a small number of evolutionary origins. In

contrast, acylglucoses were detected in DulMo and VANAns species, but not in any tested Clade

II species.

As a first step towards revealing the molecular basis of the extensive documented

acylsugar diversity, we utilized interspecific acylsugar differences and an eggplant tissue-

specific transcriptome to identify an acylinositol biosynthetic enzyme, *S. melongena* Acylsugar

AcylTransferase 3-Like 1 (SmASAT3-L1; SMEL4.1_12g015780), responsible for acetylating a

triacylinositol glycoside acyl acceptor. SmASAT3-L1 exhibits a different acyl-CoA specificity

than previously characterized ASAT3 homologs, highlighting how gene duplication and

functional divergence created acylsugar metabolic novelty in this part of the *Solanum* clade.

Results and Discussion

Eggplant glandular trichomes accumulate acylsugar-like metabolites

We began our investigation of *Solanum* acylsugar diversity with the brinjal eggplant *S.

melongena* due to its economic importance, genomic resources, and phylogenetic position within

the monophyletic Eastern Hemisphere Spiny clade of Clade II (formerly known as the 'Old

World spiny clade') (Gagnon et al., 2022). Using eight eggplant accessions (listed in Table S1),

we observed glandular trichomes on hypocotyls, cotyledons, and the first three true leaves of

young eggplant plants (Figure 1, S1). These resemble the acylsugar-producing structures found

in other *Solanum* species (Leong et al., 2020; Lou et al., 2021; Schilmiller et al., 2012). In

contrast, we observed only non-glandular stellate trichomes on leaves and stems of mature

eggplants (Figure S1), and these are types that typically do not accumulate acylsugars or other

specialized metabolites (Levin, 1973; Wagner, 1991).

We analyzed surface metabolite extracts from young and mature eggplant tissues using

LC-QToF-MS coupled with collision-induced dissociation (CID) in negative and positive ion

mode. We used these data to annotate acylsugars based on relative mass defect, molecular adduct

ion masses, retention times, and ions present in CID mass spectra. Briefly, acylsugars were

annotated from masses of fatty acid carboxylate fragment ions, as well as fragment ions

corresponding to stepwise losses of acyl chains from the pseudomolecular ion to a sugar core

fragment ion. The acylsugar annotation methods and confidence criteria are explained in detail in

the Methods. This analysis revealed abundant acylsugars in extracts from young, glandular

trichome-producing eggplant tissues, but not from mature, non-glandular trichome producing

tissues.

We annotated 38 acylsugars in leaf surface extracts of young eggplant from eight of nine

accessions (Table S1), consisting of 16 acylhexoses and 22 acyldisaccharides (Table 1). LC-MS

based acylsugar annotations are described using a shorthand nomenclature modified from

previous work (Leong et al., 2020) as follows: UX:Y:Z(A,B,C,D), where U – as a single or

multi-letter designation – represents the sugar core, X represents the number of acyl chains, Y is

the sum of acyl chain carbons, Z indicates the number of unsaturated bonds in the acyl chains

(when present), and A-D represent the number of carbon atoms in the individual acyl chains. For

example, the eggplant acylhexose I3:18(4,4,10) consists of a *myo*-inositol core with three acyl

chains with a total of 18 carbon atoms.

Curiously, eggplant acyldisaccharides contain an atypical pentose-hexose core, as

evidenced by a fragment ion mass of *m/z* 293.09 in negative-ion mode, corresponding to a fully

deacylated sugar core minus a hydroxyl group and a proton. Further evidence was provided by

positive mode CID, which promotes glycosidic bond cleavage, yielding a fragment ion

corresponding to the neutral loss of an unacylated pentose ring. All acylhexoses annotated with

medium to high confidence formed abundant fragment ions in positive mode but few fragment

ions in negative mode as illustrated in Figure S2; this pattern is characteristic of acylinositols

found in *S. quitoense* and *S. nigrum,* but has not been observed for acylglucoses, suggesting that

the detected *S. melongena* acylhexoses are acylinositols (Hurney, 2018; Leong et al., 2020; Lou

et al., 2021).

The eggplant acylsugars each contained three to four acylations, all on the hexose core,

including one medium eight-carbon (C8) to C14 acyl chain and two to three C4 or C5 acyl

chains. The medium acyl chains (C8, C10, C12, and C14) were either iso-branched or straight as

revealed by GC-MS acyl chain analysis (Figure S3). Additionally, we identified hydroxylated

C12, C14, and C16 acyl chains not previously reported in Solanaceae acylsugars (Table 1).

Although we did not observe large differences between the eight eggplant accessions, eggplant

acylsugars differ in chain length, functional groups, and acyl chain composition from the

reported *Solanum* acylinositols (Herrera-Salgado et al., 2005; Hurney, 2018; Lou et al., 2021).

Table 1. Summary of annotated acylsugars detected in *S. melongena* leaf surface extracts.
PH = pentose-hexose; AI = arabinose-inositol, I = inositol. RT = retention time; m/z_{acc} =
theoretical monoisotopic formate adduct mass; m/z_{ex} = experimental formate adduct mass; Δm
(ppm) = mass measurement error in parts per million. Compound numbers are listed for NMR
characterized compounds (Figure 1). Percent total peak area calculated by dividing acylsugar
peak area by total acylsugar peak area. Acylsugars composing ≥1% of total acylsugar peak area
are in bold. Acylsugar abundance was reported by Progenesis QI and averaged over 15 samples.
The annotation method is described in the Methods. Acylsugars are sorted by number of sugar
moieties and then by elution order.
184

Name	Compound #	Annotation confidence level	RT (min)	Neutral molecule chemical formula	m/z_{acc}	m/z_{ex}	Δm (ppm)	Average acylsugar abundance (%)
Acyldisaccharides								
PH2:12(4,8)		medium	1.98	C23H40O12	553.2496	553.2496	0.0	0.203
PH4:14(2,4,4,4)		medium	2.19	C25H40O14	609.2395	609.2445	8.2	0.0246
PH3:14(2,4,8)		medium	2.23	C25H42O13	595.2602	595.2621	3.2	0.0453
PH4:16(2,2,4,8)		medium	2.54	C27H44O14	637.2708	637.2720	1.9	0.0851
PH3:16(4,4,8)		**medium**	**2.61**	**C27H46O13**	**623.2915**	**623.2916**	**0.2**	**17.1**
AI3:16(4,4,8)	**1**	**high**	**2.66**	**C27H46O13**				
PH4:17(2,2,4,9)		medium	2.93	C28H46O14	651.2864	651.2924	9.2	0.114
PH3:17(4,5,8)		medium	2.96	C28H48O13	637.3071	637.3083	1.9	0.324
PH4:18(2,4,4,8)		medium	3.11	C29H48O14	665.3021	665.3033	1.8	0.901
AI4:18(2,4,4,8)	**2**	**high**	**3.27**	**C29H48O14**	**665.3021**	**665.3026**	**0.7**	**42.5**
PH3:18(4,4,10)		**medium**	**3.61**	**C29H50O13**	**651.3228**	**651.3205**	**-3.5**	**1.12**
PH3:18(4,4,10)		medium	3.74	C29H50O13	651.3228	651.3245	2.5	0.964
PH4:19(2,4,5,8)		medium	3.76	C30H50O14	679.3177	679.3146	-4.6	0.329
PH4:19(2,4,5,8)		medium	3.83	C30H50O14				
PH4:20(4,4,4,8)		medium	4.22	C31H52O14	693.3334	693.3350	2.3	0.0428
PH4:20(2,4,4,10)		medium	4.57	C31H52O14	693.3334	693.3351	2.5	0.0447
PH4:20(2,4,4,10)		medium	4.68	C31H52O14	693.3334	693.3349	2.1	0.127
PH4:20(2,4,4,10)		medium	4.85	C31H52O14	693.3334	693.3379	6.6	0.00422
PH3:20(4,4,12)		medium	5.05	C31H54O13	679.3541	679.3554	1.9	0.0389
PH3:20(4,4,12)		medium	5.23	C31H54O13	679.3541	679.3552	1.6	0.130
PH3:22(4,4,14)		medium	7.54	C33H58O13	707.3854	707.3879	3.5	0.0356
PH3:22(4,4,14)		medium	7.85	C33H58O13	707.3854	707.3892	5.3	0.0107
Acylhexoses								
I3:16(4,4,8)		**medium**	**3.27**	**C22H38O9**	**491.2492**	**491.2492**	**-0.1**	**3.99**
I3:20(4,4,12-OH)		medium	4.16	C26H46O10	563.3068	563.3071	0.5	0.0808
I3:20(4,4,12-OH)		medium	4.32	C26H46O10	563.3068	563.3068	0.0	0.500
I4:18(2,4,4,8)		medium	4.22	C24H40O10	533.2598	533.2604	1.1	0.0248
I3:18(4,4,10)	**3**	**high**	**4.70**	**C24H42O9**	**519.2805**	**519.2802**	**-0.5**	**10.6**
I3:18(4,4,10)		medium	4.88	C24H42O9	519.2805	519.2794	-2.1	0.693
I3:20(2,4,14-OH)		medium	4.68	C26H46O10	563.3068	563.3082	2.5	0.0588
I3:20(2,4,14-OH)		medium	4.90	C26H46O10	563.3068	563.3077	1.6	0.298
I3:22(4,4,14-OH)	**7**	**high**	**6.06**	**C28H50O10**	**591.3381**	**591.3385**	**0.7**	**4.96**
I3:22(4,4,14-OH)	**8**	**high**	**6.34**	**C28H50O10**	**591.3381**	**591.3380**	**-0.1**	**10.6**
I3:20(4,4,12)	**4**	**high**	**6.81**	**C26H46O9**	**547.3118**	**547.3119**	**0.1**	**2.24**
I3:20(4,4,12)		medium	7.09	C26H46O9	547.3118	547.3127	1.7	0.105
I3:24(4,4,16-OH)		medium	8.51	C30H54O10	619.3694	619.3707	2.0	0.0473
I3:24(4,4,16-OH)		medium	8.85	C30H54O10	619.3694	619.3683	-1.8	0.0355
I3:22(4,4,14)	**5**	**high**	**9.34**	**C28H50O9**	**575.3431**	**575.3435**	**0.7**	**0.986**

I3:22(4,4,14)	6	high	9.68	C28H50O9	575.3431	575.3433	0.3	0.722

Although LC-MS provided information about sugar core mass, it did not reveal the sugar

core structure, prompting analysis using gas chromatography-mass spectrometry (GC-MS) of

derivatized *S. melongena* acylsugar cores. When free sugar cores, produced by metabolite extract

saponification, were derivatized to form alditol acetates, GC-MS peaks corresponding to

derivatized *myo*-inositol and glucose were detected, supporting the presence of acylinositols

(Figure S4A). Because the detection of glucose might have resulted from other compounds in the

leaf surface metabolite extracts, the disaccharide sugar core composition was determined by

hydrolyzing the saponified sugar cores with formic acid to cleave the glycosidic linkage.

Comparison of the hydrolyzed and unhydrolyzed samples revealed a peak corresponding to

arabinose only in the hydrolyzed plant samples (Figure S4B). Taken together, the results of

saponification with and without hydrolysis, followed by derivatization, confirmed identification

of *myo*-inositol sugar cores and identified the pentose moiety of the hexose-pentose disaccharide

core as arabinose.

NMR analysis of eight eggplant acylsugars

While MS analysis provided valuable information about acyl chain number and length,

sugar core mass, as well as acyl chain number, complementary information about acyl chain

branch structure, sugar core stereochemistry, and acyl chain positions was obtained using NMR.

We purified and resolved the structures of eight abundant eggplant acylsugars from accession PI

555598 using a combination of 1D and 2D NMR experiments (Figure 1). All eight structures are

newly described, and because atomic connections were determined by NMR and MS data, the

proposed structures meet Metabolomics Standards Initiative level 1 criteria for metabolite

identification (Sumner et al., 2007).

A series of NMR experiments confirmed that the acylhexose and acyldisaccharide sugar

cores are *myo*-inositol and 4-*O*-β-arabinopyranosyl *myo*-inositol, respectively (Figure 1C,D;

Supplemental Data File S1). We assigned all sugar ring proton signals of each sugar core's spin

system with total correlation spectroscopy (TOCSY). Inferences from correlation spectroscopy

(COSY) data then identified the order of ring protons and revealed pyranose and cyclitol ring

structures for the pentose and hexose rings, respectively. Relative stereochemistry at ring

positions was subsequently determined through comparison of spin-spin splitting (multiplicities

and coupling constants) referenced to expected patterns inferred from chemical principles and

previously reported acylinositols (Hurney, 2018; Leong et al., 2020). The disaccharide glycosidic

linkage position was determined by heteronuclear multiple bond correlation (HMBC)

correlations to be at position 4 and position 1 of *myo*-inositol and arabinose, respectively. The

arabinose *β* anomeric configuration was inferred from the anomeric carbon $^1J_{CH}$ (162 Hz) as

revealed by a coupled-heteronuclear single quantum coherence (coupled-HSQC) experiment of

the free disaccharide sugar core after saponification. Our sugar core assignments identifying this

unusual disaccharide agreed with the previous GC-MS sugar core results, supporting the efficacy

of the sugar core GC-MS identification method (Figure S4) This disaccharide core differs in the

pentose moiety identity from a recently reported acylated 4-*O*-*ß*-xylopyranosyl *myo*-inositol in

the *Solanum* Clade II species *S. quitoense* (Hurney, 2018). To our knowledge, this is the first

report of acylated 4-*O*-*ß*-arabinosyl *myo*-inositol sugars.

Acyl chain positions, branching patterns, and hydroxyl positions were also resolved

through integration of different NMR experiments. We found that all acyl chains were confined

to *myo*-inositol, consistent with the LC-MS results. All eight acylsugars are decorated with two

short iso-branched iC4 acyl chain esters at positions 1 and 2, and one medium C8 to C14 acyl

chain ester at position 3 (Figure 1). Compound 2 (Figure 1C), the only tetraacylated acylsugar

identified in eggplant, additionally carried an acetylation at position 6 (Figure 1C). The medium-

length acyl chains at position 3 were resolved as straight (nC14) or terminally iso-branched (iC8,

iC10, iC12, iC14) based on signals characteristic of protons near the branched carbons.

Strikingly, we identified unusual hydroxylated straight and iso-branched 3-

hydroxytetradecanoate acyl chains, 3-OH-nC14 and 3-OH-iC14, in compounds 7 and 8,

respectively (Figure 1D). We assigned hydroxylation positions of 3-OH-nC14 and 3-OH-iC14 to

the third acyl carbon based on a downfield shifted signal in the ^{1}H NMR spectrum at 3.92 ppm,

corresponding to one hydrogen at that position. In contrast, the non-hydroxylated medium acyl

chains observed in compounds 1-6 (Figure 1C,D), carry two hydrogen atoms at the third acyl

carbon, and these have a characteristic signal near 1.50 ppm. We are not aware of previous

reports of 3-OH-nC14 and 3-OH-iC14 chains in Solanaceae acylsugars. While 3-OH-nC14

chains are observed in the acylsugar-like bacterial Lipid A glycolipids, the hydroxylation

position differentiates these eggplant chains from hydroxyacyl chains in castor bean (*Ricinus*

communis) seed oil, *Silene gallica* gallicasides, *Ibicella lutea* fatty acid glycosides, and

Convolvulaceae resin glycosides (Asai et al., 2010; Asai and Fujimoto, 2010; Bah and Pereda-

Miranda, 1996; Smith, 1971).

Figure 1. Profiling of *S. melongena* acylsugars. (A) LC-MS base peak intensity (BPI)
chromatogram of *S. melongena* acylsugars. Peaks characterized by NMR are indicated with their
compound numbers from Panels C and D. Metabolites were collected from seedling tissue with
glandular trichomes similar to the ones displayed in the closeup photo of a *S. melongena*
hypocotyl. (B) Annotation of HMBC experiments used to identify acylation positions of specific
acyl chains. At the top, I3:18(4,4,10) is displayed in the chair conformation along with
hydrogens important for determining acylation position. An HMBC NMR spectrum showing
couplings between hydrogens and carbonyl carbons is shown. (C) NMR-characterized

258 acyldisaccharide structures 1 and 2, with inositol ring carbons numbered. (D) NMR-
259 characterized acylinositol monosaccharide structures 3-8.
260

<u>The OH-tetradecanoate acyl chain has R stereochemistry</u>

To gain insight into the biosynthetic origin of the hydroxylated acyl chains, we tested

chirality of the 3-OH-nC14 acyl chain of compound 7 as this chain has available commercial

standards. Two-step chiral derivatization and GC-MS analysis was performed on compound 7

and two commercial standards, (3R)-OH-nC14 and the corresponding racemate (3R/S)-OH-

nC14. Fatty acids were converted to ethyl esters to aid volatilization, followed by esterification

of the acyl chain hydroxyl group with the Mosher acid (R)-(−)-α-methoxy-α-

(trifluoromethyl)phenylacetate (R-MPTA) to distinguish stereoisomers by GC separation of the

diastereomeric derivatives. GC-MS analysis of the derivatized (3R/S)-OH-nC14 standard

compounds yielded two peaks at retention times of 42.21 and 42.40 minutes, while one peak at

42.40 minutes was obtained for (3R)-OH-nC14 (Figure 2). We observed expected fragment ions

corresponding to the MPTA fragment ion, fatty acid ethyl ester, and fatty acid which supported

peak assignment (Figure S5). Thus (3S)-OH-nC14 and (3R)-OH-nC14 elute at 42.21 and 42.40

minutes, respectively. The compound 7 acyl chain was assigned as (3R)-OH-nC14 as its

derivatized 3-OH-tetradecanoate acyl chain eluted at 42.40 minutes (Figure 2).

The (3R)-OH-nC14 chain hydroxyl position and stereochemistry implicate possible

metabolic pathways leading to its biosynthesis. As fatty acid hydroxylases typically act on the

acyl chain terminal region, they are unlikely to catalyze C3 hydroxylation of a C14 fatty acid

(Pinot and Beisson, 2011). Of the other well-characterized pathways, only mitochondrial and

plastidial fatty acid metabolism mechanisms produce (3R)-OH-acyl-thioester intermediates.

Consistent with this result, previous work identified two trichome-expressed, mitochondrial

enzymes, Acylsugar Enoyl-CoA Hydratase 1 (AECH1) and Acylsugar Acyl-CoA Synthetase 1

(AACS1), and one trichome-expressed, plastidial enzyme, a beta-ketoacyl-(acyl-carrier-protein)

reductase (SpKAR1), involved in medium C10 and C12 acyl chain production in cultivated

tomato and its wild relative, *S. pennellii* (Fan et al., 2020; Ji et al., 2022).

We hypothesize that acylsugar hydroxyacyl chains evolved through one of three

scenarios. First, changes in acyl-CoA substrate availability, possibly through the actions of

thioesterases or acyl-CoA synthetases, enabled a promiscuous ASAT to esterify hydroxyacyl

chains to acylinositols. Alternatively, substrate specificity of an existing acylinositol ASAT

expanded to utilize available hydroxyacyl-CoAs. Like this second hypothesis, amino acid

substitutions between copies of ASAT2 from different tomato species impacted the accumulation

of differently branched five carbon acyl chains on acylsugars (Fan et al., 2015). A third

possibility is that a 'new' acyltransferase, unrelated to characterized ASATs and capable of

utilizing hydroxyacyl-CoA substrates, was co-opted into acylinositol biosynthesis. Future genetic

and biochemical studies should reveal the mechanisms behind (3*R*)-OH acyl chain production,

potentially providing new approaches to engineer unique acylsugars with hydroxyacyl chains

and hydroxylated fatty acids for polymeric building blocks. Acylsugar structures can impact their

function, thus we hypothesize that engineered tomato plants producing these unusual

hydroxyacyl chains on their acylsugars may exhibit different defense capabilities than the

wildtype acylsucroses.

Figure 2. Evidence that the 3-OH-n14 acyl chain on Compound 7, I3:22(i4,i4,n14-OH) is in the *R* configuration. Hydroxyacyl chains analyzed by GC-selected ion monitoring-MS (GC-SIM-MS) after ethyl transesterification and derivation to their diastereomeric (*S*)-MPTA form. Fragment ions with *m/z* of 189, 209, and 255 were monitored, and relative abundance was normalized to the highest peak. Yellow and green traces represent derivatives from commercial (3*R*/*S*)- and (3*R*)-OH-14:0 standards. The main derivatized 3-OH-14:0 acyl chain from Compound 7 (Figure 1), shown with the blue trace, comigrates with the peak corresponding to the *R* enantiomer.

Enormous *Solanum* acylsugar diversity revealed by LC-MS metabolite screening

To date, most acylsugar screening across the *Solanum* has focused on cultivated tomato and its Potato clade relatives (Fan et al., 2017; Ghosh et al., 2014; Lybrand et al., 2020; Schilmiller et al., 2016), which represent a small fraction of the species diversity in this genus (PBI Solanum Project, 2022). Though limited in number, studies profiling non-Potato clade species, including *S. nigrum* (Lou et al., 2021), *S. lanceolatum* (Herrera-Salgado et al., 2005), *S. quitoense* (Hurney, 2018; Leong et al., 2020), and *S. melongena* (this work) demonstrate acylsugar diversity across the *Solanum* is vast, and largely uncharacterized. To address this knowledge gap, we analyzed tissue surface extracts from 30 additional *Solanum* species,

including 23 Clade II, five DulMo clade, and two VANAns clade members (listed in Table S1).

While NMR is the gold standard for structural elucidation, it is prohibitively time consuming for

a largescale metabolite diversity survey. LC-MS coupled with CID provides substantial

structural information and is compatible with high-throughput screening. To strike a balance

between data quality and quantity, we performed LC-MS-CID screening on all 30 species and

annotated acylsugars as described for *S. melongena*. Twenty-five of the 30 analyzed species

produced detectable acylsugars from visible surface glandular trichomes, while species lacking

acylsugars had no or few observable glandular trichomes in the tissues analyzed. Targeted

analysis of leaf surface extracts from 31 species, including *S. melongena*, and fruit surface

extracts from two species, uncovered previously unreported acylsugars, inter-species acylsugar

differences, and identified the distributions of unusual acylsugar chemical traits in the *Solanum*

genus (Figure 3; Tables 1 and S2-27). Although we did not observe intraspecific acylsugar

variation, acylsugars varied enormously between species (Figure 4; Tables 1 and S2-27).

Figure 3. Distribution of acylsugar traits within the *Solanum* genus. Traits were mapped onto a previously published Supermatrix maximum likelihood phylogeny (Gagnon et al., 2022). The reported polytomy between Clade II, DulMO, VANANs, Regmandra, and Potato clades is not indicated (Gagnon et al., 2022). Acylsugar trait data from this report included annotations based upon LC-MS and NMR analysis (Figure 1, Tables 1 and S3-27), while data from previous studies were limited to those with NMR structural data (Ghosh et al., 2014; Herrera-Salgado et al., 2005; Hurney, 2018; King et al., 1986; Leong et al., 2020, 2019; Lou et al., 2021; Lybrand et al., 2020). (A) Distribution of sugar core types within the *Solanum* genus. Inositol and glucose cores determined by characteristic LC-MS negative mode electrospray ionization (ESI) fragmentation (Figures S2 and S10). (B) Distribution of species with acylsugars that coelute with eggplant acylsugars containing the same acyl chains and sugar core. (C) Distribution of hydroxyacyl chains and glycosylated hydroxyacyl chains.

Figure 4. *Solanum* species produce dramatically different leaf surface metabolite profiles. The seven representative species display morphological differences, as demonstrated by the flower and fruit images, and they also exhibit diverse metabolite profiles as demonstrated by base peak intensity LC-MS chromatograms from their leaf surface metabolites. Most of the peaks shown consist of acylsugars with varied sugar cores and acyl chain types and numbers. Acylsugar profiles for these species are detailed in Tables S5, S8, S18, S21, and S25-27.

Identification of diverse acyl chain types in *Solanum* acylsugars

Our survey of *Solanum* species uncovered acylsugars with a surprising diversity of acyl chain lengths, functional groups, and combinations of chain types and positions. We identified C2 to C18 acyl chains, including unsaturated and hydroxylated forms, based on analysis of fatty acid fragment carboxylate ions in negative mode and neutral losses in both negative and positive mode (Table S2). Acyl chain compositions differed between acylsugars within and across species: acylinositols with primarily medium C8 to C14 acyl chains were observed in five species (Figure 3, Tables S14, S15, S23, S25), while acylsugars with two short C4 to C6 acyl chains and one medium C8 to C14 acyl chain were observed in 17 species, an acylation pattern also observed in eggplant and wild tomato (Figure 1; Table 1) (Lybrand et al., 2020). Acylsugars with unsaturated acyl chains, including C5:1, C18:1, C18:2, and C18:3, were detected only in *S. torvum* (Table S18). To the best of our knowledge, this is the first report of unsaturated C18 chains in Solanaceae acylsugars. In contrast, hydroxylated medium-length acyl chains were surprisingly common: we annotated acylsugars bearing hydroxylated acyl chains in three DulMo clade species and 25 Clade II species (Figure 3C). These hydroxylated acyl chains are 'chemical handles' susceptible to further modifications, including glycosylations and acylations, which would result in even greater acylsugar diversity.

Indeed, we obtained evidence for glycosylated acyl chains in three Clade II species, *S. prinophyllum*, *S. sisymbriifolium*, *S. lasiophyllum*, and one DulMo species, *S. dulcamara* (Tables

S12, S13, S24 and S27). Several lines of evidence led us to conclude that these species

accumulate acylsugars with glycosyl groups attached via ether linkages to medium hydroxyacyl

chains esterified to the hexose core. First, the compounds fragmented abnormally under negative

mode CID: the glycosyl group was observed as a neutral loss from the [M-H]$^-$ ion, producing a

[sugar – H]$^-$ fragment ion (Figure S6). This contrasts with extensive observations that CID

spectra of other characterized acyldisaccharides do not exhibit negative mode disaccharide

glycosidic linkage cleavage (Figures S2 and S7) (Ghosh et al., 2014; Hurney, 2018; Lou et al.,

2021). Further evidence that the glycosylation is not on the primary acylated hexose ring was

obtained by subjecting acylsugar extracts to saponification, which induces ester linkage breakage

while retaining ether linkages. Upon analyzing saponified acylsugar extracts by LC-MS, we

detected pentose and hexose sugars with C14 or C16 acyl chains rather than glycosylated

hexoses, indicating that the glycosyl group is connected to a medium hydroxyacyl chain by an

ether linkage that is not cleaved during alkaline saponification (Figure S8). We named these

compounds glycohydroxyacylhexoses and annotated them by modifying the conventional

acylsugar nomenclature where the molecule in Figure S6 is named H3:22(4,4,14-O-p) with 14-

O-p representing the pentosylated (p) C14 hydroxyacyl chain (14-O). To our knowledge, this is

the first report of acylhexoses with glycosylated hydroxyacyl chains in Solanaceae acylsugars.

Resin glycosides produced by Convolvulaceae family species contain similar hydroxyacyl chain

glycosidic linkages. However, in resin glycosides, hydroxyacyl chains are connected to the

oligosaccharide core at two points – by an ester and ether linkage, respectively – forming a

macrocyclic structure not observed in the *Solanum* glycohydroxyacylhexoses (Bah and Pereda-

Miranda, 1996; Kruse et al., 2022; Pereda-Miranda et al., 1993).

395   The unusual glycohydroxyacylhexoses raises questions regarding their biosynthetic and

396 evolutionary origins. As these compounds are found in different lineages (Eastern Hemisphere

397 Spiny, Dulcamaroid, and Sisymbriifolium sections) – and closely related species lack detectable

398 glycohydroxyacylhexoses – this trait appears to be highly labile (Figure 3C). Because we

399 observe the cognate, non-glycosylated acylinositol in each of these species, we hypothesize that

400 glycohydroxyacylhexoses consist of a *myo*-inositol hexose core. Further analysis of acylsugars

401 from additional species may identify other types of modified acylsugar hydroxyacyl chains in

402 *Solanum* acylsugars.

403   Sixteen species accumulate acylsugars with the same molecular masses, sugar core

404 masses, and acyl chain complements as the NMR-characterized eggplant acylinositols,

405 suggesting they may be identical. To test this hypothesis, we mixed purified eggplant NMR-

406 resolved standards with metabolite extracts from each species and performed liquid co-

407 chromatography. Coelution provided one line of evidence that the eggplant-like acylsugars

408 detected in 11 Clade II species are identical to characterized eggplant acylinositols (Tables S5-

409 14, S27). In contrast, the eggplant-like acylsugars from five other species, including four DulMo

410 clade members and one Clade II member, eluted separately from characterized eggplant

411 acylinositols, indicating they are acylation positional isomers or acyl chain branching isomers

412 (Figure S9). Examples of acylinositol positional isomers were described between the DulMo

413 clade species *S. nigrum* and the Clade II species *S. quitoense* and *S. melongena*, providing

414 precedent that the non-coeluting compounds might be positional isomers of the eggplant

415 acylsugars shown in Figure 1 (Leong et al., 2020; Lou et al., 2021). Our data reveal a

416 phylogenetic pattern of acylinositol distribution in which Clade II species generally accumulate

417 eggplant-like acylinositol isomers, whereas DulMo clade species accumulate noneggplant-like

acylinositol isomers (Figure 3B). While the eggplant-like isomers coelute with eggplant

acylsugars, further analysis with complementary analytical methods is needed to confirm

whether these are structurally identical across species.

<u>Acylhexoses and acyldisaccharides accumulate throughout Clade II, DulMo, and VANAns</u>

<u>*Solanum* species</u>

Acylhexoses were detected in all analyzed acylsugar-producing species. They were

annotated as acylglucoses or acylinositols based upon differences in characteristic fragmentation

behavior under negative ion mode MS conditions (Leong et al., 2020; Lou et al., 2021).

Identification of acylglucoses was based on observation of a sugar core fragment ion of *m/z*

143.03 ($C_6H_7O_4^-$) and stepwise acyl chain losses at the lowest collision energy level (0 V)

(Figure S11). In contrast, acylinositol annotation was based on an absence of acyl chain fragment

ions at 0 V and MS^e CID negative mode functions (Figure S2). Using these rules, we identified

acylinositols in all acylsugar-producing species tested except for *S. americanum* (Tables 1, S2-

27). In contrast, acylglucoses were not detected in any Clade II species but were commonly

detected among the DulMo and VANAns species, found in all analyzed acylsugar-producing

species except *S. dulcamara* (Tables S2, S19, S20, S23, S24, S26; Figure 3A; Lou et al., 2021).

Differing from the widespread acylhexose distribution, acyldisaccharides were detected

sporadically in Clade II and DulMo clade species (Figure 3A), being observed in 14 of 21 Clade

II and 2 of 4 DulMo acylsugar-producing analyzed species. Based on negative [M-H]⁻ fragment

masses and positive mode fragmentation of the glycosidic linkage, the acyldisaccharides were

composed of hexose-hexose, pentose-hexose, or deoxyhexose-hexose sugar cores (Table S2,

Figures S2 and S7). Though complete structural information cannot be obtained from MS

fragmentation alone, two lines of evidence lead us to hypothesize that these acyldisaccharides are

composed of glycosylated inositol cores. First, most of the species with detectable

acyldisaccharides also accumulate cognate acylinositols with the same acyl chain lengths (leaf

and fruit surface extracts from 10 of 16 and two of two different species, respectively),

suggesting a shared inositol-containing core structure (Tables 1, S3-9, S12-17, S20, S22).

Second, acylinositol glycosides with varying sugar core sizes and stereochemistries were

described in *S. lanceolatum* (Herrera-Salgado et al., 2005), *S. quitoense* (Hurney, 2018), and *S.*

melongena (Figure 1), suggesting that the species in this study may accumulate similar

compounds. The ratio of acylhexose to acyldisaccharide peak number and abundance differs

across the surveyed species; for example, *S. mammosum* only accumulated detectable

acylinositols while *S. anguivi* primarily accumulated acyldisaccharides that comprised 80% of

the total acylsugar peak area (Tables S21 and S4). This is reminiscent of varied acylhexose

accumulation observed in trichomes of evolutionarily distant *Solanum* species: *S. nigrum* only

contains detectable acylinositol and acylglucose monosaccharides (Lou et al., 2021), while *S.*

pennellii accessions have mixtures of acylglucoses and acylsucroses ranging from 41-95% of

total acylsugars as acylglucoses. (Lybrand et al., 2020; Shapiro et al., 1994).

<u>Sticky fruits accumulate acylinositols</u>

We investigated the chemical basis of sticky fruit surface substances previously reported

in botanical species descriptions of the Clade II species *S. acerifolium* and *S. atropurpureum*

(Nee, 2022a, 2022b, 1991). LC-MS analysis of fruit surface extracts revealed similar acylsugar

profiles between the two species, with each extract containing what appeared to be the same 128

acylsugars. These fruit acylsugars were distinct from the 22 and 19 trichome acylsugars

identified in *S. acerifolium* and *S. atropurpureum*, respectively, as evidenced by their LC elution

and MS characteristics (Tables S14-17; Figure 5). Seventy-seven fruit acylsugars were identified

as acylinositols with two or three medium C8 to C14 acyl chains, including one to three

hydroxyacyl chains. The remaining 51 fruit acylsugars are acyldisaccharides containing an

unusual deoxyhexose-hexose core not previously reported in Solanaceae acylsugars. In contrast,

S. acerifolium and *S. atropurpureum* trichome acylsugars, composed of a pentose-hexose

disaccharide core decorated with short C5 or C6 acyl chains, were not detected in fruit surface

extracts.

Figure 5. *S. atropurpureum* trichomes and fruit exhibit dramatically different metabolite profiles. The top LC-MS base peak intensity (BPI) chromatogram displays metabolites from a leaf surface extract, while the bottom LC-MS base peak intensity chromatogram displays metabolites from a fruit surface extract. The leaf surface extract acylsugars elute from 2.2 to 7.6 min (Table S17), while fruit surface extract acylsugars elute from 4.6 to 18.4 min (Table S15).

S. mammosum and *S. capsicoides,* which are closely related to *S. acerifolium* and *S. atropurpureum*, do not produce sticky fruit surfaces, suggesting this trait evolved in the common ancestor of *S. acerifolium* and *S. atropurpureum,* perhaps as recently as 2 Mya (Särkinen et al., 2013). Fruit surface acylsugars were described previously within the *Physalis* genus and likely represent an independent evolutionary origin from the *S. acerifolium* and *S. atropurpureum* fruit acylsugars (Bernal et al., 2018; Cao et al., 2015; Cicchetti et al., 2018; Maldonado et al., 2006). Considering that, like eggplant, *S. acerifolium* and *S. atropurpureum* fruits lack trichomes (Nee, 2022a, 2022b), some other organ or cell type presumably synthesizes the fruit surface acylsugars.

Structures similar to *Solanum fernadesii* petiolar resin glands (Sampaio et al., 2021), *Hypericum*

androsaemum microscopic fruit glands (Caprioli et al., 2016), and fennel and chamomile fruit

secretory ducts (vittae) (Zizovic et al., 2007) may be involved in *S. acerifolium* and *S.*

atropurpureum fruit acylsugar secretion. Future work will be needed to understand the cellular,

biosynthetic, and evolutionary relationships between trichome and fruit acylsugars.

<u>Sugar core evolution across the Solanaceae</u>

Our screening revealed acylinositols to be broadly distributed across Clade II, DulMo,

and VANAns (Figure 3A, Table S1). Outside of this study, acylinositols were only reported in

Solanum species, including two Clade II members, *S. quitoense* and *S. lanceolatum*, and one

DulMo clade member, *S. nigrum*. The acylinositol distribution suggests that acylinositol

biosynthesis arose one or more times within the *Solanum* genus. However, our ability to

elucidate the likely number of acylinositol origins is impeded by the lack of resolution of the

phylogenetic relationships between the major *Solanum* clades, including Clade II, DulMo,

VANAns, Potato, and Regmandra (Gagnon et al., 2022). Further biochemical and genetic

analyses of acylinositol biosynthesis may reveal how this trait evolved, and whether the pathway

is conserved between Clade II, DulMo, and VANAns species.

Acylglucoses were reported to be sporadically present across the Solanaceae, including in

Salpiglossis sinuata, *Petunia*, *Nicotiana* spp., *Datura*, the DulMo clade *S. nigrum*, and Potato

clade *S. pennellii* (Castillo et al., 1989; Chortyk et al., 1997; Fiesel et al., 2022; Fobes et al.,

1985; Hurney, 2018; King and Calhoun, 1988; Lou et al., 2021; Matsuzaki et al., 1989; Schenck

et al., 2022; Van Dam and Hare, 1998). Recent work in *S. pennellii* and *S. nigrum* revealed that

acylglucoses are synthesized from acylsucroses by a neofunctionalized invertase-like enzyme,

AcylSucrose FructoFuranosidase 1 (ASFF1) (Leong et al., 2019; Lou et al., 2021). These results

show that conversion of acylsucroses to acylglucoses arose at least twice in *Solanum* because *S.*

nigrum and *S. pennellii* employ non-orthologous ASFF1 enzymes. Our current study extends

documented cases of *Solanum* acylglucose distribution to the VANAns clade, as well as

documenting it in additional DulMo species. In contrast, we found no acylglucose accumulating

species among the 23 analyzed *Solanum* Clade II species. Elucidation of acylglucose

biosynthesis outside of the *Solanum* clade will be needed to resolve the number of evolutionary

origins and biochemical mechanisms leading to synthesis of these metabolites across the

Solanaceae.

Varied acyldisaccharides were observed across DulMo and Clade II, and we predict they

are biosynthesized from acylinositols. Within the Potato clade as well as outside the *Solanum*

genus, acylsucroses are the dominant acyldisaccharide in the reported species and tissues (Fiesel

et al., 2022). In contrast, no acylsucroses were detected in any species screened. However,

acylsucrose biosynthesis might be found in DulMo and VANAns species because acylglucoses

were detected in all but one DulMo and VANAns species and that acylsucroses can be

intermediates in acylglucose biosynthesis (Leong et al., 2019; Lou et al., 2021). The lack of both

acylglucoses and acylsucroses in the analyzed Clade II species suggests a loss of acylsucrose

biosynthesis in Clade II. Rather, our LC-MS data are consistent with the hypothesis that the

acyldisaccharides observed in this study consist of glycosylated inositol cores. Recent *in vitro*

biochemistry and *in vivo* genetic evidence in *S. quitoense* suggests that acylinositols are the

precursors to acylinositol disaccharides in Clade II *Solanum* species (Leong et al., 2022, 2019).

In this scenario, acylinositols would be converted to their cognate glycosides by one or more

glycosyltransferases, with different sugars added either by a conserved promiscuous

glycosyltransferase or multiple glycosyltransferases. *S. quitoense* and *S. melongena*, with their

acylinositol glycosides bearing different glycosyl moieties, are promising systems for addressing

the biochemical origins of these unusual *Solanum* acyldisaccharides.

Figure 6. SmASAT3-L1 acetylates AI3:16 to produce AI4:18. Trichome expressed ASAT
homologs were identified and tested for AI3:16 acetyl transferase activity. (A) LC-MS analysis

of AI3:16(4,4,8) and AI4(2,4,4,8) in *S. aethiopicum* (red) and *S. melongena* (purple) leaf surface
extracts. *S. aethiopicum* does not produce detectable levels of AI4:18. (B) Shown is a clade III
BAHD tree subset from a phylogeny including 106 predicted BAHDs (PF002458) in the
eggplant Smel_V4.1 reference genome, published reference BAHD sequences for clades I-VII,
characterized ASAT sequences from other selected Solanaceae species, and the SaASAT3 and
SaASAT3-L1 candidates from *S. aethiopicum* (see Figure S12 for full phylogeny). The
maximum likelihood tree was inferred from amino acid sequences using the Jones-Taylor-
Thornton algorithm with seven rate categories in IQ-TREE v2.1.3. Values at nodes indicate
bootstrap support calculated from 100,000 ultrafast bootstrap replicates. Heatmap shows absolute
transcript abundance (log2 TPM) across trichomes, trichomeless hypocotyls, and roots from 7-
day-old eggplant seedlings. Expression data for non-eggplant sequences are not included in the
heatmap. Color gradient provides a visual marker to rank the transcript abundance from high
(purple) to low (white) or absent (grey). ASAT, acylsugar acyltransferase. log2TPM, log2
transformed transcripts per million. Genes denoted with "*" were selected for biochemical
characterization based on expression in trichomes and sequence similarity to characterized
ASAT enzymes in other species. (C) Of the seven enzymes tested *in vitro*, only SmASAT3-L1
successfully acetylated AI3:16(4,4,8) to form AI4:18(2,4,4,8). The extracted ion chromatograms
on the left display products from reverse enzyme assays and the formate adduct of AI3:16, *m/z*
623.29. The extracted ion chromatograms on the right display products from forward enzyme
assays and the formate adduct of AI4:18, *m/z* 665.30. (D) SmASAT3-L1 acetylates AI3:16 *in*
vitro to produce AI4:18. Forward assay extracted ion chromatograms display the formate adduct
of AI4:18, *m/z* 665.30, and reverse assay extracted ion chromatograms display the formate
adduct of AI3:16, *m/z* 623.29. The reverse assay chromatograms display peaks for both AI3:16
and AI4:18 due to in source fragmentation of remaining AI4:18.

S. aethiopicum is defective in expression of a trichome acylsugar acyltransferase enzyme

We began analyzing the biochemical mechanisms underlying the observed *Solanum*

acylsugar diversity in seven of 23 members from the Eggplant clade and related Anguivi grade,

two small subclades within Clade II that includes brinjal eggplant and its close relatives (Figure

S11) (Aubriot et al., 2018). In contrast to the other six analyzed eggplant species, *S. aethiopicum*

(scarlet or Ethiopian eggplant) is unique in that it had unacetylated AI3:16(4,4,8) (Compound 1)

but did not accumulate detectable levels of the acetylated form, AI4:18(2,4,4,8) – the highest

abundance *S. melongena* acylsugar (Compound 2; Figures 6A, S11, Tables 1, S2, S5). We

hypothesized that the absence of AI4:18(2,4,4,8) in *S. aethiopicum* is either due to loss-of-

function or altered expression of an acetylating enzyme in this species. The availability of *S.*

melongena and *S. aethiopicum* genomic sequences (Barchi et al., 2021; Song et al., 2019)

provided the opportunity to seek a trichome expressed acetyltransferase responsible for

producing AI4:18 and the mechanism behind the lack of detectable AI4:18 in *S. aethiopicum*.

To identify enzyme candidates, we sequenced transcriptomes of isolated trichomes,

trichome-depleted shaved hypocotyls, and whole roots collected from 7-day-old *S. melongena*

seedlings and performed differential gene expression analysis (see Methods). Of the 23,251

eggplant genes expressed in at least one of the three tissues, 745 were significantly enriched

(log2 fold-change >2, *p*-value < 0.05) in trichomes compared to hypocotyls and roots, including

20 BAHDs (Tables S37 and S38). As shown in Table S39, we selected for further testing seven

BAHDs that were abundantly expressed in trichomes (TPM > 200) and homologous to

characterized *Solanum* ASATs (sequences in colored text and denoted with "*" in Figure 6B)

(D'Auria, 2006; Fan et al., 2015; Leong et al., 2022, 2020; Lou et al., 2021; Schenck et al.,

2022). We expressed each of the seven candidates in *Escherichia coli* and tested for acetylation

of Compound 1, AI3:16(4,4,8). An SlASAT3 outparalog (Koonin, 2005) SMEL4.1_12g015780,

ACYLSUGAR ACYLTRANSFERASE3-LIKE1 (SmASAT3-L1) was the only enzyme to

exhibit detectable forward activity acetylating Compound 1, AI3:16(4,4,8), to Compound 2,

AI4:18(2,4,4,8) (Figure 6C). As shown in Figure S13, LC-MS analysis of the SmASAT3-L1 *in*

vitro assay product revealed that it had identical molecular masses and elution times to plant-

derived AI4:18(2,i4,i4,i8) (Compound 1) providing evidence that SmASAT3-L1 acetylates

AI3:16(i4,i4,i8). Characterization of reverse activities of the seven BAHD acyltransferases

(Leong et al., 2020; Lou et al., 2021), in which an acyl chain can be removed from an acylsugar

by incubating it with the enzyme and free Coenzyme A, confirmed that SmASAT3-L1 was the

only tested enzyme that removed an acetyl chain from AI4:18(2,4,4,8) (Figures 6D and S13).

Taken together, these results indicate that trichome-expressed *SmASAT3-L1* encodes an

acylinositol acetyltransferase responsible for AI4:18(2,4,4,8) biosynthesis.

We tested the hypothesis that *S. aethiopicum* trichome extracts lack detectable acetylated

AI4:18(2,4,4,8) due to a defect in *SmASAT3-L1* ortholog expression. Indeed, reverse

transcription (RT)-PCR comparing isolated trichome and trichomeless shaved hypocotyl cDNA

revealed that *SaASAT3-L1* (GenBank ID: OQ547782) transcript was undetectable in *S.*

aethiopicum trichomes (Figure S14). This result implicates the defect in ASAT3-L1 enzyme

gene expression as responsible for lack of AI4:18 in *S. aethiopicum*. The interspecific variation

in *ASAT3-L1* expression is reminiscent of the ASAT4 expression differences among accessions

of the wild tomato *Solanum habrochaites*. In that case, a subgroup of *S. habrochaites* accessions

possessed a functional ASAT4 copy, but its low levels of gene expression correlated with

reduced accumulation of acetylated acylsugars in these 'subgroup E' accessions (Kim et al.,

2012).

611

Figure 7. Characterized *in vitro* acyl-CoA usage by ASATs. The maximum likelihood phylogeny was inferred from amino acid sequences using a time-reversible algorithm specifying a plant-specific empirical substitution matrix, invariant sites, and four rate categories (Q.plant+I+G4) in IQ-TREE v2.1.3. Bootstrap support was calculated from 100,000 ultrafast bootstrap replicates; values >=95 indicate strong support. Other than SmASAT3-L1, enzyme activities were described previously (Fan et al., 2017, 2015; Leong et al., 2022, 2020; Lou et al., 2021; Moghe et al., 2017; Nadakuduti et al., 2017; Schenck et al., 2022; Schilmiller et al., 2015, 2012). Only reported, positive activities are displayed with the black squares. Lack of activity or untested activities are displayed with blank squares.

<u>Natural variation potentiates novel activities and uses</u>

Until recently, it appeared that acylsucroses dominate acylsugar profiles of Solanaceae species with a small number of documented examples of acylglucose- and acylinositol-producing *Solanum* species (Fiesel et al., 2022; Fobes et al., 1985; Herrera-Salgado et al., 2005; Hurney, 2018; King and Calhoun, 1988; Leong et al., 2020; Liu et al., 2017; Lou et al., 2021; Moghe et al., 2017; Schenck et al., 2022). Our chemotaxonomic survey of nearly three dozen *Solanum* species from the sparsely sampled Clade II, DulMo, and VANAns clades revealed widespread occurrence of glucose-, inositol-, and non-sucrose disaccharide-based acylsugars decorated with unusual acyl chains, including medium-length (C12-C16) hydroxyacyl chains, glycosylated hydroxyacyl chains, and unsaturated chains. Considering that this work and previously published analyses together cover <3% of *Solanum* species, this presumably is only a subset of trichome acylsugar diversity within this large genus. Adding to interspecific variation we observe here, the recent identification of acylsugars in cultivated tomato root exudates supports the value of metabolite screening in additional tissue types as well as species (Korenblum et al., 2020).

With 33 acyl chains (Lou et al., 2021; Leong et al., 2020; Herrera-Salgado et al., 2005) and six inositol acylation positions, there are 1.3 billion theoretically possible acylinositol structures, not accounting for glycosylated acylinositols and glycohydroxyacylhexoses. This is in addition to the 820 million theoretically possible acylsucrose structures (Fan et al., 2019), and

demonstrates that this study greatly expands our knowledge of known and potential acylsugar

structural diversity.

Knowledge of *Solanum* acylsugar diversity provides a framework for investigating the

molecular basis of metabolic pathway evolution. For example, the widespread occurrence of

acylglucoses across the DulMo clade leads us to hypothesize that the neofunctionalized invertase

responsible for *S. nigrum* acylglucose synthesis (Lou et al., 2021) is shared by other DulMo

clade species. Similarly, documented acylsugar variation between cultivated eggplant, *S.*

melongena, and its relative, *S. aethiopicum*, led to the discovery of a terminal acylsugar

acetyltransferase in eggplant, SmASAT3-L1. The characteristics of ASAT3-L1, the enzyme

absent from the *S. aethiopicum* trichome transcriptome, in turn reveals the dynamic nature of

plant specialized metabolism. First, SmASAT3-L1 is unique among characterized ASATs in its

acyl acceptor specificity, acetylating an acylinositol glycoside rather than an acylated sucrose,

glucose, or *myo*-inositol. Second, acetyl-CoA donor activity was previously only described for a

subset of Clade III BAHDs, which include ASAT4 and ASAT5 clade members (Figure 7).

SmASAT3-L1 breaks a pattern seen across the Solanaceae, including trichome ASATs from

species of early evolving lineages, *Petunia axillaris* and *Salpiglossis sinuata* (Moghe et al., 2017;

Nadakuduti et al., 2017), *Nicotiana attenuata* (Schenck et al., 2022), as well as cultivated and

wild tomatoes (Kim et al., 2012; Schilmiller et al., 2012). This theme was also observed in

trichomes of two other *Solanum* plants, more closely related to the species screened in this study:

the Clade II *S. quitoense* SqTAIAT (Leong et al., 2020) and DulMo *S. nigrum* SnAGAT1 (Lou et

al., 2021). Both enzymes reside outside of the ASAT3 clade, and thus are phylogenetically

distinct from SmASAT3-L1. SmASAT3-L1 characteristics and those of recently published *S.*

quitoense enzymes (Leong et al., 2022, 2020) also suggest that acylinositols are synthesized

through one or more pathway(s) distinct from that of acylsucroses. Existing genetic resources in

S. nigrum, *S. quitoense*, and *S. melongena* can be utilized to explore when and how acylinositols

arose and to identify the underlying mechanisms behind acylation position and chain type

differences between these species.

The remarkably varied structures of acylsugars strongly suggest that they have evolved

distinct bioactivities, likely in coevolutionary arms races between plants and their insect and

microbial pests. For example, hydrophobic acylsugars with longer acyl chains and few free

hydroxyl groups may disrupt membranes, reminiscent of less polar triterpene saponin variants

(Augustin et al., 2011; Baumann et al., 2000). Variation in acyl chain linkage chemistries also

likely influence acylsugar mode of action. For example, ester linked short chains acylsucroses

are digested by *Manduca sexta* (hawkmoth) larvae and the volatile organic acids released into the

environment, attracting ant predators (Weinhold and Baldwin, 2011). In contrast, longer chain

hydroxylated and ether-linked glycohydroxyacyl chains observed in the study are likely to have

quite different metabolic fates, perhaps persisting in the digestive systems of herbivores, and

alternative metabolic fates in microbes. Uncovering the enzymes responsible for synthesis of the

wide variety of natural acylsugars will enable transgenic production of isogenic variants and

rigorous testing of their ecological activities. Similarly, access to a wider variety of often

promiscuous ASAT BAHD acyltransferases (Moghe et al., 2023) creates opportunities for

producing novel acylsugars with a wide range of physical properties and bioactivities (Schenck

et al., 2022).

Hydroxyacyl chains also present promising opportunities to expand upon natural

acylsugar diversity through synthetic biology and/or synthetic chemistry. The hydroxyl acts as a

reactive chemical handle, which was exploited in a subset of *Solanum* species trichomes to create

glycohydroxyacylhexoses. This handle can also be exploited with synthetic chemistry using

these chains as new feedstocks for creating polymers. These chains may also be useful in

creating synthetic acylsugars for use as food-grade emulsifiers. We can also further modify the

hydroxyacyl chains to add unusual sugars, acyl chains, and aromatic groups through

promiscuous enzymes or synthetic chemistry. These structural changes would likely impact

biological activities and add another asset to developing more pest resistant plants.

Characterizing more natural acylsugar diversity will further enable synthetic biology approaches

to create completely new acylsugar structures.

Methods

<u>Plant material and growth conditions</u>

Solanum spp. seeds were obtained from the sources described in Table S1. Seeds were

treated with 10% (v/v) bleach for 10 min while being rocked at 24 rpm with a GyroMini nutating

mixer (Labnet, Edison, NJ, USA), and then rinsed 5-6 times with distilled water. Unless

otherwise noted, seeds were germinated on Whatman filter paper (MilliporeSigma, Burlington,

MA, USA) at 28°C and in the dark. Germinated seedlings were transferred to peat pots (Jiffy,

Zwijndrecht, Netherlands), and grown at 25°C, 16/8-h day/night light cycle, and ~70 µmol $m^{-2}s^{-2}$

photosynthetic photon flux density with cool white fluorescent bulbs. Mature plants were grown

in controlled environment growth chambers or in a greenhouse. The growth chamber conditions

consisted of 25°/18°C day/night temperatures, 16/8-h day/night light cycle, and ~100 µmol $m^{-2}s^{-2}$

photosynthetic photon flux density under LED bulbs. The greenhouse conditions consisted of a

16/8-h day/night light cycle achieved with supplemental sodium iodide lighting. Plants were

fertilized weekly with 0.5X Hoagland's solution.

Surface metabolite extractions

Surface metabolites were extracted as described previously (Leong et al., 2019; Lou and

Leong, 2019). Briefly, 0.1-1 g of leaf, stem, hypocotyl, or cotyledon tissue was collected into a

1.5 mL screw-cap tube (Dot Scientific, Inc., Burton, MI, USA), 1 mL extraction solvent (3:3:2

acetonitrile:isopropanol:water, 0.1% formic acid, 10 µM propyl 4-hydroxybenzoate (internal

standard)) was added and the sample was rocked at 24 rpm by a GyroMini nutating mixer for

two min. After extraction, the supernatant was transferred into a glass 2 mL autosampler vial

(Restek, Bellefonte, PA, USA) and sealed with a 9 mm cap containing a PTFE/silicone septum

(J.G. Finneran, Vineland, NJ, USA). For fruit surface metabolite extractions, we modified the

protocol by placing 1-2 fruit into a 15 mL polypropylene conical tube (Corning Inc., Corning,

NY, USA) containing 5 mL extraction solvent then proceeded to the nutation step above.

Bulk eggplant acylsugar extraction and purification for NMR analysis

Surface metabolites were bulk extracted from approximately 1000 *S. melongena* PI

555598 seedlings. Seeds were treated with 10% (v/v) bleach (Clorox, Oakland, CA, USA) for 10

min while being gently rocked at 24 rpm with a GyroMini nutating mixer and subsequently

rinsed 5-6 times with distilled water. Seeds were sown in moist SUREMIX soil (Michigan

Grower Products, Galesburg, MI, USA) in Perma-Next plant trays, 22 x 11 x 2.5 inches

(Growers Supply Company, Dexter, MI, USA), covered with a humidity dome, 22 x 11 x 3

inches (Growers Supply Company), then transferred to the growth chamber immediately.

Seedlings were harvested when 2-3 true leaves were observed, approximately one week, by

cutting them at the base and placing them into two 2L beakers each containing 1L 100%

acetonitrile. Surface metabolites were extracted with gentle agitation with a metal spatula for five

min at room temperature. Plant material and sediment was removed by vacuum filtration through

a Büchner funnel lined with Whatman filter paper (MilliporeSigma). solvent was removed *in*

vacuo by rotary evaporation and dried residue was dissolved in 20 mL of acetonitrile with 0.1%

formic acid. Solvent was removed again using a vacuum centrifuge and the dried residue was

dissolved in 1 mL of acetonitrile:water:formic acid (80:20:0.001). The semi-preparative LC

method is described in detail in Table S28 and is described in brief here. *S. melongena*

acylsugars were separated with a Waters 2795 HPLC (Waters Corporation, Milford, MA, USA)

equipped with an Acclaim 120 C18 HPLC column (4.6 x 150 mm, 5 μm; Thermo Fisher

Scientific, Waltham, MA, USA). Solvent A was water with 0.1% formic acid and Solvent B was

acetonitrile. The reverse-phase linear gradient was as follows: 5% B at 0 min, 60% B at 2 min,

80% B at 40 min, 100% B at 42 min, 5% B at 42.01 min, held at 5% B until 44 min. Flow rate

was 1.5 mL/min, injection volume was 100 μL, and the column temperature was 40°C. Fractions

were collected automatically at 0.25 min intervals by a 2211 Superrac fraction collector (LKB

Bromma, Stockholm, Sweden) and assessed for acylsugar presence and purity by LC-MS

analysis, as described below. Column fractions were collected in the same tubes for each method

run which worked to pool the fractions between each run.

<u>LC-MS acylsugar analysis</u>

Acylsugars were analyzed by LC-MS with each method described below and in Tables

S29-33. For each LC method, the mobile phase consisted of aqueous 10 mM ammonium

formate, adjusted to pH 2.8 with formic acid (Solvent A) and 100% acetonitrile (Solvent B). The

flow rate was maintained at 0.3 mL/min.

756  Acylsugar extracts were analyzed with a 22 min LC gradient using a Waters Acquity

757 UPLC coupled to a Waters Xevo G2-XS QToF mass spectrometer (Waters Corporation, Milford,

758 MA) equipped with electrospray ionization (ESI) operating in positive (ESI[+]) and negative (ESI[-])

759 modes. Ten microliter acylsugar extracts were separated on an Acquity UPLC BEH C18 column

760 (10 cm x 2.1 mm, 130 Å, 1.7 μm; Waters), kept at 40°C, using a binary solvent gradient. The 22

761 min linear gradient was as follows: 5% B at 0 min, 60% B at 2 min, 100% B at 16 min, held at

762 100% B until 20 min, 5% B at 20.01 min, held at 5% B until 22 min. Acylsugar extracts were

763 analyzed with both ESI[-], and ESI[+]. For ESI[-], the following parameters were used: capillary

764 voltage, 2 kV; sampling cone voltage, 35 V; source temperature, 100°C; desolvation temperature

765 350°C; cone gas flow, 50 L/Hr; desolvation gas flow, 600 L/Hr. For ESI[+], the following

766 parameters were used: capillary voltage, 3 kV; sampling cone voltage, 35 V; source temperature,

767 100°C; desolvation temperature 300°C; cone gas flow, 50 L/Hr; desolvation gas flow, 600 L/Hr.

768 Acylsugars were fragmented in either MS[E] or data-dependent acquisition (DDA) MS/MS modes

769 as described previously (Lou et al., 2021; Lybrand et al., 2020). DDA survey and MS/MS

770 functions acquired over *m/z* 50 to 1500 with scan times of 0.2 s. Ions selected for MS/MS were

771 fragmented with a ramped collision energy where voltage varies based on ion mass. Collision

772 energies followed a ramp with the voltage changing linearly for ions between the low mass

773 setting (*m/z* 50, 15 to 30 V) and high mass setting (*m/z* 1500, 30 to 60 V). To increase mass

774 accuracy, lock mass correction was performed during data collection, with leucine enkephalin as

775 the reference.

776  Semi-preparative LC fractions were analyzed with direct infusion and a 14-min LC

777 gradient using a LC-20ADvp ternary pump (Shimadzu, Kyoto, Japan) coupled to a Waters Xevo

778 G2-XS QToF mass spectrometer (Waters Corporation). The direct infusion method quickly

screened fractions for acylsugar presence while the 14-min LC method tested fraction purity. Ten

microliters of acylsugar fractions were injected into an Ascentis Express C18 HPLC column (10

cm x 2.1 mm, 2.7 μm; Supelco, Bellefonte, PA, USA), kept at 40°C. The 14-min gradient was as

follows: 5% B at 0 min, 60% B at 2 min, 100% B at 10 min held until 12 min, 5% B at 12.01

min, and held at 5% until 14 min. Fractions were analyzed under ESI$^+$ with the following

parameters: capillary voltage, 3 kV; sampling cone voltage, 40 V; source temperature, 100°C;

desolvation temperature 350°C; cone gas flow, 20 L/hr; desolvation gas flow, 500 L/hr. For

direct infusion analysis, ions were acquired from m/z 50 to 1500 with a scan time of 0.1 s and

one acquisition function with 0 V collision potential. For 14-min LC analysis, ions were acquired

from m/z 50 to 1200 with a scan time of 0.1 s and three acquisition functions with different

collision potentials (0, 25, 60 V). Lock mass calibration referenced to the leucine enkephalin

[M+H]$^+$ ion was applied during data acquisition.

Enzyme assays products were analyzed with a 7-min gradient using Waters Acquity

UPLC coupled to a Waters Xevo G2-XS QToF mass spectrometer (Waters Corporation, Milford,

MA) equipped with electrospray ionization (ESI) operating in negative (ESI$^-$) mode. Reaction

products were separated with an Ascentis Express C18 HPLC column (10 cm x 2.1 mm, 2.7 μm;

Supelco), kept at 40°C, using a binary solvent gradient. The 7 min linear gradient was as follows:

5% B at 0 min, 60% B at 1 min, 100% B at 5 min, held at 100% B until 6 min, 5% B at 6.01 min,

held at 5% B until 7 min. The ESI$^-$ parameters described for acylsugar extract analysis were

used. Ions were acquired from m/z 50 to 1200 with a scan time of 0.1 s and three acquisition

functions with different collision potentials (0, 25, 60 V). Lock mass calibration referenced to the

leucine enkephalin [M+H]$^-$ ion was applied during data acquisition.

For coelution analysis between enzymatically- and plant-produced AI3:16 and AI4:18, a

24-min linear gradient was used with an Acquity UPLC BEH C18 column (10 cm x 2.1 mm, 130

Å, 1.7 μm; Waters), kept at 40ºC, on the same instrument used for enzyme assay analysis. The

binary solvent, linear gradient was as follows: 5% B at 0 min, 60% B at 18 min, 100% B at 18.01

min, held at 100% B until 22 min, 5% B at 22.01 min, held at 5% B until 24 min. ESI$^-$

parameters and MS method from the above enzyme assay analysis were used.

Saponified acylsugars were analyzed using a Waters Acquity UPLC coupled to a Waters

Xevo G2-XS QToF mass spectrometer (Waters Corporation, Milford, MA). Ten microliters of

saponified acylsugars were injected into either an Acquity UPLC BEH C18 column (10 cm x 2.1

mm, 130 Å, 1.7 μm; Waters) or an Acquity UPLC BEH Amide column (10 cm x 2.1 mm, 130 Å,

1.7 μm; Waters), kept at 40°C. Samples were analyzed on the C18 column with the 22-min

method described above. To detect free sugars resulting from saponification, samples were

analyzed on the BEH amide column with a 9 min method which was as follows: 95% B at 0 min

held until 1 min, 60% B at 6 min, 5% B at 7 min, 95% B at 7.01 min, held at 95% B until 9 min.

Both methods used the ESI- parameters and MS acquisition parameters described for enzyme

assays.

<u>Acylsugar annotation</u>

We inferred acylsugar structures utilizing negative and positive mode MS and MS/MS

collision-induced dissociation (CID) as previously described (Ghosh et al., 2014; Hurney, 2018;

Leong et al., 2019; Lou et al., 2021; Lybrand et al., 2020). We employed modified confidence

criteria developed by Lou and coworkers (Lou et al., 2021). Putative acylsugars only meeting

criterion A were annotated as low confidence. Putative acylsugars meeting criteria A-D were

annotated as medium confidence. We also added two additional criteria, G and H, described

below. Acylsugars meeting criteria A-F and either G or H were annotated with highest

confidence. Confidence levels for each reported acylsugar are specified in their species-specific

annotation table (Tables S2-27).

(G) **NMR structural determination.** Acylsugars from *S. melongena* were further characterized

by NMR experiments in which sugar core relative stereochemistry, acyl chain positions, and acyl

chain branching were resolved.

(H) Coelution with NMR characterized compounds. Acylsugars with the same acyl chains

and sugar core size were analyzed by coelution with *S. melongena* acylsugars. If acylsugars

coeluted with an NMR characterized peak and shared the same acyl chains and fragmentation

patterns, they met this criterion and were annotated with high confidence. This criterion was

employed for acylsugars from species other than *S. melongena*.

Acylsugar metabolomic analysis

To collect accurate formate adduct masses and relative abundances from putative

acylsugars, negative mode LC-MS raw data from metabolite extracts were analyzed with

Progenesis QI software v3.0 (Waters, Milford, MA, USA) using retention time alignment, peak

detection, adduct grouping, and deconvolution. Lock mass correction was performed during data

collection. We used the following analysis parameters: peak picking sensitivity, default;

retention time range, 0-22 min; adducts detected, M+FA-H, M+Cl, M-H, 2M-H, and 2M+FA-H.

Average percent peak abundance was calculated for each putative acylsugar by dividing its raw

abundance by the sum of the total acylsugar raw abundance and then averaging that value across

all samples for a species.

NMR analysis of acylsugars

Purified acylsugars were dissolved in acetonitrile-d_3 (99.96 atom % D; MilliporeSigma, Burlington, MA, USA) and transferred to Kontes NMR tubes (MilliporeSigma, Burlington, MA, USA). Samples were analyzed at the Michigan State University Max T. Rogers NMR Core with a Varian Inova 600 MHz spectrometer (Agilent , Santa Clara, CA, USA) equipped with a Nalorac 5 mm PFG switchable probe, a DirectDrive2 500 MHz spectrometer (Agilent, Santa Clara, CA, USA) equipped with a OneNMR probe, Bruker Avance NEO 800 MHz spectrometer (Bruker, Billerica, MA, USA) equipped with a 5 mm helium cryogenic HCN probe, or a Bruker Avance NEO 600 MHz spectrometer equipped with a 5 mm nitrogen cryogenic HCN Prodigy probe. All acylsugars were analyzed with a series of 1D (^{1}H, and ^{13}C) and 2D (gCOSY, TOCSY, gHSQC, gHMBC, gH2BC, J-resolved) experiments. Resulting spectra were referenced to acetonitrile-d_3 ($^{\delta}$H = 1.94 and $^{\delta}$C = 118.70 ppm). NMR spectra were processed and analyzed with MestReNova 12.0 software (MestreLab, A Coruña, Spain). For full NMR metadata, see Supplementary Datafile 1.

Sugar core composition analysis

S. melongena PI 555598 acylsugar extracts were dried down *in vacuo* with centrifugation (Savant, ThermoFisher Scientific). Dried acylsugar extracts were dissolved in 1 mL of methanol. One mL of 3 M ammonium hydroxide was added, and the solution was mixed vigorously. This initial saponification reaction proceeded at room temperature for 48 hours. Solvent was removed by vacuum centrifugation. Saponified sugar cores were reduced and derivatized with acetate groups as previously described (Sassaki et al., 2008). Sugars were dissolved in 50 µL of 1 M

ammonium hydroxide for 15 min at room temperature. Addition of 50 µL 20 mg/mL NaBH$_4$ and

incubation at 100°C for 10 min converted aldoses and ketoses to polyols. Excess sodium

borohydride was quenched with 100 µL 1M trifluoroacetic acid. Two volumes of methanol were

added, and the sample was dried down by a stream of nitrogen gas. Dried down reaction products

were redissolved in 200 µL of methanolic 0.5 M HCl and heated at 100°C for 15 min in a closed

½ dram vial, 12 x 35 mm (Kimble Chase, Vineland, NJ, USA). Solvent was removed by a stream

of nitrogen gas. To acetylate the sugar cores, residue was dissolved in 200 µL of pyridine:acetic

anhydride (1:1) and incubated at 100°C for 30 min. Acetylated polyols were then dried down *in*

vacuo in a vacuum centrifuge and redissolved in 100 µL hexane for GC-MS analysis.

To determine the composition of disaccharide sugar cores, saponified sugars were

hydrolyzed prior to derivatization. Dried down sugars were redissolved in 98 µL water in a ½

dram vial, 12 x 35 mm (Kimble Chase). After addition of 102 µL of 88% formic acid, the vial

was sealed and heated at 100°C for 15 hours in a heat block. Hydrolyzed sugars were dried down

in vacuo with a vacuum centrifuge and derivatized to alditol acetates following the method

described above.

<u>Hydroxyl acyl chain stereochemistry analysis</u>

We derivatized hydroxyacyl chains and extracted the derivatives as previously described

(Jenske and Vetter, 2007). The commercial standards (3*R*)-OH-tetradecanoate (Cayman

Chemical, Ann Arbor, MI, USA) and (3*R/S*)-OH-tetradecanoate (Cayman Chemical, Ann Arbor,

MI, USA) were first derivatized to their respective ethyl ester derivatives. 600 µg of each fatty

acid was dissolved in 0.5 mL of 0.5 M ethanolic KOH and incubated at 80°C for 5 min. After the

solution was cooled on ice, 1 mL of ethanolic BF$_3$ was added, and reactions were incubated at

80°C for five min. Two-phase partitioning with hexane and saturated sodium chloride yielded

ethyl ester derivatives in the hexane partition.

The 3-OH-nC14 acyl chain from I3:22(iC4,iC4,3-OH-nC14) was derivatized to an ethyl

ester following a previously published method (Fan et al., 2020; Ning et al., 2015). Half of the

purified I3:22(iC4,iC4,3-OH-nC14) was dissolved in acetonitrile with 0.1% formic acid and

transferred to a 1.5 mL microcentrifuge tube. The purified acylsugar was dried down *in vacuo*

with vacuum centrifugation. The resulting dry, purified I3:22(iC4,iC4,3-OH-nC14) was

dissolved in 300 µL of 21% (w/w) sodium ethoxide in ethanol (MilliporeSigma, Burlington,

MA, USA). The reaction proceeded at room temperature for 30 min with constant rocking at 24

rpm by a GyroMini nutating mixer (Labnet, Edison, NJ, USA) and occasional vortexing. 400 µL

hexane with 55 ng/µL tetradacane (MilliporeSigma, Burlington, MA, USA) as an internal

standard was then added followed by vigorous vortexing. 500 µL of aqueous saturated sodium

chloride was added to the hexane-ethanol mixture, and vortexed. The hexane layer was pipetted

to another tube and two more two-phase partitions with saturated sodium chloride were

completed. The final hexane layer containing the acyl chain ethyl esters was extracted and dried

down by a stream of nitrogen gas.

The resulting fatty acid ethyl esters from the commercial standards and purified

compound were derivatized with (*R*)-(-)-α-methoxy-α-trifluoromethylphenylacetyl chloride ((*R*)-

(-)-MTPA-Cl; MilliporeSigma, Burlington, MA, USA) following a previously published

procedure (Jenske and Vetter, 2007). Dried fatty acid ethyl esters were redissolved in 400 µL

pyridine and 15 µL of (*R*)-(-)-MPTA-Cl. The reaction proceeded at room temperature for two

hours, after which, 5 mL of water and tert-butyl methyl ether (TBME) were added along with

solid K_2CO_3 (one spatula tip). The TBME phase was collected after three successive phase

separations with 5 mL of water. TBME was evaporated to 1 mL and subjected to GC-MS

analysis.

GC-MS analysis

All GC-MS analyses employed an Agilent 5890 GC and an Agilent 5975 single

quadrupole MS equipped with a FactorFour VF-5ms column (30 m x 0.25 mm, 0.25 um;

Agilent) and 10 m EZ-Guard column (Agilent, Santa Clara, CA, USA). Helium was used as the

carrier gas with a constant flow rate of 1 mL/min. Electron energy was set at 70 eV. MS source

and quadrupole were maintained at 230°C and 150°C, respectively. Parameters specific to each

analysis are listed below.

For analysis of sugar core alditol acetate derivatives, we followed previously published

GC-MS parameters (Sassaki et al., 2008). Inlet temperature was maintained at 275°C. GC oven

temperature was held at 60°C for one min and then was ramped at a rate of 40°C/min to 180°C.

Oven temperature was then ramped to 240°C at a rate of 5°C/min and held for three min. Total

run time was 19 min. The MS detector transfer line was maintained at 280°C. Split ratio was

10:1 with a split flow rate of 10 mL/min. A three min solvent delay was used. MS data was

collected in full scan mode, m/z 50-600. Analysis parameters are also described in Table S34.

For analysis of MPTA fatty acid ethyl ester derivatives, we followed previously

published GC-MS parameters (Jenske and Vetter, 2007). GC oven temperature was held at 60°C

for 1.5 min and then was ramped at a rate of 40°C/min to 180°C. 180°C was held for two min

after which the oven temperature was ramped to 230°C at a rate of 2°C/min. After 230°C was

held for nine min, the oven temperature was ramped to 300°C at 10°C/min and held for 7.5 min.

Total run time was 55 min. A 5-min solvent delay was applied. The MS detector transfer line

was maintained at 280°C. Selective ion monitoring detected the ions *m/z* 189, 209, and 255 with

a dwell time of 50 ms. Analysis parameters are also described in Table S35.

<u>Sample collection and transcriptome sequencing</u>

We sequenced 18 transcriptomes: six biological replicates each of trichomes isolated

from hypocotyls (trichomes), hypocotyls stripped of their trichomes (stripped hypocotyls), and

whole roots (Supplemental data file 1). We sampled trichomes and stripped hypocotyls from 7-

day-old *S. melongena* 67/3 seedlings following methods developed for root hair cell isolation

(Bucher et al., 1997) with modifications. We grew lawns of eggplant seedlings in soil flats as

described above for bulk leaf surface metabolite extraction. At 7 days post germination, we

removed roots and cotyledons from seedlings, then transferred hypocotyls to liquid nitrogen in a

plastic 2 L Dewar flask. Frozen hypocotyls were gently stirred with a glass rod for 20 minutes to

physically shear trichomes from hypocotyls. After confirming under a dissecting scope that a

sample of three hypocotyls had been stripped bare, we filtered trichomes into a 2 L glass beaker

by slowly pouring the contents of the Dewar flask through a 500 μm wire mesh sieve

(MilliporeSigma, Burlington, MA, USA). To maximize trichome recovery, stripped hypocotyls

were returned to the Dewar, rinsed with liquid nitrogen, and filtered through the 500 μm sieve

six more times. Stripped hypocotyls were divided into six pre-weighed 50 mL conical tubes,

quickly weighed, then transferred to -80ºC for storage. Trichomes were subsequently filtered

through a 150 μm sieve (MilliporeSigma, Burlington, MA, USA) into a 500 mL beaker to

increase sample purity then transferred to a 50 mL conical tube to allow the excess liquid

nitrogen to evaporate. Finally, filtered trichomes were divided into six pre-weighed 2 mL screw-

cap tubes (Dot Scientific, Inc., Burton, MI, USA), quickly weighed, then transferred to -80ºC for

storage.

We extracted total RNA using the RNeasy Plant Mini Kit (Qiagen, Hilden, Germany)

following the manufacturer's directions, measured RNA concentration using a Qubit 1.0

instrument (Thermo Fisher Scientific, Waltham, MA, USA) with the RNA HS assay, and the

samples were processed by the Michigan State University Research Technology Support Facility

Genomics Core (East Lansing, MI, USA) for library preparation and high-throughput

sequencing. The core checked RNA quality using a Bioanalyzer 2100 (Agilent Technologies,

Santa Clara, CA, USA), constructed sequencing libraries using an Illumina Stranded mRNA

Prep kit (Illumina, Inc. San Diego, CA, USA), and sequenced all 18 libraries on a NovaSeq 6000

(Illumina, Inc. San Diego, CA) using a single S4 flow cell lane, producing 150-base pair paired-

end reads (Supplemental datafile 1). The raw fastq files are available to download from the

National Center for Biotechnology Information Sequence Read Archive database under

BioProject PRJNA935765.

<u>Transcriptome alignment and differential gene expression analysis</u>

To prepare the 150-bp paired-end sequences for alignment, we employed Trimmomatic

(Bolger et al., 2014) to trim adapters and low-quality bases, then filter reads shorter than 75 bp,

removing an average of 3.7% of sequences (range: 3.3% - 4.2%). We mapped the resulting 75–

150-bp (average length: 149 bp) paired-end RNAseq reads separately to the eggplant V4.1 and

HQ reference genomes using STAR in two-pass mode, which enhances splice junction discovery

and mapping sensitivity (Dobin et al., 2013; Dobin and Gingeras, 2015). Using this approach, an

average of 79.2% (range: 60.1 - 87.1%) and 82.2% (range: 62.0 - 90.7%) of reads mapped to a

unique genomic location in V4.1 and HQ, respectively (Table S36). We filtered the resulting

transcriptome alignments according to best practices as defined by the Genome Analysis Toolkit

(GATK) (DePristo et al., 2011; Van der Auwera et al., 2013). Briefly, we removed optical and

PCR duplicates with MarkDuplicates from the Picard toolkit

(http://broadinstitute.github.io/picard), parsed reads into exon segments and removed intron-

spanning bases using SplitNCigarReads from GATK (McKenna et al., 2010). Finally, we

selected unique alignments by eliminating reads with a mapping quality score below Q60 with

the view command from SAMtools (Li et al., 2009). To generate raw read counts, we used the

HTSeq command, htseq-count with the nonunique parameter set to all (Anders et al., 2015).

Raw read counts generated by HTSeq-count were used to perform differential gene

expression analysis in edgeR (Robinson et al., 2010). To restrict comparisons to expressed genes,

only transcripts with at least one read count-per-million (CPM) in at least one sample were

retained for further analysis. This filtering step removed 11,665 (33.4%) and 14,544 (39.8%)

annotated genes in the V4.1 and HQ eggplant genomes, respectively, yielding 23,251 and 22,024

expressed transcripts for differential gene expression analysis. Next, we normalized transcript

abundances across the 18 transcriptomes in our eggplant V4.1 and HQ alignments using the

default trimmed mean of M-values (TMM) method in the calcNormFactors function, then

performed multidimensional scaling (MDS) with the plotMDS function to compare global gene

expression profiles (Figure S15). This showed that our samples cluster tightly by tissue (i.e.,

trichome, shaved hypocotyl, or root), with no obvious differences between V4.1 and HQ

alignments. To test differences in gene expression across tissues, we implemented a generalized

linear model (GLM) using a quasi-likelihood (QL) approach: we generated an experimental

design matrix specifying the three tissues (i.e., trichomes, trichomeless hypocotyls, and roots)

with the model.matrix function, then used the glmQLFit function to fit our data to a QL-GLM

model. To identify genes with a log2 fold-change (FC) > 2 between tissues, we used the

glmTreat function, which performs threshold hypothesis testing, a rigorous statistical approach

that evaluates variance and magnitude to detect expression differences greater than the specified

value (e.g., log2 FC > 2), then applies false discovery rate (FDR) p-value corrections. Genes

were classified as significantly differentially expressed between two tissues if log2 FC > 2 and

FDR-corrected p-value < 0.05. We calculated absolute transcript abundance for all expressed

genes as transcripts per million (TPM) with the calculateTPM function in scateR (McCarthy et

al., 2017). Transcriptome expression values are in Tables S37 and S38.

1016

<u>Phylogenetic analyses</u>

Characterized ASAT enzymes fall into clade III of the BAHD family (Fan et al., 2017,

2015; Leong et al., 2022, 2020; Lou et al., 2021; Moghe et al., 2017; Nadakuduti et al., 2017;

Schenck et al., 2022; Schilmiller et al., 2015, 2012). To identify ASAT candidates in eggplant,

we searched for sequences containing the Pfam Transferase domain (PF02458) (Finn et al.,

2010), associated with all characterized catalytically active BAHD proteins. The PF02458 HMM

profile was obtained from the Pfam website (http://pfam.sanger.ac.uk/) and queried against the

V4.1 and HQ eggplant proteomes using the hmmsearch tool from HMMER v3.2.1 (hmmer.org),

revealing 106 and 108 putative BAHD sequences, respectively. Using MAFFT v7.471 in E-INS-

i mode, we built multiple sequence alignments (MSA) of amino acid sequences from four

sources: 1) V4.1 or HQ eggplant PF02458 hits, 2) published reference sequences for clades I-VII

(Moghe et al., 2023), 3) characterized ASAT sequences from other Solanaceae species, and 4)

the SaASAT3 and SaASAT3-L1 candidates from *S. aethiopicum* (Katoh and Standley, 2013).

The E-INS-i algorithm implements local alignment with a generalized affine gap cost (Altschul,

1998), which aligns conserved regions (e.g., the BAHD transferase domain) and essentially

ignores nonconserved regions. Phylogenetic reconstruction was performed using IQ-TREE

v2.1.3 (Minh et al., 2020). The ModelFinder tool was implemented to identify the best maximum

likelihood model for estimating evolutionary relationships (Kalyaanamoorthy et al., 2017),

leading to selection of Jones-Taylor-Thornton (JTT)+F with seven rate categories and Q.plant+F

with seven rate categories for the V4.1 and HQ MSAs, respectively. Phylogenetic trees were

inferred by maximum likelihood using the chosen model, and branch support was obtained from

100,000 ultrafast bootstrap iterations (Hoang et al., 2018). The resulting BAHD phylogenies

were visualized using the ggtree package in R (Yu et al., 2017). To generate the clade III BAHD

heatmap-tree, we used the viewClade function in ggtree to subset the phylogeny and used

gheatmap to visualize transcript abundance (log2 TPM) for the eggplant BAHDs.

1042

<u>BAHD acyltransferase cloning, expression, and purification</u>

Candidate ASATs were cloned into pET28b(+) (MilliporeSigma, Burlington, MA, USA).

Open reading frames from genes were either synthesized by Twist Biosciences (South San

Francisco, CA, USA) with or without codon-optimization for *E. coli* expression (Table S39) or

amplified from genomic DNA or cDNA with primers listed in Table S40. Q5 2X Hotstart master

mix (New England Biolabs, Ipswich, MA, USA) was used for cloning PCRs. The source of

cloned DNA and whether a gene was codon optimized is described in Table S39. Amplified

genes were purified by agarose gel electrophoresis and extraction with the Monarch DNA Gel

Extraction Kit (New England Biolabs, Ipswich, MA, USA). Both the synthesized genes and PCR

amplified genes were then inserted into a doubly digested BamHI/XhoI pET28b(+) through

Gibson assembly using the 2X Gibson Assembly Master Mix (New England Biolabs, Ipswich,

MA, USA) according to the manufacturer's instructions. The constructs using synthesized genes

were transformed into BL21(DE3) (MilliporeSigma, Burlington, MA, USA) and constructs with

PCR amplified genes were transformed into BL21 Rosetta(DE3) cells (MilliporeSigma,

Burlington, MA, USA). Constructs were verified with colony PCR and Sanger sequencing using

T7 terminator and promoter primers (Table S40). Sanger sequencing was completed by the

Michigan State University Research Technology Support Facility Genomics Core (East Lansing,

MI, USA).

Protein expression occurred as previously described (Leong et al., 2022; Lou et al.,

2021). Briefly, 50 mL cultures of picked transformation colonies were grown overnight at 37°C,

shaking at 225 rpm in Luria-Bertani (LB) media (Neogen, Lansing, MI, USA) supplemented

with 1% glucose (w/v). Fifteen mL of the overnight cultures were inoculated into 1 L of fresh LB

medium, which was incubated at 37°C shaking at 225 rpm until an OD600 of 0.5 was reached.

The cultures were incubated on ice for 25 min, after which, isopropylthio-β-galactoside was

added to a final concentration of 50-500 μM. Then, cultures were incubated at 16°C shaking at

180 rpm overnight for 16 hours before cells were harvested by centrifugation at 4,000 rpm for 10

minutes at 4°C.

S. melongena BAHDs were purified as previously described (Leong et al., 2020) with the

following modifications. The extraction buffer contained 10 mM imidazole, the wash buffer

contained 20 mM imidazole, and the elution buffer contained 500 mM imidazole. Protein eluent

was concentrated with 30-kD Amicon Ultra centrifugal filter units (MilliporeSigma).

1074

Enzyme assays

Enzyme assays were conducted in 100 mM sodium phosphate buffer at pH 6. For

forward assays, acetyl-CoA (MilliporeSigma, Burlington, MA, USA) was added to a final

concentration of 0.1 mM, and for reverse assays free CoA (MilliporeSigma, Burlington, MA,

USA) was added to a final concentration of 1 mM. Purified AI3:16 and AI4:18 substrates were

dried down using a vacuum centrifuge and redissolved in ethanol:water:formic acid (1:1:0.001).

One microliter of the prepared acylsugars were used as acyl acceptors. Six microliters of enzyme

were added to a final volume of 60 µL. For negative controls, 6 µL of enzyme that was heat

inactivated at 95°C for 10 minutes was substituted in place of untreated enzyme. Assays were

incubated at 30°C for 30 minutes after which 120 µL of acetonitrile:isopropanol:formic acid

(1:1:0.001) with 1.5 µM telmisartan (MilliporeSigma, Burlington, MA, USA) stop solution was

added. Reactions were then spun at 17,000 x *g* for 10 minutes to remove precipitate. Supernatant

was placed in autosampler vials and analyzed by LC-MS.

1088

RT-PCR of *S. aethiopicum* BAHDs

We employed semi-quantitative RT-PCR to test *S. aethiopicum* BAHD expression in

glandular trichomes with cDNA (Figure S14) and genomic DNA (gDNA) (Figure S16). Total

RNA was isolated with the RNeasy Plant Mini Kit (Qiagen, Hilden, Germany), including an on-

column DNase digestion (Qiagen, Hilden, Germany), from shaved hypocotyls from accession *S.*

aethiopicum PI 666075 and glandular trichomes from accessions PI 666075 and Grif 14165

isolated as described above for *S. melongena* glandular trichomes. RNA was quantified with a

Nanodrop 2000c instrument (Thermo Fisher Scientific, Waltham, MA, USA). We synthesized

cDNA using 10 ng of RNA and SuperScript III Reverse Transcriptase (Invitrogen, Waltham,

MA, USA). gDNA was isolated with the DNeasy Plant Mini Kit (Qiagen, Hilden, Germany)

from young leaf tissue collected from mature *S. aethiopicum* PI 666075 plants. PCR reactions

(25 µL) were set up with GoTaq Green Master Mix (Promega, Madison, WI, USA), 200 nM of

forward and reverse primers (Table S40), and 1 µL of cDNA or gDNA. PCR was performed

under these conditions: 2 min at 95°C followed by 22, 30, and 35 cycles of 30s at 95°C, 30s at

58°C, and 1 min at 72°C.

We identified the putative *S. aethiopicum* orthologs of the *S. melongena* candidate

BAHDs by querying *S. aethiopicum* annotated transcripts (Song et al., 2019) with *S. melongena*

candidate ASAT DNA sequences using BLASTn. SaASAT3 and SaASAT3-L1 (GenBank

accession: OQ547782) were not annotated in the *S. aethiopicum* genome and were identified by

querying *S. aethiopicum* scaffolds with BLASTn and then identifying open reading frames with

Geneious software v9.1.8 (Dotmatics, Boston, MA, USA). The *S. aethiopicum* ef1α gene was

identified by querying a putative *S. melongena* ef1α, SMEL4.1_06g005890, against the *S.*

aethiopicum scaffolds.

1112

S. melongena acyl chain composition analysis

S. melongena acyl chain composition was assessed with a previously developed

esterification and GC-MS methods (Fan et al., 2020; Ning et al., 2015; Schenck et al., 2022).

Acyl chain identifies were determined through authentic reference standards for nC8, nC10,

nC12, and nC14 fatty acid ethyl esters (MilliporeSigma). The QuanLynx function of MassLynx

v4.1 (Waters Corporation) integrated peaks from extracted ion chromatograms. Extracted ion

chromatograms for the medium acyl chains iC8, nC8, iC10, nC10, iC12, nC12, iC14, and nC14

were generated for the *m/z* of 88 with a mass window of *m/z* of 0.50. Extracted ion

chromatograms for the short chains iC4, aiC5, and iC5 were generated for with the *m/z* values of

71, 102, and 101, respectively, with a mass window of *m/z* of 0.50.

1123

1124   <u>Accession numbers</u>

1125   Raw RNA sequence data can be found in the National Center for Biotechnology

1126   Information (NCBI) Sequence Read Archive database under BioProject PRJNA935765.

1127   Sequence data for SaASAT3-L1 is available from the NCBI GenBank database under accession

1128   number OQ547782.

1129

Acknowledgements

We thank Dr. Christopher T. Martine for providing Australian *Solanum* ssp. seeds, Dr.

Joyce van Eck for providing *Solanum prinophyllum* seeds, Dr. Guiseppe Leonardo Rotino and

the Consiglio per la ricercar in agricoltura e l'analisi dell'economia agraria for providing *S.*

melongena 67/3 and 305E40 seeds, and USDA-GRIN for providing *Solanum* ssp. seeds. We

acknowledge the Michigan State University RTSF Mass Spectrometry and Metabolomics Core

Facilities for LC-MS analysis support. We also acknowledge Dr. Daniel Holmes and Dr. Li Xie

at the Michigan State University Max T. Rogers NMR Facility for experimental design and data

analysis support. We are grateful to current and past members of the Last lab for feedback on the

research and especially Drs. Jaynee Hart and Xingxing Li for comments on the manuscript.

1140

Funding

This research was funded by the National Science Foundation grant IOS-PGRP-1546617

(to R. L. L. and A. D. J.), National Science Foundation grant IOS-PGRP-2218206 (to R. L. L.

and R. E. K.) and National Institute of General Medical Sciences of the National Institutes of

Health graduate training grant no. T32-GM110523 (to P. D. F.). A. D. J. is supported in part by

USDA National Institute of Food and Agriculture Hatch project number MICL02474.

1147

References

Agati, G., Tattini, M., 2010. Multiple functional roles of flavonoids in photoprotection. New
Phytol. 186, 786–793.

Altschul, S.F., 1998. Generalized affine gap costs for protein sequence alignment. Proteins
Struct. Funct. Bioinforma. 32, 88–96. https://doi.org/10.1002/(SICI)1097-
0134(19980701)32:1<88::AID-PROT10>3.0.CO;2-J

Anders, S., Pyl, P.T., Huber, W., 2015. HTSeq—a Python framework to work with high-
throughput sequencing data. Bioinformatics 31, 166–169.
https://doi.org/10.1093/bioinformatics/btu638

Asai, T., Fujimoto, Y., 2010. Cyclic fatty acyl glycosides in the glandular trichome exudate of
Silene gallica. Phytochemistry 71, 1410–1417. https://doi.org/10.1016/j.phytochem.2010.05.008

Asai, T., Hara, N., Fujimoto, Y., 2010. Fatty acid derivatives and dammarane triterpenes from
the glandular trichome exudates of *Ibicella lutea* and *Proboscidea louisiana*. Phytochemistry 71,
877–894. https://doi.org/10.1016/j.phytochem.2010.02.013

Aubriot, X., Knapp, S., Syfert, M.M., Poczai, P., Buerki, S., 2018. Shedding new light on the
origin and spread of the brinjal eggplant (*Solanum melongena* L.) and its wild relatives. Am. J.
Bot. 105, 1175–1187. https://doi.org/10.1002/ajb2.1133

Augustin, J.M., Kuzina, V., Andersen, S.B., Bak, S., 2011. Molecular activities, biosynthesis and
evolution of triterpenoid saponins. Phytochemistry 72, 435–457.
https://doi.org/10.1016/j.phytochem.2011.01.015

Bah, M., Pereda-Miranda, R., 1996. Detailed FAB-mass spectrometry and high resolution NMR
investigations of tricolorins A-E, individual oligosaccharides from the resins of *Ipomoea tricolor*
(Convolvulaceae). Tetrahedron 52, 13063–13080. https://doi.org/10.1016/0040-4020(96)00789-
2

Barchi, L., Rabanus-Wallace, M.T., Prohens, J., Toppino, L., Padmarasu, S., Portis, E., Rotino,
G.L., Stein, N., Lanteri, S., Giuliano, G., 2021. Improved genome assembly and pan-genome
provide key insights into eggplant domestication and breeding. Plant J. 107, 579–596.
https://doi.org/10.1111/tpj.15313

Baumann, E., Stoya, G., Völkner, A., Richter, W., Lemke, C., Linss, W., 2000. Hemolysis of
human erythrocytes with saponin affects the membrane structure. Acta Histochem. 102, 21–35.
https://doi.org/10.1078/0065-1281-00534

Bernal, C.-A., Castellanos, L., Aragón, D.M., Martínez-Matamoros, D., Jiménez, C., Baena, Y.,
Ramos, F.A., 2018. Peruvioses A to F, sucrose esters from the exudate of *Physalis peruviana*
fruit as α-amylase inhibitors. Carbohydr. Res. 461, 4–10.
https://doi.org/10.1016/j.carres.2018.03.003

Bohs, L., 2004. Major clades in Solanum based on ndhF sequence data. Monogr. Syst. Bot. 27–49.

Bohs, L., Olmstead, R.G., 1997. Phylogenetic relationships in Solanum (Solanaceae) based on ndhF sequences. Syst. Bot. 22, 5–17. https://doi.org/10.2307/2419674

Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30, 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Cao, C.-M., Wu, X., Kindscher, K., Xu, L., Timmermann, B.N., 2015. Withanolides and sucrose esters from *Physalis neomexicana*. J. Nat. Prod. 78, 2488–2493. https://doi.org/10.1021/acs.jnatprod.5b00698

Caprioli, G., Iannarelli, R., Cianfaglione, K., Fiorini, D., Giuliani, C., Lucarini, D., Papa, F., Sagratini, G., Vittori, S., Maggi, F., 2016. Volatile profile, nutritional value and secretory structures of the berry-like fruits of *Hypericum androsaemum* L. Food Res. Int. 79, 1–10. https://doi.org/10.1016/j.foodres.2015.11.021

Castillo, M., Connolly, J.D., Ifeadike, P., Labbé, C., Rycroft, D.S., Woods, N., 1989. Partially acylated glucose and sucrose derivatives from *Salpiglossis sinuata* (Solanaceae). J. Chem. Res. Synop. 398–399.

Chortyk, O.T., Kays, S.J., Teng, Q., 1997. Characterization of insecticidal sugar esters of *Petunia*. J. Agric. Food Chem. 45, 270–275. https://doi.org/10.1021/jf960322f

Cicchetti, E., Duroure, L., Le Borgne, E., Laville, R., 2018. Upregulation of skin-aging biomarkers in aged NHDF cells by a sucrose ester extract from the agroindustrial waste of *Physalis peruviana* calyces. J. Nat. Prod. 81, 1946–1955. https://doi.org/10.1021/acs.jnatprod.7b01069

D'Auria, J.C., 2006. Acyltransferases in plants: a good time to be BAHD. Curr. Opin. Plant Biol. 9, 331–340. https://doi.org/10.1016/j.pbi.2006.03.016

De Moraes, C.M., Mescher, M.C., Tumlinson, J.H., 2001. Caterpillar-induced nocturnal plant volatiles repel conspecific females. Nature 410, 577–580. https://doi.org/10.1038/35069058

DePristo, M.A., Banks, E., Poplin, R., Garimella, K.V., Maguire, J.R., Hartl, C., Philippakis, A.A., del Angel, G., Rivas, M.A., Hanna, M., McKenna, A., Fennell, T.J., Kernytsky, A.M., Sivachenko, A.Y., Cibulskis, K., Gabriel, S.B., Altshuler, D., Daly, M.J., 2011. A framework for variation discovery and genotyping using next-generation DNA sequencing data. Nat. Genet. 43, 491–498. https://doi.org/10.1038/ng.806

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., Gingeras, T.R., 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29, 15–21. https://doi.org/10.1093/bioinformatics/bts635

Dobin, A., Gingeras, T.R., 2015. Mapping RNA-seq reads with STAR. Curr. Protoc. Bioinforma. 51, 11.14.1-11.14.19. https://doi.org/10.1002/0471250953.bi1114s51

Fan, P., Leong, B.J., Last, R.L., 2019. Tip of the trichome: evolution of acylsugar metabolic diversity in Solanaceae. Curr. Opin. Plant Biol., Physiology and metabolism 49, 8–16. https://doi.org/10.1016/j.pbi.2019.03.005

Fan, P., Miller, A.M., Liu, X., Jones, A.D., Last, R.L., 2017. Evolution of a flipped pathway creates metabolic innovation in tomato trichomes through BAHD enzyme promiscuity. Nat. Commun. 8, 2080. https://doi.org/10.1038/s41467-017-02045-7

Fan, P., Miller, A.M., Schilmiller, A.L., Liu, X., Ofner, I., Jones, A.D., Zamir, D., Last, R.L., 2015. In vitro reconstruction and analysis of evolutionary variation of the tomato acylsucrose metabolic network. Proc. Natl. Acad. Sci. U.S.A https://doi.org/10.1073/pnas.1517930113

Fan, P., Wang, P., Lou, Y.R., Leong, B.J., Moore, B.M., Schenck, C.A., Combs, R., Cao, P., Brandizzi, F., Shiu, S.H., Last, R.L., 2020. Evolution of a plant gene cluster in solanaceae and emergence of metabolic diversity. eLife 9, 1–26. https://doi.org/10.7554/eLife.56717

Fiesel, P.D., Parks, H.M., Last, R.L., Barry, C.S., 2022. Fruity, sticky, stinky, spicy, bitter, addictive, and deadly: evolutionary signatures of metabolic complexity in the Solanaceae. Nat. Prod. Rep. 39, 1438–1464. https://doi.org/10.1039/D2NP00003B

Fobes, J.F., Mudd, J.B., Marsden, M.P.F., 1985. Epicuticular lipid accumulation on the leaves of *Lycopersicon pennellii* (Corr.) D'Arcy and Lycopersicon esculentum Mill. Plant Physiol. 77, 567–570. https://doi.org/10.1104/pp.77.3.567

Gagnon, E., Hilgenhof, R., Orejuela, A., McDonnell, A., Sablok, G., Aubriot, X., Giacomin, L., Gouvêa, Y., Bragionis, T., Stehmann, J.R., Bohs, L., Dodsworth, S., Martine, C., Poczai, P., Knapp, S., Särkinen, T., 2022. Phylogenomic discordance suggests polytomies along the backbone of the large genus Solanum. Am. J. Bot. 109, 580–601. https://doi.org/10.1002/ajb2.1827

Ghosh, B., Westbrook, T.C., Jones, A.D., 2014a. Comparative structural profiling of trichome specialized metabolites in tomato (*Solanum lycopersicum*) and *S. habrochaites*: acylsugar profiles revealed by UHPLC/MS and NMR. Metabolomics 10, 496–507. https://doi.org/10.1007/s11306-013-0585-y

Goffreda, J.C., Mutschler, M.A., Avé, D.A., Tingey, W.M., Steffens, J.C., 1989. Aphid deterrence by glucose esters in glandular trichome exudate of the wild tomato, *Lycopersicon pennellii*. J. Chem. Ecol. 15, 2135–2147. https://doi.org/10.1007/BF01207444

Herrera-Salgado, Y., Garduño-Ramírez, M.L., Vázquez, L., Rios, M.Y., Alvarez, L., 2005. Myo-inositol-derived glycolipids with anti-inflammatory activity from *Solanum lanceolatum*. J. Nat. Prod. 68, 1031–1036. https://doi.org/10.1021/np050054s

Hoang, D.T., Chernomor, O., von Haeseler, A., Minh, B.Q., Vinh, L.S., 2018. UFBoot2: improving the ultrafast bootstrap approximation. Mol. Biol. Evol. 35, 518–522. https://doi.org/10.1093/molbev/msx281

Hurney, S.M., 2018. Strategies for profiling and discovery of acylsugar specialized metabolites (Ph.D.). Michigan State University, United States -- Michigan.

Jenske, R., Vetter, W., 2007. Highly selective and sensitive gas chromatography–electron-capture negative-ion mass spectrometry method for the indirect enantioselective identification of 2- and 3-hydroxy fatty acids in food and biological samples. J. Chromatogr. A 1146, 225–231. https://doi.org/10.1016/j.chroma.2007.01.102

Ji, W., Mandal, S., Rezenom, Y.H., McKnight, T.D., 2022. Specialized metabolism by trichome-enriched Rubisco and fatty acid synthase components. Plant Physiol. kiac487. https://doi.org/10.1093/plphys/kiac487

Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., von Haeseler, A., Jermiin, L.S., 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. Nat. Methods 14, 587–589. https://doi.org/10.1038/nmeth.4285

Katoh, K., Standley, D.M., 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. Mol. Biol. Evol. 30, 772–780. https://doi.org/10.1093/molbev/mst010

Kim, J., Kang, K., Gonzales-Vigil, E., Shi, F., Daniel Jones, A., Barry, C.S., Last, R.L., 2012. Striking natural diversity in glandular trichome acylsugar composition is shaped by variation at the acyltransferase2 locus in the wild tomato *Solanum habrochaites*. Plant Physiol. 160, 1854–1870. https://doi.org/10.1104/pp.112.204735

King, R.R., Calhoun, L.A., 1988. 6 2,3-Di-O- and 1,2,3-tri-O-acylated glucose esters from the glandular trichomes of *Datura metel*. Phytochemistry 27, 3761–3763. https://doi.org/10.1016/0031-9422(88)83013-9

King, R.R., Pelletier, Y., Singh, R.P., Calhoun, L.A., 1986. 3,4-Di-O-isobutyryl-6-O-caprylsucrose: the major component of a novel sucrose ester complex from the type B glandular trichomes of *Solanum berthaultii* Hawkes (Pl 473340). J. Chem. Soc. Chem. Commun. 1078–1079. https://doi.org/10.1039/C39860001078

Koonin, E.V., 2005. Orthologs, paralogs, and evolutionary genomics. Annu. Rev. Genet. 39, 309–338. https://doi.org/10.1146/annurev.genet.39.073003.114725

Korenblum, E., Dong, Y., Szymanski, J., Panda, S., Jozwiak, A., Massalha, H., Meir, S., Rogachev, I., Aharoni, A., 2020. Rhizosphere microbiome mediates systemic root metabolite exudation by root-to-root signaling. Proc. Natl. Acad. Sci. U. S. A. 117, 3874–3883. https://doi.org/10.1073/pnas.1912130117

Kretschmar, J.A., Baumann, T.W., 1999. Caffeine in Citrus flowers. Phytochemistry 52, 19–23. https://doi.org/10.1016/S0031-9422(99)00119-3

Kruse, L.H., Bennett, A.A., Mahood, E.H., Lazarus, E., Park, S.J., Schroeder, F., Moghe, G.D., 2022. Illuminating the lineage-specific diversification of resin glycoside acylsugars in the

morning glory (Convolvulaceae) family using computational metabolomics. Hortic. Res. 9, uhab079. https://doi.org/10.1093/hr/uhab079

Landry, L.G., Chapple, CCS., Last, R.L., 1995. Arabidopsis mutants lacking phenolic sunscreens exhibit enhanced ultraviolet-B injury and oxidative damage. Plant Physiol. 109, 1159–1166. https://doi.org/10.1104/pp.109.4.1159

Leckie, B.M., D'Ambrosio, D.A., Chappell, T.M., Halitschke, R., De Jong, D.M., Kessler, A., Kennedy, G.G., Mutschler, M.A., 2016. Differential and synergistic functionality of acylsugars in suppressing oviposition by insect herbivores. PLoS ONE. https://doi.org/10.1371/journal.pone.0153345

Leong, B.J., Hurney, S., Fiesel, P., Anthony, T.M., Moghe, G., Jones, A.D., Last, R.L., 2022. Identification of BAHD acyltransferases associated with acylinositol biosynthesis in *Solanum quitoense* (naranjilla). Plant Direct 6, e415. https://doi.org/10.1002/pld3.415

Leong, B.J., Hurney, S.M., Fiesel, P.D., Moghe, G.D., Jones, A.D., Last, R.L., 2020. Specialized metabolism in a nonmodel nightshade: trichome acylinositol biosynthesis. Plant Physiol. 183, 915–924. https://doi.org/10.1104/pp.20.00276

Leong, B.J., Lybrand, D.B., Lou, Y.R., Fan, P., Schilmiller, A.L., Last, R.L., 2019. Evolution of metabolic novelty: a trichome-expressed invertase creates specialized metabolic diversity in wild tomato. Sci. Adv. 5, 1–14. https://doi.org/10.1126/sciadv.aaw3754

Levin, D.A., 1973. The role of trichomes in plant defense. Q. Rev. Biol. 48, 3–15.

Levin, R.A., Myers, N.R., Bohs, L., 2006. Phylogenetic relationships among the "spiny solanums" (Solanum subgenus Leptostemonum, Solanaceae). Am. J. Bot. 93, 157–169. https://doi.org/10.3732/ajb.93.1.157

Li, D., Qian, J., Li, Weiliu, Yu, N., Gan, G., Jiang, Y., Li, Wenjia, Liang, X., Chen, R., Mo, Y., Lian, J., Niu, Y., Wang, Y., 2021. A high-quality genome assembly of the eggplant provides insights into the molecular basis of disease resistance and chlorogenic acid synthesis. Mol. Ecol. Resour. 21, 1274–1286. https://doi.org/10.1111/1755-0998.13321

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., 1000 Genome Project Data Processing Subgroup, 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics 25, 2078–2079. https://doi.org/10.1093/bioinformatics/btp352

Liu, X., Enright, M., Barry, C.S., Jones, A.D., 2017. Profiling, isolation and structure elucidation of specialized acylsucrose metabolites accumulating in trichomes of Petunia species. Metabolomics 13, 85. https://doi.org/10.1007/s11306-017-1224-9

Lou, Y.-R., Anthony, T.M., Fiesel, P.D., Arking, R.E., Christensen, E.M., Jones, A.D., Last, R.L., 2021. It happened again: convergent evolution of acylglucose specialized metabolism in black nightshade and wild tomato. Sci. Adv. 7, eabj8726. https://doi.org/10.1126/sciadv.abj8726

Lou, Y.-R., Leong, B., 2019. Leaf surface acylsugar extraction and LC-MS profiling - v1.0.

Luu, V.T., Weinhold, A., Ullah, C., Dressel, S., Schoettner, M., Gase, K., Gaquerel, E., Xu, S., Baldwin, I.T., 2017. O-acyl sugars protect a wild tobacco from both native fungal pathogens and a specialist herbivore. Plant Physiol. 174, 370–386. https://doi.org/10.1104/pp.16.01904

Lybrand, D.B., Anthony, T.M., Jones, A.D., Last, R.L., 2020. An integrated analytical approach reveals trichome acylsugar metabolite diversity in the wild tomato *Solanum pennellii*. Metabolites 10, 1–25. https://doi.org/10.3390/metabo10100401

Maldonado, E., Torres, F.R., Martínez, M., Pérez-Castorena, A.L., 2006. Sucrose esters from the fruits of *Physalis nicandroides* var. attenuata. J. Nat. Prod. 69, 1511–1513. https://doi.org/10.1021/np060274l

Matsuzaki, T., Shinozaki, Y., Suhara, S., Ninomiya, M., Shigematsu, H., Koiwai, A., 1989. Isolation of glycolipids from the surface lipids of *Nicotiana bigelovii* and their distribution in *Nicotiana* species. Agric. Biol. Chem. 53, 3079–3082. https://doi.org/10.1271/bbb1961.53.3079

McCarthy, D.J., Campbell, K.R., Lun, A.T.L., Wills, Q.F., 2017. Scater: pre-processing, quality control, normalization and visualization of single-cell RNA-seq data in R. Bioinformatics 33, 1179–1186. https://doi.org/10.1093/bioinformatics/btw777

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K., Altshuler, D., Gabriel, S., Daly, M., DePristo, M.A., 2010. The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. Genome Res. 20, 1297–1303. https://doi.org/10.1101/gr.107524.110

Mennella, G., Rotino, G.L., Fibiani, M., D'Alessandro, A., Francese, G., Toppino, L., Cavallanti, F., Acciarri, N., Lo Scalzo, R., 2010. Characterization of health-related compounds in eggplant (*Solanum melongena* L.) lines derived from introgression of allied species. J. Agric. Food Chem. 58, 7597–7603. https://doi.org/10.1021/jf101004z

Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A., Lanfear, R., 2020. IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. Mol. Biol. Evol. 37, 1530–1534. https://doi.org/10.1093/molbev/msaa015

Moghe, G., Kruse, L.H., Petersen, M., Scossa, F., Fernie, A.R., Gaquerel, E., D'Auria, J.C., 2023. BAHD company: the ever-expanding roles of the BAHD acyltransferase gene family in plants. Annu. Rev. Plant Biol. 74, annurev-arplant-062922-050122. https://doi.org/10.1146/annurev-arplant-062922-050122

Moghe, G.D., Leong, B.J., Hurney, S.M., Jones, A.D., Last, R.L., 2017. Evolutionary routes to biochemical innovation revealed by integrative analysis of a plant-defense related specialized metabolic pathway. eLife 6, 1–33. https://doi.org/10.7554/eLife.28468

Nadakuduti, S.S., Uebler, J.B., Liu, X., Jones, A.D., Barry, C.S., 2017. Characterization of trichome-expressed BAHD acyltransferases in *Petunia axillaris* reveals distinct acylsugar assembly mechanisms within the Solanaceae. Plant Physiol. 175, 36–50. https://doi.org/10.1104/pp.17.00538

Nee, M., 2022a. *Solanum acerifolium* [Dunal] [WWW Document]. URL
https://solanaceaesource.myspecies.info/taxonomy/term/105367/descriptions (accessed 10.3.22).

Nee, M., 2022b. *Solanum atropurpureum* [Schrank] [WWW Document]. URL
https://solanaceaesource.myspecies.info/taxonomy/term/105749/descriptions (accessed 10.3.22).

Nee, M., 1991. Synopsis of Solanum section Acanthophora: a group of interest for
glycoalkaloids, in: Solanaceae III, taxonomy, chemistry, evolution. Royal Botanic Gardens for
the Linnean Society of London, Kew, Richmond, Surrey, UK, pp. 257–266.

Ning, J., Moghe, G.D., Leong, B., Kim, J., Ofner, I., Wang, Z., Adams, C., Jones, A.D., Zamir,
D., Last, R.L., 2015. A feedback-insensitive isopropylmalate synthase affects acylsugar
composition in cultivated and wild tomato. Plant Physiol. 169, 1821–1835.
https://doi.org/10.1104/pp.15.00474

PBI Solanum Project, 2022. Solanaceae Source [WWW Document]. URL
https://solanaceaesource.org (accessed 10.3.22).

Pereda-Miranda, R., Mata, R., Anaya, A.L., Wickramaratne, D.B.M., Pezzuto, J.M., Kinghorn,
A.D., 1993. Tricolorin A, major phytogrowth inhibitor from *Ipomoea tricolor*. J. Nat. Prod. 56,
571–582. https://doi.org/10.1021/np50094a018

Pinot, F., Beisson, F., 2011. Cytochrome P450 metabolizing fatty acids in plants:
characterization and physiological roles. FEBS J. 278, 195–205. https://doi.org/10.1111/j.1742-
4658.2010.07948.x

Robinson, M.D., McCarthy, D.J., Smyth, G.K., 2010. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics 26, 139–140.
https://doi.org/10.1093/bioinformatics/btp616

Sampaio, V. da S., Coutinho, Í.A.C., Särkinen, T., Loiola, M.I.B., 2021. Secretory and ecological
function of petiolar glands in *Solanum fernandesii*: first description of resin glands in the genus
Solanum. Aust. J. Bot. 70, 32–41. https://doi.org/10.1071/BT21001

Särkinen, T., Bohs, L., Olmstead, R.G., Knapp, S., 2013. A phylogenetic framework for
evolutionary study of the nightshades (Solanaceae): a dated 1000-tip tree. BMC Evol. Biol. 13,
214. https://doi.org/10.1186/1471-2148-13-214

Sassaki, G.L., Souza, L.M., Serrato, R.V., Cipriani, T.R., Gorin, P.A.J., Iacomini, M., 2008.
Application of acetate derivatives for gas chromatography–mass spectrometry: novel approaches
on carbohydrates, lipids and amino acids analysis. J. Chromatogr. A 1208, 215–222.
https://doi.org/10.1016/j.chroma.2008.08.083

Schenck, C.A., Anthony, T.M., Jacobs, M., Jones, A.D., Last, R.L., 2022. Natural variation
meets synthetic biology: promiscuous trichome-expressed acyltransferases from Nicotiana. Plant
Physiol. kiac192. https://doi.org/10.1093/plphys/kiac192

Schilmiller, A.L., Charbonneau, A.L., Last, R.L., 2012. Identification of a BAHD acetyltransferase that produces protective acyl sugars in tomato trichomes. Proc. Natl. Acad. Sci. U.S.A. 109, 16377–16382. https://doi.org/10.1073/pnas.1207906109

Schilmiller, A.L., Gilgallon, K., Ghosh, B., Jones, A.D., Last, R.L., 2016. Acylsugar acylhydrolases: carboxylesterase-catalyzed hydrolysis of acylsugars in tomato trichomes. Plant Physiol. 170, 1331–1344. https://doi.org/10.1104/pp.15.01348

Schilmiller, A.L., Moghe, G.D., Fan, P., Ghosh, B., Ning, J., Jones, A.D., Last, R.L., 2015. Functionally divergent alleles and duplicated Loci encoding an acyltransferase contribute to acylsugar metabolite diversity in Solanum trichomes. Plant Cell 27, 1002–17. https://doi.org/10.1105/tpc.15.00087

Shapiro, J.A., Steffens, J.C., Mutschler, M.A., 1994. Acylsugars of the wild tomato *Lycopersicon pennellii* in relation to geographic distribution of the species. Biochem. Syst. Ecol. 22, 545–561. https://doi.org/10.1016/0305-1978(94)90067-1

Smith, C.R., 1971. Occurrence of unusual fatty acids in plants. Prog. Chem. Fats Other Lipids 11, 137–177. https://doi.org/10.1016/0079-6832(71)90005-X

Song, B., Song, Y., Fu, Y., Kizito, E.B., Kamenya, S.N., Kabod, P.N., Liu, H., Muthemba, S., Kariba, R., Njuguna, J., Maina, S., Stomeo, F., Djikeng, A., Hendre, P.S., Chen, X., Chen, W., Li, X., Sun, W., Wang, S., Cheng, S., Muchugi, A., Jamnadass, R., Shapiro, H.-Y., Van Deynze, A., Yang, H., Wang, J., Xu, X., Odeny, D.A., Liu, X., 2019. Draft genome sequence of *Solanum aethiopicum* provides insights into disease resistance, drought tolerance, and the evolution of the genome. GigaScience 8, giz115. https://doi.org/10.1093/gigascience/giz115

Stern, S., Agra, M. de F., Bohs, L., 2011. Molecular delimitation of clades within New World species of the "spiny solanums" (Solanum subg. Leptostemonum). TAXON 60, 1429–1441. https://doi.org/10.1002/tax.605018

Sumner, L.W., Amberg, A., Barrett, D., Beale, M.H., Beger, R., Daykin, C.A., Fan, T.W.-M., Fiehn, O., Goodacre, R., Griffin, J.L., Hankemeier, T., Hardy, N., Harnly, J., Higashi, R., Kopka, J., Lane, A.N., Lindon, J.C., Marriott, P., Nicholls, A.W., Reily, M.D., Thaden, J.J., Viant, M.R., 2007. Proposed minimum reporting standards for chemical analysis. Metabolomics 3, 211–221. https://doi.org/10.1007/s11306-007-0082-2

Tepe, E.J., Anderson, G.J., Spooner, D.M., Bohs, L., 2016. Relationships among wild relatives of the tomato, potato, and pepino. TAXON 65, 262–276. https://doi.org/10.12705/652.4

Van Dam, N.M., Hare, J.D., 1998. Biological activity of *Datura wrightii* glandular trichome exudate against *Manduca sexta* larvae. J. Chem. Ecol. 24, 1529–1549. https://doi.org/10.1023/A:1020963817685

Van der Auwera, G.A., Carneiro, M.O., Hartl, C., Poplin, R., del Angel, G., Levy-Moonshine, A., Jordan, T., Shakir, K., Roazen, D., Thibault, J., Banks, E., Garimella, K.V., Altshuler, D., Gabriel, S., DePristo, M.A., 2013. From FastQ data to high-confidence variant calls: the genome

analysis toolkit best practices pipeline. Curr. Protoc. Bioinforma. 43, 11.10.1-11.10.33.
https://doi.org/10.1002/0471250953.bi1110s43

Wagner, G.J., 1991. Secreting glandular trichomes: more than just hairs. Plant Physiol. 96, 675–
679. https://doi.org/10.1104/pp.96.3.675

Weese, T.L., Bohs, L., 2007. A three-gene phylogeny of the genus Solanum (Solanaceae). Syst.
Bot. 32, 445–463.

Weinhold, A., Baldwin, I.T., 2011. Trichome-derived O-acyl sugars are a first meal for
caterpillars that tags them for predation. Proc. Natl. Acad. Sci. 108, 7855–7859.
https://doi.org/10.1073/pnas.1101306108

Yu, G., Smith, D.K., Zhu, H., Guan, Y., Lam, T.T.-Y., 2017. ggtree: an r package for
visualization and annotation of phylogenetic trees with their covariates and other associated data.
Methods Ecol. Evol. 8, 28–36. https://doi.org/10.1111/2041-210X.12628

Yu, P., He, X., Baer, M., Beirinckx, S., Tian, T., Moya, Y.A.T., Zhang, X., Deichmann, M.,
Frey, F.P., Bresgen, V., Li, C., Razavi, B.S., Schaaf, G., von Wirén, N., Su, Z., Bucher, M.,
Tsuda, K., Goormachtig, S., Chen, X., Hochholdinger, F., 2021. Plant flavones enrich
rhizosphere Oxalobacteraceae to improve maize performance under nitrogen deprivation. Nat.
Plants 7, 481–499. https://doi.org/10.1038/s41477-021-00897-y

Zizovic, I., Stamenić, M., Orlović, A., Skala, D., 2007. Supercritical carbon dioxide extraction of
essential oils from plants with secretory ducts: mathematical modelling on the micro-scale. J.
Supercrit. Fluids 39, 338–346. https://doi.org/10.1016/j.supflu.2006.03.009

Supplementary Figures for

Trading acyls and swapping sugars: metabolic innovations in *Solanum* trichomes

Paul D. Fiesel[1], Rachel E. Kerwin[1], A. Daniel Jones[1], and Robert L. Last[1,2]

[1]Department of Biochemistry and Molecular Biology, Michigan State University, East Lansing, MI 48823 USA

[2]Department of Plant Biology, Michigan State University, East Lansing, MI 48823 USA

Corresponding Author: Robert L. Last, Michigan State University, Department of Biochemistry and Molecular Biology, 603 Wilson Road, East Lansing, MI 48824; Phone: 517-432-3278; email: lastr@msu.edu

Figure S1. *S. melongena* produces single stalked glandular trichomes on seedling above ground tissue but stellate nonglandular trichomes on non-seedling above ground tissue. (A) Close up photo of a *S. melongena* hypocotyl displaying glandular trichomes with similar morphology to the acylsugar-producing *S. lycopersicum* Type I/IV trichomes (Luckwill, 1943; Schilmiller et al., 2012). (B) Close up photo of a *S. melongena* leaf from a reproductive stage plant displaying non-glandular stellate trichomes. Leaf surface metabolite extracts from this tissue do not have detectable acylsugars by LC-MS.

Figure S2. Annotation of I3:18(4,4,10) and AI3:16(4,4,8) from *S. melongena* 67/3 using negative and positive mode MS and CID MS/MS fragmentation. (A) I3:18(4,4,10) negative and positive mode MS and MS/MS fragmentation. (B) AI3:16(4,4,8) negative and positive MS and MS/MS fragmentation. For each compound, ESI- MS (top panels) display the formate adduct accurate mass which was used to determine chemical formulas. ESI- MS/MS (second from the top panels) exhibit acyl chain carboxylate fragment ions for both compounds and stepwise loss of acyl chains for only AI3:16. ESI+ MS/MS exhibit stepwise losses of acyl chains for both compounds. Positive mode CID of AI3:16 also produces a fragment ion (*m/z* 429.2492) corresponding to the neutral loss of the pentose moiety. This indicates that all acyl chains reside on the hexose ring. The acylsugar LC-MS annotation method is described in the Materials and Methods. fa = fatty acid; k = ketene.

Figure S3. Acyl chain composition of *S. melongena* acylsugars. Relative abundance of acyl chains shown between three *S. melongena* PI 555598 leaf surface extracts from different seedlings. Straight acyl chains were identified with authentic reference standards and iso-branched chains were identified with NIST mass spectral library searches. Hydroxylated acyl chains were not included in this analysis and only detected acyl chains are included.

Figure S4: Identification of *S. melongena* acylsugar core composition through GC-MS analysis of alditol acetate sugar derivatives. *S. melongena* acylsugars collected from surface extracts were first saponified to remove acyl chains, and then, with or without acid hydrolysis to break the glycosidic linkages, sugar cores were derivatized to alditol acetates. (A) Alditol acetate derivatization of saponified *S. melongena* acylsugars yield a peak that comigrated with that of a *myo*-inositol standard. The traces displayed are GC-MS total ion chromatograms (TICs). (B) Alditol acetate pentose derivatives of saponified and hydrolyzed acylsugar cores comigrate with

that of a derivatized arabinose standard. The traces displayed are GC-MS TICs. Vertical scales for both panels are normalized to the largest signal within the displayed region.

Figure S5. GC-MS full scan mass spectra of 3-OH-14:0 ethyl ester stereoisomers as their MPTA derivatives. The mass spectra contain ions at *m/z* 189, 209, and 255 corresponding to the expected fragmentation of the derivatized fatty acid. EE = ethyl ester. Peak numbers correspond to the peaks displayed in Figure 2.

Figure S6. Negative mode MS/MS fragmentation of H3:22(4,4,16-O-p) from *S. sisymbriifolium* results in neutral loss of a pentose moiety. This neutral loss of a sugar group ($C_5H_{10}O_5$ plus formic acid) from the [M+formate]$^-$ ion *m/z* 751.4025 to form *m/z* 555.3522 is not observed with negative CID of other acyldisaccharides (Figures S2 and S7) suggesting an unusual glycosidic linkage on the acyl chain. fa = fatty acid; k = ketene; 16-O-p = pentosylated 16 carbon acyl chain.

Figure S7. Negative and positive mode MS and CID MS/MS fragmentation of dHH3:36(10,12-OH,14-OH) from a *S. atropurpureum* fruit surface extract and HH3:18(2,8,8) from a *S. heteropodium* leaf surface extract. Formate and ammonium adduct ions were selected by data-dependent acquisition software and fragmented with a ramped collision energy detailed in the Methods. Under negative mode CID MS/MS, each compound produces fragment ions corresponding to stepwise acyl chain loss down to a free sugar core fragment ion. In contrast, under positive mode CID MS/MS each compound produces fragment ions corresponding to neutral loss of a sugar moiety. (A) dHH3:36(10,12-OH,14-OH) MS and MS/MS fragmentation. Positive mode CID produces a fragment ion (*m/z* 569.4069) corresponding to a neutral loss of a deoxyhexose acylated with a ten carbon acyl chain. dHH = deoxyhexose-hexose (B) HH3:18(2,8,8) MS and MS/MS fragmentation. Positive mode CID produces a fragment ion (*m/z* 331.1836) corresponding to a neutral loss of a hexose acylated with an eight carbon acyl chain. HH = hexose-hexose; fa = fatty acid; k = ketene.

Figure S8. Acylsugar saponification produces free glycosylated fatty acids in four *Solanum* species supporting the identification of glycohydroxyacyl hexoses. All traces are extracted ion LC/MS chromatograms for pentosylated hydroxytetradecanoic acid, *m/z* 375.24±0.05 (retention time 2.90 min) from saponified leaf surface extracts. *S. melongena* does not contain detectable glycosylated fatty acids and acts as a negative control. Vertical scale is normalized to the largest signal within the displayed region (values in the upper right of each chromatogram).

Figure S9. Coelution analysis reveals that five *Solanum* species accumulate acylsugar positional or branching isomers that differ from *S. melongena* acylsugars. Extracted ion chromatograms were generated for combined signals of *m/z* 591.34 and 519.28 corresponding to I3:22(4,4,14-OH) and I3:18(4,4,10) [M+formate]$^-$ adducts, respectively. The yellow traces (bottom row) represent *S. melongena*, the blue traces represent the five other *Solanum* species (*S. prinophylum, S. abutiloides, S. dulcamara, S. villosum*, and *S. scabrum*), and the middle row green traces represent acylsugar mixtures between *S. melongena* and the species in the trace below. The mixed samples allow for corrections of retention time drift that may occur between samples.

Figure S10. Negative and positive mode MS and CID MS/MS fragmentation of G3:15(2,5,8) from *S. americanum*. Acylglucoses fragment characteristically in negative mode MS and MS/MS functions producing fragment ions corresponding to stepwise acyl chain loss. This is in stark contrast to acylinositol negative mode fragmentation which does not produce major fragment ions corresponding to stepwise acyl chain loss (Figure S2). This fragmentation difference distinguishes the two sugar cores and enables their annotation by LC-MS. G3:15 formate and ammonium adduct ions were selected by data-dependent acquisition software and fragmented with a ramped collision energy detailed in the Methods. fa = fatty acid; k = ketene.

Figure S11. Phylogenetic analysis of Eggplant Clade and Anguivi Grade species reveals *S. aethiopicum* is the only analyzed acylsugar-producing species to not accumulate detectable AI4:18(2,4,4,8). AI3:16(4,4,8) and AI4:18(2,4,4,8) presence and absence in surface extracts as detected by LC-MS are plotted on a phylogeny of the Eggplant Clade and Anguivi Grade. Species that either do not accumulate detectable acylsugars (*S. macrocarpon* and *S. virginianum)* or were not analyzed did not have AI3:16 and AI4:18 presence plotted. The phylogeny was modified from a previously published version (Aubriot et al., 2018). Full acylsugar profiles are detailed Tables 1, S3-8.

Figure S12. *S. melongena* BAHD tree. Shown is a phylogeny including 106 predicted BAHDs (PF002458) in the eggplant Smel_V4.1 reference genome, published reference BAHD sequences for clades I-VII, characterized ASAT sequences from other Solanaceae species, and the SaASAT3 and SaASAT3-L1 candidates from *S. aethiopicum*. The maximum likelihood tree was inferred from amino acid sequences using the Jones-Taylor-Thornton algorithm with seven rate categories in IQ-TREE v2.1.3. Bootstrap support calculated from 100,000 ultrafast bootstrap replicates and values >=95 indicate strong support. Clade III BAHDs are highlighted in purple,

with a purple dot marking the node. Branch lengths represent substitution rates (substitutions per site) as denoted by scale bar.

Figure S13. SmASAT3-L1 forward and reverse assay produced AI4:18 and AI3:16, respectively, coelute with plant produced AI4:18 and AI3:16. (A) Forward assay produced AI4:18 (top) coelutes with AI4:18 from a *S. melongena* 555598 leaf surface extract (bottom). The extracted ion chromatograms display the formate adduct of AI4:18, *m/z* 665.30. (B) Reverse assay produced AI3:16 (top) coelutes with AI3:16 from a *S. melongena* 555598 leaf surface extract (bottom). The extracted ion chromatograms display the formate adduct of AI3:16, *m/z* 623.20.

Figure S14. Semi-quantitative RT-PCR analysis of *S. aethiopicum* BAHD expression in glandular trichomes. *Elongation factor1α* (EF1α) was used as a positive control. P1 = primer pair 1; P2 = primer pair 2; S = shaved hypocotyl; T1 = PI 666075 glandular trichomes; T2 = Grif 14165 glandular trichomes.

Figure S15. MDS plot of *S. melongena* RNAseq data. Genome-wide expression patterns across eggplant trichomes, trichomeless hypocotyls, and roots. Depicted is a multidimensional scaling (MDS) plot demonstrating that our 18 eggplant RNAseq samples cluster tightly by tissue identity. Distance between points illustrates expression differences between pairs of samples, calculated as leading (i.e., largest absolute) log2 fold-change (FC) in two dimensions, dim 1 and dim 2. Along the y-axis (dim 2), samples are separated into three tissue-specific groups. Along the x-axis (dim 1), samples are separated into two groups representing aerial (trichomes and trichomeless hypocotyls) and subterranean (roots) tissues, respectively. Colors and shapes represent the three tissues. T, trichomes (green circles); H, trichomeless hypocotyls (yellow squares); R, roots (brown triangles).

Figure S16. *S. aethiopicum* **RT-PCR primers validated with gDNA controls.** Soaet10024792 primers amplified a region containing an intron resulting in the larger amplicon length.