Natural genetic variation underlying the negative effect of elevated CO₂ on ionome composition in *Arabidopsis thaliana*

Océane Cassan[1], Léa-Lou Pimparé[1], Timothy Mozzanino[1], Cécile Fizames[1], Sébastien Devidal[2], Fabrice Roux[3], Alexandru Milcu[2,4], Sophie Lèbre[5], Alain Gojon[1], Antoine Martin[1*]

1   IPSiM, Univ Montpellier, CNRS, INRAE, Institut Agro, 34060, Montpellier, France

2   Montpellier European Ecotron, Univ Montpellier, CNRS, Campus Baillarguet, 34980, Montferrier-sur-Lez, France

3   Laboratoire des Interactions Plantes-Microbes-Environnement, Institut National de Recherche pour l'Agriculture, l'Alimentation et l'Environnement, CNRS, Université de Toulouse, Castanet-Tolosan, France

4   CEFE, Univ Montpellier, CNRS, EPHE, IRD, 34293, Montpellier, France

5   IMAG, Univ Montpellier, CNRS, Montpellier, France

*Author for correspondence: antoine.martin@cnrs.fr

1 **Abstract**

2 The elevation of atmospheric CO_2 leads to a decline in the plant mineral content, which might

3 pose a significant threat to food security in the coming decades. To date, very few genes have

4 been identified as having a role in the negative effect of elevated CO_2 on plant mineral

5 composition. Yet, several studies have shown a certain degree of diversity in the ionome's

6 response to elevated CO_2, associated with genotypic variation. This suggests the existence of

7 genetic factors controlling the effect of CO_2 on ionome composition. However, no large-scale

8 studies have been carried out to date to explore the genetic diversity of the ionome responses

9 to elevated CO_2. Here, we used six hundred *Arabidopsis thaliana* accessions, representing

10 geographical distributions ranging from worldwide to regional and local environments, to

11 analyze the natural genetic variation underlying the negative effect of elevated CO_2 on the

12 ionome composition in plants. We show that the growth under elevated CO_2 leads to a global

13 and important decrease of the ionome content whatever the geographic distribution of the

14 population. We also observed a high range of genetic diversity in the response of the ionome

15 composition to elevated CO_2, and we identified sub-populations, showing effects on their

16 ionome ranging from the most pronounced to resilience or even to a benefit in response to

17 elevated CO_2. Using genome-wide association mapping on the response of each mineral

18 element to elevated CO_2 or on integrative traits, we identified a large set of QTLs and genes

19 associated with the ionome response to elevated CO_2. Finally, we demonstrate that the

20 function of one of these genes is associated to the negative effect of elevated CO_2 on the plant

21 mineral composition. This resource will contribute to understand the genetic mechanisms

22 underlying the negative effect of elevated CO_2 on plant mineral nutrition, and could help

23 towards the development of crops adapted to a high-CO_2 world.

Introduction

The elevation of atmospheric CO_2 concentration leads to a decline in the mineral composition of C3 plants (Gojon et al., 2023). The negative effect of elevated CO_2 on plant mineral composition has been observed worldwide, and alters the content of nutrients that are essential for human nutrition, such as nitrogen (N) and proteins, iron (Fe) or zinc (Zn) (Loladze, 2014). Therefore, the rise in atmospheric CO_2 poses a significant threat to food security in the coming decades. Indeed, several modeling approaches predict a decrease in plant-based nutrient availability due to the negative effect of elevated CO_2 on the mineral status of plants, leading to an additional risk of nutritional deficiency for hundreds of millions of people (Medek et al., 2017; Ebi and Loladze, 2019). However, the reasons why elevated CO_2 leads to the degradation of plant mineral composition are far from being well understood. To date, only a few genes with a potential regulatory role on this effect have been identified (Gao et al., 2019; Umnajkitikorn et al., 2020; Yang et al., 2020; Bouain et al., 2022; Sun et al., 2022; Cassan et al., 2023). Despite their limited number, these studies show that the detrimental effect of high CO_2 on the plant mineral status has genetic bases. In addition to this, several reports suggest that exploring the natural genetic variability of plants represents a major opportunity to understand the mechanisms by which high CO_2 leads to a decline in plant mineral composition (Myers et al., 2014; Zhu et al., 2018; Marcos-Barbero et al., 2021). Indeed, a significant diversity in the response of mineral composition to high CO_2 has been observed in several plant species. For protein and therefore N content, as well as for Fe or Zn content, substantial variations have been observed between small panels of genotypes from different species (Myers et al., 2014; Zhu et al., 2018; Marcos-Barbero et al., 2021). This implies the presence of a genetic diversity reservoir, which can facilitate the understanding of the ionome's response to high CO_2 and subsequently provide an opportunity to alleviate this negative

Figure 1: Elevated CO₂ negatively impacts the ionome content at the population-scale level in *Arabidopsis thaliana*. A. Representation of the experimental design used in this study. The content of eight mineral elements was assessed for around 600 *Arabidopsis thaliana* accessions coming from the REGMAP (B), LANGUEDOC (C) and TOU-A (D) populations. Each dot represents the value of the content of a mineral element for one accession (yellow: ambient CO₂ (aCO₂, ~420 ppm), blue: elevated CO₂ (eCO₂, 900 ppm). N (% of dry weight), Fe (μg.g⁻¹ dry weight), Zn (μg.g⁻¹ dry weight), Cu (μg.g⁻¹ dry weight), Mg (μg.g⁻¹ dry weight), Mn (μg.g⁻¹ dry weight), Na (μg.g⁻¹ dry weight), C (% of dry weight). Asterisks indicate significant differences (Paired Wilcoxson test; *, P<0.05; **, P<0.005; ***, P<0.0005). ns; not significant.

impact. However, in order to identify the genetic determinants of this negative response of

the ionome to high CO_2, large-scale approaches are necessary, but are still lacking for the

moment. The objective of this work was to fill the aforementioned knowledge gap by using a

large collection of natural genotypes of the model plant *Arabidopsis thaliana.* This allowed to

explore in depth the natural variation of the ionome response to elevated CO_2, and to

generate a resource of phenotypic data that can be used in association genetics approaches.

To this end, we used several hundreds of accessions from different geographic scales of *A.*

thaliana, and analyzed their leaf mineral composition under contrasted conditions of CO_2

concentration. This allowed us to extract the general trends in the leaf ionome response to

high CO_2, and to identify a large set of genes associated with the variation in the mineral

composition of plants in response to high CO_2. By combining this information with genome

expression data under elevated CO_2, we end up by functionally validating one of these genes

for its importance in the reduction of Zn content under elevated CO_2. Our results open the

way for a better understanding of the genetic and molecular mechanisms involved in the

regulation of plant mineral nutrition by the elevation of atmospheric CO_2.

Results

In order to explore the natural variation and identify its underlying genetic basis associated

with the negative effect of elevated CO_2 on plant ionome, we used three populations of *A.*

thaliana representing different geographic scales (i.e., the worldwide REGMAP population,

the LANGUEDOC regional population and the local TOU-A population from east of France) and

displaying different levels of genetic diversity (Fig. 1A). These populations were grown under

ambient or elevated CO_2, and we measured in each accession the composition of their ionome

in rosettes, including C, N, Na, Fe, Mg, Mn, Zn and Cu content.

Figure 2: Elevated CO$_2$ leads to high phenotypic diversity of ionome response in *Arabidopsis thaliana.* Distributions of the relative change (%) of the content of 8 mineral elements between elevated CO$_2$ and ambient CO$_2$, in each population (A: REGMAP, B: LANGUEDOC, C: TOU-A). Each dot represents the value of the relative change of the content a mineral element for one accession. The name of the element appears in bold if the mean of the element in elevated CO$_2$ is significantly different from the mean of the element in ambient CO$_2$ (Paired wilcoxon test, significance threshold of 0.05).

Elevated CO$_2$ globally decreases ionome content at the population level, whatever the

geographic scale.

In the three *A. thaliana* populations, we observed a global and important decrease of the

ionome content when plants were grown under elevated CO$_2$ as compared to ambient CO$_2$.

This was particularly the case for N and Fe, for which the decrease in content was very robust

and important in each of the population analyzed (Fig. 1B-D). Zn, Cu and Mg content were also

negatively affected to a significant extent by the growth under elevated CO$_2$ in the REGMAP

and in the TOU-A populations (Fig. 1B, C), although not significantly in the LANGUEDOC

population (Fig. 1D). More variability for the effect of elevated CO$_2$ was observed on Mn and

Na content, which were decreased in the REGMAP population, but not significantly changed

in the TOU-A and LANGUEDOC populations, respectively. In parallel, the C content of these

populations increased under elevated CO$_2$, by very significant factors for the REGMAP and the

LANGUEDOC populations. Altogether, these observations demonstrate that elevated CO$_2$ has

on average a negative impact on the mineral content of natural genotypes of *A. thaliana* at

the population-scale, whatever their geographic distribution.

The ionome of *Arabidopsis thaliana* natural accessions displays a high range of phenotypic

diversity in response to elevated CO$_2$.

To explore the effect of elevated CO$_2$ in each accession, we calculated the relative change in

nutrient composition of *A. thaliana* accessions from the three populations in response to

elevated CO$_2$. In agreement with the results previously mentioned, we observed that the

median relative change of most nutrient content at the population-level was negatively

affected by elevated CO$_2$ (Fig. 2). But the most striking observation was the genetic diversity

of ionome response observed in these populations. Indeed, while most the natural accessions

Figure 3: Elevated CO₂ results in a general pattern of ionome variation common to most accessions constituting natural populations of *Arabidopsis thaliana*. Principal Component Analysis (PCA) was performed using the variation of each element in response to elevated CO₂. A. Natural accessions were positioned on the PCA and colored based on population. B. Contribution of each element to the PCA axis.

were negatively affected by elevated CO_2 (with a negative relative ratio of their nutrient

content between ambient and elevated CO_2), a considerable number of accessions were

rather not affected by elevated CO_2, or even positively affected, therefore showing an

improved nutrient composition under elevated CO_2. For macronutrients like N, the relative

change of concentration between ambient and elevated CO_2 varied from 20% to -50%, and for

micronutrients like Cu, Fe or Zn, the relative change of concentration between ambient and

elevated CO_2 varied from 100% to -60% (Fig. 2). In addition, some differences among nutrients

were observed between populations. For instance, a smaller dispersion of Fe relative change

in the LANGUEDOC population, against a higher distribution of Mn relative change.

In order to explore the behavior of the different elements in response to elevated CO_2 and to

observe the structure of phenotypic variation, we performed a principal component analysis

(PCA) of the relative change in the 8 elements for the accessions from the three populations.

The accessions from all populations seem to have globally similar responses to elevated CO_2,

as suggested by the overlap of the three populations in the two first principal components

(Fig. 3A). The first component of the PCA described a clear antagonistic trend between C

content and the change of other mineral elements (Fig. 3B), suggesting that most of the

variation between accessions in term of mineral response (almost 40%) could be driven by

one or a few mechanisms resulting in an inverse variation between the whole ionome and C

change (Fig. 3B). Interestingly, the second component, explaining almost 15% of the variation

among accessions in term of mineral response, was mainly driven jointly by change in N and

C concentration. Altogether, these results show that there is a marked and large variability

among accessions in their mineral concentration in response to elevated CO_2, illustrated by

accessions negatively affected by elevated CO_2 and others positively affected by elevated CO_2.

Figure 4: Variation in the response of the ionome to elevated CO$_2$ identifies contrasting subpopulations inside the REGMAP panel. K-means clustering was performed in the REGMAP accessions to identify different subpopulations. Each accession is represented by a dot, connected by a line between each element. Cluster 1:65 accessions. Cluster 2: 25 accessions. Cluster 3: 69 accessions.

In order to explore specific behavior of sub-populations, we clustered the accessions from the

REGMAP panel via a k-means approach. This multivariate clustering resulted in the

partitioning of accessions in three groups (Fig.4 – Suppl. Table 1). Cluster 1 displayed the most

negative pattern of ionome response to elevated CO_2. Inversely, accessions included in Cluster

2 displayed a globally positive response, with the highest relative change for almost all mineral

elements, except for C content. These accessions did not appear to be clustered

geographically with respect to their collection origin in the REGMAP panel (Suppl. Fig. 1),

which is in line with the high genetic diversity of response to elevated CO_2 observed at smaller

geographical scales (Fig.2). Finally, Cluster 3 displayed a resilient pattern, with accessions

showing a globally attenuated response to elevated CO_2. Interestingly, the large phenotypic

diversity of the ionome observed in the three populations in response to high CO_2, as well as

the presence of contrasted subpopulations in the REGMAP panel, suggests the presence of

genetic determinants associated with this response.

Genetic architecture of the ionome response to elevated CO_2, and identification of genetic

determinants

We ran Genome-Wide Association (GWA) mapping to describe the genetic architecture of the

ionome response to elevated CO_2, and to fine-map candidate genes underlying the detected

quantitative trait loci (QTLs). We focused here on the phenotypic data collected on the

REGMAP population, and used the sequencing data available for this population (Arouisse et

al., 2020). We included in this analysis the level of each mineral under ambient and under

elevated CO_2, as well as the relative change between ambient and elevated CO_2 for each

element. We also included a trait corresponding for each accession to the coordinate on the

first and on the second PCA axes (PCA1 and PCA2) explaining collectively more than 50% of

Figure 5: Genetic architecture of the response of the ionome to elevated CO_2 in the REGMAP panel of *Arabidopsis thaliana*. A. Manhattan plots for the content of eight mineral elements under elevated CO_2, or for the relative change of the content of mineral elements between elevated CO_2 and ambient CO_2. For each Manhattan plot, SNPs with the 50 most significant P-value, located above the horizontal line, are colored. Bar plots showing the number of SNPs identified by GWAs for traits under elevated CO_2 (B) or for the relative change of the content of mineral elements between elevated CO_2 and ambient CO_2 (C) that are unique to one element or shared between 2 or 3 traits.

ionomic variation (Fig. 3). Therefore, these values correspond to traits driving and

summarizing a large part of the ionome variation under elevated CO_2. This resulted as a whole

in running GWA mapping on 30 different single-trait GWAS. The overall approach was first

validated by observing expected results for traits phenotyped under ambient CO_2. For

instance, we observed a very strong peak for the Na content at the locus of the *HKT1* gene

(Suppl. Fig. 2A), which is known to be involved in the natural genetic variation of Na content

in *Arabidopsis thaliana* (Baxter et al., 2010), or a strong peak for the N content at the locus of

the *NIA1* gene (Suppl. Fig. 2B), encoding for an isoform of the nitrate reductase required for

the first step of nitrate reduction and associated with natural genetic variation of N content

in *A. thaliana* (North et al., 2009).

GWA mapping revealed a polygenic architecture for each phenotypic trait, although its

complexity largely differs among traits. For instance, very few and neat peaks of association

were detected Na and Mn content under elevated CO2, or of Fe and Cu relative change

between ambient and elevated CO_2 (Fig.5A, Suppl. Fig. 3). On the other hand, a more complex

genetic architecture with the detection of a large number of QTLs was observed for traits

related to N or C content (Fig.5A, Suppl. Fig. 3). For each of the traits that have been analyzed

under elevated CO_2 or corresponding to the relative change of their content between ambient

and elevated CO_2, we isolated the 50 SNPs with the most significant p-value, hereafter named

top SNPs (Fig. 5A, Suppl. Tables 2 and 3). In order to identify the overlap between the genetic

architecture of each trait, we looked whether some of the top SNPs were shared among traits.

While the large majority of SNPs were specific to one trait, 30 and 21 SNPs were shared

between two traits for the content under elevated CO_2 or for the relative change between

ambient and elevated CO_2, respectively (Fig. 5B and C, Supplemental Tables 2 and 3). In

addition, 8 and 2 SNPs were shared between three traits for the content under elevated CO_2

Figure 6: Identification of genes detected by GWA mapping and differentially regulated by elevated CO$_2$. Intersection between elevated CO$_2$-DEG in shoot (A) or root (B) and genes identified by GWA mapping. UpSet plots display the number of elevated CO$_2$-DEG that are associated to a locus identified for the content or the relative change of one or several mineral elements under elevated CO$_2$. Illustration of the pattern of elevated CO$_2$-DEG in shoot (C) or root (D) also identified by GWA mapping.

or for the relative change between ambient and elevated CO_2, respectively (Fig. 5B and C,

Supplemental Tables 2 and 3). Most of the shared SNPs were associated with micronutrients

(Fe, Mn, Zn and Mg content) and with N and/or with the first component of the PCA axis. An

interesting QTL located on chromosome 1 was notably associated with 6 traits, displaying

SNPs shared between Mn, Zn and N relative change and SNPs shared between Mn, N and PC1

content under elevated CO_2 (Fig. 5A, Suppl. Tables 2 and 3). Another QTL located on

chromosome 3 encompasses SNPs shared between Fe, Zn and PC1 content under elevated

CO_2 (Fig. 5A, Suppl. Table 2).

We next identified for each trait a list of the genes located at ±25 kb from the top 50 SNPs,

which corresponds to the rough estimate of the decay of linkage disequilibrium identified in

A. thaliana at the worldwide scale (Kim et al., 2007). This resulted in a list of genes for each

element, ranging from 154 to 422 genes depending on the element (Suppl. Tables 2 and 3).

Among others, several genes associated with top 50 SNPs were identified as obvious

candidates of the effect of elevated CO_2 on plant nutrition and ionome content. This was the

case of *ZINC INDUCED FACILITATOR 1* (*ZIF1*, *AT5G13740*) and *ZIF-LIKE1* (*AT5G13750*), linked

with SNPs identified for Zn content under elevated CO_2, and involved Zn sequestration

mechanisms (Lee et al., 2021). We also noticed the link between SNPs identified for Zn relative

change and *TIP2;2* (*AT4G17340*), known to be involved in Zn root-to-shoot translocation

(Wang et al., 2022). Concerning N relative change, some of the top 50 SNPs were linked to

ASN1 (*AT3G47340*), which is an actor of N status and remobilization (Lam et al., 2003;

Gaufichon et al., 2017). Some of the top 50 SNPs identified for Fe relative change were linked

to *MCO2* (*AT5G21100*) and *MCO3* (*AT5G21105*) genes, which have been recently

characterized as actors of the regulation of Fe homeostasis (Brun et al., 2022). Finally, it is

interesting to note that the QTL located on chromosome 3 mentioned above displaying

Figure 7: Natural variation of the *TIP2;2* gene is associated with improved responses of Zn content to elevated CO$_2$. A. Manhattan plot of the relative change of Zn content between elevated CO$_2$ and ambient CO$_2$ showing the presence of a peak closed to the *TIP2;2* locus. **B.** Comparison of haplotypes and their relative change of Zn content between elevated CO$_2$ and ambient CO$_2$. Three SNPs located at the *TIP2;2* locus are associated to an improvement of Zn content under elevated CO$_2$ for accessions that possess them (haplotype 1) compared to the rest of the population (haplotype 0). **C, D.** Relative expression of *TIP2;2* in the roots under ambient (**C**) or elevated (**D**) CO$_2$ for accessions belonging to haplotype 0 or haplotype 1. Relative expression levels were calculated based on *UBQ10* as internal control. Horizontal black line represented the median of each group of haplotypes. ***P < 0,001, **P < 0,01, unpaired Mann-Whitney test. **E.** Shoot Zn content under ambient or elevated CO$_2$ for WT (Columbia) and *tip2;2-1* mutant. Data are presented as the mean (with SD) of 5 and 6 biological repeats for the WT and *tip2;2-1*, respectively. **P < 0.01, unpaired Mann-Whitney test.

significant shared SNPs identified for Fe, Zn and PC1 content under elevated CO_2 was

associated among other genes with *ISU2* (*AT3G01020*), coding for one of the Fe-S clusters in

Arabidopsis thaliana, which are known to be essential for photosynthesis and metabolism

(Balk and Schaedler, 2014). Altogether, this demonstrated that genes identified through this

approach represent a large and valuable reservoir of candidates to study and to counteract

the effect of elevated CO_2 on plant nutrition and ionome content.

To analyze how these genes identified by GWA mapping are regulated by elevated CO_2, we

performed RNA-seq from shoots and roots grown under ambient and elevated CO_2.

Differentially expressed genes (DEG) associated to the effect of elevated CO_2 were identified

from shoots and roots (Suppl. Table 4). We compared the list of shoots or roots elevated CO_2-

DEG with the list of genes identified by GWA mapping for each element, which resulted in a

list of 182 genes identified by GWA mapping and differentially regulated by elevated CO_2 in

shoot or in roots (Suppl. Table 5), making them relevant candidates to be involved in the

response of the mineral composition of plants to elevated CO_2. Most of these genes were

deregulated by elevated CO_2 in shoot (Fig. 6A, B). In shoot or in roots, these genes mainly

showed an association with C-, Mg- or Zn-related traits (Fig. 6A, B). Several of these genes,

identified by GWA mapping and whose expression is deregulated in response to high CO_2,

were known for their role in nutrient homeostasis. This was the case for the *ASN1* and *DUR3*

genes, encoding an asparagine synthase and a urea transporter involved in N metabolism and

remobilization, both associated here with a N-related peak of association, and whose

expression is modulated by high CO_2 in leaves (Fig. 6C). We also observed in the leaves an

interesting profile for several genes related to C metabolism and photosynthesis. This was the

case for the *BGP3* gene, involved in chloroplast development, or for the carbonic anhydrase

CA1, both showing a decreased expression in response to high CO_2 and both associated with

a peak in C-related GWA mapping under elevated CO_2 (Fig. 6C). In roots, the gene most

deregulated in response to high CO_2 was *AT1G64710*, encoding a GroES-type alcohol

dehydrogenase, which interestingly is also deregulated in leaves (Fig. 6D). We also observed

in the roots that the expression of the *TIP2;2* gene, associated here with a peak detected for

Zn relative change GWA mapping, was deregulated in response to elevated CO_2 (Fig. 6D).

To go further, we selected one of the association peaks identified by GWA mapping, and

sought to functionally validate the importance of this QTL in response to elevated CO_2, in order

to demonstrate the value of our data set and GWA mapping analyses. To do so, we selected

an association peak located on chromosome 4 and associated with Zn relative change (Fig.

7A). More precisely, this association peak displayed the SNPs with the most significant *p*-

values and more largely three SNPs that fell into the top 10 SNPs of the trait corresponding to

the Zn relative change between ambient and elevated CO_2. The SNPs corresponding to the

alternative alleles were associated to an increase of Zn content under elevated CO_2 (Fig. 7B).

These SNPs are located very close to the *TIP2;2* (*AT4G17340*) gene, which has been recently

characterized as an actor of Zn root-to-shoot translocation (Wang et al., 2022). We thus

selected a set of accessions from haplotype 0 (reduced Zn content under elevated CO_2) or

haplotype 1 (increased Zn content under elevated CO_2), and analyzed *TIP2;2* expression in the

roots under ambient and elevated CO_2. This analysis revealed a haplotype-specific difference

in *TIP2;2* expression: under ambient and elevated CO_2, accessions from haplotype 1

(correlated with a higher Zn content in the shoot (Fig. 7B)) show a reduced *TIP2;2* expression

in the roots compared to those from haplotype 1 (Fig. 7C & D), with a reduction being more

pronounced under elevated CO_2. To test the effect of *TIP2;2* expression of Zn content under

elevated CO_2, we used the *tip2;2-1* knock-out mutant and compared its Zn content under

ambient and elevated CO_2 to this of the WT. We observed that the *tip2;2-1* mutant line did

not present any decrease in Zn shoot content in response to elevated CO_2, in opposition to

what is observed for the WT (Fig. 7E). In the *tip2;2-1* mutant, Zn content under elevated CO_2

was even slightly higher than under ambient CO_2. Altogether, these results demonstrated that

these data sets generated in this study and the associated analyses are a valuable resource to

identify genes and associated mechanisms involved in the effect of elevated CO_2 on the

mineral composition of plants.

Discussion

The natural variation of ionome response to elevated CO_2 in *Arabidopsis thaliana* displays a

high degree of genetic variation

In the present work, we analyzed the diversity of leaf ionome response to elevated CO_2

present in the natural variation of *Arabidopsis thaliana*. In agreement with several other

phenotypic traits related to phenology and disease resistance (Brachi et al., 2013; Huard-

Chauveau et al., 2013; Roux and Frachon, 2022), we observed a wide range of responses at

complementary geographical scales, from accessions with an ionome negatively affected by

high CO_2 to accessions with an ionome benefiting from high CO_2. This confirms for the first

time on large and complementary sets of natural genotypes what has been observed by meta-

analysis on isolated groups of plants worldwide (Loladze, 2014; Myers et al., 2014). The global

analysis of the distribution of each mineral element studied suggests firstly a trend where the

whole ionome would evolve in a unified manner in response to high CO_2, and in an opposite

manner to C. This is in line with a number of studies that have proposed that the accumulation

of carbohydrates due to the stimulation of photosynthesis by high CO_2 would be the cause of

the decrease in plant mineral composition (Ainsworth and Long, 2005; Thompson et al., 2017;

Dusenge et al., 2019; Tausz-Posch et al., 2020). However, the reading of the genetic

architecture performed here by a genome-wide association genetics approach suggests that

the majority of the genetic mechanisms underlying the negative effect of elevated CO_2 on the

ionome are specific to each mineral element. Some specific cases, such as the QTL detected

on chromosome 1 and associated with the natural genetic variation of 6 traits among the 20

considered, will certainly deserve a more in-depth analysis.

By clustering globally distributed accessions according to their ionome sensitivity to high CO_2,

we were able to observe that the geographic origin of the accessions likely did not determine

their response to CO_2. This suggests that inherent genetic factors, more than those due to

local adaption, direct the response of plants to elevated CO_2. This seems consistent since the

CO_2 elevation applied here to natural *Arabidopsis thaliana* variants does not correspond to

any environment experienced by plants yet, at least for several tens of millions of years

(Pearson and Palmer, 2000; Luthi et al., 2008). In this context of brutal and highly impactful

environmental change, the presence of cryptic genetic variation often explains the

appearance of relatively rapid adaptive mechanisms (Pauls et al., 2013; Cortés and López-

Hernández, 2021). Although not formally tested here, it would be interesting to examine

whether the variation in the ionome in response to elevated CO_2 shows evidence of cryptic

variation. In any case, the presence of high phenotypic diversity in these natural populations

of *A. thaliana* demonstrates very clearly the possibility of taking advantage of this genetic

variation to understand and alleviate the negative response of plant mineral composition to

high CO_2.

GWA mapping of ionome variation under elevated CO_2 identified a large number of genes

to understand and mitigate the negative effect of high CO_2 on plant mineral composition

In order to understand the genetic mechanisms underlying the effect of high CO_2 on plant

mineral composition, and to enable future breeding approaches, we adopted an association

genetics approach. This led to the identification of a large number of candidate genes

associated to the variation of nutrients under elevated CO_2. Several genes in this list can easily

attract attention. In particular, we can note the identification of *ASN1* and *DUR3* genes in two

of the loci associated with N content variation under elevated CO_2. *ASN1*, and to a lesser

extent *DUR3*, play an important role in the remobilization and the reallocation of N within the

plant, and their manipulation can lead to variation in N use efficiency (Lam et al., 2003; Bohner

et al., 2015; Gaufichon et al., 2017). This is interesting because for the moment, root N uptake

and N assimilation seemed to be the key targets of the negative effect of high CO_2 on plant N

content (Bloom et al., 2010; Cassan et al., 2023), but these results suggest that remobilization

of N may also be involved. We also identified the *CA1* gene, coding for a carbonic anhydrase,

in the vicinity of a QTL associated with C variation under high CO_2. *CA1* is involved in the

regulation of stomatal opening by elevated CO_2 (Hu et al., 2015), and the β carbonic anhydrase

family of which *CA1* belongs is involved in the regulation of photosynthetic efficiency,

although *CA1* shows no significant effect under standard conditions (Sharma et al.). It would

be therefore interesting to assess the role of *CA1* natural genetic variation under elevated CO_2.

If *CA1* regulates the C variation of the ionome under elevated CO_2, this could, according to our

observations, significantly influence the global mineral composition of plants. Interestingly,

the genes identified by GWA mapping in the ionome response to high CO_2, including those

mentioned above, showed substantial variation at the gene expression level. We ended this

study with the functional validation of an association peak identified by GWA mapping for the

relative change of Zn content between ambient and elevated CO_2. Zn is an essential element

for a large number of metabolic processes in humans, and Zn deficiency, found in up to one

third of the world's population, leads to severe health problems. We demonstrated that *TIP2;2*

gene expression varied in a haplotype-specific manner, in both ambient and elevated CO_2. In

parallel to this, we show that loss of *TIP2;2* expression using a knock-out mutant can abolish

the Zn decrease observed under high CO_2. It therefore seems that variation in *TIP* expression

is associated with the effect that atmospheric CO_2 can have on zinc levels. A recent study

demonstrated that TIP2;2 was responsible for Zn retention in the roots (Wang et al., 2022). It

therefore seems consistent that natural accessions with the lowest expression levels of this

gene are those with the highest Zn content in aerial parts, due to low retention in their roots.

This example illustrates the potential of the resource we have generated here towards the

identification of genes involved in the variation of leaf ionome in response to rising CO_2, and

towards the characterization of the associated mechanisms. The understanding of these

mechanisms represents a considerable challenge in view of the current rise in atmospheric

CO_2, which might be useful to the coming breeding programs for crops adapted to a

forthcoming CO_2-rich atmosphere (Shahzad and Rouached, 2022). It must however be recalled

that leaf ionome is not always correlated with seed ionome, in particular in *Arabidopsis*

thaliana accessions (Campos et al., 2021). Therefore, the effect of elevated CO_2 on seed

ionome, certainly in major crops, must be the next target to analyze in order to pave the way

for the development of nutritious crops adapted to elevated CO_2.

Methods

Data and code availability

Data and R notebooks containing the analyses performed in this article can be found at

https://src.koda.cnrs.fr/groups/ipsim/sirene-team. RNA-seq data generated for this study are

available at https://www.ebi.ac.uk/biostudies/arrayexpress/studies using the accession no E-

MTAB-13661.

Plant Material

A subset of the REGMAP panel, the LANGUEDOC panel and the TOU-A panel were used in this

study. These populations were previously described here (Horton et al., 2012; Brachi et al.,

2013; Frachon et al., 2017). These populations were grown on Jiffy-7 peat pellets (Jiffy

Products International, NL) under ambient (~420 ppm) or elevated (900 ppm) CO_2 in the

growth chambers of the Microcosms experimental platform at the Montpellier European

Ecotron CNRS. Five replicates of each accession were randomly distributed in the growth

chambers. The concentration of 900 ppm of CO_2 was chosen as current CO_2 emissions align

with the IPCC's RCP8.5 model, which predicts a CO2 concentration of around 900 ppm in 2100.

Growth conditions were 6-h/22-h light (22°C) / dark (20°) photoperiod, with 200 µmol m^{-2} s^{-1}

light intensity and 65% of hygrometry. Plants were watered twice a week with a growth

solution containing KH_2PO_4 1 mM, $MgSO_4$ 1 mM, K_2SO_4 250 µM, $CaCl_2$ 250 µM, Na-Fe-EDTA

100 µM, KNO_3 10 mM, KCl 50 µM, H_3BO_3 30 µM, $MnSO_4$ 5 µM, $ZnSO_4$ 1 µM, $CuSO_4$ 1 µM,

$(NH_4)_6Mo_7O_{24}$ 0,1 µM, as described by (Gansel et al., 2001). The entire rosettes were collected

three weeks after sowing. The *tip2;2-1* mutant line corresponds to the *SALK_152463* allele

(Wang et al., 2022).

Ionome analysis

From 3 to 5 replicates per accession were pooled and used for each ionome analysis. Total C

and N content was obtained from dried shoot tissue using an Elementar Pyrocube analyzer.

Cu, Fe, Mg, Mn, Na and Zn content was obtained from dry shoot tissue mixed with 750 µl of

nitric acid (65% [v/v]) and 250 µl of hydrogen peroxide (30% [v/v]). After one night at room

temperature, samples were mineralized at 85°C during 24 hours. Once mineralized, 4 ml of

milliQ water was added to each sample. Mineral contents present in the samples were then

measured by microwave plasma atomic emission spectroscopy (MP-AES, Agilent

Technologies).

Removal of outlier observations

Prior to GWAS and multivariate analyses such as PCA or clustering, mineral composition

measures were pre-processed to remove technical outliers. For a given element and CO_2

condition, the values positioned more than 5 median absolute deviations away from the

median were removed from the dataset. The number of outliers removed from each dataset

is indicated in Supplemental Table 7.

PCA and Clustering

Principal Component Analysis was performed using the R *ade4* package after the prior scaling

of the variables to a z-score. Clustering of the REGMAP panel based on the relative changes of

the mineral composition of each accession has been done using a k-means clustering with the

R *kmeans* function. For this step, the variables were also scaled to a z-score. The number of

clusters in the k-means algorithm was chosen by the elbow method on the criteria of cluster

homogeneity (within-sum of squares).

GWAs

Genome-Wide Association mapping was performed using the R *statgenGWAs* package.

Genotype data was prepared using the *codeMarkers* function, removing duplicated SNPs and

filtering for a minimum allele relative frequency of 0.04. Associations were performed by the

runSingleTraitGwas function, that implements the EMMA algorithm. Population structure was

modeled via a kinship matrix built from the Astle method. Manhattan plots were drawn using

the *manPlotFast* function of the *ramwas* R package.

RNA-seq experiments

Plants from the Columbia accession were grown in hydroponics to have access to the roots in

addition to the shoot, as previously described in (Cassan et al., 2023). Shoot or root from 5

plants were pooled into one biological replicate, flash frozen in liquid nitrogen, and stored at

-80◦C. RNA of three biological replicates were extracted from shoot or root tissues using

Direct-zol RNA Miniprep (Zymo Research, CA, USA), according to the manufacturer

recommendations. RNA-sequencing libraries were done from shoot or root total RNA using

standard RNA-Seq protocol method (Poly-A selection for mRNA species) by the Novogene

company. RNA-sequencing was performed using Illumina technology on a NovaSeq6000

system providing PE150 reads. The quality control and adapter trimming of raw paired-end

fastq files was done with *fastp* and its default parameters. Mapping to the TAIR10 reference

genome was performed with STAR, and using the following options:

--outSAMtype BAM SortedByCoordinate

--outFilterMismatchNmax 1

--outFilterMismatchNoverLmax 0.15

--alignIntronMin 30

--alignIntronMax 5000

Quantification of the bam files against the TAIR10 GFF3 annotation file was done using *htseq-*

count with options:

-f bam --type gene -r pos

--idattr=Name --stranded=no

Normalization and differential expression were performed using *DIANE* R package (Cassan et

al., 2021), with no fold change constraint, and an adjusted p-value threshold (FDR) of 0.05.

Lowly expressed genes with an average value across conditions under 25 reads were excluded

from the analysis.

Quantitative real-time PCR

Plants were grown in hydroponics to have access to the roots, as previously described in

(Cassan et al., 2023). Root tissue from 5 plants were pooled into one biological replicate, flash

frozen in liquid nitrogen, and stored at -80∘C. RNA were extracted from shoot or root tissues

using TRIZOL (Invitrogen, USA), according to the manufacturer recommendations, and DNAse

treated using RQ1 (Promega, USA). Reverse transcription was achieved from 1 μg of total RNA

with M-MLV reverse transcriptase (RNase H minus, Point Mutant, Promega, USA) using an

anchored oligo(dT)20 primer. Accumulation of transcripts was measured by qRT-PCR

(LightCycler 480, Roche Diagnostics, USA) using the SYBR Premix Ex TaqTM (TaKaRa, Japan).

Gene expression was normalized using *UBQ10* and *ACT2* as internal standards. Results are

presented as the expression relative to *UBQ10*. Sequences of primers used in RT-qPCR for

gene expression analysis are listed in Supplemental Table 6.

Acknowledgments

This work was supported by the I-Site Montpellier Université d'Excellence (MUSE; project

ECO2THREATS), the CNRS through the Mission for Transversal and Interdisciplinary Initiatives

(MITI) 80 PRIME program, and the Plant Biology and Breeding department of INRAE. O.C. was

recipient of a PhD fellowship from the CNRS. T.M. was recipient of a PhD fellowship from

INRAE and Région Occitanie. We thank Jiping Liu from Cornell University for the gift of *tip2;2-*

1 mutant line. This study benefited from the CNRS human and technical resources allocated

to the Ecotron Research Infrastructure and from the state allocation "Investissement

d'Avenir" AnaEE France ANR-11-INBS-0001. A CC-BY public copyright license has been applied

by the authors to the present document.

References

Ainsworth, E.A., and Long, S.P. (2005). What have we learned from 15 years of free-air CO_2

enrichment (FACE)? A meta-analytic review of the responses of photosynthesis,

canopy properties and plant production to rising CO_2. New Phytol **165,** 351-371.

Arouisse, B., Korte, A., van Eeuwijk, F., and Kruijer, W. (2020). Imputation of 3 million SNPs

in the Arabidopsis regional mapping population. The Plant Journal **102,** 872-882.

Balk, J., and Schaedler, T.A. (2014). Iron cofactor assembly in plants. Annu Rev Plant Biol **65,**

125-153.

Baxter, I., Brazelton, J.N., Yu, D., Huang, Y.S., Lahner, B., Yakubova, E., Li, Y., Bergelson, J.,

Borevitz, J.O., Nordborg, M., Vitek, O., and Salt, D.E. (2010). A Coastal Cline in Sodium

Accumulation in *Arabidopsis thaliana* Is Driven by Natural Variation of the Sodium

Transporter AtHKT1;1. PLOS Genetics **6,** e1001193.

Bloom, A.J., Burger, M., Rubio Asensio, J.S., and Cousins, A.B. (2010). Carbon dioxide

enrichment inhibits nitrate assimilation in wheat and Arabidopsis. Science **328,** 899-

903.

Bohner, A., Kojima, S., Hajirezaei, M., Melzer, M., and von Wirén, N. (2015). Urea

retranslocation from senescing Arabidopsis leaves is promoted by DUR3-mediated

urea retrieval from leaf apoplast. The Plant Journal **81,** 377-387.

Bouain, N., Cho, H., Sandhu, J., Tuiwong, P., Prom, U.T.C., Zheng, L., Shahzad, Z., and

Rouached, H. (2022). Plant growth stimulation by high CO_2 depends on phosphorus

homeostasis in chloroplasts. Curr Biol.

Brachi, B., Villoutreix, R., Faure, N., Hautekèete, N., Piquot, Y., Pauwels, M., Roby, D.,

Cuguen, J., Bergelson, J., and Roux, F. (2013). Investigation of the geographical scale

of adaptive phenological variation and its underlying genetics in *Arabidopsis thaliana*.

Molecular Ecology **22,** 4222-4240.

Brun, A., Smokvarska, M., Wei, L., Chay, S., Curie, C., and Mari, S. (2022). MCO1 and MCO3,

two putative ascorbate oxidases with ferroxidase activity, new candidates for the

regulation of apoplastic iron excess in Arabidopsis. Plant Direct **6,** e463.

Campos, A., van Dijk, W.F.A., Ramakrishna, P., Giles, T., Korte, P., Douglas, A., Smith, P., and

Salt, D.E. (2021). 1,135 ionomes reveal the global pattern of leaf and seed mineral

nutrient and trace element diversity in *Arabidopsis thaliana*. Plant J **106,** 536-554.

Cassan, O., Lebre, S., and Martin, A. (2021). Inferring and analyzing gene regulatory networks

from multi-factorial expression data: a complete and interactive suite. BMC Genomics

22, 387.

Cassan, O., Pimpare, L.L., Dubos, C., Gojon, A., Bach, L., Lebre, S., and Martin, A. (2023). A

gene regulatory network in Arabidopsis roots reveals features and regulators of the

plant response to elevated CO(2). New Phytol **239,** 992-1004.

Cortés, A.J., and López-Hernández, F. (2021). Harnessing Crop Wild Diversity for Climate

Change Adaptation. Genes **12,** 783.

Dusenge, M.E., Duarte, A.G., and Way, D.A. (2019). Plant carbon metabolism and climate

change: elevated CO_2 and temperature impacts on photosynthesis, photorespiration

and respiration. New Phytol **221,** 32-49.

Ebi, K.L., and Loladze, I. (2019). Elevated atmospheric CO(2) concentrations and climate

change will affect our food's quality and quantity. Lancet Planet Health **3,** e283-e284.

Frachon, L., Libourel, C., Villoutreix, R., Carrere, S., Glorieux, C., Huard-Chauveau, C.,

Navascues, M., Gay, L., Vitalis, R., Baron, E., Amsellem, L., Bouchez, O., Vidal, M., Le

Corre, V., Roby, D., Bergelson, J., and Roux, F. (2017). Intermediate degrees of

synergistic pleiotropy drive adaptive evolution in ecological time. Nat Ecol Evol **1,** 1551-
1561.

Gansel, X., Munos, S., Tillard, P., and Gojon, A. (2001). Differential regulation of the NO_3^- and
479 NH4+ transporter genes *AtNrt2.1* and *AtAmt1.1* in Arabidopsis: relation with long-
480 distance and local controls by N status of the plant. Plant J **26,** 143-155.

Gao, Y., de Bang, T.C., and Schjoerring, J.K. (2019). Cisgenic overexpression of cytosolic
482 glutamine synthetase improves nitrogen utilization efficiency in barley and prevents
483 grain protein decline under elevated CO_2. Plant Biotechnology Journal **17,** 1209-1221.

Gaufichon, L., Marmagne, A., Belcram, K., Yoneyama, T., Sakakibara, Y., Hase, T., Grandjean,
485 **O., Clément, G., Citerne, S., Boutet-Mercey, S., Masclaux-Daubresse, C., Chardon, F.,**
486 **Soulay, F., Xu, X., Trassaert, M., Shakiebaei, M., Najihi, A., and Suzuki, A.** (2017).
487 ASN1-encoded asparagine synthetase in floral organs contributes to nitrogen filling in
488 Arabidopsis seeds. The Plant Journal **91,** 371-393.

Gojon, A., Cassan, O., Bach, L., Lejay, L., and Martin, A. (2023). The decline of plant mineral
490 nutrition under rising CO_2: physiological and molecular aspects of a bad deal. Trends
491 in Plant Science **28,** 185-198.

Horton, M.W., Hancock, A.M., Huang, Y.S., Toomajian, C., Atwell, S., Auton, A., Muliyati,
493 **N.W., Platt, A., Sperone, F.G., Vilhjálmsson, B.J., Nordborg, M., Borevitz, J.O., and**
494 **Bergelson, J.** (2012). Genome-wide patterns of genetic variation in worldwide
495 Arabidopsis thaliana accessions from the RegMap panel. Nature Genetics **44,** 212-216.

Hu, H., Rappel, W.-J., Occhipinti, R., Ries, A., Böhmer, M., You, L., Xiao, C., Engineer, C.B.,
Boron, W.F., and Schroeder, J.I. (2015). Distinct Cellular Locations of Carbonic
Anhydrases Mediate Carbon Dioxide Control of Stomatal Movements Plant Physiology
169, 1168-1178.

Huard-Chauveau, C., Perchepied, L., Debieu, M., Rivas, S., Kroj, T., Kars, I., Bergelson, J.,

Roux, F., and Roby, D. (2013). An atypical kinase under balancing selection confers

broad-spectrum disease resistance in Arabidopsis. PLoS Genet **9,** e1003766.

Kim, S., Plagnol, V., Hu, T.T., Toomajian, C., Clark, R.M., Ossowski, S., Ecker, J.R., Weigel, D.,

and Nordborg, M. (2007). Recombination and linkage disequilibrium in *Arabidopsis*

thaliana. Nature Genetics **39,** 1151-1155.

Lam, H.-M., Wong, P., Chan, H.-K., Yam, K.-M., Chen, L., Chow, C.-M., and Coruzzi, G.M.

(2003). Overexpression of the *ASN1* Gene Enhances Nitrogen Status in Seeds of

Arabidopsis. Plant Physiology **132,** 926-935.

Lee, S., Ricachenevsky, F.K., and Punshon, T. (2021). Functional overlap of two major

facilitator superfamily transporter, ZIF1, and ZIFL1 in zinc and iron homeostasis.

Biochemical and biophysical research communications **560,** 7-13.

Loladze, I. (2014). Hidden shift of the ionome of plants exposed to elevated CO_2 depletes

minerals at the base of human nutrition. Elife **3,** e02245.

Luthi, D., Le Floch, M., Bereiter, B., Blunier, T., Barnola, J.M., Siegenthaler, U., Raynaud, D.,

Jouzel, J., Fischer, H., Kawamura, K., and Stocker, T.F. (2008). High-resolution carbon

dioxide concentration record 650,000-800,000 years before present. Nature **453,** 379-

382.

Marcos-Barbero, E.L., Pérez, P., Martínez-Carrasco, R., Arellano, J.B., and Morcuende, R.

(2021). Screening for Higher Grain Yield and Biomass among Sixty Bread Wheat

Genotypes Grown under Elevated CO_2 and High-Temperature Conditions. Plants **10,**

1596.

Medek, D.E., Schwartz, J., and Myers, S.S. (2017). Estimated Effects of Future Atmospheric

CO_2 Concentrations on Protein Intake and the Risk of Protein Deficiency by Country

and Region. Environ Health Perspect **125,** 087002.

Myers, S.S., Zanobetti, A., Kloog, I., Huybers, P., Leakey, A.D., Bloom, A.J., Carlisle, E.,

Dietterich, L.H., Fitzgerald, G., Hasegawa, T., Holbrook, N.M., Nelson, R.L., Ottman,

M.J., Raboy, V., Sakai, H., Sartor, K.A., Schwartz, J., Seneweera, S., Tausz, M., and

Usui, Y. (2014). Increasing CO_2 threatens human nutrition. Nature **510,** 139-142.

North, K.A., Ehlting, B., Koprivova, A., Rennenberg, H., and Kopriva, S. (2009). Natural

variation in Arabidopsis adaptation to growth at low nitrogen conditions. Plant

Physiology and Biochemistry **47,** 912-918.

Pauls, S.U., Nowak, C., Balint, M., and Pfenninger, M. (2013). The impact of global climate

change on genetic diversity within populations and species. Mol Ecol **22,** 925-946.

Pearson, P.N., and Palmer, M.R. (2000). Atmospheric carbon dioxide concentrations over the

past 60 million years. Nature **406,** 695-699.

Roux, F., and Frachon, L. (2022). A Genome-Wide Association study in *Arabidopsis thaliana* to

decipher the adaptive genetics of quantitative disease resistance in a native

heterogeneous environment. PLoS One **17,** e0274561.

Shahzad, Z., and Rouached, H. (2022). Protecting plant nutrition from the effects of climate

change. Curr Biol **32,** R725-R727.

Sharma, N., Froehlich, J.E., Rillema, R., Raba, D.A., Chambers, T., Kerfeld, C.A., Kramer, D.M.,

Walker, B., and Brandizzi, F. (2023). Arabidopsis stromal carbonic anhydrases exhibit

non-overlapping roles in photosynthetic efficiency and development. Plant J 2023 **115,**

386-397.

Sun, P., Isner, J.C., Coupel-Ledru, A., Zhang, Q., Pridgeon, A.J., He, Y., Menguer, P.K., Miller,

A.J., Sanders, D., McGrath, S.P., Noothong, F., Liang, Y.K., and Hetherington, A.M.

(2022). Countering elevated CO_2 induced Fe and Zn reduction in Arabidopsis seeds.

New Phytol **235,** 1796-1806.

Tausz-Posch, S., Tausz, M., and Bourgault, M. (2020). Elevated [CO_2] effects on crops:

Advances in understanding acclimation, nitrogen dynamics and interactions with

drought and other organisms. Plant Biol (Stuttg) **22 Suppl 1,** 38-51.

Thompson, M., Gamage, D., Hirotsu, N., Martin, A., and Seneweera, S. (2017). Effects of

Elevated Carbon Dioxide on Photosynthesis and Carbon Partitioning: A Perspective on

Root Sugar Sensing and Hormonal Crosstalk. Front Physiol **8,** 578.

Umnajkitikorn, K., Sade, N., Rubio Wilhelmi, M.D.M., Gilbert, M.E., and Blumwald, E. (2020).

Silencing of OsCV (chloroplast vesiculation) maintained photorespiration and N

assimilation in rice plants grown under elevated CO(2). Plant Cell Environ **43,** 920-933.

Wang, Y., Kang, Y., Yu, W., Lyi, S.M., Choi, H.W., Xiao, E., Li, L., Klessig, D.F., and Liu, J. (2022).

AtTIP2;2 facilitates resistance to zinc toxicity via promoting zinc immobilization in the

root and limiting root-to-shoot zinc translocation in *Arabidopsis thaliana*.

Ecotoxicology and Environmental Safety **233,** 113333.

Yang, A., Li, Q., Chen, L., and Zhang, W.-H. (2020). A rice small GTPase, Rab6a, is involved in

the regulation of grain yield and iron nutrition in response to CO_2 enrichment. Journal

of Experimental Botany **71,** 5680-5688.

Zhu, C., Kobayashi, K., Loladze, I., Zhu, J., Jiang, Q., Xu, X., Liu, G., Seneweera, S., Ebi, K.L.,

Drewnowski, A., Fukagawa, N.K., and Ziska, L.H. (2018). Carbon dioxide (CO_2) levels

this century will alter the protein, micronutrients, and vitamin content of rice grains

with potential health consequences for the poorest rice-dependent countries. Sci Adv

4, eaaq1012.

Figure legends :

Figure 1: Elevated CO$_2$ negatively impacts the ionome content at the population-scale level in *Arabidopsis thaliana*. A. Representation of the experimental design used in this study. The content of eight mineral elements was assessed for around 600 *Arabidopsis thaliana* accessions coming from the REGMAP (B), LANGUEDOC (C) and TOU-A (D) populations. Each dot represents the value of the content of a mineral element for one accession (yellow: ambient CO$_2$ (aCO$_2$, ~420 ppm), blue: elevated CO$_2$ (eCO$_2$, 900 ppm). N (% of dry weight), Fe (µg.g^{-1} dry weight), Zn (µg.g^{-1} dry weight), Cu (µg.g^{-1} dry weight), Mg (µg.g^{-1} dry weight), Mn (µg.g^{-1} dry weight), Na (µg.g^{-1} dry weight), C (% of dry weight). Asterisks indicate significant differences (Paired Wilcoxson test; *, $P < 0.05$; **, $P < 0.005$; ***, $P < 0.0005$). ns; not significant.

Figure 2: Elevated CO$_2$ leads to high phenotypic diversity of ionome response in *Arabidopsis thaliana*. Distributions of the relative change (%) of the content of 8 mineral elements between elevated CO$_2$ and ambient CO$_2$, in each population (A: REGMAP, B: LANGUEDOC, C: TOU-A). Each dot represents the value of the relative change of the content a mineral element for one accession. The name of the element appears in bold if the mean of the element in elevated CO$_2$ is significantly different from the mean of the element in ambient CO$_2$ (Paired wilcoxon test, significance threshold of 0.05).

Figure 3: Elevated CO$_2$ results in a general pattern of ionome variation common to most accessions constituting natural populations of *Arabidopsis thaliana*. Principal Component Analysis (PCA) was performed using the variation of each element in response to elevated CO$_2$.

A. Natural accessions were positioned on the PCA and colored based on population. B. Contribution of each element to the PCA axis.

Figure 4: Variation in the response of the ionome to elevated CO_2 identifies contrasting subpopulations inside the REGMAP panel. K-means clustering was performed in the REGMAP accessions to identify different subpopulations. Each accession is represented by a dot, connected by a line between each element. Cluster 1:65 accessions. Cluster 2: 25 accessions. Cluster 3: 69 accessions.

Figure 5: Genetic architecture of the response of the ionome to elevated CO_2 in the REGMAP panel of *Arabidopsis thaliana*. A. Manhattan plots for the content of eight mineral elements under elevated CO_2, or for the relative change of the content of mineral elements between elevated CO_2 and ambient CO_2. For each Manhattan plot, SNPs with the 50 most significant P-value, located above the horizontal line, are colored. Bar plots showing the number of SNPs identified by GWAs for traits under elevated CO_2 (B) or for the relative change of the content of mineral elements between elevated CO_2 and ambient CO_2 (C) that are unique to one element or shared between 2 or 3 traits.

Figure 6: Identification of genes detected by GWA mapping and differentially regulated by elevated CO_2. Intersection between elevated CO_2-DEG in shoot (A) or root (B) and genes identified by GWA mapping. UpSet plots display the number of elevated CO_2-DEG that are associated to a locus identified for the content or the relative change of one or several mineral elements under elevated CO_2. Illustration of the pattern of elevated CO_2-DEG in shoot (C) or root (D) also identified by GWA mapping.

Figure 7: Natural variation of the *TIP2;2* gene is associated with improved responses of Zn content to elevated CO_2. A. Manhattan plot of the relative change of Zn content between elevated CO_2 and ambient CO_2 showing the presence of a peak closed to the *TIP2;2* locus. **B.** Comparison of haplotypes and their relative change of Zn content between elevated CO_2 and ambient CO_2. Three SNPs located at the *TIP2;2* locus are associated to an improvement of Zn content under elevated CO_2 for accessions that possess them (haplotype 1) compared to the rest of the population (haplotype 0). **C, D.** Relative expression of *TIP2;2* in the roots under ambient (**C**) or elevated (**D**) CO_2 for accessions belonging to haplotype 0 or haplotype 1. Relative expression levels were calculated based on *UBQ10* as internal control. Horizontal black line represented the median of each group of haplotypes. ***P < 0,001, **P < 0,01, unpaired Mann-Whitney test. **E.** Shoot Zn content under ambient or elevated CO_2 for WT (Columbia) and *tip2;2-1* mutant. Data are presented as the mean (with SD) of 5 and 6 biological repeats for the WT and *tip2;2-1*, respectively. **P < 0.01, unpaired Mann-Whitney test.