Negative cell cycle regulation by Calcineurin is necessary for proper beta cell regeneration in zebrafish

Laura Massoz[1*], David Bergemann[1], Arnaud Lavergne[1,2], Célia Reynders[1], Caroline Désiront[1], Chiara Goossens[1], Lydie Flasse[1], Bernard Peers[1], Marianne L. Voz[1] and Isabelle Manfroid[1]

Author Affiliations

[1]Zebrafish Development and Disease Models laboratory, GIGA-Stem Cells, University of Liège, Liège, Belgium

[2]GIGA-Genomics core facility, GIGA, University of Liège, Liège, Belgium

*Corresponding author:

Email address: laura.massoz@doct.uliege.be

Abstract

Stimulation of pancreatic beta cell regeneration could be a therapeutic lead to treat diabetes. Unlike humans, the zebrafish can efficiently regenerate beta cells, notably from ductal pancreatic progenitors. To gain insight into the molecular pathways involved in this process, we established the transcriptomic profile of the ductal cells after beta cell ablation in the adult zebrafish. These data highlighted the protein phosphatase calcineurin as a new potential modulator of beta cell regeneration. We showed that calcineurin overexpression abolished the regenerative response, leading to glycemia dysregulation. On the opposite, calcineurin inhibition increased ductal cell proliferation and subsequent beta cell regeneration. Interestingly, the enhanced proliferation of the progenitors was paradoxically coupled with their exhaustion. This suggests that the proliferating progenitors are next entering in differentiation. Calcineurin appears as a guardian which prevents an excessive progenitor proliferation to preserve the pool of progenitors. Altogether, our findings reveal calcineurin as a key player in the balance between proliferation and differentiation to enable a proper beta cell regeneration.

Introduction

Blood glucose homeostasis is tightly controlled by pancreatic endocrine cells. Insulin-producing beta cells work in close association with alpha cells, which secrete glucagon, and delta cells secreting somatostatin, to maintain normal glycemia. The insulin plays a critical role in this process as it is the only hormone able to lower the glycemia. Beta cell loss is a hallmark of type 1 and of late stages of type 2 diabetes, leading to chronic hyperglycemia. Beta cell destruction is largely irreversible in human and in mammals, making the disease incurable nowadays. Nevertheless, studies in diabetic mice models uncovered promising evidences of slight recovery of beta cells via regeneration. For example, new beta cells can arise from the replication of remaining beta cells [1]. Different models of pancreas injuries revealed the plasticity of mammalian pancreatic cells. Actually, differentiated endocrine cells such as alpha cells[2] and delta cells[3] can reprogram and convert into insulin-producing cells, a process that is age dependent. Another possible source of beta cells could be de *novo* formation from pancreatic progenitors residing in the ductal compartment of the adult pancreas [4]. Lineage tracing experiments could not clearly highlight new beta cells arising from the ductal cells in the adult [4]–[6]. However, expression of the pro-endocrine marker Neurog3 was detected in the ducts in different mouse models of pancreas regeneration [4], [7], [8]. In addition, a rare population of ductal cells expressing Neurog3 has been reported to contribute to beta cell neogenesis during diabetes [9]. A recent study of single cell RNA sequencing reveal a sub-population of human ductal cells which are

able to give rise to all the pancreatic cell types, including beta cells, following implantation in mice [10].

Altogether, these findings suggest that mammalian pancreatic ducts possess the intrinsic capacity to

(re)generate beta cells even though this process is poorly efficient and slow, especially in adults. In

contrast to mammals, zebrafish possess remarkable capacity of regeneration, independently of its age

[11]–[16]. This model can therefore be exploited to identify and characterize regenerative mechanisms

and ultimately induce regeneration in mammals. Based on regenerative mechanisms identified in

zebrafish, several studies succeed to improve regeneration in mammals, underlying the possibility of

translation from zebrafish to mammals ([17]–[21] and [22], for review).

In zebrafish, the nitroreductase (NTR)/nitroaromatic prodrug system is widely used. In this technique,

the combination of cell type-specific NTR expression and nitroaromatic prodrug exposure allow for

controlled and targeted cell ablation [23], [24]. When NTR is expressed under the control of the *insulin*

(*ins*) promoter (*Tg*(*ins:NTR-mCherry*) [25]), beta cell destruction is complete within 3 days following

the treatment in adult fish, which correlate with a peak of hyperglycemia [26], [27]. The regeneration

of beta cells upon ablation is spontaneous and fast and the glycemia is normalized within two weeks

[13], [29]. Similar to mice, new beta cells can arise through proliferation of surviving beta cells [28] as

well as through the contribution of alpha cells [29], [30] and delta cells[31], [32], underscoring the

overall conservation of these processes from zebrafish to mammals. Nevertheless, unlike mammals,

the presence of pancreatic progenitors in the ducts is well established in both larval [33] and adult

zebrafish [26], [27]. Lineage tracing experiments pointed out that ductal cells and centroacinar cells

(CACs) give rise to new beta cells [26], [33]. In adults, duct-derived beta cells start to be detected

between 7 and 10 days following beta cell ablation [26], [27].

Notch signaling is a key player regulating the differentiation of duct-associated progenitors into

endocrine cells. Larval duct cells as well as adult CACs display strong Notch activity [26], [34]. This

signaling pathway has a central role in beta cell genesis during both development [35] and

regeneration [33] by repressing endocrine differentiation. In zebrafish, different levels of Notch activity

determine the behavior of the pancreatic progenitors. While a high level of Notch activity maintains

cells in quiescence, a moderate level induces the entry in the cell cycle and proliferation whereas a low

level drives endocrine differentiation of the progenitor cells [35]. A steep decrease of Notch activity

pushes the progenitors to differentiate prematurely, bypassing the amplification step and leading to

their depletion [35]. Repression of the Notch signaling by mTor, activated by glucose and nutrients,

hence promotes beta cell formation and regeneration from ductal progenitors [33]. While Notch and

mTor signaling are crucial for this process, there is still a need to establish a global view of the

molecular mechanisms regulating beta cell regeneration.

To identify early events regulating ductal-derived beta cell regeneration, we determined the

transcriptomic signature of ductal cells from adult zebrafish following beta cell destruction. Our data

highlighted an upregulation of the Calcineurin (CaN) pathway. To elucidate CaN function in beta cell

regeneration, we both repressed and activated CaN pathway. We showed that CaN regulates beta cell

neogenesis in the ducts during regeneration, by acting on progenitor proliferation. Together, our

findings underline that CaN fine tunes the balance between progenitor proliferation and beta cell

differentiation to guarantee proper regeneration.

Results

Transcriptomic profiling of ductal cells after beta cell destruction highlights regulation of calcineurin
pathway

To gain a better understanding of the molecular mechanisms underlying the regeneration of beta cells

from the ducts, we determined the transcriptional landscape of ductal cells by RNA-sequencing after

beta cell ablation in the adult zebrafish. To selectively ablate the beta cells, we used the *Tg*(*ins:NTR-*

mCherry) transgenic fish. The ductal cells were labelled thanks to the *Tg*(*nkx6.1:GFP*) reporter line [27]

in which GFP marks the ductal tree and associated multipotent pancreatic progenitors [27]. More

precisely, 3 to 4 fishes were treated with the pro-drug MTZ at 10mM overnight, to induce ablation of

beta cells or with DMSO for the non-ablated controls. Ablation was confirmed by blood glucose

measurement before collection of the pancreas. To capture the early events triggered by the

destruction of beta cells, we generated the transcriptome of the ducts 3 days post ablation treatment

(dpt), *i.e.* before beta cell neogenesis (**Figure 1A supplemental**). Differential gene expression analysis

revealed that 1866 genes are up-regulated and 1515 genes down-regulated in the ductal cell of fish

treated with MTZ compared to control (pAdj < 0.05). According to Gene Ontology analysis, the most

enriched pathways among the up-regulated genes were DNA replication and cell cycle (**Figure 1A**). This

further corroborates our previous findings and those of others, regarding the activation of duct-

associated progenitors' proliferation in response to beta cell ablation [26], [27]. As expected, the Notch

pathway was enriched in the down-regulated genes [33], [35] (**Figure 1B**). On a contrasting note, in a

context marked by a robust proliferative response, we found it intriguing to observe an enrichment in

the cellular senescence signature (**Figure 1A**). Subsequently, we investigated deeper into the genes

associated with this specific signature. We found among them several components of the calcineurin

signaling pathway such as *nfatc3b*, *ppp3ccb* (the catalytic subunit of Calcineurin*), itpr2* and *calm3b*

(Figure 1C). In addition to these genes related to the cellular senescence signature, our transcriptomic

studies revealed the modulation of other genes from the CaN canonical pathway (**Figure 1D**),

underlying its potential role in beta cell regeneration. Calcineurin (CaN) is a highly conserved

calcium/calmodulin dependent Ser/Thr phosphatase, involved in numerous biological process

including fin regeneration and beta cell function [36]–[39]. This prompted us to investigate the role of

CaN in beta cell regeneration.

Previous RNAseq data performed in our laboratory indicate that CaN (*ppp3cca/b*) and NFATc3

(*nfatc3a/b*) are mainly express in endocrine cells [40] (**Figure 1B supplemental**), which is in accordance

with the role of CaN/NFAT signalling in beta cells [41]. CaN genes (*ppp3cca/b*) as well as *nfatc3b,* are

express at lower levels in the ducts at basal state but their expression is induced in response to beta

cell destruction (**Figure 1C supplemental**).

Calcineurin activity regulates the ductal regenerative response

To evaluate the role of CaN in beta cell regeneration, more specifically derived from ductal progenitors,

we have chosen to use young larvae, where regenerated beta cells in the pancreatic tail arise

exclusively from the ducts [33]. In response to beta cells ablation, the intrapancreatic ducts undergo a

ductal regenerative response whereby differentiation toward the endocrine fate is increased [33]. We

first determined the rate of beta cell neogenesis from the ducts in response to a single acute ablation

of beta cells as we performed in adults. We treated *Tg*(*ins:NTR-P2A-mCherry*); *Tg*(*nkx6.1:GFP*) larvae

with nifurpirinol (NFP) from 3 to 4 dpf and mCherry+ beta cells were quantified in the GFP+ ducts in

the pancreatic tail at several time points: 4, 7, 10 and 14 dpt (**Figure 1E**). Duct-associated beta cells

started to be detected in non-ablated larvae between 7 and 10 dpt (**Figure 1F**) and the number of beta

cells slowly increased until 14 dpt (**Figure 1F, 1G**). In ablated larvae, the increase became more

pronounced from 10 dpt onwards (**Figure 1F, 1G**), indicating faster endocrine differentiation. This

experiment establishes that the ductal regenerative response is detectable between 10 and 14 days

after the beta cell ablation, performed at 3 dpf. We next wanted to determine if ductal cell proliferation

is activated in response to beta cell destruction in larvae as in adult fish. We exposed *Tg(ins:NTR-P2A-*

mCherry); Tg(nkx6.1:GFP) larvae to EdU the second day following ablation (**Figure 1H**). In the ablated

larvae, the proportion of GFP+ ductal cells EdU+ (in S-phase) was higher compared to non-ablated

larvae (**Figure 1I, 1J**). This result shows that acute beta cell ablation in larvae rapidly activates ductal

cell proliferation, as previously reported in adult zebrafish.

As our transcriptomic data from adult zebrafish revealed modulation of the CaN pathway at 3 dpt

(**Figure 1D**), we treated *Tg*(*ins:NTR-P2A-mCherry*); *Tg*(*nkx6.1:GFP*) larvae from 1 to 3 dpt after beta cell

ablation, with a CaN inhibitor, the Cyclosporin A (CsA) [39] (**Figure 2A**). The number of newly formed

beta cells in the tail was monitored from 4 to 14 dpt. CsA enhanced beta cell formation at 10 dpt (**Figure**

2B-D). However, this effect appears to be transient since no discernible difference was observed

between the control and CsA-treated larvae in regeneration at the latest time point (14dpt), suggesting

an acceleration of the regenerative response (**Figure 2B, Figure 2A supplemental**). Interestingly, CsA did not affect beta cell differentiation in non-ablated larvae, indicating that CsA only acts in a regenerative context (**Figure 2B-D)**. Of note, CsA increased as well the number of regenerated beta cells in the principal islet (**Figure 2B, 2C supplemental**). We and others, previously showed that the other major source of beta cell regeneration, besides the ducts, are the bi-hormonal sst1.1+/ins+ cells [31], [32], [42]. However, CsA did not affect bi-hormonal cell formation **(Figure 2D supplemental**), suggesting that the additional cells in the principal islet could also originate from the ducts.

We next tested the effect of CsA on endocrine progenitors in a regenerative context. We induced regeneration in Tg(*neurod1:GFP*) larvae where the GFP is expressed in both endocrine progenitors and mature endocrine cells. We first assessed generation of GFP cells at different time points (**Figure 2E supplemental)** and showed that CsA induced an increase of neurod1+ cells from 4 dpt. The increase was still detectable at least until 10dpt (**Figure 2E, Figure 2F supplemental**). To determine if these additional cells result from their own proliferation, we performed a pulse of EdU just before analysis (**Figure 2E supplemental**). We observed that CsA did not affect the neurod1+ cell proliferation rate ,which is very low at these stages (**Figure 2F**). As a consequence, the effect of CsA cannot be explained by endocrine cell proliferation but rather by neogenesis from progenitors. As CsA affects pro-endocrine cells formation, we next wondered if the increased cell formation induced by CsA is specific to beta cells. Treatment with CsA was performed as previously and delta1.1 and alpha cells were detected by immunofluorescence. Interestingly, CsA did not affect alpha nor delta1.1 cells neogenesis in response to beta cell ablation (**Figure 2G-H supplemental**). Overall, these experiments showed that CsA affects specifically the beta cells and their endocrine progenitors.

CaN over-activation abolishes the regenerative response

We then wondered if an opposite regulation of CaN *i.e.* its activation impacts as well the regenerative response. To that end, we generated a transgenic line Tg(*hsp70:GFP-P2A-ppp3ccCA*) that allows ubiquitous expression of a constitutively active form of CaN^{CA} (*ppp3cc*^{CA}) upon heat shocks. Beta cell ablation was triggered in Tg(*hsp70:GFP-P2A-ppp3cc*^{CA}); Tg(*ins:NTR-P2A-mCherry*); Tg(*nkx6.1:GFP*) larvae from 3 to 4 dpf and CaN^{CA} expression was induced by four successive heat-shocks from 1 to 3 dpt (**Figure 3A**). The overexpression of CaN^{CA} after ablation impaired the regenerative response at 14 dpt (**Figure 3B-C**). Similar results were obtained with Tg(*UAS:GFP-P2A-ppp3cc*^{CA}); Tg(*cftr:gal4*)[45] ; Tg(*ins:NTR-P2A-mCherry*) larvae in which CaN^{CA} is continuously and specifically overexpressed in the ducts within the pancreas (**Figure 3D-E**). Importantly, the structure of the ducts was similar in CaN^{CA} overexpressing larvae compared to Tg(*nkx6.1:GFP*) controls (**Figure 3D**), suggesting that the suppression of the regenerative response in CaN^{CA}-overexpressing larvae was not due to

morphogenetic defects during ductal growth. This result shows that CaN acts directly in the ducts to decrease beta cell regeneration while it is not necessary for normal beta cell differentiation.

CaN regulates beta cell differentiation induced by Notch inhibition in absence of regeneration

Our transcriptomic data showed that the Notch pathway is downregulated in ductal cells during beta cell regeneration (**Figure 1**). As the level of Notch activity determines the behavior of ductal cells [33] from quiescence to proliferation and subsequently to beta cell differentiation, we tested whether calcineurin acts together with the Notch pathway on a common pool of ductal progenitors. To inhibit the Notch pathway, we treated larvae with several concentrations of the gamma-secretase inhibitor LY411575 from 3 to 4dpf in absence of regeneration. The activity of CaN was inhibited by CsA during the same timeframe (**Figure 4A**). As previously, we used reporter lines for beta and ductal cells *Tg*(*ins:NTR-P2A-mCherry*); *Tg*(*nkx6.1:GFP*) and the number of secondary beta cells was analyzed at 6dpf (**Figure 4A**). As expected, the number of beta cells progressively rose as the concentration of the Notch inhibitor increased (**Figure 4B**). Combined treatment with CsA fostered the differentiation of beta cells between 1 and 10 µM LY411575 but did not result in further increase at 15 µM LY411575 (**Figure 4B**), suggesting that CaN is important within a permissive window of Notch activity. Since we overserved the highest synergistic effect at 5µM of LY411575, we used this concentration for the following experiments (**Figure 4C-D**). It is worth noting that combined treatment of LY411575 (5uM) and another CaN inhibitor, FK506, resulted in the same synergistic increase of beta cell differentiation (**Figure 4A-B supplemental**), confirming that the effect is due to CaN inhibition. The combined effect of Notch and CaN inhibition is transient as it is not observed at 7dpf anymore (**Figure 4E**). Therefore, as observed in regenerative conditions (**Figure 2B**), CaN inhibition accelerates beta cell neogenesis induced by Notch repression.

To determine to which extent CaN pathway can modulate Notch induced beta cell neogenesis, we activated CaN^{CA} overexpression in *Tg*(*hsp70:GFP-P2A-ppp3cc*^{CA}); *Tg*(*ins:NTR-P2A-mCherry*) larvae by an heat-shock at 3 dpf and treated them with LY411575 (**Figure 4C supplemental)**. CaN^{CA} overexpression resulted in a lowered beta cell formation induced by Notch inhibition (**Figure 4D supplemental**), revealing that CaN activation counterbalanced the effects of Notch inhibition. Using our previous settings of Notch and CaN inhibition, we next wondered if the canonical pathway downstream of CaN was involved in the enhancement of beta cell differentiation. To activate NFAT, a well-known target of CaN, we used CHIR99021 allowing a stabilization of the active form of NFAT (**Figure 1D**). We found that CHIR99021 rescued the effect of CsA (**Figure 4E supplemental**), suggesting that CaN inhibition increases beta cell neogenesis at least partially by the regulation of NFAT. Overall, these results reveal that CaN impacts beta cell formation in pro-endocrinogenic context, such as induced by a low level of

Notch activity. Moreover, it suggests that both CaN and Notch pathways act on a common pool of
ductal progenitors to govern beta cell neogenesis.

CaN controls the proliferation of duct-associated progenitors induced by Notch inhibition

Given that we observed the most significant increase in beta cell formation with CsA when Notch
activity was mildly repressed (**Figure 4B-D**), and since mild Notch activity has been shown to promote
progenitor amplification [35], it suggests that CaN acts at this level. To explore this possibility, we
exposed briefly *Tg*(*nkx6.1:GFP*) larvae to EdU after mild Notch (LY411575 5µM) and CaN inhibition
(**Figure 5A**) and analyzed at 4dpf and 6dpf. As expected, Notch inhibition increased the proportion of
proliferating EdU+ GFP+ duct cells at 4dpf (**Figure 5B-C**) and decreased the amount of ductal
progenitors 2 days later (at 6dpf) (**Figure 5D**), which is concomitant with the increase of beta cell
differentiation (**Figure 4C-D**). Interestingly, the combined inhibition with CsA further increased these
proportions, while CaN inhibition alone had no effect (**Figure 5B-D**). Of note, we detected beta cells
EdU+ at 6dpf (**Figure 5A supplemental**), underlying that these cells originate from the proliferation of
ductal progenitors.

Incidentally, at 4dpf, while the proliferation is increased (**Figure 5C**), the number of ductal cells
remained the same in all conditions (**Figure 5B supplemental**), suggesting that ductal cells have not
yet left the cell cycle to differentiate. In comparison, after stronger Notch inhibition (15µM), the ductal
cells are already depleted at 4dpf (**Figure 5C supplemental**), as they directly differentiate without
entering the cell cycle [35]. In these conditions, CsA could therefore not enhance ductal progenitor
proliferation and thus beta cell formation (**Figure 4B**). These results show that CaN and Notch
pathways act together on the proliferation of the ductal progenitors to prevent their exhaustion.

CaN prevents exhaustion of Notch responsive progenitors during beta cell regeneration

Taken together, our results indicate that CaN plays a role in the proliferation of ductal progenitors in
contexts that are permissive for beta cell differentiation. To demonstrate that CaN acts on ductal
progenitor proliferation in a similar manner to Notch inhibition but during regeneration, we exposed
Tg(ins:NTR-P2A-mCherry); *Tg(nkx6.1:GFP)* larvae after ablation and CsA treatment (**Figure 5E**). CaN
inhibition enhanced the proliferation of ductal cells in ablated larvae (**Figure 5F-G**). To next determine
if CaN acts on the Notch-responsive progenitors in beta cell regeneration, we then used
Tg(tp1:VenusPest) Notch reporter line. At larval stages the vast majority of ductal cells are Notch

responsive and the reporter line marks all the progenitors within the ductal tree [34]. We treated

Tg(ins:NTR-P2A-mCherry); *Tg(tp1:VenusPest)* as described above (**Figure 5E**). In ablated-larvae, CsA

increased tp1+ ductal cell proliferation (**Figure 5H-I**). Moreover, CsA induced a reduction of tp1+ ductal

cells in ablated larvae (**Figure 5 H-J**), suggesting an exhaustion of the Notch responsive progenitors, in

accordance with premature beta cell differentiation we observed (**Figure 2B-4D**). Lastly, CsA does not

affect tp1+ cells in non-ablated larvae showing that CaN inhibition did not directly affect Notch

signaling (**Figure 5J**). Those results suggest that CaN fine tunes the balance between proliferation of

the progenitors and their differentiation to prevent their exhaustion during beta cell regeneration.

CaN regulation is functionally relevant in adult zebrafish

To further expand upon our findings and investigate their relevance in a non-developmental context,

we next ought to determine whether CaN function is maintained in older zebrafish. We used 2 months

old *Tg*(*ins:NTR-P2A-mCherry*); *Tg*(*nkx6.1:GFP*) juveniles fish to perform beta cell ablation followed by

CsA treatments. We analyzed the number of small islets (up to 5 cells) at 7 and 10dpt. At 7dpt, CsA

increased the number of small islets in ablated juveniles (**Figure 6A-B**) showing that CaN inhibition

enhances beta cell regeneration in juvenile zebrafish, as in larvae. It is noteworthy that, as in larvae,

we highlighted an acceleration of beta cell regeneration. The increase number of small islets is indeed

transient as it not observe anymore at 10dpt (**Figure 6C**).

Next, we determined the functional impact of CaN overexpression or inhibition by assessing the

glycemia at 7, 10 and 14 after beta cell ablation. Overexpression of CaN^{CA} using either *Tg(hsp70:GFP-*

P2A-ppp3cc^{CA}) or *Tg*(*cftr:gal4*); *Tg*(*UAS:GFP-P2A-ppp3cc^{CA}*)*, led to an increased glycemia at both 7 and

10dpt (**Figure 6D-E**)*, before recovery at 14 dpt (**Figure 6A supplemental**). This indicates that the

overexpression of CaN delayed glycemia recovery induced by beta cell regeneration. However, CaN

inhibition did not seem to further improve the glycemia (**Figure 6D-E**) probably because the glycemia

was already low at 7dpt (77 mg/dL on average) compared to non-ablated control (50 mg/dl on average)

(**Figure 6D-E**). Altogether, these results show that in adult zebrafish also, CaN regulation is necessary

to enable beta cell regeneration and for proper recovery of the glycemia after beta cell loss.

Discussion

The present study reveals the importance of the protein phosphatase CaN as a new player in beta cell

regeneration in zebrafish. We demonstrate that CaN acts on ductal-associated progenitor cells by

balancing proliferation and endocrine differentiation. In addition, we show the interplay between CaN

and Notch signalling, a master regulator of beta cell regeneration.

Previous drug and genetic screening using zebrafish larvae enabled the identification of several

regulators of beta cell regeneration from different pancreatic cellular sources. For example, adenosine

has been shown to stimulate beta cell replication [28], igfbp1a [43] and TGFb suppression [30] promote

alpha-to-beta cell transdifferentiation. As for cdk5 inhibition and folinic acid/Folr1, they promote beta

cell regeneration from the pancreatic ducts [44], [45]. Here, to identify novel regulators of beta cell

regeneration specifically from pancreatic ducts, we carried out a transcriptomic profiling of duct cells

following beta cell ablation in the adult zebrafish. Transcriptomic analyses show that the regulated

genes encompass most of the genes and pathways identified in previous studies (igfbp1, mTor, Notch,

etc.), underlying the importance of those actors in beta cell regeneration. Our data reveal also that

DNA replication is the most enriched signature attesting that duct cells undergo a potent proliferative

response after the destruction of beta cells.

Besides these expected signatures, our transcriptomic data uncover the unanticipated upregulation of

numerous genes implicated in DNA repair and cell cycle arrest. These signatures might indicate that

highly proliferating ductal cells activate counteracting mechanisms. Among the genes regulated in

those signatures, we focused on CaN and determined its role in beta cell regeneration.

Pharmacological inhibition of CaN increases the proliferation of duct cells induced by beta cell ablation,

resulting in an acceleration of beta cell regeneration in the ducts. Consistently, transgene-mediated

CaN overactivation abolishes the regenerative response. Importantly, the inhibition of regeneration is

observed when CaN is overexpressed either ubiquitously or selectively in *cftr*-expressing ductal cells

indicating that the role of CaN in beta cell regeneration is intrinsic to the ducts. Based on functional

assays in larvae, we not only confirm the activation of the proliferation of ductal cells soon after beta

cell ablation, but also that the rate of progenitor proliferation is carefully controlled by CaN in order to

achieve proper and timely regeneration of beta cells. Our data are consistent with earlier studies

reporting a role of CaN in proliferation dynamics during fin regeneration. In the regenerating fin, low

CaN activity is found in the proximal region of the blastema characterized by a high rate of proliferation

and regeneration and its activity increases distally where lower proliferation is observed [36], [38],

[39]. It was suggested that CaN control blastemal cell progeny divisions [36]. In human, the importance

of CaN in proliferation is also highlighted in organ transplanted patients. When patients are treated

with Cyclosporin A (*i.e.* the CaN inhibitor we used in this study) as immunosuppressive drug they

indeed present an increased risk of skin cancer, notably due to keratinocyte senescence inhibition [46].

It has been shown that Notch inhibitory treatments switch progenitors from proliferative self-renewing

to premature differentiation, leading to progenitor depletion [34], [35]. Our study reveals that this

phenomenon is further exacerbated by CaN inhibition. Importantly, during normal larval development

in absence of Notch inhibitory treatment, CaN does not affect basal ductal proliferation nor beta cell

differentiation. Hence, Notch signalling has to be repressed to detect the effect of CaN on the

progenitors, suggesting that CaN acts downstream of Notch pathway. In differentiating keratinocytes,

CaN cooperates with Notch signalling to regulate p21/*cdkn1a* (which is upregulated in the ducts at 3

dpt), cell cycle withdrawal and differentiation [47]. These studies show that CaN acts in association

with Notch signalling on progenitors proliferation and on their differentiation.

Based on our data, we build the model depicted in **Figure 7A**. Our study suggests that CaN acts in

competent progenitor cells and that this competence is determined by Notch signalling. When Notch

is repressed to a mild level, the progenitor enter into the cell cycle and acquire a pro-endocrinogenic

competence [35]. CaN acts on these progenitors to tone down an excessive proliferation and avoid the

exhaustion of these progenitors. CaN is therefore a guardian of the progenitor population. CaN

inhibition both increases progenitor proliferation and induces their depletion, suggesting a switch to a

symmetric division resulting in two daughter cells entering in endocrine differentiation.

More precisely, a previous study in mice uncover the existence of three division modes of the

pancreatic progenitors during embryonic development, *i.e.* symmetric self-renewing resulting in two

progenitors cells ; asymmetric resulting in a progenitor and a endocrine cell and symmetric

differentiative resulting in two differentiated cells [48]. The authors actually show that the type of

division is defined by the timing of induction of endocrine program by NEUROG3 [48]. Interestingly,

NEUROG3 seems to be the link between proliferation and differentiation, and its expression is

regulated by Notch signalling [49]. In zebrafish, endocrine differentiation is not induced by *Neurog3*

but by *Ascl1b* and *Neurod1* [50]. Concerning that subject, CaN inhibition accelerates the formation of

neurod1+ cells. Hence CaN could possibly act via the determination of the type of division *i.e.*

symmetric vs asymmetric. This model is supported by previous observation in others systems. In stem

cells and neuronal and hematopoietic progenitors, premature differentiation results from a switch in

the mode of cellular division, from symmetric amplifying division to asymmetric differentiating division

[51], [52]. Notch determines the choice between both types of divisions [53], [54].

Calcineurin is known to be implicated in cellular senescence [46]. Usually thought as negative

regulators of development and cellular growth, DNA repair [55] and cellular senescence [56] appear

to be required in both developmental and regenerative processes. Our transcriptomic data suggest

that these mechanisms are required for beta cell regeneration. Therefore, it would be interesting to

determine the contribution of these cellular processes in the ductal progenitors and determine if CaN

acts via cellular senescence in this case.

Overall, this study brings new insights on beta cell regeneration and highlights the ductal progenitor

cell cycle as a cornerstone in the process. Some studies report an increase of proliferation of some

ductal cell population in diabetic patients [10], [57], implying a regenerative response. However, these

ductal cells cannot efficiently reform the beta cell mass, suggesting a dormant mechanism of

regeneration. As such, the balance between proliferation and induction of endocrine differentiation

could be a key to improve beta cell neogenesis. However, as CaN is also important for beta cell

function, this approach would require to be transient to induce neogenesis. Therefore, it should be

combined with methods to induce beta cell proliferation to ultimately reconstitute the beta cell mass.

Overall, this study brings a better understanding on the regulation of the balance between ductal

progenitors proliferation and endocrine differentiation. These results should provide new hints to help

improve regenerative competences in mammals.

Material and Methods

Key resources table

Reagent type or resource	Designation	Source or reference	Identifier	Additional information
Genetic reagent (Danio rerio)	*TgBAC(nkx6.1:eGFP)^ulg004*	PMID: 26329351	ZFIN: ZDB-ALT-160205-1	
Genetic reagent (Danio rerio)	*Tg(ins:NTR-P2A-mCherry)^ulg034*	PMID: 29663654	ZFIN: ZDB-ALT-171122-9	
Genetic reagent (Danio rerio)	*Tg(cftr:gal4)*	PMID: 25592226	ZFIN : ZDB-FISH-150901-25442	
Genetic reagent (Danio rerio)	*Tg(tp1:VenusPest)*	PMID: 22492351	ZFIN: ZDB-FISH-150901-8023	
Genetic reagent (Danio rerio)	*Tg(hsp70:eGFP-P2A-ppp3ccaCA)^ulg068*	This paper		See Zebrafish husbandry and generation of the *Tg(hsp70:eGFP-P2A-ppp3ccaCA) zebrafish line*
Genetic reagent (Danio rerio)	*Tg(UAS:eGFP-P2A-ppp3ccaCA)^ulg069*	This paper		See Zebrafish husbandry and generation of the *Tg(UAS:eGFP-P2A-ppp3ccaCA) zebrafish line*
Antibody	Anti-GFP (chicken polyclonal)	Aves Labs	GFP-1020	1:1000

Antibody	anti-mCherry/dsRed (Living Colors Polyclonal)	Clontech	632496	1:500
Antibody	Anti-glucagon (mouse polyclonal)	Sigma	G2654	1:300
Antibody	Goat anti-Chicken IgY (H+L), Alexa Fluor™ 488	Invitrogen	A-11039	1:750
Antibody	Goat anti-dsred 568	Invitrogen		1:750
Antibody	Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 633	Invitrogen		1:750
Chemical compound	Nifurpirinol (NFP)	Sigma-Aldrich	32439	
Chemical compound	Metronidazole (MTZ)	Sigma-Aldrich	M1547	
Chemical compound	Cyclosporine A (CsA)	Selleckchem	S2286	
Chemical compound	LY411575	Sigma-Aldrich	SML0506	
Chemical compound	CHIR990211	Sellekchem	CT99021	
Commercial assay or kit	Gateway™ LR Clonase™ II Enzyme mix	Invitrogen	11791020	
Commercial assay or kit	Gateway™ BP Clonase™ II Enzyme mix	Invitrogen	11789020	
Sequence based reagent	IM369	This paper	PCR primer	gaagaaaaccccggtcctatgtc gacgaaagagccgaaag
Sequence based reagent	IM380	This paper	PCR primer	ccttacacattcccgtcagtgc
Sequence based reagent	IM371	This paper	PCR primer	CGGCTCTTTCGTCGACAT AGGACCGGGGTTTTCTTC CACG
Sequence based reagent	O226	This paper	PCR primer	GCCACCATGGTGAGCAA GGGCGAGGA
Sequence based reagent	IM370	This paper	PCR primer	ttattagatcttatttctgatcacc tcctt
Sequence based reagent	IM459	This paper	PCR primer	cacacgaattcgccgccaccAT GGTGAGCAAGGGCGAG
Sequence based reagent	IM460	This paper	PCR primer	ggatcggtcgagatccttacGA TCTTATTTCTGATCACCTC CTTACG
Sequence based reagent	IM457	This paper	PCR primer	GTAAGGATCTCGACCGAT CCTG
Sequence based reagent	IM458	This paper	PCR primer	GGTGGCGGCGAATTCGT G

Commercial assay or kit	Nextera® XT DNA Library kit	Illumina	FC-131-1024	
Commercial assay or kit	Click-iT™ EdU Cell Proliferation Kit for Imaging, Alexa Fluor™ 647 dye	Invitrogen	C10340	
Software, algorithm	Imaris	Bitplane (http://www.bitplane.com/imaris/imaris)	RRID:SCR _007370	Version 9.5
Software, algorithm	GraphPad Prism	GraphPad Prism (https://graphpad.com)	RRID:SCR _015807	Version 8
Software, algorithm	DESeq2	DESeq2 (https://bioconductor.org/packages/release/bioc/html/DESeq2.html)	RRID:SCR _015687	
Software, algorithm	WebGestalt	WebGestalt (http://www.webgestalt.org/)	RRID:SCR _006786	

Zebrafish husbandry and generation of the Tg(hsp70:eGFP-P2A-ppp3ccaCA)[ulg068] *and*

Tg(UAS:eGFP-P2A-ppp2ccaCA)[ulg069] *zebrafish lines*

Tg BAC(nkx6.1:eGFP)[ulg004] [58] *;   Tg(ins:NTR-P2A-mCherry)*[ulg034] [59] *; Tg(cftr:gal4)* and
Tg(tp1:VenusPest) were used. Zebrafish were raised in standard conditions at 28°C. All experiments

were carried out in compliance with the European Union and Belgian law and with the approval of the

ULiège Ethical Committee for experiments with laboratory animals (approval number : 2075).

The *hsp70:GFP-P2A-ppp3ccaCA* transgene has been generated by cloning a PCR fragment containing

the Gateway vector pCR8/GW/TOPO. Firstly, we amplified the full length of ppp3ccaCA with primers

IM369/IM380 and amplified *GFP-P2A* with overlapping regions with IM371/O226. The overlapping PCR

used the primers O226/IM380. Then to obtain a truncated *ppp3ccaCA* lacking the calmodulin biding

and the autoinhibitory domain, resulting in a constantly active form of calcineurin, we amplified the

last fragments with IM370/O226 and cloned into PCR8 vector. The promoter was assembled by LR

recombination with pE5-hsp70 into pDestTol2p2A from the Tol2kit [60]. *Tg (hsp70:GFP-P2A-*
ppp3ccaCA) fish have been generated using the Tol2 mediated transgenesis [61]. *The Tg(UAS:GFP-*

385 *P2A-ppp3ccacA*) has been generated by ligation (KLD kit, NEB) of PCR fragments *GFP-P2A-ppp3ccacA*

386 *(*IM459/IM460) with the UAS sequences (IM457/IM458) in plasmid from (Distel 2009) and then

387 inserted into pDestTol2p2A from the Tol2kit. Final constructions has been injected with transposase

388 into wild type (WT) AB embryos.

Beta-cell ablation

Adults fish for RNA-sequencing experiment were treated with freshly prepared metronidazole (MTZ)

(Sigma M1547) at 10mM with 0.2% DMSO in fish water. Control treatments consisted of fish water

containing 0.2% DMSO. Fish were treated for 18 hours in the dark. Nifurpirinol (NFP) (32439, Sigma-

Aldrich) stock solution was dissolved at 2.5 mM in DMSO. b-cell ablation in *Tg(nkx6.1:eGFP);*

Tg(ins:NTR-P2A-mCherry) larvae was induced by treatment with 2.5 µM NFP in E3. Control treatments

consisted of E3 containing 0.2% DMSO. Larvae were treated for 18 hours in the dark.

Drug treatments

Cyclosporine A (Selleckchem, S2286), CHIR99021 and LY411575 (Sigma-Aldrich, SML0506) stock

solution were dissolved at 10mM in DMSO. Larvae treatment were respectively performed at 0.1 µM

and 5µM in E3. Control treatments consisted of E3 containing the same amount of DMSO than drug

treatment. Larvae were treated for 18 hours in the dark.

5-ethynyl-2'-deoxyuridine (EdU) incorporation assay

Zebrafish larvae were incubated in 4 mM EdU dissolved in E3 water for 8 hours, they were then directly

euthanised in tricaine and fixed in 4% PFA. EdU was detected according to the protocol of Click-iT™

EdU Cell Proliferation Kit for Imaging, Alexa Fluor™ 647 (ThermoFisher C10340) after whole mount

immunodetection.

Heat Shock

Successive heat shocks of 30 minutes and 12 hours apart were performed at 39°C for larvae and 37°C

for juveniles and adults zebrafish.

Whole mount immunodetection

Larvae were euthanized in tricaine and fixed in 4% PFA at 4 °C for IHC. The digestive tract of juveniles

was dissected prior immunodetection and kept in methanol for at least 18 hours. After depigmentation

with 3% H2O2/1% KOH, larvae were permeabilised 30 min in PBS/ Triton X-100 and incubated for two

hours in blocking buffer (4% goat serum/1% BSA/PBS/0.1% Triton X-100). Primary and secondary

antibodies were incubated at 4 °C overnight.

	Fixation duration	Depigmentation duration	Permeabilization solution	Permeabilization duration
5-10 dpf	18 hours	15 min	PBTr 0.05%	30 min
13-17 dpf	36 hours	20 min	PBTr 2%	30 min
2months (digestive tract)	18 hours	15 min	/	/

Primary antibodies: Living Colors Polyclonal anti-mCherry/dsRed (rabbit, 1:500, Clontech 632496),

anti-GFP (chicken, 1:1000),

Secondary antibodies: Alexa Fluor-488, -568, -633 (goat, 1:750, Molecular Probes).

Flow cytometry and FACS

The whole pancreas from 3-4 fish of *Tg(nkx6.1:eGFP); Tg(ins:NTR-P2A-mCherry)* adult fish (6–10

months old, males and females) were dissected, collected and washed in HBSS without Ca^{2+}/Mg^{2+}. Live

cell dissociation was performed in Tryple Select 1x solution (GIBCO) supplemented with 100 U/mL

collagenase IV (Life Technologies 17104-019) and 40 µg/mL proteinase K (Invitrogen, 25530031) for 10

min at 28 °C.

The GFP+ cells were selected on FACS Aria III and sorted under purity mode and after exclusion of the

doublets. The purity of the sorted cells was confirmed by epifluorescence microscopy (~95 %). Cells

(about 1000-5000/fish depending on the cell type) were immediately lysed with 0.5% Triton X-100

containing 2U/µl RNAse inhibitor and stored at –80 °C.

mRNA sequencing of FACSed cells and bioinformatic analyses

cDNAs were prepared from lysed cells according to SMART-Seq2.0 (Picelli et al., 2014) for low input

RNA sequencing and libraries were prepared with Nextera® DNA Library kit (Illumina). Independent

biological replicates of each cell type sequenced using Illumina NextSeq500 and obtained ~20 million

100 bp paired-end reads. Reads were mapped and aligned to the zebrafish genome GRCz11 from

Ensembl gene annotation version using STAR version 2.6.1 (Dobin et al., 2013). Gene expression levels

were calculated with featureCounts (http://bioinf.wehi.edu.au/featureCounts/) and differential

expression determined with DESeq2 (Love, Huber, & Anders, 2014). Expression values are given as

normalized read counts. Poorly expressed genes with mean normalized expression counts <10 were

excluded from the subsequent analyses. DESeq2 uses Wald test for significance with posterior

adjustment of P values (Padj) using Benjamini and Hochberg multiple testing. The differentially

expressed (DE) genes identified with a Padj cutoff of 0.05 were submitted for GO analysis using

WebGestalt tool (Liao, Wang, Jaehnig, Shi, &Zhang, 2019).

Confocal microscopy and image analysis

Images were acquired using Leica SP5 confocal microscope. We used ImageJ to count the cells and

Imaris to do the pictures.

Glycemia measurement

Glycemia measurement were performed as described in [59].

Data availability

The sequences that support the findings are of this study have been deposit in Geo with the accessions

codes GSE212124, https://www.ncbi.nlm.nih.gov/geo/info/linking.html. The authors declare that all

other data supporting the findings of this study are available within the paper and its supplementary

information files.

Authors contribution

L.M. conducted experiments, acquired and analysed data, analysed transcriptomic data, wrote the

manuscript. **D.B.** conducted RNA-seq experiment and revised the manuscript **A.L.** implemented

bioinformatic programs and analysed transcriptomic data. **C.R., C.G. and C.D.** conducted experiments.

L.F. revised the manuscript. **B.P.** provided resources and advised on research studies. **M.L.V.** provided

resources, advised on research studies and revised the manuscript **I.M.** supervised the whole work,

acquired funding, designed the research studies, verified data and assisted in writing the manuscript.

Acknowledgments

The authors thank the EOS (Excellence of Sciences) program from FNRS and more especially Dr.

Emmanuel Dejardin for support. We also thank the GIGA technology platforms, GIGA-Genomics and

GIGA-Imaging. The authors also thanks Jérémie Zappia for critically reading the manuscript, Jordane

Bourdouxhe and Marie Alice Dupont for their help with the constructions.

Duality of interest

No potential conflicts of interest.

References

[1] Y. Dor, J. Brown, O. I. Martinez, and D. a Melton, "Adult pancreatic beta-cells are formed by

self-duplication rather than stem-cell differentiation.," *Nature*, vol. 429, no. model 3, pp. 41–

46, 2004, doi: 10.1038/nature02520.

[2] F. Thorel *et al.*, "Conversion of Adult Pancreatic alpha-cells to Beta-cells after Extreme Beta

Cell Loss," *Nature*, vol. 464, no. 7292, pp. 1149–1154, 2010, doi:

10.1038/nature08894.Conversion.

[3] S. Chera *et al.*, "Diabetes recovery by age-dependant conversion of pancreatic delta-cells into

insulin producers," *Nature*, vol. 514, no. 7523, pp. 503–507, 2014, doi:

10.1038/nature13633.Diabetes.

[4]   X. Xu *et al.*, "Beta Cells Can Be Generated from Endogenous Progenitors in Injured Adult

Mouse Pancreas," *Cell*, vol. 132, pp. 197–207, 2008, doi: 10.1016/j.cell.2007.12.015.

[5]   M. Solar *et al.*, "Pancreatic Exocrine Duct Cells Give Rise to Insulin-Producing β Cells during

Embryogenesis but Not after Birth," *Dev. Cell*, vol. 17, pp. 849–860, 2009, doi:

10.1016/j.devcel.2009.11.003.

[6]   H. Zhao *et al.*, "Pre-existing beta cells but not progenitors contribute to new beta cells in the

adult pancreas," *Nat. Metab.*, vol. 3, no. 3, pp. 352–365, 2021, doi: 10.1038/s42255-021-

00364-0.

[7]   J. L. Kopp *et al.*, "Sox9+ ductal cells are multipotent progenitors throughout development but

do not produce new endocrine cells in the normal or injured adult pancreas," *Development*,

vol. 138, pp. 653–665, 2011, doi: 138/4/653 [pii].

[8]   M. Courtney *et al.*, "The Inactivation of Arx in Pancreatic α-Cells Triggers Their Neogenesis and

Conversion into Functional β-Like Cells," *PLoS Genet.*, vol. 9, no. 10, pp. 1–18, 2013, doi:

10.1371/journal.pgen.1003934.

[9]   C. Gribben *et al.*, "Ductal Ngn3-expressing progenitors contribute to adult β cell neogenesis in

the pancreas," *Cell Stem Cell*, vol. 28, no. 11, pp. 2000-2008.e4, 2021, doi:

10.1016/j.stem.2021.08.003.

[10]   M. M. F. Qadir *et al.*, "Single-cell resolution analysis of the human pancreatic ductal

progenitor cell niche," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 117, no. 20, pp. 10876–10887, 2020,

doi: 10.1073/pnas.1918314117.

[11]   J. B. Moss, P. Koustubhan, M. Greenman, M. J. Parsons, I. Walter, and L. G. Moss,

"Regeneration of the pancreas in adult zebrafish," *Diabetes*, vol. 58, no. August, pp. 1844–

1851, 2009, doi: 10.2337/db08-0628.

[12]   K. D. Poss, L. G. Wilson, and M. T. Keating, "Heart regeneration in zebrafish," *Science (80-.).*,

vol. 298, no. 5601, pp. 2188–2190, 2002, doi: 10.1126/science.1077857.

[13]   T. Y. Choi, N. Ninov, D. Y. R. Stainier, and D. Shin, "Extensive conversion of hepatic biliary

epithelial cells to hepatocytes after near total loss of hepatocytes in zebrafish,"

Gastroenterology, vol. 146, no. 3, pp. 776–788, 2014, doi: 10.1053/j.gastro.2013.10.019.

[14]   Y. Goldshmit, T. E. Sztal, P. R. Jusuf, T. E. Hall, M. Nguyen-Chi, and P. D. Currie, "Fgf-dependent

glial cell bridges facilitate spinal cord regeneration in zebrafish.," *J. Neurosci.*, vol. 32, no. 22, pp. 7477–7492, May 2012, doi: 10.1523/JNEUROSCI.0758-12.2012.

[15] V. Kroehne, D. Freudenreich, S. Hans, J. Kaslin, and M. Brand, "Regeneration of the adult zebrafish brain from neurogenic radial glia-type progenitors," *Development*, vol. 138, no. 22, pp. 4831–4841, 2011, doi: 10.1242/dev.072587.

[16] M. Gemberling, T. J. Bailey, D. R. Hyde, and K. D. Poss, "The zebrafish as a model for complex tissue regeneration," *Trends Genet.*, vol. 29, no. 11, pp. 611–620, 2013, doi: 10.1016/j.tig.2013.07.003.

[17] C. Papadimitriou *et al.*, "3D Culture Method for Alzheimer's Disease Modeling Reveals Interleukin-4 Rescues Aβ42-Induced Loss of Human Neural Stem Cell Plasticity," *Dev. Cell*, vol. 46, no. 1, pp. 85-101.e8, 2018, doi: 10.1016/j.devcel.2018.06.005.

[18] V. Mashkaryan *et al.*, "Type 1 Interleukin-4 Signaling Obliterates Mouse Astroglia in vivo but Not in vitro," *Front. Cell Dev. Biol.*, vol. 8, no. February, pp. 1–16, 2020, doi: 10.3389/fcell.2020.00114.

[19] S. Ko, T. Choi, J. O. Russell, J. So, and D. Shin, "Bromodomain and extraterminal (BET) proteins regulate biliary-driven liver regeneration," *J. Hepatol.*, vol. 64, no. 2, pp. 316–325, 2016, doi: 10.1016/j.jhep.2015.10.017.Bromodomain.

[20] Y. Goldshmit *et al.*, "Decreased anti-regenerative effects after spinal cord injury in spry4-/- mice.," *Neuroscience*, vol. 287, pp. 104–112, Feb. 2015, doi: 10.1016/j.neuroscience.2014.12.020.

[21] B. V. Fausett, J. D. Gumerson, and D. Goldman, "The proneural basic helix-loop-helix gene Ascl1a is required for retina regeneration," *J. Neurosci.*, vol. 28, no. 5, pp. 1109–1117, 2008, doi: 10.1523/JNEUROSCI.4853-07.2008.

[22] L. Massoz, M. A. Dupont, and I. Manfroid, "Zebra-fishing for regenerative awakening in mammals," *Biomedicines*, vol. 9, no. 1, pp. 1–18, 2021, doi: 10.3390/biomedicines9010065.

[23] H. Pisharath, "Validation of nitroreductase, a prodrug-activating enzyme, mediated cell death in embryonic zebrafish (Danio rerio)," *Comp. Med.*, vol. 57, no. 3, pp. 241–246, 2007.

[24] S. Curado, R. M. Anderson, B. Jungblut, J. Mumm, E. Schroeter, and D. Y. R. Stainier, "Conditional targeted cell ablation in zebrafish: A new tool for regeneration studies," *Dev. Dyn.*, vol. 236, no. February, pp. 1025–1035, 2007, doi: 10.1002/dvdy.21100.

[25] H. Pisharath, J. M. Rhee, M. a Swanson, S. D. Leach, and M. J. Parsons, "Targeted ablation of
beta cells in the embryonic zebrafish pancreas using E.coli nitroreductase," *Mech Dev.*, 2007,
doi: 10.1016/j.mod.2006.11.005.Targeted.

[26] F. Delaspre *et al.*, "Centroacinar cells are progenitors that contribute to endocrine pancreas
regeneration," *Diabetes*, vol. 64, no. October, pp. 3499–3509, 2015, doi: 10.2337/db15-0153.

[27] A. P. Ghaye *et al.*, "Progenitor potential of nkx6.1-expressing cells throughout zebrafish life
and during beta cell regeneration," *BMC Biol.*, vol. 13, p. 70, 2015, doi: 10.1186/s12915-015-
0179-4.

[28] O. Andersson *et al.*, "Adenosine signaling promotes regeneration of pancreatic β- cells in
vivo," *Cell Metab.*, 2012, doi: 10.1016/j.cmet.2012.04.018.Adenosine.

[29] L. Ye, M. A. Robertson, D. Hesselson, D. Y. R. Stainier, and R. M. Anderson, "Glucagon is
essential for alpha cell transdifferentiation and beta cell neogenesis," *Dev.*, vol. 142, no. 8, pp.
1407–1417, 2015, doi: 10.1242/dev.117911.

[30] C. S. M. Helker *et al.*, "A whole organism small molecule screen identifies novel regulators of
pancreatic endocrine development," *Dev.*, vol. 146, no. 14, 2019, doi: 10.1242/dev.172569.

[31] C. Carril Pardo *et al.*, "A delta-cell subpopulation with a pro-beta-cell identity contributes to
efficient age-independent recovery in a zebrafish model of diabetes," *Elife*, 2022.

[32] S. P. Singh *et al.*, "A single-cell atlas of de novo β-cell regeneration reveals the contribution of
hybrid β/δ-cells to diabetes recovery in zebrafish," *Development*, vol. 149, no. 2, 2022, doi:
10.1242/dev.199853.

[33] N. Ninov, D. Hesselson, P. Gut, A. Zhou, K. Fidelin, and D. Y. R. Stainier, "Metabolic regulation
of cellular plasticity in the pancreas," *Curr. Biol.*, vol. 23, no. 13, pp. 1242–1250, 2013, doi:
10.1016/j.cub.2013.05.037.

[34] M. J. Parsons *et al.*, "Notch-responsive cells initiate the secondary transition in larval zebrafish
pancreas," *Mech. Dev.*, vol. 126, no. 10, pp. 898–912, 2009, doi: 10.1016/j.mod.2009.07.002.

[35] N. Ninov, M. Borius, and D. Y. R. Stainier, "Different levels of Notch signaling regulate
quiescence, renewal and differentiation in pancreatic endocrine progenitors," *Development*,
vol. 139, no. 9, pp. 1557–1567, 2012, doi: 10.1242/dev.076000.

[36] V. A. Tornini *et al.*, "Live Monitoring of Blastemal Cell Contributions During Appendage
Regeneration," *Curr. Biol.*, 2016, doi: 10.1016/j.cub.2016.08.072.Live.

[37]   S. C. McMillan *et al.*, "A regulatory pathway involving retinoic acid and calcineurin demarcates
567   and maintains joint cells and osteoblasts in regenerating fin," *Dev.*, vol. 145, no. 11, 2018, doi:
568   10.1242/DEV.161158.

569   [38]   Z. Cao *et al.*, "Calcineurin controls proximodistal blastema polarity in zebrafish fin
570   regeneration," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 118, no. 2, pp. 2–9, 2021, doi:
571   10.1073/pnas.2009539118.

572   [39]   S. Kujawski *et al.*, "Calcineurin Regulates Coordinated Outgrowth of Zebrafish Regenerating
573   Fins," *Dev. Cell*, vol. 28, no. 5, pp. 573–587, 2014, doi: 10.1016/j.devcel.2014.01.019.

574   [40]   E. Tarifeño-Saldivia *et al.*, "Transcriptome analysis of pancreatic cells across distant species
575   highlights novel important regulator genes," *BMC Biol.*, vol. 15, no. 1, pp. 1–19, 2017, doi:
576   10.1186/s12915-017-0362-x.

577   [41]   J. J. Heit *et al.*, "Calcineurin/NFAT signalling regulates pancreatic β-cell growth and function,"
578   *Nature*, vol. 443, no. 7109, pp. 345–349, 2006, doi: 10.1038/nature05097.

579   [42]   J. Mi, K. C. Liu, and O. Andersson, "Decoding pancreatic endocrine cell differentiation and β
580   cell regeneration in zebrafish," *Sci. Adv.*, vol. 9, no. 33, p. eadf5142, 2023, doi:
581   10.1126/sciadv.adf5142.

582   [43]   J. Lu *et al.*, " IGFBP 1 increases β-cell regeneration by promoting α- to β-cell
583   transdifferentiation ," *EMBO J.*, vol. 35, no. 18, pp. 2026–2044, 2016, doi:
584   10.15252/embj.201592903.

585   [44]   C. Karampelias *et al.*, "Reinforcing one-carbon metabolism via folic acid/Folr1 promotes β-cell
586   differentiation," *Nat. Commun.*, vol. 12, no. 1, pp. 1–13, 2021, doi: 10.1038/s41467-021-
587   23673-0.

588   [45]   K. C. Liu *et al.*, "Inhibition of Cdk5 promotes β-cell differentiation from ductal progenitors,"
589   *Diabetes*, vol. 67, no. 1, pp. 58–70, 2017, doi: 10.2337/db16-1587.

590   [46]   X. Wu *et al.*, "Calcineurin and ATF3: opposite roles in squamous skin cancer," *Nature*, 2010,
591   doi: 10.1038/nature08996.Calcineurin.

592   [47]   C. Mammucari *et al.*, "Integration of notch 1 and calcineurin/NFAT signaling pathways in
593   keratinocyte growth and differentiation control," *Dev. Cell*, vol. 8, no. 5, pp. 665–676, 2005,
594   doi: 10.1016/j.devcel.2005.02.016.

[48]   Y. H. Kim, H. L. Larsen, P. Rué, L. A. Lemaire, J. Ferrer, and A. Grapin-Botton, "Cell Cycle–

Dependent Differentiation Dynamics Balances Growth and Endocrine Differentiation in the

Pancreas," *PLoS Biol.*, vol. 13, no. 3, 2015, doi: 10.1371/journal.pbio.1002111.

[49]   N. A. J. Krentz *et al.*, "Phosphorylation of NEUROG3 Links Endocrine Differentiation to the Cell

Cycle in Pancreatic Article Phosphorylation of NEUROG3 Links Endocrine Differentiation to the

Cell Cycle in Pancreatic Progenitors," *Dev. Cell*, vol. 41, no. 2, pp. 129-142.e6, 2017, doi:

10.1016/j.devcel.2017.02.006.

[50]   L. C. Flasse, D. G. Stern, J. L. Pirson, I. Manfroid, B. Peers, and M. L. Voz, "The bHLH

transcription factor Ascl1a is essential for the specification of the intestinal secretory cells and

mediates Notch signaling in the zebrafish intestine," *Dev. Biol.*, vol. 376, no. 2, pp. 187–197,

2013, doi: 10.1016/j.ydbio.2013.01.011.

[51]   W. B. Huttner and Y. Kosodo, "Symmetric versus asymmetric cell division during neurogenesis

in the developing vertebrate central nervous system," *Curr. Opin. Cell Biol.*, vol. 17, no. 6, pp.

648–657, 2005, doi: 10.1016/j.ceb.2005.10.005.

[52]   A. D. Ho and W. Wagner, "The beauty of asymmetry: Asymmetric divisions and self-renewal in

the haematopoietic system," *Curr. Opin. Hematol.*, vol. 14, no. 4, pp. 330–336, 2007, doi:

10.1097/MOH.0b013e3281900f12.

[53]   R. S. Bultje, D. R. Castaneda-Castellanos, L. Y. Jan, Y. N. Jan, A. R. Kriegstein, and S. H. Shi,

"Mammalian Par3 Regulates Progenitor Cell Asymmetric Division via Notch Signaling in the

Developing Neocortex," *Neuron*, vol. 63, no. 2, pp. 189–202, 2009, doi:

10.1016/j.neuron.2009.07.004.

[54]   M. Guo, L. Y. Jan, and Y. N. Jan, "Control of daughter cell fates during asymmetric division:

Interaction of Numb and Notch," *Neuron*, vol. 17, no. 1, pp. 27–41, 1996, doi: 10.1016/S0896-

6273(00)80278-0.

[55]   K. Sousounis *et al.*, "Eya2 promotes cell cycle progression by regulating DNA damage response

during vertebrate limb regeneration," 2020.

[56]   S. Da *et al.*, "Cell senescence contributes to tissue regeneration in zebrafish," no. July 2019,

pp. 1–5, 2020, doi: 10.1111/acel.13052.

[57]   A. S. Md Moin, P. C. Butler, and A. E. Butler, "Increased proliferation of the pancreatic duct

gland compartment in type 1 diabetes," *J. Clin. Endocrinol. Metab.*, vol. 102, no. 1, pp. 200–

209, 2017, doi: 10.1210/jc.2016-3001.

[58]   A. P. Ghaye *et al.*, "Progenitor potential of nkx6.1-expressing cells throughout zebrafish life

and during beta cell regeneration," *BMC Biol.*, vol. 13, no. 1, 2015, doi: 10.1186/s12915-015-
0179-4.

[59] D. Bergemann *et al.*, "Nifurpirinol: A more potent and reliable substrate compared to
metronidazole for nitroreductase-mediated cell ablations," *Wound Repair Regen.*, vol. 26, no.
2, pp. 238–244, 2018, doi: 10.1111/wrr.12633.

[60] K. M. Kwan *et al.*, "The Tol2kit: a multisite gateway-based construction kit for Tol2 transposon
transgenesis constructs.," *Dev. Dyn. an Off. Publ. Am. Assoc. Anat.*, vol. 236, no. 11, pp.
3088–3099, Nov. 2007, doi: 10.1002/dvdy.21343.

[61] K. Kawakami, "Tol2: a versatile gene transfer vector in vertebrates.," *Genome Biol.*, vol. 8
Suppl 1, p. S7, 2007, doi: 10.1186/gb-2007-8-s1-s7.

Legends

<u>Figure 1 :</u> *Transcriptomic profiling of ductal cells during beta cell regeneration and validation in larvae*

A-B) Enrichment ratio of selected non-redundant signatures of KEGG pathways overrepresented
in ductal cells after beta cells ablation (UP – A and DOWN -B) compared to ductal cells without
beta cells ablation. Gene Ontology (GO) terms were identified using ORA analysis by WebGestalt
(Liao et al., 2019) using the list of DE genes provided by DESeq. The light color for Notch pathway
means p-value = 0.11.

C) List of genes associated with the signature of cellular senescence from A-B. Genes related to
CaN pathway are in bold.

D) Calcineurin canonical pathway with up-regulated genes in green and down-regulated genes in
red in transcriptomic data from A-B.

E) Experimental design for regeneration test in larvae. Briefly, after nifurpirinol treatment from 3
to 4dpf, larvae were fixed and analyzed at 4-7-10 and 14 days post treatment (dpt).

F) Graph representing the mean number of mCherry+ beta cells in the pancreatic tail of *Tg(ins:NTR-
P2A-mCherry) ; Tg(nkx6.1:GFP)* at 0-4-7-10 and 14 dpt. The gray spheres represent non-ablated
conditions and the pink triangles the ablated condition. Data are presented as mean values ± SEM.
One-way ANOVA test with Tukey's multiple comparison test, ****p-value<0.0005.

G) Whole mount fluorescent immunohistochemistry (GFP and mCherry) of the pancreas of
Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP) larvae at 14dpt. 3D projection (stack) of one non-
ablated and one ablated representative samples. The principal islet (PI) and the pancreatic tail are
shown. Arrows point out mCherry+ beta cells in the pancreatic tail. Scale 100µM.

H) Experimental design for EdU assay in larvae. After NFP treatment for 3 to 4dpf, larvae were
exposed to EdU at 2dpt before fixation for analysis.

I) Whole mount fluorescent immunohistochemistry (GFP and EdU) of the pancreatic tail of *Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP)* larvae at 2dpt. 3D projection (stack) of one non-ablated and one ablated representative samples. Arrows point out GFP+ duct cells EdU+ in the pancreatic tail. Scale 50µM.

J) Barplot representing the percentage of GFP+ ductal cells which incorporated EdU+ in non-ablated and ablated conditions. Data are presented as mean values ± SD. T-test. **p-value<0.005.

Figure 2 : *Calcineurin inhibition with CsA increases the ductal regenerative response*

A) Experimental design for regeneration test in larvae with CsA treatment. Briefly, after nifurpirinol treatment from 3 to 4dpf, larvae were treated with CsA from 1 to 3dpt and fixed and analyzed at 4-7-10 and 14 days post treatment (dpt).

B) Graph representing the mean number of mCherry+ beta cells in the pancreatic tail of *Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP)* at 0-4-7-10 and 14 dpt. The gray spheres represent non-ablated condition ; the pink triangles represent the ablated condition ; the black squares CsA condition and inverted green triangles ablated + CsA condition. Data are presented as mean values ± SEM. Two-way ANOVA test with Sidak's multiple comparisons test, *p-value<0.05.

C) Whole mount fluorescent immunohistochemistry (GFP and mCherry) of the pancreas of *Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP)* larvae at 10dpt. 3D projection (stack) of non-ablated and ablated larvae treated with DMSO or CsA representative samples. The principal islet (PI) and the pancreatic tail are shown. Arrows point out mCherry+ beta cells in the pancreatic tail. Scale 100µM.

D) Barplot representing the number of number of mCherry+ beta cells in the pancreatic tail of *Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP)* larvae at 10dpt. The gray spheres represent non-ablated condition ; the pink triangles represent the ablated condition ; the black squares CsA condition and inverted green triangles ablated + CsA condition. Data are presented as mean values ± SD. Two-way ANOVA with Tukey's multiple comparison test, *p-value<0.05.

E) Graph representing the mean number of GFP+ neurod1+ cells in the pancreatic tail of *Tg(ins:NTR-P2A-mCherry) ; Tg(neurod1:GFP)* at 0-4-7 and 10 dpt. The gray spheres represent non-ablated condition ; the pink triangles represent the ablated condition ; the black squares CsA condition and inverted green triangles ablated + CsA condition. Data are presented as mean values ± SEM. Two-way ANOVA test with Sidak's multiple comparisons test, **p-value<0.005; *** p-value<0.0005; **** p-value<0.00005.

F) Graph representing the mean number of GFP+ neurod1 EdU+ cells in the pancreatic tail of *Tg(ins:NTR-P2A-mCherry) ; Tg(neurod1:GFP)* at 0-4-7 and 10 dpt. The gray spheres represent non-ablated condition ; the pink triangles represent the ablated condition ; the black squares CsA condition and inverted green triangles ablated + CsA condition. Data are presented as mean values ± SEM.

Figure 3: *Transgenic mediated overexpression of calcineurin abolish the ductal regenerative response*

A) Experimental design for regeneration test in larvae with heat-shocks. Briefly, after nifurpirinol treatment from 3 to 4dpf, four heat shock were performed from 1 to 3dpt and larvae were fixed and analyzed at 14dpt.

B) Whole mount fluorescent immunohistochemistry (GFP and mCherry) of the pancreas of *Tg(hsp70:CaN^{CA})*; *Tg(ins:NTR-P2A-mCherry); Tg(nkx6.1:GFP)* larvae at 14dpt. 3D projection (stack) of one non-ablated and one ablated with or without heat-shock representative samples. The principal islet (PI) and the pancreatic tail are showed. Arrows point out mCherry+ beta cells in the pancreatic tail. Scale 100µM.

C) Barplot representing the mean number of mCherry+ beta cells in the pancreatic tail of *Tg(hsp70:CaN^{CA})*; *Tg(ins:NTR-P2A-mCherry); Tg(nkx6.1:GFP)* larvae at 14dpt. The gray spheres represent non-ablated condition ; the pink triangles represent the ablated condition ; the black squares heat-shock condition and inverted green triangles ablated + heat-shock condition. Data are presented as mean values ± SD. Two-way ANOVA with Tukey's multiple comparisons test, *p-value<0.05, *** p-value<0.0005, ns = non-significant.

D) Whole mount fluorescent immunohistochemistry (GFP and mCherry) of the pancreas of larvae at 14dpt. 3D projection (stack) of *Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP)* one non-ablated and one ablated representative control samples and *Tg(UAS:CaN^{CA})*; *Tg(ins:NTR-P2A-mCherry) ; Tg(cftr:gal4)* one non-ablated and one ablated representative samples. The principal islet (PI) and the pancreatic tail are showed. Arrows point out mCherry+ beta cells in the pancreatic tail. Scale 100µM.

E) Barplot representing the mean number of mCherry+ beta cells in the pancreatic tail of larvae at 14dpt. The gray spheres represent non-ablated *Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP)* condition ; the pink triangles represent the ablated *Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP)* condition ; the black squares non-ablated *Tg(UAS:CaN^{CA})*; *Tg(ins:NTR-P2A-mCherry) ; Tg(cftr:gal4)* condition and inverted green triangles ablated *Tg(UAS:CaN^{CA})*; *Tg(ins:NTR-P2A-mCherry) ; Tg(cftr:gal4)* condition. Data are presented as mean values ± SD. Two-way ANOVA with Tukey's multiple comparison test, *** means p-value<0.0005, **** p-value<0.00005, ns = non-significant.

Figure 4: *CaN repression potentializes the effect of Notch inhibition on beta cell formation*

A) Experimental design for Notch inhibition test in non-ablated condition. Larvae were treated concomitantly with LY411575 (Notch inhibitor) and CsA from 3 to 4dpf and were fixed and analyzed at 6dpf.

B) Graph representing the mean number of mCherry+ beta cells in the pancreatic tail of *Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP)* larvae at 6 dpf depending the concentration of LY411575. The blue dots represent LY411575; and purple combination of LY411575 and CsA. Data are presented as mean values ± SEM. Two-way ANOVA test with Sidak's multiple comparison test, *p-value<0.05, **p-value<0.05, ns = non-significant.

C) Whole mount fluorescent immunohistochemistry (GFP and mCherry) of the pancreas of *Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP)* larvae at 6dpf. 3D projection (stack) of one control (without any treatment) ; one CsA-treated ; one LY411757-treated and one with both CsA and LY411575 treated larvae. The principal islet (PI) and the pancreatic tail are showed. Arrows point out mCherry+ beta cells in the pancreatic tail. Scale 50µM.

D) Barplot representing the mean number of mCherry+ beta cells in the pancreatic tail of *Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP)* larvae at 6dpf. The black dots represent the control; gray CsA treatment ;blue LY411575; and purple combination of LY411575 and CsA. Data are presented as mean values ± SD. Two-way ANOVA with Tukey multiple comparison test, ***p-value<0.0005, ****p-value<0.00005.

E) Graph representing the mean number of mCherry+ beta cells in the pancreatic tail of *Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP)* larvae at 5-6-7 dpf. The black dots represent the control; gray CsA treatment ;blue LY411575; and purple combination of LY411575 and CsA.

Data are presented as mean values ± SEM. Two-way ANOVA test with Sidak's multiple comparison test, *p-value<0.05.

Figure 5: *CaN repression increases the proportion of ductal proliferating cells*

A) Experimental design for EdU assay in Notch test. Larvae were treated concomitantly with LY411575 (Notch inhibitor) and CsA from 3 to 4dpf and then briefly treated with EdU before fixation and analysis at 4 dpf or at 6dpf.

B) Whole mount fluorescent immunohistochemistry (GFP and EdU) of the pancreatic tail of *Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP)* larvae at 4dpf. 3D projection (stack) of one control (without any treatment), one with CsA only, one with LY411575 only and one with both CsA and LY411757 representative samples. Scale 50μM.

C) Barplot representing the percentage of GFP+ ductal cells which incorporated EdU+ in pancreatic tail of *Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP)* larvae for the Notch test. The black dots represent the control; gray CsA treatment ;blue LY411575; and purple combination of LY411575 and CsA. Data are presented as mean values ± SD. T-test. Two-way ANOVA test with Tukey's multiple comparisons test, *p-value<0.05 ; ****p-value<0.00005 ; ns = non-significant.

D) Barplot representing the number of GFP+ ductal cells which in pancreatic tail of *Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP)* larvae at 6dpf for the Notch test. The black dots represent the control; gray CsA treatment ;blue LY411575; and purple combination of LY411575 and CsA. Data are presented as mean values ± SD. T-test. Two-way ANOVA test with Tukey's multiple comparisons test, *p-value<0.05 ; ****p-value<0.00005 ; ns = non-significant.

E) Experimental design for EdU assay in regeneration. Larvae were treated with nifurpirinol for beta cell ablation from 3 to 4dpf then with CsA from 4 to 5dpf and then briefly treated with EdU before fixation and analysis.

F) Whole mount fluorescent immunohistochemistry (GFP and EdU) of the pancreatic tail of *Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP)* larvae at 5dpf. 3D projection (stack) of one representative sample of non-ablated or ablated with or without CsA are shown. Scale 50μM.

G) Barplot representing the percentage of GFP+ ductal cells which incorporated EdU+ in pancreatic tail of *Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP)* larvae at 5dpf. The gray spheres represent non-ablated condition ; the pink triangles the ablated condition ; the black squares CsA condition and inverted green triangles ablated + CsA condition. Data are presented as mean values ± SD. Two-way ANOVA test with Tukey's multiple comparisons test, *p-value<0.05 ; ****p-value<0.00005 ; ns = non-significant.

H) Whole mount fluorescent immunohistochemistry (VenusPest and EdU) of the pancreatic tail of *Tg(ins:NTR-P2A-mCherry) ; Tg(tp1:VenusPest)* larvae at 5dpf. 3D projection (stack) of one representative sample of non-ablated or ablated with or without CsA are shown. Scale 50μM.

I) Barplot representing the percentage of GFP+ ductal cells which incorporated EdU+ in pancreatic tail of *Tg(ins:NTR-P2A-mCherry) ; Tg(tp1:VenusPest)* larvae at 5dpf. The gray spheres represent non-ablated condition ; the pink triangles the ablated condition ; the black squares CsA condition and inverted green triangles ablated + CsA condition. Data are presented as mean values ± sd. Two-way ANOVA test with Tukey multiple comparisons test, *p-value<0.05 ; ****p-value<0.00005 ; ns means non-significant.

J) Barplot representing the number of VenusPest+ ductal cells which incorporated EdU+ in pancreatic tail of *Tg(ins:NTR-P2A-mCherry) ; Tg(tp1:VenusPest)* larvae at 5dpf. The gray spheres represent non-ablated condition ; the pink triangles the ablated condition ; the black squares CsA condition and inverted green triangles ablated + CsA condition. Data are

presented as mean values ± SD. Two-way ANOVA test with Tukey's multiple comparisons test,
**p-value<0.005.

Figure 6: *CaN regulation is important in juveniles/adults and necessary for correct glycemia recovery*

A) Whole mount fluorescent immunohistochemistry (GFP and mCherry) of the pancreas of
Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP) 2 months old zebrafish at 7dpt. 3D projection (stack)
of non-ablated and ablated larvae treated with DMSO or CsA representative samples. The
principal islet (PI) and the pancreatic tail are shown. Zoom from section of the pancreatic tail
are shown, arrows point out mCherry+ beta cells in these zoom. Scale 200µM

B-C) Barplot representing the number of number of mCherry+ small secondary islets (=<5 cells) in
the pancreatic tail of *Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP)* 2 months old zebrafish at 7 **(B)** and
10dpt **(C).** The gray spheres represent non-ablated condition ; the pink triangles represent the
ablated condition ; the black squares CsA condition and inverted green triangles ablated + CsA
condition. Data are presented as mean values ± SD. Two-way ANOVA with Tukey's multiple
comparison test, *p-value<0.05.

D-E) Barplot representing the glycemia (mg/dL) of *Tg(ins:NTR-P2A-mCherry) ;* months old adult
zebrafish at 7 **(D)** and 10dpt **(E).** The pink triangles represent the ablated condition ; the inverted
green triangles ablated + CsA condition; the blue squares *Tg(hsp70:CaN^{CA})* after heat shocks ; the
orange lozenges *Tg(UAS:CaN^{CA})*; *Tg(cftr:gal4)*. The grey line represents the mean glycemia of
controls (non-ablated) fish. Data are presented as mean values ± SD. One-way ANOVA with Tukey's
multiple comparison test, *p-value<0.05; **p-value<0.005.

Figure 7: *Model of CaN action on ductal progenitors to regenerate beta cells*

A) Under physiological conditions, the behavior of the ductal progenitors is determined by
Notch signaling. Calcineurin is active in these progenitors and enable a proper control
between proliferation and differentiation. When CaN in repressed, more ductal
progenitors enter in the cell cycle (2dpt) and switch to a mode of proliferation leading to
differentiation of the two daughter cells (4dpt), as more pro-endocrine cells are formed.
The result is a exhaustion of the progenitors and a premature beta cell differentiation
(10dpt).

Figure 1 supplemental : *Transcriptomic profiling of ductal cells during beta cell regeneration and*
validation in larvae

A) Design of the experimental pipeline for transcriptomic experiment

B) Expression of *ppp3cca; ppp3ccb ; nfatc3a ; nfatc3b* in acinar, alpha, beta, delta or ductal cells
population from the zebrafish pancreas [40].

C) Calcineurin (*ppp3ccb and ppp3cca)* and NFATc3 (*nfatc3a and nfatc3b)* expression in ductal cells
from zebrafish in non-ablated and ablated conditions.

Figure 2 supplemental : *Calcineurin inhibition with CsA increases the ductal regenerative response*

A) Whole mount fluorescent immunohistochemistry (mCherry) of the pancreas of *Tg(ins:NTR-P2A-mCherry)* larvae at 4-7 and 14dpt. 3D projection (stack) of ablated larvae treated with DMSO or CsA representative samples. The principal islet (PI) and the pancreatic tail are shown. Scale 50 or 100μM.

B-C) Barplot representing the number of mCherry+ beta cells in the principal islet (PI) **(B)** and the number of secondary islets of mCherry+ beta cells in the pancreatic tail **(C)** of Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP) larvae at 10dpt. Data are presented as mean values ± SD. T-test or Two-way ANOVA with Tukey's multiple comparison test. **p-value<0.005

D) Barplot representing the number of sst1.1+ mCherry+ bi-hormonal cells in the principal islet (PI) of Tg(ins:NTR-P2A-mCherry) ; Tg(sst1.1:GFP) larvae at 6dpt. Data are presented as mean values ± SD. T-test.

E) Experimental design for EdU assay in larvae. After NFP treatment for 3 to 4dpf, larvae were exposed to EdU before fixation for analysis.

F) Whole mount fluorescent immunohistochemistry (GFP and EdU) of the pancreas of Tg(ins:NTR-P2A-mCherry) ; Tg(neurod1:GFP) larvae at 2-4-7 and 10 dpt. 3D projection (stack) of non-ablated and ablated larvae treated with DMSO or CsA representative samples. The principal islet (PI) and the pancreatic tail are shown. Scale 50μM.

G-H) Barplot representing the number of gcg+ alpha cells **(G)**; the number of GFP+ sst1.1 delta cells **(H)**; of Tg(*ins:NTR-P2A-mCherry*) ; Tg(*sst1.1:GFP*) larvae at 10dpt. Gcg was detected by IHC. The gray spheres represent non-ablated condition ; the pink triangles represent the ablated condition ; the black squares CsA condition and inverted green triangles ablated + CsA condition. Data are presented as mean values ± SD. Two-way ANOVA with Tukey's multiple comparison test. **p-value<0.005

Figure 4 supplemental : *CaN repression potentializes the effect of Notch inhibition on beta cell formation*

A) Experimental design for Notch inhibition test in non-ablated condition. Larvae were treated concomitantly with LY411575 (Notch inhibitor) and FK506 from 3 to 4dpf and were fixed and analyzed at 5dpf.

B) Barplot representing the mean number of mCherry+ beta cells in the pancreatic tail of *Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP)* larvae at 5dpf. The black dots represent the control; gray FK506 treatment ;blue LY411575; and purple combination of LY411575 and FK506. Data are presented as mean values ± SD. Two-way ANOVA with Tukey multiple comparison test, *p-value<0.05,**p-value<0.005, ****p-value<0.00005.

C) Experimental design for Notch inhibition test in non-ablated condition. Larvae were heat-shocked and then directly treated with LY411575 (Notch inhibitor) from 3 to 4dpf and were fixed and analyzed at 6dpf.

D) Barplot representing the mean number of mCherry+ beta cells in the pancreatic tail of *Tg(hsp70:CaN^{CA})*; *Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP)* larvae at 5dpf. The black dots represent the control; gray heat-shock ;blue LY411575; and purple combination of LY411575 and heat-shock. Data are presented as mean values ± SD. Two-way ANOVA with Tukey multiple comparison test, ****p-value<0.00005.

E) Barplot representing the mean number of mCherry+ beta cells in the pancreatic tail of *Tg(ins:NTR-P2A-mCherry) ; Tg(nkx6.1:GFP)* larvae at 6dpf after Notch inhibition (LY411575),

884   CaN inhibition (CsA) and NFATc inhibition (CHIR99021). Data are presented as mean values ±
885   SD. Two-way ANOVA with Tukey multiple comparison test, *p-value<0.05,**p-value<0.005,
886   ****p-value<0.00005.

888   **Figure 5 supplemental:** *CaN repression increases the proportion of duct proliferating cells*

889   **A)** Z-plan pictures of Tg (nkx6.1:GFP); Tg (ins:NTR*-mCherry) larvae that were treated at 3dpf with
890   both CsA 1µM and LY411575 5µM. At 4dpf, the larvae were exposed to EdU 4mM during 8
891   hours, before analysis at 6 dpf.

B-C) Barplot representing the number of GFP+ ductal cells which in pancreatic tail of *Tg(ins:NTR-*
P2A-mCherry) ; Tg(nkx6.1:GFP) larvae at 4dpf for the Notch test. **(B)** Mild Notch inhibition with
LY411575 5µM and **(C)** stronger Notch inhibition with LY411575 15µM. The black dots represent
the control; gray CsA treatment ;blue LY411575; and purple combination of LY411575 and CsA.
Data are presented as mean values ± SD. T-test. Two-way ANOVA test with Tukey's multiple
comparisons test, *p-value<0.05 ; ****p-value<0.00005 ; ns = non-significant.

Figure 6 supplemental: *CaN regulation is important in juveniles/adults and necessary for correct*
glycemia recovery

A) Barplot representing the glycemia (mg/dL) of *Tg(ins:NTR-P2A-mCherry) ;* months old adult
901   zebrafish at 14dpt**.** The pink triangles represent the ablated condition ; the inverted green
902   triangles ablated + CsA condition; the blue squares *Tg(hsp70:CaN^{CA})* after heat shocks. The grey
903   line represents the mean glycemia of controls (non-ablated) fish. Data are presented as mean
904   values ± SD.
905   B) Barplot representing the glycemia (mg/dL) of *Tg(ins:NTR-P2A-mCherry) ; and Tg(UAS:CaN^{CA})*;
906   *Tg(cftr:gal4)* in non-ablated adult zebrafish. Data are presented as mean values ± SD.

Figure 5

Figure 4

A

B

Tg(nkx6.1:GFP) ; *Tg(ins:NTR-P2A-mCherry)* 6dpf

Figure 1

A UP-REGULATED GENES

B DOWN-REGULATED GENES

C chek1;rbl1;hipk2;ccne1;ccna2;hipk1a;nbn;cdk1;cdkn1a;mybl2b;**nfatc3b**;ccnb1;e2f1;ccne2;tgfb3;ccnd2a;mapk1;foxm1;cdk2;zfp36l1a;ccnb2;**ppp3ccb;itpr2**;mcu;rb1;ccnb3;rela;e2f3;**calm3b**;chek2;si:ch211-160f23.5;mapk13

D

E

F

G *Tg(nkx6.1:GFP)* ; *Tg(ins:NTR-P2A-mCherry)* 14 dpt

H

I *Tg(nkx6.1:GFP)* ; *Tg(ins:NTR-P2A-mCherry)* 2dpt

J

Figure 2

Figure 3

Figure 6

Figure 7

Legend