Broaden the application of *Yarrowia Lipolytica* synthetic biology tools to explore the potential of Yarrowia clade biodiversity

Young-Kyoung Park[1] and Tristan Rossignol[1]

[1]Université Paris-Saclay, INRAE, AgroParisTech, Micalis Institute, 78350, Jouy-en-Josas, France.

Abstract

Yeasts have established themselves as prominent microbial cell factories, and the availability of synthetic biology tools has led to breakthroughs in the rapid development of industrial chassis strains. The selection of a suitable microbial host is critical in metabolic engineering applications, but it has been largely limited to a few well-defined strains. However, there is growing consideration for evaluating strain diversity, as a wide range of specific traits and phenotypes have been reported even within a specific yeast genus or species. Moreover, with the advent of synthetic biology tools, non-type strains can now be easily and swiftly reshaped.

The yeast *Yarrowia lipolytica* has been extensively studied for various applications such as fuels, chemicals, and food. Additionally, other members of the Yarrowia clade are currently being evaluated for their industrial potential. In this study, we demonstrate the versatility of synthetic biology tools originally developed for *Y. lipolytica* by repurposing them for engineering other yeasts belonging to the Yarrowia clade. Leveraging the GoldenGate *Y. lipolytica* tool kit, we successfully expressed fluorescent proteins as well as the carotenoid pathway in at least six members of the clade, serving as proof of concept.

This research lays the foundation for conducting more comprehensive investigations into the under-characterized strains within the Yarrowia clade and exploring their potential applications in biotechnology.

Introduction

Bio-based products through the microbial fermentation process to replace petroleum-based
chemical production are extensively studied as an essential way to develop sustainable and
environmentally friendly bioproduction. A key point for microbial cell factory success is the choice of
the chassis microorganism for its metabolite production and its metabolic rewiring capacity as well as
the availability of synthetic biology tools. Yeasts are particularly attractive eukaryotic production
hosts with their capacity to grow on a broad variety of renewable carbon sources, to produce
complex target molecules, and their well-established scale-up process.

Apart from *Saccharomyces cerevisiae*, non-conventional yeasts have been investigated due to their
particular industrial traits. Among them, *Yarrowia lipolytica* has been extensively used and studied
for many applications from fuels and chemicals to foods. This yeast is genetically amenable, can grow
on a large range of substrates, has the GRAS status, and can produce high yields of various
biomolecules, in particular lipids and organic acids, (Park and Ledesma-Amaro, 2023). Target
molecules produced by this yeast have been recently expanded to plant-based derivative products
like terpenes (Zhang, et al., 2022). This continuously growing interest in using this yeast as an
industrial chassishas led to an exponential advancements in the development of synthetic biology
tools including fast and modular cloning systems and CRISPR/Cas9 methods (Larroude, et al., 2018).

Yi et *al.* recently reviewed the importance of considering strain diversity for establishing microbial
cell factories by highlighting the fact that a wide range of strain variations already exists even within
a specific yeast genus and species (Yi and Alper, 2022). An illustrative example has been provided by
a primary screening of the potential producer of triacetic acid lactone by expressing a 2-pyrone
synthase in 13 industrial *S. cerevisiae* strains with different genetic background (Saunders, et al.,
2015). The production levels varied up to 63-fold among different strains, highlighting the
importance of considering the strain variation for optimal microbial cell factories. Moreover, basal
production is not always indicative of the rewiring potential (Yi and Alper, 2022), and phenotypes and
physiology differ for non-native metabolites. Therefore, heterologous expression must be screened
in various backgrounds to evaluate the best strains/species for further genetic engineering. This
emphasizes the growing importance of considering strain-level characteristics when selecting host
microorganism. However, this requires extensive study of variants or extensive screening with the
possible requirements of genetic tools adaptations to each strain.

Most studies focus on a limited set of industrial or type strains and rarely extend to other strains or
related species, limiting the knowledge on the physiologic and metabolic state of such species or
strains. Nevertheless, for some traits of interest like lipid production, physiological data of clade or
set of strains have been explored. The diversity of closely related species in the Yarrowia clade has

been recently investigated through comparative studies of their oleaginous properties (Kurtzman,
2011). These species, *Y. lipolytica, Yarrowia alimentaria, Yarrowia deformans, Yarrowia galli, Candida*
hispaniensis, Yarrowia hollandica, Yarrowia oslonensis, Yarrowia phangngensis and *Yarrowia*
yakushimensis exhibited diverse salt stress tolerance, optimal temperature for growth (Groenewald,
et al., 2014) and maximum lipid content, with significant differences ranging from 30% of cell dry
weight in *C. oslonensis* to 67% in *C. hispaniensis* (Michely, et al., 2013). In the past decade, the clade
has grown up to 15 potential species including now *Yarrowia keelungensis, Yarrowia divulgata,*
Yarrowia porcina, Yarrowia bubula, Yarrowia brassicae and *Yarrowia parophonii,* (Quarterman, et
al., 2017, Chang, et al., 2013, Péter, et al., 2019). Quarterman et *al.* investigated the biomass and
lipid production of 13 species of the Yarrowia clade by using a nondetoxified acid-pretreated
switchgrass hydrolysate as a feedstock (Quarterman, et al., 2017) with further characterization of a
subset of them for inhibitor tolerance. This study highlighted the diversity in terms of behaviour,
production and resistance, with for example, *Y. hollandica* and *Y. phangngensis* reached higher cell
biomass and lipid titer compared to *Y. lipolytica* W29 in these conditions. In other studies, up to
thirteen of these yeast species from the Yarrowia clade were screened for erythritol, arabitol
mannitol, citric acid, lipids and protein production (Rakicka, et al., 2016, Rakicka-Pustułka, et al.,
2021). In particular, *Y. divulgata* and *Y. oslonensis* were identified as particularly robust producers of
polyol compared to *Y. lipolytica* when grown on glycerol (Rakicka-Pustułka, et al., 2021). Protease
production has been evaluated within the Yarrowia clade has confirmed the potential of certain
Yarrowia species to secrete significantly high amounts of proteases and revealed the influence of
culture media when benchmarking species for a particular trait (Ciurko, et al., 2023).

Therefore, the physiological aspect and diversity of the Yarrowia clade are now well-documented,
highlighting the variation in technological traits among different species, which can outperform *Y.*
lipolytica. Following this, genetic engineering of the other species of the clade starts to be
considered. A first transformation method for the most distant species *C. hispaniensis* allows using
sucrose as a carbon source by expressing the *S. cerevisiae* invertase as a marker and a heterologous
enzyme. *C. hispaniensis* is resistant to antibiotics like gentamycin, nourseothricin, and hygromycin,
which are typically employed as selective markers in yeast genetic engineering. This limits the
development of genetic tools in this species (Morin, et al., 2020). Moreover, this yeast is refractory to
the standard transformation method and requires the use of biolistic approach for transformation
(Morin, et al., 2020). Similarly, *Y. phangngensis* has been transformed for the first time by using
genetic tools and the standard transformation method developed for *Y. lipolytica* with hygromycin
and zeocin as markers for selections (Quarterman, et al., 2018). The recycling of *Y. lipolytica* tools in

this species is shows possibility for more complex genetic engineering. However, it is worth noting

that the hygromycin concentration required for selection in *Y. phangngensis* needs to be 3 to 5 times

higher compared to *Y. lipolytica* transformants selection. It opens the way for reusing the existed

tools or developing the standardized tools in non-conventional and non-type organisms, allowing

rapid screening of numerous potential hosts for diverse applications.

Here we evaluated the functionality of a GoldenGate modular synthetic biology tool developed for *Y.*

lipolytica (Larroude, et al., 2019) in 6 other species from the Yarrowia clade by using two different

selection markers and the expression of a fluorescent protein as a validation system with a fast

transformation method. As a proof of concept, the genes in the carotenoid pathway were expressed

in 5 of the 6 species.

Results

108 **Results**

Evaluation of selection markers

Evaluation of selection markers

The GoldenGate tool kit developed for *Y. lipolytica* contains 2 dominant selection markers,

hygromycin and nourseothricin, for selecting transformants. This toolkit enables the expression of up

to three transcription units in a single transformation (Larroude, et al., 2019). The minimum

concentration of hygromycin and nourseothricin required for growth inhibition was first evaluated

on 8 strains of the Yarrowia clade (*Y. lipolytica*, *Y. alimentaria*, *Y. deformans*, *Y. galli*, *Y. hollandica*, *Y.*

oslonensis, *Y. phangngensis* and *Y. yakushimensis*) identified in Michely et *al.* (Michely, et al., 2013).

All the strains used in this study are listed in Table 1. *C. hispaniensis* was excluded due to its

previously established resistance to the major antibiotics (Morin, et al., 2020) and the antibiotics

available in the *Y. lipolytica* GoldenGate toolkit would not be effective for this species. The growth of

all the strains were inhibited with 200µg/ml hygromycin or 500µg/ml nourseothricin except for *Y.*

phangngensis which required higher concentration already demonstrated (Quarterman, et al., 2018).

In our hands, we had residual growth of *Y. phangngensis* with 500 µg/mL of hygromycin and 1mg/mL

of nourseothricin. Therefore, this specie was not selected thereafter for evaluation of

transformability with the *Y. lipolytica* GoldenGate vectors.

ID	Yeast strain
Y2183	*Yarrowia lipolytica W29*
Y8112	*Yarrowia oslonensis CBS 10146*
Y8113	*Yarrowia deformans CBS 2071*
Y8114	*Yarrowia galli CBS 9722*
Y8115	*Yarrowia phangngensis CBS 10407*
Y8116	*Yarrowia yakushimensis CBS 10253*
Y8117	*Yarrowia alimentaria CBS 10151*
Y8119	*Yarrowia hollandica CBS 4855*

129   *Table1.* List of wild-type yeast strains of the Yarrowia clade used in this study.

Evaluation of transformation capacity

To evaluate the transformation of each strain, we used integrative expression vectors assembled

using the GoldenGate method containing the expression cassette of the Redstar2 fluorescent protein

with either the selective marker hygromycin or nourseothricin, and ZETA sequences that are known

for their random integration in the *Y. lipolytica* genome (see material and methods). All the building

block have been optimized for *Y. lipolytica* and the details are described in Larroude *et al.* (Larroude,

et al., 2019). The two genes present in each construct, the gene coding for the Redstar2 and the gene

coding for antibiotic resistance, are under the *Y. lipolytica* pTEF promoter. Therefore, we first

evaluated the level of homology between the pTEF promoter sequences of the different strains of

the clade. The *Y. lipolytica* pTEF promoter was used to retrieve promoter ortholog from Whole

Genome Sequences of the other species. All sequences retrieved are available in supplementary file

1. Sequence alignment showed a strong homology, particularly in the 220 base pair region proximal

to the start codon, which suggest the potential for cross-species functionality of the promoter. *Y*

alimantaria appears the most distant and is missing a homology region in the distal part of the

promoter (supplementary file 2).

The transformation of seven species including *Y. lipolytica* as a positive control were carried out by

using a simple and rapid method with a yeast transformation kit (see material and methods section),

in line with the objectives of rapid screening of different host without specific transformation setup.

Both plasmids expressing the Redstar2 with hygromycin or nourseothricin were tested. We were able

to obtain from 1 to around 50 clones for all strains in one transformation experiment except for *Y.*

alimentaria for which no clones were obtained even after several attempts and the extended

incubation for more than a week which usually helps potential transformants to recover. The

relatively lower promoter sequence homology between *C. alimentaria* and *Y. lipolytica,* compared to
other species within the clade, might explain the poor expression of selective marker gene resulting
in unsuccessful transformation. Individual clones from all transformants were then evaluated for
their fluorescence in a microtiter plate reader. All the strains constructed are listed in supplementary
Table 1.

Fig.1. Relative fluorescence reported to OD for wild-type strains and each transformant expressing
the Redstar2 protein after 72h into 96-well microplates. Cultures were performed at least in
duplicate. The upper illustration correspond to schematic draw of the vector used for
transformation. Hgh (vector Hygromycin –pTEF-Redstar2); Nat (vector Nourseothricin –pTEF-
Redstar2). lipo (*Y. lipolytica*); oslo (*Y. oslonensis*); defo (*Y. deformans*); galli (*Y. galli*); yaku (*Y.*
yakushimensis); holl (*Y. hollandica*). All the strains are listed in supplementary Table 1.

All the clones tested were expressing the Redstar2 fluorescent protein with some variation in
intensity depending on the species and marker. As the genomic integration takes place randomly, we
cannot formally compare the level of fluorescence as genome integration locus may influence the
level of heterologous expression. *Y. lipolytica* and *Y. galli* are the species expressing the highest level
of fluorescence in average with both markers while *Y. oslonensis* is the being the lowest (Fig. 1).

Those differences are not due to the variation in fitness between species as they present a similar

growth rate (Fig. 2).

Fig. 2. Representative growth curves in 96 well plates for each species. The growth analysis was

performed using a microtiter plate reader with a reading every 30 min for 72 h. LIPO (*Y. lipolytica*);

OSLO (*Y. oslonensis*); DEFO (*Y. deformans*); GALL (*Y. galli*); YAKU (*Y. yakushimensis*); HOLL (*Y.*

hollandica).

Heterologous pathway expression

The evaluation was pushed forward by the attempt to express a heterologous pathway already

established in *Y. lipolytica* allowing the production of carotenoids (Larroude, et al., 2022) with a

plasmid containing three transcription units (see material and methods section). This pathway is

regularly used as a proof of concept as it allows rapid screening of positive clones with the orange

colour. This heterologous pathway-expressing cassette was transformed into seven species,

respectively.. As we have seen from the transformation of Redstar2 gene expression cassette, we

were not able to obtain transformant *for Y. alimentaria*. We obtained the orange transformants for

all the others except for *Y. yakushimensis* (Fig. 3). All the transformants are listed in supplementary

Table 1. Again, we got slight differences in colour intensity with *Y. hollandica* having the lowest

intensity (Fig. 3). The level of carotenoid could either be due to differences in heterologous

expression or metabolic capacity between species for this particular pathway or locus used for the

integration even if multiple clones had the same trends in colour intensity. The fact that we have

different trends between fluorescent protein expression and carotenoid pathway expression for the

same strain is in favour of a metabolic capacity variability.

Fig. 3. A) Schematic draw of the vector GGE115 used for transformation. B) Growth on YNB plates of

strains transformed with the GG115 plasmid for carotenoid pathway expression. LIPO (*Y. lipolytica*);

OSLO (*Y. oslonensis*); DEFO (*Y. deformans*); GALL (*Y. galli*); YAKU (*Y. yakushimensis*); HOLL (*Y.*

hollandica); W29 (*Y. lipolytica* wild-type strain not transformed). All the strains are listed in

supplementary Table 1.

Conclusion

In this study, we demonstrate that the synthetic biology tools developed for the yeast *Y. lipolytica*

can be effectively repurposed for genetic engineering of other members of the clades, in particular

for *Y. oslonensis*, *Y. deformans*, *Y. galli*, *Y. yakushimensis* and *Y. hollandica* for which we were able to

express a multi-gene heterologous pathway. *Y. phangngensis* and *C. hispaniensis* exhibit a distinct GC
content compared to other members of the clades and significant reduction in size for some protein
families (Groenewald, et al., 2014), indicating a greater sequence divergence. Furthermore, both
strains are highly resistant to typical antibiotics, which may suggest lower compatibility, if any, with
Y. lipolytica's dedicated tools. On the other hand, *Y. alimentaria* is much closer *to Y. lipolytica* in term
of GC content, and its protein expression failure could help determine the homology threshold for
compatible expression, especially for promoters, although further investigation is required.

While it is widely recognized that the choice of host is crucial in metabolic engineering applications,
there's still limited understanding of the strain diversity among non-type strains, particularly for non-
conventional microorganisms. Therefore, exploring this aspect of diversity is a critical route to
achieving success in microbial cell factories. By evaluating other relevant traits, we anticipate that
the results presented in this study will facilitate new biotechnological applications involving
previously overlooked and under-characterized strains within the Yarrowia clade.

Material and methods

Strains and media

All the wild-type yeast strains used in this study are listed in Table 1. All constructed strains are listed
in supplementary Table 1.

Chemically competent *Escherichia coli* DH5α cells were used for cloning and plasmid propagation. *E.*
coli cells were grown at 37°C with constant shaking on 5 ml of LB medium with ampicillin (100 µg ml⁻
¹) or kanamycin (50 µg ml⁻¹) for plasmid selection. Yeasts were grown on rich medium YPD (10 g l⁻¹ of
yeast extract, 10 g l⁻¹ of Peptone, 10 g l⁻¹ of glucose) complemented with hygromycin (200 µg ml⁻¹) or
nourseothricin (500 µg ml⁻¹) for the selection of transformants. Solid media were prepared by adding
1.5% agar. For growth on 96 well plates, yeasts were grown on YNB minimal medium containing 10 g
l⁻¹ glucose, 10 g l⁻¹, 1.7 g l⁻¹ yeast nitrogen base without amino acids, 5.0 g l⁻¹ NH_4Cl and 50 mM
phosphate buffer (pH 6.8).

Plasmid construction

Plasmids expressing the Redstar2 fluorescent protein were assembled using GoldenGate method as
described in (Larroude, et al., 2022) using hygromycin or nourseothricin marker, pTEF promoter,

tLip1-3 terminator, ZETA upstream and downstream integration sequences, Redstar2 gene and
pSB1A3 backbone vector. All building parts are described in (Larroude, et al., 2019).

Plasmid GGE115 allowing expression of the carotenoid pathway was described in (Larroude, et al.,
2022). It contains the ZETA sequences for random integration, the hygromycin marker, and the three
genes involved, GGS1 (geranylgeranyl diphosphate synthase) from *Y. lipolytica*, carPR (phytoene
synthase/lycopene cyclase) from *Mucor circinelloides*, and carB (phytoene dehydrogenase) from *M.*
circinelloides (Larroude, et al., 2018).

Promoter sequences alignment
The *Y. lipolytica* pTEF promoter was used to retrieve promoter ortholog from Whole Genome
Sequences of the other species (*Yarrowia_osloensis* ULGU01000000; *Yarrowia_deformans*
ULGY01000000; *Yarrowia_galli* ULGS01000000; *Yarrowia_yakushimensis* ULGW01000000;
Yarrowia_alimentaria ULGN01000000; *Yarrowia_hollandica* ULGV000000). Multiple sequences
alignment was performed using T-Coffee web server (Di Tommaso, et al., 2011)

Transformation
EZ-Yeast transformation kit (MP Biomedicals) was used for all yeast transformations following the
manufacturer's instruction. The kit does not require competent cells and allows using colony growing
on plates followed by a 30 min transformation step. Plasmids were first digested with restriction
enzyme NotI to release the expression cassette and 2 µg of digested DNA was used for
transformation. For the release of the expression cassette of plasmid GGE115 the restriction enzyme
SfiI was used.

Growth curves and fluorescence measurements
Yeasts were precultured for 24 h in YNB medium and diluted to an OD_{600nm} of 0.1 in fresh medium
and 200 µl was transferred into 96-well microplates. The growth analysis was performed using a
microtiter plate reader (Synergy Mx; BioTek). The settings were 28°C and constant shaking with a
reading every 30 min for 72 h for OD_{600nm} as well as red fluorescence (excitation 558 nm /emission
586 nm). Fluorescence was expressed as a ratio of relative fluorescence divided by OD. Cultures were
performed at least in duplicate.

References

Park, Y.-K., and Ledesma-Amaro, R. (2023) What makes Yarrowia lipolytica well suited for industry?, *Trends Biotechnol.* **41**: 242-254.https://doi.org/10.1016/j.tibtech.2022.07.006.

Zhang, G., Wang, H., Zhang, Z., Verstrepen, K.J., Wang, Q., and Dai, Z. (2022) Metabolic engineering of Yarrowia lipolytica for terpenoids production: advances and perspectives, *Crit. Rev. Biotechnol.* **42**: 618-633.10.1080/07388551.2021.1947183.

Larroude, M., Rossignol, T., Nicaud, J.M., and Ledesma-Amaro, R. (2018) Synthetic biology tools for engineering Yarrowia lipolytica, *Biotechnol. Adv.* **36**: 2150-2164.10.1016/j.biotechadv.2018.10.004.

Yi, X., and Alper, H.S. (2022) Considering Strain Variation and Non-Type Strains for Yeast Metabolic Engineering Applications, *Life* **12**: 510.

Saunders, L.P., Bowman, M.J., Mertens, J.A., Da Silva, N.A., and Hector, R.E. (2015) Triacetic acid lactone production in industrial Saccharomyces yeast strains, *J. Ind. Microbiol. Biotechnol.* **42**: 711-721.10.1007/s10295-015-1596-7.

Kurtzman, C.P. (2011) Chapter 82 - Yarrowia van der Walt & von Arx (1980). In: *The Yeasts (Fifth Edition)*. London: Elsevier. 927-929.https://doi.org/10.1016/B978-0-444-52149-1.00082-3.

Groenewald, M., Boekhout, T., Neuvéglise, C., Gaillardin, C., van Dijck, P.W.M., and Wyss, M. (2014) *Yarrowia lipolytica*: Safety assessment of an oleaginous yeast with a great industrial potential, *Crit. Rev. Microbiol.* **40**: 187-206.doi:10.3109/1040841X.2013.770386.

Michely, S., Gaillardin, C., Nicaud, J.-M., and Neuvéglise, C. (2013) Comparative Physiology of Oleaginous Species from the Yarrowia Clade, *PLoS ONE* **8**: e63356.10.1371/journal.pone.0063356.

Quarterman, J., Slininger, P.J., Kurtzman, C.P., Thompson, S.R., and Dien, B.S. (2017) A survey of yeast from the Yarrowia clade for lipid production in dilute acid pretreated lignocellulosic biomass hydrolysate, *Appl. Microbiol. Biotechnol.* **101**: 3319-3334.10.1007/s00253-016-8062-y.

Chang, C.-F., Chen, C.-C., Lee, C.-F., and Liu, S.-M. (2013) Identifying and characterizing Yarrowia keelungensis sp. nov., an oil-degrading yeast isolated from the sea surface microlayer, *Antonie Van Leeuwenhoek* **104**: 1117-1123.10.1007/s10482-013-0033-z.

Péter, G., Nagy, E.S., and Dlauchy, D. (2019) Systematics, Diversity and Ecology of the Genus Yarrowia and the Methanol-Assimilating Yeasts. In: *Non-conventional Yeasts: from Basic Research to Application*. Sibirny, A. (ed). Cham: Springer International Publishing. 297-339.10.1007/978-3-030-21110-3_9.

Rakicka, M., Kieroń, A., Hapeta, P., Neuvéglise, C., and Lazar, Z. (2016) Sweet and sour potential of yeast from the Yarrowia clade, *Biomass Bioenergy* **92**: 48-54.https://doi.org/10.1016/j.biombioe.2016.06.004.

Rakicka-Pustułka, M., Miedzianka, J., Jama, D., Kawalec, S., Liman, K., Janek, T., et al. (2021) High value-added products derived from crude glycerol via microbial fermentation using Yarrowia clade yeast, *Microbial Cell Factories* **20**: 195.10.1186/s12934-021-01686-0.

Ciurko, D., Neuvéglise, C., Szwechłowicz, M., Lazar, Z., and Janek, T. (2023) Comparative Analysis of the Alkaline Proteolytic Enzymes of Yarrowia Clade Species and Their Putative Applications, *Int J Mol Sci* **24**: 6514.

Morin, N., Czerwiec, Q., Nicaud, J.M., Neuveglise, C., and Rossignol, T. (2020) Transforming Candida hispaniensis, a promising oleaginous and flavogenic yeast, *Yeast* **37**: 348-355.10.1002/yea.3466.

Quarterman, J.C., Slininger, P.J., Hector, R.E., and Dien, B.S. (2018) Engineering Candida phangngensis—an oleaginous yeast from the Yarrowia clade—for enhanced detoxification of lignocellulose-derived inhibitors and lipid overproduction, *FEMS Yeast Res.* **18**.10.1093/femsyr/foy102.

Larroude, M., Park, Y.-K., Soudier, P., Kubiak, M., Nicaud, J.-M., and Rossignol, T. (2019) A modular Golden Gate toolkit for Yarrowia lipolytica synthetic biology, *Microb Biotechnol* **12**: 1249-1259.10.1111/1751-7915.13427.

Larroude, M., Nicaud, J.-M., and Rossignol, T. (2022) Golden Gate Multigene Assembly Method for Yarrowia lipolytica. In: *Yeast Metabolic Engineering: Methods and Protocols*. Mapelli, V., and Bettiga, M. (eds). New York, NY: Springer US. 205-220.10.1007/978-1-0716-2399-2_12.

Larroude, M., Celinska, E., Back, A., Thomas, S., Nicaud, J.M., and Ledesma-Amaro, R. (2018) A
synthetic biology approach to transform Yarrowia lipolytica into a competitive biotechnological
producer of beta-carotene, *Biotechnol. Bioeng.* **115**: 464-472.10.1002/bit.26473.
Di Tommaso, P., Moretti, S., Xenarios, I., Orobitg, M., Montanyola, A., Chang, J.-M., et al. (2011) T-
Coffee: a web server for the multiple sequence alignment of protein and RNA sequences using
structural information and homology extension, *Nucleic Acids Res.* **39**: W13-
W17.10.1093/nar/gkr245.

335