Inhibition of the glucocorticoid-activating enzyme 11β-hydroxysteroid dehydrogenase type 1 drives

concurrent 11-oxygenated androgen excess

Lina Schiffer[1,#], Imken Oestlund[2,#], Jacky Snoep[2,3], Lorna C. Gilligan[1], Angela E. Taylor[1], Alexandra J.

Sinclair[1], Rishi Singhal[4], Adrian Freeman[5], Ramzi Ajjan[6,7], Ana Tiganescu[6,7], Wiebke Arlt[1,8,9] and Karl-

Heinz Storbeck[1,2]

[1] Institute of Metabolism and Systems Research, University of Birmingham, Birmingham, UK

[2] Department of Biochemistry, Stellenbosch University, Stellenbosch, South Africa

[3] Molecular Cell Physiology, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

[4] Upper GI Unit and Minimally Invasive Unit, Heartlands Hospital, University Hospitals Birmingham NHS

Foundation Trust, Birmingham, UK

[5]Emerging Innovations Unit, Discovery Sciences, BioPharmaceuticals R&D, AstraZeneca, Cambridge,

UK

[6] Leeds Institute of Cardiovascular and Metabolic Medicine, University of Leeds, Leeds, UK

[7] NIHR Leeds Biomedical Research Center, Leeds Teaching Hospitals, NHS Trust, Leeds, UK

[8] Institute of Clinical Sciences, Faculty of Medicine, Imperial College, London, UK

[9] Medical Research Council London Institute of Medical Sciences, London, UK

[#]Shared first authors.

Correspondence:

Karl-Heinz Storbeck, Department of Biochemistry, Stellenbosch University, Private Bag X1,

Matieland, 7602, South Africa

Email: storbeck@sun.ac.za

Phone: +27 21 808 5862

Abstract

Aldo-keto reductase 1C3 (AKR1C3) is a key enzyme in the activation of both classic and 11-oxygenated androgens. In adipose tissue, AKR1C3 is co-expressed with 11β-hydroxysteroid dehydrogenase type 1 (HSD11B1), which catalyses the local activation of glucocorticoids but also the inactivation of 11-oxygenated androgens, and thus has the potential to counteract AKR1C3. Using a combination of *in vitro* assays and *in silico* modelling we show that HSD11B1 attenuates the biosynthesis of the potent 11-oxygenated androgen, 11-ketotestosterone, by AKR1C3. Employing *ex vivo* incubations of human female adipose tissue samples we show that inhibition of HSD11B1 results in the increased peripheral biosynthesis of 11-ketotestosterone. Moreover, circulating 11KT increased 2-3 fold in individuals with type 2 diabetes after receiving the selective oral HSD11B1 inhibitor AZD4017 for 35 days, thus confirming that HSD11B1 inhibition results in systemic increases in 11KT concentrations. Our findings show that HSD11B1 protects against excess 11KT production by adipose tissue, a finding of particular significance when considering the evidence for adverse metabolic effects of androgens in women. Therefore, when targeting glucocorticoid activation by HSD11B1 inhibitor treatment in women, the consequently increased generation of 11-ketotestosterone may offset beneficial effects of decreased glucocorticoid activation.

Keywords: Aldo-keto reductase 1C3 (AKR1C3); 11-ketotestosterone; intracrinology; adipose tissue; metabolic syndrome

 Graphical Abstract

Introduction

The 11-oxygenated androgens are a group of adrenal-derived and peripherally activated androgens that make a significant contribution to the circulating androgen pool, particularly in women (>>>). The most potent 11-oxygenated androgen in circulation is 11-ketotestosterone (11KT), which has been shown to bind to and activate the androgen receptor with an affinity and potency similar to testosterone (5). Moreover, several studies have also shown that 11KT circulates at concentrations equal to or higher than those of testosterone in women (3, 6, 7). Unlike the classic androgens, 11-oxygenated androgens do not decline with age, with the result that 11KT is the most abundant active androgen in the circulation of postmenopausal women (3, 6). While the role of 11-oxygenated androgens in healthy individuals has yet to be defined, these androgens have been shown to play a role in disease states such as polycystic ovary syndrome (PCOS), congenital adrenal hyperplasia (CAH), Cushing's syndrome and castration resistant prostate cancer (CRPC) (8–12).

The biosynthesis of 11KT begins in the adrenal, where cytochrome P450 11B1 (CYP11B1) catalyses the 11β-hydroxylation of locally produced androstenedione to yield 11β-hydroxyandrostenedione (11OHA4). 11OHA4 is released into circulation, and serves as the primary precursor for the biosynthesis of 11KT by peripheral tissues (**Figure 1**). Specifically, 11OHA4 is converted to 11-ketoandrostenedione (11KA4) by mineralocorticoid target tissues expressing 11β-hydroxysteroid dehydrogenase type 2 (HSD11B2), such as the kidney (1>). 11KA4 then subsequently serves as the substrate for the key androgen activating enzyme, aldo-keto reductase 1C3 (AKR1C3), which is abundantly expressed in adipose tissue (14–17). Notably, AKR1C3 catalyses the conversion of 11KA4 to 11KT significantly more efficiently than that of androstenedione to testosterone (16, 17). The biosynthesis of 11KT is, however, more complex than that of testosterone as the presence of the 11-keto group makes both 11KA4 and 11KT substrates for 11β-hydroxysteroid dehydrogenase type 1 (HSD11B1)—an enzyme best known for amplification of glucocorticoid signalling, which it achieves by converting the inactive cortisone to active cortisol in key metabolic tissues including the liver, adipose tissue and skeletal muscle (11, 18–20). Given that AKR1C3 and HSD11B1 are co-expressed in adipose tissue, we propose that HSD11B1 regulates the amount of 11KT produced in two ways. First, by

competing with AKR1C3 for 11KA4, thereby limiting the amount of 11KA4 converted to 11KT by

AKR1C3. Secondly, by converting some of the 11KT formed to the far less androgenic 11β-

hydroxytestosterone (11OHT) (11) (**Figure 1**). Indeed, while investigating the metabolism of 11KA4 in

differentiated Simpson-Golabi-Behmel syndrome preadipocyte (SGBS) cells Paulukinas et. al. noted

that HSD11B1 may be protective of androgen excess (17).

This proposed mechanism is important when considering that HSD11B1 has long been a therapeutic

target for prevention of tissue-specific excess glucocorticoid exposure and treatment of related

adverse metabolic effects (19, 21–24). Promising preclinical studies in rodents, which crucially do not

produce 11-oxygenated androgens (25–30), were unable to accurately model competing HSD11B1

and AKR1C3 effects in humans. Our proposed hypothesis could provide an explanation for the more

limited effect sizes observed in clinical trials with HSD11B1 inhibitors (31, 32, 41, 42, 33–40), as

increased 11KT production resulting from HSD11B1 inhibition could counteract beneficial effects of

reduced glucocorticoid signalling, by increasing androgen signalling. This is particularly significant in

women where elevated androgens are linked to insulin resistance, type 2 diabetes, hypertension,

cardiovascular disease and metabolic dysfunction associated steatotic liver disease (MASLD), in

particular in the context of polycystic ovary syndrome, for which androgen excess is a defining feature

(43–52).

Here, we characterised the interplay between HSD11B1 and AKR1C3 as co-regulators of 11KT

biosynthesis using a combination of *in vitro*, *in silico*, and human *ex vivo* and *in vivo* approaches

including selective HSD11B1 inhibitor treatment in individuals with type 2 diabetes.

Materials and Methods

Steroids and inhibitors

Androstenedione (4-androstene-3,17-dione), carbenoxolone ((3β,20β)-3-(3-carboxy-1-oxopropoxy)-

11-oxo-olean-12-en-29-oic acid, disodium salt; CBX), cortisol (11β,17α,21-trihydroxypregn-4-ene-3,20-

dione), cortisone (17α,21-dihydroxy-4-pregnene-3,11,20-trione), DHEA (5-androsten-3β-ol-17-one),

DHEA sulfate (5-androsten-3β-ol-17-one-3-sulfate) and testosterone (4-androsten-17β-ol-3-one) were

purchased from Sigma Aldrich. 11β-hydroxyandrostenedione (11β-hydroxy-4-androstene-3,17-dione),

11β-hydroxytestosterone (11β,17β-dihydroxyandrost-4-en-3-one), 11-ketoandrostenedione (4-

androsten-3,11,17-trione) and 11-ketotestosterone (4-androsten-17β-ol-3,11-dione) were from

Steraloids. The deuterated internal standards 11β-hydroxyandrostenedione-2,2,4,6,6,16,16-D7 (D7-

11OHA4), 11-ketotestosterone-16,16,17-D3 (D3-11KT), androstenedione-2,2,4,6,6,16,16-D7 (D7-A4),

cortisol-9,11,12,12-D4 (D4-F), DHEAS-16,16-D2 (D2-DHEAS) and testosterone-16,16,17-D3 (D3-T) were

all purchased from Cambridge Isotope Laboratories. Dehydroepiandrosterone-2,2,3,4,4,6-D6 (D6-

DHEA) and cortisone-2,2,4,6,6,9,12,12-D8 (D8-E) were purchased from Sigma Aldrich, while 11-

ketoandostenedione-2,2,4,6,6,9,12,12,16,16-D10 (D10-11KA4) was custom synthesized by Toronto

Research Chemicals. The selective HSD11B1 inhibitor, 2-[(3S)-1-[5-(cyclohexylcarbamoyl)-6-

propylsulfanylpyridin-2-yl]-3-piperidyl] acetic acid (AZD-4017) was provided by AstraZeneca.

Plasmids

The pcDNA3 vector containing AKR1C3 was provided by Prof. J. Adamski (Helmholtz Zentrum

München, Germany). The pcDNA3 vectors containing HSD11B1 and hexose 6-phosphate

dehydrogenase (H6PDH) were from Prof. K. Chapman (University of Edinburgh, UK). The pTAT-GRE-

Elb-luc luciferase promoter reporter construct, containing two copies of the GRE was from Prof. G.

Jenster (Erasmus University of Rotterdam, Netherlands), while the GR expressing plasmid (pRShGRα)

was from Prof. R. Evans (Howard Hughes Medical Institure, La Jolla, USA). The pCIneo plasmid

expressing the wild-type human AR was from Prof. N. Sharifi (Cleveland Clinic, Cleveland, USA). A

pCIneo plasmid, containing no cDNA insert, was purchased from Promega (Wisconsin, USA).

Cell lines

HEK293 cells were purchased from the American Type Culture Collections (ATCC) and cultured in high

glucose Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% FBS, 1.5 g/L NaHCO$_3$ and

1% penicillin-streptomycin and maintained at 37 °C in 90% humidity and 5% CO^2. The cells were

authenticated by short-tandem repeat profiling (NorthGene) and were regularly tested for

mycoplasma contamination.

Determining kinetic parameters for HSD11B1 and computational model construction

The kinetic parameters ($K_{m,\,app}$ and $V_{max,\,app}$) for HSD11B1 were determined using the methodology

previously reported for AKR1C3 (1>). Briefly, HEK293 cells were seeded into 10 cm^2 tissue culture

dishes at a density of 2x10^5 cells per mL (10 mL per dish). After a 24 hour incubation period, cells were

co-transfected with 0.5 μg HSD11B1 and 0.5 μg H6PDH DNA using X-tremeGENE™ HP DNA transfection

reagent (Roche) according to manufacturer's protocol. Following a 24 hour incubation the transfected

cells were re-plated into 48-well Corning® CELLBIND® plates at a density of 2x10^5 cells per mL (300 μL

per well). Cells were incubated for an additional 24 hours to allow for attachment before being treated

with serum free media, containing appropriate steroid substrate (cortisone, 11KA4, or 11KT) at

concentrations ranging from 0.1 µM to 10 µM. Aliquots (250 µL) for the quantification of steroid

metabolism by ultra-high performance liquid chromatography tandem mass spectrometry (UHPLC-

MS/MS) were collected at specific time intervals. Kinetic parameters were determined in Wolfram

Mathematica (Version 13) by assuming Michaelis Menten kinetics and fitting the $K_{m, app}$ and $V_{max, app}$

values by minimizing the sum of the squared differences between the experimentally determined

progress curves and model, weighted by the variance using the NMinimize function. The model was

fitted to triplicate data from three independent experiments. The transfection efficiency of each

individual experiment was taken into account by determining the initial rate of the conversion of 100

nM 11KA4 for each independent experiment. A computational model containing the determined

kinetic for HSD11B1 and those previously determined for AKR1C3 was subsequently constructed. This

model is comprised of ordinary differential equations (ODEs) for the metabolites, and numerical

integration by making use of the NDSolve routine in Mathematica allowed for experimentally

comparable metabolite profiles to be obtained. Additionally, the model was used to predict the

conversion of cortisone or 11KA4 for varying HSD11B1:AKR1C3 ratios, see Supplementary material for

a complete model description and simulation details.

Determining the effect of different HSD11B1:AKR1C3 ratios on cortisone and 11KA4 metabolism

HEK293 cells were seeded into 10 cm^2 tissue culture dishes at a density of 2x10^5 cells per mL (10 mL

per dish). After a 24 hour incubation period, dishes were co-transfected with either 0.5 µg HSD11B1

and 0.5 µg H6PDH or 0.5 µg AKR1C3 and 0.5 µg pCINeo using X-tremeGENE™ HP DNA transfection

reagent (Roche). Additional dishes were transfected with 1 µg of the pCINeo (no insert) as to allow for

the subsequent plating of an equal number of transfected cells at all HSD11B1:AKR1C3 ratios. After 24

hours cells were counted and combined in different ratios of HSD11B1:AKR1C3, before being re-plated

into 12-well Corning® CELLBIND® surface plates at a density of 4x10^5 cells per mL (1 mL per well). After

an additional incubation period of 24 hours, the media was replaced with serum free DMEM containing

100 nM substrate (11KA4 or cortisone). Aliquots (500 µL) were collected 24 hours after the addition of

substrate for analysis by UHPLC-MS/MS, as described below.

Determining the effect of CBX on HSD11B1 and AKR1C3 co-expression

163 HSD11B1 and AKR1C3 transfections were performed as described above. Transfected cells were

164 re-plated at a 1:1 ratio of HSD11B1:AKR1C3 (2x10^5 cells/mL, 300 µL) in 48-well Corning® CELLBIND®

165 surface plates. After 24 hours, media was replaced with serum free media containing 100 nM cortisone

166 in combination with either 10 nM or 100 nM 11KA4 in the absence and presence of 10 µM CBX.

167 Aliquots (200 µL) were collected over a 24 hour period for analysis by UHPLC-MS/MS.

Promoter reporter assays

To determine the effect of the HSD11B1 inhibitor CBX on AR and GR transactivation, HSD11B1 and AKR1C3 transfections were performed 24 hours before co-transfections of HEK293 cells with either a AR expression plasmid (0.09 µg) and the pTAT-GRE-Elb-luc luciferase promoter reporter construct (0.9 µg) or a GR expression plasmid (0.09 µg) and the pTAT-GRE-Elb-luc luciferase promoter reporter construct (0.9 µg). HSD11B1 and AKR1C3 transfected cells were re-plated at a 1:1 ratio ($4x10^5$ cells/mL, 1 mL) in 12-well Corning® CELLBIND® surface plates. Following 24 hours, media was replaced with serum free media containing 100 nM cortisone and 10 nM 11KA4 in the absence and presence of 10 µM CBX. At the same time the AR-GRE transfected cells and GR-GRE transfected cells were re-plated into 48-well Corning® CELLBIND® plates ($3x10^5$ cells/mL, 300 µL). Following a 24 hour incubation with steroid, aliquots (200 µL) from the HSD11B1:AKR1C3 expressing cells were collected for analysis of steroid metabolism by UHPLC-MS/MS. A further 300 µL conditioned media was transferred to each of the AR-GRE and GR-GRE transfected cells. These cells were incubated for a further 24 hours after which cell lysates were harvested in passive lysis buffer. Reporter activity was then measured by making use of the Luciferase Assay System (Promega). The luciferase values were normalised to the protein concentration of each individual sample determined using a Thermo Scientific™ Pierce™ BCA Protein Assay kit.

Determining the effect of ADZ4017 on 11KA4 conversion in human adipose tissue

Paired subcutaneous and omental adipose tissue was collected from eight female participants (age range 32-59 years, BMI range 42-57 kg/m^2) undergoing elective abdominal non-cancer surgery (Black Country Research Ethics Committee references 14/WM/0011 and 19/WM/0183). Written, informed consent was obtained prior to surgery and sample collection. Adipose tissue samples were transported to the laboratory at room temperature in DMEM/F12 supplemented with 33 µM biotin, 17 µM panthotenic acid and 1% penicillin-streptomycin. Connective tissue and vessels were removed and the adipose tissue washed in PBS. Tissue samples of 100-300 mg were cut and weighed. Each sample was cut into four smaller pieces for incubations with 100 nM cortisone or 100 nM 11KA4 in the presence and absence of 100 nM HSD11B1 inhibitor AZD4017 in DMEM/F12-Ham (supplemented as described above). Final solvent concentrations were adjusted to 0.00302% MeOH and 0.1% DMSO in all incubations. Tissue incubations were constantly rotated in a hybrid oven at 37 °C for 72 hours. Medium samples were centrifuged for 10 minutes at 16,000g and 4 °C and the supernatant was used for analysis of steroid metabolism by UHPLC-MS/MS. The tissue was washed in PBS, snap frozen in liquid nitrogen and stored at -80 °C for RNA extraction.

RNA isolation and gene expression analysis from HEK293 cells

Total RNA was isolated from cells using Tri-Reagent® (Sigma) and cDNA synthesis carried out using the GoScript™ Reverse Transcription System kit (Promega). Quantitative PCR was performed using a Roche LightCycler® 96 rapid thermal cycler instrument and the KAPA SYBR® FAST qPCR Master Mix for LightCycler®. Primer sequences as follows, AKR1C3 (forward) 5'- AGCCAGGTGAGGAACTTTC-3' (53), AKR1C3 (reverse) 5'- ATCACTGTTAAAATAGTGGAG-3' (53), HSD11B1 (forward) 5'- CTGCCTGCTTAGGAGGTTGT-3', HSD11B1 (reverse) 5'- CCTTGGAGCATCTCTGGTCTG-3', the reference gene ubiquitin conjugating enzyme (UBC) (forward) 5'- CCGGGATTTGGGTCGCAG-3', UBC (reverse) 5'- TCACGAAGATCTGCATTGTCAAG-3'. Primer efficiencies were between 90% and 110% for all primer sets. Relative gene expression was calculated using a modified form of the model described by Pfaffl (54). Inter-experimental variations were adjusted using a mean-centering (55).

RNA isolation and gene expression analysis from adipose tissue

Adipose tissue was homogenised in Tri-Reagent® and total RNA was isolated after phenol-chloroform extraction using the Qiagen RNeasy Mini Kit using an established protocol (56). Reverse transcription was performed using Applied Biosystems™ TaqMan™ Reverse Transcription Reagents according to the manufacturer's protocol. qPCR was performed on an ABI 7900HT sequence detection system (Perkin Elmer, Applied Biosystems) using TaqMan™ Gene Expression Assays (FAM-labelled) and the SensiFAST™ Probe Hi-ROX kit (Bioline). The following TaqMan™ Gene Expression Assays were used: 18S Hs99999901_s1, AKR1C3 Hs00366267_m1, HSD11B1 Hs01547870_m1.

Determining the effect of ADZ4017 glucocorticoid and androgen concentrations *in vivo*

A total of 28 participants (22 male and 6 female), all with type 2 diabetes, were recruited for a randomized, double-blind, parallel-group, placebo-controlled trial. Full ethical approval was acquired from North West Greater Manchester Central Research Ethics Committee 17/NW/0283 prior to the initiation of recruitment for the study. Informed consent was obtained from all participants after the nature and possible consequences of the study had been explained. Participants received either oral AZD4017 (400 mg) or a matched oral placebo twice daily for 35 days. Blood samples and urine were collected from individuals before first treatment and after completion of the trial and were analysed by UHPLC-MS/MS and GC-MS, respectively. Refer to Ajjan *et al.* (40) for further details on study design and participants.

Steroid extraction

Steroids were extracted from HEK293 tissue culture media using Tert-butyl methyl ether (MTBE) (57). In brief, following the addition of an internal standard mix (100 µL containing 2 ng D3-T and 5 ng D7-

A4, D7-11OHA4, D3-11KT and D4-F in acetonitrile) and MTBE (3:1 ratio of MTBE:media), samples were

vortexed at 1 500 rpm for 10 minutes, after which they were incubated at -80 °C for 1 hour, allowing

for the aqueous layer to freeze. The resulting organic phase was subsequently transferred to a clean

test tube and evaporated under a stream of nitrogen gas at 45 °C. Extracted steroids were then

reconstituted by the addition of 50% (v/v) methanol and stored at -20 °C until analysis by UHPLC-

MS/MS. Serum steroids and steroids in medium adipose tissue incubations were extracted using a

similar method. Samples (200 μL for serum, 500 μL for adipose tissue conditioned medium) were mixed

with 10 μL internal standards in 50% (v/v) methanol in water (D4-F, D8-E, D6-DHEA, D7-A4, D3-T, D7-

11OHA4, D10-11KA4, D3-11KT; 5 ng each for serum, 20 ng each for adipose tissue). For serum samples

50 μL acetonitrile were added for protein precipitation prior to extraction and the samples vortexed.

MTBE was added to each sample (1 mL for serum; 3 mL for medium from adipose tissue incubations)

and the samples were vortexed at 1000 rpm for 10 min. Following incubation at room temperature for

30 minutes the organic phase was transferred, dried under a stream of nitrogen gas at 45 °C and

reconstituted with 50% (v/v) methanol in water prior to analysis by UHPLC-MS/MS. For the

measurement of serum DHEAS, 200 ng D2-DHEAS in 20 uL of 50% methanol was added to 20 μL of

serum followed by 20 μL of 1 mM $ZnSO_4$ and 100 μL acetonitrile. The samples were centrifuged and

100 μL of the supernatant was dried and reconstituted in 50% (v/v) methanol in water prior to UHPLC-

MS/MS analysis.

Quantification of steroids by ultra-high performance liquid chromatography tandem mass
spectrometry (UHPLC-MS/MS)

Steroids from assays performed in HEK293 cells were separated and quantified using a UPLC high

strength silica (HSS) T3 column (2.1 mm X 50 mm, 1.8 μm particle size) coupled to an ACQUITY UPLC

and Xevo TQ-S triple quadrupole mass spectrometer (Waters Corporation, Milford, USA) as previously

reported (57). Steroids extracted from serum or adipose tissue incubations were separated and

quantified using Phenomenex Luna Omega polar C18 column (2.1 X 50 mm, 1.6 μm particle size;

Phenomenex, Macclesfield, UK) coupled to an ACQUITY UPLC and a Waters Xevo TQ-XS triple

quadrupole mass spectrometer as previously described (58). Serum DHEAS was quantified using a HSS

T3 column (2.1 X 50 mm, 1.8 μm particle size; Waters) coupled to an ACQUITY UPLC and a Waters Xevo

TQ-XS triple quadrupole mass spectrometer as previously described (59, 60).

Quantification of urinary steroids by GC-MS

263   Measurement of urinary steroid metabolite excretion was carried out by a long-established gas

264   chromatography-mass spectrometry (GC-MS) method (61). In brief, free and conjugated steroids were

265   extracted from 1 mL urine by solid-phase extraction. Steroid conjugates were enzymatically

hydrolyzed, reextracted, and chemically derivatized to form methyloxime trimethyl silyl ethers. An Agilent 5975 instrument operating in selected-ion-monitoring (SIM) mode achieved sensitive and specific detection and quantification of selected steroid metabolites.

Statistical analysis

We performed unpaired t-tests to compare the effect of HSD11B1 inhibition on 11-oxygenated androgen production in HEK293 cells expressing AKR1C3 and HSD11B1 and on the transactivation of the androgen receptor. Paired t-tests were performed to assess the effect of HSD11B1 inhibition in adipose tissue, while Wilcoxon signed-rank tests were used to assess the effect in serum and urine. Correlations between serum steroid concentrations and clinical outcomes were investigated using a Spearman's Rho test.

Results

HSD11B1 efficiently catalyses the 11β-reduction of 11-oxygenated androgens.

Non-steroidogenic HEK293 cells were transiently transfected to express HSD11B1 and activity assayed. Rate equations were fitted to the progress curves for three independent experiments as shown in **Supplementary Figure 1**. A low but detectable reverse reaction rate was observed for the cortisone to cortisol and the 11KA4 to 11OHA4 reactions. Reversible kinetics were therefore used for these reactions. Our results show that the apparent K_m ($K_{m, app}$) of HSD11B1 for cortisone and 11KA4 are similar (0.9 μM and 1 μM), while the $K_{m, app}$ for 11KT (0.21 μM) is >4-fold lower, indicating a higher affinity towards 11KT (**Table 1**). However, the apparent V_{max} ($V_{max, app}$) for 11KT (0.05 μM/h) was >2-fold lower than that for cortisone (0.12 μM/h) and 7-fold lower than that for 11KA4 (0.35 μM/h). Considering the V_{max}/K_m values, an estimate of enzyme efficiency, both 11KA4 and 11KT are more efficiently converted to their respective products than the traditionally recognised HSD11B1 substrate cortisone.

Co-expression of HSD11B1 with AKR1C3 modulates the activation of 11-oxygenated androgens.

AKR1C3 is the key enzyme catalysing the peripheral conversion of androgen precursors into active androgens (1>). We have previously shown that AKR1C3 catalyses the conversion of 11KA4 to the

potent androgen 11KT 8-fold more efficiently than the conversion of androstenedione to testosterone

(16). To determine how the co-expression of HSD11B1 with AKR1C3 would affect 11KT production, we

mixed transfected HEK293 cells expressing either AKR1C3 or HSD11B1 to obtain increasing ratios of

HSD11B1:AKR1C3, while keeping the number of cells expressing AKR1C3 constant. The total cell count

was also kept constant by including cells transfected with an empty plasmid as required. The relative

expression of HSD11B1:AKR1C3 was confirmed by qPCR (**Figure 2A**). Increased ratios of

HSD11B1:AKR1C3 resulted in increased concentrations of cortisol being produced upon addition of

100 nM cortisone (**Figure 2B**), thereby confirming the increased HSD11B1 expression. Similarly,

increasing the ratio of HSD11B1:AKR1C3 resulted in the concomitantly increased production of

11OHA4 from 11KA4, with 11OHA4 becoming the most abundant product when the HSD11B1:AKR1C3

ratio was increased beyond 0.1:1 (**Figure 2C**). Biosynthesis of 11KT was highest when AKR1C3 was

expressed in the absence of HSD11B1, with increasing HSD11B1:AKR1C3 ratios resulting in a rapid

decrease in the concentrations of 11KT. 11OHT initially increased as the ratio of HSD11B1:AKR1C3

increased, peaking at a ratio of 0.25:1. Further increases in the HSD11B1:AKR1C3 ratio resulted in

decreased concentrations of 11OHT being produced.

Using the parameterised rate equations for HSD11B1 and those previously determined by us for

AKR1C3 (1>), a computational model consisting of a set of ordinary differential equations was

constructed to simulate steroid metabolism in cells with both enzymes expressed:

$$\frac{d\,\mathrm{ka4}(t)}{d\,\mathrm{t}} = -v_{\mathrm{HSD\_ka4}} - v_{\mathrm{AKR\_ka4}}; \quad \frac{d\,\mathrm{oha4}(t)}{d\,\mathrm{t}} = v_{\mathrm{HSD\_ka4}}; \quad \frac{d\,\mathrm{kt}(t)}{d\,\mathrm{t}} = v_{\mathrm{AKR\_ka4}} - v_{\mathrm{HSD\_kt}}; \quad \frac{d\,\mathrm{oht}(t)}{d\,\mathrm{t}} = v_{\mathrm{HSD\_kt}}$$

The rate equations for HSD11B1 and AKR1C3 are given in the Supplementary material. The model was

able to predict the experimentally determined effect of increasing HSD11B1:AKR1C3 ratios on the

metabolism of cortisone and 11KA4, respectively (**Figure 2**). Note that the kinetic parameters in the

mathematical model were not fitted to the experimental data for the different HSD11B1:AKR1C3

ratios, but were determined in the progress curves for the individual incubations (**Supplementary**

Figure 1). Taken together these data clearly show that HSD11B1 modulates the biosynthesis of 11KT

by AKR1C3, and that the model can predict the effect of varying HSD11B1:AKR1C3 ratios.

321 *In vitro* **inhibition of HSD11B1 results in the accumulation of 11-ketotestosterone.**

322 Next, we set out to determine the effect of HSD11B1 inhibition on 11KT biosynthesis. We mixed

323 HEK293 cells transiently transfected with either AKR1C3 or HSD11B1 at a ratio of 1:1 and

324 simultaneously treated the cells with cortisone and 11KA4 in the absence and presence of the HSD11B1

325 inhibitor, carbenoxolone (CBX). We used either equimolar concentrations of cortisone and 11KA4 (100

326 nM each) or 10-fold excess of cortisone (100 nM) over 11KA4 (10 nM) to more closely mimic circulating

327 concentrations (>). Our results show that CBX reduced the production of cortisol from cortisone from

328 56.5 nM to 3.2 nM after 24 hours, demonstrating effective HSD11B1 inhibition (**Figure 3A & B**). In the

329 absence of CBX the majority of 11KA4 (69%) was converted to 11OHA4 by HSD11B1, with only 3.8 nM

330 of 11KT produced from 100 nM 11KA4 by AKR1C3 (**Figure 3C**). However, inhibition of HSD11B1 resulted

331 in a 7-fold increase in 11KT production after 24 hours (**Figure 3D**). Similarly, in cells treated with 100

332 nM cortisone and 10 nM 11KA4, inhibition of HSD11B1 by CBX resulted in a 12-fold increase in 11KT

333 production (0.15 nM without CBX to 1.8 nM with CBX) (**Figure 3E & F**). 11KT was the by far most

334 abundant 11-oxygenated androgen following inhibition of HSD11B1. Inhibition of both the conversions

335 from 11KA4 to 11OHA4 and 11KT to 11OHT, respectively, allowed AKR1C3 to catalyse the conversion

336 of 11KA4 to 11KT without the need to compete for substrate and prevented the inactivation of 11KT

337 to the less potent 11OHT. The computational model was able to independently predict the outcome

338 of HSD11B1 inhibition under the experimental conditions (**Figure 3**).

339

340 *In silico* **analysis of the combined effect of HSD11B1:AKR1C3 co-expression and HSD11B1 inhibition**

341 **on 11KT biosynthesis.**

342 After having shown that the computational model was able to individually predict both the effect of

343 increasing HSD11B1:AKR1C3 ratios (**Figure 2**) and the effect of HSD11B1 inhibition (**Figure 3**), we used

344 the model to simultaneously predict the effect of differing HSD11B1 and AKR1C3 expression and

345 differing degrees of HSD11B1 inhibition on the conversion of 10 nM 11KA4 to 11OHA4, 11KT and

346 11OHT after 24 hours (**Figure 4**). These results confirmed that HSD11B1 strongly modulates the

347 biosynthesis of 11KT by competing for the precursor, 11KA4, and by converting the 11KT that is formed

to the less potent androgen 11OHT. The inhibition of HSD11B1 reduced these effects under all HSD11B1:AKR1C3 ratios tested leading to increased production of 11KT. Notably, lower ratios of HSD11B1:AKR1C3 were more sensitive to HSD11B1 inhibition, more quickly leading to increased 11KT concentrations as the level of inhibition was increased. Higher absolute concentrations of 11KT were also produced by lower ratios of HSD11B1:AKR1C3 (AKR1C3 was kept constant) when HSD11B1 was inhibited up to the maximum simulated I/Ki value of ten. Near complete inhibition of HSD11B1 (I/Ki = 100), however, negated the effect of the HSD11B1:AKR1C3 ratio with 11KT production dependent only on AKR1C3 (**Supplementary Figure 2**). The effect of HSD11B1 inhibition is therefore dependent on the degree of inhibition, the ratio of HSD11B1:AKR1C3 and the absolute expression level of HSD11B1 and AKR1C3.

To investigate the potential biological impact of HSD11B1 inhibition, we combined HEK293 cells expressing AKR1C3 or HSD11B1 at a 1:1 ratio and incubated the cells with 100 nM cortisone and 10 nM 11KA4 in the absence and presence of CBX for 24 hours. The conditioned media were then transferred to HEK293 cells transiently transfected with either a plasmid expressing the glucocorticoid receptor (GR) or the androgen receptor (AR), and a luciferase reporter construct. The GR- and AR-mediated induction of luciferase was subsequently measured after an additional 24 hour incubation. As expected, HSD11B1 mediated the conversion of cortisone to cortisol, with the resulting cortisol increasing the transactivation of the GR (94-fold induction). Inhibition of HSD11B1 with CBX reduced cortisol production 32-fold (**Figure 5A**) and resulted in a 34-fold decrease in GR transactivation (**Figure 5B**). Conversely, inhibition of HSD11B1 resulted in a 7-fold increase in transactivation of the AR due to significantly increased 11KT biosynthesis (**Figure 5C & D**). Basal levels of AR transactivation in the absence of CBX and 11KT were due to the presence of low concentrations of 11OHT (0.81 nM; **Supplementary Figure 3**), which is a partial agonist of the AR (11, 62). It should be noted that we observed that CBX weakly antagonises the AR (**Supplementary Figure 4**), hence, the induction of the AR was likely underestimated. Nonetheless, taken together, these data demonstrate that while inhibition of HSD11B1 results in the desired decrease in GR transactivation, this is accompanied by a shift towards AR transactivation.

Selective HSD11B1 inhibition increases 11KT production in female human adipose tissue *ex vivo*.

We next set out to determine the effect of HSD11B1 inhibition on 11KT biosynthesis in paired subcutaneous and omental adipose tissue samples from women undergoing elective bariatric surgery. For these experiments we employed the selective HSD11B1 inhibitor AZD4017 (63). The conversion of cortisone to cortisol was reduced to 3% of the uninhibited conversion rate in subcutaneous adipose tissue treated with AZD4017, thereby confirming effective HSD11B1 inhibition (**Figure 6A**). Following incubation of subcutaneous adipose tissue with 11KA4 (100 nM), the most abundant product formed in the absence of AZD4017 was 11OHT (60.1% of total products formed), followed by 11OHA4 (23.4%) and 11KT (16.5%) (**Figure 6F**). The biosynthesis of both 11KT and 11OHT is dependent on AKR1C3 activity and the observed sum of 11KT and 11OHT (76.6% of products formed) suggested substantial AKR1C3 expression. This was confirmed by qPCR analysis, which showed higher AKR1C3 expression relative to that of HSD11B1 (**Figure 6C**). However, HSD11B1 converted 78.5% of the 11KT formed to the less active 11OHT. Inhibition of HSD11B1 by AZD4017, resulted in a 5-fold increase in 11KT, with 11KT becoming the most abundant product (85.6%) (**Figure 6G**). This was accompanied by significant decreases in the production of both 11OHA4 and 11OHT (21-fold and 4.5-fold, respectively).

Marginally lower conversion of cortisone to cortisol was observed in omental adipose tissue as compared to subcutaneous tissue, with AZD4017 again effectively inhibiting HSD11B1 activity (**Figure 6B**). Treatment with AZD4017 again resulted in a significant decrease in 11OHA4 and 11OHT concentrations, with an accompanying 4-fold increase in 11KT production (**Figure 6H & I**).

The sum of 11KT and 11OHT (66.0%) was lower in the omental tissue than in subcutaneous tissue (76.6%) illustrating lower AKR1C3 expression as confirmed by qPCR (**Figure 6C & D**). The ratio of HSD11B1 over AKR1C3 expression was lower in subcutaneous than omental adipose tissue (**Figure 6E**), favouring androgen activation in subcutaneous tissue vs glucocorticoid activation in omental tissue.

The metabolism of 11KA4 and the effect of AZD4017 inhibition in human adipose tissue was simulated with the mathematical model that was constructed using conversion data obtained from transfected HEK293 cells. The expression levels of the HSD11B1 and AKR1C3 in the adipose tissue relative to that

in transfected HEK293 cells therefore needed to be estimated. This was achieved by fitting the expression levels of HSD11B1 to the cortisone to cortisol conversion data (**Figure 6A & B**) and then using an AKR1C3 activity based on the AKR1C3/HSD11B1 expression ratio as determined by qPCR (**Figure 6C & D**). This allowed the model to predict the conversion of 11KA4 to the respective products over time. These simulation revealed the initial accumulation of 11KT, prior to the time dependent conversion of this product to 11OHT by HSD11B1 (**Figure 6F &H**). Inhibition of HSD11B1 eliminated the latter conversion yielding only 11KT and illustrated that the amount of 11KT formed over time was dependent on the absolute expression levels of AKR1C3.

Taken together, these data indicate that inhibition of HSD11B1 in adipose tissue leads to a shift in the metabolism of 11-oxygenated androgens resulting in increased 11KT biosynthesis.

***In vivo* treatment with a selective HSD11B1 inhibitor increases circulating 11KT concentrations.**

After having shown that inhibition of HSD11B1 in adipose tissue shifts the metabolism of 11-oxygenated androgens towards 11KT biosynthesis, we next set out to determine if inhibition of HSD11B1 would alter circulating concentrations of 11KT. We therefore measured a panel of 11-oxygenated androgens as well as cortisol and cortisone in the serum of individuals with type 2 diabetes who had received treatment with the selective HSD11B1 inhibitor AZD4017 during a randomized, double-blind, parallel-group, placebo-controlled phase 2b pilot trial (40). Paired (day 0 and day 35) samples were available for 18 participants (13 men and 5 women; median BMI 32.2 kg/m^2 range 25.0-47.5 kg/m^2; median age 64.5, range 28-84 years), with 11 randomised to HSD11B1 inhibitor treatment (7 men and 4 women) and 7 (6 men and 1 woman) to placebo.

Our measurements confirmed that AZD4017 significantly reduced the serum cortisol/cortisone ratio, indicating effective HSD11B1 inhibition as previously reported for selected urinary glucocorticoid metabolites by LC-MS/MS (40), and further confirmed here by GC-MS (**Supplementary Figure 5**). However, concomitantly, circulating 11KT concentrations significantly increased 2.5-fold (from 1.7 to 4.3 nM) following treatment with 400 mg AZD4017 twice daily for 35 days. This was accompanied by a significant, 2-fold decrease in 11OHT concentrations and a 4-fold decrease in the 11OHT/11KT ratio

(Figure 7). While the circulating concentrations of the 11-oxygenated androgen precursors 11OHA4 and 11KA4 did not change, a small, but significant decrease was also observed for the 11OHA4/11KA4 ratio. These changes were also reflected by a significant increase in the urinary metabolite 11-oxoandrosterone, which is derived from 11KA4 and 11KT (64), and a 6-fold decrease in the 11β-hydroxyandrosterone/11-oxoandrosterone ratio **(Figure 7)**. No changes in the serum concentrations of 11-oxygenated androgens were observed in the placebo-treated group (**Supplementary Figure 6**).

When considering correlations between the changes in circulating steroid concentrations, it is interesting to note that the cortisol/cortisone ratio is highly correlated with the change in both the ratios of 11OHA4/11KA4 (R = 0.893; P < 0.001) and 11OHT/11KT (R = 0.686; P < 0.01), but not with the change in circulating 11KT (R = -0.375; P = 0.126) (**Supplementary Table 1**).

Discussion

The relatively recent identification of 11-oxygenated androgens as important contributors to the human androgen pool has prompted a re-evaluation of existing paradigms. One such paradigm is the idea that the two HSD11B isoforms are predominantly involved in the peripheral metabolism of glucocorticoids. We and others have previously shown that HSD11B1 and HSD11B2 are essential for the interconversion of the 11β-hydroxyl and 11-keto forms of the 11-oxygenated androgens (11, 20). We have also previously shown that 11KA4, but not 11OHA4, is a substrate for the key androgen-activating enzyme AKR1C3 (16, 17). Therefore, HSD11B2 plays a vital role in catalysing the conversion of adrenal-derived 11OHA4 to 11KA4, which can then be converted to the potent androgen 11KT by AKR1C3 (13). One of the main sites of AKR1C3 expression and peripheral androgen activation is adipose tissue (14, 15) which also abundantly expresses HSD11B1 (19, 22, 65). Paulukinas et. al. have previously observed that co-expression of HSD11B1 with AKR1C3 in differentiated SGBS adipocyte cells may prevent the accumulation of 11KT, thus suggesting that HSD11B1 may protect from androgen excess (17). Here, using a combination of *in vitro*, *in silico* and *ex vivo* studies, we demonstrate that co-expression of HSD11B1 with AKR1C3 modulates peripheral biosynthesis of 11KT, first, by competing

with AKR1C3 for the substrate 11KA4, and, second, by converting much of the resulting 11KT to the

less potent androgen 11OHT. We demonstrated that inhibition of HSD11B1 results in significantly

increased 11KT biosynthesis as shown by us *in vitro*, *in silico*, *ex vivo* and in a phase 2a study comparing

the effects of 35 days of treatment with the selective HSD11B1 inhibitor AZD4017 to placebo

treatment. Our data show that HSD11B1 inhibition does not only alter local, adipose tissue-specific

11KT concentrations, but also systemic levels, thus emphasising that peripheral tissues expressing

AKR1C3 are essential for 11KT biosynthesis (13). Collectively, our findings show that under normal

circumstances HSD11B1 plays an important novel role in preventing 11KT accumulation, in addition to

its well-known role in local glucocorticoid activation.

Our findings are of immediate clinical relevance, particularly in women, where circulating

concentrations of 11KT are similar to testosterone concentrations (3, 6, 66, 67). Therefore, increasing

11KT due to HSD11B1 inhibition markedly increases the total circulating pool of active androgens in

women. This increase in androgen load is significant given that 11KT has shown functional activity

comparable to testosterone (5, 62, 68) and increased circulating androgens have been linked to a

significantly greater risk of type 2 diabetes in women (43). Moreover, there is growing evidence that

androgens drive the risk of common metabolic disorders, including type 2 diabetes, hypertension,

cardiovascular disease and MASLD (15, 43, 69–72, 44, 46–52), in women with PCOS where androgen

excess is a hallmark (72). Notably, O'Reilly et. al. have previously shown that local AKR1C3-mediated

androgen activation in adipose tissue drives lipotoxicity in women with PCOS (15). They propose a

vicious cycle in which increased androgen generation in adipose tissue results in lipid accumulation,

increased fat mass and as a result systemic insulin resistance. This cycle is completed by the insulin-

induced stimulation of AKR1C3 expression in adipose, leading to increased androgen activation. Given

that 11KA4 is the preferred substrate for AKR1C3, increased insulin-induced AKR1C3 expression would

therefore lead to significantly more 11KT being produced, which would be further increased by

HSD11B1 inhibition (16, 17, 73).

Our finding of increased 11KT biosynthesis to clinically highly relevant concentrations occurring after

HSD11B1 inhibition therefore needs to be taken into account when considering HSD11B1 inhibition as

treatment for type 2 diabetes or metabolic syndrome in women, as the beneficial effects of reduced

local glucocorticoid activation could be offset by opposing metabolic effects due to increased 11KT

concentrations and AR signalling. This is highly relevant for the recently proposed use of HSD11B1

inhibitors in ameliorating the metabolically adverse consequences of mild autonomous cortisol excess

(MACS) due to adrenal adenomas (74) as affected patients are predominantly female (75).

Notably, quantification of AKR1C3 and HSD11B1 expression levels in subcutaneous and omental

adipose tissue revealed a degree of interpersonal variation. Our computational simulations revealed

that variations in the relative, as well as the absolute expression levels of HSD11B1 and AKR1C3 would

result in individual differences in the amount of 11KT produced as well as the magnitude of change

upon HSD11B1 inhibition. Indeed, this biological variation was reflected in both the *ex vivo* adipose

tissue conversion data and the serum concentrations of 11KT before and after administration of

AZD4017 in individuals with type 2 diabetes. Our simulations suggest that when considering 11KT

biosynthesis, individuals with higher HSD11B1 expression relative to AKR1C3 would be less responsive

to partial HSD11B1 inhibition. However, if HSD11B1 activity is completely abolished by inhibition then

the amount of 11KT produced is dependent of the absolute levels of AKR1C3 expressed. Increased

11KT concentrations are therefore not solely dependent on the inhibition of HSD11B1, but also on the

amount of AKR1C3 expressed. Notably, AKR1C3 expression correlates with BMI in subcutaneous, but

not omental, adipose tissue and its expression is upregulated by insulin (14, 15, 17). Women with

insulin resistance and obesity due to metabolic syndrome are therefore likely to have high expression

levels of AKR1C3 (14, 15), which would result in greater increases in 11KT upon inhibition of HSD11B1.

Thus, while such women may benefit from the local reduction of glucocorticoid activation by HSD11B1

inhibition in adipose tissue, they may be primed to generate more 11KT. Indeed, we noted that the

expression of AKR1C3 was higher than that of HSD11B1 in adipose tissue obtained from women with

obesity undergoing elective bariatric surgery. As a result, the conversion of 11KA4 yielded more 11KT

and 11OHT combined (both requiring AKR1C3 activity) than 11OHA4. In subcutaneous adipose tissue,

which had a lower HSD11B1:AKR1C3 ratio compared to omental adipose tissue, 11OHT was the most

abundant product, demonstrating effective conversion of 11KA4 to 11KT by AKR1C3 and subsequent

conversion of 11KT to 11OHT by HSD11B1. Similarly, we noted that in our clinical cohort, which consisted of individuals with type 2 diabetes (BMI mostly in the obese range), the mean circulating 11OHT concentration was ≥2-fold higher than previously reported for healthy individuals thus suggesting high peripheral AKR1C3 expression (3). Interestingly, a previous *ex vivo* study found that androgen exposure upregulates HSD11B1 expression in adipose tissue (76) emphasising that inactivation of 11-oxygenated androgens by HSD11B1 might be a protective mechanism, that no longer suffices once AKR1C3 is significantly upregulated, e.g. in the context of PCOS (15).

Increased 11KT concentrations following HSD11B1 inhibition are unlikely to be a concern in men as the metabolic effects of androgens are sexually dimorphic. In men, reduced androgen concentrations are associated with adverse metabolic consequences, including insulin resistance, type 2 diabetes and metabolic syndrome, and as such increased 11KT levels have the potential to be beneficial (43, 71, 77). Glucocorticoids are, however, also known to have sexually dimorphic effects, which are further complicated by crosstalk between glucocorticoids and androgens (78–80). Unfortunately, the vast majority of HSD11B1 inhibitor clinical studies did not report sex-specific outcomes for HSD11B1 inhibition (34–38). Larger sex-specific studies are therefore needed to fully assess the benefits of HSD11B1 inhibition in relation to metabolic conditions, while taking into account potential opposing androgenic effects in women, resulting from increased 11KT.

One recent study by Hardy *et al.* investigated the effects of HSD11B1 inhibition in women with obesity and Idiopathic Intracranial Hypertension (41). Twice daily administration of 400 mg AZD4017 for 12 weeks had no effect on fasting glucose, fasting insulin, insulin resistance or HbA1c, in agreement with our hypothesis that androgenic effects may counter beneficial metabolic effects of reduced glucocorticoid concentrations. Furthermore, this study found no effect of HSD11B1 inhibition on adipose or bone composition, but did report reduced cholesterol and increased HDL with HSD11B1 inhibition suggesting that women do exhibit some cardioprotective responses to HSD11B1 inhibition. Importantly, Hardy *et al*. also reported increased lean muscle mass with HSD11B1 inhibition, which correlated with increased testosterone and 11-oxygenated androgen precursors (11OHA4 and 11KA4) levels; however, they did not report 11KT concentrations.

Increased adrenal output of 11OHA4 is not unexpected with HSD11B1 inhibition, which leads to

reduced negative feedback on the HPA-axis, increased ACTH levels and increased adrenal output of

cortisol and adrenal androgen precursors (41). While we did not observe increased serum 11OHA4

concentrations in the current study, we cannot exclude that increased adrenal output of this precursor

contributed to the observed increase in 11KT biosynthesis. In fact, this is likely the case as circulating

11OHA4 concentrations were not reduced despite inhibition of HSD11B1 and the resulting significant

decrease in the 11OHA4/11KA4 ratio. Furthermore, the original clinical trial observed a significant

increase in serum DHEAS, a marker of adrenal C19 steroid biosynthesis, thus confirming the stimulation

of adrenal steroidogenesis (40). Despite the potential increase in adrenal 11OHA4 biosynthesis, our *in*

vitro, *ex vivo* and *in silico* data unequivocally show that HSD11B1 modulates the peripheral

biosynthesis of 11KT and that inhibition of HSD11B1 results in increased peripheral 11KT production

irrespective of adrenal 11OHA4 output.

This study highlights that HSD11B1 inhibition may not be as effective as previously anticipated for the

treatment of metabolic conditions in women due to the concomitant increase in the active androgen

pool. However, the current data does not call into question the safety or efficacy of HSD11B1 inhibitors

such as AZD4017, which has been shown to be well tolerated (39–41, 63, 81), but rather highlights

that the condition targeted by the use of HSD11B1 inhibitors needs to be carefully considered. To date,

AZD4017 has shown promise in the treatment of Idiopathic Intracranial Hypertension and wound

healing in adults with type 2 diabetes, thus warranting further studies (40, 41). It may also be

interesting to consider HSD11B1 inhibition in oophorectomized women. In this setting, the increased

biosynthesis of 11KT due to HSD11B1 inhibition may be beneficial in offsetting the androgen deficiency

due to oophorectomy, while reduced local glucocorticoid activation has the potential to protect

against glucocorticoid induced osteoporosis, especially against the backdrop of reduced estrogen

biosynthesis (82–84).

While the current study focusses on the regulation of 11-ketotestosterone biosynthesis by HSD11B1

in adipose tissue, HSD11B1 is also abundantly expressed in liver and as such the hepatic conversion of

11KA4 to 11OHA4 and 11KT to 11OHT may also regulate systemic 11KT levels, with inhibition of hepatic

HSD11B1 by AZD4017 contributing to the observed increase in 11KT levels (18, 19).

In summary, our findings indicate that HSD11B1 plays an essential role in modulating the amount of

11KT produced by AKR1C3 expressed in adipose tissue. As a result, inhibition of HSD11B1 does not just

reduce local glucocorticoid activation, but also alters the biosynthesis of the potent 11-oxygenated

androgen 11KT. In women, the significant increase in 11KT concentrations may have metabolically

opposing effects that attenuate the beneficial reduction of glucocorticoid activation in metabolic

tissue. It is therefore imperative that future HSD11B1 inhibitor studies consider both the glucocorticoid

and androgen pools and that sex-specific effects are systematically investigated and considered in the

design of future studies.

Data Availability

The data that support the findings of this study are available in the results and/or supplementary

material of this article. The computational model is available in SBML format upon request.

Conflict-of-interest

The authors declare no conflicts of interest

Author contributions

KHS, WA and LS were responsible for study concept and design. LS and IO conducted experiments. JS

performed all computational modelling. LS, IO, JS, LCG, AET, and KHS analysed data. AJC and RS

provided human adipose tissue. AT, RA and AF provided serum and urine from a phase 2b pilot trial

with AZD4017. AET and LCG conducted the analysis of clinical samples by LC-MS/MS and GC-MS,

respectively. KHS, LS and IO drafted the manuscript. WA, AT and JS critically reviewed the manuscript.

All authors reviewed and approved the final version of the manuscript. KS and WA obtained funding.

These authors contributed equally: LS and IO. LS was selected as the first of the two authors as she

was involved in the conceptualisation of the study.

Funding

This work has been supported by the National Research Foundation of South Africa (Grant Number

132503, to KHS); the Academy of Medical Sciences UK (Newton Advanced Fellowship NAF004\1002,

to KHS); the Wellcome Trust (Investigator Grant WT209492/Z/17/Z, to WA); a Medical Research

Council Confidence in Concept Award to AT (MC_PC_15046); and a SARChI DST/NRF grant (82813, to

JS).

References

1. Pretorius, E., Arlt, W., and Storbeck, K.-H. (2017) A new dawn for androgens: Novel lessons from 11-oxygenated C19 steroids. *Mol. Cell. Endocrinol.* **441**, 76–85

2. Turcu, A. F., Rege, J., Auchus, R. J., and Rainey, W. E. (2020) 11-Oxygenated androgens in health and disease. *Nat. Rev. Endocrinol.* **16**, 284–296

3. Schiffer, L., Kempegowda, P., Sitch, A. J., Adaway, J. E., Shaheen, F., Ebbehoj, A., Singh, S., McTaggart, M. P., O'Reilly, M. W., Prete, A., Hawley, J. M., Keevil, B. G., Bancos, I., Taylor, A. E., and Arlt, W. (2023) Classic and 11-oxygenated androgens in serum and saliva across adulthood: a cross-sectional study analyzing the impact of age, body mass index, and diurnal and menstrual cycle variation. *Eur. J. Endocrinol.* **188**, 1–15

4. Storbeck, K.-H. and O'Reilly, M. W. (2023) The clinical and biochemical significance of 11-oxygenated androgens in human health and disease. *Eur. J. Endocrinol.* **188**, R98–R109

5. Pretorius, E., Africander, D. J., Vlok, M., Perkins, M. S., Quanson, J. L., and Storbeck, K.-H. (2016) 11-ketotestosterone and 11-ketodihydrotestosterone in castration resistant prostate cancer: potent androgens which can no longer be ignored. *PLoS One* **11**, e0159867

6. Nanba, A. T., Rege, J., Ren, J., Auchus, R. J., Rainey, W. E., and Turcu, A. F. (2019) 11-Oxygenated C19 Steroids Do Not Decline with Age in Women. *J. Clin. Endocrinol. Metab.* **104**, 2615–2622

7. Skiba, M. A., Bell, R. J., Islam, R. M., Handelsman, D. J., Desai, R., and Davis, S. R. (2019) Androgens During the Reproductive Years: What Is Normal for Women? *J Clin Endocrinol Metab* **104**, 5382–5392

8. O'Reilly, M. W., Kempegowda, P., Jenkinson, C., Taylor, A. E., Quanson, J. L., Storbeck, K.-H. H., and Arlt, W. (2017) 11-oxygenated C19 steroids are the predominant androgens in polycystic ovary syndrome. *J. Clin. Endocrinol. Metab.* **102**, 840–848

9. Turcu, A. F., Nanba, A. T., Chomic, R., Upadhyay, S. K., Giordano, T. J., Shields, J. J., Merke, D. P., Rainey, W. E., and Auchus, R. J. (2016) Adrenal-derived 11-oxygenated 19-carbon steroids are the dominant androgens in classic 21-hydroxylase deficiency. *Eur. J. Endocrinol.* **174**, 601–

609

10.   Nowotny, H. F., Braun, L., Vogel, F., Bidlingmaier, M., Reincke, M., Tschaidse, L., Auer, M. K.,

Lottspeich, C., Wudy, S. A., Hartmann, M. F., Hawley, J., Adaway, J. E., Keevil, B., Schilbach, K.,

and Reisch, N. (2022) 11-Oxygenated C19 steroids are the predominant androgens

responsible for hyperandrogenemia in Cushing's disease. *Eur. J. Endocrinol.* **187**, 663–673

11.   Storbeck, K.-H., Bloem, L. M., Africander, D. J., Schloms, L., Swart, P., and Swart, A. C. (2013)

11β-hydroxydihydrotestosterone and 11-ketodihydrotestosterone, novel C19 steroids with

androgenic activity: a putative role in castration resistant prostate cancer? *Mol. Cell.*

Endocrinol. **377**, 135–146

12.   Snaterse, G., van Dessel, L. F., van Riet, J., Taylor, A. E., van der Vlugt-Daane, M., Hamberg, P.,

de Wit, R., Visser, J. A., Arlt, W., Lolkema, M. P., and Hofland, J. (2021) 11-Ketotestosterone is

the predominant active androgen in prostate cancer patients after castration. *JCI Insight* **6**,

e148507

13.   Storbeck, K.-H. (2022) A commentary on the origins of 11-ketotestosterone. *Eur J Endocrinol*

187, C5-8

14.   Quinkler, M., Sinha, B., Tomlinson, J. W., Bujalska, I. J., Stewart, P. M., and Arlt, W. (2004)

Androgen generation in adipose tissue in women with simple obesity - A site-specific role for

17β-hydroxysteroid dehydrogenase type 5. *J. Endocrinol.* **183**, 331–342

15.   O'Reilly, M. W., Kempegowda, P., Walsh, M., Taylor, A. E., Manolopoulos, K. N., Allwood, J.

W., Semple, R. K., Hebenstreit, D., Dunn, W. B., Tomlinson, J. W., Arlt, W., Reilly, M. W. O.,

Manolopoulos, N., and Warwick, B. (2017) AKR1C3-Mediated Adipose Androgen Generation

Drives Lipotoxicity in Women With Polycystic Ovary Syndrome. *J Clin Endocrinol Metab* **102**,

3327–3339

16.   Barnard, M., Quanson, J. L., Mostaghel, E., Pretorius, E., Snoep, J. L., and Storbeck, K.-H.

(2018) 11-Oxygenated androgen precursors are the preferred substrates for aldo-keto

reductase 1C3 (AKR1C3): Implications for castration resistant prostate cancer. *J. Steroid*

Biochem. Mol. Biol. **183**, 192–201

17.   Paulukinas, R. D., Mesaros, C. A., and Penning, T. M. (2022) Conversion of Classical and 11-

651   oxygenated Androgens by Insulin Induced AKR1C3 in a Model of Human PCOS Adipocytes.

652   *Endocrinology* **163**, bqac068

653   18.   Tomlinson, J. W., Walker, E. A., Bujalska, I. J., Draper, N., Lavery, G. G., Cooper, M. S.,

654   Hewison, M., and Stewart, P. M. (2004) 11β-Hydroxysteroid Dehydrogenase Type 1: A Tissue-

655   Specific Regulator of Glucocorticoid Response. *Endocr. Rev.* **25**, 831–866

656   19.   Gathercole, L. L., Lavery, G. G., Morgan, S. A., Cooper, M. S., Sinclair, A. J., Tomlinson, J. W.,

657   and Stewart, P. M. (2013) 11β-Hydroxysteroid Dehydrogenase 1: Translational and

658   Therapeutic Aspects. *Endocr. Rev.* **34**, 525–555

659   20.   Gent, R., du Toit, T., Bloem, L. M., and Swart, A. C. (2019) The 11β-hydroxysteroid

660   dehydrogenase isoforms: pivotal catalytic activities yield potent C11-oxy C 19 steroids with

661   11βHSD2 favouring 11-ketotestosterone, 11-ketoandrostenedione and 11-ketoprogesterone

662   biosynthesis. *J. Steroid Biochem. Mol. Biol.* **189**, 116–126

663   21.   Morgan, S. A., McCabe, E. L., Gathercole, L. L., Hassan-Smith, Z. K., Larner, D. P., Bujalska, I. J.,

664   Stewart, P. M., Tomlinson, J. W., and Lavery, G. G. (2014) 11β-HSD1 is the major regulator of

665   the tissue-specific effects of circulating glucocorticoid excess. *Proc. Natl. Acad. Sci. U. S. A.*

666   **111**, E2482–E2491

667   22.   Gregory, S., Hill, D., Grey, B., Ketelbey, W., Miller, T., Muniz-Terrera, G., and Ritchie, C. W.

668   (2020) 11β-hydroxysteroid dehydrogenase type 1 inhibitor use in human disease-a systematic

669   review and narrative synthesis. *Metabolism.* **108**, 154246

670   23.   Tiganescu, A., Hupe, M., Uchida, Y., Mauro, T., Elias, P. M., and Holleran, W. M. (2018) Topical

671   11β-Hydroxysteroid Dehydrogenase Type 1 Inhibition Corrects Cutaneous Features of

672   Systemic Glucocorticoid Excess in Female Mice. *Endocrinology* **159**, 547–556

673   24.   Doig, C. L., Fletcher, R. S., Morgan, S. A., McCabe, E. L., Larner, D. P., Tomlinson, J. W., Stewart,

674   P. M., Philp, A., and Lavery, G. G. (2017) 11β-HSD1 Modulates the Set Point of Brown Adipose

675   Tissue Response to Glucocorticoids in Male Mice. *Endocrinology* **158**, 1964–1976

25.   Masuzaki, H., Paterson, J., Shinyama, H., Morton, N. M., Mullins, J. J., Seckl, J. R., and Flier, J. S.

(2001) A transgenic model of visceral obesity and the metabolic syndrome. *Science (80-.).*

294, 2166–2170

26.   Masuzaki, H., Yamamoto, H., Kenyon, C. J., Elmquist, J. K., Morton, N. M., Paterson, J. M.,

Shinyama, H., Sharp, M. G. F., Fleming, S., Mullins, J. J., Seckl, J. R., and Flier, J. S. (2003)

Transgenic amplification of glucocorticoid action in adipose tissue causes high blood pressure

in mice. *J. Clin. Invest.* **112**, 83–90

27.   Kotelevtsev, Y., Holmes, M. C., Burchell, A., Houston, P. M., Schmoll, D., Jamieson, P., Best, R.,

Brown, R., Edwards, C. R., Seckl, J. R., and Mullins, J. J. (1997) 11beta-hydroxysteroid

dehydrogenase type 1 knockout mice show attenuated glucocorticoid-inducible responses

and resist hyperglycemia on obesity or stress. *Proc. Natl. Acad. Sci. U. S. A.* **94**, 14924–14929

28.   Hermanowski-Vosatka, A., Balkovec, J. M., Cheng, K., Chen, H. Y., Hernandez, M., Koo, G. C.,

Le Grand, C. B., Li, Z., Metzger, J. M., Mundt, S. S., Noonan, H., Nunes, C. N., Olson, S. H.,

Pikounis, B., Ren, N., Robertson, N., Schaeffer, J. M., Shah, K., Springer, M. S., Strack, A. M.,

Strowski, M., Wu, K., Wu, T. J., Xiao, J., Zhang, B. B., Wright, S. D., and Thieringer, R. (2005)

11β-HSD1 inhibition ameliorates metabolic syndrome and prevents progression of

atherosclerosis in mice. *J. Exp. Med.* **202**, 517–527

29.   Morgan, S. A., Gathercole, L. L., Hassan-Smith, Z. K., Tomlinson, J., Stewart, P. M., and Lavery,

G. G. (2022) 11β-HSD1 contributes to age-related metabolic decline in male mice. *J.*

Endocrinol. **255**, 117–129

30.   Rege, J., Garber, S., Conley, A. J., Elsey, R. M., Turcu, A. F., Auchus, R. J., and Rainey, W. E.

(2019) Circulating 11-oxygenated androgens across species. *J. Steroid Biochem. Mol. Biol.* **190**,

242–249

31.   Walker, B. R., Connacher, A. A., Lindsay, R. M., Webb, D. J., and Edwards, C. R. W. (1995)

Carbenoxolone increases hepatic insulin sensitivity in man: a novel role for 11-oxosteroid

reductase in enhancing glucocorticoid receptor activation. *J. Clin. Endocrinol. Metab.* **80**,

3155–3159

32.   Andrews, R. C., Rooyackers, O., and Walker, B. R. (2003) Effects of the 11 beta-hydroxysteroid

dehydrogenase inhibitor carbenoxolone on insulin sensitivity in men with type 2 diabetes. *J.*

Clin. Endocrinol. Metab. **88**, 285–291

33. Sandeep, T. C., Andrew, R., Homer, N. Z. M., Andrews, R. C., Smith, K., and Walker, B. R. (2005)

Increased in vivo regeneration of cortisol in adipose tissue in human obesity and effects of the

11beta-hydroxysteroid dehydrogenase type 1 inhibitor carbenoxolone. *Diabetes* **54**, 872–879

34. Rosenstock, J., Banarer, S., Fonseca, V. A., Inzucchi, S. E., William, S., Wenqing, Y., Hollis, G.,

Flores, R., Levy, R., Williams, W. V., Seckl, J. R., and Huber, R. (2010) The 11-beta-

hydroxysteroid dehydrogenase type 1 inhibitor INCB13739 improves hyperglycemia in

patients with type 2 diabetes inadequately controlled by metformin monotherapy. *Diabetes*

Care **33**, 1516–1522

35. Feig, P. U., Shah, S., Hermanowski-Vosatka, A., Plotkin, D., Springer, M. S., Donahue, S., Thach,

C., Klein, E. J., Lai, E., and Kaufman, K. D. (2011) Effects of an 11β-hydroxysteroid

dehydrogenase type 1 inhibitor, MK-0916, in patients with type 2 diabetes mellitus and

metabolic syndrome. *Diabetes. Obes. Metab.* **13**, 498–504

36. Shah, S., Hermanowski-Vosatka, A., Gibson, K., Ruck, R. A., Jia, G., Zhang, J., Hwang, P. M. T.,

Ryan, N. W., Langdon, R. B., and Feig, P. U. (2011) Efficacy and safety of the selective 11β-

HSD-1 inhibitors MK-0736 and MK-0916 in overweight and obese patients with hypertension.

J. Am. Soc. Hypertens. **5**, 166–176

37. Heise, T., Morrow, L., Hompesch, M., Häring, H. U., Kapitza, C., Abt, M., Ramsauer, M.,

Magnone, M. C., and Fuerst-Recktenwald, S. (2014) Safety, efficacy and weight effect of two

11β-HSD1 inhibitors in metformin-treated patients with type 2 diabetes. *Diabetes. Obes.*

Metab. **16**, 1070–1077

38. Yadav, Y., Dunagan, K., Khot, R., Venkatesh, S. K., Port, J., Galderisi, A., Cobelli, C., Wegner, C.,

Basu, A., Carter, R., and Basu, R. (2022) Inhibition of 11β-Hydroxysteroid dehydrogenase-1

with AZD4017 in patients with nonalcoholic steatohepatitis or nonalcoholic fatty liver disease:

A randomized, double-blind, placebo-controlled, phase II study. *Diabetes. Obes. Metab.* **24**,

881–890

731 39. Abbas, A., Schini, M., Ainsworth, G., Brown, S. R., Oughton, J., Crowley, R. K., Cooper, M. S.,

732   Fairclough, R. J., Eastell, R., and Stewart, P. M. (2022) Effect of AZD4017, a Selective 11β-HSD1

733   Inhibitor, on Bone Turnover Markers in Postmenopausal Osteopenia. *J. Clin. Endocrinol.*

734   *Metab.* **107**, 2026–2035

735 40. Ajjan, R. A., Hensor, E. M. A., Del Galdo, F., Shams, K., Abbas, A., Fairclough, R. J., Webber, L.,

736   Pegg, L., Freeman, A., Taylor, A. E., Arlt, W., Morgan, A. W., Tahrani, A. A., Stewart, P. M.,

737   Russell, D. A., and Tiganescu, A. (2022) Oral 11β-HSD1 inhibitor AZD4017 improves wound

738   healing and skin integrity in adults with type 2 diabetes mellitus: a pilot randomized

739   controlled trial. *Eur. J. Endocrinol.* **186**, 441–455

740 41. Hardy, R. S., Botfield, H., Markey, K., Mitchell, J. L., Alimajstorovic, Z., Westgate, C. S. J.,

741   Sagmeister, M., Fairclough, R. J., Ottridge, R. S., Yiangou, A., Storbeck, K. H. H., Taylor, A. E.,

742   Gilligan, L. C., Arlt, W., Stewart, P. M., Tomlinson, J. W., Mollan, S. P., Lavery, G. G., and

743   Sinclair, A. J. (2021) 11βHSD1 Inhibition with AZD4017 Improves Lipid Profiles and Lean

744   Muscle Mass in Idiopathic Intracranial Hypertension. *J. Clin. Endocrinol. Metab.* **106**, 174–187

745 42. Bianzano, S., Nordaby, M., Plum-Mörschel, L., Peil, B., and Heise, T. (2023) Safety, tolerability,

746   pharmacodynamics and pharmacokinetics following once-daily doses of BI 187004, an

747   inhibitor of 11 beta-hydroxysteroid dehydrogenase-1, over 28 days in patients with type 2

748   diabetes mellitus and overweight or obesity. *Diabetes, Obes. Metab.* **25**, 832–843

749 43. O'Reilly, M. W., Glisic, M., Kumarendran, B., Subramanian, A., Manolopoulos, K. N., Tahrani, A.

750   A., Keerthy, D., Muka, T., Toulis, K. A., Hanif, W., Thomas, G. N., Franco, O. H., Arlt, W., and

751   Nirantharakumar, K. (2019) Serum testosterone, sex hormone-binding globulin and sex-

752   specific risk of incident type 2 diabetes in a retrospective primary care cohort. *Clin.*

753   *Endocrinol. (Oxf).* **90**, 145–154

754 44. Fauser, B. C. J. M., Tarlatzis, Fauser, Chang, Aziz, Legro, Dewailly, Franks, Balen, Bouchard,

755   Dahlgren, Devoto, Diamanti, Dunaif, Filicori, Homburg, Ibanez, Laven, Magoffin, Nestler,

756   Norman, Pasquali, Pugeat, Strauss, Tan, Taylor, Wild, Wild, Guzik, Ehrmann, and Lobo. (2004)

757   Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic

ovary syndrome (PCOS). *Hum. Reprod.* **19**, 41–47

45. Legro, R. S., Arslanian, S. A., Ehrmann, D. A., Hoeger, K. M., Murad, M. H., Pasquali, R., and

Welt, C. K. (2013) Diagnosis and treatment of polycystic ovary syndrome: An endocrine

society clinical practice guideline. *J. Clin. Endocrinol. Metab.* **98**, 4565–4592

46. McCartney, C. R. and Marshall, J. C. (2016) Polycystic Ovary Syndrome. *N. Engl. J. Med.* **375**,

54–64

47. Legro, R. S., Kunselman, A. R., Dodson, W. C., and Dunaif, A. (1999) Prevalence and Predictors

of Risk for Type 2 Diabetes Mellitus and Impaired Glucose Tolerance in Polycystic Ovary

Syndrome: A Prospective, Controlled Study in 254 Affected Women. *J. Clin. Endocrinol. Metab.*

84, 165–169

48. Randeva, H. S., Tan, B. K., Weickert, M. O., Lois, K., Nestler, J. E., Sattar, N., and Lehnert, H.

(2012) Cardiometabolic Aspects of the Polycystic Ovary Syndrome. *Endocr. Rev.* **33**, 812–841

49. Jones, H., Sprung, V. S., Pugh, C. J. A., Daousi, C., Irwin, A., Aziz, N., Adams, V. L., Thomas, E. L.,

Bell, J. D., Kemp, G. J., and Cuthbertson, D. J. (2012) Polycystic ovary syndrome with

hyperandrogenism is characterized by an increased risk of hepatic steatosis compared to

nonhyperandrogenic PCOS phenotypes and healthy controls, independent of obesity and

insulin resistance. *J. Clin. Endocrinol. Metab.* **97**, 3709–3716

50. Rocha, A. L. L., Faria, L. C., Guimarães, T. C. M., Moreira, G. V., Cândido, A. L., Couto, C. A., and

Reis, F. M. (2017) Non-alcoholic fatty liver disease in women with polycystic ovary syndrome:

systematic review and meta-analysis. *J. Endocrinol. Invest.* **40**, 1279–1288

51. Cai, J., Wu, C. H., Zhang, Y., Wang, Y. Y., Xu, W. D., Lin, T. C., Li, S. X., Wang, L. H., Zheng, J.,

Sun, Y., Liu, W., and Tao, T. (2017) High-free androgen index is associated with increased risk

of non-alcoholic fatty liver disease in women with polycystic ovary syndrome, independent of

obesity and insulin resistance. *Int. J. Obes.* **41**, 1341–1347

52. Kumarendran, B., O'Reilly, M. W., Manolopoulos, K. N., Toulis, K. A., Gokhale, K. M., Sitch, A.

J., Wijeyaratne, C. N., Coomarasamy, A., Arlt, W., and Nirantharakumar, K. (2018) Polycystic

ovary syndrome, androgen excess, and the risk of nonalcoholic fatty liver disease in women: A

longitudinal study based on a United Kingdom primary care database. *PLOS Med.* **15**,

e1002542

53.   Matsuura, K., Shiraishi, H., Hara, A., Sato, K., Deyashiki, Y., Ninomiya, M., and Sakai, S. (1998)

Identification of a Principal mRNA Species for Human 3 -Hydroxysteroid Dehydrogenase

Isoform (AKR1C3) That Exhibits High Prostaglandin D2 11-Ketoreductase Activity. *J. Biochem.*

124, 940–946

54.   Pfaffl, M. W. (2001) A new mathematical model for relative quantification in real-time RT-PCR.

Nucleic Acids Res. **29**, e45

55.   Willems, E., Leyns, L., and Vandesompele, J. (2008) Standardization of real-time PCR gene

expression data from independent biological replicates. *Anal. Biochem.* **379**, 127–129

56.   Guan, H. and Yang, K. (2008) RNA isolation and real-time quantitative RT-PCR. *Methods Mol.*

Biol. **456**, 259–270

57.   Quanson, J. L., Stander, M. A., Pretorius, E., Jenkinson, C., Taylor, A. E., and Storbeck, K.-H. H.

(2016) High-throughput analysis of 19 endogenous androgenic steroids by ultra-performance

convergence chromatography tandem mass spectrometry. *J. Chromatogr. B Anal. Technol.*

Biomed. Life Sci. **1031**, 131–138

58.   Schiffer, L., Shaheen, F., Gilligan, L. C., Storbeck, K. H., Hawley, J. M., Keevil, B. G., Arlt, W., and

Taylor, A. E. (2022) Multi-steroid profiling by UHPLC-MS/MS with post-column infusion of

ammonium fluoride. *J. Chromatogr. B Anal. Technol. Biomed. Life Sci.* **1209**, 123413

59.   Foster, M. A., Taylor, A. E., Hill, N. E., Bentley, C., Bishop, J., Gilligan, L. C., Shaheen, F., Bion, J.

F., Fallowfield, J. L., Woods, D. R., Bancos, I., Midwinter, M. M., Lord, J. M., and Arlt, W. (2020)

Mapping the Steroid Response to Major Trauma From Injury to Recovery: A Prospective

Cohort Study. *J. Clin. Endocrinol. Metab.* **105**, 925–937

60.   Chadwick, C. A., Owen, L. J., and Keevil, B. G. (2005) Development of a method for the

measurement of dehydroepiandrosterone sulphate by liquid chromatography-tandem mass

spectrometry. *Ann. Clin. Biochem.* **42**, 468–474

61.   Shackleton, C. H. L. (1986) Profiling steroid hormones and urinary steroids. *J Chromatogr* **379**,

91–156

62.   Snaterse, G., Mies, R., van Weerden, W. M., French, P. J., Jonker, J. W., Houtsmuller, A. B., van

Royen, M. E., Visser, J. A., and Hofland, J. (2022) Androgen receptor mutations modulate

activation by 11-oxygenated androgens and glucocorticoids. *Prostate Cancer Prostatic Dis.*

63.   Scott, J. S., Bowker, S. S., Deschoolmeester, J., Gerhardt, S., Hargreaves, D., Kilgour, E., Lloyd,

A., Mayers, R. M., McCoull, W., Newcombe, N. J., Ogg, D., Packer, M. J., Rees, A., Revill, J.,

Schofield, P., Selmi, N., Swales, J. G., and Whittamore, P. R. O. (2012) Discovery of a potent,

selective, and orally bioavailable acidic 11β-hydroxysteroid dehydrogenase type 1 (11β-HSD1)

inhibitor: discovery of 2-[(3S)-1-[5-(cyclohexylcarbamoyl)-6-propylsulfanylpyridin-2-yl]-3-

piperidyl]acetic acid (AZD4017). *J. Med. Chem.* **55**, 5951–5964

64.   Barnard, L., Nikolaou, N., Louw, C., Schiffer, L., Gibson, H., Gilligan, L. C., Gangitano, E., Snoep,

J., Arlt, W., Tomlinson, J. W., and Storbeck, K. H. (2020) The A-ring reduction of 11-

ketotestosterone is efficiently catalysed by AKR1D1 and SRD5A2 but not SRD5A1. *J. Steroid*

Biochem. Mol. Biol. **202**, 105724

65.   Tomlinson, J. W., Walker, E. A., Bujalska, I. J., Draper, N., Lavery, G. G., Cooper, M. S.,

Hewison, M., and Stewart, P. M. (2004) 11beta-hydroxysteroid dehydrogenase type 1: a

tissue-specific regulator of glucocorticoid response. *Endocr Rev* **25**, 831–866

66.   Davio, A., Woolcock, H., Nanba, A. T., Rege, J., O'Day, P., Ren, J., Zhao, L., Ebina, H., Auchus, R.,

Rainey, W. E., and Turcu, A. F. (2020) Sex Differences in 11-Oxygenated Androgen Patterns

Across Adulthood. *J. Clin. Endocrinol. Metab.* **105**, e2921–e2929

67.   Skiba, M. A., Bell, R. J., Islam, R. M., Handelsman, D. J., Desai, R., and Davis, S. R. (2019)

Androgens During the Reproductive Years: What Is Normal for Women? *J. Clin. Endocrinol.*

Metab. **104**, 5382–5392

68.   Handelsman, D. J., Cooper, E. R., and Heather, A. K. (2022) Bioactivity of 11 keto and hydroxy

androgens in yeast and mammalian host cells. *J. Steroid Biochem. Mol. Biol.* **218**, 106049

69.   Legro, R. S., Arslanian, S. A., Ehrmann, D. A., Hoeger, K. M., Murad, M. H., Pasquali, R., and

Welt, C. K. (2013) Diagnosis and Treatment of Polycystic Ovary Syndrome: An Endocrine

Society Clinical Practice Guideline. *J. Clin. Endocrinol. Metab.* **98**, 4565–4592

70.   O'Reilly, M. W., Taylor, A. E., Crabtree, N. J., Hughes, B. A., Capper, F., Crowley, R. K., Stewart,

P. M., Tomlinson, J. W., and Arlt, W. (2014) Hyperandrogenemia predicts metabolic

phenotype in polycystic ovary syndrome: The utility of serum androstenedione. *J. Clin.*

Endocrinol. Metab. **99**, 1027–1036

71.   Schiffer, L., Kempegowda, P., Arlt, W., and O'Reilly, M. W. (2017) MECHANISMS IN

ENDOCRINOLOGY: The sexually dimorphic role of androgens in human metabolic disease. *Eur.*

J. Endocrinol. **177**, R125–R143

72.   Kempegowda, P., Melson, E., Manolopoulos, K. N., Arlt, W., and O'Reilly, M. W. (2020)

Implicating androgen excess in propagating metabolic disease in polycystic ovary syndrome.

Ther. Adv. Endocrinol. Metab. **11**, 2042018820934319

73.   Paulukinas, R. D. and Penning, T. M. (2023) Insulin-Induced AKR1C3 Induces Fatty Acid

Synthase in a Model of Human PCOS Adipocytes. *Endocrinology* **164**, bqad033

74.   Oda, S., Ashida, K., Uchiyama, M., Sakamoto, S., Hasuzawa, N., Nagayama, A., Wang, L.,

Nagata, H., Sakamoto, R., Kishimoto, J., Todaka, K., Ogawa, Y., Nakanishi, Y., and Nomura, M.

(2021) An Open-label Phase I/IIa Clinical Trial of 11β-HSD1 Inhibitor for Cushing's Syndrome

and Autonomous Cortisol Secretion. *J. Clin. Endocrinol. Metab.* **106**, E3865–E3880

75.   Prete, A., Subramanian, A., Bancos, I., Chortis, V., Tsagarakis, S., Lang, K., Macech, M.,

Delivanis, D. A., Pupovac, I. D., Reimondo, G., Marina, L. V., Deutschbein, T., Balomenaki, M.,

O'Reilly, M. W., Gilligan, L. C., Jenkinson, C., Bednarczuk, T., Zhang, C. D., Dusek, T.,

Diamantopoulos, A., Asia, M., Kondracka, A., Li, D., Masjkur, J. R., Quinkler, M., Ueland, G.,

Dennedy, M. C., Beuschlein, F., Tabarin, A., Fassnacht, M., Ivović, M., Terzolo, M., Kastelan, D.,

Young, W. F., Manolopoulos, K. N., Ambroziak, U., Vassiliadi, D. A., Taylor, A. E., Sitch, A. J.,

Nirantharakumar, K., Arlt, W., Glöckner, S., Sinnott, R. O., Stell, A., Fragoso, M. C. B. V.,

Simunov, B., Cazenave, S., Haissaguerre, M., Bertherat, J., Libé, R., Kienitz, T., Eisenhofer, G.,

Brugger, C., Reincke, M., Riester, A., Spyroglou, A., Burger-Stritt, S., Hahner, S., Kroiss, M.,

Ronchi, C. L., Palimeri, S., Tsirou, I., Basile, V., Ingargiola, E., Canu, L., Mannelli, M., Ettaieb, H.,

Haak, H. R., Kerkhofs, T. M., Biehl, M., Feelders, R. A., Hofland, J., Hofland, L. J., Grytaas, M. A.,

Husebye, E. S., Zawierucha, M., Paiva, I., Sajwani, A., Sherlock, M., Crowley, R. K., Deeks, J. J.,

Hughes, B. A., Ivison, H. E., O'Neil, D. M., Papathomas, T. G., Ronchi, C. L., Shackleton, C. H. L.,

Ronchi, C. L., Sutcliffe, R. P., Guest, P., Skordilis, K., Bancos, C., Chang, A., Davidge-Pitts, C. J.,

Erickson, D., Natt, N., Nippoldt, T. B., Thomas, M., and Shackleton, C. H. L. (2022)

Cardiometabolic Disease Burden and Steroid Excretion in Benign Adrenal Tumors : A Cross-

Sectional Multicenter Study. *Ann. Intern. Med.* **175**, 325–334

76. Zhu, L., Hou, M., Sun, B., Burén, J., Zhang, L., Yi, J., Hernell, O., and Li, X. (2010) Testosterone

stimulates adipose tissue 11beta-hydroxysteroid dehydrogenase type 1 expression in a depot-

specific manner in children. *J. Clin. Endocrinol. Metab.* **95**, 3300–3308

77. Pivonello, R., Menafra, D., Riccio, E., Garifalos, F., Mazzella, M., De Angelis, C., and Colao, A. A.

(2019) Metabolic disorders and male hypogonadotropic hypogonadism. *Front. Endocrinol.*

(Lausanne). **10**, 345

78. Kroon, J., Pereira, A. M., and Meijer, O. C. (2020) Glucocorticoid Sexual Dimorphism in

Metabolism: Dissecting the Role of Sex Hormones. *Trends Endocrinol. Metab.* **31**, 357–367

79. Ruiz, D., Padmanabhan, V., and Sargis, R. M. (2020) Stress, sex, and sugar: Glucocorticoids and

sex-steroid crosstalk in the sex-specific misprogramming of metabolism. *J. Endocr. Soc.* **4**,

bvaa087

80. Kaikaew, K., Grefhorst, A., and Visser, J. A. (2021) Sex Differences in Brown Adipose Tissue

Function: Sex Hormones, Glucocorticoids, and Their Crosstalk. *Front. Endocrinol. (Lausanne).*

12, 357

81. Markey, K., Mitchell, J., Botfield, H., Ottridge, R. S., Matthews, T., Krishnan, A., Woolley, R.,

Westgate, C., Yiangou, A., Alimajstorovic, Z., Shah, P., Rick, C., Ives, N., Taylor, A. E., Gilligan, L.

C., Jenkinson, C., Arlt, W., Scotton, W., Fairclough, R. J., Singhal, R., Stewart, P. M., Tomlinson,

J. W., Lavery, G. G., Mollan, S. P., and Sinclair, A. J. (2020) 11β-Hydroxysteroid dehydrogenase

type 1 inhibition in idiopathic intracranial hypertension: a double-blind randomized controlled

trial. *Brain Commun.* **2**, fcz050

82.   Shifren, J. L. (2002) Androgen deficiency in the oophorectomized woman. *Fertil. Steril.* **77**, 60–

894   62

895   83.   Stanczyk, F. Z., Chaikittisilpa, S., Sriprasert, I., Rafatnia, A., Nadadur, M., and Mishell, D. R.

896   (2019) Circulating androgen levels before and after oophorectomy in premenopausal and

897   postmenopausal women. *Climacteric* **22**, 169–174

898   84.   Hardy, R. S., Zhou, H., Seibel, M. J., and Cooper, M. S. (2018) Glucocorticoids and Bone:

899   Consequences of Endogenous and Exogenous Excess and Replacement Therapy. *Endocr. Rev.*

900   **39**, 519–548

Table and figures:

Figure 1: Schematic showing the peripheral metabolism of cortisol and 11OHA4. Cortisol is converted

to its inactive form, cortisone, by HSD11B2 expressed in mineralocorticoid target tissues. Cortisone is

in turn converted back to cortisol by HSD11B1 expressed in glucocorticoid target tissues (e.g., adipose).

Like cortisol, adrenal derived 11OHA4 is a substrate for HSD11B2, yielding 11KA4. 11KA4 is a substrate

for both the androgen activating enzyme AKR1C3 and HSD11B1, both of which are expressed in

adipose tissue. Conversion of 11KA4 to 11OHA4 by HSD11B1 prevents the conversion of 11KA4 to the

potent 11-oxygenated androgen, 11KT. The 11KT that is produced by AKR1C3 is also a substrate for

HSD11B1, yielding the substantially less active androgen, 11OHT.

Figure 2: Increased HSD11B1:AKR1C3 ratios modulate the biosynthesis of 11KT. (A) qPCR data
913

confirming the increase in HSD11B1 expression relative to that of AKR1C3 at the different
914

HSD11B1:AKR1C3 ratios. **(B)** The conversion of cortisone (0.1 µM) to cortisol by increasing ratios of
915

HSD11B1:AKR1C3 after 24 hours **(C)**. The conversion of 11KA4 (0.1 µM) to 11OHA4, 11KT and 11OHT
916

by increasing ratios of HSD11B1:AKR1C3 after 24 hours. For **B** and **C** the experimental data from the
917

ratio experiments are shown as points with error bars, while the computational models' predictions
918

are shown with the solid connecting lines. All results are shown as means ± SEM of three independent
919

experiments.
920

Figure 3: Inhibition of HSD11B1 by carbenoxolone (CBX) inhibits the biosynthesis of cortisol, but allows for increased AKR1C3-mediated production of 11KT when HSD11B1 and AKR1C3 are co-expressed. HEK293 cells expressing HSD11B1 and AKR1C3 were combined at a ratio of 1:1 and treated with either 100 nM cortisone **(A & B)** and 100 nM 11KA4 **(C & D)** with and without CBX (10 μM) or 100 nM cortisone and 10 nM 11KA4 **(E & F)** with or without CBX (10 μM). The experimental data from the time course experiment is shown as points with error bars, while the computational models' predictions are shown with the solid lines. All results are shown as means ± SD of a single biological experiment performed in triplicate. This enabled the model to predict the outcomes from one experiment with the same transfection efficiency.

Figure 4: Computational model of the effect of varying HSD11B1:AKR1C3 ratios and different levels of HSD11B1 inhibition on the resulting concentrations of 11OHA4 (A), 11KT (B) and 11OHT (C). The model varies that ratio of HSD11B1:AKR1C3 as in Figure 2 and estimates the effect of different levels of HSD11B1 inhibition on the conversion of 10 nM 11KA4 after 24 hours as in Figures 2 and 3.

Figure 5: Inhibition of HSD11B1 by carbenoxolone (CBX) results in decreased cortisol production (A) and an accompanying decrease in transactivation of the glucocorticoid receptor (B), but increased production of 11KT (C) and transactivation of the androgen receptor (D). HEK293 cells expressing HSD11B1 and AKR1C3 were combined at a ratio of 1:1 and treated with 100 nM cortisone and 10 nM 11KA4 with or without CBX (10 µM). After 24 hours the conditioned media was transferred to HEK293 cells transiently transfected with either a plasmid expressing the glucocorticoid receptor (GR) and a luciferase reporter construct or a plasmid expressing the androgen receptor (AR) and the same luciferase reporter construct. Luciferase was assayed after an additional 24 hour incubation and is shown as the fold-induction relative to a vehicle control. All results are shown as means ± SEM of three biological experiments. P values were calculated using unpaired t-tests (*P <0.05; **P<0.01; ***P<0.001; ****P <0.0001).

Figure 6: Inhibition of HSD11B1 by AZD4017 in human subcutaneous (A, F, G) and omental (B, H, I) adipose tissue results in the increased production of 11KT. Adipose tissue was incubated *ex vivo* with 100 nM cortisone (**A, B**) or 11KA4 (**F-I**) with or without AZD4017 (100 nM) for 72 hours. The mathematical model's prediction of the conversion over time is shown next to the experimentally determined end point data after the 72 hour incubation. Expression of AKR1C3 and HSD11B1 in subcutaneous (**C**) and omental (**D**) adipose tissue was measured by qPCR. The expression ratio of HSD11B1:AKR1C3 were determined for both adipose tissue subtypes (**E**). P values were calculated using Wilcoxon signed-rank tests (*P <0.05; **P<0.01; ***P<0.001).

Figure 7: Inhibition of HSD11B1 by oral administration of AZD4017 (400 mg twice daily) for 35 days

(n=11) changes the serum (A-F) and urinary (G-I) profile of 11-oxygenated androgens and their

metabolites, with significant increases in the circulating concentration of the potent androgen 11KT.

P values were calculated using Wilcoxon signed-rank tests (*P <0.05; **P<0.01; ***P<0.001). Serum

concentrations of classic androgens are shown in **Supplementary Figure 7**.

964 **Table 1: Kinetic constants ($K_{m, app}$; $V_{max, app}$; V_{max}/K_m) determined for HSD11B1.**

Reaction	$K_{m, app}$ (substrate) (µM)	$K_{m, app}$ (product) (µM)	$V_{max, app}$ (forward) (µM/h)[*]	$V_{max, app}$ (reverse) 1/h[a] or (µM/h)[b]	V_{max}/K_m[#]
Cortisone -> Cortisol	0.9	-	0.12	0.01[a]	0.13
11KA4 -> 11OHA4	1	2.46	0.35	0.03[b]	0.35
11KT -> 11OHT	0.21	-	0.05	-	0.24

965 [*]$V_{max, app}$ values are dependent on transfection efficiency and differ between experiments. We included the
966 conversion of 100 nM 11KA4 to 11OHA4 in all experiments and used the initial rate to normalise the transfection
967 efficiency, which was set to 1 for the values reported above.
968 [#]V_{max}/K_m values are reported for the substrate and forward reaction.