Title

Broca's area, variation and taxic diversity in early *Homo* from Koobi Fora (Kenya)

Authors

Amélie Beaudet[1,2,3], Edwin de Jager[1]

Author affiliations

[1]Laboratoire de Paléontologie, Évolution, Paléoécosystèmes et Paléoprimatologie

(PALEVOPRIM), UMR 7262 CNRS & University of Poitiers, 6 Rue Michel Brunet, 86000

Poitiers, France

[2]Department of Archaeology, University of Cambridge, Henry Wellcome Building,

Fitzwilliam St, Cambridge CB2 1QH

[3]School of Geography, Archaeology and Environmental Studies, University of the

Witwatersrand, Private Bag 3, WITS 2050, South Africa

Corresponding author

*Amélie Beaudet

Email: beaudet.amelie@gmail.com

Abstract

Because brain tissues rarely fossilize, pinpointing when and how modern human cerebral traits

emerged in the hominin lineage is particularly challenging. The fragmentary nature of the fossil

material, coupled with the difficulty of characterizing such a complex organ, have been the

source of long-standing debates. Prominent among them is the uncertainties around the derived

or primitive state of the brain organization in the earliest representatives of the genus *Homo*,

more particularly in key areas such as the Broca's area. By revisiting a particularly well-

preserved fossil endocast from the Turkana basin (Kenya) attributed to early *Homo*, here we

confirm that humans in Africa had a primitive organization of the Broca's area ca. 1.9 million

years ago. Additionally, our description of KNM-ER 3732 adds further information about the

variation pattern of the inferior frontal gyrus in fossil hominins, with implications for early

Homo taxic diversity (i.e., one or two *Homo* species at Koobi Fora) and the nature of the

mechanisms involved in the emergence of derived cerebral traits.

Main text

Introduction

The modern human brain is an exceptionally complex, highly specialized and extremely costly machinery. Because of the fragmentary nature of the hominin fossil record, assessing when and how changes in the brain of our ancestors happened, and inferring any related functional, behavioral and metabolic consequences, is particularly challenging (Zollikofer and Ponce de León, 2013). Nonetheless, reconstructing the chronological and taxonomical context of the emergence of derived cerebral traits is a prerequisite for disentangling underlying evolutionary processes. For instance, local reorganization of specific areas in an overall primitive hominin brain would support a mosaic-like evolutionary pattern (e.g., Holloway et al., 2004), and raise essential questions on the role of selection pressure (or absence of; rev. in Beaudet, 2021). Beyond the value of such information on the origins of the human brain, the assumption that *Homo* developed a uniquely complex brain organization (e.g., Tobias, 1987) requires further evidence.

Although brains rarely fossilize, it is possible to glean structural information about the evolutionary history of the hominin brain by studying sulcal imprints in fossil brain endocasts (Neubauer et al., 2014). In this regard, the Broca's cap has been the focus of much interest in paleoneurology due to striking structural differences between extant human and chimpanzee brains and endocasts, and the implication of this area in articulated language (rev. in Beaudet, 2017). Ponce de León et al. (2021) thoroughly examined brain endocasts of *Homo* specimens in eastern Africa and Eurasia and demonstrated that the organization of the Broca's area in the earliest representatives of the genus before 1.5 Ma was primitive. Because the imprints of the Broca's cap in endocasts are not always readable, they used the coronal suture and surrounding sulcal imprints as a proxy to identify frontal lobe expansion (i.e., derived condition). Unfortunately, in some of the oldest and key specimens from Africa that could represent the >1.5 Ma condition (e.g., KNM-ER 1470) the interpretations remain inconclusive.

Testing the hypothesis of Ponce de León et al. (2021) of a primitive brain in the earliest representatives of the genus *Homo* before 1.5 Ma thus necessitates (i) an excellent preservation of very fine neuroanatomical details in fossil endocasts, and (ii) reliable information on their taxonomic identity (i.e., *Homo*) and stratigraphic context (i.e., before/after 1.5 Ma). The hominin specimen KNM-ER 3732 from Kenya, which fulfills these requirements, has the potential to shed new light on this conundrum. KNM-ER 3732 was discovered during the 1974-1975 field programme in the Koobi Fora Formation (area 115), east of Lake Turkana in Kenya

(Leakey, 1976; Wood, 1991). KNM-ER 3732 consists of a calotte, left zygoma and a natural endocast. The expanded neurocranium and robust upper face supports an attribution to *Homo* sp. indet. (Wood, 1991). The specimen was lying below the KBS Tuff of the upper Burgi Member that is dated to 1.87 million years ago, which provides a minimum age (Feibel et al., 1983, 2009). Accordingly, if KNM-ER 3732 shows a derived cerebral condition, the hypothesis of >1.5 Ma *Homo* being associated to a primitive organization of the Broca's area is falsified. Here we provide a comparative study of the natural endocast of pre-1.5 Ma *Homo* specimen KNM-ER 3732 to test the hypothesis of a late emergence of a modern Broca's cap in the hominin lineage.

Results

KNM-ER 3732 offers a glimpse of early *Homo* prefrontal organization. The natural endocast of KNM-ER 3732, initially described by Holloway et al. (2004), preserves the dorsal part but misses the frontal pole, the occipital and temporal lobes, as well as the entire ventral surface (Fig. 1). Wood (1991: 132) identified an "irregular bony excrescence on the surface of the right parietal at the level of lambda and 32 mm from the midline. It projects 6 mm from the surface of the bone, and may be an example of myositis ossificans associated with damage to the overlying temporal muscle". Because of the location (right parietal) and nature (traumatic) of the outgrowth, it does not affect our area of study (i.e., left frontal bone) (Walczak et al., 2015). The gyral and sulcal details are well-preserved, particularly in the prefrontal region. The superior, middle and inferior frontal sulci are visible on both hemispheres. On the left hemisphere, the Broca's cap is prominent. A vertical groove, identified as the precentral sulcus, separates the Broca's cap in half (Fig. 2A). On both hemispheres, the central sulcus intersects the inter-hemispheric scissure and the post-central sulcus seems to be connected to the lateral fissure. The superior temporal sulcus nearly intersects the lateral fissure on the right hemisphere. The intra-parietal sulcus can be found in the parieto-occipital region of both hemispheres. The examination of the internal surface of the braincase (Fig. 2B) confirms the pattern described on the natural endocast.

KNM-ER 3732 has a primitive prefrontal cortex. When compared to extant human and chimpanzee sulcal patterns, KNM-ER 3732 closely resembles the latter (Fig. 2C-D). In extant humans, the inferior frontal sulcus often transects the Broca's region, while the ascending ramus of the lateral fissure caudally borders the prominence (Figure 2C-D; Connolly, 1950; de Jager

et al., 2022). In chimpanzees, the central sulcus is placed more rostrally, and the inferior portion of the precentral sulcus bisects the orbital cap that is bordered rostrally by the inferior frontal sulcus (Figure 2E-F; Connolly, 1950). The coronal suture runs in between the precentral and central sulci in KNM-ER 3732, which points towards the primitive configuration described in Ponce de León et al. (2021). The sulcal pattern seen in the prefrontal cortex of KNM-ER 3732, and more particularly the deeply marked precentral sulcus that incised the Broca's cap, is not found in any of the early *Homo* specimens described in Ponce de León et al. (2021) but approximates the condition seen in the Dmanisi cranium D2282 (Fig. S1B in Ponce de León et al., 2021). However, this pattern is seen in a contemporaneous non-*Homo* hominin specimen in South Africa, i.e., *Australopithecus sediba* (Falk, 2014).

Discussion

Paleoneuroanatomy supports taxic diversity within early *Homo*. Overall, the present study supports Ponce de León et al.'s (2021) hypothesis of a primitive brain of early *Homo* but also adds information about the variation pattern of the inferior frontal gyrus. In particular, the diversity of the prefrontal sulcal pattern of hominin endocasts at Koobi Fora revives the debate about the possible presence of two early *Homo* species in this locality (i.e., *Homo habilis sensu stricto* and *Homo rudolfensis*). In 1983, Falk (1983) published the description of the endocasts of two *Homo* specimens from Koobi Fora, KNM-ER 1470 and KNM-ER 1805. Her analysis supported the co-existence of two morphs, KNM-ER 1470 representing a more derived human-like sulcal pattern. Interestingly, cerebral evidence brought up by her analysis matched other studies that emphasized the more derived craniodental anatomy of specimens attributed to *Homo rudolfensis* as opposed to the more primitive (*Australopithecus*-like) traits identified within *Homo habilis sensu stricto* (Leakey et al., 2012). While describing the external morphology of the neurocranium, Leakey (1976: 575) noted that KNM-ER 3732 was "strikingly similar to KNM-ER 1470". This resemblance is not reflected in their cerebral organization since the present study rather suggests a primitive organization of the Broca's cap in KNM-ER 3732. If KNM-ER 1470 had indeed a derived brain, taxic diversity as a source of variation cannot be discarded. If we go further down that route, the similarities between KNM-ER 3732 and *Australopithecus sediba* revealed by our study could be an argument supporting the presence of *Australopithecus* in Koobi Fora (Wood and Collard, 1999).

The evolutionary history of the human Broca's area unravelled. Beyond the taxonomic aspect, variation detected in Koobi Fora could provide information about underlying evolutionary mechanisms, and more specifically the process of fixation of an adaptive variant, i.e., a new organization of the Broca's cap and the increase of neural interconnectivity in this region (Van Essen, 1987). Such neurological changes might have had deep implication in the emergence of novel behavior, such as articulated language (Beaudet, 2017, 2021). Within this scenario, the study from Ponce de León et al. (2021) would suggest that this trait became stabilized by 1.5 million years ago. The identification of protracted brain growth as early as in *Australopithecus afarensis* (Gunz et al., 2020), inducing longer exposure to social environment during brain maturation (rev. in Hublin et al., 2014), could be consistent with a derived organization of the Broca's cap being selected as a response to social environmental stimuli through developmental plasticity, culminating in this variant becoming dominant within *Homo*.

Materials and Methods

Materials. KNM-ER 3732 is currently housed in the National Museums of Kenya in Nairobi (Kenya). Comparative extant human (N = 1, adult female) and chimpanzee (N = 1, adult female) specimens are from the Pretoria Bone Collection at the University of Pretoria (South Africa) and in the Royal Museum for Central Africa in Tervuren (Belgium), respectively (Beaudet et al., 2018).

Digitization. KNM-ER 3732 was scanned using an Artec Space Spider 3D scanner and reconstructed with the software Artec Studio 16X. The 3D mesh can be viewed on MorphoSource (https://www.morphosource.org/concern/media/000497752?locale=en). The comparative specimens were imaged by microfocus X-ray tomography (21) at the South African Nuclear Energy Corporation in Pelindaba (South Africa) and at the Centre for X-ray Tomography of Ghent University (UGCT) in Ghent (Belgium). Virtual endocasts were generated using Endex software (Subsol et al., 2010).

Detection and identification of sulcal imprints. Sulcal imprints were automatically detected through a geometry-based method using curvature lines computed from the surface. Sulcal imprints, considered as variation points of the surface on a triangle mesh, were detected through a geometry-based method using curvature lines defined as salient subsets of the extrema of the principal curvatures on surfaces (Yoshizawa et al., 2008; Beaudet et al., 2016). Vascular

imprints and non-anatomical structures (e.g., fractures) were manually removed through a
customized script written in MATLAB R2013a (Mathworks).

Acknowledgements

We thank Emmanuel Ndiema and the curatorial staff of the National Museums of Kenya for
collection access. For scientific/technical discussion we are grateful to: G. Castelli
(Cambridge), M. Mirazón-Lahr (Cambridge), R. Holloway (New York), F. Spoor (London).
AB is funded by the National Research Foundation of South Africa (Research Development
Grants for Y-Rated Researchers, grant number 129336) and the University of Cambridge. EdJ
is funded by the University of Cambridge Harding Distinguished Postgraduate Scholars
Programme. AB and EdJ are funded by the South Africa/France (PROTEA) Joint Research
Programme (grant number 129923) and the McDonald Institute for Archaeological Research.

Competing Interest Statement

The authors declare no competing interest.

References

Beaudet A. 2017. The emergence of language in the hominin lineage: perspectives from fossil
endocasts. *Frontiers in Human Neuroscience* **11**:427.

Beaudet A. 2021. The enigmatic origins of the human brain. *Science* **372**:124-125.

Beaudet A, Dumoncel J, de Beer,F, Duployer B, Durrleman S, Gilissen E, Hoffman J,
Tenailleau C, Thackeray JF, Braga J. 2016. Morphoarchitectural variation in South
African fossil cercopithecoid endocasts. *Journal of Human Evolution* **101**:65-78.

Beaudet A, Carlson KJ, Clarke RJ, de Beer F, Dhaene J, Heaton J, Pickering TR, Stratford D.
2018. Cranial vault thickness variation and inner structural organization in the StW 578
hominin cranium from Jacovec Cavern, South Africa. *Journal of Human Evolution*
121:204-220.

Connolly CJ. 1950. External Morphology of the Primate Brain. CC Thomas.

de Jager EJ, Risser L, Mescam M, Fonta C, Beaudet A. 2022. Sulci 3D mapping from human
cranial endocasts: A powerful tool to study hominin brain evolution. *Human Brain*
Mapping **43**:4433-4443.

Falk D. 1983. Cerebral cortices of East African early hominids. *Science* **221**:1072-1074.

Falk D. 2014. Interpreting sulci on hominin endocasts: old hypotheses and new findings. *Frontiers in Human Neuroscience* **8**:134.

Feibel CS, Brown FH. McDougall I. 1983. Stratigraphic context of fossil hominids from the Omo groups deposits: northern Turkana Basin, Kenya and Ethiopia. *American Journal of Physical Anthropology* **78**:595-622.

Feibel CS, Lepre CJ, Quinn RL. 2006. Stratigraphy, correlation, and age estimates for fossils from Area 123, Koobi Fora. *Journal of Human Evolution* **57**: 112-122.Gunz P, Neubauer S, Falk D, Tafforeau P, Le Cabec A, Smith TM, Kimbel WH, Spoor F, Alemseged Z. 2020. *Australopithecus afarensis* endocasts suggest ape-like brain organization and prolonged brain growth. *Science Advances* **6**:eaaz4729.

Holloway RL, Broadfield DC, Yuan MS. 2004. The Human Fossil Record: Brain Endocasts, the Paleoneurological Evidence. Wiley-Liss.

Hublin JJ, Neubauer S, Gunz P. 2015. Brain ontogeny and life history in Pleistocene hominins. *Philosophical Transactions of the Royal Society B: Biological Sciences* **2370**:20140062.

Leakey REF. 1976. New hominid fossils from the Koobi Fora formation in Northern Kenya. *Nature* **261**:574-576.

Leakey MG, Spoor F, Dean MC, Feibel CS, Antón SC, Kiarie C, Leakey LN. 2012. New fossils from Koobi Fora, northern Kenya, confirm taxonomic diversity in early *Homo*. *Nature* **488**:201-204.

Neubauer S. 2014. Endocasts: Possibilities and limitations for the interpretation of human brain evolution. *Brain, Behavior and Evolution* **84**:117-134.

Ponce de León MS, Bienvenu T, Marom A, Engel S, Tafforeau P, Alatorre Warren JL, Lordkipanidze D, Kurniawan I, Murti DB, Suriyanto RA, Koesbardiati T, Zollikofer CPE. 2021. The primitive brain of early *Homo*. *Science* **372**:165-171.

Subsol G, Gesquière G, Braga J, Thackeray F. 2010. 3D automatic methods to segment "virtual" endocasts: state of the art and future directions. *American Journal of Physical Anthropology* **141**(suppl. 50):226-227.

Tobias PV. 1987. The brain of *Homo habilis*: a new level of organization in cerebral evolution. *Journal of Human Evolution* **16**:741-761.

Van Essen DC. 1987. A tension-based theory of morphogenesis and compact wiring in the central nervous system. *Nature* **385**:313.

Walczak BE, Johnson CN, Howe BM. 2015. Myositis ossificans. *Journal of the American Academy of Orthopaedic Surgeons* **23**: 612-622

Wood B. 1991. Koobi Fora Research Project Volume 4: Hominid Cranial Remains. Clarendon
Press.

Wood B, Collard M. 1999. The human genus. *Science* **284:**65-71.

Yoshizawa S, Belyaev A, Yokota H, Seidel HP. 2008. Fast, robust, and faithful methods for
detecting crest lines on meshes. *Computer Aided Geometric Design* **25:**545-560.

Zollikofer CPE, Ponce de León MSP. 2013. Pandora's growing box: Inferring the evolution and
development of hominin brains from endocasts. *Evolutionary Anthropology* **22:**20-33.

Figures

Figure 1. 3D virtual rendering of the natural endocast of KNM-ER 3732 and identification of sulcal imprints. KNM-ER 3732 is shown in lateral right (**A**), lateral left (**B**) and dorsal (**C**) views. Images not to scale. ar: ascending ramus of the lateral fissure; c: central sulcus; CO: coronal suture; fi: inferior frontal sulcus; fm: middle frontal sulcus; ip: intra-parietal sulcus; pc: pre-central sulcus; pt: post-central sulcus.

Figure 2. Comparison of the sulcal patterns identified in the inferior frontal area of KNM-ER 3732. The natural endocast (A) and braincase (**B**) of KNM-ER 3732 are compared to the virtual endocasts of extant human (C) and chimpanzee (D) individuals. Images not to scale. ar: ascending ramus of the lateral fissure; c: central sulcus; co: coronal suture; fi: inferior frontal sulcus; fm: middle frontal sulcus; ip: intra-parietal sulcus; pc: pre-central sulcus; pt: post-central sulcus.