Enhanced fungal specificity and *in vivo* therapeutic efficacy of a C-22 modified FK520 analog

against *C. neoformans*

List of Authors

Angela Rivera[1], Won Young Lim[2†], Eunchong Park[3,†], Patrick A. Dome[2, †], Michael J. Hoy[4], Ivan Spasojevic[5], Sheng Sun[4], Anna Floyd Averette[4], Sergio Pina-Oviedo[6], Praveen R. Juvvadi[7], William J. Steinbach[7], Maria Ciofani[3,4], Jiyong Hong[1,2*], and Joseph Heitman[1,4*]

Affiliations

1. Department of Pharmacology and Cancer Biology, Duke University Medical Center, Durham, North Carolina, USA

2. Department of Chemistry, Duke University, Durham, North Carolina, USA

3. Department of Integrative Immunobiology, Duke University Medical Center, Durham, North Carolina, USA

4. Department of Molecular Genetics and Microbiology, Duke University Medical Center, Durham, North Carolina, USA

5. Department of Medicine, Duke University Medical Center, Durham, North Carolina, USA

6. Department of Pathology, Duke University Medical Center, Durham, North Carolina, USA

7. Department of Pediatrics, University of Arkansas for Medical Sciences, Little Rock, Arkansas, USA

[†] These authors contributed equally to this work

*Address correspondence to Joseph Heitman, heitm001@duke.edu and Jiyong Hong,

jiyong.hong@duke.edu.

Abstract

Fungal infections are of mounting global concern, and the current limited treatment arsenal poses challenges when treating such infections. In particular, infections by *Cryptococcus neoformans* are associated with high mortality, emphasizing the need for novel therapeutic options. Calcineurin is a protein phosphatase that mediates fungal stress responses, and calcineurin inhibition by the natural product FK506 blocks *C. neoformans* growth at 37°C. Calcineurin is also required for pathogenesis. However, because calcineurin is conserved in humans, and inhibition with FK506 results in immunosuppression, the use of FK506 as an anti-infective agent is precluded. We previously elucidated the structures of multiple fungal calcineurin-FK506-FKBP12 complexes and implicated the C-22 position on FK506 as a key point for differential modification of ligand inhibition of the mammalian versus fungal target proteins. Through *in vitro* antifungal and immunosuppressive testing of analogs of FK520 (natural analog of FK506) analogs, we identified JH-FK-08 as a lead candidate for further antifungal development. JH-FK-08 exhibited significantly reduced immunosuppressive activity and both reduced fungal burden and prolonged survival of infected animals. JH-FK-08 exhibited additive activity in combination with fluconazole *in vivo*. These findings further advance calcineurin inhibition as an antifungal therapeutic approach.

Importance

Fungal infections cause significant morbidity and mortality globally. The therapeutic armamentarium against these infections is limited and development of antifungal drugs has been hindered by the evolutionary conservation between fungi and the human host. With rising resistance to the current antifungal arsenal and an increasing at-risk population, there is an urgent need for the development of new antifungal compounds. The FK520 analogs described in this study display potent antifungal activity as a novel class of antifungals centered on modifying an existing orally-active FDA approved therapy. This research advances the development of much needed newer antifungal treatment options with novel mechanisms of action.

Introduction

Fungal infections, which result in more than 1.5 million deaths per year, are recognized as a serious global health threat (1). The World Health Organization recently released a prioritized list of fungal pathogens for global research (2). The critical priority group consists of 4 life-threatening fungal pathogens: *Cryptococcus neoformans, Candida albicans, Candida auris,* and *Aspergillus fumigatus* (2). Underscoring the global severity of *C. neoformans* infection, cryptococcal meningoencephalitis, is responsible for ~20% of all AIDS-related deaths, and is among the deadliest fungal infections of global impact (3). These infections have a further devastating impact on sub-Saharan African populations (1, 3, 4). The current treatment for cryptococcal infection consists of combination therapy with amphotericin B and flucytosine in high-income countries; however, those most impacted by this health burden have fewer and less effective treatment options (5). Flucytosine is typically not available in these regions, and often fluconazole alone is prescribed due to its oral availability. However, fluconazole antifungal activity against *C. neoformans* is only fungistatic*,* and resistance frequently develops in the treated patient as a result of aneuploidy (6)*.* Resistance to fluconazole is rapidly emerging in multiple other fungal species including *Candida auris* and *Aspergillus fumigatus* that is in part due to environmental use of azoles in agriculture, further limiting this therapeutic option and driving the need for newer antifungal drugs (7–9).

One promising antifungal drug target under investigation is the protein phosphatase calcineurin. Calcineurin is highly conserved across pathogenic organisms including fungi and parasites (10–12). Calcineurin mediates stress responses in *C. neoformans, C. albicans,* and *A. fumigatus* (13–16). Calcineurin inhibition by FK506 prevents *C. neoformans* growth at high temperature (17, 18). In fact, calcineurin is essential for virulence in *C. neoformans* and also in several other pathogenic fungal species including *Candida* and *Aspergillus*, and therefore is an attractive antifungal drug target (15, 19, 20). The high degree of conservation of calcineurin across human fungal pathogens supports the approach of targeting calcineurin to develop a pan-antifungal therapeutic.

However, calcineurin is also conserved in humans (21). Inhibiting the calcineurin phosphatase in humans results in immunosuppression. Specifically, the natural calcineurin inhibitor FK506 (also

known as tacrolimus) induces immune suppression by inhibiting T cell activation and interleukin-2-dependent responses mediated by the nuclear factor of activated T cells (NFAT) transcription factor (22–28). The FK506 ligand first binds to the FK506-binding protein (FKBP12) and the FKBP12-FK506 complex then binds to and inhibits calcineurin (29–31). FK506 is an FDA approved therapeutic initially deployed to prevent graft rejection following organ transplants (32, 33). FK506 has also been researched for use in additional applications including during treatment of hepatic fibrosis, glioblastoma, as well as aging and bone degeneration in Alzheimer's (34–37). Currently, FK506 has received additional FDA approval for use in atopic dermatology applications (38).

Humans and fungi share a high degree of conservation of cellular machinery making it difficult to identify fungal-specific drug targets. As such, immunosuppression by FK506 precludes its use as an antifungal therapeutic as immune suppression is a factor that increases the risk of fungal infection (39). Our goal is to modify the structure of the calcineurin inhibitor FK520 (a natural product analog of FK506 or ascomycin) to preserve antifungal activity and reduce or eliminate immunosuppressive activity. Both FK506 and FK520 inhibit *C. neoformans* growth under stress conditions, and synthesis of less or non-immunosuppressive analogs that retain antifungal activity is a validated antifungal drug development approach (40–43). We previously solved high resolution X-ray crystal structures of the calcineurin-FK506-FKBP12 complexes from fungal pathogens including *A. fumigatus, C. albicans,* and *C. neoformans.* Using structure-guided drug design, the carbon at position 22 (C-22) on FK506 was implicated as a critical point for chemical modification to achieve differential ligand targeting of human and fungal calcineurin (44). Thus far, it was demonstrated how modifications at this site on both FK506 and its natural analog FK520, differing by a single allyl to ethyl group substitution at the C-21 position, resulted in decreased immune suppression while maintaining sufficient antifungal activity to be of benefit in animal infection models (45, 46). Developing FK506 analogs as an antifungal therapeutic approach is further bolstered by the research describing synergistic activity of calcineurin inhibitors with additional antifungal agents currently being investigated in clinical trials (47).

Here we have developed an expanded library of FK506/FK520 analogs modified at the C-22 position. These efforts have resulted in the generation of compound JH-FK-08 as a novel antifungal

therapeutic effective at significantly reducing lung fungal burden in a murine cryptococcosis model compared to untreated infected animals. Furthermore, treatment with JH-FK-08 significantly prolonged murine survival and surpassed the antifungal efficacy achieved in previous efforts. These advances identify an FK520 analog with potential to be utilized to treat invasive fungal infections.

Results

Chemical synthesis of novel FK520 analogs modified at the C-22 position

Previous studies determined the C-22 position on FK506/FK520 as a key point of differentiation between fungal versus human calcineurin-FK506-FKBP12 complexes (44, 45). Based on this, 12 new FK520 analogs were synthesized through one-step syntheses that can be readily scaled up for studies in animal infection models for pre-clinical analysis (Fig. 1) (44, 48). The synthesis of FK520 analogs with C-22 modifications features a condensation of FK520 with a hydrazide or a substituted hydrazine either under reflux or with acidic conditions. After completion of the reaction, the reaction mixture was concentrated *in vacuo* and purified by silica gel column chromatography to give the desired C-22-modified FK520 analogs (33-52% yield). Each analog was validated through ^{1}H nuclear magnetic resonance (NMR) and high-resolution mass spectrometry (HRMS) (Text S1, Table S1).

FK520 analogs display conserved antifungal activity

Eleven of these FK520 analogs were found to retain some level of antifungal activity. To assess antifungal activity, the panel of compounds was tested for fungal growth inhibition using minimum inhibitory concentration (MIC) assays against *C. neoformans* and *C. albicans* and minimum effective concentration (MEC) assays against *A. fumigatus*. The results from this broth microdilution assay showed a decrease in antifungal activity of the modified compounds compared to FK506 and FK520 (Table 1). Ten of these analogs exhibited an MIC value between 0.4 to 1.6 µg/mL which is an 8- to 32-fold decrease in activity compared to FK506 or FK520 in *C. neoformans* and *C. albicans.* MEC values against *A. fumigatus* ranged from 0.6 to 2.5 µg/mL for 10 of the analogs tested. Notably, analogs JH-FK-13, JH-FK-15, and JH-FK-17 exhibited solubility issues during microdilution broth assays (Fig. S1) (49–51). Overall, eleven analogs maintain relatively potent antifungal activity across species and represent an active library for further investigation.

Modifications at C-22 position lead to enhanced fungal specificity

Mammalian calcineurin inhibition was then assessed and found to be reduced for these FK520 analogs. The inhibition of mammalian calcineurin was assessed by measuring the reduction of interleukin-2 (IL-2) expression from primary mouse $CD4^+$ T cells cultured in the presence of FK506, FK520, or an analog. Calcineurin activity is required for T cell activation and cytokine production, and IL-2 production serves as an effective measure of mammalian calcineurin activity. After culturing mouse $CD4^+$ T cells for 72 hours, IL-2 expression was analyzed by flow cytometry to evaluate immunosuppressive activity of the analogs. The results showed an overall reduction in *in vitro* immunosuppressive activity of the analogs compared to FK506 and FK520 as denoted by the increased concentration required to inhibit cytokine production (Fig. 2A). IC_{50} values were determined from these data to quantify the reduction in immunosuppression due to the structural modification of FK520 (Table 2). The IC_{50} values of the analogs demonstrate a minimum 65-fold reduction in mammalian calcineurin inhibition. Notably, IC_{50} values increased from 0.09 nM for FK506 up to 70.5 nM for analog JH-FK-23 and to 42.6 nM for JH-FK-08. Analog JH-FK-13 showed an IC_{50} greater than 200 nM and an MIC value greater than 50 µg/mL, indicating targeting of both mammalian and fungal calcineurin has been compromised. Our results demonstrate that structurally modified analogs have reduced capacity to inhibit mammalian calcineurin, and therefore their immunosuppressive activity has been reduced.

JH-FK-08 has been identified as a lead compound

Based on the *in vitro* fungal and mammalian calcineurin inhibition data, the panel of FK520 analogs was assessed and ranked. Therapeutic index scores were used to quantify the effects of structural modification on the compounds (Table 2). Therapeutic index scores (TI) were calculated for each analog as the ratio of relative change in immunosuppressive and antifungal activities compared to FK506. This ratio of therapeutic benefit and host toxicity, or fungal and mammalian calcineurin inhibition, allowed us to quantify and rank these analogs. Using the formula TI = ($IC_{50[compound]}$/$IC_{50[FK506]}$) / ($MIC_{[compound]}$/$MIC_{[FK506]}$) and the *in vitro* immunosuppressive and antifungal activities against *C.*

neoformans, the TI for JH-FK-08 is 29.6. This is the greatest TI found within the library and allowed us to identify JH-FK-08 as a lead candidate for *in vivo* study. Tolerability of JH-FK-08 was assessed up to a maximum of 40 mg/kg given twice per day (unpublished data). It should be noted that analogs JH-FK-12 and JH-FK-23 are also promising candidates with respective TIs of 24.7 and 24.5.

For the lead compound JH-FK-08, the mechanism of action resulting in antifungal activity was confirmed to be via calcineurin inhibition. To confirm the target of JH-FK-08, MICs of JH-FK-08 were determined with fungal mutants that lack the drug binding protein FKBP12 or have mutations in the binding site on calcineurin that prevent FKBP12-FK506 binding and inhibition. The results show that JH-FK-08 has no antifungal activity against these mutant fungal strains (Table S2). This supports the conclusion that JH-FK-08 binds FKBP12 and the resulting FKBP12-JH-FK-08 complex binds to and inhibits calcineurin.

The length of time that drugs are active *in vivo* is critical information relevant to drug development, and the time frame of drug activity can have a drastic impact on treatment regimen and efficacy. Therefore, the pharmacokinetic profiles of FK506 and JH-FK-08 were investigated. Three male and three female CD1 mice were administered a single dose of JH-FK-08 or FK506 at 40 mg/kg. Whole blood samples were taken over a 24-hour period and the drug concentration remaining in the blood was determined. Pharmacokinetic parameters were calculated by a non-compartmental approach. No significant differences across groups were observed in whole blood samples of drug concentration over time, indicating a strong conservation of the pharmacokinetic profile of JH-FK-08 compared to that of FK506 (Fig. 2B). When assessing specific pharmacokinetic measures sex differences began to emerge. When looking at the area under the curve (AUC), females showed roughly twice the value of males, and male clearance of both FK506 and JH-FK-08 was higher compared to female treated animals (Table S3). Despite this, the half-life across all groups ranged from 1.14-1.48 hours. Overall, JH-FK-08 exhibits a pharmacokinetic profile very similar to that of FK506 including sex-dependent trends seen in clearance and AUC.

In vivo immunosuppressive activity of JH-FK-08 is reduced

Whole animal studies were conducted to assess the *in vivo* implications of the reduced mammalian calcineurin inhibition demonstrated during *in vitro* immune cell testing. The *in vivo* immunosuppressive effect of JH-FK-08 was assessed by studying T follicular helper (Tfh) cell production in mice. Tfh cells play a role in protection against and response to fungal infections (52, 53). For this study, mice received 8 days of twice daily dosing of vehicle, FK506 (2.5 mg/kg), or JH-FK-08 (20 or 40 mg/kg). Twenty-four hours after the first dose, mice were immunized with NP-OVA to induce an immune response. Cells were analyzed 7 days post immunization, and cell types were analyzed by flow cytometry (Fig. 3B).

Treatment with JH-FK-08 at 20 mg/kg had no significant effect on the percent of Tfh cells produced compared to vehicle (Fig. 3A). However, the production of Tfh cells was reduced at a dosage of 40 mg/kg of JH-FK-08. The immune response was also significantly reduced in mice treated with 2.5 mg/kg FK506. In germinal center B cells, JH-FK-08 showed similar levels of immunosuppressive activity compared to FK506 (Fig. S2). Overall, these findings indicate that Tfh cells are impacted by JH-FK-08 in a dose-dependent manner, and FK506 significantly inhibits Tfh cell production at 1/16[th] of the dose of JH-FK-08. These results in whole animals are well correlated with the finding that the analog JH-FK-08 has significantly reduced mammalian calcineurin inhibition *in vitro* compared to FK506.

JH-FK-08 reduces fungal burden during murine cryptococcal infection

In vivo antifungal activity and therapeutic benefit was assayed by investigating the impact of the FK520 analog on organ fungal burden during cryptococcosis. Specifically, *in vivo* efficacy of JH-FK-08, on fungal burden was assessed using A/J female mice. Treatment groups included vehicle of 10% Kolliphor : ethanol (1:1) in phosphate buffered saline (PBS), 6 mg/kg fluconazole, 40 mg/kg JH-FK-08, or the fluconazole and JH-FK-08 combination, all given via intraperitoneal (i.p.) injection twice per day. Single day dosing with vehicle, 12 mg/kg fluconazole, 40 mg/kg JH-FK-08, and JH-FK-08 plus fluconazole combination resulted in limited efficacy reducing lung fungal burden and dosing twice per day was implemented to augment treatment efficacy (Fig. S3). Mice were intranasally infected with *C.*

neoformans strain H99, and fungal burden in the lung, spleen, and brain was assessed after 14 days of treatment. Treatment with fluconazole, JH-FK-08, or a combination resulted in no detectable fungal burden in the brain or spleen. The average lung fungal burden in untreated animals was approximately 3.7 million fungal cells per gram of lung tissue. This lung fungal burden was significantly reduced by ~6-fold, ~200-fold, and by >400-fold in treatments with fluconazole, JH-FK-08, or the combination, respectively (Fig. 4). Additionally, recruitment of immune cells to the site of infection in lung tissue occurred based on histopathology analysis. The expected granulomatous inflammatory response admixed with mature lymphocytes and neutrophils was seen during JH-FK-08 and combination treatment (Fig. S4). This indicates that reduction of JH-FK-08 immunosuppression was sufficient such that immune responses were still mounted during infection, and JH-FK-08 treatment had a direct impact reducing organ fungal burden during cryptococcal infection.

Treatment with JH-FK-08 extends survival during cryptococcal infection

Finally, whether the antifungal efficacy translates into extended survival during infection was assessed. To test treatment impact on survival, murine survival was monitored in the cryptococcosis infection model. Following the same protocol of infection and dosing as in the fungal burden experiment, animal survival was monitored following infection and two weeks of intraperitoneal dosing. Fluconazole treatment extended median survival from 16 to 27 days. JH-FK-08 treatment alone more than doubled median survival from 16 days in vehicle treated mice to 33 days in treated animals. This beneficial impact on survival was further enhanced by concomitant fluconazole treatment, and this resulted in an increased median survival from 16 days to 36 days (Fig. 5). These results further emphasize the therapeutic benefit JH-FK-08 offers by demonstrating an extended survival of animals treated with JH-FK-08 alone or in combination with fluconazole.

Discussion

In this study, an expanded library of structurally validated FK520 analogs was generated by one-step chemical synthesis and screened for their antifungal and immunosuppressive activity to develop non-immunosuppressive fungal-specific calcineurin inhibitors. The research and development of novel antifungal therapeutics is relevant due to rapidly emerging resistance to the current antifungal arsenal (6, 7). Additionally, cryptococcal infections are associated with high mortality, and the treatment options for the population most devastated by this human fungal pathogen are typically limited to orally available but fungistatic fluconazole (1, 3). FK506 is an orally active drug, making the development of fungal-specific analogs a potential avenue to better serve these patients.

By first assessing *in vitro* antifungal and immunosuppressive activity of the various C-22 modified FK520 analogs, we were able to confirm fungal specificity. During antifungal *in vitro* screening, fungal calcineurin inhibition was measured through microdilution broth growth assays. These results demonstrated many compounds were able to maintain potent antifungal activity indicating these structural modifications did not prevent fungal calcineurin targeting. Reduced mammalian calcineurin inhibition by the FK520 analogs was confirmed in cytokine production studies. During this *in vitro* immunosuppressive assay, production of the cytokine IL-2, which is known to be inhibited by FK520, was shown to be suppressed to a lesser degree by the FK520 analogs. This finding confirms reduced targeting of the mammalian calcineurin by the FK520 analogs. Using this information, therapeutic index scores were calculated to quantify the degree to which fungal selectivity was achieved.

JH-FK-08 was identified as the lead compound following *in vitro* screening. The introduced structural modifications did not significantly alter the pharmacokinetic (PK) properties (e.g., C_{max}, half-life, whole-blood drug concentration over time) compared to FK506 in both male and female mice. A pattern of higher plasma exposure (higher AUC, decreased clearance) in females was also observed in JH-FK-08- and FK506-treated animals relative to male treated animals (Table S3). To further study the antifungal effect of calcineurin inhibition during infection, JH-FK-08 was then investigated for *in vivo* antifungal efficacy. Animals were intranasally infected with *C. neoformans* to replicate the conventional infection route in humans. In this study, JH-FK-08 was found to significantly reduce lung fungal burden

by approximately 200-fold during two weeks of treatment and more than doubled murine survival time during infection compared to untreated infected animals. These benefits were further enhanced when animals were treated with JH-FK-08 in combination with fluconazole.

Overall, the preclinical efforts presented here advance JH-FK-08 as a promising candidate for potential clinical study. The activity of JH-FK-08 far exceeds the previous two libraries' top candidates APX879 and JH-FK-05 based on therapeutic index score comparison and *in vivo* efficacy (44, 45). These results support further study of calcineurin inhibition as a potential effective antifungal approach. Exciting research avenues for future study include: 1) investigating oral activity of JH-FK-08, 2) researching how JH-FK-08 can interact with and cross the blood brain barrier, and 3) investigating possible combination therapies. Current treatments for fungal infections make use of a combinations of therapies, and using multiple drugs during one therapeutic application can increase treatment efficacy as well as reduce emerging resistance and mitigate potential side effects (5, 49). One potential combination partner is APX001A and its analogs as this compound is under clinical investigation as a novel antifungal class (47, 50, 51). Furthermore, this approach of designing fungal specific analogs of existing therapies can be utilized with other natural FKBP12 binding products such as rapamycin, which also demonstrates potent antifungal activity against high priority fungal pathogens (52, 53).

Materials and methods

Synthesis of FK520 analogs

General procedure A: To a solution of FK520 (APIChem Technology) in EtOH (0.15 M) was added a hydrazide (6 eq) in a sealed vial. The reaction mixture was stirred at 90 °C for 48 h. After cooling to 25 °C, the reaction mixture was concentrated *in vacuo*. The crude mixture was reconstituted in EtOAc and washed with H_2O and brine. The combined organic layers were dried over anhydrous Na_2SO_4 and concentrated *in vacuo*. The residue was purified by column chromatography (silica gel, CH_2Cl_2/MeOH) to afford a substituted hydrazone analog.

General procedure B: To a solution of FK520 in MeOH (0.10 M) was added a hydrazide (6 eq) or a substituted hydrazine (6 eq) and TFA (1 µL per 100 mg of FK520). The reaction was stirred at 25 °C for 48 h. The reaction mixture was concentrated *in vacuo* and reconstituted in EtOAC. The resulting reaction mixture was washed with 10% aqueous $NaHCO_3$ solution, H_2O, and brine. The combined organic layers were dried over anhydrous Na_2SO_4 and concentrated *in vacuo*. The residue was purified by column chromatography (silica gel, CH_2Cl_2/MeOH) to afford a substituted hydrazone analog.

Antifungal susceptibility testing

In vitro antifungal activity was assessed for all fungal strains in RPMI 1640 (Sigma-Aldrich R1383). MICs were measured using Clinical and Laboratory Standards Institute (CLSI) M38-A2 and M27-A3 standard *in vitro* antifungal susceptibility protocols. *C. neoformans* and *A. fumigatus* testing was performed at 37°C. MECs for *A. fumigatus* were recorded after 48 h growth. *C. albicans* testing was performed at 30°C with a uniform concentration of 2 µg/mL of fluconazole to sensitize the cells to calcineurin inhibition.

Fungal strains

C. neoformans clinical isolate H99 and congenic derived strain KN99α strain were used (54). KN99α derived *frr1*Δ deletion mutant with nourseothricin resistance marker as part of the Madhani deletion collection (NIH funding R01AI100272) was used. *C. deneoformans* strains used include JEC21

and derived C21F2 and C21F3 strains (20, 55, 56). *C. albicans* strain SC5314 is the wild type strain from which mutants YAG171 (*rbp1/rbp1*) and YAG237 (*CNB1-1/CNB1*) were derived (57). *A. fumigatus* strain *akuB*[Ku80] and a derived FKBP12 mutant *fkbp12Δ* were used during antifungal screening (58).

In vitro immunosuppressive activity testing

Spleen and lymph nodes from C57BL/6 mice were collected and homogenized through a 40 µm filter. Pan-CD4$^+$ T cells were enriched using MagniSort Mouse CD4 T cell Enrichment Kit (Invitrogen). Thereafter, CD4$^+$ CD25$^-$ CD62L^{hi} CD44^{lo} naïve CD4$^+$ T cells were FACS-sorted using a MoFlo Astrios cell sorter (Beckman Coulter). Naïve CD4$^+$ T cells were cultured in Iscove's Modified Dulbecco's Medium (IMDM) supplemented with 10% FBS, penicillin (10 U/ml), streptomycin (10 µg/ml), glutamine (2 mM), gentamicin (50 µg/ml), and β-mercaptoethanol (55 µM). Cells were cultured for 72 h on anti-hamster IgG-coated plates with hamster anti-CD3ε antibodies, hamster anti-CD28 antibodies, neutralizing anti-IL-4 antibody, recombinant IL-12 (10 ng/mL), and recombinant IL-2 (50 U/mL). During this culture, cells were incubated with serially diluted drug suspended in DMSO.

Flow cytometry

Cell surface staining of single cell suspension was performed in staining buffer (PBS supplemented with 2 mM EDTA and 0.5% BSA). Anti-FcγR (93; Biolegend) was treated to block Fc receptor before surface staining of primary cells isolated from lymph nodes of NP-OVA-immunized mice. Cells were stained with fixable viability dye (eBioscience) before surface staining to exclude dead cells from analysis. For the detection of intracellular cytokines, cells were additionally incubated with phorbol 12-myristate 13-acetate (PMA), ionomycin in the presence of GolgiStop (BD Biosciences) during the final 4 h of incubation. For intracellular staining, cells were fixed and permeabilized with Fixation/Permeabilization buffer (eBioscience). Flow cytometric analysis was performed utilizing BD FACSCantoII or BD LSRFortessaX20 (BD Biosciences) and FlowJo software (BD Biosciences).

Compound preparation for *in vivo* studies

For *in vivo* studies, fluconazole was purchased from Duke Pharmacy Storeroom. Vehicle was a sterile solution of 10% Kolliphor EL (Sigma-Aldrich C5315), 10% ethanol, and 80% phosphate-buffered saline (PBS, Sigma-Aldrich D8537). For administration to animals, all other drugs were obtained in powder form and diluted in a sterile mixture of 50% Kolliphor EL 50% ethanol before dilution to 80% PBS.

In vivo immunosuppressive assessment

Groups of 5 female C57BL/6 mice were treated twice daily for 8 days with vehicle, FK506 (2.5 mg/kg), JH-FK-08 (20mg/kg), or JH-FK-08 (40 mg/kg) via intraperitoneal (i.p.) injection. 24 h after the first dose, animals were immunized with NP-OVA (100 µg) in alum via subcutaneous injection. 7 days post-immunization, mice were sacrificed, and draining lymph nodes were harvested for the analysis of Tfh cells and GC B cells by flow cytometry.

C. neoformans murine infection

C. neoformans inoculum was prepared by growing a 5 ml overnight YPD culture spinning in a tissue culture roller drum at 30°C. Cells were pelleted by centrifugation and washed in sterile PBS 3 times, concentration was determined by hemocytometer, and cells were diluted to 2×10^6 cells/mL. Groups of 10 mice (10 to 20 g) were infected with 10^5 cells of *C. neoformans* H99. For inhalation infection, animals were anesthetized with isoflurane before 50 µl of inoculum was pipetted onto the upright nasal passages of each animal. Following inhalation of the inoculum, animals were monitored for post anesthesia recovery.

Histology preparation

Lungs were inflated with and immersed in 10% neutral buffered formalin, processed routinely to paraffin embedment and sectioned at 5 mm. Sections were stained either with Hematoxylin & Eosin (H&E) or Periodic-acid Schiff (PAS). All tissue sections were evaluated and photographed with an

Olympus BX51 microscope with a DP70 camera using Olympus DP Manager software v.3.3.112. Samples processed by the Department of Pathology.

Pharmacokinetics (PK) experiment

Male (n=3; avg. 28 g) and female (n=3, avg. 25 g) CD-1 mice were injected intraperitoneally with 40 mg/kg of either JH-FK-08 or FK506. The drug formulation was 10% Kolliphor/10% ethanol/80% PBS given as 10 µl/g body weight (i.e., 200 µl per 20 g mouse). Blood (~30 µl) was collected serially ("tail snip") at 10, 20, 45 min, 1.5, 3, 6, 12, and 24 h into vials containing 1 ml of 75 mg/ml K_2EDTA in water, and immediately frozen until the day of LC/MS/MS analysis. Blood concentration/time data was utilized to calculate relevant PK parameters by non-compartmental approach and extravascular administration model within WinNonlin (2.1) software.

LC/MS/MS analysis

In 0.5 ml polypropylene vial, 10 µl blood and 20 µl of acetonitrile fortified with 100 ng/ml FK520 (internal standard), were vigorously agitated in FastPrep FP120 apparatus (Thermo-Savant) at speed 4 for 45 s. After centrifugation for 5 min at 16,000 rcf, 20 µl of supernatant was transferred to polypropylene autosampler vial. Liquid chromatography - tandem-mass spectroscopy (LC/MS/MS; Agilent 1200 series HPLC and Sciex/Applied Biosystems API 5500 QTrap) was utilized to quantify JH-FK-08 and FK506 in mouse blood. Analytical column: Agilent Eclipse Plus (C18, 1.8 µm, 50 × 4.6 mm), at 40°C. Mobile phase: (A) 0.1% formic acid, 10 mM ammonium acetate, 2% acetonitrile in water, (B) methanol. Elution gradient: 0-0.1 min 30-98% B, 0.1-1.5 min 98% B, 1.5-1.6 min 98-30% B, 1.6 -2.1 min 30% B. Run time: 5 min. Injection volume: 10 µl. Mass spectrometer parameters (voltages, gas flow, and temperature) were optimized by infusion of 100 ng/mL of analytes in mobile phase at 10 µl/min using Analyst 1.6.2 software tuning module. The MS/MS (m/z) transitions used for quantification: 864.4/846.5 (JH-FK-08), 821.4/768.5 (FK506), and 809.4/756.5 (FK520, internal standard). A set of calibrator samples in drug-free blood was prepared by adding appropriate amounts of pure analyte (JH-FK-08 or FK506) in 2.43 - 1000 µl/ml range. The calibration samples were analyzed alongside the

experimental samples. Accuracy acceptance criteria was 85% for each but the lowest level (2.43 ng/mL, 80%, LLOQ). Analyst 1.6.2 software was used for data acquisition, integration of the chromatograms, calibration curve calculation, and quantification of the study samples.

Cryptococcus fungal burden and survival analysis

For survival curve analysis, animals were monitored until reaching defined humane end points, including social isolation, hunched posture, weight loss, lack of grooming, and labored breathing, and were then euthanized via CO_2 inhalation. For fungal burden analysis, mice were euthanized following 2-weeks of treatment with designated antifungal, vehicle, or combination. Brain, lung, and spleen were harvested from animals, weighed, and homogenized in sterile PBS using steel beads and a bead beater. Each organ homogenate was serial diluted 10-fold to 10^{-4} in PBS. 100 µl of each dilution was plated to YPD plates supplemented with YPD 50 µg/ml Ampicillin, 30 µg/ml chloramphenicol and incubated at 30°C for 2 days and assessed for colony forming units. CFU per organ weight was calculated and plotted using GraphPad Prism 9.

Acknowledgements

We thank Catherine Denning-Jannace and Jeffrey Everitt for their expertise and assistance.

I.S. (PK/PD Core Laboratory) acknowledges support of the NCI/NiH Comprehensive Cancer Center Core Grant P30CA014236. M.C. acknowledges support for R01 GM115474. This material is based upon work supported by the National Science Foundation Graduate Research Fellowship under Grant No. DGF 2139754. This research is supported by NIH/NIAID R01 grant AI172451-01 and R56 grant AI112595-05 and the Gilhuly Accelerator Fund.

Joseph Heitman is co-director and fellow of the CIFAR Program Fungal Kingdom: Threats & Opportunities.

Ethics statement

Experiments utilizing mice were conducted in compliance with guidelines issued by the US Animal Welfare Act and by the Duke Institutional Animal Care and Use Committee (IACUC). Duke's Animal Care and Use Program has been accreditation by the Association for Assessment and Accreditation of Laboratory Animal Care (AAALAC), and its animal care and housing facilities are maintained by Duke's Division of Laboratory Animal Resources (DLAR). All procedures involving animals were approved by Duke IACUC under protocol no. A098-22-05.

References

1. Bongomin F, Gago S, Oladele RO, Denning DW. 2017. Global and multi-national prevalence of fungal diseases-estimate precision. J Fungi 3:57.

2. World Health Organization. 2022. WHO fungal priority pathogens list to guide research, development and public health action. Geneva: World Health Organization. License: CC BY-NC-SA 3.0 IGO.

3. Rajasingham R, Govender NP, Jordan A, Loyse A, Shroufi A, Denning DW, Meya DB, Chiller TM, Boulware DR. 2022. The global burden of HIV-associated cryptococcal infection in adults in 2020: a modelling analysis. Lancet Infect Dis 22:1748–1755.

4. Park BJ, Wannemuehler KA, Marston BJ, Govender N, Pappas PG, Chiller TM. 2009. Estimation of the current global burden of cryptococcal meningitis among persons living with HIV/AIDS. AIDS 23:525–530.

5. Perfect JR, Dismukes WE, Dromer F, Goldman DL, Graybill JR, Hamill RJ, Harrison TS, Larsen RA, Lortholary O, Nguyen M-H, Pappas PG, Powderly WG, Singh N, Sobel JD, Sorrell TC. 2010. Clinical practice guidelines for the management of cryptococcal disease: 2010 update by the Infectious Diseases Society of America. Clin Infect Dis Off Publ Infect Dis Soc Am 50:291–322.

6. Sionov E, Lee H, Chang YC, Kwon-Chung KJ. 2010. *Cryptococcus neoformans* overcomes stress of azole drugs by formation of disomy in specific multiple chromosomes. PLoS Pathog 6:e1000848.

7. Rocha MFG, Alencar LP, Paiva MAN, Melo LM, Bandeira SP, Ponte YB, Sales JA, Guedes GMM, Castelo-Branco DSCM, Bandeira TJPG, Cordeiro RA, Pereira-Neto WA, Brandine GS, Moreira JLB, Sidrim JJC, Brilhante RSN. 2016. Cross-resistance to fluconazole induced by exposure to the agricultural azole tetraconazole: an environmental resistance school? Mycoses 59:281–290.

8. Berkow EL, Lockhart SR. 2017. Fluconazole resistance in *Candida* species: a current perspective. Infect Drug Resist 10:237.

9. Arendrup MC. 2014. Update on antifungal resistance in *Aspergillus* and *Candida*. Clin Microbiol Infect 20:42–48.

10. Naderer T, Dandash O, McConville MJ. 2011. Calcineurin is required for *Leishmania* major stress response pathways and for virulence in the mammalian host. Mol Microbiol 80:471–480.

11. Matsumoto Y, Yoshikawa A, Nagamachi T, Sugiyama Y, Yamada T, Sugita T. 2022. A critical role of calcineurin in stress responses, hyphal formation, and virulence of the pathogenic fungus *Trichosporon asahii*. Sci Rep 12:16126.

12. Kurakado S, Matsumoto Y, Yamada T, Shimizu K, Wakasa S, Sugita T. 2023. Tacrolimus inhibits stress responses and hyphal formation via the calcineurin signaling pathway in *Trichosporon asahii*. Microbiol Immunol 67:49-57.

13. Chow EWL, Clancey SA, Billmyre RB, Averette AF, Granek JA, Mieczkowski P, Cardenas ME, Heitman J. 2017. Elucidation of the calcineurin-Crz1 stress response transcriptional network in the human fungal pathogen *Cryptococcus neoformans*. PLoS Genet 13:e1006667.

14. Lev S, Desmarini D, Chayakulkeeree M, Sorrell TC, Djordjevic JT. 2012. The Crz1/Sp1 transcription factor of *Cryptococcus neoformans* is activated by calcineurin and regulates cell wall integrity. PLoS One 7:e51403.

15. Cruz MC, Goldstein AL, Blankenship JR, Del Poeta M, Davis D, Cardenas ME, Perfect JR, McCusker JH, Heitman J. 2002. Calcineurin is essential for survival during membrane stress in *Candida albicans*. EMBO J 21:546–559.

16. Shwab EK, Juvvadi PR, Waitt G, Soderblom EJ, Barrington BC, Asfaw YG, Moseley MA, Steinbach WJ. 2019. Calcineurin-dependent dephosphorylation of the transcription factor CrzA at specific sites controls conidiation, stress tolerance, and virulence of *Aspergillus fumigatus*. Mol Microbiol 112:62–80.

17. Arndt C, Cruz MC, Cardenas ME, Heitman J. 1999. Secretion of FK506/FK520 and rapamycin by *Streptomyces* inhibits the growth of competing *Saccharomyces cerevisiae* and *Cryptococcus neoformans*. Microbiol 145:1989–2000.

18. Odom A, Del Poeta M, Perfect J, Heitman J. 1997. The immunosuppressant FK506 and its nonimmunosuppressive analog L-685,818 are toxic to *Cryptococcus neoformans* by inhibition of a common target protein. Antimicrob Agents Chemother 41:156-161.

19. Breuder T, Hemenway CS, Movva NR, Cardenas ME, Heitman J. 1994. Calcineurin is essential in cyclosporin A- and FK506-sensitive yeast strains. Proc Natl Acad Sci USA 91:5372–5376.

20. Odom A, Muir S, Lim E, Toffaletti DL, Perfect J, Heitman J. 1997. Calcineurin is required for virulence of *Cryptococcus neoformans*. EMBO J 16:2576–2589.

21. Klee CB, Crouch TH, Krinks MH. 1979. Calcineurin: a calcium- and calmodulin-binding protein of the nervous system. Proc Natl Acad Sci USA 76:6270–6273.

22. Kino T, Hatanaka H, Hashimoto M, Nishiyama M, Goto T, Okuhara M, Kohsaka M, Aoki H, Imanaka H. 1987. FK-506, a novel immunosuppressant isolated from a *Streptomyces*. I. Fermentation, isolation, and physico-chemical and biological characteristics. J Antibiot 40:1249–1255.

23. Tanaka H, Kuroda A, Marusawa H, Hatanaka H, Kino T, Goto T, Hashimoto M, Taga T. 1987. Structure of FK506, a novel immunosuppressant isolated from *Streptomyces*. J Am Chem Soc 109:5031–5033.

24. Dumont FJ, Staruch MJ, Koprak SL, Melino MR, Sigal NH. 1990. Distinct mechanisms of suppression of murine T cell activation by the related macrolides FK-506 and rapamycin. J Immunol 144:251–258.

25. O'Keefe SJ, Tamura J, Kincaid RL, Tocci MJ, O'Neill EA. 1992. FK-506- and CsA-sensitive activation of the interleukin-2 promoter by calcineurin. Nature 357:692–694.

26. Clipstone NA, Crabtree GR. 1992. Identification of calcineurin as a key signalling enzyme in T-lymphocyte activation. Nature 357:695–697.

27. Bierer BE, Mattila PS, Standaert RF, Herzenberg LA, Burakoff SJ, Crabtree G, Schreiber SL. 1990. Two distinct signal transmission pathways in T lymphocytes are inhibited by complexes formed between an immunophilin and either FK506 or rapamycin. Proc Natl Acad Sci USA 87:9231–9235.

28. Gwack Y, Feske S, Srikanth S, Hogan PG, Rao A. 2007. Signalling to transcription: store-operated $Ca2^+$ entry and NFAT activation in lymphocytes. Cell Calcium 42:145–156.

29. Liu J, Albers MW, Wandless TJ, Luan S, Alberg DG, Belshaw PJ, Cohen P, MacKintosh C, Klee CB, Schreiber SL. 1992. Inhibition of T cell signaling by immunophilin-ligand complexes correlates with loss of calcineurin phosphatase activity. Biochemistry 31:3896–3901.

30. Foor F, Parent SA, Morin N, Dahl AM, Ramadan N, Chrebet G, Bostian KA, Nielsen JB. 1992. Calcineurin mediates inhibition by FK506 and cyclosporin of recovery from alpha-factor arrest in yeast. Nature 360:682–684.

31. Liu J, Farmer JD, Lane WS, Friedman J, Weissman I, Schreiber SL. 1991. Calcineurin is a common target of cyclophilin-cyclosporin A and FKBP-FK506 complexes. Cell 66:807–815.

32. Starzl TE, Todo S, Fung J, Demetris AJ, Venkataramman R, Jain A. 1989. FK 506 for liver, kidney, and pancreas transplantation. Lancet 2:1000–1004.

33. Ochiai T, Nakajima K, Nagata M, Hori S, Asano T, Isono K. 1987. Studies of the induction and maintenance of long-term graft acceptance by treatment with FK506 in heterotopic cardiac allotransplantation in rats. Transplantation 44:734–738.

34. Z. Manojlovic, J. Blackmon, B. Stefanovic. 2013. Tacrolimus (FK506) prevents early stages of ethanol induced hepatic fibrosis by targeting LARP6 dependent mechanism of collagen synthesis. PLoS One 8:e65897.

35. Ge J, Wang Y, Yan Q, Wu C, Yu H, Yang H, Zou J. 2020. FK506 induces the TGF-β1/Smad 3 pathway independently of calcineurin inhibition to prevent intervertebral disk degeneration. Front Cell Dev Biol 8:608308.

36. Torres Á, Arriagada V, Erices JI, Toro MD los Á, Rocha JD, Niechi I, Carrasco C, Oyarzún C, Quezada C. 2018. FK506 attenuates the MRP1-mediated chemoresistant phenotype in glioblastoma stem-like cells. Int J Mol Sci 19:2697.

37. Gant JC, Blalock EM, Chen KC, Kadish I, Porter NM, Norris CM, Thibault O, Landfield PW. 2014. FK506-binding protein 1b/12.6: a key to aging-related hippocampal Ca2 [+] dysregulation? Eur. J. Pharmacol. 739:74-82.

38. Alaiti S, Kang S, Fiedler VC, Ellis CN, Spurlin DV, Fader D, Ulyanov G, Gadgil SD, Tanase A, Lawrence I, Scotellaro P, Raye K, Bekersky I. 1998. Tacrolimus (FK506) ointment for atopic dermatitis: a phase I study in adults and children. J Am Acad Dermatol 38:69–76.

39. Stein DK, Sugar AM. 1989. Fungal infections in the immunocompromised host. Diagn Microbiol Infect Dis 12:221–228.

40. Odom A, Del Poeta M, Perfect J, Heitman J. 1997. The immunosuppressant FK506 and its nonimmunosuppressive analog L-685,818 are toxic to *Cryptococcus neoformans* by inhibition of a common target protein. Antimicrob Agents Chemother 41:156–161.

41. Beom JY, Jung JA, Lee K-T, Hwangbo A, Song MC, Lee Y, Lee SJ, Oh JH, Ha S-J, Nam S-J, Cheong E, Bahn Y-S, Yoon YJ. 2019. Biosynthesis of nonimmunosuppressive FK506 analogues with antifungal activity. J Nat Prod 82:2078–2086.

42. Lee Y, Lee K-T, Lee SJ, Beom JY, Hwangbo A, Jung JA, Song MC, Yoo YJ, Kang SH, Averette AF, Heitman J, Yoon YJ, Cheong E, Bahn Y-S. 2018. *In vitro* and *in vivo* assessment of FK506 analogs as novel antifungal drug candidates. Antimicrob Agents Chemother 62:e01627-18.

43. Rotonda J, Burbaum JJ, Chan HK, Marcy AI, Becker JW. 1993. Improved calcineurin inhibition by yeast FKBP12-drug complexes. Crystallographic and functional analysis. J Biol Chem 268:7607–7609.

44. Juvvadi PR, Fox D, Bobay BG, Hoy MJ, Gobeil SMC, Venters RA, Chang Z, Lin JJ, Averette AF, Cole DC, Barrington BC, Wheaton JD, Ciofani M, Trzoss M, Li X, Lee SC, Chen Y-L, Mutz M, Spicer LD, Schumacher MA, Heitman J, Steinbach WJ. 2019. Harnessing calcineurin-FK506-FKBP12 crystal structures from invasive fungal pathogens to develop antifungal agents. Nat Commun 10:4275.

45. Aria T, Kouama Y, Suenaga T, Honda H. 1962. Ascomycin, an antifungal antibiotic. J Antibiot 15:231-232.

46. Hoy MJ, Park E, Lee H, Lim WY, Cole DC, DeBouver ND, Bobay BG, Pierce PG, Fox D 3rd, Ciofani M, Juvvadi PR, Steinbach W, Hong J, Heitman J. 2022. Structure-guided synthesis of

FK506 and FK520 analogs with increased selectivity exhibit *in vivo* therapeutic efficacy against *Cryptococcus*. mBio 13:e01049-22.

47. S. D. Liston, L. Whitesell, M. Kapoor, K. J. Shaw, L. E. Cowen. 2022. Calcineurin inhibitors synergize with manogepix to kill diverse human fungal pathogens. J Fungi 8:1102.

48. Rothbard JB, Wender PA, McGrane PL, Sista LVS, Kirschberg TA. 2012. Compositions and methods for enhancing drug delivery across and into ocular tissues. US8278264B2.

49. Ascierto PA, Marincola FM. 2011. Combination therapy: the next opportunity and challenge of medicine. J Transl Med 9:115.

50. Shaw KJ, Schell WA, Covel J, Duboc G, Giamberardino C, Kapoor M, Moloney M, Soltow QA, Tenor JL, Toffaletti DL, Trzoss M, Webb P, Perfect JR. 2018. *In vitro* and *in vivo* evaluation of APX001A/APX001 and other Gwt1 inhibitors against *Cryptococcus*. Antimicrob Agents Chemother 62:e00523-18.

51. Shaheen SK, Juvvadi PR, Allen J, Shwab EK, Cole DC, Asfaw YG, Kapoor M, Shaw KJ, Steinbach WJ. *In vitro* activity of APX2041, a new Gwt1 inhibitor, and *in vivo* efficacy of the prodrug APX2104 against *Aspergillus fumigatus*. Antimicrob Agents Chemother 65:e00682-21.

52. Wong GK, Griffith S, Kojima I, Demain AL. 1998. Antifungal activities of rapamycin and its derivatives, prolylrapamycin, 32-desmethylrapamycin, and 32-desmethoxyrapamycin. J Antibiot 51:487–491.

53. Cruz MC, Cavallo LM, Görlach JM, Cox G, Perfect JR, Cardenas ME, Heitman J. 1999. Rapamycin antifungal action is mediated via conserved complexes with FKBP12 and TOR kinase homologs in *Cryptococcus neoformans*. Mol Cell Biol 19:4101–4112.

54. Nielsen K, Cox GM, Wang P, Toffaletti DL, Perfect JR, Heitman J. 2003. Sexual cycle of *Cryptococcus neoformans var. grubii* and virulence of congenic **a** and α isolates. Infect Immun 71:4831–4841.

55. Moore TD, Edman JC. 1993. The alpha-mating type locus of *Cryptococcus neoformans* contains a peptide pheromone gene. Mol Cell Biol 13:1962–1970.

56. Fox DS, Cruz MC, Sia RA, Ke H, Cox GM, Cardenas ME, Heitman J. 2001. Calcineurin regulatory subunit is essential for virulence and mediates interactions with FKBP12-FK506 in *Cryptococcus neoformans.* Mol Microbiol 39:835–849.

57. Cruz MC, Goldstein AL, Blankenship J, Del Poeta M, Perfect JR, McCusker JH, Bennani YL, Cardenas ME, Heitman J. 2001. Rapamycin and less immunosuppressive analogs are toxic to *Candida albicans* and *Cryptococcus neoformans* via FKBP12-dependent inhibition of TOR. Antimicrob Agents Chemother 45:3162–3170.

58. Falloon K, Juvvadi PR, Richards AD, Vargas-Muñiz JM, Renshaw H, Steinbach WJ. 2015. Characterization of the FKBP12-encoding genes in *Aspergillus fumigatus.* PLoS One 10:e0137869.

Fig. 1. Chemical structures of novel C-22 modified FK520 analogs synthesized in this study. (*A*) One-step synthesis for C-22 modified FK520 analogs. Condition **A**: RNHNH$_2$, EtOH, 90 °C, 48 h; Condition **B**: RNHNH$_2$, MeOH, TFA (cat), 25 °C, 48 h. (*B*) Structures of 12 analogs modified with a hydrazide or a substituted hydrazine at the C-22 position of FK520 are depicted.

Table 1 MICs for FK506, FK520, and analogs[a]

Compound	MIC/ MEC (µg/mL) against:		
	Cryptococcus neoformans (H99)	*Candida albicans* (SC5314)	*Aspergillus fumigatus* (akuB^Ku80)
FK506	0.05	0.02	0.04
FK520	0.05	0.02	0.16
JH-FK-08	0.8	0.4	1.3
JH-FK-09	0.8	0.8	1.3
JH-FK-12	0.8	0.8	2.5
JH-FK-13	>50	>50	>10
JH-FK-14	1.6	0.8	0.6
JH-FK-15	3.1	0.4	0.3
JH-FK-16	0.4	0.4	1.3
JH-FK-17	0.4	0.4	1.3
JH-FK-20	0.8	0.8	0.6
JH-FK-21	1.6	0.8	2.5
JH-FK-22	1.6	1.6	2.5
JH-FK-23	1.6	1.6	5

[a] MIC_{90} were tested at 37°C for *C. neoformans.* MIC_{80} were tested at 30°C with 2 µg/mL supplementary fluconazole for *C. albicans.* MEC was collected for *A. fumigatus* strains.

Fig. 2. *In vitro* immunosuppressive activity of FK506, FK520, and analogs and pharmacokinetic profile of JH-FK-08 compared to FK506. (*A*) The percentage of IL-2-producing CD4$^+$ T cells is plotted vs. drug concentration as a measure of immunosuppressive activity of calcineurin inhibitors. Newly synthesized analogs demonstrate reduced immunosuppression compared to controls FK506 and FK520. (n=3, error bars represent standard error of the mean (SEM), IC$_{50}$ values determined by nonlinear regression with variable slope analysis). (*B*) Whole blood drug concentrations are plotted over time following single dose of 40 mg/kg of FK506 or JH-FK-08 (n=3, no statistically significant difference between groups, repeated measures ANOVA with Greisser Greenhouse correction test).

Table 2. Therapeutic indices for FK506, FK520, and analogs.

FK506: R´ = CHCH$_2$
FK520: R´ = CH$_3$

C22-modified F520 analogs

Compound	R	H99 MIC$_{90}$ (μg/mL)	IL-2 IC$_{50}$ (nM)	Therapeutic Index
FK506	-	0.05	0.09	1.0
FK520	-	0.05	0.3	3.3
JH-FK-08	CO$_2$Me	0.8	42.6	29.6
JH-FK-09	C(=O)-4-pyridine	0.8	11.2	7.8
JH-FK-12	CONH$_2$	0.8	35.6	24.7
JH-FK-13	SO$_2$-4-Me-Ph	>50	>200	undefined
JH-FK-14	C(=O)-2-CONHNH$_2$-Ph	1.6	14.0	4.9
JH-FK-15	C(=O)-2-naphthalene	3.1	17.4	3.1
JH-FK-16	Et	0.4	5.9	8.2
JH-FK-17	C(=O)-4-F-Ph	0.4	9.4	13.1
JH-FK-20	C(=O)-4-NH$_2$-Ph	0.8	17.8	12.3
JH-FK-21	CONHNH$_2$	1.6	54.5	18.9
JH-FK-22	COCH$_2$OH	1.6	45.0	15.6
JH-FK-23	SO$_2$Me	1.6	70.5	24.5

Fig. 3. *In vivo* immunosuppression activity of JH-FK-08 is reduced compared to FK506. (*A*) Shown here are the production of T follicular helper (Tfh) cells during immune response in C57BL/6 female mice. (*B*) Shown here are representative flow cytometry plots showing the frequencies of Tfh cells among CD44hi CD4+ T cells. Mice received twice daily treatment of vehicle, FK506 (2.5 mg/kg), JH-FK-08 (20 mg/kg), or JH-FK-08 (40 mg/kg) (n=5; **, p<0.001, ordinary One-way ANOVA with Tukey's test).

Fig. 4. JH-FK-08 treatment significantly reduced organ fungal burden in infected animals. Female A/J mice (n=5) were infected with *C. neoformans* strain H99 through intranasal inoculation and treated twice daily with vehicle, 6 mg/kg fluconazole, 40 mg/kg JH-FK-08, or combination through i.p. dosing. Fungal burden levels in lung (*A*), brain (*B*), and spleen (*C*) were assessed after 14 days of treatment. (*** , p<0.0001; ** , p=0.0018, ordinary 1-way ANOVA with Dunnet's test, error bars represent SEM)

Fig. 5. Murine survival is significantly extended by treatment with JH-FK-08 alone or in combination with fluconazole. Female A/J mice (n=10) were intranasally infected with *C. neoformans* strain H99 and received twice daily dosing of 6 mg/kg fluconazole, 40 mg/kg JH-FK-08, or a combination of the two drugs for the first two weeks post infection. Survival was monitored following the treatments (* , p=0.0065 found between JH-FK-08 and combination treatment groups; *** , p<0.0001 found between all other groups, Mantel Cox test).