1

4   Metabolomic analysis of *Drosophila melanogaster* larvae lacking Pyruvate kinase

6   **Authors**

7   Yasaman Heidarian[1], Jason P. Tourigny[1], Tess D. Fasteen[1], Nader H.

8   Mahmoudzadeh[1], Alexander J. Hurlburt[1], Travis Nemkov[2], Julie A. Reisz[2], Angelo

9   D'Alessandro[2], and Jason M. Tennessen[1]

11   **Affiliations**

[1]Department of Biology, Indiana University, Bloomington, IN 47405, USA

[2]Department of Biochemistry and Molecular Genetics, Anschutz Medical Campus,

University of Colorado School of Medicine, Aurora, CO 80045, USA

ABSTRACT

Pyruvate kinase (Pyk) is a rate-limiting enzyme that catalyzes the final metabolic

reaction in glycolysis. The importance of this enzyme, however, extends far beyond

ATP production, as Pyk is also known to regulate tissue growth, cell proliferation, and

development. Studies of this enzyme in *Drosophila melanogaster*, however, are

complicated by the fact that the fly genome encodes six Pyk paralogs whose functions

remain poorly defined. To address this issue, we used sequence distance and

phylogenetic approaches to demonstrate that the gene *Pyk* encodes the enzyme most

similar to the mammalian Pyk orthologs, while the other five *Drosophila* Pyk paralogs

have significantly diverged from the canonical enzyme. Consistent with this observation,

metabolomic studies of two different *Pyk* mutant backgrounds revealed that larvae

lacking Pyk exhibit a severe block in glycolysis, with a buildup of glycolytic intermediates

upstream of pyruvate. However, our analysis also unexpectedly reveals that steady

state pyruvate levels are unchanged in *Pyk* mutants, indicating that larval metabolism

maintains pyruvate pool size despite severe metabolic limitations. Consistent with our

metabolomic findings, a complementary RNA-seq analysis revealed that genes involved

in lipid metabolism and peptidase activity are elevated in *Pyk* mutants, again indicating

that loss of this glycolytic enzyme induces compensatory changes in other aspects of

metabolism. Overall, our study provides both insight into how Drosophila larval

metabolism adapts to disruption of glycolytic metabolism as well as immediate clinical

relevance, considering that Pyk deficiency is the most common congenital enzymatic

defect in humans.

INTRODUCTION

The enzyme Pyruvate kinase (Pyk; E.C. 2.7.1.40) catalyzes the final reaction of glycolysis (phosphoenolpyruvate + ADP + Pi to pyruvate + ATP). Since the Pyk-catalyzed reaction is essentially irreversible under normal cellular conditions (SCHORMANN *et al.* 2019), this enzyme controls a key rate-limiting, "pay-off" glycolytic step that controls ATP production, and dictates the rate at which glycolytic intermediates enter the metabolic pathways that branch off both glycolysis and the TCA cycle. In so doing, Pyk regulates fluxes towards serine synthesis, which indirectly influences redox balance and nucleoside anabolism via the amino acids cysteine (de novo glutathione synthesis) and glycine (one-carbon metabolism for de novo purine synthesis) (MADDOCKS *et al.* 2013). Thus, changes in Pyk activity can influence cell growth, proliferation, and differentiation by balancing the flux of glycolytic metabolites between mitochondrial energy production and the synthesis of metabolites involved in chromatin remodeling, one-carbon metabolism, and biomass production (ISRAELSEN AND VANDER HEIDEN 2015; HSU AND HUNG 2018).

The importance of Pyk in human health is clear from the myriad of diseases that are caused by altered activity of the two human orthologs, Pyruvate Kinase, Liver and Red Blood Cell (PKLR) and Pyruvate Kinase, Muscle (PKM). For example, mutations in the *PKLR* gene result in a common congenital disorder known as Pyk deficiency (PKD) (FATTIZZO *et al.* 2022; LUKE *et al.* 2023). Individuals with PKD experience a wide spectrum of clinical and biochemical symptoms and have a highly variable prognoses that depend on environmental factors, early interventions, and the severity of the causal mutations (DOLAN *et al.* 2002; GRACE *et al.* 2015). Similarly, changes in splicing of PKM

transcripts are well-documented in a number of disease states, with a switch from the

PKM1 isoform towards the PKM2 isoform associated with pulmonary hypertension

(CARUSO *et al.* 2017; ZHANG *et al.* 2017), sporadic Alzheimer's disease (TRAXLER *et al.*

2022), and enhancement of the Warburg effect in cancer cells (ISRAELSEN AND VANDER

HEIDEN 2015). Finally, population genetic studies support a model in which *PKLR*

mutations are protective against malaria (MIN-OO *et al.* 2007; AYI *et al.* 2008; MACHADO *et*

al. 2012; VAN BRUGGEN *et al.* 2015), suggesting that partial inhibition of the enzyme could

be exploited toward anti-malarial efforts. Basic studies of PYK function in the context of

endogenous metabolic networks thus hold the potential to inform better interventions for

a wide range of human diseases.

The fruit fly *Drosophila melanogaster* is a powerful model for studying the

function of metabolic enzymes within the context of both normal physiology and human

disease models (DRUMMOND-BARBOSA AND TENNESSEN 2020; KIM *et al.* 2021). In this

regard, previous studies of *Drosophila* Pyk have demonstrated roles for this enzyme in

fly development and behavior, with reduced *Pyk* levels being associated with defects in

muscle and wing development, neuron and glial function, and olfactory memory (TIXIER

et al. 2013; VOLKENHOFF *et al.* 2015; WU *et al.* 2018; SPANNL *et al.* 2020). However, the

effects of decreased Pyk activity in the fly have not been examined at a metabolic level.

Moreover, the fly genome encodes six predicted Pyk paralogs (Pyk, CG7069, CG2964,

CG7362, CG11249, and CG12229,GARAPATI *et al.* 2019; GRAMATES *et al.* 2022b), thus

highlighting a need to characterize the *in vivo* functional potential of individual enzymes

within this gene class.

Here, we use sequence identities and phylogenetics in combination with

metabolomics and transcriptomics to analyze *Pyk* (FBgn0267385), which encodes the

most abundantly and widely expressed member of the *Drosophila melanogaster* Pyk

family (GRAVELEY *et al.* 2011). Our studies indicate that the gene *Pyk* encodes the

protein most analogous to the mammalian PKM and PKLR enzymes. Subsequent

metabolomic analysis of two genetically distinct *Pyk* mutant backgrounds reveals a

severe block in glycolysis, with a buildup of glycolytic intermediates and a depletion of

lactate, 2-hydroxyglutarate (2HG), and the amino acid proline. Moreover, the loss of Pyk

activity also induced a significant upregulation of genes involved in lipid metabolism and

protease activity. Overall, our study validates the presumed function of Pyk within

Drosophila metabolism and identifies the transcriptional and metabolic networks that

respond to loss of Pyk activity during larval development.

METHODS

Drosophila husbandry and larval collection

Fly stocks were maintained on Bloomington *Drosophila* Stock Center media at 25ºC. Larvae were collected on molasses agar plates as previously described (LI AND TENNESSEN 2017; LI AND TENNESSEN 2018). Briefly, 50 virgin females and 25 males of the appropriate genotypes were placed in six-ounce plastic bottles (Genesee Scientific; Cat #: 32-130) with holes punched in the side using a 20-gauge needle. A 35 mm molasses agar plate (Corning 353001) with a smear of yeast paste on the surface was taped in place in the mouth of the bottle and the bottle inverted and placed in a 25ºC incubator. The molasses agar/yeast plate was replaced at least once per day. For sample collection, eggs were collected on the molasses agar/yeast plate for 6 hours, individual plates were placed inside a 60 mm cell culture dish, and the collected embryos were placed in a 25ºC incubator and allowed to develop for 48 hours. Larvae were collected at the early L2 stage for subsequent metabolomics analysis.

Generation of *Pyk* alleles using CRISPR/Cas9

The *Pyk* deletions Pyk^{23} and Pyk^{31} were generated using a previously described approach for CRISPR/Cas9 mutagenesis (GRATZ *et al.* 2013; SEBO *et al.* 2014). Briefly, oligos containing gRNA sequences #1 (5'- GTGCCCCATGTGCGTCTGTC-3') and #2 (5'- GCTGCTGGAGGCAGGTCCGA-3') were inserted into pU6-BbsI-chiRNA (DGRC Stock #1362) and the resulting plasmids were independently injected into BDSC Stock #52669 (y^1 *M{RFP[3xP3.PB] GFP[E.3xP3]=vas-Cas9.S}ZH-2A* w^{1118}) by Rainbow Transgenics (Camarillo, CA). Injected females and F1 progeny were crossed to *w*; ry^{506}*

Dr^1/TM3, P{Dfd-GMR-nvYFP}3, Sb^1. Putative F2 *Pyk^Δ/ TM3, P{Dfd-GMR-nvYFP}3, Sb^1*

siblings were mated and the F3 generation screened for animals lacking the *TM3*

balancer, indicating the presence of a lethal mutation. Any strain that failed to generate

homozygous mutant adults was crossed with *Pyk^{61}* mutants and the F1 progeny

analyzed for failure to complement. Mutant strains that failed to complement *Pyk^{61}* were

further analyzed for the presence of *Pyk* mutations. *Pyk^{23}* was isolated from injections

using gRNA#1 and *Pyk^{31}* was isolated from injections using gRNA#2. The deletions

were identified by amplifying and sequencing a region of the *Pyk* gene using oligos 5'-

CACGCACTTTGTTTACATCAGC-3' and 5'-GCACCAGTCCACGGTAGAGA-3'.

Generation of *Pyk* alleles using p-element excision

Both the *Pyk^{60} and Pyk^{61}* alleles were generated using standard techniques for

imprecise p-element excision (ROBERTSON *et al.* 1988; BELLEN *et al.* 2004). Briefly, virgin

females containing the transposon insertions *Pyk^{DG05605}/TM3, Sb^1, Ser^1* and

Pyk^{EY10213}/TM3, Sb^1, Ser^1 (BDSC stock 20088) were independently crossed with *ry506*

p{Δ2-3}99B males. F1 virgin females lacking the TM3 balancer were crossed with *w*;*

ry^{506} Dr^1/TM3, P{Dfd-GMR-nvYFP}3, Sb^1 males (BDSC stock 23231). F2 males with

white eyes were then crossed to *w*; ry^{506} Dr^1/TM3, P{Dfd-GMR-nvYFP}3, Sb^1* females

and F3 *Pyk^Δ/ TM3, P{Dfd-GMR-nvYFP}3, Sb^1* siblings were used to establish mutant

lines. The *Pyk^{60}* allele was generated by the imprecise excision of *Pyk^{EY21058}* and the

Pyk^{61} deletion was generated by the imprecise excision of *Pyk^{DG05605}*. Endpoints of the

deletions were mapped using a PCR-based approach. The *Pyk^{61}* deletion was

sequenced by isolating homozygous mutant larvae, amplifying across the deletion using

142 the oligos 5'-GATTTCCTTCAGAGCATTTGCGTC-3' and 5'-

143 TTCACCGTGCAGCAAGACATC-3' and sequencing the resulting PCR product. The

144 Pyk^{61} deletion is 6400 bp long starting with the sequence 5'-

145 GATGCCTTTGTTGCCGCCCT-3' and ending with the sequence 5'-

146 GTATGAACTTCTCTCACGGC-3' and includes an 18 bp insertion of the sequence 5'-

147 AAGTTCAAGTTCTGGATT-3'. The Pyk^{60} allele represents a large deletion or

148 rearrangement between the sequences 5'-CTTCTGTTCTATCCGATTGCCGG-3' and

149 5'-TTGACGCGCTCATTGGGTTTC-3'. However, we were unable to produce a PCR

150 product spanning the lesion. The Pyk^{prec} allele is a precise excision of $Pyk^{DG05605}$. For all

151 analyses, the Pyk^{60}/TM3, P{Dfd-GMR-nvYFP}3, Sb^1 and Pyk^{61}/TM3, P{Dfd-GMR-

152 nvYFP}3, Sb^1 strains were crossed together and trans-heterozygous $Pyk^{60/61}$ mutant

153 larvae were selected for based on the absence of YFP.

154  Since the transposon excision events that produced the Pyk^{60} and Pyk^{61} deletion

155 alleles also disrupted neighboring genes, we further analyzed the effects of these

156 deletions on expression of neighboring genes. When these deletions are placed in

157 *trans*, the resulting heterozygote lacks sequence corresponding to nearly the entire *Pyk*

158 coding region and the 5' exons of the neighboring *Pyk* homolog *CG7069*. Northern blot

159 analysis of trans-heterozygous L2 larvae revealed that $Pyk^{60/61}$ larvae fail to accumulate

160 detectable *Pyk* mRNA transcripts (Figure S1). The mRNA of neighboring genes *Polr3F*

161 and *CG18596*, however, were present at similar levels in both Pyk^{Prec} and $Pyk^{60/61}$,

162 indicating that the phenotypes arising from the *Pyk* mutant background are specific to

163 loss of *Pyk* expression. Of note, the second pyruvate kinase homolog disrupted by

164 these deletions, *CG7069*, is not expressed during the embryonic and larval stages, and

165   since *Pyk*[60/61] larvae die prior to the L3 stage, the effects of these deletions on *CG7069*

166   expression could not be examined.

167
Northern Blot Analysis
RNA was extracted from L2 larvae with TriPure isolation reagent and northern blot

analysis was conducted as previously described (KARIM AND THUMMEL 1991; SULLIVAN

AND THUMMEL 2003). The following PCR oligos were used to generate northern blot

probes:

Pyk: 5'-GCTGACCACCAACAAGGAAT-3' and 5'-GCACCAGTCCACGGTAGAGA-3'

Polr3F: 5'-ACCAACGATGACCTGACCAAG-3' and 5'-ATTGTTTCCAGATCGGCCTCC-

3'

rp49: 5'-ACAAATGTGTATTCCGACCACG-3' and 5'-TCAAGATGACCATCCGCCCAG-

3'

Phylogenetic Analysis of Pyk orthologs and paralogs

Protein sequences for each of the Pyk homologs were retrieved from FlyBase

(GRAMATES *et al.* 2022a), Saccharomyces Genome Database (ENGEL *et al.* 2014; WONG

et al. 2023), WormBase (DAVIS *et al.* 2022), NCBI RefSeq (O'LEARY *et al.* 2016), TAIR

(BERARDINI *et al.* 2015), and UniProt (2023). The longest verified isoform of each was

selected for multiple sequence alignment (MSA; see Table S1 for protein sequence

information), except for mosquito which was a predicted entry (PETCHAMPAI *et al.* 2019).

For the sequence identity and similarity heatmaps, an ensemble MSA was performed

via the M-Coffee web server (WALLACE *et al.* 2006; MORETTI *et al.* 2007; NGUYEN *et al.*

2015; MINH *et al.* 2020), running the default command: `t_coffee -

in=Pyk_homologs_E_coli_out.fa Mpcma_msa Mmafft_msa Mclustalw_msa

Mdialigntx_msa Mpoa_msa Mmuscle_msa Mprobcons_msa Mt_coffee_msa -

output=score_html clustalw_aln fasta_aln score_ascii phylip -tree -maxnseq=150 -

maxlen=2500 -case=upper -seqnos=off -outorder=input -run_name=result -

multi_core=4 -quiet=stdout'. The resulting MSA was analyzed using the `seqidentity()'

function of the Bio3D R package to extract pairwise matrices (GRANT *et al.* 2006), which

were visualized with corrplot (WEI AND V. 2021).

For the phylogenetic analysis, we first performed MSA with MAFFT v7.520 using

the E-INS-I algorithm to tolerate potentially large gaps between conserved segments

(KATOH AND STANDLEY 2013), with the command `mafft –maxiterate 1000 –thread 4 –

genafpair Pyk_homologs_incl_Aa_and_E_coli_outgroup.fa >

mafft_einsi_output.msa.fa'. We then reconstructed the phylogenetic tree with IQ-TREE

2 using the WAG model of protein substitution (WHELAN AND GOLDMAN 2001; NGUYEN *et*

al. 2015; HOANG *et al.* 2018; MINH *et al.* 2020), run with the command `iqtree2 -s

mafft_einsi_output.msa.fa –redo -B 1000 -T 4 -m WAG –prefix mafft_einsi_WAG'. The

resulting phylogeny was rooted to *E. coli*, formatted, and visualized using FigTree v1.4.4

(RAMBAUT 2018).

Triglyceride Assays and Nile Red Staining

Triglyceride levels in *Pyk* control and mutant L2 larvae were measured as

previously described (TENNESSEN *et al.* 2014). Briefly, synchronized populations of

embryos were collected for 4 hrs on a molasses agar plate with yeast paste smeared on

the surface, as described (LI AND TENNESSEN 2017). Embryos were allowed to hatch on

the surface of the collection plate and larvae were reared on the same plate in a 25 ºC

incubator. Sixty hours after egg-laying, heterozygous controls and homozygous mutant

larvae were identified based on lack of GFP expression from the *TM3, P{Dfd-GMR-*

nvYFP}3, Sb^1 balancer chromosome. Twenty-five larvae of the appropriate genotypes

were then collected in 1.5 mL microfuge tubes, washed three times with phosphate-

buffered saline pH 7.0 (PBS), homogenized in 100 µl of cold PBS + 0.05% Tween 20

(PBST), and heat-treated for 10 min at 70ºC. The resulting homogenate was assayed

for triglyceride and soluble protein levels as previously described (TENNESSEN *et al.*

2014).

Nile red staining was conducted on L2 fat bodies as previously described

(GRÖNKE *et al.* 2005). Briefly, dissected tissues were fixed with 4% paraformaldehyde in

PBS for 30 min and washed with PBST three times for 5 minutes each, and then

incubated for 1 hr at room temperature in a 1:1000 dilution of Nile Red stock (10mg/mL

in acetone) in PBS. Tissues were washed three times for 5 minutes each in PBST,

mounted in Vectashield (Vector Laboratories), and imaged on a Leica SP8 Confocal at

568nm.

Gas Chromatography Mass Spectrometry (GC-MS)-based Metabolomics

Pyk mutant and control larvae were collected at the early L2 stage (~48 hours after egg-

laying at 25ºC) as previously described (LI AND TENNESSEN 2018). All samples collected

for analysis represent biological replicates obtained from independent mating bottles

and each sample contained 25 larvae. Analysis of the *Pyk^{60} and Pyk^{61}* alleles was

conducted by the University of Utah metabolomics core facility as previously described

(Cox *et al.* 2017). *Pyk*$^{23/31}$ mutants and *Pyk*$^{23/+}$ control samples were analyzed by the

University of Colorado Metabolomics core facility. Data were normalized to sample

mass and internal standards.

Ultra High-pressure Liquid Chromatography - Mass Spectrometry (UHPLC-MS)-

based Metabolomics UHPLC-MS metabolomics analyses were performed at the

University of Colorado Anschutz Medical Campus, as previously described (Nемкоv *et*

al. 2019). Briefly, the analytical platform employs a Vanquish UHPLC system (Thermo

Fisher Scientific, San Jose, CA, USA) coupled online to a Q Exactive mass

spectrometer (Thermo Fisher Scientific, San Jose, CA, USA). The (semi)polar extracts

were resolved over a Kinetex C18 column, 2.1 x 150 mm, 1.7 µm particle size

(Phenomenex, Torrance, CA, USA) equipped with a guard column (SecurityGuard^{TM}

Ultracartridge – UHPLC C18 for 2.1 mm ID Columns – AJO-8782 – Phenomenex,

Torrance, CA, USA) using an aqueous phase (A) of water and 0.1% formic acid and a

mobile phase (B) of acetonitrile and 0.1% formic acid for positive ion polarity mode, and

an aqueous phase (A) of water:acetonitrile (95:5) with 1 mM ammonium acetate and a

mobile phase (B) of acetonitrile:water (95:5) with 1 mM ammonium acetate for negative

ion polarity mode. The Q Exactive mass spectrometer (Thermo Fisher Scientific, San

Jose, CA, USA) was operated independently in positive or negative ion mode, scanning

in Full MS mode (2 µscans) from 60 to 900 m/z at 70,000 resolution, with 4 kV spray

voltage, 45 sheath gas, 15 auxiliary gas. Calibration was performed prior to analysis

using the Pierce^{TM} Positive and Negative Ion Calibration Solutions (Thermo Fisher

Scientific).

Statistical Analysis of Metabolomics Data

Both metabolomics datasets were analyzed using Metaboanalyst 5.0 (Pang *et al.* 2021), with data first preprocessed using log normalization and Pareto scaling.

RNA-seq Analysis

RNA from *Pyk*$^{23/+}$ heterozygous controls and *Pyk*$^{23/31}$ mutants was extracted from L2 larvae with TriPure isolation reagent and further purified using a Qiagen RNeasy kit (Catalog #74004). RNA-seq was performed on three whole-fly biological replicates for *Pyk*$^{23/+}$ heterozygous controls and *Pyk*$^{23/31}$ mutants. Samples were paired-end sequenced on the NextSeq 550 platform 75 cycles to a depth of 15-20 million reads each, using standard Illumina TruSeq Stranded mRNA libraries, at the IU Center for Genomics and Bioinformatics .

RNA-seq read quality was assessed with FastQC (Andrews 2010) and MultiQC (Ewels *et al.* 2016), with adequate reads and no significant issues noted. Reads were pseudo-aligned and quantified using Kallisto v0.46.0 (Bray *et al.* 2016), the *D. melanogaster* BDGP6.32 reference assembly and annotation retrieved through Ensembl (Cunningham *et al.* 2022), and the gffread utility from the Cufflinks suite to generate the transcriptome from the assembly and annotation (Trapnell *et al.* 2010).

Differential expression analysis was performed with DESeq2 v1.30.1 (Love *et al.* 2014) running in RStudio v1.4.1717 on R v4.0.4. Genes with average expression below 2 counts per sample (i.e. 48 counts per row) were filtered out before determining genes with significant expression difference in at least one sample with the likelihood-ratio test (LRT) and LRT *p*-value ≤ 0.05. Subsequent analysis and visualization were performed

in R using a variety of tools and packages. Correlation plots were generated using base

R and the corrplot package v0.92 (WEI AND SIMKO 2021). Principal components were

analyzed and visualized with PCAtools v2.2.0 (BLIGHE AND LUN 2022). Heatmaps,

including clustering analysis performed by the hclust method (from base R {stats}

package), were generated by pheatmap v1.0.12 (KOLDE 2019).

Gene Ontology Analysis was carried out using the GOrilla tool for genes that

were significantly (*P*-value <0.05) up-regulated or down-regulated by more than two-fold

in *Pyk* mutants (EDEN *et al.* 2009). Tissue enrichment analysis was conducted using

default settings for the "Preferred tissue (modEncode RNA_seq)" tool in PANGEA

version 1.1 beta (HU *et al.* 2023).

Data Availability

All strains and reagents are available upon request. All targeted metabolomics data

described herein are included in Tables S2 and S3. Processed RNA-seq data is

presented in Table S4 and available in NCBI Gene Expression Omnibus (GEO).

RESULTS

299
The *Drosophila* gene *Pyk* encodes the Pyk enzyme orthologous to human PKs

The *Drosophila* genome harbors six genes predicted to encode pyruvate kinases

(GARAPATI *et al.* 2019; GRAMATES *et al.* 2022b). To better understand the relationships

between the *Drosophila* Pyk paralogs and Pyk orthologs in other organisms, we

compared their pairwise protein sequence identities and generated a phylogenetic gene

tree to infer their evolutionary history (Figure 1, Table S1). Species examined included

human and mouse, *Aedes aegypti, Caenorhabditis elegans*, *Saccharomyces cerevisiae*,

and *Arabidopsis thaliana,* with *Escherichia coli* as the outgroup for tree rooting (Table

S1). The pairwise sequence comparisons indicate that *Drosophila* Pyk shares strongest

identity with that of the *Aedes aegypti* protein PK (XP_001649983.1), followed by those

of its mammalian orthologs, the worm paralogs, and fly paralog CG7069 (Figure 1A).

The other putative paralogs show identity to a lesser extent, with two of the *Drosophila*

proteins (CG11249 and CG12229) displaying less than 20% identity with all other

analyzed Pyk orthologs (Figure 1A). The sequence similarities, allowing for

conservative residue substitutions, show a similar pattern (Figure S2). These results

indicate that, at the sequence level, *Drosophila* Pyk is as or more similar to the yeast

orthologs as it is to 4 of the 5 other putative fly paralogs.

Sequence distances may not correspond to actual evolutionary or functional

relationships, so we next sought to reconstruct the Pyk gene tree using a maximum

likelihood model of protein evolution. This phylogeny indicates that the *Drosophila* Pyk

is sister to its *Aedes aegypti* ortholog (Figure 1B), which composes a larger clade with

the more distantly related *Drosophila* paralogs. This insect Pyk/PK clade is in turn sister

to that of the mammalian orthologs, which share a common ancestor. We note strong

ultrafast bootstrap support for the insect and mammalian ortholog subclades, the yeast

and worm paralogs, and two pairs of the fly paralogs, with weak support for the fly

paralog internal branches and the relationships between species (Figure 1B).

Aedes PK1 was previously demonstrated to exhibit unique allosteric regulation

when compared with mammalian enzymes (PETCHAMPAI *et al.* 2019). Given the high

sequence similarity and close evolutionary relationship between these two Dipteran

proteins, our findings suggest that *Drosophila* Pyk functions in a similar manner to that

of mosquito PK. Overall, these results support the hypothesis that the fly gene *Pyk*

encodes the canonical glycolytic enzyme, which is consistent with previous studies

demonstrating that *Pyk* is the most widely expressed member of the *Drosophila*

melanogaster Pyk gene class (RUST AND COLLIER 1985; GRAVELEY *et al.* 2011).

334
335
Pyk mutants die during the second larval instar

To determine how loss of Pyk activity affects *Drosophila* development, we used

CRISPR/Cas9 to generate two small deletions in the *Pyk* locus, *Pyk*^{23} and *Pyk*^{31}, both of

which induce frameshifts in the first coding exon and are predicted to be null alleles

(Figure 2A). When these two *Pyk* mutant alleles were placed in trans, the resulting

Pyk$^{23/31}$ mutants survived embryogenesis but died during the mid-L2 stage (Figure 2B).

Moreover, when compared with *Pyk*$^{23/+}$ heterozygous controls, *Pyk* mutants displayed a

significant decrease in both body mass and triglyceride stores (Figure 2C and 3A, B),

thus demonstrating that loss of Pyk activity significantly disrupts larval development. We

would note, however, that although *Pyk* mutants accumulate far less triglycerides (TAG)

than controls, the density of lipid droplets within the fat body as determined by Nile Red

staining (Figure 3A,B) remains similar to controls. Overall, these findings reveal that *Pyk*

mutant larvae exhibit many of the same phenotypes observed in *dERR* and *Pfk*

mutants, which also exhibit severe disruptions of glycolytic metabolism (RUST AND

COLLIER 1985; TENNESSEN *et al.* 2011).

Metabolomic analysis of *Pyk* mutants

To better understand how loss of *Pyk* alters larval metabolism, we used a

targeted metabolomics approach to compare $Pyk^{23/31}$ mutants with $Pyk^{23/+}$ heterozygous

controls during the L2 larval stage (Table S2). Partial Least Squares Discriminant

Analysis (PLS-DA) of the resulting data revealed that control and mutant samples

clustered in distinct groups (Figure S3). Subsequent analysis of the data revealed

increased levels of the metabolites immediately upstream of Pyk, 2/3-phosphoglycerate

and phosphoenolpyruvate, and a decrease in lactate, and 2-hydroxyglutarate, and most

amino acids that can be used as anaplerotic fuel in the citric acid cycle (Figure 4A-B,

Table S2). Overall, these changes are indicative of a block in glycolysis at the reaction

catalyzed by Pyk.

While the metabolite changes observed in our initial metabolomics dataset were

largely expected, we were surprised to find that steady state pyruvate levels were

unchanged in *Pyk* mutants (Table S2). To verify that pyruvate levels remained constant

even in the absence of Pyk, we analyzed a second series of *Pyk* mutants using a

different methodology in an independent metabolomics core facility. In this case, a

trans-heterozygous combination of two *Pyk* deletion alleles, $Pyk^{60/61}$, were compared

with a genetically-matched control strain that was homozygous for a precise excision

allele (Table S3; see Methods for a description of the alleles used). Although these

deletions also removed portions of *CG7069* (Figure 2A), which encodes another

member of the Pyk family, this gene is not expressed during most of larval

development, and thus our analysis likely reflects phenotypes caused by the loss of the

Pyk locus. Similar to the metabolomic profile of *Pyk*$^{23/31}$ mutants, *Pyk*$^{60/61}$ mutant L2

larvae exhibited increased levels of glycolytic metabolites immediately upstream of

pyruvate kinase activity (2-phosphoglycerate, 3-phosphoglycerate and

phosphoenolpyruvate), and decreased levels of the downstream metabolites lactate,

2HG, and several amino acids (Figure 5A,B; Table S3). Moreover, these *Pyk* mutant

larval samples also displayed normal pyruvate levels (Figure 5A), suggesting that

steady state levels of larval pyruvate are maintained by compensatory changes in other

metabolic pathways.

As an extension of our metabolomic studies, we compared our datasets to

identify those metabolites that were similarly altered in both *Pyk* mutant backgrounds.

While we would note that this comparative analysis is somewhat limited in that we used

two different targeted metabolomics protocols that were optimized to detect different

subsets of metabolites, molecules that appear in both analyses highlight metabolic

pathways that are significantly altered by loss of Pyk activity in different genetic

backgrounds. Despite the many metabolic changes observed in each individual dataset,

only a handful of metabolites were common among the two experiments: 2/3-

phosphoglycerate, phosphoenolpyruvate, lactate, 2-hydroxyglutarate, and proline

(Figure 6A-C). While the changes in 2/3-phosphoglycerate, phosphoenolpyruvate, and

lactate were to be expected, the decrease in 2HG and proline present an interesting

model for how metabolic flux could be rewired in the absence of Pyk activity. The

decrease in 2HG likely represents L-2HG, which is abundant in larvae (LI *et al.* 2017;

MAHMOUDZADEH *et al.* 2020). Since L-2HG is synthesized from α-ketoglutarate by LDH

and is sensitive to changes in lactate production (LI *et al.* 2017; LI *et al.* 2018), we

hypothesize that the decreased abundance of 2HG represents a smaller lactate pool

combined with a potential decrease in TCA cycle flux. Meanwhile, the decreased proline

levels present in *Pyk* mutants are quite informative because this amino acid is

commonly used by insects as an anaplerotic fuel for the TCA cycle (BURSELL 1981).

Based on this observation, future studies should determine if *Pyk* mutants are using

proline to sustain both pyruvate levels and early larval growth.

Gene Expression Analysis of *Pyk* mutants

Our metabolomic studies suggest that loss of Pyk activity induces compensatory

changes in other metabolic pathways that allow the larvae to survive early development.

To better understand if this metabolic rewiring is recapitulated at the level of gene

expression, we used RNA-seq to compare the expression profiles of $Pyk^{23/31}$ mutant

larvae with $Pyk^{23/+}$ heterozygous controls (Table S4). This analysis revealed significant

differences in the gene expression between mutant and control samples (Figure

S5A,B), with 755 genes up-regulated and 214 genes downregulated more than two-fold

in the *Pyk* mutant dataset (adjusted *P* value <= 0.05; Table S4, Figure 7). Notably, the

expression of the *Pyk* gene itself was down-regulated 8.3-fold when compared with the

control strain, thus demonstrating the severity of these deletion alleles (Table S4, Figure

7). The data also revealed a significant upregulation of the gene *Thor* (FBgn0261560),

the fly homolog of 4E-BP (BERNAL AND KIMBRELL 2000), which functions as a metabolic

brake that dampens mRNA translation (HAY AND SONENBERG 2004; TELEMAN *et al.*

2005). The significant upregulation of *Thor* transcript levels hints at one possible

mechanism by which *Pyk* mutants both rewire larval metabolism and slow growth in

response to a major metabolic insult.

To further analyze the *Pyk* mutant RNA-seq data, we used Gene Ontology (GO)

to better define the biological processes most affected by loss of Pyk activity (Figure

8A,B and Table S5). Our analysis revealed that *Pyk* mutant samples exhibited a

significant enrichment in GO categories associated with the digestion and metabolism of

chitin and glucosamines, lipids, and proteins (Figure 8A). Notably, analysis of the RNA-

seq data set using the "Preferred tissue" tool in PANGEA indicated that the digestive

system displays the largest number of significantly up-regulated genes in *Pyk* mutants

(Table S6) (HU *et al.* 2023), with many of the those genes being involved in lipid and

protein metabolism (Figure 8C). For example, lipid metabolism was the second-most

enriched GO category, and; most of the significantly up-regulated genes that fit within

this GO category encode intestinal lipases. Similarly, *Pyk* mutants exhibit significant

increases in genes that encode proteases (Figure 8A, C), many of which represent

digestive proteases predicted to be expressed within the intestine. Overall, our results

suggest that the metabolism of *Pyk* mutants attempts to compensate for its loss by

increasing expression of genes involved in the breakdown of macromolecules within the

digestive tract, perhaps as a means of increasing the uptake of dietary nutrients.

DISCUSSION

Here we demonstrate that *Pyk* mutants exhibit a larval lethal phenotype with defects in larval growth and a severe block in glycolysis. While the phenotypic consequences of mutating *Pyk* are largely expected, our study contains several important findings for both the *Drosophila* and biomedical communities. First, we demonstrate that while the *Drosophila* genome encodes several pyruvate kinase orthologs, the gene *Pyk* encodes the canonical enzyme most analogous to Pyk in other organisms. In fact, *Drosophila* Pyk is more similar to the two yeast orthologs than to the other five pyruvate kinase orthologs present in the fly. Future studies should more closely examine these paralogous enzymes and determine why *Drosophila melanogaster* evolution produced such a specialized set of pyruvate kinases.

A second key finding from our sequence analysis is that *Drosophila* Pyk exhibits nearly equivalent sequence similarity to both the mammalian PKLR and PKM orthologs. This observation is important because PLKR and PKM are connected with distinct disease phenotypes. Thus, future studies in the fly should be cautious when making comparisons with disease states that are unique to either of the human enzymes. Instead, the *Pyk* mutants should be used as a tool to understand how cells and tissues respond to a general deficiency in Pyk activity, similar to previous studies that manipulated *Drosophila* Pyk activity in imaginal disc development and neurons/glia (TIXIER *et al.* 2013; VOLKENHOFF *et al.* 2015; WU *et al.* 2018; SPANNL *et al.* 2020). One exciting possibility would be to use our newly described mutants as a background for expressing disease causing variants of the human enzymes. After all, pyruvate kinase deficiency is the most common enzymatic disorder detected in newborns, and moreover, a wide variety of disease-causing PKLR variants have been identified (DOLAN

et al. 2002; GRACE *et al.* 2015; FATTIZZO *et al.* 2022; LUKE *et al.* 2023). Expression of the

wild-type and mutated enzyme variants in the fly would allow for a rapid genetic analysis

of how PKLR mutations alter cellular metabolism while also providing an *in vivo* platform

to test potential therapeutics.

Our studies of the *Pyk* mutant also reveal how a block in glycolysis alters

systemic metabolism in unexpected ways. Previous studies in the fly have examined

how disruption of glycolytic metabolism affects both the metabolome and transcriptome.

For example, the *Drosophila* Estrogen-Related Receptor (dERR) is a central regulator

of glycolytic metabolism throughout the fly lifecycle (TENNESSEN *et al.* 2011; BEEBE *et al.*

2020). Our studies reveal that the developmental defects exhibited by *Pyk* mutants

largely phenocopy those observed in *dERR* mutants although the *Pyk* and *dERR*

mutant metabolomic and transcriptomic profiles differ significantly. Unlike *dERR*

mutants, which display a buildup of sugars due to the down-regulated expression of

nearly every glycolytic enzyme (TENNESSEN *et al.* 2011; BEEBE *et al.* 2020), *Pyk* mutants

instead accumulate three glycolytic metabolites immediately upstream of the Pyk-

catalyzed reaction. Thus, our studies of *Pyk* highlight how the *dERR* mutant metabolic

profile represents a broad disruption of carbohydrate metabolism when compared with

the loss of a single *dERR* target gene.

While most aspects of the *Pyk* mutant metabolomics profile were expected based

on the known role of Pyk, several interesting observations emerged. First, steady state

levels of pyruvate were unchanged in *Pyk* mutants as compared with controls. Since the

other metabolites in glycolysis are changed in a manner that is indicative of a loss of

Pyk activity, and both lactate and TCA cycle intermediates exhibit a severe decrease,

flux into and out of the larval pyruvate pool must be severely compromised. Future

studies should explore this possibility using stable-isotope tracers and further examine

how larval metabolism maintains steady state pyruvate levels even under severe

metabolic stress.

Our analysis also raises questions about how *Pyk* mutants, as well as *dERR*

mutants, survive until the mid-L2 stage. Glycolysis is severely disrupted in both mutant

backgrounds, yet these animals continue to develop for almost three days when raised

under standard culture conditions. While this impressive resilience might be partially

explained by maternal loading of Pyk and other enzymes, both metabolomics and

transcriptomics analysis suggest that *Pyk* mutants switch fuel sources in an attempt to

maintain developmental progress. Notably, *Pyk* mutants exhibit a significant decrease in

the fermentation products lactate and 2HG, as well as a decrease in the amino acid

proline. All three changes signal a major reorganization of larval metabolism, as lactate

and L-2HG are present at high levels within normal larvae and represent pools of

carbon that are not immediately used for energy production and growth (LI *et al.* 2017).

Meanwhile, proline is commonly used for energy production in insects (TEULIER *et al.*

2016). Thus the depletion of lactate, 2HG, and proline steady state pools suggests that

Pyk mutant metabolism has become reliant on alternative energy sources, such as

proline, and lacks adequate resources to ferment pyruvate and α-ketoglutarate into

lactate and L-2HG, respectively. This hypothesis is also supported by RNA-seq

analysis, which uncovered a significant up-regulation of enzymes involved in the

digestion and metabolism of complex lipids and proteins. Overall, further studies of *Pyk*

mutants are warranted to understand how these metabolic adaptations are induced and

regulated.

Our results also revealed that *Pyk* mutants exhibit more severe developmental

phenotypes when compared with *Mpc1* mutants, which are unable to properly transport

pyruvate into the mitochondria (BRICKER *et al.* 2012). Considering that Pyk and Mpc1

both directly affect pyruvate metabolism, one might predict that mutations in these

genes would result in similar phenotypes. Instead, *Mpc1* mutants are viable when raised

on standard *Drosophila* media and only exhibit a lethal phenotype when placed on a

sugar-only diet (Bricker et al., 2012). Moreover, the metabolomics profile of *Mpc1* and

Pyk mutants display significant differences. Notably, serine levels are significantly

increased in *Mpc1* mutants – an observation that would be consistent with an

uncoupling of glycolysis and the TCA cycle causing a rerouting of glycolytic

intermediates into the serine/glycine biosynthetic pathway (Bricker et al.,2012). In contrast,

Pyk mutants exhibit no increase in serine production despite harboring very high levels

of 3-phosphoglycerate, which serves as the precursor for serine biosynthesis (YE *et al.*

2012). This difference is surprising because studies of PKM2 in cancer cells suggests

that decreased Pyk activity induces shunting of glycolytic intermediates into serine

production (YE *et al.* 2012).

One explanation for the dramatic phenotypic differences between *Pyk* and *Mpc1*

mutants stems from differences in lactate metabolism. Unlike *Pyk* mutants, *Mpc1*

mutants accumulate excess lactate. Thus, *Mpc1* mutants can generate two more ATPs

from each glucose molecule when compared with *Pyk* mutants, while also regenerating

ATP independent of mitochondrial activity. Future studies should test this model by

examining if *Mpc1;Ldh* double mutant exhibits metabolic and developmental

phenotypes that more closely mimic those observed in *Pyk* mutants.

In conclusion, we have here described a metabolomics profile of *Drosophila*

larvae lacking Pyk activity. While the resulting changes in central carbon metabolism

are largely expected, such an analysis of fly metabolism is essential as *Drosophila*

geneticists increasingly study the role of metabolism across a wide range of contexts.

Moreover, *Drosophila* metabolism is highly adaptable and compensates for metabolic

insult – as seen here by depletion of the anaplerotic amino acids proline, glutamate, and

glutamine. A careful exploration of these metabolic networks will be essential for

studying metabolism in the context of *Drosophila* growth, development, and models of

human disease.

541
542

ACKNOWLEDGEMENTS

Thanks to the Bloomington *Drosophila* Stock Center (NIH P40OD018537) for providing

fly stocks, the *Drosophila* Genomics Resource Center (NIH 2P40OD010949) for

genomic reagents, and Flybase (NIH 5U41HG000739). Metabolomics analysis

performed at the Metabolomics Core Facility at the University of Utah is supported by

spectrometry equipment obtained through NCRR Shared Instrumentation Grants

1S10OD016232-01, 1S10OD018210-01A1 and 1S10OD021505-01. Sequencing was

performed at Indiana University's Center for Genomics and Bioinformatics. All

computation was performed on Indiana University's Carbonate and Research Desktop

HPC clusters. We thank members of the Matt Hahn lab at IU for helpful guidance on the

phylogeny. J.M.T. is supported by the National Institute of General Medical Sciences of

the National Institutes of Health under a R35 Maximizing Investigators' Research Award

(MIRA; 1R35GM119557).

FIGURE LEGENDS

Figure 1. Sequence identity and phylogeny of Pyk homologs. (**A**) Heatmap of pairwise sequence identities extracted from the ensemble MSA. (**B**) Phylogenetic maximum likelihood gene tree. Ultrafast bootstrap support is shown on internal branches. The dark grey highlighted clades include the canonical *Drosophila melanogaster*, *Aedes aegypti*, and human/mouse orthologs, while the light grey region encompasses the divergent fly paralogs. The tree is rooted to *E. coli* homolog PykA as the outgroup. Species abbreviations: Aa, *A. aegypti*; Dm, *D. melanogaster*; Hs, *H. sapiens*; Mm, *M. musculus*; Ce, *C. elegans*; Sc, *S. cerevisiae*; At, *A. thaliana*; Ec, *E. coli.* See Supplemental Table 1 for a list of isoforms used in this analysis.

Figure 2. Deletions in the *Pyk* coding region induce larval lethality and growth defects. (A) An illustration of the *Drosophila Pyk* locus and surrounding genes, as well as the *Pyk* mutant alleles. The *Pyk^{23}* and *Pyk^{31}* alleles were generated using a CRISPR/Cas9-based strategy and induce small deletions that result in frameshifts. The *Pyk^{60}* and *Pyk^{61}* alleles represent deletions that remove the entire PYK coding region as well as portions of neighboring genes. See Figure S1 for analysis of *Pol3f* and *CG7069* gene expression in trans-heterozygous *Pyk* mutant backgrounds. (B) A histogram illustrating the effects of *Pyk* mutations on larval viability. Note that nearly all *Pyk$^{23/31}$* trans-heterozygous mutant animals die during the second larval instar – phenocopying *dERR* mutants. $P<0.001$ (C) *Pyk$^{23/31}$* mutants are significantly smaller than heterozygous controls. n=6 samples containing 25 mid-L2 larvae. **$P<0.01$. ***$P<0.001$. *P*-values calculated using a Mann Whitney test.

Figure 3. *Pyk* mutants exhibit decreased TAG levels. TAG levels were assayed

during L2 development in *Pyk* mutants and heterozygous controls. (A) *Pyk*$^{23/31}$ mutants

exhibit significantly decreased triglycerides (TAG) as compared with *Pyk*$^{21/+}$ controls.

Data are normalized to soluble protein. n=6 biological replicates containing 25 mid-L2

larvae per sample. **P<0.01. *P*-values calculated using a Mann Whitney test. (B,C) Nile

Red was used to stain lipid droplets in L2 fat bodies of both (B) *Pyk*$^{21/+}$ controls and (C)

Pyk$^{23/31}$ mutants. Scale bar in (B) applies to (C).

Figure 4. Targeted metabolomic analysis of *Pyk* mutant larvae. *Pyk*$^{23/31}$ mutant

larvae and *Pyk*$^{21/+}$ controls were analyzed using a targeted GC-MS approach. (A)

Differences in metabolite abundance between control and mutant samples are

represented as a volcano plot. Dashed vertical line represents a log2 fold change (FC)

of > 1.5. Dashed horizontal line represents p < 0.01. (B) A heat map illustrating the top

25 most significantly altered compounds in *Pyk*$^{23/31}$ mutants as compared with *Pyk*$^{21/+}$

controls. Both figures were generated using Metaboanalyst 5.0, as described in the

methods. Abbreviated compounds are 2/3-phosphoglycerate (2/3PG),

phosphoenolpyruvate (PEP), 2-hydroxyglutarate (2HG), 3-hydroxybutyrate (2HB), 5,10-

Methenyltetrahydrofolate (5,10-CH=THF), alanine (Ala), Asparagine (Asn), Glutamic

acid (Glu), Glutamine (Gln), 5'−Phosphoribosyl−N-formylglycinamide (FGAR), proline

(Pro), trehalose (Tre).

Figure 5. Targeted metabolomic analysis of *Pyk* mutant larvae. *Pyk*$^{60/61}$ mutant larvae and precise excision controls (*Pyk*^{Prec}) were analyzed using a targeted GC-MS approach. (A) Differences in metabolite abundance between control and mutant samples are represented as a volcano plot. Dashed vertical line represents a fold change (FC) of > 1.5. Dashed horizontal line represents *P*< 0.01. (B) A heat map illustrating the top 25 most significantly altered compounds in *Pyk*$^{60/61}$ mutants as compared with *Pyk*^{Prec} controls. Both figures were generated using Metaboanalyst 5.0, as described in the methods. Abbreviated compounds are 3-phosphoglycerate (3PG), 2-phosphoglycerate (2PG), phosphoenolpyruvate (PEP), 2-hydroxyglutarate (2HG), 3-hydroxybutyrate (2HB), alanine (Ala), Asparagine (Asn), Glutamic acid (Glu), Glutamine (Gln), 5'−Phosphoribosyl−N-formylglycinamide (FGAR), proline (Pro), trehalose (Tre).

Figure 6. A limited set of metabolites are significantly altered in both *Pyk* mutant backgrounds. (A) An illustration of glycolysis and the TCA cycle with metabolites that are significantly altered in both *Pyk* mutant backgrounds highlighted in dark red (up-regulated) and light blue (down-regulated). (B,C) Boxplots illustrating the change in abundance of metabolites that are significantly altered in both (B) *Pyk*$^{60/61}$ and (C) *Pyk*$^{23/31}$ mutant backgrounds. All boxplots were generated using Metaboanalyst 5.0, as described in the methods. Black dots represent individual samples, the horizontal bar in the middle represents the median and the yellow diamond represents the mean concentration. For all boxplots, the metabolite fold-change was >2 fold and *P*<0.01. Abbreviated compounds are 3-phosphoglycerate (3PG), 2-phosphoglycerate (2PG), phosphoenolpyruvate (PEP), 2-hydroxyglutarate (2HG).

Figure 7. RNA-seq analysis of $Pyk^{23/31}$ mutants as compared with $Pyk^{23/+}$ controls.

A volcano plot illustrating gene expression changes in $Pyk^{23/31}$ mutants as compared

with $Pyk^{23/+}$ controls. Black vertical lines indicate absolute log fold change (LFC) ≥ 1.

Black horizontal line represents adjusted P value ≤ 0.05. The top 15 most significantly

changed genes are labeled, with all significant genes (abs(LFC) ≥ 1 and adj p ≤ 0.05)

shaded darker.

Figure 8. Gene Ontology (GO) enrichment analysis of genes that are significantly

misregulated in $Pyk^{23/31}$ mutants. GOrilla was used to determine GO enrichment

among those genes that were either (A) significantly up-regulated or (B) significantly

down-regulated in $Pyk^{23/31}$ mutants when compared to the $Pyk^{23/+}$ control. (C)

Significantly up-regulated genes in the GO categories "lipid metabolic process" and

"proteolysis" are grouped by enzymatic function, with the increased fold-change in gene

expression represented by a color gradient.

SUPPLEMENTAL FIGURE LEGENDS

Figure S1. *Pyk* mRNA transcript levels are significantly reduced in *Pyk* mutant

larvae. Total RNA from stage w^{1118}; Pyk^{prec} control larvae and w^{1118}; $Pyk^{60/61}$ mutant

larvae were analyzed by northern blot hybridization to detect transcripts encoding Pyk,

CG18596, and Polr3F. Hybridization to detect *rp49* mRNA is included as a loading

control.

Figure S2. A comparison of sequence similarities between Pyk homologs. A

heatmap of pairwise sequence identities extracted from the ensemble MSA. Species

abbreviations: Aa, *A. aegypti; Dm*, *D. melanogaster*; Hs, *H. sapiens*; Mm, *M. musculus*;

Ce, *C. elegans*; Sc, *S. cerevisiae*; At, *A. thaliana*; Ec, *E. coli.* See Supplemental Table

1 for a list of isoforms used in this analysis.

Figure S3. A comparison of the metabolomic data from *Pyk*$^{23/31}$ mutant and *Pyk*$^{21/+}$

control samples using principal component (PC) analysis. Targeted metabolomics

data from Table S1 was analyzed using principal component analysis. Analysis was

conducted using Metaboanalyst 5.0.

Figure S4. A comparison of the metabolomic data from *Pyk*$^{60/61}$ mutant and *Pyk*^{prec}

control samples using principal component (PC) analysis. Targeted metabolomics

data from Table S2 was analyzed using principal component analysis. Analysis was

conducted using Metaboanalyst 5.0.

Figure S5. RNA-seq Principal Component Analysis and Correlations of

Experimental Replicates. (A) PCA biplot of principal component 1 (PC1) versus PC2

for the top 5,000 genes by variance, with replicates labeled. (B) Spearman correlation

plot of the top 5,000 genes by variance after filtering out genes with either low/no

expression.

SUPPLEMENTAL TABLE LEGENDS

Table S1. Protein isoform information for the Pyk orthologs used in Figure 1 and Figure
S2.

Table S2. Metabolomic analysis of $Pyk^{23/31}$ mutants as compared with $Pyk^{23/+}$ controls.

Table S3. Metabolomic analysis of imprecise excision Pyk mutants as compared with
precise excision controls.

Table S4. RNA-seq results comparing gene expression between Pyk23/31 mutants and

$Pyk^{23/+}$.

Table S5. GO analysis of significantly mis-regulated genes from the RNAseq data

represented in Table S4 using GOrilla.

Table S6. Analysis of RNA-seq data from Table S4 using the Preferred tissue

(modEncode RNA_seq) tool in PANGEA.

LITERATURE CITED

2023 UniProt: the Universal Protein Knowledgebase in 2023. Nucleic Acids Res 51**:** D523-d531.

Andrews, S., 2010 FastQC: a quality control tool for high throughput sequence data., pp.

Ayi, K., G. Min-Oo, L. Serghides, M. Crockett, M. Kirby-Allen *et al.*, 2008 Pyruvate kinase deficiency and malaria. N Engl J Med 358**:** 1805-1810.

Beebe, K., M. M. Robins, E. J. Hernandez, G. Lam, M. A. Horner *et al.*, 2020 Drosophila estrogen-related receptor directs a transcriptional switch that supports adult glycolysis and lipogenesis. Genes Dev 34**:** 701-714.

Bellen, H. J., R. W. Levis, G. Liao, Y. He, J. W. Carlson *et al.*, 2004 The BDGP gene disruption project: single transposon insertions associated with 40% of Drosophila genes. Genetics 167**:** 761-781.

Berardini, T. Z., L. Reiser, D. Li, Y. Mezheritsky, R. Muller *et al.*, 2015 The Arabidopsis information resource: Making and mining the "gold standard" annotated reference plant genome. Genesis 53**:** 474-485.

Bernal, A., and D. A. Kimbrell, 2000 Drosophila Thor participates in host immune defense and connects a translational regulator with innate immunity. Proc Natl Acad Sci U S A 97**:** 6019-6024.

Blighe, K., and A. Lun, 2022 PCAtools: Everything Principal Components Analysis. R package version 2.8.0, pp.

Bray, N. L., H. Pimentel, P. Melsted and L. Pachter, 2016 Near-optimal probabilistic RNA-seq quantification. Nat Biotechnol 34**:** 525-527.

Bricker, D. K., E. B. Taylor, J. C. Schell, T. Orsak, A. Boutron *et al.*, 2012 A mitochondrial pyruvate carrier required for pyruvate uptake in yeast, Drosophila, and humans. Science 337**:** 96-100.

Bursell, E., 1981 The role of proline in energy metabolism, pp. 135-154 in *Energy metabolism in insects*. Springer.

Caruso, P., B. J. Dunmore, K. Schlosser, S. Schoors, C. Dos Santos *et al.*, 2017 Identification of MicroRNA-124 as a Major Regulator of Enhanced Endothelial Cell Glycolysis in Pulmonary Arterial Hypertension via PTBP1 (Polypyrimidine Tract Binding Protein) and Pyruvate Kinase M2. Circulation 136**:** 2451-2467.

Cox, J. E., C. S. Thummel and J. M. Tennessen, 2017 Metabolomic Studies in Drosophila. Genetics 206**:** 1169-1185.

Cunningham, F., J. E. Allen, J. Allen, J. Alvarez-Jarreta, M. R. Amode *et al.*, 2022 Ensembl 2022. Nucleic Acids Res 50**:** D988-d995.

Davis, P., M. Zarowiecki, V. Arnaboldi, A. Becerra, S. Cain *et al.*, 2022 WormBase in 2022-data, processes, and tools for analyzing Caenorhabditis elegans. Genetics 220.

Dolan, L. M., M. Ryan and J. Moohan, 2002 Pyruvate kinase deficiency in pregnancy complicated by iron overload. Bjog 109**:** 844-846.

Drummond-Barbosa, D., and J. M. Tennessen, 2020 Reclaiming Warburg: using developmental biology to gain insight into human metabolic diseases. Development 147.

Eden, E., R. Navon, I. Steinfeld, D. Lipson and Z. Yakhini, 2009 GOrilla: a tool for
discovery and visualization of enriched GO terms in ranked gene lists. BMC
Bioinformatics 10**:** 48.
Engel, S. R., F. S. Dietrich, D. G. Fisk, G. Binkley, R. Balakrishnan *et al.*, 2014 The
reference genome sequence of Saccharomyces cerevisiae: then and now. G3
(Bethesda) 4**:** 389-398.
Ewels, P., M. Magnusson, S. Lundin and M. Käller, 2016 MultiQC: summarize analysis
results for multiple tools and samples in a single report. Bioinformatics 32**:** 3047-
3048.
Fattizzo, B., F. Cavallaro, A. Marcello, C. Vercellati and W. Barcellini, 2022 Pyruvate
Kinase Deficiency: Current Challenges and Future Prospects. J Blood Med 13**:**
461-471.
Garapati, P. V., J. Zhang, A. J. Rey and S. J. Marygold, 2019 Towards comprehensive
annotation of Drosophila melanogaster enzymes in FlyBase. Database (Oxford)
2019.
Grace, R. F., A. Zanella, E. J. Neufeld, D. H. Morton, S. Eber *et al.*, 2015 Erythrocyte
pyruvate kinase deficiency: 2015 status report. Am J Hematol 90**:** 825-830.
Gramates, L. S., J. Agapite, H. Attrill, B. R. Calvi, M. A. Crosby *et al.*, 2022a FlyBase: a
guided tour of highlighted features. Genetics 220.
Gramates, L. S., J. Agapite, H. Attrill, B. R. Calvi, M. A. Crosby *et al.*, 2022b FlyBase: a
guided tour of highlighted features. Genetics 220.
Grant, B. J., A. P. Rodrigues, K. M. ElSawy, J. A. McCammon and L. S. Caves, 2006
Bio3d: an R package for the comparative analysis of protein structures.
Bioinformatics 22**:** 2695-2696.
Gratz, S. J., A. M. Cummings, J. N. Nguyen, D. C. Hamm, L. K. Donohue *et al.*, 2013
Genome engineering of Drosophila with the CRISPR RNA-guided Cas9
nuclease. Genetics 194**:** 1029-1035.
Graveley, B. R., A. N. Brooks, J. W. Carlson, M. O. Duff, J. M. Landolin *et al.*, 2011 The
developmental transcriptome of Drosophila melanogaster. Nature 471**:** 473-479.
Grönke, S., A. Mildner, S. Fellert, N. Tennagels, S. Petry *et al.*, 2005 Brummer lipase is
an evolutionary conserved fat storage regulator in Drosophila. Cell Metab 1**:** 323-
330.
Hay, N., and N. Sonenberg, 2004 Upstream and downstream of mTOR. Genes Dev 18**:**
1926-1945.
Hoang, D. T., O. Chernomor, A. von Haeseler, B. Q. Minh and L. S. Vinh, 2018
UFBoot2: Improving the Ultrafast Bootstrap Approximation. Mol Biol Evol 35**:**
518-522.
Hsu, M. C., and W. C. Hung, 2018 Pyruvate kinase M2 fuels multiple aspects of cancer
cells: from cellular metabolism, transcriptional regulation to extracellular
signaling. Mol Cancer 17**:** 35.
Hu, Y., A. Comjean, H. Attrill, G. Antonazzo, J. Thurmond *et al.*, 2023 PANGEA: A New
Gene Set Enrichment Tool for Drosophila and Common Research Organisms.
bioRxiv.
Israelsen, W. J., and M. G. Vander Heiden, 2015 Pyruvate kinase: Function, regulation
and role in cancer. Semin Cell Dev Biol 43**:** 43-51.

Karim, F. D., and C. S. Thummel, 1991 Ecdysone coordinates the timing and amounts
of E74A and E74B transcription in Drosophila. Genes Dev 5: 1067-1079.
Katoh, K., and D. M. Standley, 2013 MAFFT multiple sequence alignment software
version 7: improvements in performance and usability. Mol Biol Evol 30: 772-780.
Kim, S. K., D. D. Tsao, G. S. B. Suh and I. Miguel-Aliaga, 2021 Discovering signaling
mechanisms governing metabolism and metabolic diseases with Drosophila. Cell
Metab 33: 1279-1292.
Kolde, R., 2019 pheatmap: Pretty Heatmaps – an implementation of heatmaps that
offers more control over dimensions and appearance. (Version 1.0.12), pp.
Li, H., G. Chawla, A. J. Hurlburt, M. C. Sterrett, O. Zaslaver et al., 2017 Drosophila
larvae synthesize the putative oncometabolite L-2-hydroxyglutarate during
normal developmental growth. Proc Natl Acad Sci U S A 114: 1353-1358.
Li, H., A. J. Hurlburt and J. M. Tennessen, 2018 A Drosophila model of combined D-2-
and L-2-hydroxyglutaric aciduria reveals a mechanism linking mitochondrial
citrate export with oncometabolite accumulation. Dis Model Mech 11.
Li, H., and J. M. Tennessen, 2017 Methods for studying the metabolic basis of
Drosophila development. Wiley Interdiscip Rev Dev Biol 6.
Li, H., and J. M. Tennessen, 2018 Preparation of Drosophila Larval Samples for Gas
Chromatography-Mass Spectrometry (GC-MS)-based Metabolomics. J Vis Exp.
Love, M. I., W. Huber and S. Anders, 2014 Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol 15: 550.
Luke, N., K. Hillier, H. Al-Samkari and R. F. Grace, 2023 Updates and advances in
pyruvate kinase deficiency. Trends Mol Med 29: 406-418.
Machado, P., L. Manco, C. Gomes, C. Mendes, N. Fernandes et al., 2012 Pyruvate
kinase deficiency in sub-Saharan Africa: identification of a highly frequent
missense mutation (G829A;Glu277Lys) and association with malaria. PLoS One
7: e47071.
Maddocks, O. D., C. R. Berkers, S. M. Mason, L. Zheng, K. Blyth et al., 2013 Serine
starvation induces stress and p53-dependent metabolic remodelling in cancer
cells. Nature 493: 542-546.
Mahmoudzadeh, N. H., A. J. Fitt, D. B. Schwab, W. E. Martenis, L. M. Nease et al.,
2020 The oncometabolite L-2-hydroxyglutarate is a common product of dipteran
larval development. Insect Biochem Mol Biol 127: 103493.
Min-Oo, G., M. Tam, M. M. Stevenson and P. Gros, 2007 Pyruvate kinase deficiency:
correlation between enzyme activity, extent of hemolytic anemia and protection
against malaria in independent mouse mutants. Blood Cells Mol Dis 39: 63-69.
Minh, B. Q., H. A. Schmidt, O. Chernomor, D. Schrempf, M. D. Woodhams et al., 2020
IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the
Genomic Era. Mol Biol Evol 37: 1530-1534.
Moretti, S., F. Armougom, I. M. Wallace, D. G. Higgins, C. V. Jongeneel et al., 2007 The
M-Coffee web server: a meta-method for computing multiple sequence
alignments by combining alternative alignment methods. Nucleic Acids Res 35:
W645-648.
Nemkov, T., J. A. Reisz, S. Gehrke, K. C. Hansen and A. D'Alessandro, 2019 High-
Throughput Metabolomics: Isocratic and Gradient Mass Spectrometry-Based
Methods. Methods Mol Biol 1978: 13-26.

Nguyen, L. T., H. A. Schmidt, A. von Haeseler and B. Q. Minh, 2015 IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol 32**:** 268-274.

O'Leary, N. A., M. W. Wright, J. R. Brister, S. Ciufo, D. Haddad *et al.*, 2016 Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. Nucleic Acids Res 44**:** D733-745.

Pang, Z., J. Chong, G. Zhou, D. A. de Lima Morais, L. Chang *et al.*, 2021 MetaboAnalyst 5.0: narrowing the gap between raw spectra and functional insights. Nucleic Acids Res 49**:** W388-w396.

Petchampai, N., C. Murillo-Solano, J. Isoe, J. C. Pizarro and P. Y. Scaraffia, 2019 Distinctive regulatory properties of pyruvate kinase 1 from Aedes aegypti mosquitoes. Insect Biochem Mol Biol 104**:** 82-90.

Rambaut, A., 2018 FigTree, pp.

Robertson, H. M., C. R. Preston, R. W. Phillis, D. M. Johnson-Schlitz, W. K. Benz *et al.*, 1988 A stable genomic source of P element transposase in Drosophila melanogaster. Genetics 118**:** 461-470.

Rust, K. J., and G. E. Collier, 1985 Localization of a dosage sensitive region for pyruvate kinase in Drosophila melanogaster. J Hered 76**:** 39-44.

Schormann, N., K. L. Hayden, P. Lee, S. Banerjee and D. Chattopadhyay, 2019 An overview of structure, function, and regulation of pyruvate kinases. Protein Sci 28**:** 1771-1784.

Sebo, Z. L., H. B. Lee, Y. Peng and Y. Guo, 2014 A simplified and efficient germline-specific CRISPR/Cas9 system for Drosophila genomic engineering. Fly (Austin) 8**:** 52-57.

Spannl, S., T. Buhl, I. Nellas, S. A. Zeidan, K. V. Iyer *et al.*, 2020 Glycolysis regulates Hedgehog signalling via the plasma membrane potential. Embo j 39**:** e101767.

Sullivan, A. A., and C. S. Thummel, 2003 Temporal profiles of nuclear receptor gene expression reveal coordinate transcriptional responses during Drosophila development. Mol Endocrinol 17**:** 2125-2137.

Teleman, A. A., Y. W. Chen and S. M. Cohen, 2005 4E-BP functions as a metabolic brake used under stress conditions but not during normal growth. Genes Dev 19**:** 1844-1848.

Tennessen, J. M., K. D. Baker, G. Lam, J. Evans and C. S. Thummel, 2011 The Drosophila estrogen-related receptor directs a metabolic switch that supports developmental growth. Cell Metab 13**:** 139-148.

Tennessen, J. M., W. E. Barry, J. Cox and C. S. Thummel, 2014 Methods for studying metabolism in Drosophila. Methods 68**:** 105-115.

Teulier, L., J. M. Weber, J. Crevier and C. A. Darveau, 2016 Proline as a fuel for insect flight: enhancing carbohydrate oxidation in hymenopterans. Proc Biol Sci 283.

Tixier, V., L. Bataillé, C. Etard, T. Jagla, M. Weger *et al.*, 2013 Glycolysis supports embryonic muscle growth by promoting myoblast fusion. Proc Natl Acad Sci U S A 110**:** 18982-18987.

Trapnell, C., B. A. Williams, G. Pertea, A. Mortazavi, G. Kwan *et al.*, 2010 Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. Nat Biotechnol 28**:** 511-515.

Traxler, L., J. R. Herdy, D. Stefanoni, S. Eichhorner, S. Pelucchi *et al.*, 2022 Warburg-
like metabolic transformation underlies neuronal degeneration in sporadic
Alzheimer's disease. Cell Metab 34**:** 1248-1263.e1246.
van Bruggen, R., C. Gualtieri, A. Iliescu, C. Louicharoen Cheepsunthorn, P.
Mungkalasut *et al.*, 2015 Modulation of Malaria Phenotypes by Pyruvate Kinase
(PKLR) Variants in a Thai Population. PLoS One 10**:** e0144555.
Volkenhoff, A., A. Weiler, M. Letzel, M. Stehling, C. Klämbt *et al.*, 2015 Glial Glycolysis
Is Essential for Neuronal Survival in Drosophila. Cell Metab 22**:** 437-447.
Wallace, I. M., O. O'Sullivan, D. G. Higgins and C. Notredame, 2006 M-Coffee:
combining multiple sequence alignment methods with T-Coffee. Nucleic Acids
Res 34**:** 1692-1699.
Wei, T., and V. Simko, 2021 R package 'corrplot': Visualization of a Correlation Matrix.
(Version 0.92), pp.
Wei, T., and S. V., 2021 R package 'corrplot': Visualization of a Correlation Matrix., pp.
Whelan, S., and N. Goldman, 2001 A general empirical model of protein evolution
derived from multiple protein families using a maximum-likelihood approach. Mol
Biol Evol 18**:** 691-699.
Wong, E. D., S. R. Miyasato, S. Aleksander, K. Karra, R. S. Nash *et al.*, 2023
Saccharomyces genome database update: server architecture, pan-genome
nomenclature, and external resources. Genetics 224.
Wu, C. L., C. C. Chang, J. K. Wu, M. H. Chiang, C. H. Yang *et al.*, 2018 Mushroom
body glycolysis is required for olfactory memory in Drosophila. Neurobiol Learn
Mem 150**:** 13-19.
Ye, J., A. Mancuso, X. Tong, P. S. Ward, J. Fan *et al.*, 2012 Pyruvate kinase M2
promotes de novo serine synthesis to sustain mTORC1 activity and cell
proliferation. Proc Natl Acad Sci U S A 109**:** 6904-6909.
Zhang, H., D. Wang, M. Li, L. Plecitá-Hlavatá, A. D'Alessandro *et al.*, 2017 Metabolic
and Proliferative State of Vascular Adventitial Fibroblasts in Pulmonary
Hypertension Is Regulated Through a MicroRNA-124/PTBP1 (Polypyrimidine
Tract Binding Protein 1)/Pyruvate Kinase Muscle Axis. Circulation 136**:** 2468-
2485.
897

Figure 1. Sequence identity and phylogeny of Pyk homologs. (**A**) Heatmap of pairwise sequence identities extracted from the ensemble MSA. (**B**) Phylogenetic maximum likelihood gene tree. Ultrafast bootstrap support is shown on internal branches. The dark grey highlighted clades include the canonical *Drosophila melanogaster*, *Aedes aegypti*, and human/mouse orthologs, while the light grey region encompasses the divergent fly paralogs. The tree is rooted to *E. coli* homolog PykA as the outgroup. Species abbreviations: Aa, *A. aegypti; Dm*, *D. melanogaster*; Hs, *H. sapiens*; Mm, *M. musculus*; Ce, *C. elegans*; Sc, *S. cerevisiae*; At, *A. thaliana*; Ec, *E. coli.* See Supplemental Table 1 for a list of isoforms used in this analysis.

Figure 2. Deletions in the *Pyk* coding region induce larval lethality and growth defects. (A) An illustration of the *Drosophila Pyk* locus and surrounding genes, as well as the *Pyk* mutant alleles. The *Pyk²³* and *Pyk³¹* alleles were generated using a CRISPR/Cas9-based strategy and induce small deletions that result in frameshifts. The *Pyk⁶⁰* and *Pyk⁶¹* alleles represent deletions that remove the entire PYK coding region as well as portions of neighboring genes. See Figure S1 for analysis of *Pol3f* and *CG7069* gene expression in trans-heterozygous *Pyk* mutant backgrounds. (B) A histogram illustrating the effects of *Pyk* mutations on larval viability. Note that nearly all *Pyk²³/³¹* trans-heterozygous mutant animals die during the second larval instar – phenocopying *dERR* mutants. $P<0.001$ (C) *Pyk²³/³¹* mutants are significantly smaller than heterozygous controls. n=6 samples containing 25 mid-L2 larvae. **$P<0.01$. ***$P<0.001$. P-values calculated using a Mann Whitney test.

Figure 3. *Pyk* **mutants exhibit decreased TAG levels.** TAG levels were assayed during L2 development in *Pyk* mutants and heterozygous controls. (A) *Pyk²³/³¹* mutants exhibit significantly decreased triglycerides (TAG) as compared with *Pyk²¹/⁺* controls. Data are normalized to soluble protein. n=6 biological replicates containing 25 mid-L2 larvae per sample. **P<0.01. P-values calculated using a Mann Whitney test. (B,C) Nile Red was used to stain lipid droplets in L2 fat bodies of both (B) *Pyk²¹/⁺* controls and (C) *Pyk²³/³¹* mutants. Scale bar in (B) applies to (C).

Figure 4. Targeted metabolomic analysis of *Pyk* mutant larvae. *Pyk²³/³¹* mutant larvae and *Pyk²¹/⁺* controls were analyzed using a targeted GC-MS approach. (A) Differences in metabolite abundance between control and mutant samples are represented as a volcano plot. Dashed vertical line represents a log2 fold change (FC) of > 1.5. Dashed horizontal line represents p < 0.01. (B) A heat map illustrating the top 25 most significantly altered compounds in *Pyk²³/³¹* mutants as compared with *Pyk²¹/⁺* controls. Both figures were generated using Metaboanalyst 5.0, as described in the methods. Abbreviated compounds are 2/3-phosphoglycerate (2/3PG), phosphoenolpyruvate (PEP), 2-hydroxyglutarate (2HG), 3-hydroxybutyrate (2HB), 5,10-Methenyltetrahydrofolate (5,10-CH=THF), alanine (Ala), Asparagine (Asn), Glutamic acid (Glu), Glutamine (Gln), 5'-Phosphoribosyl−N-formylglycinamide (FGAR), proline (Pro), trehalose (Tre).

Figure 5. Targeted metabolomic analysis of *Pyk* mutant larvae. *Pyk^60/61^* mutant larvae and precise excision controls (*Pyk^Prec^*) were analyzed using a targeted GC-MS approach. (A) Differences in metabolite abundance between control and mutant samples are represented as a volcano plot. Dashed vertical line represents a fold change (FC) of > 1.5. Dashed horizontal line represents *P*< 0.01. (B) A heat map illustrating the top 25 most significantly altered compounds in *Pyk^60/61^* mutants as compared with *Pyk^Prec^* controls. Both figures were generated using Metaboanalyst 5.0, as described in the methods. Abbreviated compounds are 3-phosphoglycerate (3PG), 2-phosphoglycerate (2PG), phosphoenolpyruvate (PEP), 2-hydroxyglutarate (2HG), 3-hydroxybutyrate (2HB), alanine (Ala), Asparagine (Asn), Glutamic acid (Glu), Glutamine (Gln), 5'−Phosphoribosyl−N-formylglycinamide (FGAR), proline (Pro), trehalose (Tre).

Figure 6. A limited set of metabolites are significantly altered in both *Pyk* mutant backgrounds. (A) An illustration of glycolysis and the TCA cycle with metabolites that are significantly altered in both *Pyk* mutant backgrounds highlighted in dark red (up-regulated) and light blue (down-regulated). (B,C) Boxplots illustrating the change in abundance of metabolites that are significantly altered in both (B) *Pyk²³/³¹* and (C) *Pyk²³/³¹* mutant backgrounds. All boxplots were generated using Metaboanalyst 5.0, as described in the methods. Black dots represent individual samples, the horizontal bar in the middle represents the median and the yellow diamond represents the mean concentration. For all boxplots, the metabolite fold-change was >2 fold and *P*>0.01. Abbreviated compounds are 3-phosphoglycerate (3PG), 2-phosphoglycerate (2PG), phosphoenolpyruvate (PEP), 2-hydroxyglutarate (2HG).

Figure 7. RNAseq analysis of *Pyk^23/31* mutants as compared with *Pyk^23/+* controls. A volcano plot illustrating gene expression changes in *Pyk^23/31* mutants as compared with *Pyk^23/+* controls. Vertical lines indicate log fold change (LFC). Horizontal lines represent adjusted *P* value. The top 15 most significantly changed genes are labeled.

A

Gene categories upregulated in *Pyk* mutants		
GO Category	**Enrichment (N, B, n, b)**	**P-value**
chitin metabolic process	3.45 (8151,101,421,18)	3.35E-06
lipid metabolic process	2.15 (8151,342,421,38)	5.51E-06
glucosamine-containing compound metabolic process	3.17 (8151,110,421,18)	1.16E-05
aminoglycan metabolic process	3.02 (8151,122,421,19)	1.39E-05
amino sugar metabolic process	3.11 (8151,112,421,18)	0.000015
cuticle development	2.66 (8151,160,421,22)	2.28E-05
fatty-acyl-CoA metabolic process	6.20 (8151,25,421,8)	2.35E-05
chitin-based cuticle development	2.64 (8151,154,421,21)	3.96E-05
fatty acid derivative metabolic process	5.74 (8151,27,421,8)	4.41E-05
lipid catabolic process	3.16 (8151,92,421,15)	0.000065
proteolysis	1.71 (8151,578,421,51)	9.53E-05
drug metabolic process	2.20 (8151,220,421,25)	0.000162
long-chain fatty-acyl-CoA metabolic process	8.07 (8151,12,421,5)	0.00021
regulation of chitin-based cuticle tanning	7.45 (8151,13,421,5)	0.000327
acyl-CoA metabolic process	4.19 (8151,37,421,8)	0.000486
thioester metabolic process	4.19 (8151,37,421,8)	0.000486
tyrosine metabolic process	8.60 (8151,9,421,4)	0.000718
central nervous system formation	8.60 (8151,9,421,4)	0.000718

B

Gene categories downregulated in *Pyk* mutants		
GO Category	**Enrichment (N, B, n, b)**	**P-value**
peptidyl-proline hydroxylation to 4-hydroxy-L-proline	27.70 (8151,11,107,4)	8.63E-06
4-hydroxyproline metabolic process	27.70 (8151,11,107,4)	8.63E-06
peptidyl-proline hydroxylation	27.70 (8151,11,107,4)	8.63E-06
protein hydroxylation	23.44 (8151,13,107,4)	1.83E-05
pyrimidine ribonucleoside monophosphate metabolic process	76.18 (8151,2,107,2)	1.71E-04
pyrimidine ribonucleoside monophosphate biosynthetic process	76.18 (8151,2,107,2)	1.71E-04
cellular amino acid metabolic process	5.04 (8151,121,107,8)	1.79E-04
response to organic substance	4.01 (8151,190,107,10)	1.85E-04
sensory perception of smell	8.86 (8151,43,107,5)	2.30E-04
peptidyl-proline modification	10.51 (8151,29,107,4)	5.18E-04
cellular modified amino acid metabolic process	5.86 (8151,78,107,6)	5.31E-04
alpha-amino acid metabolic process	5.57 (8151,82,107,6)	6.94E-04

Figure 8. Gene Ontology (GO) enrichment analysis of genes that are significantly misregulated in *Pyk$^{23/31}$* mutants. GOrilla was used to determine GO enrichment of biolog among those genes that were either (A) significantly up-regulated or (B) significantly down-regulated in *Pyk$^{23/31}$* mutants when compared to the *Pyk$^{23/+}$* control. (C) Significantly up-regulated genes in the GO categories "lipid metabolic process" and "proteolysis" are grouped by enzymatic function and the fold-change in gene expression is represented using a color gradient.