**Expanded diversity of hemoplasmas in rare and undersampled Neotropical bats**

Dmitriy V. Volokhov[1], Lauren R. Lock[2], Kristin E. Dyer[2], Isabella K. DeAnglis[3,4], Benjamin R.

Andrews[3], Cynthia J. Downs[3], M. Brock Fenton[5], Nancy B. Simmons[6], Daniel J. Becker[2*]

[1] Center for Biologics Evaluation and Research, Food and Drug Administration, Silver Spring,

MD, USA

[2] Department of Biology, University of Oklahoma, Norman, OK, USA

[3] Department of Environmental Biology, SUNY College of Environmental Science and Forestry,

Syracuse, NY, USA

[4] Department of Biological Sciences, University of Arkansas, Fayetteville, AR, USA

[5] Department of Biology, University of Western Ontario, London, Ontario, Canada

[6] Department of Mammalogy, Division of Vertebrate Zoology, American Museum of Natural

History, New York, NY, USA

*Corresponding author, [danbeck@ou.edu](mailto:danbeck@ou.edu)

**Keywords:** hemotropic *Mycoplasma*; Chiroptera; Belize; 16S rRNA gene; 23S rRNA gene

**Running head:** Novel Neotropical bat hemoplasmas

**Abstract**

Hemotropic mycoplasmas are emerging as a model system for studying bacterial pathogens in

bats, but taxonomic coverage of sampled host species remains poor. We leveraged a long-term

field program in Belize to uncover novel hemoplasma diversity in bats by analyzing samples

from species infrequently encountered in typical surveys. PCR targeting the partial 16S rRNA

and 23S rRNA genes and subsequent phylogenetic analyses identified two putative novel host

shifts of hemoplasmas, four entirely novel hemoplasma genotypes, and the first detections of

hemoplasmas in four bat species. Our analyses demonstrate that additional genetic diversity of

hemoplasmas in bats can be uncovered by targeting rare or otherwise undersampled host species.

**Main text**

Several decades of research have characterized the diversity of viruses in bats, for which some

species host viruses of notable human health concern [1]. Although bats remain relatively

understudied for other pathogens, the diversity of their bacterial pathogens has been of increasing

interest [2]. Hemotropic mycoplasmas (hemoplasmas) are facultative intracellular bacteria that are

emerging as a model group of bat pathogens [3]. Sampling globally has uncovered substantial

genetic diversity of bat hemoplasmas [4–6], including high similarity to human hemoplasmas that

may indicate zoonotic transmission [7,8]. However, most sampling has focused on relatively

common bat species, leaving taxonomic gaps in understanding how hemoplasma diversity is

distributed globally and which bats may harbor potentially zoonotic hemoplasmas [2,3]. Here, we

leveraged an ongoing field program from which we previously identified 24 new hemoplasma

genotypes in 23 bat species to test additional under- and unsampled species for these bacteria [4,9].

During April–May 2019, November 2021, and April–May 2022, we sampled a diverse

bat community in Orange Walk District, Belize [10]. Bats were captured using mist nets and harp

traps along flight paths and at roosts, held in individual bags until processing, and identified to

species based on morphology [11]. Blood volumes under 1% body mass were sampled by lancing

the propatagial vein using sterile needles and collected with heparinized capillary tubes. Blood

was preserved on Whatman FTA cards and held at room temperature until storage at -20C at the

University of Oklahoma. Field methods were approved by the Institutional Animal Care and Use

Committees of the American Museum of Natural History (AMNHIACUC-20190129 and -

20210614) and University of Oklahoma (2022-0197). Bat sampling was authorized by the Belize

Forest Department by permits FD/WL/1/19(06), FD/WL/1/19(09), and FD/WL/1/21(12).

We used Qiagen QIAamp DNA Investigator Kits to extract genomic DNA from 61 blood

samples from 17 bat species. Our prior studies and those of other groups had mostly minimally

surveyed 14 of these species [4,6,12], whereas *Lophostoma nicaraguae*, *Micronycteris microtis*, and

*Molossus alvarezi* remained wholly unsampled. To determine the presence of hemoplasmas, we

used PCR targeting the partial 16S rRNA gene with primers and procedures described previously

[4,9]. We also attempted to amplify the partial 23S rRNA gene for 16S rRNA–positive samples to

establish a broader library of this gene for hemoplasmas [13]. Amplicons were purified and directly

sequenced at Psomagen. We used NCBI BLASTn to identify related sequences, followed by

MUSCLE for sequence alignment and MrBayes for phylogenetic analysis (10,000,000

generations with a GTR+G+I model) using the NGPhylogeny.fr platform [14]. We delineated

hemoplasma genotypes based on analysis of the partial 16S rRNA gene (850–860 bp) and

clustering on the hemoplasma phylogeny, using previously established criteria for novelty [4,9].

We detected hemoplasmas via amplification of the 16S rRNA gene in 41% of samples

(95% CI = 30–54%; Table S1 and Figure 1), with additional amplification of the 23S rRNA gene

for six bats (GenBank: OQ456384–86, OQ456392, OR055987–88). Of the 11 bat species that

tested positive, four had identical genotypes to those we earlier described in Belize [4]: *Myotis*

*elegans* and the MYE genotype, *Carollia sowelli* and genotypes CS1 and CS2, *Glossophaga*

*mutica* (formerly *G. soricina*) and the GLS genotype, and *Pteronotus fulvus* and the PPM1

genotype. All three *Phyllostomus discolor* had sequences 100% identical to partial sequences

from *P. discolor* in Brazil [6], which we classify as the PD1 genotype. Finding the same genotype

at the extremes of the *P. discolor* range (sites 5,000 km apart) and in two subspecies (*P. discolor*

*discolor* and *verrucosus*) suggests the ancestral spread of this pathogen across the Americas.

The remaining six bat species were infected with novel genotypes in that host (i.e., host

shifts or a new host-specific genotype) or represented the first hemoplasma detections for that

species. For the former, we detected two genotypes in *Dermanura phaeotis* that were previously

identified as specific to *Carollia sowelli* (CS1) and *Sturnira parvidens* (SP1, group B); the later

genotype was also previously detected once in *Artibeus lituratus* [4], and we recently identified the

group A variant in *Pteronotus mesoamericanus* [15]. Whereas the former four bat species belong to

the family Phyllostomidae, *Dermanura* and *Carollia* belong to different subfamilies in that clade,

suggesting putative inter-generic host shifts likely restricted by broader evolutionary history [4].

Novel hemoplasmas were detected in *Eptesicus furinalis*, *Molossus alvarezi*, *Lophostoma*

*nicaraguae*, *Gardnerycteris keenani*, and *Noctilio leporinus*, with the latter four representing the

first hemoplasma detections in these species and the latter three representing entirely novel

hemoplasma genotypes. Although we previously characterized two genotypes in *Eptesicus*

*furinalis* (EF1 and EF2, with the former also identified here) [4], we identified a third, novel

genotype (EF3) in a single *E. furinalis*, with 99.4% identity to EF1. A single *Molossus alvarezi*

tested positive for the MR1 genotype (99.8% identity), previously identified only in the closely

related *M. nigricans* in Belize [4]. We identified the novel NL1 genotype in *Noctilio leporinus*, one

of two members of the family Noctilionidae; this genotype shared only 96.1% identity to the

MYE genotype found in *Myotis elegans*, a distantly related vespertilionid [4]. We identified the

novel LN1 genotype in a single *Lophostoma nicaraguae*, which shared 99.2% identity to the LE

genotype previously identified in the closely related *L. evotis*[4] and 99.1% identity to the PD1

genotype in *P. discolor*[6]; all three bats are members of the Phyllostominae subfamily. Lastly, we

detected a novel genotype (GK1) in a single additional phyllostomine (*Gardnerycteris keenani*),

but GK1 was most related to the GLS genotype (97.6%) earlier found in *Glossophaga mutica* [4].

Our analyses demonstrate additional genetic diversity of hemoplasmas in bats uncovered

by targeting less-sampled species. We encourage additional analyses of hemoplasmas in other

unsampled bat taxa and geographies [2,3]. Further, longitudinal studies are needed to understand

how these hemoplasmas circulate in bats, transmission risk to other species, and impacts on bats.

**Acknowledgements**

This work was supported by the National Geographic Society (NGS-55503R-19), SUNY-ESF

Honors Program, Explorer's Club, and Research Corporation for Science Advancement (RCSA).

This work was conducted as part of Subaward No. 28365, part of a USDA Non-Assistance

Cooperative Agreement with RCSA Federal 341Award No. 58-3022-0-005. We thank Mark

Howells and staff of the Lamanai Field Research Center for assistance with field logistics and

permits as well as the many colleagues who helped net bats during 2019, 2021, and 2022

research in Belize. Lastly, we also thank Konstantin Chumakov for laboratory support.

## 110    **Conflicts of interest**

111    The authors declare no conflicts of interest.

 **Figure 1.** Consensus Bayesian phylogeny of 16S rRNA hemoplasma sequences from this study

113 (highlighted in bold; see Table S1 for genotype assignments) and reference sequences from bats

114 and other host species. Nodes are colored by posterior probability (nodes with less than 50%

115 support are not shown), and sequences from bats are colored by bat family. The inset shows

116 hemoplasma prevalence per bat species sampled here (*n* = 17) and associated 95% confidence

117 intervals (Wilson score interval). Photograph of *Noctilio leporinus* by Sherri and Brock Fenton.

**References**

Letko *et al*. Bat-borne virus diversity, spillover and emergence. *Nat Rev Microbiol* 2020; **18**: 461–471.

Szentivanyi *et al*. Trends in bacterial pathogens of bats: global distribution and knowledge gaps. *Transbound Emerg Dis* 2023; **2023**.

Millán *et al*. Worldwide occurrence of haemoplasmas in wildlife: insights into the patterns of infection, transmission, pathology and zoonotic potential. *Transbound Emerg Dis* 2020.

Becker *et al.* Ecological and evolutionary drivers of haemoplasma infection and bacterial genotype sharing in a Neotropical bat community. *Mol Ecol* 2020; **29**: 1534–1549.

Wang *et al.* High prevalence and genetic diversity of hemoplasmas in bats and bat ectoparasites from China. *One Health* 2023; **16**: 100498.

Ikeda *et al.* Molecular detection and genotype diversity of hemoplasmas in non-hematophagous bats and associated ectoparasites sampled in peri-urban areas from Brazil. *Acta Trop* 2022; **225**: 106203.

Millán *et al*. Widespread infection with hemotropic mycoplasmas in bats in Spain, including a hemoplasma closely related to 'Candidatus Mycoplasma hemohominis'. *Comp Immunol Microbiol Infect Dis* 2015; **39**: 9–12.

Descloux *et al.* Flying fox haemolytic fever, description of a new zoonosis caused by 'Candidatus Mycoplasma haemohominis'. *Clin Infect Dis* 2020.

Volokhov *et al*. Novel hemotropic mycoplasmas are widespread and genetically diverse in vampire bats. *Epidemiology & Infection* 2017.

Herrera *et al*. Disassembly of fragmented bat communities in Orange Walk District, Belize. *Acta Chiropterologica*. 2018; **20**: 147.

Reid. A field guide to the mammals of Central America and Southeast México: Oxford University Press. *New York* 1997; **334**.

Ikeda *et al.* Evidence and molecular characterization of *Bartonella* spp. and hemoplasmas in neotropical bats in Brazil. *Epidemiol Infect* 2017; **145**: 2038–2052.

Volokhov *et al.* Novel hemotrophic mycoplasma identified in naturally infected California sea lions (*Zalophus californianus*). *Vet Microbiol* 2011; **149**: 262–268.

Lemoine *et al.* NGPhylogeny.fr: new generation phylogenetic services for non-specialists. *Nucleic Acids Res* 2019; **47**: W260–W265.

DeAnglis *et al.* Bat cellular immunity varies by year and dietary habit in an increasingly fragmented landscape. *bioRxiv*. 2023.