Environmental variability can promote parasite diversity within hosts and

transmission among hosts

Madeline Jarvis-Cross[1], Martin Krkosek[1]

[1] The University of Toronto, Department of Ecology and Evolutionary Biology

25 Willcocks Street, Toronto, ON, M5S 3B2

Corresponding author: Madeline Jarvis-Cross madeline.jarvis.cross@mail.utoronto.ca

10 1.0.0. Abstract

While the mechanisms that govern disease emergence and spread among hosts are

increasingly well-described, the mechanisms that promote parasite diversity within-

hosts, affecting host outcomes and spillover potential, have been comparatively

understudied. Furthermore, while attention has been paid to the effects of increasing

temperatures on disease systems, the effects of environmental variability have been left

underexplored, despite rising climatic variability and internal temperature variability in a

prolific reservoir for disease. To investigate the impacts of environmental variability on

parasite diversity within-hosts, we analyzed a model of within-host population dynamics

wherein two parasites indirectly compete through the host's immune response. We

simulated the model under constant, demographically stochastic, environmentally

stochastic, and demographically and environmentally stochastic conditions, and

analysed the viability and longevity of non-equilibrium parasite co-occurrence. We found

that environmental stochasticity increased the viability and longevity of parasite co-

occurrence, suggesting that thermal variability arising from climatic change and as a

physiological trait may promote parasite diversity within ectotherms and help explain

bats' propensity to support diverse communities of parasites. Further, we found that

under certain conditions, the transmissibility of co-occurring parasites can surpass the

transmissibility of single parasites, suggesting that thermal variability may increase the

transmissibility of co-occurring parasites.

2.0.0. Introduction

In recent years, it has become clear that climate and land-use change can facilitate

novel species interactions and increase contact rates between wild populations and

humans, resulting in devastating epidemics and enabling spillover events and disease

emergence among wild and human hosts alike (Daszak et al., 2000, 2001; Patz et al.,

2004; R. Plowright et al., 2020; R. K. Plowright et al., 2017). However, despite the

consequential interplay between within- and among-host processes, the mechanisms

that promote parasite diversity within hosts and create kindling for events observed

among hosts have been understudied. In particular, while stabilizing mechanisms like

niche differentiation have been employed to explain the maintenance of parasite

diversity within hosts, the contributions of equalizing mechanisms, which decrease

average fitness differences between competitors, have yet to be examined (Bashey,

2015; Chesson, 2000; Chesson & Huntly, 1997; Karvonen et al., 2006; Rynkiewicz et

al., 2015).

An intermediate level of disturbance may promote species diversity by forestalling

the competitive advantages allowed by low levels of disturbance, and the extinction

risks brought on by high levels of disturbance (Chesson, 1994; Connell, 1978).

Expanding on the work of Connell (1978) and Chesson (1994), Roxburgh et al. (2004)

posited that because different species respond to disturbances in different ways, there

are myriad ways in which diversity may be generated, thereby interpreting the

intermediate disturbance hypothesis not as a mechanism unto itself, but as a collection

of mechanisms through which intermediately sized or timed disturbances in resource

availability or environmental conditions may promote diversity (Roxburgh et al., 2004).

Though the applicability of the intermediate disturbance hypothesis has been debated,

positive and negative effects of environmental variability on diversity have been well-

documented across systems (Laprise & Dodson, 1994; Righetti et al., 2019; Shurin et

al., 2010; Vega-Rodriguez et al., 2015).

When considering the effects of environmental variability on parasite diversity

within hosts, there arise a number of ways to conceptualize environmental variability.

First, one may consider rising environmental variability as a feature of climate change

(Stocker et al., 2013). While much attention has been paid to the effects of rising

temperatures on disease systems, far less has been invested in exploring the effects of

rising temperature variability, despite evidence of its outsized impacts on species

outcomes (Dobson & Carper, 1992; Lafferty, 2009; Lawson et al., 2015; Vasseur et al.,

2014). In ectothermic hosts, increasingly erratic external conditions will affect the

internal conditions of the host, thereby also affecting any parasites the host may

harbour (Cizauskas et al., 2017). A relationship between thermal variability and within-

host parasite diversity may be of particular interest when considered alongside the long-

debated relationship between climate change and widespread disease-mediated

amphibian declines (Cohen et al., 2019; Daszak et al., 1999, 2003; Lips et al., 2008;

Pounds et al., 1999; Rohr et al., 2008; Rohr & Raffel, 2010).

Second, one may consider environmental variability as a physiological trait of

hosts. In particular, bats (order *Chiroptera*) have long been acknowledged as prolific

reservoirs for disease and have oft been implicated in zoonotic spillover events (Brook

& Dobson, 2015; Calisher et al., 2006; Irving et al., 2021). Bats are also unique among

mammals in their ability to vary their internal temperature according to their activity level

(e.g. torpor, flight) (Fumagalli et al., 2021). It has been theorized that these variations in

internal temperature may aid bats' immune responses, allowing them to carry

microparasites without suffering outsized morbidity (Fumagalli et al., 2021; Luo et al.,

2021). Extending on that idea, it seems pertinent to question how internal temperature

variation may contribute to bats' propensity to support large communities of parasites.

We used a model to investigate the impacts of environmental variability on the

viability and maintenance of parasite diversity within hosts. We began by modifying

Antia et al.'s (1994) model of within-host population dynamics to include two parasite

strains or species that indirectly interact via the host's immune system. We analyzed

deterministic, demographically stochastic, environmentally stochastic, and

demographically-and-environmentally stochastic implementations of the model. Within

the environmentally stochastic implementations of the model, we varied the magnitude

and frequency of thermal disturbance within the system. Between and among the

deterministic and stochastic simulations of the model, we compared the ability of the

secondary parasite to invade a system containing the primary parasite, and the duration

of time for which the primary and secondary parasites co-occurred. Further, we

compared the transmission potentials of co-occurring parasites with the transmission

potential of a single parasite. We found that compared to a constant environment, a

variable environment tended to increase the ability of the secondary parasite to invade

the system, facilitated longer periods of co-occurrence between the primary and

secondary parasites, and increased the transmissibility of co-occurring parasites. These

results (1) suggest that climatic changes may have significant effects on parasite

diversity within ectothermic hosts, (2) provide insight on a host's ability to support large

communities of parasites, and propose that variability within-hosts may influence

transmission dynamics among-hosts.

3.0.0. Methods

3.1.0. Model Description

3.1.1. Within-Host Population Dynamics Model

We began with a model of within host population dynamics based on Antia et al.,

(1994) that represents the interaction between an infecting parasite and its host's

immune system as analogous to the interaction between a prey item and its predator.

The model assumes that the replication rate of the host's immune system is proportional

to parasite density at low parasite densities, and saturates at high parasite densities,

allowing the host's immune system to either extirpate or succumb to the infection (Antia

et al., 1994). We adapted the model to include a second parasite that interacts indirectly

with the primary parasite through the host's immune response (Eqns. 1-3, Table 1).

Parameters values were not chosen to describe a specific system, but to replicate a

dynamical behaviour, as described above and shown in Figure 3. Given that an infinite

number of parameter value combinations could have produced this behaviour, we opted

to maintain the initial conditions and select parameter values proposed by Antia et al.

(1994). However, we increased the rate at which parasites are destroyed by the host's

immune system and decreased the maximum replication rates of the primary and

secondary parasites to favour extirpation, and eliminate host death as an infection

outcome (Table 1). We also conducted a sensitivity analysis to determine how

variations in (1) the value of k and (2) the shape of the relationship between parasite

replication rate and the temperature of the system would affect our results, and found

the model to be sufficiently robust (see pages 2, 4-8 of supplement for complete

results).

(1) $P_i(t+1) = (1 + r_i - (1 - e^{-kI(t)}))P_i(t)$

(2) $P_j(t+1) = (1 + r_j - (1 - e^{-kI(t)}))P_j(t)$

(3) $I(t+1) = (1 + \rho \left(\dfrac{P_i(t)}{P_i(t) + \phi} + \dfrac{P_j(t)}{P_j(t) + \phi} \right))I(t)$

Table 1: Parameter definitions and values.

Symbol	Parameter	Value used in Antia et al. (1994)	Value Used
P_i and P_j	Parasite abundance	1*	1*
I	Host immune cell abundance	1*	1*
r_i and r_j	Replication rates of parasites	0.1 – 10.0	0.1 – 2.0†
k	Rate at which parasites are destroyed by host's immune system	10^{-3}	10^{-2}
ρ	Maximum replication rate of host's immune system	1.0	1.0
ϕ	Parasite density at which replication rate of host's immune system is half its maximum	10^3	10^3
D	Lethal within-host parasite density	10^9	10^9

128 * Initial conditions

129 † Dependent on the temperature of system in environmentally stochastic simulations
130

To compare the impacts of different types of stochasticity on parasite diversity, we wrote

deterministic (D), demographically stochastic (S_D), environmentally stochastic (S_E), and

demographically-and-environmentally stochastic (S_DE) implementations of the model.

Within the demographically stochastic implementation of the model, we wrote the

model's growth terms as Poisson processes, and mortality term as a binomial process

(Eqns. 4-6).

(4) $P_i(t + 1) = P_i(t) + B_{P_i}(t) - D_{P_i}(t)$

(5) $P_j(t + 1) = P_j(t) + B_{P_j}(t) - D_{P_j}(t)$

(6) $I(t + 1) = I(t) + B_I(t)$

In Eqns. (4) and (5), $B_{P_i}(t)$ and $B_{P_j}(t)$ are Poisson random variables such that

$B_{P_i}(t) \sim Poisson(1, \lambda_i)$, where $\lambda_i = r_i \cdot P_i(t)$, and $B_{P_j}(t) \sim Poisson(1, \lambda_j)$, where $\lambda_j = r_j \cdot$

$P_j(t)$. In Eqn. (6), $B_I(t)$ is a Poisson random variable such that $I_B(t) \sim Poisson(1, \lambda)$,

where $\lambda = \rho \left(\frac{P_i(t)}{P_i(t) + \phi} + \frac{P_j(t)}{P_j(t) + \phi} \right) I(t)$. In Eqns. (4) and (5), $P_{i,D}(t)$ and $P_{j,D}(t)$ are binomial

random variables such that $D_{P_i}(t) \sim Binomial(1, P_i(t), p)$, where $p = 1 - e^{-kI(t)}$, and

$D_{P_j}(t) \sim Binomial(1, P_j(t), p)$. To write the environmentally stochastic implementation of

the model, we generated a stochastic temperature series and allowed the replication

rates of each parasite to vary in response to the temperature of the system (Eqns. 7-9).

(7) $P_i(t + 1) = (1 + r_i(T(t + 1)) - (1 - e^{-kI(t)}))P_i(t)$

(8) $P_j(t + 1) = (1 + r_j(T(t + 1)) - (1 - e^{-kI(t)}))P_j(t)$

(9) $I(t + 1) = (1 + \rho \left(\frac{P_i(t)}{P_i(t) + \phi} + \frac{P_j(t)}{P_j(t) + \phi} \right))I(t)$

We used first-order autocorrelated Gaussian noise to generate stochastic temperature

series. We began with a value of 0, and generated the next value, $T(t + 1)$, by

multiplying the previous value by the coefficient of autocorrelation (AC = 0.25), and

summing the resulting value with a value drawn from a normal random variable with a

mean of zero. We varied the standard deviation of the normal random variable, as

described below. We then added the average temperature of the system to each value

in the series. Resulting values outside the thermal range of the system were re-drawn.

We then converted the stochastic temperature series into series of replication rates

using various parameterizations of the Sharpe-Schoolfield sub-model, as detailed below

(Eqns. 10.1, 10.2; Table 2, Table S2).

(10.1) $r_i(T(t+1)) = r_{i,0}e^{\frac{-E_{r_i}}{k}(\frac{1}{T(t+1)}-\frac{1}{T_0})} \cdot \left(1 + e^{\frac{E^L_{r_i}}{k}\left(\frac{1}{T^L_{r_i}}-\frac{1}{T(t+1)}\right)} + e^{\frac{E^H_{r_i}}{k}\left(\frac{1}{T^H_{r_i}}-\frac{1}{T(t+1)}\right)}\right)^{-1}$

(10.2) $r_j(T(t+1)) = r_{j,0}e^{\frac{-E_{r_j}}{k}(\frac{1}{T(t+1)}-\frac{1}{T_0})} \cdot \left(1 + e^{\frac{E^L_{r_j}}{k}\left(\frac{1}{T^L_{r_j}}-\frac{1}{T(t+1)}\right)} + e^{\frac{E^H_{r_j}}{k}\left(\frac{1}{T^H_{r_j}}-\frac{1}{T(t+1)}\right)}\right)^{-1}$

Table 2: Parameter definitions. See Table A1 for parameter values.

Symbol	Unit	Parameter
T	K	Temperature of the system at time t
r_0		Baseline values of r_i or r_j at a reference temperature T_0
T_0	K	Reference temperature (set arbitrarily such that $T^L \ll T_0 \ll T^L$)
E_r	eV	Activation energy of r_i or r_j
E^L_r, E^H_r	eV	Low- and high-temperature inactivation energies of r_i or r_j
T^L_r, T^H_r	K	Low- and high-temperature inactivation thresholds for r_i or r_j
k		Boltzmann's constant

165   **3.1.2. Sharpe-Schoolfield Sub-Model**

166   The Metabolic Theory of Ecology (MTE) predicts how metabolic rates control

ecological processes across levels of organization, and explains how metabolic rates

vary with body size and temperature (Brown et al., 2004). Building off the MTE, the

Boltzmann-Arrhenius (BA) relation describes metabolism as a function of temperature.

The Sharpe-Schoolfield model characterizes the performance of a trait as a function of

temperature through a thermal performance curve (TPC) (Dell et al., 2011; Molnár et al.,

2017; Schoolfield et al., 1981). When applied to disease systems, the Sharpe-

Schoolfield model has been shown to describe relationships between temperature and

species traits, and has proven useful in investigating the relationship between rising

temperatures and disease transmission (Kirk et al., 2018; Molnár et al., 2013).

We used various parameterizations of the Sharpe-Schoolfield model to generate

paired thermal performance curves relating the temperature of the system to the

replication rates of two parasites (r_i and r_j) (Figure 1). We generated four paired thermal

performance curves, representing four thermal scenarios: (1) the identical responses

scenario, (2) the offset responses scenario, (3) the P_i-advantage scenario, and (4) the

P_j-advantage scenario (Table S1). Each thermal scenario describes a different

relationship between the temperature of the system and parasite replication rates. The

identical responses scenario describes r_i and r_j as responding identically to the

temperature of the system (Fig. 1). Under the offset responses scenario, r_i and r_j peak

on opposite sides of the average temperature of the system (Fig. 1). To develop the P_i-

advantage scenario, we decreased the average temperature of the system to give the

primary parasite, P_i, a thermal advantage (Fig. 1). We considered two versions, (I) and

(II), of the P_j-advantage scenario, each of which gave the secondary parasite, P_j, a

thermal advantage. To develop (I), we increased the average temperature of the system

to give the secondary parasite a quantitatively comparable thermal advantage (Fig. 1).

To develop (II), we replicated the P_i-advantage scenario, and assigned the TPC with the

lower optimum to P_j, to give the secondary parasite the same thermal advantage given

to the primary parasite and isolate the effects of asymmetry (Fig. S9). Given that (I) and

(II) produced qualitatively comparable results, we will only discuss (I) in the main text

(see supplement for complete results). Within each thermal scenario, the average

temperature of the system was chosen to maintain quantitatively comparable (1)

parasite replication rates between thermal scenarios and (2) differences between

replication rates at the average temperature of the system.

199
200 Figure 1: Sharpe-Schoolfield thermal performance curves relating the temperature of the system to the replication
201 rates (r_i and r_j) of two parasites (P_i and P_j). Yellow lines show the relationship between the temperature of the system
202 and the replication rate (r_i) of P_i. Green lines show the relationship between the temperature of the system and the
203 replication rate (r_j) of P_j. Pink lines denote the average temperature of the system. Each panel (A-E) shows a
204 different thermal scenarios.

205
3.2.0. Model Simulations

Within each thermal scenario, we simulated the deterministic model (D), and

each of the three stochastic models (S$_D$, S$_E$, S$_{DE}$) for fifty days with a discretization time

step of 0.1, which was sufficient to capture the full course of parasite infection and

clearance across all simulations. We began each set of simulations with the primary

parasite and the host's immune system in the system, at initial abundances of 1 as the

initial condition. We then considered all cases for the introduction of the secondary

parasite from its introduction at the onset of infection, to its introduction on the fourty-

ninth day of infection.

For each environmentally stochastic model (S_E, S_{DE}) simulation, we generated a

new temperature series and two corresponding series of replication rates. When

determining the ways in which new temperatures would be selected, we considered two

axes of thermal disturbance: magnitude and frequency. To vary magnitude of

disturbance, we varied the standard deviation of the normal random variable in the

temperature autocorrelation function, starting from zero (no thermal disturbance) and

increasing by two until the effect of magnitude on co-occurrence was fully characterized

(SD = 30), resulting in fifteen discrete magnitudes of disturbance. To vary frequency of

disturbance, we re-assigned the temperature of the system every 10, 9, 8, 7, 6, 5, 4, 3,

and 2 days, once per day, and 2, 5, and 10 times per day, resulting in thirteen discrete

frequencies of disturbance. We conducted 1000 simulations per stochastic model and

treatment (Fig. 2). Two hundred such simulations are shown in Figure 3.

227
228 Figure 2: Schematic detailing simulation experiment. We considered four thermal scenarios, and four models
229 (deterministic, demographically stochastic, environmentally stochastic, and demographically and environmentally
230 stochastic). When simulating the environmentally stochastic models, we considered 195 disturbance regimes. When
231 simulating each of the four models, we considered 50 days on which the secondary parasite could be introduced. We
232 conducted 1000 simulations per stochastic model and treatment.

233

3.3.0. Co-Occurrence Analyses

We quantified longevities of co-occurrence by counting the number of time steps

for which primary and secondary parasite population abundances were equal to or

greater than one within each simulated timeseries. We summarised these results as

distributions for each stochastic model and treatment. Using the S_D model as a

baseline, we investigated the effects of environmental variability on the viability and

maintenance of parasite diversity by comparing the ability of the secondary parasite to

invade a single-parasite system, and the duration of time for which the primary and

secondary parasites co-occurred across stochastic treatments. Additionally, we

analysed (1) the proportion of stochastic simulations per model and treatment that

resulted in longer periods of co-occurrence relative to the deterministic model to

quantify rarity, and (2) the competitive outcomes of each treatment to gain insight on

underlying mechanisms.

3.4.0. Transmissibility Analyses

To explore the impacts of co-infection on transmission potential, we compared

the lifetime replicative outputs of co-occurring parasites with those of a single parasite in

constant and variable environments. As per Antia et al., (1994), we determined the total

transmission (q_i, q_j, q, where q is the total transmission of the single parasite) of a

parasite (P_i, P_j, P, where P is the single parasite) from an infected host by taking the

integral of the rate of transmission (u) of each parasite from the host over the duration

of infection (Eqns. 11.1-11.3). For simplicity, we set the rate of transmission to one.

(11.1) $\quad q_i = \int_0^\infty uP_i(t)dt$

(11.2) $\quad q_j = \int_0^\infty uP_j(t)dt$

(11.3) $\quad q = \int_0^\infty uP(t)dt$

262 To compare constant and variable environments, we analysed multi-parasite and single-

263 parasite versions of the S_D and S_{DE} models.

265 4.0.0. Results

266   Analytically, the secondary parasite could invade $(P_{j(t+1)} > P_{j(t)})$ when $I(t) <$

267 $\frac{-\ln(1-r_j)}{k}$. Our parameterization of the S_D model allowed the secondary parasite to

268 invade when introduced before the tenth to twelfth day of infection, resulting in 8.4 to 9.2

269 days of co-occurrence, depending on thermal scenario (Fig. 4, Supplementary Datafile

270 1-1:10).

271
272 Figure 4: Longevity of co-occurrence as per the deterministic model. The plot shows the number of time steps for
273 which P_i and P_j have population abundances greater than one, by introduction day, as per the deterministic model.
274 The black line denotes the first day on which the introduction of the secondary parasite, P_j, does not result in any co-
275 occurrence between the two parasites.

276 4.1.0. Co-occurrence Analysis

Simulation outputs show that compared to a deterministic baseline, the addition

of demographic stochasticity has little effect on co-occurrence (Supplementary Datafile

2-1:10), while the addition of thermal variability increases the ability of the secondary

parasite to invade the system and facilitates longer periods of co-occurrence between

the primary and secondary parasites. While longevities of co-occurrence are maximised

under the P_j-advantage scenario, the difference between deterministic and

environmentally stochastic implementations is not consistently maximised by any one

thermal scenario. Additionally, while larger and more frequent disturbances tend to

promote longer periods of co-occurrence, increasingly large disturbances produce a

less steep effect than do increasingly frequent disturbances. These findings are

common among thermal scenarios.

When both parasites were introduced simultaneously, S_D model simulations

resulted in periods of co-occurrence that were, on average, 4.5% to 3.1% shorter (0.4 to

0.3 days) than deterministically simulated periods of co-occurrence (Figs. 5, S10).

Under the same conditions, simulations of the environmentally stochastic models (S_E,

S_{DE}) resulted in periods of co-occurrence that were up to 46.2% (S_E) and 36.0% (S_{DE})

longer (3.9 (S_E) and 3.0 (S_{DE}) days) than deterministically simulated periods of co-

occurrence (Figs. 5, S10). Across all 195 disturbance regimes, simulations of the S_E

model resulted in mean periods of co-occurrence that were between 2.7% shorter (0.2

days) and 46.2% longer (3.9 days) than their deterministic counterparts (Figs. 5, S10).

The identical responses scenario promoted the shortest and longest mean periods of

co-occurrence (Figs. 5-6, S10-S11). Simulations of the S_{DE} model resulted in mean

periods of co-occurrence that were between 6.8% shorter (0.6 days) and 36.0% longer

(3.0 days) than deterministically simulated periods of co-occurrence (Figs. 5, S10).

Again, the identical responses scenario promoted the shortest and longest mean

periods of co-occurrence (Figs. 5-6, S10-S11). Across thermal scenarios, mean periods

of co-occurrence were shortest when disturbances were large (standard deviation of the

normal random variable in the temperature autocorrelation function = 22 to 30) and

frequent (once every 7 to 9 days) (Figs. 6, S11). Mean periods of co-occurrence were

longest when disturbances were large (SD = 24 to 30) and infrequent (five to ten times

per day) (Figs. 6, S11).

When isolating the effect of size of disturbance (frequency of disturbance fixed at

an intermediate value (once per day)), simulations of the S_E model resulted in mean

periods of co-occurrence that were between 2.2% and 37.8% longer (0.2 and 3.2 days)

than deterministically simulated periods of co-occurrence (Figs. 5-7, S10-S12, S14).

The P_i-advantage and identical responses scenarios promoted the shortest and longest

mean periods of co-occurrence, respectively (Figs. 5-7, S10-S12, S14). Simulations of

the S_{DE} model resulted in mean periods of co-occurrence that were between 2.9%

shorter (0.3 days) and 32.3% longer (2.7 days) than deterministically simulated periods

of co-occurrence (Figs. 5-7, S10-S12, S14). The P_j-advantage and identical responses

scenarios promoted the shortest and longest mean periods of co-occurrence,

respectively (Figs. 5-7, S10-S12, S14). Periods of co-occurrence were shortest when

disturbances were small (SD = 2), and longest when disturbances were large (SD = 24

to 30) (Figs. 5-7, S10-S12, S14).

When isolating the effect of frequency of disturbance (magnitude of disturbance

fixed at an intermediate value (SD = 10)), simulations of the S_E model resulted in mean

periods of co-occurrence that were between 2.3% shorter (0.2 days) and 35.2% longer

(3.0 days) than deterministically simulated periods of co-occurrence (Figs. 5-7, S10-

S12, S15). The identical responses scenario promoted the shortest and longest mean

periods of co-occurrence (Figs. 5-7, S10-S12, S15). Simulations of the S_{DE} model

resulted in mean periods of co-occurrence that were between -6.0% shorter (0.5 days)

and 25.3% longer (2.1 days) than deterministically simulated periods of co-occurrence

(Figs. 5-7, S10-S12, S15). Again, the identical responses scenario promoted the

shortest and longest mean periods of co-occurrence (Figs. 5-7, S10-S12, S15). Periods

of co-occurrence were shortest when disturbances were infrequent (once every 7 to 8

days), and longest when disturbances were frequent (five to ten times per day) (Fig. 5-

7, S10-S12, S15).

More generally, as disturbances increased in size, periods of co-occurrence

increased in length towards a saturation point (standard deviation of the normal random

variable in the temperature autocorrelation function ≈ 24-30, depending on thermal

scenario) (Figs. 7A, S12, S16, Supplementary Datafiles 3-1:5, 4-1:5). Additionally, when

disturbances were infrequent (once every seven to ten days), periods of co-occurrence

were similar in length to deterministically simulated periods of co-occurrence. Per the

deterministic model, P_i and P_j can only co-occur for a maximum of 9.2 days. Thus,

systems subjected to infrequent disturbances were functionally similar to systems

subjected to a constant environment, making the outcome of a simulation reliant on the

initial conditions of the system. Interestingly, when disturbances were intermediately

frequent (once per day to once every few days), mean periods of co-occurrence were

shorter than deterministically simulated periods of co-occurrence (Fig. 7B, S12, S19).

When disturbances were frequent (twice to ten times per day), mean periods of co-

occurrence were longer than deterministically simulated periods of co-occurrence.

Finally, within a given disturbance regime, distributions of longevities of co-occurrence

were normally distributed (Figs. 7, S12-S13, S20).

Figure 5: The mean percent difference in longevity of co-occurrence relative to deterministically simulated periods of
co-occurrence, when both parasites are introduced at the same time. Each point represents the mean percent
difference in longevity of co-occurrence relative to deterministically simulated periods of co-occurrence across 1000
stochastic simulations, as per a given environmental disturbance regime. As such, there are 195 points above each
"S_E Model"/"S_{DE} Model" label.

356
357 Figure 6: Fully factorial representations of the impacts of varyingly sized and timed disturbances on the longevity of
358 co-occurrence. The plot shows the mean number of time steps (10 time steps/day) for which P_i and P_j have
359 population abundances greater than one as per the S_{DE} model. The circled numbers denote the number of time steps
360 for which P_i and P_j can co-occur, per the deterministic model.

361
Figure 7: Distributions of outcomes across select disturbance regimes when both parasites are introduced at the
same time. "S_D" refers to longevities of co-occurrence as per the S_D model. "S_{DE}" refers to longevities of co-
occurrence as per the S_{DE} model. Panel A shows the effects of increasingly large disturbances (frequency of
disturbance fixed (once per day)). Panel B shows the effects of increasingly frequent disturbances (magnitude of
disturbance fixed (SD = 10)). Black vertical lines represent deterministic reference points. This figure shows
longevities of co-occurrence under the identical responses scenario. See Fig. S12 for full results.

368
4.2.0. Rarity and Competitive Outcomes

Analyzing the proportion of environmentally stochastic simulations in which

longevities of co-occurrence exceeded deterministically simulated periods of co-

occurrence revealed that the effect of environmental variability on the viability and

maintenance of parasite co-occurrence was not rare. When the secondary parasite was

introduced before or on the invasion threshold (10th or 12th day of infection, depending

on thermal scenario), 29%-71% (S$_E$) and 46%-58% (S$_{DE}$) of environmentally stochastic

simulations resulted in longer periods of co-occurrence relative to deterministic

simulations. When the secondary parasite was introduced after the invasion threshold,

the likelihood that an environmentally stochastic simulation resulted in some period of

co-occurrence sharply declined. When the secondary parasite was introduced one to

twelve days after the invasion threshold, 1%-31% (S$_E$) and 1%-46% (S$_{DE}$) of

environmentally stochastic simulations resulted in some period of co-occurrence. Thus,

environmental variability extended the invasion threshold by 50%-75% (S$_E$) and 100%-

120% (S$_{DE}$), depending on thermal scenario (Fig. 8).

384
Figure 8: Proportion of environmentally stochastic simulations (SD = 10, frequency of disturbance = once per day)
resulting in longer periods of co-occurrence relative to deterministic simulations by the day on which the secondary
parasite was introduced. Vertical lines show the first day on which the introduction of the secondary parasite does not
result in any co-occurrence between the two parasites, as per the deterministic model (D).

Analyzing competitive outcomes at intermediate levels of disturbance (SD = 10,

frequency of disturbance = once per day) revealed that under the offset responses

scenario, when P_i and P_j were introduced at the same time, P_j outlasted P_i 65.5% (S$_E$)

and 63.4% (S_{DE}) of the time (Fig. S20-S21). Additionally, under the P_i-advantage

scenario, when P_i and P_j were introduced at the same time, P_i and P_j were equally likely

to outlast one another (P_j outlasts P_i 49.9% (S_E) and 51.0% (S_{DE}) of the time), but under

the P_j-advantage, P_j was much more likely to outlast P_i (P_j outlasts P_i 76.6% (S_E) and

70.4% (S_{DE}) of the time) (Fig. S19-S20). Across thermal scenarios, when the secondary

parasite, P_j, was introduced in the middle of the infection period, P_i was more likely to

outlast P_j. When P_j was introduced late in the infection period, P_j was more likely to

outlast P_i (Fig. S20-S21). Regardless of when P_j was introduced, simulations resulting

in especially long periods of co-occurrence are equally likely to show either parasite

outlasting the other (Fig. S20-S21).

4.3.0. Transmissibility

Comparing the transmission potentials of co-occurring and single parasites

revealed that under certain conditions, the transmissibility of co-occurring parasites can

exceed the transmissibility of single parasites. Across thermal scenarios, simulations of

the multi-parasite S_D model produced lifetime replicative outputs (LROs) (median LRO

of P_i ranged between 1.49E+04 and 9.12E+05; median LRO of P_j ranged between

2.59E+04 and 6.09E+05) that were one to two orders of magnitude lower than those

produced by simulations of the single-parasite S_D model (median LRO of P ranged

between 3.62E+06 and 5.13E+07) (Supplementary Datafiles 5-1:16). Similarly, under

the first three thermal scenarios, simulations of the multi-parasite S_{DE} model produced

LROs (median LRO of P_i ranged between 1.02E+04 and 1.05E+06; median LRO of P_j

ranged between 8.36E+03 and 3.00E+05) that were one to two orders of magnitude

lower than those produced by simulations of the single-parasite S_{DE} model (median

LRO of P ranged between 6.45E+04 and 3.41E+07). However, under the P_j-advantage

scenario and among 62 of 195 (32%) disturbance regimes, the average transmission

potential of the co-occurring secondary parasite, P_j, exceeded that of a single parasite

(Fig. 9B). Among these disturbance regimes, the transmission potential of P_j exceeded

that of a single parasite by an average of 2% to 135%. The transmission potential of P_j

was maximised when disturbances were large (SD = 20) and infrequent (once every

eight days) (Fig. 9A, Supplementary Datafiles 5-8). The transmission potentials of co-

occurring parasites peaked when disturbances were infrequent (once every ten days) to

intermediately frequent (once every six days) (Fig. S22). While large and frequent

disturbances maximally increased longevities of co-occurrence, such infections did not

result in particularly high transmission potentials, indicating that especially long periods

of co-occurrence may be facilitated by lower lifetime replicative outputs that do not

trigger a large immune response, and thus allow for more persistent infections.

428
Figure 9: The impacts of environmental variability on transmission potential under the P_j-advantage scenario. Panel A
shows the total transmission of parasites by model (S$_D$, S$_{DE}$) and structure (multi-parasite, single-parasite), when
disturbances are large (SD = 20) and infrequent (once every eight days). Panel B shows a fully factorial
representation of the impacts of thermal disturbance on the transmission potential of P_j, relative to the equivalent
single-parasite treatment. The black line indicates disturbance regimes under which the average transmission
potential of P_j was equal to that of a single parasite. Colours warmer than denoted by the black line indicate
disturbance regimes under which the average transmission potential of P_j exceeded that of a single parasite; colours
cooler than denoted by the black line indicate the opposite.

5.0.0. Discussion

Despite increasing rates of disease emergence, the mechanisms that promote

parasite diversity within hosts and create kindling for emergences and epidemics among

hosts have been understudied (Baker et al., 2022; Jones et al., 2008; Taylor et al.,

2001). Here, we investigated the potential for environmental variability, both as a feature

of climate change and as a physiological trait unique to a prolific reservoir for disease to

affect the viability and maintenance of within-host parasite diversity. Our work provides

theoretical evidence that relative to a constant environment, a variable environment can

increase the ability of a secondary parasite to invade a single-parasite system, promote

longer periods of co-occurrence between parasites, and increase the transmissibility of

co-occurring parasites. In more detail, thermal disturbances of various sizes and

frequencies allowed a secondary parasite to invade a single-parasite system up to

twelve days after the invasion threshold, extending the invasion threshold by 100% to

120% (S_{DE} model, Fig. 8). Large and frequent thermal disturbances enabled parasites

to co-occur for up to 37.8% longer, relative to a constant environment (S_{DE} model, Figs.

5, S10).

Across thermal scenarios and disturbance regimes, the relationship between

disturbance size and longevity of co-occurrence was initially positive and eventually

saturating (Fig. S16). We determined that increasing magnitudes of disturbance

produced a saturating response rather than the expected parabolic response due to the

structure of the Sharpe-Schoolfield sub-model, which cannot generate negative

replication rates. When we introduced negative replication rates, larger disturbances

began to have negative effects on parasite populations, resulting in the expected

parabolic response, and implying that intermediate magnitudes of disturbance

maximised the viability and longevity of parasite co-occurrence (Figs. S17-S18).

The relationship between climate change and disease-mediated amphibian

declines has long been debated. While some argue that climatic changes may impede

immune function or result in performance gaps between parasites and their hosts, and

thus make amphibians more susceptible to infection, others argue that insufficient

evidence should preclude a causal link (Cohen et al., 2019, 2019; Daszak et al., 1999,

2003; Lips et al., 2008; Pounds et al., 1999; Rohr et al., 2008). While much of the

literature proposes mean-climate hypotheses, some evidence suggests that intensifying

El Niño events are increasing thermal variability, which may in turn affect amphibians'

susceptibility to or ability to defend against infection (Rohr & Raffel, 2010). Similarly,

emerging evidence suggests that in addition to rising mean temperatures, daily

temperature fluctuations may affect the incubation period of malaria parasites and

related vital rates, altering malaria transmission patterns (Paaijmans et al., 2010, 2013).

Congruently, our findings provide theoretical evidence that thermal variability promotes

the viability and maintenance of within-host parasite diversity, and thus suggest that as

the climate continues to change, attention should be paid to shifts in microparasite

communities and their impacts on ectothermic hosts. In particular, well-documented

climate-driven increases in disease incidence among a wide variety of marine

organisms may constitute an important starting point for future work (Bates et al., 2009;

Bruno et al., 2007; Carilli et al., 2012; Drinkwater et al., 2003; Harvell et al., 2001; Tracy

et al., 2019).

Across thermal scenarios and disturbance regimes, intermediate and high

frequencies of disturbance impeded and promoted the viability and longevity of parasite

co-occurrence, respectively (Figs. 6, S11, S19). Given the transient nature of infection

specified by the underlying model, intermediate frequencies of disturbance (once every

seven days, for example) resulted in the temperature of the system being disturbed

towards the end of the infection period, when parasite populations were small and

especially susceptible to unfavourable conditions. Conversely, high frequencies of

disturbance resulted in the temperature of the system being disturbed throughout the

duration of the infection period, and as such, neither favourable nor unfavorable

conditions lasted long enough for one parasite to gain a decisive advantage over and

exclude the other. While it has been theorized that variations in internal temperature

may aid bats' immune responses, allowing them to carry microparasites without

suffering outsized morbidity (Fumagalli et al., 2021; Luo et al., 2021), these findings

suggest that variations in internal temperature may also facilitate the co-occurrence of

parasites within bats, and thus may help explain bats' propensity to support diverse

communities of parasites. Further, these results, combined with the regularity with which

bats are implicated in spillover events and epidemics, suggest that the observed effects

of environmental variability on parasite diversity within hosts will scale up to have effects

on between-host processes like transmission, as discussed below.

With regards to competitive outcomes, under the offset responses scenario, P_j

often outlasted P_i, while under the P_i-advantage scenario, P_i and P_j were equally likely

to outlast each other (Fig. S20-S21). Under the P_j-advantage scenario, P_j was much

more likely to outlast P_i (Fig. S20-S21). These outcomes are consequences of the

asymmetrical structure of the thermal performance curves (Sinclair et al., 2016). As an

example, within the offset responses scenario, colder than average temperatures

resulted in a slightly higher replication rate for P_i and a slightly lower replication rate for

P_j, while warmer than average temperatures resulted in a much lower replication rate for

P_i and a slightly higher replication rate for P_j. In short, P_i was more disadvantaged by

unfavourable conditions than was P_j. Further, regardless of when P_j was introduced,

simulations resulting in especially long periods of co-occurrence were equally likely to

show either parasite outlasting the other. Taken together, these results suggest that to

predict the competitive outcome of a multiple infection, one may need to consider the

interplay between how well adapted a parasite is to its host, priority effects, and the

responses of biological rates to environmental factors. While one might expect a

specialist or well-adapted parasite (here, a thermally advantaged parasite) to exploit its

host more efficiently than, and thus outcompete, a generalist or novel parasite, the

results of this work suggest that environmental variability can even the scales, and allow

a generalist or novel parasite to establish and persist, as predicted by the intermediate

disturbance hypothesis (Chesson, 1994; Connell, 1978; Roxburgh et al., 2004).

Similarly, while one might additionally expect a colonizing parasite (here, the primary

parasite) to benefit from an initial lack of competition, the results of this work again

suggest that environmental variability and disturbance may contribute to the structuring

of communities by allowing latecomers to establish and persist more often and for

longer than they otherwise would (Jiang & Patel, 2008; Munguia et al., 2010; Symons &

Arnott, 2014).

While the metabolic theory of ecology posits that certain biological rates (e.g.

metabolic rate, replication/growth rate, etc.) will exhibit an asymmetrical and unimodal

response to temperature, other vital rates may exhibit symmetrical or bimodal

responses to temperature (Dell et al., 2014; Molnár et al., 2017; Sinclair et al., 2016).

This work suggests that the shapes of thermal performance curves can meaningfully

affect the nature and outcome of an infection. As anthropogenic changes alter host-

parasite interaction networks, understanding how evolutionary and environmental

factors combine to affect interactions between parasites will be essential in predicting

and managing spillover and emergence in wildlife and human populations (Paull et al.,

2012; Pedersen & Fenton, 2007).

From an array of parasites circulating among ectotherms to Hendra virus,

emerging coronaviruses, and other parasites carried by bats, developing an

understanding of how the impacts of environmental variability on parasite diversity

within hosts may scale to affect among-host processes and events, including

transmission, spillover, and emergence will be essential to the maintenance of wildlife,

human, and ecosystem health (Allocati et al., 2016; Daszak et al., 2003; Halpin et al.,

2000; Johnson, 2006). While increased within-host parasite diversity provides more

strains or species of a parasite or parasites an opportunity to transmit, within-host

competition may result in decreased parasite loads, thus decreasing the transmission

probability of any one strain or species, and altering expected transmission patterns

(Johnson et al., 2013; Pedersen & Fenton, 2007). Similarly, our findings provide

theoretical evidence that when environmental variations are very large and very

frequent, within-host competition may decrease the transmission rates of coinfecting

parasites, thereby decreasing transmission probability. However, our findings also

suggest that when thermal disturbances are variably sized and infrequent, the lifetime

replicative outputs of coinfecting parasites can exceed those of single parasites,

indicating that the effects of environmental variability may not be limited to the

promotion of within-host parasite diversity, but may also promote transmission between

hosts (Fig. 9B). Additionally, we found that while large and frequent disturbances may

maximally increase the amount of time for which two parasites co-occur, such infections

do not result in particularly high transmission potentials, indicating that especially long

periods of co-occurrence may be facilitated by lower replicative outputs that do not

trigger a large immune response, and thus allow for more persistent infections.

Conversely, disturbances of intermediate size and low to intermediate frequency may

constitute an anomalous region within which environmental variation can increase the

amount of time for which two parasites co-occur, while allowing parasites to achieve

high replicative outputs, resulting in longer and more transmissible infections (Fig. 9B).

While both Antia et al.'s (1994) model and our modified model should be broadly

applicable to an array of within-host multi-parasite systems, including multi-strain or

cross-species SARS-CoV-2 and influenza infections, both models are relatively simple

as to remain generalizable, and thus cannot represent chronic infections, systems within

which parasites exhibit more complex interactions with their host or their host's immune

system (ex. malaria, HIV), nor systems within which parasites directly interact with one

another (as is often observed in trematode coinfection and in hosts coinfected with

multiple strains of the bacterium *Pseudomonas aeruginosal*) (Al Khatib et al., 2020;

Bashey, 2015; Kim et al., 2021; Saenz et al., 2010). As such, it may be interesting to

validate the theoretical insights of this work in an empirical system. It may be of

particular interest to explore the impacts of environmental variability on within-host

parasite diversity in amphibians, mosquitos, ectothermic marine organisms, or, given

the risk they pose to public health, bats (Bates et al., 2009; Bruno et al., 2007; Carilli et

al., 2012; Drinkwater et al., 2003; Paaijmans et al., 2010; Rohr & Raffel, 2010; Tracy et

al., 2019).

This work is also limited by the scope with which thermal dependency was

considered. Given the thermal sensitivity of immune function in ectotherms and

endotherms, it will be worthwhile to consider the thermal dependency of immune

processes like response time and replication rate in tandem with the thermal

dependency of parasite replication rates (Butler et al., 2013; Wright & Cooper, 1981).

Similarly, using nonlinear averaging to assess the response of a nonlinear process to

variability is contentious. While many theoretical works and some empirical studies

support the use of nonlinear averaging and find evidence for Jensen's inequality, borne

out as disproportionate effects of thermal variability on performance, many empirical

studies have contested an implicit assumption of nonlinear averaging: that traits should

respond instantaneously to change, without acclimation (Bernhardt et al., 2018; Estay et

al., 2011; Krichel et al., 2023; Paaijmans et al., 2010; Sinclair et al., 2016). Thermal

acclimation through phenotypic plasticity alters thermal performance curves, and as

such, produces a more complex relationship between biological rates and temperature

that may not be well-characterized by nonlinear averaging (Altman et al., 2016; Schulte

et al., 2011; Sckrabulis et al., 2022). With regards to host-parasite interactions, the

temperature variability hypothesis posits that owing to their small relative size, parasites

should acclimate more quickly than their hosts, though the speed at which various

parasites may acclimate remains unclear (Kunze et al., 2022; Raffel et al., 2013; Rohr

et al., 2013). Interestingly, some empirical studies of the impacts of environmental

variability on host-parasite interactions have considered thermal acclimation, yielding a

range of conclusions including decreased pathogen prevalence, reduced infection

prevalence, and decreased resistance in hosts (Dallas & Drake, 2016; Duncan et al.,

2011; Raffel et al., 2013).

This work provides theoretical evidence that through climate change and as a

physiological trait, environmental variability may promote parasite diversity within hosts

and transmission among hosts, and thus create kindling for the spillover events, disease

emergences, and epidemics we observe among wild and human hosts alike.

Anthropogenic changes have allowed the next pandemic to leave the hypothetical and

become a looming spectre. Camus wrote that plagues, despite their commonality, find

us unprepared (Camus, Albert, 1991). Improving our understanding of how

environmental variability will impact parasite diversity within- and between-hosts will

allow us to expand our understanding of the mechanisms that can facilitate disease

spread and emergence, which will allow us to make robust predictions, and develop and

implement mitigation strategies as we move into an increasingly uncertain future. In

brief, we will be found *less* unprepared.

6.0.0. Acknowledgements

This work was supported by The Emerging and Pandemic Infections Consortium

Doctoral Award (MJ-C), an NSERC Discovery Grant (MK) and Canada Research Chair

(MK).

7.0.0. References

Al Khatib, H. A., Benslimane, F. M., Elbashir, I. E., Coyle, P. V., Al Maslamani, M. A.,
Al-Khal, A., Al Thani, A. A., & Yassine, H. M. (2020). Within-Host Diversity of
SARS-CoV-2 in COVID-19 Patients With Variable Disease Severities. *Frontiers
in Cellular and Infection Microbiology*, *10*, 575613.
https://doi.org/10.3389/fcimb.2020.575613

Allocati, N., Petrucci, A. G., Di Giovanni, P., Masulli, M., Di Ilio, C., & De Laurenzi, V.
(2016). Bat–man disease transmission: Zoonotic pathogens from wildlife
reservoirs to human populations. *Cell Death Discovery*, *2*(1), 16048.
https://doi.org/10.1038/cddiscovery.2016.48

Altman, K. A., Paull, S. H., Johnson, P. T. J., Golembieski, M. N., Stephens, J. P.,
LaFonte, B. E., & Raffel, T. R. (2016). Host and parasite thermal acclimation
responses depend on the stage of infection. *Journal of Animal Ecology*, *85*(4),
1014–1024. https://doi.org/10.1111/1365-2656.12510

Antia, R., Levin, B. R., & May, R. M. (1994). Within-Host Population Dynamics and the
Evolution and Maintenance of Microparasite Virulence. *The American Naturalist*,
144(3), 457–472. https://doi.org/10.1086/285686

Baker, R. E., Mahmud, A. S., Miller, I. F., Rajeev, M., Rasambainarivo, F., Rice, B. L.,
Takahashi, S., Tatem, A. J., Wagner, C. E., Wang, L.-F., Wesolowski, A., &
Metcalf, C. J. E. (2022). Infectious disease in an era of global change. *Nature
Reviews Microbiology*, *20*(4), 193–205. https://doi.org/10.1038/s41579-021-
00639-z

Bashey, F. (2015). Within-host competitive interactions as a mechanism for the

maintenance of parasite diversity. *Philosophical Transactions of the Royal*

Society B: Biological Sciences, *370*(1675), 20140301.

https://doi.org/10.1098/rstb.2014.0301

Bates, A., Hilton, B., & Harley, C. (2009). Effects of temperature, season and locality on

wasting disease in the keystone predatory sea star Pisaster ochraceus. *Diseases*

of Aquatic Organisms, *86*, 245–251. https://doi.org/10.3354/dao02125

Bernhardt, J. R., Sunday, J. M., Thompson, P. L., & O'Connor, M. I. (2018). Nonlinear

averaging of thermal experience predicts population growth rates in a thermally

variable environment. *Proceedings of the Royal Society B: Biological Sciences*,

285(1886), 20181076. https://doi.org/10.1098/rspb.2018.1076

Brook, C. E., & Dobson, A. P. (2015). Bats as 'special' reservoirs for emerging zoonotic

pathogens. *Trends in Microbiology*, *23*(3), 172–180.

https://doi.org/10.1016/j.tim.2014.12.004

Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., & West, G. B. (2004). Toward a

metabolic theory of ecology. *Ecology*, *85*(7), 1771–1789.

https://doi.org/10.1890/03-9000

Bruno, J. F., Selig, E. R., Casey, K. S., Page, C. A., Willis, B. L., Harvell, C. D.,

Sweatman, H., & Melendy, A. M. (2007). Thermal Stress and Coral Cover as

Drivers of Coral Disease Outbreaks. *PLoS Biology*, *5*(6), e124.

https://doi.org/10.1371/journal.pbio.0050124

Butler, M. W., Stahlschmidt, Z. R., Ardia, D. R., Davies, S., Davis, J., Guillette, L. J.,

Johnson, N., McCormick, S. D., McGraw, K. J., & DeNardo, D. F. (2013).

Thermal Sensitivity of Immune Function: Evidence against a Generalist-

Specialist Trade-Off among Endothermic and Ectothermic Vertebrates. *The*

American Naturalist, *181*(6), 761–774. https://doi.org/10.1086/670191

Calisher, C. H., Childs, J. E., Field, H. E., Holmes, K. V., & Schountz, T. (2006). Bats:

Important Reservoir Hosts of Emerging Viruses. *Clinical Microbiology Reviews*,

19(3), 531–545. https://doi.org/10.1128/CMR.00017-06

Camus, A. (1991). *The Plague* (T. Stuart Gilbert.). Vintage. (Original work published

1948).

Carilli, J., Donner, S. D., & Hartmann, A. C. (2012). Historical Temperature Variability

Affects Coral Response to Heat Stress. *PLoS ONE*, *7*(3), e34418.

https://doi.org/10.1371/journal.pone.0034418

Chesson, P. (1994). Multispecies Competition in Variable Environments. *Theoretical*

Population Biology, *45*(3), 227–276. https://doi.org/10.1006/tpbi.1994.1013

Chesson, P. (2000). Mechanisms of Maintenance of Species Diversity. *Annual Review*

of Ecology and Systematics, *31*(1), 343–366.

https://doi.org/10.1146/annurev.ecolsys.31.1.343

Chesson, P., & Huntly, N. (1997). The Roles of Harsh and Fluctuating Conditions in the

Dynamics of Ecological Communities. *The American Naturalist*, *150*(5), 519–553.

https://doi.org/10.1086/286080

Cizauskas, C. A., Carlson, C. J., Burgio, K. R., Clements, C. F., Dougherty, E. R.,

Harris, N. C., & Phillips, A. J. (2017). Parasite vulnerability to climate change: An

evidence-based functional trait approach. *Royal Society Open Science*, *4*(1),

160535. https://doi.org/10.1098/rsos.160535

Cohen, J. M., Civitello, D. J., Venesky, M. D., McMahon, T. A., & Rohr, J. R. (2019). An

interaction between climate change and infectious disease drove widespread

amphibian declines. *Global Change Biology*, *25*(3), 927–937.

https://doi.org/10.1111/gcb.14489

Connell, J. H. (1978). Diversity in Tropical Rain Forests and Coral Reefs: High diversity

of trees and corals is maintained only in a nonequilibrium state. *Science*,

199(4335), 1302–1310. https://doi.org/10.1126/science.199.4335.1302

Dallas, T., & Drake, J. M. (2016). Fluctuating temperatures alter environmental

pathogen transmission in a *Daphnia* –pathogen system. *Ecology and Evolution*,

6(21), 7931–7938. https://doi.org/10.1002/ece3.2539

Daszak, P., Berger, L., Cunningham, A. A., Hyatt, A. D., Green, D. E., & Speare, R.

(1999). Emerging Infectious Diseases and Amphibian Population Declines.

Emerging Infectious Diseases, *5*(6), 735–748.

https://doi.org/10.3201/eid0506.990601

Daszak, P., Cunningham, A. A., & Hyatt, A. D. (2000). Emerging Infectious Diseases of

Wildlife—Threats to Biodiversity and Human Health. *Science*, *287*(5452), 443–

449. https://doi.org/10.1126/science.287.5452.443

Daszak, P., Cunningham, A. A., & Hyatt, A. D. (2001). Anthropogenic environmental

change and the emergence of infectious diseases in wildlife. *Acta Tropica*, *78*(2),

103–116. https://doi.org/10.1016/S0001-706X(00)00179-0

Daszak, P., Cunningham, A. A., & Hyatt, A. D. (2003). Infectious disease and amphibian

population declines. *Diversity <html_ent Glyph="@amp;" Ascii="&"/>*

Distributions, *9*(2), 141–150. https://doi.org/10.1046/j.1472-4642.2003.00016.x

Dell, A. I., Pawar, S., & Savage, V. M. (2011). Systematic variation in the temperature

dependence of physiological and ecological traits. *Proceedings of the National*

Academy of Sciences, *108*(26), 10591–10596.

https://doi.org/10.1073/pnas.1015178108

Dell, A. I., Pawar, S., & Savage, V. M. (2014). Temperature dependence of trophic

interactions are driven by asymmetry of species responses and foraging strategy.

Journal of Animal Ecology, *83*(1), 70–84. https://doi.org/10.1111/1365-

2656.12081

Dobson, A. P., & Carper, R. E. (1992). Global Warming and Potential Changes in Host-

Parasite and Disease-Vector Relationships. In R. L. Peters & T. E. Lovejoy

(Eds.), *Global Warming and Biological Diversity* (pp. 201–207). Yale University

Press.

Drinkwater, K. F., Belgrano, A., Borja, A., Conversi, A., Edwards, M., Greene, C. H.,

Ottersen, G., Pershing, A. J., & Walker, H. (2003). The response of marine

ecosystems to climate variability associated with the North Atlantic Oscillation. In

J. W. Hurrell, Y. Kushnir, G. Ottersen, & M. Visbeck (Eds.), *Geophysical*

Monograph Series (Vol. 134, pp. 211–234). American Geophysical Union.

https://doi.org/10.1029/134GM10

Duncan, A. B., Fellous, S., & Kaltz, O. (2011). Temporal variation in temperature

determines disease spread and maintenance in *Paramecium* microcosm

populations. *Proceedings of the Royal Society B: Biological Sciences*, *278*(1723),

3412–3420. https://doi.org/10.1098/rspb.2011.0287

Estay, S. A., Clavijo-Baquet, S., Lima, M., & Bozinovic, F. (2011). Beyond average: An

experimental test of temperature variability on the population dynamics of

Tribolium confusum. *Population Ecology*, *53*(1), 53–58.

https://doi.org/10.1007/s10144-010-0216-7

Fumagalli, M. R., Zapperi, S., & La Porta, C. A. M. (2021). Role of body temperature

variations in bat immune response to viral infections. *Journal of The Royal*

Society Interface, *18*(180), 20210211. https://doi.org/10.1098/rsif.2021.0211

Halpin, K., Young, P. L., Field, H. E., & Mackenzie, J. S. (2000). Isolation of Hendra

virus from pteropid bats: A natural reservoir of Hendra virus. *Journal of General*

Virology, *81*(8), 1927–1932. https://doi.org/10.1099/0022-1317-81-8-1927

Harvell, D., Kim, K., Quirolo, C., Weir, J., & Smith, G. (2001). Coral bleaching and

disease: Contributors to 1998 mass mortality in Briareum asbestinum

(Octocorallia, Gorgonacea). In J. W. Porter (Ed.), *The Ecology and Etiology of*

Newly Emerging Marine Diseases (pp. 97–104). Springer Netherlands.

https://doi.org/10.1007/978-94-017-3284-0_8

Irving, A. T., Ahn, M., Goh, G., Anderson, D. E., & Wang, L.-F. (2021). Lessons from the

host defences of bats, a unique viral reservoir. *Nature*, *589*(7842), 363–370.

https://doi.org/10.1038/s41586-020-03128-0

Jiang, L., & Patel, S. N. (2008). Community assembly in the presence of disturbance: A

microcosm experiment. *Ecology*, *89*(7), 1931–1940. https://doi.org/10.1890/07-

1263.1

Johnson, P. T. J. (2006). Amphibian diversity: Decimation by disease. *Proceedings of*

the National Academy of Sciences, *103*(9), 3011–3012.

https://doi.org/10.1073/pnas.0600293103

Johnson, P. T. J., Preston, D. L., Hoverman, J. T., & LaFonte, B. E. (2013). Host and

parasite diversity jointly control disease risk in complex communities.

Proceedings of the National Academy of Sciences, *110*(42), 16916–16921.

https://doi.org/10.1073/pnas.1310557110

Jones, K. E., Patel, N. G., Levy, M. A., Storeygard, A., Balk, D., Gittleman, J. L., &

Daszak, P. (2008). Global trends in emerging infectious diseases. *Nature*,

451(7181), 990–993. https://doi.org/10.1038/nature06536

Karvonen, A., Terho, P., Seppälä, O., Jokela, J., & Valtonen, E. T. (2006). Ecological

divergence of closely related Diplostomum (Trematoda) parasites. *Parasitology*,

133(02), 229. https://doi.org/10.1017/S0031182006000242

Kim, K. S., Ejima, K., Iwanami, S., Fujita, Y., Ohashi, H., Koizumi, Y., Asai, Y., Nakaoka,

S., Watashi, K., Aihara, K., Thompson, R. N., Ke, R., Perelson, A. S., & Iwami, S.

(2021). A quantitative model used to compare within-host SARS-CoV-2, MERS-

CoV, and SARS-CoV dynamics provides insights into the pathogenesis and

treatment of SARS-CoV-2. *PLOS Biology*, *19*(3), e3001128.

https://doi.org/10.1371/journal.pbio.3001128

Kirk, D., Jones, N., Peacock, S., Phillips, J., Molnár, P. K., Krkošek, M., & Luijckx, P.

(2018). Empirical evidence that metabolic theory describes the temperature

dependency of within-host parasite dynamics. *PLOS Biology*, *16*(2), e2004608.

https://doi.org/10.1371/journal.pbio.2004608

Krichel, L., Kirk, D., Pencer, C., Hönig, M., Wadhawan, K., & Krkošek, M. (2023). Short-

term temperature fluctuations increase disease in a Daphnia-parasite infectious

disease system. *PLOS Biology*, *21*(9), e3002260.

https://doi.org/10.1371/journal.pbio.3002260

Kunze, C., Luijckx, P., Jackson, A. L., & Donohue, I. (2022). Alternate patterns of

temperature variation bring about very different disease outcomes at different

mean temperatures. *eLife*, *11*, e72861. https://doi.org/10.7554/eLife.72861

Lafferty, K. D. (2009). The ecology of climate change and infectious diseases. *Ecology*,

90(4), 888–900. https://doi.org/10.1890/08-0079.1

Laprise, F., & Dodson, J. (1994). Environmental variability as a factor controlling spatial

patterns in distribution and species diversity of zooplankton in the St. Lawrence

Estuary. *Marine Ecology Progress Series*, *107*, 67–81.

https://doi.org/10.3354/meps107067

Lawson, C. R., Vindenes, Y., Bailey, L., & van de Pol, M. (2015). Environmental

variation and population responses to global change. *Ecology Letters*, *18*(7),

724–736. https://doi.org/10.1111/ele.12437

Lips, K. R., Diffendorfer, J., Mendelson, J. R., & Sears, M. W. (2008). Riding the Wave:

Reconciling the Roles of Disease and Climate Change in Amphibian Declines.

PLoS Biology, *6*(3), e72. https://doi.org/10.1371/journal.pbio.0060072

Luo, J., Greif, S., Ye, H., Bumrungsri, S., Eitan, O., & Yovel, Y. (2021). Flight rapidly

modulates body temperature in freely behaving bats. *Animal Biotelemetry*, *9*(1),

45. https://doi.org/10.1186/s40317-021-00268-6

Molnár, P. K., Kutz, S. J., Hoar, B. M., & Dobson, A. P. (2013). Metabolic approaches to

understanding climate change impacts on seasonal host-macroparasite

dynamics. *Ecology Letters*, *16*(1), 9–21. https://doi.org/10.1111/ele.12022

Molnár, P. K., Sckrabulis, J. P., Altman, K. A., & Raffel, T. R. (2017). Thermal

Performance Curves and the Metabolic Theory of Ecology—A Practical Guide to

Models and Experiments for Parasitologists. *Journal of Parasitology*, *103*(5), 423.

https://doi.org/10.1645/16-148

Munguia, P., Osman, R., Hamilton, J., Whitlatch, R., & Zajac, R. (2010). Modeling of

priority effects and species dominance in Long Island Sound benthic

communities. *Marine Ecology Progress Series*, *413*, 229–240.

https://doi.org/10.3354/meps08764

Paaijmans, K. P., Blanford, S., Bell, A. S., Blanford, J. I., Read, A. F., & Thomas, M. B.

(2010). Influence of climate on malaria transmission depends on daily

temperature variation. *Proceedings of the National Academy of Sciences*,

107(34), 15135–15139. https://doi.org/10.1073/pnas.1006422107

Paaijmans, K. P., Heinig, R. L., Seliga, R. A., Blanford, J. I., Blanford, S., Murdock, C.

C., & Thomas, M. B. (2013). Temperature variation makes ectotherms more

sensitive to climate change. *Global Change Biology*, *19*(8), 2373–2380.

https://doi.org/10.1111/gcb.12240

Patz, J. A., Daszak, P., Tabor, G. M., Aguirre, A. A., Pearl, M., Epstein, J., Wolfe, N. D.,

Kilpatrick, A. M., Foufopoulos, J., Molyneux, D., Bradley, D. J., & Members of the

Working Group on Land Use Change Disease Emergence. (2004). Unhealthy

Landscapes: Policy Recommendations on Land Use Change and Infectious

Disease Emergence. *Environmental Health Perspectives*, *112*(10), 1092–1098.

https://doi.org/10.1289/ehp.6877

Paull, S. H., LaFonte, B. E., & Johnson, P. T. J. (2012). Temperature-driven shifts in a

host-parasite interaction drive nonlinear changes in disease risk. *Global Change*

Biology, *18*(12), 3558–3567. https://doi.org/10.1111/gcb.12018

Pedersen, A. B., & Fenton, A. (2007). Emphasizing the ecology in parasite community

ecology. *Trends in Ecology & Evolution*, *22*(3), 133–139.

https://doi.org/10.1016/j.tree.2006.11.005

Plowright, R. K., Parrish, C. R., McCallum, H., Hudson, P. J., Ko, A. I., Graham, A. L., &

Lloyd-Smith, J. O. (2017). Pathways to zoonotic spillover. *Nature Reviews*

Microbiology, *15*(8), 502–510. https://doi.org/10.1038/nrmicro.2017.45

Plowright, R., Reaser, J., Locke, H., Woodley, S. J., Patz, J. A., Becker, D., Oppler, G.,

Hudson, P., & Tabor, G. M. (2020). *A call to action: Understanding land use-*

induced zoonotic spillover to protect environmental, animal, and human health

[Preprint]. EcoEvoRxiv. https://doi.org/10.32942/osf.io/cru9w

Pounds, J. A., Fogden, M. P. L., & Campbell, J. H. (1999). Biological response to

climate change on a tropical mountain. *Nature*, *398*(6728), 611–615.

https://doi.org/10.1038/19297

Raffel, T. R., Romansic, J. M., Halstead, N. T., McMahon, T. A., Venesky, M. D., &

Rohr, J. R. (2013). Disease and thermal acclimation in a more variable and

unpredictable climate. *Nature Climate Change*, *3*(2), 146–151.

https://doi.org/10.1038/nclimate1659

Righetti, D., Vogt, M., Gruber, N., Psomas, A., & Zimmermann, N. E. (2019). Global

pattern of phytoplankton diversity driven by temperature and environmental

variability. *Science Advances*, *5*(5), eaau6253.

https://doi.org/10.1126/sciadv.aau6253

Rohr, J. R., & Raffel, T. R. (2010). Linking global climate and temperature variability to

widespread amphibian declines putatively caused by disease. *Proceedings of the*

National Academy of Sciences, *107*(18), 8269–8274.

https://doi.org/10.1073/pnas.0912883107

Rohr, J. R., Raffel, T. R., Blaustein, A. R., Johnson, P. T. J., Paull, S. H., & Young, S.

(2013). Using physiology to understand climate-driven changes in disease and

their implications for conservation. *Conservation Physiology*, *1*(1), cot022–

cot022. https://doi.org/10.1093/conphys/cot022

Rohr, J. R., Raffel, T. R., Romansic, J. M., McCallum, H., & Hudson, P. J. (2008).

Evaluating the links between climate, disease spread, and amphibian declines.

Proceedings of the National Academy of Sciences, *105*(45), 17436–17441.

https://doi.org/10.1073/pnas.0806368105

Roxburgh, S. H., Shea, K., & Wilson, J. B. (2004). The intermediate disturbance

hypothesis: Patch dynamics and mechanisms of species coexistence. *Ecology*,

85(2), 359–371. https://doi.org/10.1890/03-0266

Rynkiewicz, E. C., Pedersen, A. B., & Fenton, A. (2015). An ecosystem approach to

understanding and managing within-host parasite community dynamics. *Trends*

in Parasitology, *31*(5), 212–221. https://doi.org/10.1016/j.pt.2015.02.005

Saenz, R. A., Quinlivan, M., Elton, D., MacRae, S., Blunden, A. S., Mumford, J. A.,

Daly, J. M., Digard, P., Cullinane, A., Grenfell, B. T., McCauley, J. W., Wood, J.

L. N., & Gog, J. R. (2010). Dynamics of Influenza Virus Infection and Pathology.

Journal of Virology, *84*(8), 3974–3983. https://doi.org/10.1128/JVI.02078-09

Schoolfield, R. M., Sharpe, P. J. H., & Magnuson, C. E. (1981). Non-linear regression of

biological temperature-dependent rate models based on absolute reaction-rate

theory. *Journal of Theoretical Biology*, *88*(4), 719–731.

https://doi.org/10.1016/0022-5193(81)90246-0

Schulte, P. M., Healy, T. M., & Fangue, N. A. (2011). Thermal Performance Curves,

Phenotypic Plasticity, and the Time Scales of Temperature Exposure. *Integrative*

and Comparative Biology, *51*(5), 691–702. https://doi.org/10.1093/icb/icr097

Sckrabulis, J. P., Altman, K. A., & Raffel, T. R. (2022). Using Metabolic Theory to

Describe Temperature and Thermal Acclimation Effects on Parasitic Infection.

The American Naturalist, *199*(6), 789–803. https://doi.org/10.1086/719409

Shurin, J. B., Winder, M., Adrian, R., Keller, W. (Bill), Matthews, B., Paterson, A. M.,

Paterson, M. J., Pinel-Alloul, B., Rusak, J. A., & Yan, N. D. (2010). Environmental

stability and lake zooplankton diversity—Contrasting effects of chemical and

thermal variability. *Ecology Letters*, *13*(4), 453–463.

https://doi.org/10.1111/j.1461-0248.2009.01438.x

Sinclair, B. J., Marshall, K. E., Sewell, M. A., Levesque, D. L., Willett, C. S., Slotsbo, S.,

Dong, Y., Harley, C. D. G., Marshall, D. J., Helmuth, B. S., & Huey, R. B. (2016).

Can we predict ectotherm responses to climate change using thermal

performance curves and body temperatures? *Ecology Letters*, *19*(11), 1372–

1385. https://doi.org/10.1111/ele.12686

Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A.,

Xia, Y., Bex, V., & Midgley, P. M. (Eds.). (2013). *Climate change 2013: The*

physical science basis. In: Contribution of Working Group I to the Fifth

Assessment Report of the Intergovernmental Panel on Climate Change.

Cambridge University Press.

Symons, C. C., & Arnott, S. E. (2014). Timing is everything: Priority effects alter

community invasibility after disturbance. *Ecology and Evolution*, *4*(4), 397–407.

https://doi.org/10.1002/ece3.940

Taylor, L. H., Latham, S. M., & woolhouse, M. E. J. (2001). Risk factors for human

disease emergence. *Philosophical Transactions of the Royal Society of London.*

Series B: Biological Sciences, *356*(1411), 983–989.

https://doi.org/10.1098/rstb.2001.0888

Tracy, A. M., Pielmeier, M. L., Yoshioka, R. M., Heron, S. F., & Harvell, C. D. (2019).

Increases and decreases in marine disease reports in an era of global change.

Proceedings of the Royal Society B: Biological Sciences, *286*(1912), 20191718.

https://doi.org/10.1098/rspb.2019.1718

Vasseur, D. A., DeLong, J. P., Gilbert, B., Greig, H. S., Harley, C. D. G., McCann, K. S.,

Savage, V., Tunney, T. D., & O'Connor, M. I. (2014). Increased temperature

variation poses a greater risk to species than climate warming. *Proceedings of*

the Royal Society B: Biological Sciences, *281*(1779), 20132612.

https://doi.org/10.1098/rspb.2013.2612

Vega-Rodriguez, M., Müller-Karger, F., Hallock, P., Quiles-Perez, G., Eakin, C., Colella,

M., Jones, D., Li, J., Soto, I., Guild, L., Lynds, S., & Ruzicka, R. (2015). Influence

of water-temperature variability on stony coral diversity in Florida Keys patch

reefs. *Marine Ecology Progress Series, 528*, 173–186.

https://doi.org/10.3354/meps11268

Wright, R. K., & Cooper, E. L. (1981). Temperature effects on ectotherm immune

responses. *Developmental and Comparative Immunology, 5*, 117–122.

917