Environmental variability can promote parasite diversity within hosts and

transmission among hosts

Madeline Jarvis-Cross[1], Martin Krkosek[1]

[1] The University of Toronto, Department of Ecology and Evolutionary Biology

25 Willcocks Street, Toronto, ON, M5S 3B2

Corresponding author: Madeline Jarvis-Cross madeline.jarvis.cross@mail.utoronto.ca

1.0.0. Abstract

While the mechanisms that govern disease emergence and spread among hosts are

increasingly well-described, the mechanisms that promote parasite diversity within-

hosts, affecting host outcomes and spillover potential, have been comparatively

understudied. Furthermore, while attention has been paid to the effects of increasing

temperatures on disease systems, the effects of environmental variability have been left

underexplored, despite rising climatic variability and internal temperature variability in a

prolific reservoir for disease. To investigate the impacts of environmental variability on

parasite diversity within-hosts, we analyzed a model of within-host population dynamics

wherein two parasites indirectly compete through the host's immune response. We

simulated the model under constant, demographically stochastic, environmentally

stochastic, and demographically and environmentally stochastic conditions, and

analysed the viability and longevity of non-equilibrium parasite co-occurrence. We found

that environmental stochasticity increased the viability and longevity of parasite co-

occurrence, suggesting that thermal variability arising from climatic change and as a

physiological trait may promote parasite diversity within ectotherms and help explain

bats' propensity to support diverse communities of parasites. Further, we found that

under certain conditions, the transmissibility of co-occurring parasites can surpass the

transmissibility of single parasites, suggesting that thermal variability may increase the

transmission potential of co-occurring parasites.

2.0.0. Introduction

In 2019, SARS-CoV-2 spilled into the human population, sparking a global

pandemic from which we've yet to emerge. Worryingly, SARS-CoV-2 is not the first

zoonotic disease we've encountered, nor is it predicted to be the last (Baker et al.,

2022; Jones et al., 2008; Taylor et al., 2001). In recent years, it has become clear that

climate and land-use change can facilitate novel species interactions and increase

contact rates between wild populations and humans, resulting in devastating epidemics

and enabling spillover events and disease emergence among wild and human hosts

alike (Daszak et al., 2000, 2001; Patz et al., 2004; R. Plowright et al., 2020; R. K.

Plowright et al., 2017). However, despite the consequential interplay between within-

and among-host processes, the mechanisms that promote parasite diversity within

hosts and create kindling for events observed among hosts have been understudied. In

particular, while stabilizing mechanisms like niche differentiation have been employed to

explain the maintenance of parasite diversity within hosts, the contributions of

equalizing mechanisms, which decrease average fitness differences between

competitors, have yet to be examined (Bashey, 2015; Chesson, 2000; Chesson &

Huntly, 1997; Karvonen et al., 2006; Rynkiewicz et al., 2015).

An intermediate level of disturbance may promote species diversity by forestalling

the competitive advantages allowed by low levels of disturbance, and the extinction

risks brought on by high levels of disturbance (Chesson, 1994; Connell, 1978).

Expanding on the work of Connell (1978) and Chesson (1994), Roxburgh et al. (2004)

posited that because different species respond to disturbances in different ways, there

are myriad ways in which diversity may be generated, thereby interpreting the

intermediate disturbance hypothesis not as a mechanism unto itself, but as a collection

of mechanisms through which intermediately sized or timed disturbances in resource

availability or environmental conditions may promote diversity (Roxburgh et al., 2004).

Though the applicability of the intermediate disturbance hypothesis has been debated,

positive and negative effects of environmental variability on diversity have been well-

documented across systems (Laprise & Dodson, 1994; Righetti et al., 2019; Shurin et

al., 2010; Vega-Rodriguez et al., 2015).

When considering the effects of environmental variability on parasite diversity

within hosts, there arise a number of ways to conceptualize environmental variability.

First, one may consider rising environmental variability as a facet of climate change

(Stocker et al., 2013). While much attention has been paid to the effects of rising

temperatures on disease systems, far less has been invested in exploring the effects of

rising temperature variability, despite evidence of its outsized impacts on species

outcomes (Dobson & Carper, 1992; Lafferty, 2009; Lawson et al., 2015; Vasseur et al.,

2014). In ectothermic hosts, increasingly erratic external conditions will affect the

internal conditions of the host, thereby also affecting any parasites the host may

harbour (Cizauskas et al., 2017). A relationship between thermal variability and within-

host parasite diversity may be of particular interest when considered alongside the long-

debated relationship between climate change and widespread disease-mediated

amphibian declines (Cohen et al., 2019; Daszak et al., 1999, 2003; Lips et al., 2008;

Pounds et al., 1999; Rohr et al., 2008; Rohr & Raffel, 2010).

Second, one may consider environmental variability as a physiological trait of

hosts. In particular, bats (order *Chiroptera*) have long been acknowledged as prolific

reservoirs for disease and have oft been implicated in zoonotic spillover events (Brook

& Dobson, 2015; Calisher et al., 2006; Irving et al., 2021). Bats are also unique among

mammals in their ability to vary their internal temperature according to their activity level

(e.g. torpor, flight) (Fumagalli et al., 2021). It has been theorized that these variations in

internal temperature may aid bats' immune responses, allowing them to carry

microparasites without suffering outsized morbidity (Fumagalli et al., 2021; Luo et al.,

2021). Extending on that idea, it seems pertinent to question how internal temperature

variation may contribute to bats' propensity to support diverse communities of parasites.

We used a model to investigate the impacts of environmental variability on the

viability and maintenance of parasite diversity within hosts. We began by modifying

Antia et al.'s (1994) model of within-host population dynamics to include two parasite

species or strains that indirectly interact via the host's immune system. We analyzed a

deterministic version of the model, a demographically stochastic version of the model,

an environmentally stochastic version of the model, and a demographically and

environmentally stochastic version of the model. Within the environmentally stochastic

versions of the model, we varied the magnitude and frequency of thermal disturbance

within the system. Between and among the deterministic and stochastic simulations of

the model, we compared the ability of the secondary parasite to invade a system

containing the primary parasite and the host's immune system, and the duration of time

for which the primary and secondary parasites co-occurred. Further, we compared the

transmission potentials of co-occurring parasites with the transmission potential of a

single parasite. We found that compared to a constant environment, a variable

environment tended to increase the ability of the secondary parasite to invade the

system, facilitated longer periods of co-occurrence between the primary and secondary

parasites, and increased the transmissibility of co-occurring parasites. These results

suggest that climatic changes may have significant effects on parasite diversity within

ectothermic hosts, provide insight on bats' propensity to support diverse communities of

parasites, and influence transmission dynamics at the among-host level.

3.0.0. Methods

3.1.0. Model Description

3.1.1. Within-Host Population Dynamics Model

We began with a model of within host population dynamics based on Antia et al.,

(1994) that represents the interaction between an infecting parasite and its host's

immune system as analogous to the interaction between a prey item and its predator.

The model assumes that the replication rate of the host's immune system is proportional

to parasite density at low parasite densities, and saturates at high parasite densities,

allowing the host's immune system to either extirpate or succumb to the infection (Antia

et al., 1994). We adapted the model to include a second parasite that interacts indirectly

with the primary parasite through the host's immune response (Eqns. 1-3, Table 1).

(1) $P_i(t+1) = \left(1 + r_i - \left(1 - e^{-kI(t)}\right)\right)P_i(t)$

(2) $P_j(t+1) = (1 + r_j - (1 - e^{-kI(t)}))P_j(t)$

(3) $I(t+1) = (1 + \rho\left(\dfrac{P_i(t)}{P_i(t) + \phi} + \dfrac{P_j(t)}{P_j(t) + \phi}\right))I(t)$

Table 1: Parameter definitions and values.

Symbol	Parameter	Value in Antia et al. (1994)	Value Used
P_i and P_j	Parasite abundance	1*	1*
I	Host immune cell abundance	1*	1*
r_i and r_j	Replication rates of parasites	0.1 – 10.0	0.1 – 2.0†
k	Rate at which parasites are destroyed by host's immune system	10^{-3}	10^{-2}
ρ	Maximum replication rate of host's immune system	1.0	1.0
ϕ	Parasite density at which replication rate of host's immune system is half its maximum	10^3	10^3
D	Lethal within-host parasite density	10^9	10^9

119   * Initial conditions.
120   † Dependent on the temperature of system in environmentally stochastic simulations.
121

To compare the impacts of different types of stochasticity on parasite diversity, we wrote

a demographically stochastic version of the model (S$_D$), an environmentally stochastic

version of the model (S$_E$), and a demographically and environmentally stochastic

version of the model (S$_{DE}$). For the demographically stochastic version of the model, we

wrote the model's growth terms as Poisson processes, and mortality term as a binomial

process (Eqns. 4-6).

(4) $P_i(t+1) = P_i(t) + P_{i,B}(t) - P_{i,D}(t)$

(5) $P_j(t+1) = P_j(t) + P_{j,B}(t) - P_{j,D}(t)$

(6) $I(t+1) = I(t) + I_B(t)$

In Eqns. (4) and (5), $P_{i,B}(t)$ and $P_{j,B}(t)$ are Poisson random variables such that

$P_{i,B}(t) \sim Poisson(1, \lambda_i)$, where $\lambda_i = r_i \cdot P_i(t)$, and $P_{j,B}(t) \sim Poisson(1, \lambda_j)$, where $\lambda_j = r_j \cdot$

$P_j(t)$. In Eqn. (6), $I_B(t)$ is a Poisson random variable such that $I_B(t) \sim Poisson(1, \lambda)$,

where $\lambda = \rho \left(\frac{P_i(t)}{P_i(t)+\phi} + \frac{P_j(t)}{P_j(t)+\phi} \right) I(t)$. In Eqns. (4) and (5), $P_{i,D}(t)$ and $P_{j,D}(t)$ are binomial

random variables such that $P_{i,D}(t) \sim Binomial(1, P_i(t), p)$, where $p = 1 - e^{-kI(t)}$, and

$P_{j,D}(t) \sim Binomial(1, P_j(t), p)$. To write the environmentally stochastic version of the

model, we generated a stochastic temperature series and allowed the replication rates

of each parasite to vary in response to the temperature of the system (Eqns. 7-9).

(7) $P_i(t+1) = (1 + r_i(T(t+1)) - (1 - e^{-kI(t)}))P_i(t)$

(8) $P_j(t+1) = (1 + r_j(T(t+1)) - (1 - e^{-kI(t)}))P_j(t)$

(9) $I(t+1) = (1 + \rho \left(\frac{P_i(t)}{P_i(t)+\phi} + \frac{P_j(t)}{P_j(t)+\phi} \right))I(t)$

We used first-order autocorrelated Gaussian noise to generate stochastic temperature

series. We began with a value of 0, and generated the next value, $T(t+1)$, by

multiplying the previous value by the coefficient of autocorrelation (AC = 0.25), and

summing the resulting value with a value drawn from a normal random variable with a

mean of zero. When considering the effects of rising and frequent environmental

variability and variations, we varied and held constant the standard deviation of the

normal random variable, respectively. We then added the average temperature of the

system to each value in the series. Values outside the thermal range of the system were

re-drawn. We then converted the stochastic temperature series into series of replication

rates using various parameterizations of the Sharpe-Schoolfield sub-model, as detailed

below (Eqns. 7.1, 8.1; Table 2, Table A1).

(7.1) $r_i(T(t+1)) = r_{i,0} e^{\frac{-E_{r_i}}{k}\left(\frac{1}{T(t+1)}-\frac{1}{T_0}\right)} \cdot \left(1 + e^{\frac{E_{r_i}^L}{k}\left(\frac{1}{T_{r_i}^L}-\frac{1}{T(t+1)}\right)} + e^{\frac{E_{r_i}^H}{k}\left(\frac{1}{T_{r_i}^H}-\frac{1}{T(t+1)}\right)}\right)^{-1}$

(8.1) $r_j(T(t+1)) = r_{j,0} e^{\frac{-E_{r_j}}{k}\left(\frac{1}{T(t+1)}-\frac{1}{T_0}\right)} \cdot \left(1 + e^{\frac{E_{r_j}^L}{k}\left(\frac{1}{T_{r_j}^L}-\frac{1}{T(t+1)}\right)} + e^{\frac{E_{r_j}^H}{k}\left(\frac{1}{T_{r_j}^H}-\frac{1}{T(t+1)}\right)}\right)^{-1}$

Table 2: Parameter definitions. See Table A1 for parameter values.

Symbol	Unit	Parameter
T	K	Temperature of the system at time t
r_0		Baseline values of r_i or r_j at a reference temperature T_0
T_0	K	Reference temperature (set arbitrarily such that $T^L << T_0 << T^L$)
E_r	eV	Activation energy of r_i or r_j
E_r^L, E_r^H	eV	Low- and high-temperature inactivation energies of r_i or r_j
T_r^L, T_r^H	K	Low- and high-temperature inactivation thresholds for r_i or r_j
k		Boltzmann's constant

3.1.2. Sharpe-Schoolfield Sub-Model

The Metabolic Theory of Ecology (MTE) predicts how metabolic rates control

ecological processes across levels of organization, and explains how metabolic rates

vary with body size and temperature (Brown et al., 2004). Building off the MTE, the

Boltzmann-Arrhenius (BA) relation describes metabolism as a function of temperature.

As an extension of the BA relation, the Sharpe-Schoolfield model characterizes the

performance of a trait as a function of temperature through a thermal performance curve

(TPC) (Dell et al., 2011; Molnár et al., 2017; Schoolfield et al., 1981). When applied to

disease systems, the Sharpe-Schoolfield model has been shown to accurately describe

the relationships between temperature and species traits, and has proven useful in

investigating the relationship between rising temperatures and disease transmission

(Kirk et al., 2018; Molnár et al., 2013).

We used various parameterizations of the Sharpe-Schoolfield model to generate

a suite of paired thermal performance curves relating the temperature of the system to

the replication rates of two parasites (r_i and r_j) (see Figure 1 for a schematic of the

experimental design). We generated two sets of four paired thermal performance

curves, representing two thermal environments, and four thermal scenarios per

environment (Table S1). The first thermal environment exhibits a lower average

temperature (min. = 0°C, mean = 18 to 20.8°C, max. = 30°C), while the second exhibits

a higher average temperature (min. = 0°C, mean = 25.5 to 28°C, max. = 40°C),

representing the thermal conditions within ectotherms and bats, respectively (Fumagalli

et al., 2021). Each thermal scenario describes a different relationship between the

temperature of the system and the parasites' replication rates. Within the first scenario,

r_i and r_j respond identically to the temperature of the system (Fig. 1A, 2A). Within the

second scenario, r_i and r_j peak on opposite sides of the average temperature of the

system (Fig. 1B, 2B). To develop the third scenario, we shifted the average temperature

of the system to give the first parasite, P_i, a thermal advantage (Fig. 1C, 2C). We

considered two versions of the fourth scenario, each of which give the secondary

parasite a thermal advantage. To develop the first version, (A), we shifted the average

temperature of the system to give the secondary parasite a quantitatively comparable

thermal advantage (Fig. 1D, 2D). For the second version, (B), we reversed the curves

resulting from the third scenario to give the secondary parasite the same thermal

advantage given to the primary parasite and isolate the effect of asymmetrical thermal

performance curves (Fig. 1E, 2E). Given that versions (A) and (B) produced qualitatively

comparable results, we will only discuss version (A) moving forward (see supplement

for complete results). Within each scenario, the average temperature of the system was

chosen to maintain quantitatively comparable (1) parasite replication rates between

scenarios and (2) differences between parasite replication rates at the average

temperature of the system between the third and fourth scenarios.

196
Figure 1: Sharpe-Schoolfield thermal performance curves relating the temperature of the system to the replication
rates (r_i and r_j) of two parasites (P_i and P_j) in the low-average temperature environment, representing the conditions
within ectotherms. Yellow lines show the relationship between the temperature of the system and the replication rate
(r_i) of P_i. Green lines show the relationship between the temperature of the system and the replication rate (r_j) of P_j.
Pink lines denote the average temperature of the system. Each panel (A-E) shows a different thermal scenario.
Panels A, B, and C show the first, second, and third thermal scenarios. Panels D and E show versions A and B of the
fourth thermal scenario.

204
Figure 2: Sharpe-Schoolfield thermal performance curves relating the temperature of the system to the replication
rates (r_i and r_j) of two parasites (P_i and P_j) in the low-average temperature environment, representing the conditions
within ectotherms. Yellow lines show the relationship between the temperature of the system and the replication rate
(r_i) of P_i. Green lines show the relationship between the temperature of the system and the replication rate (r_j) of P_j.
Pink lines denote the average temperature of the system. Each panel (A-E) shows a different thermal scenario.
Panels A, B, and C show the first, second, and third thermal scenarios. Panels D and E show versions A and B of the
fourth thermal scenario.

3.2.0. Model Simulations

Within each thermal environment and scenario, we simulated the deterministic

model (D), and each of the three stochastic models (S_D, S_E, S_{DE}) for fifty days with a

discretization time step of 0.1, which was sufficient to capture the full course of parasite

infection and clearance across all simulations. We began each set of simulations with

the primary parasite, P_i, and the host's immune system in the system, at initial

abundances of 1 as the initial condition. We then considered all cases for the

introduction time of the secondary parasite, P_j, from its introduction at the onset of

infection, to its introduction on the 49[th] day of infection.

For each environmentally stochastic model (S_E, S_{DE}) simulation, we generated a

new stochastic temperature series and the corresponding series of replication rates for

each parasite. When determining the ways in which new temperatures would be

selected, we considered two axes of thermal disturbance: magnitude and frequency.

To vary the magnitude of disturbance, we varied the standard deviation of the normal

random variable in the temperature autocorrelation function, starting from zero (no

thermal disturbance) and increasing by two until the effect of magnitude on co-

occurrence plateaued (where SD ≈ 30-40), resulting in 15 and 20 magnitudes of

disturbance in the low and high average temperature environments, respectively. To

vary the frequency of thermal disturbance, we ran the model with one temperature

change every 10, 9, 8, 7, 6, 5, 4, 3, 2, and 1 days, and 2, 5, and 10 temperature

changes per day (preliminary simulations indicated this was sufficient to characterize

the results), resulting in thirteen frequencies of disturbance. We ran each permutation of

each stochastic simulation one thousand times (Fig. 3).

235
Figure 3: Schematic detailing simulations. Within the low- and high- average temperature environments, we
considered four thermal scenarios. We then considered four models (deterministic, demographically stochastic,
environmentally stochastic, and demographically and environmentally stochastic). When simulating the
environmentally stochastic models, we considered a range of magnitudes and frequencies of disturbance. When
simulating each of the four models, we considered 50 introduction days. When simulating the stochastic models, we
conducted 1000 simulation per permutation.

3.3.0. Co-Occurrence Analyses

For each simulation, we evaluated the duration of time for which the primary and

secondary parasites co-occurred (i.e. the number of time steps for which P_i and P_j had

population abundances greater than one). We then summarised these outputs to

quantify distributions of co-occurrence for each stochastic treatment. Using the

deterministic simulation outputs as a baseline, we investigated the effects of

environmental variability on the viability and maintenance of parasite diversity by

comparing the ability of the secondary parasite to invade a single-parasite system, and

the duration of time for which the primary and secondary parasites co-occurred across

stochastic treatments. Additionally, we analysed (1) the proportion of stochastic

simulations per treatment that resulted in longer periods of co-occurrence than the

deterministic simulation to quantify rarity of outcome, and (2) competitive outcomes of

each treatment to gain insight on the biological drivers of said outcomes.

3.4.0. Transmissibility Analyses

To explore the impacts of co-infection on transmission potential, we compared

the transmission potentials of co-occurring parasites with the transmission potential of a

single parasite in constant and variable environments. As per Antia et al., (1994), we

determined the total transmission (q_i, q_j, q, where q is the total transmission of the single

parasite) of a parasite (P_i, P_j, P, where P is the single parasite) from an infected host by

taking the integral of the rate of transmission (u) of each parasite from the host over the

duration of infection (Eqns. 10.1-10.3). For simplicity, we set the rate of transmission to

one.

(10.1) $q_i = \int_0^\infty u P_i(t) dt$

(10.2) $q_j = \int_0^\infty u P_j(t) dt$

(10.3) $q = \int_0^\infty u P(t) dt$

To compare constant and variable environments, we analysed multi-parasite and single-

parasite versions of the S_D and S_{DE} models. When analysing outputs from the multi-

parasite and single-parasite versions of the S_{DE} model, we considered both thermal

environments, and all four thermal scenarios. Within each thermal environment and

scenario, we considered a fully factorial design with regards to the magnitude and

frequency of disturbance.

4.0.0. Results

274   Within the deterministic model, the secondary parasite could invade ($P_{j(t+1)} >$

275     $P_{j(t)}$) when $I(t) < \frac{-\ln(1-r_j)}{k}$. Our parameterization of the model allowed the secondary

276   parasite to invade when introduced before the tenth to twelfth day of infection, resulting

277   in 8.4 to 9.2 days of co-occurrence, depending on thermal environment and scenario

278   (Fig. 4, Supplementary Datafile 1-1:10).

279
Figure 4: Longevity of co-occurrence as per the deterministic model. Both panels show the number of time steps for
which P_i and P_j have population abundances greater than one, by introduction day, as per the deterministic model.
The red, yellow, green, and blue lines show results from the first, second, third, and fourth thermal scenarios,
respectively. The black line denotes the first day on which the introduction of the secondary parasite, P_j, does not
result in any co-occurrence between the two parasites. Panel A shows the results from the low-average temperature
environment. Panel B shows the results from the high-average temperature environment.

Within both thermal environments, simulation outputs show that compared to

deterministic conditions, the introduction of demographic stochasticity has little effect on

co-occurrence (Supplementary Datafile 2-1:10), while a thermally variable environment

increases the ability of the secondary parasite to invade the system and facilitates

longer periods of co-occurrence between the primary and secondary parasites. More

specifically, the parasites co-occur for longer when the secondary parasite is introduced

earlier (Figs. 5, S1A-S1E).

293
Figure 5: Simulation results and summary. Panel A shows the number of time steps for which P_i and P_j have
population abundances greater than one, by introduction day, as per the demographically and environmentally
stochastic model (in the low-average temperature environment, under the first thermal scenario, when the magnitude
and frequency of disturbance are intermediate). Each dot shows the result of a single simulation. The grey lines
represent medians, and red diamonds represent means. Red stars show the number of time steps for which P_i and P_j
have population abundances greater than one as per the deterministic model. The blue line denotes the first day on
which the introduction of the secondary parasite, P_j, does not result in any co-occurrence between the two parasites.
Panel B shows a summarised version of Panel A, where each line represents mean longevities of co-occurrence by
introduction day. The black and blue lines show results from the deterministic and demographically stochastic model
simulations, respectively. The solid and dashed red lines show results from the environmentally stochastic and
demographically and environmentally stochastic model simulations, respectively.

Additionally, while duration of co-occurrence is maximised under the fourth thermal

scenario (P_j has a thermal advantage over P_i), the difference between the deterministic

and environmentally stochastic models is not consistently maximised by any one

thermal scenario. With respect to the combined effects of magnitude and frequency of

thermal disturbance on the longevity of co-occurrence between the primary and

secondary parasites, while higher magnitudes and frequencies of disturbance tend to

promote longer periods of co-occurrence, increasing magnitudes of disturbance

produce a less steep effect than do increasing frequencies of disturbance. These

findings are common among thermal environments and scenarios.

4.1.0. Ectotherms: Low Average Temperature Environment (LATE)

When the frequency of thermal disturbance is fixed and the magnitude of thermal

disturbance varies, environmental variation can allow for co-occurrence beyond the

deterministically simulated threshold (DST) and can promote longer periods of co-

occurrence compared with the deterministic and demographically stochastic

environments. More specifically, when the frequency of thermal disturbance is fixed at

an intermediate value of ten (once per day) and the magnitude of thermal disturbance

varies, environmental variation allows for some amount of co-occurrence when the

secondary parasite is introduced 10 to 25 (S_E) and 17 to 20 days (S_{DE}) after the DST (9

to 11 days, depending on thermal scenario), and allows parasites to co-occur for a

maximum average of 2.95 to 3.84 days (S_E) and 2.56 to 3.66 days (S_{DE}) past the

deterministic maximum (8.4 to 9.2 days) and an absolute maximum of 17.5 to 22.8 days

(S_E) and 18.1 to 22.4 days (S_{DE}), depending on thermal scenario. Further, as the size of

thermal disturbances increases, periods of co-occurrence increase in length to a

saturation point (standard deviation of the normal random variable in the temperature

autocorrelation function ≈ 24-30, depending on thermal environment and scenario)

(Figs. 6, S2, S4A-S4E, Supplementary Datafiles 3-1:5, 4-1:5).

331
Figure 6: The impacts of varying magnitudes of disturbance on longevity of co-occurrence in the low-average
temperature environment. Each panel shows the mean number of time steps for which P_i and P_j have population
abundances greater than one, by introduction day, when the magnitude of disturbance varies, and the frequency of
disturbance is intermediate (once per day). The black and blue lines show results from the deterministic and
demographically stochastic model simulations. Solid and dashed lines show results from the environmentally and
demographically and environmentally model simulations. Pink, orange, yellow, and green lines show results when
magnitudes of disturbance are low (SD = 2), low-intermediate (SD = 10), intermediate-high (SD = 20), and high (SD =
30), respectively. Panels A-D show results from the first, second, third, and fourth thermal scenarios.

4.2.0. Bats: High Average Temperature Environment (HATE)

When the magnitude of thermal disturbance is fixed and the frequency of thermal

disturbance varies, environmental variation can allow for co-occurrence beyond the

deterministically simulated threshold and can promote longer periods of co-occurrence

compared with the deterministic and demographically stochastic environments. More

specifically, when the magnitude of thermal disturbance is fixed at an intermediate value

(standard deviation of the normal random variable in the temperature autocorrelation

function = 10) and the frequency of thermal disturbance varies, environmental variation

allows for some amount of co-occurrence when the secondary parasite is introduced 23

to 35 (S_E) and 17 to 36 days (S_{DE}) after the DST (9 to 10 days, depending on thermal

scenario). When the frequency of disturbance is low (once every seven to ten days),

periods of co-occurrence are similar in length to those resulting from deterministic

simulations. Given that deterministically, P_i and P_j can co-occur for a maximum of nine

days, environments subject to infrequent disturbances are functionally similar to a

deterministic environment, making the outcome of a simulation reliant on the initial

thermal conditions of the system. Interestingly, when the frequency of disturbance is

intermediate (once every few days), periods of co-occurrence are routinely shorter than

those resulting from deterministic simulations, to an average minimum of 0.96 to 1.53

(S_E) and 1.21 to 1.95 (S_{DE}) days fewer than the deterministic maximum (8.4 to 9.0

days). Periods of co-occurrence are longest when disturbances are very frequent (once

every two days to ten times per day). When disturbances are very frequent, P_i and P_j

co-occur for an average maximum of 2.63 to 4.57 (S_E) and 2.87 and 4.13 (S_{DE}) days

past the deterministic maximum and an absolute maximum of 24.7 to 36.7 (S_E) and 23.5

to 28.5 (S_{DE}) days, depending on thermal scenario (see supplement for complete

results) (Figs. 7, S3, S7A-S7E, Supplementary Datafiles 3-6:10, 4-6:10).

365
Figure 7: The impacts of varying frequencies of disturbance on longevity of co-occurrence in the high-average
temperature environment. Each panel shows the mean number of time steps for which P_i and P_j have population
abundances greater than one, by introduction day, when the frequency of disturbance varies, and the magnitude of
disturbance is intermediate (SD = 10). The black and blue lines show results from the deterministic and
demographically stochastic model simulations. Solid and dashed lines show results from the environmentally and
demographically and environmentally model simulations. Pink, orange, yellow, and green lines show results when
frequencies of disturbance are low (once every 10 days), low-intermediate (once every 5 days), intermediate-high
(once per day), and high (ten times per day), respectively. Panels A-D show results from the first, second, third, and
fourth thermal scenarios.

4.3.0. Rarity and Competitive Outcomes

Analyzing the number of simulations per treatment that resulted in a non-zero

period of co-occurrence revealed that the effect of environmental variability on the

viability and maintenance of parasite co-occurrence is not rare. When the secondary

parasite is introduced early (before the 10th day of infection), ⪞50% of environmentally

stochastic simulations result in longer periods of co-occurrence than deterministic

simulations. While the likelihood that an environmentally stochastic simulation will result

in co-occurrence sharply declines when the secondary parasite is introduced after the

deterministically simulated threshold, four to nine percent of environmentally stochastic

simulations result in some amount of co-occurrence four to six days after said threshold,

depending on thermal scenario (Fig. S9).

Analyzing the competitive outcomes of each treatment revealed that under the

second thermal scenario (r_i and r_j peak on opposite sides of the average temperature

of the system), when P_i and P_j are introduced at the same time, P_j outlasts P_i 65.5%

(LATE, S_E model) and 63.4% (LATE, S_{DE} model) of the time. Additionally, under the

third thermal scenario (P_i has a thermal advantage), when P_i and P_j are introduced at

the same time, P_i and P_j are equally likely to outlast each other (P_j outlasts P_i 49.9%

(LATE, S_E model) and 51.0% (LATE, S_{DE} model) of the time), but under the fourth

thermal scenario (P_j has a thermal advantage), P_j is much more likely to outlast P_i (P_j

outlasts P_i 76.6% (LATE, S_E model) and 70.4% (LATE, S_{DE} model) of the time).

Regardless of thermal environment or scenario, when the secondary parasite, P_j, is

introduced later, P_i is more likely to outlast P_j, until the point at which P_j is introduced

when P_i is about to be or has been extirpated (Fig. S11A-F). However, regardless of

when P_j is introduced, simulations resulting in especially long periods of co-occurrence

are equally likely to show either parasite outlasting the other (Fig. S10A-F).

4.4.0. Transmissibility

Comparing the transmission potentials (TPs) of co-occurring and single parasites

revealed that under certain conditions, the transmissibility of co-occurring parasites can

surpass the transmissibility of single parasites. Across thermal environments and

scenarios, TPs from the multi-parasite version of the S_D model (median TP of P_i ranged

between 1.49E+04 and 9.12E+05, median TP of P_j ranged between 2.59E+04 and

6.09E+05) were one to two orders of magnitude lower than TPs from the single-parasite

versions of the S_D model (median TP of P ranged between 3.62E+06 and 5.13E+07)

(Supplementary Datafiles 5-1:16). Similarly, under the first three thermal scenarios, TPs

from the multi-parasite version of the S_{DE} model (median TP of P_i ranged between

1.02E+04 and 1.05E+06, median TP of P_j ranged between 8.36E+03 and 3.00E+05)

were one to two orders of magnitude lower than TPs from the single-parasite versions

of the S_{DE} model (median TP of P ranged between 6.45E+04 and 3.41E+07). However,

under the fourth thermal scenario, intermediate magnitudes and frequencies of

disturbance often resulted in the TP of the secondary parasite, P_j (median TP ranged

between 2.50E+04 and 7.76E+05), exceeding that of a single parasite (median TP

ranged between 1.20E+05 and 1.07E+06. More specifically, in the low and high

average temperature environments under the fourth thermal scenario, the TP of the

secondary parasite, P_j, exceeded that of a single parasite in 62 of 195 and 60 of 260

disturbance regimes, or, in about 20-30% of disturbance regimes. In the low average

temperature environment, the TP of P_j exceeded that of a single parasite by a maximum

of 3.40E+05, when the magnitude of disturbance was equal to 20 and the frequency of

disturbance was equal to 80 (Figs. 8A, S13A, Supplementary Datafiles 5-8). In the high

average temperature environment, the TP of P_j exceeded that of a single parasite by a

maximum of 2.55E+05, when the magnitude of disturbance was equal to 30 and the

frequency of disturbance was equal to 80 (Figs. 8B, S13B, Supplementary Datafiles 5-

16). Regardless of environment and scenario, the transmission potentials of co-

occurring parasites peaked across a range of magnitudes of disturbance when the

frequency of disturbance was low (once every ten days) to intermediate (once every six

days) (Fig. S12A-H).

Figure 8: The impacts of environmental variability on the transmission potential of parasites. Panels A and B show the transmission potentials of parasites by model (S_D, S_{DE}) and formulation (1 Parasite, 2 Parasites). Further, Panels A and B show the two cases in which the inclusion of environmental variability resulted in the transmission potentials of co-occurring parasites exceeding that of a single parasite. Yellow (single parasite) and orange (co-occurring parasites) boxes show results from the demographically stochastic model simulations. Pink (single parasite) and magenta (co-occurring parasites) boxes show results from the demographically and environmentally stochastic model simulations. Panel A shows results from simulations in the low-average temperature environment, under the fourth thermal scenario when the magnitude of disturbance was intermediate (SD = 20), and disturbances occurred every eight days. Panel B shows results from simulations in the high-average temperature environment, under the fourth thermal scenario when the magnitude of disturbance was intermediate (SD = 30), and disturbances occurred every eight days.

5.0.0. Discussion

Despite increasing rates of disease emergence, the mechanisms that promote parasite diversity within hosts and create kindling for events observed among hosts, have been understudied (Baker et al., 2022; Jones et al., 2008; Taylor et al., 2001). It has been posited that by forestalling the competitive advantages allowed by low levels of disturbance, and the extinction risks brought on by high levels of disturbance, intermediately sized or timed disturbances in resource availability or environmental conditions may promote species diversity (Chesson, 1994; Connell, 1978; Roxburgh et al., 2004). Here, we investigated the potential for environmental variability, both as a facet of climate change and as a physiological trait unique to a prolific reservoir for disease, bats (order *Chiroptera*), to affect the viability and maintenance of within-host parasite diversity. The results of this work suggest that compared to a constant environment, demographic stochasticity has little impact on parasite co-occurrence, while the addition of environmental variability, and more specifically thermal variability,

can increase the ability of a secondary parasite to invade a single-parasite system,

promote longer periods of co-occurrence between parasites, and increase the

transmissibility of co-occurring parasites. The results of this work also indicate that the

primary and secondary parasites were able to co-occur for longer when the secondary

parasite was introduced earlier, and that the longevity of co-occurrence was maximised

when the secondary parasite had a thermal advantage.

In the low average temperature environment, representing the thermal conditions

within ectotherms subject to rising environmental variability, the relationship between

the magnitude of thermal disturbance and the longevity of co-occurrence was initially

positive and eventually saturating. It was determined that increasing magnitudes of

disturbance produced a saturating response rather than the expected parabolic

response due to the structure of the Sharpe-Schoolfield sub-model, which cannot

generate negative replication rates. When we allowed for negative replication rates,

larger disturbances began to have negative effects on parasite populations, resulting in

the expected parabolic response, and implying that intermediate magnitudes of

disturbance maximised the viability and longevity of parasite co-occurrence (Fig. S5A-D,

Fig. S6A-D).

The relationship between climate change and disease-mediated amphibian

declines has long been debated. While some argue that climatic changes may impede

immune function or result in performance gaps between parasites and their hosts, and

thus make amphibians more susceptible to infection, others argue that insufficient

evidence should preclude a causal link (Cohen et al., 2019, 2019; Daszak et al., 1999,

2003; Lips et al., 2008; Pounds et al., 1999; Rohr et al., 2008). While much of the

literature proposes mean-climate hypotheses, some evidence suggests that intensifying

El Niño events are increasing thermal variability, which may in turn affect amphibians'

susceptibility to or ability to defend against infection (Rohr & Raffel, 2010). Similarly,

emerging evidence suggests that in addition to rising mean temperatures, daily

temperature fluctuations may affect the incubation period of malaria parasites and

related vital rates, altering malaria transmission patterns (Paaijmans et al., 2010, 2013).

Congruently, our findings provide theoretical evidence that thermal variability promotes

the viability and maintenance of within-host parasite diversity, and thus suggest that as

the climate continues to change, close attention should be paid to shifts in microparasite

communities and their impacts on ectothermic hosts. In particular, well-documented

climate-driven increases in disease incidence among a wide variety of marine

organisms may constitute an important starting point for future work (Bates et al., 2009;

Bruno et al., 2007; Carilli et al., 2012; Drinkwater et al., 2003; Harvell et al., 2001; Tracy

et al., 2019).

In the high average temperature environment, intermediate and high frequencies

of disturbance impeded and promoted the viability and longevity of parasite co-

occurrence, respectively. Given the transient nature of infection specified by the

underlying model, intermediate frequencies of disturbance (once every seven days, for

example) resulted in the temperature of the system being disturbed towards the end of

the infection period, when parasite populations were small and especially susceptible to

unfavourable conditions. Conversely, high frequencies of disturbance resulted in the

temperature of the system being disturbed throughout the duration of the infection

period, and as such, neither favourable nor unfavorable conditions lasted long enough

for one parasite to gain a decisive advantage over and exclude the other. While it has

been theorized that variations in internal temperature may aid bats' immune responses,

allowing them to carry microparasites without suffering outsized morbidity (Fumagalli et

al., 2021; Luo et al., 2021), these findings suggest that variations in internal temperature

may also facilitate the co-occurrence of parasites within bats, and thus may help explain

bats' propensity to support diverse communities of parasites. Further, these results,

combined with the regularity with which bats are implicated in spillover events and

epidemics, suggest that the observed effects of environmental variability on parasite

diversity within hosts will scale up to have effects on between-host processes like

transmission, as discussed below.

With regards to competitive outcomes, under the second thermal scenario (r_i and

r_j peak on opposite sides of the average temperature of the system), P_j often outlasted

P_i, while under the third scenario, P_i and P_j were equally likely to outlast each other, and

under the fourth scenario, P_j was much more likely to outlast P_i. These outcomes are, in

part, consequences of the asymmetrical structure of the thermal performance curves.

Within the second thermal scenario, colder than average temperatures resulted in a

slightly higher replication rate for P_i and a slightly lower replication rate for P_j, while

warmer than average temperatures resulted in a much lower replication rate for P_i and a

slightly higher replication rate for P_j. In short, P_i was more disadvantaged by

unfavourable conditions than was P_j. Further, regardless of when P_j was introduced,

simulations resulting in especially long periods of co-occurrence were equally likely to

show either parasite outlasting the other. Taken together, these results suggest that to

predict the competitive outcome of a multiple infection, one may need to consider the

interplay between how well adapted a parasite is to its host, priority effects, and

environmental factors. While one might expect a specialist or well-adapted parasite

(here, a thermally advantaged parasite) to exploit its host more efficiently than, and thus

outcompete, a generalist or novel parasite, the results of this work suggest that

environmental variability can even the scales, and allow a generalist or novel parasite to

establish and persist, as predicted by the intermediate disturbance hypothesis

(Chesson, 1994; Connell, 1978; Roxburgh et al., 2004). Similarly, while one might

additionally expect a colonizing parasite (here, the primary parasite) to benefit from

priority effects, the results of this work again suggest that environmental variability and

disturbance may contribute to the structuring of communities by allowing latecomers to

establish and persist more often and for longer than they otherwise would (Jiang &

Patel, 2008; Munguia et al., 2010; Symons & Arnott, 2014). As anthropogenic changes

alter host-parasite interaction networks, understanding how evolutionary and

environmental factors combine to affect interactions between parasites will be essential

in predicting and managing spillover and emergence in wildlife and human populations

(Paull et al., 2012; Pedersen & Fenton, 2007).

From an array of parasites circulating among ectotherms to Hendra virus,

emerging coronaviruses, and other parasites carried by bats, developing an

understanding of how the impacts of environmental variability on parasite diversity

within hosts may scale to affect among-host processes and events, including

transmission, spillover, and emergence will be essential to the maintenance of wildlife,

human, and ecosystem health (Allocati et al., 2016; Daszak et al., 2003; Halpin et al.,

2000; Johnson, 2006). While increased within-host parasite diversity provides more

strains or species of a parasite or parasites an opportunity to transmit, within-host

competition may result in decreased parasite loads, thus decreasing the transmission

probability of any one strain or species, and altering expected transmission patterns

(Johnson et al., 2013; Pedersen & Fenton, 2007). Similarly, our findings provide

theoretical evidence that when environmental variations are very large and very

frequent, within-host competition may decrease the transmission rates of coinfecting

parasites, thereby decreasing transmission probability. However, our findings also

suggest that when environmental variations are low to intermediately sized and timed,

the transmission rates of coinfecting parasites are comparable to, or in excess of, the

transmission rates of single parasites, indicating that the effects of environmental

variability may not be limited to the promotion of within-host parasite diversity, but may

also promote transmission between hosts. Additionally, we found that while large and

frequent disturbances may maximally increase the amount of time for which two

parasites co-occur, such infections do not result in particularly high transmission

potentials, indicating that especially long periods of co-occurrence may be facilitated by

lower replicative outputs that do not trigger a large immune response, and thus allow for

more persistent infections. Conversely, low to intermediately sized and timed

disturbances may constitute an anomalous region within which environmental variation

can increase the amount of time for which two parasites co-occur, while allowing

parasites to achieve high replicative outputs, resulting in longer and more transmissible

infections.

Some studies have posited that high host diversity may decrease transmission

risk via the dilution effect and increase parasite diversity among hosts by enhancing

colonization opportunities, while others have posited that high host diversity may

increase transmission risk via the amplification effect (Hechinger & Lafferty, 2005;

Johnson et al., 2015; Ostfeld & Keesing, 2012). Our findings suggest that by promoting

transmission between hosts, low to intermediately sized and timed environmental

variations may also increase parasite diversity among hosts. It is important to

emphasize that while the impacts of global change on parasite diversity and of

increasing or decreasing parasite diversity on wildlife and human global health remain

disputed, parasites, like predators, are essential to the communities of which they are a

part (Altizer et al., 2013; Anderson & May, 1979; Carlson et al., 2017; Cizauskas et al.,

2017; Dobson & Hudson, 1986; Rohr et al., 2011). As such, the maintenance of parasite

diversity should be, and increasingly is, recognized as an essential aspect of

conservation ecology (Carlson et al., 2013; Dougherty et al., 2016).

While both Antia et al.'s (1994) model and our modified model should be broadly

applicable to an array of within-host multi-parasite systems, including multi-strain or

cross-species SARS-CoV-2 and influenza infections, both models are relatively simple

as to remain generalizable, and thus cannot represent chronic infections, systems within

which parasites exhibit more complex interactions with their host or their host's immune

system (ex. malaria, HIV), nor systems within which parasites directly interact with one

another, as is often observed in trematode coinfection and in hosts coinfected with

multiple strains of the bacterium *Pseudomonas aeruginosa* (Al Khatib et al., 2020;

Bashey, 2015; Kim et al., 2021; Saenz et al., 2010). As such, it may be interesting to

validate the theoretical insights of this work in an empirical system. It may be of

particular interest to explore the impacts of environmental variability on within-host

parasite diversity in amphibians, mosquitos, or marine organisms, as discussed above,

in bats, given the risk posed to public health, or in rodents, another group of mammals

that tends to host diverse communities of parasites (Bates et al., 2009; Bruno et al.,

2007; Carilli et al., 2012; Drinkwater et al., 2003; Paaijmans et al., 2010; Rohr & Raffel,

2010; Tracy et al., 2019). This work is also limited by the scope with which thermal

dependency was considered. Given the thermal sensitivity of immune function in

ectotherms and endotherms, it will be worthwhile to consider the thermal dependency of

immune processes like response time and replication rate in tandem with the thermal

dependency of parasite replication rates (Butler et al., 2013; Wright & Cooper, 1981).

This work provides theoretical evidence that through climate change and as a

physiological trait, environmental variability may promote parasite diversity within hosts

and transmission among hosts, and thus create kindling for the spillover events, disease

emergences, and epidemics we observe among wild and human hosts alike.

Anthropogenic changes have allowed the next pandemic to leave the hypothetical and

become a looming spectre. Camus wrote that plagues, despite their commonality, find

us unprepared (Camus, Albert, 1991). Improving our understanding of how

environmental variability will impact parasite diversity within- and between-hosts will

allow us to expand our understanding of the mechanisms that can facilitate disease

spread and emergence, which will allow us to make robust predictions, and develop and

implement mitigation strategies as we move into an increasingly uncertain future. In

brief, we will be found *less* unprepared.

6.0.0. Acknowledgements

This work was supported by The Emerging and Pandemic Infections Consortium

Doctoral Award (MJ-C), an NSERC Discovery Grant (MK) and Canada Research Chair

(MK).

621   **7.0.0. References**

Al Khatib, H. A., Benslimane, F. M., Elbashir, I. E., Coyle, P. V., Al Maslamani, M. A., Al-

Khal, A., Al Thani, A. A., & Yassine, H. M. (2020). Within-Host Diversity of SARS-

CoV-2 in COVID-19 Patients With Variable Disease Severities. *Frontiers in*

Cellular and Infection Microbiology, *10*, 575613.

https://doi.org/10.3389/fcimb.2020.575613

Allocati, N., Petrucci, A. G., Di Giovanni, P., Masulli, M., Di Ilio, C., & De Laurenzi, V.

(2016). Bat–man disease transmission: Zoonotic pathogens from wildlife

reservoirs to human populations. *Cell Death Discovery*, *2*(1), 16048.

https://doi.org/10.1038/cddiscovery.2016.48

Altizer, S., Ostfeld, R. S., Johnson, P. T. J., Kutz, S., & Harvell, C. D. (2013). Climate

Change and Infectious Diseases: From Evidence to a Predictive Framework.

Science, *341*(6145), 514–519. https://doi.org/10.1126/science.1239401

Anderson, R. M., & May, R. M. (1979). Population biology of infectious diseases: Part I.

Nature, *280*(5721), 361–367. https://doi.org/10.1038/280361a0

Antia, R., Levin, B. R., & May, R. M. (1994). Within-Host Population Dynamics and the

Evolution and Maintenance of Microparasite Virulence. *The American Naturalist*,

144(3), 457–472. https://doi.org/10.1086/285686

Baker, R. E., Mahmud, A. S., Miller, I. F., Rajeev, M., Rasambainarivo, F., Rice, B. L.,

Takahashi, S., Tatem, A. J., Wagner, C. E., Wang, L.-F., Wesolowski, A., &

Metcalf, C. J. E. (2022). Infectious disease in an era of global change. *Nature*

Reviews Microbiology, *20*(4), 193–205. https://doi.org/10.1038/s41579-021-

00639-z

Bashey, F. (2015). Within-host competitive interactions as a mechanism for the

maintenance of parasite diversity. *Philosophical Transactions of the Royal*

Society B: Biological Sciences, *370*(1675), 20140301.

https://doi.org/10.1098/rstb.2014.0301

Bates, A., Hilton, B., & Harley, C. (2009). Effects of temperature, season and locality on

wasting disease in the keystone predatory sea star Pisaster ochraceus. *Diseases*

of Aquatic Organisms, *86*, 245–251. https://doi.org/10.3354/dao02125

Brook, C. E., & Dobson, A. P. (2015). Bats as 'special' reservoirs for emerging zoonotic

pathogens. *Trends in Microbiology*, *23*(3), 172–180.

https://doi.org/10.1016/j.tim.2014.12.004

Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., & West, G. B. (2004). Toward a

metabolic theory of ecology. *Ecology*, *85*(7), 1771–1789.

https://doi.org/10.1890/03-9000

Bruno, J. F., Selig, E. R., Casey, K. S., Page, C. A., Willis, B. L., Harvell, C. D.,

Sweatman, H., & Melendy, A. M. (2007). Thermal Stress and Coral Cover as

Drivers of Coral Disease Outbreaks. *PLoS Biology*, *5*(6), e124.

https://doi.org/10.1371/journal.pbio.0050124

Butler, M. W., Stahlschmidt, Z. R., Ardia, D. R., Davies, S., Davis, J., Guillette, L. J.,

    Johnson, N., McCormick, S. D., McGraw, K. J., & DeNardo, D. F. (2013).

    Thermal Sensitivity of Immune Function: Evidence against a Generalist-

    Specialist Trade-Off among Endothermic and Ectothermic Vertebrates. *The*

    American Naturalist, *181*(6), 761–774. https://doi.org/10.1086/670191

Calisher, C. H., Childs, J. E., Field, H. E., Holmes, K. V., & Schountz, T. (2006). Bats:

    Important Reservoir Hosts of Emerging Viruses. *Clinical Microbiology Reviews*,

    19(3), 531–545. https://doi.org/10.1128/CMR.00017-06

Camus, A. (1991). *The Plague* (T. Stuart Gilbert.). Vintage. (Original work published

    1948).

Carilli, J., Donner, S. D., & Hartmann, A. C. (2012). Historical Temperature Variability

    Affects Coral Response to Heat Stress. *PLoS ONE*, *7*(3), e34418.

    https://doi.org/10.1371/journal.pone.0034418

Carlson, C. J., Burgio, K. R., Dougherty, E. R., Phillips, A. J., Bueno, V. M., Clements,

    C. F., Castaldo, G., Dallas, T. A., Cizauskas, C. A., Cumming, G. S., Doña, J.,

    Harris, N. C., Jovani, R., Mironov, S., Muellerklein, O. C., Proctor, H. C., & Getz,

    W. M. (2017). Parasite biodiversity faces extinction and redistribution in a

    changing climate. *Science Advances*, *3*(9), e1602422.

    https://doi.org/10.1126/sciadv.1602422

Carlson, C. J., Cizauskas, C. A., Burgio, K. R., Clements, C. F., & Harris, N. C. (2013).

    The More Parasites, the Better? *Science*, *342*(6162), 1041–1041.

    https://doi.org/10.1126/science.342.6162.1041-a

Chesson, P. (1994). Multispecies Competition in Variable Environments. *Theoretical*

Population Biology, *45*(3), 227–276. https://doi.org/10.1006/tpbi.1994.1013

Chesson, P. (2000). Mechanisms of Maintenance of Species Diversity. *Annual Review*

of Ecology and Systematics, *31*(1), 343–366.

https://doi.org/10.1146/annurev.ecolsys.31.1.343

Chesson, P., & Huntly, N. (1997). The Roles of Harsh and Fluctuating Conditions in the

Dynamics of Ecological Communities. *The American Naturalist*, *150*(5), 519–553.

https://doi.org/10.1086/286080

Cizauskas, C. A., Carlson, C. J., Burgio, K. R., Clements, C. F., Dougherty, E. R.,

Harris, N. C., & Phillips, A. J. (2017). Parasite vulnerability to climate change: An

evidence-based functional trait approach. *Royal Society Open Science*, *4*(1),

160535. https://doi.org/10.1098/rsos.160535

Cohen, J. M., Civitello, D. J., Venesky, M. D., McMahon, T. A., & Rohr, J. R. (2019). An

interaction between climate change and infectious disease drove widespread

amphibian declines. *Global Change Biology*, *25*(3), 927–937.

https://doi.org/10.1111/gcb.14489

Connell, J. H. (1978). Diversity in Tropical Rain Forests and Coral Reefs: High diversity

of trees and corals is maintained only in a nonequilibrium state. *Science*,

199(4335), 1302–1310. https://doi.org/10.1126/science.199.4335.1302

Daszak, P., Berger, L., Cunningham, A. A., Hyatt, A. D., Green, D. E., & Speare, R.

(1999). Emerging Infectious Diseases and Amphibian Population Declines.

Emerging Infectious Diseases, *5*(6), 735–748.

https://doi.org/10.3201/eid0506.990601

Daszak, P., Cunningham, A. A., & Hyatt, A. D. (2000). Emerging Infectious Diseases of

Wildlife—Threats to Biodiversity and Human Health. *Science*, *287*(5452), 443–

449. https://doi.org/10.1126/science.287.5452.443

Daszak, P., Cunningham, A. A., & Hyatt, A. D. (2001). Anthropogenic environmental

change and the emergence of infectious diseases in wildlife. *Acta Tropica*, *78*(2),

103–116. https://doi.org/10.1016/S0001-706X(00)00179-0

Daszak, P., Cunningham, A. A., & Hyatt, A. D. (2003). Infectious disease and amphibian

population declines. *Diversity <html_ent Glyph="@amp;" Ascii="&"/>*

Distributions, *9*(2), 141–150. https://doi.org/10.1046/j.1472-4642.2003.00016.x

Dell, A. I., Pawar, S., & Savage, V. M. (2011). Systematic variation in the temperature

dependence of physiological and ecological traits. *Proceedings of the National*

Academy of Sciences, *108*(26), 10591–10596.

https://doi.org/10.1073/pnas.1015178108

Dobson, A. P., & Carper, R. E. (1992). Global Warming and Potential Changes in Host-

Parasite and Disease-Vector Relationships. In R. L. Peters & T. E. Lovejoy

(Eds.), *Global Warming and Biological Diversity* (pp. 201–207). Yale University

Press.

Dobson, A. P., & Hudson, P. J. (1986). Parasites, disease and the structure of

ecological communities. *Trends in Ecology & Evolution*, *1*(1), 11–15.

https://doi.org/10.1016/0169-5347(86)90060-1

Dougherty, E. R., Carlson, C. J., Bueno, V. M., Burgio, K. R., Cizauskas, C. A.,

Clements, C. F., Seidel, D. P., & Harris, N. C. (2016). Paradigms for parasite

conservation: Parasite Conservation. *Conservation Biology*, *30*(4), 724–733.

https://doi.org/10.1111/cobi.12634

Drinkwater, K. F., Belgrano, A., Borja, A., Conversi, A., Edwards, M., Greene, C. H.,

Ottersen, G., Pershing, A. J., & Walker, H. (2003). The response of marine

ecosystems to climate variability associated with the North Atlantic Oscillation. In

J. W. Hurrell, Y. Kushnir, G. Ottersen, & M. Visbeck (Eds.), *Geophysical*

Monograph Series (Vol. 134, pp. 211–234). American Geophysical Union.

https://doi.org/10.1029/134GM10

Fumagalli, M. R., Zapperi, S., & La Porta, C. A. M. (2021). Role of body temperature

variations in bat immune response to viral infections. *Journal of The Royal*

Society Interface, *18*(180), 20210211. https://doi.org/10.1098/rsif.2021.0211

Halpin, K., Young, P. L., Field, H. E., & Mackenzie, J. S. (2000). Isolation of Hendra

virus from pteropid bats: A natural reservoir of Hendra virus. *Journal of General*

Virology, *81*(8), 1927–1932. https://doi.org/10.1099/0022-1317-81-8-1927

Harvell, D., Kim, K., Quirolo, C., Weir, J., & Smith, G. (2001). Coral bleaching and

disease: Contributors to 1998 mass mortality in Briareum asbestinum

(Octocorallia, Gorgonacea). In J. W. Porter (Ed.), *The Ecology and Etiology of*

Newly Emerging Marine Diseases (pp. 97–104). Springer Netherlands.

https://doi.org/10.1007/978-94-017-3284-0_8

Hechinger, R. F., & Lafferty, K. D. (2005). Host diversity begets parasite diversity: Bird

final hosts and trematodes in snail intermediate hosts. *Proceedings of the Royal*

Society B: Biological Sciences, *272*(1567), 1059–1066.

https://doi.org/10.1098/rspb.2005.3070

Irving, A. T., Ahn, M., Goh, G., Anderson, D. E., & Wang, L.-F. (2021). Lessons from the

host defences of bats, a unique viral reservoir. *Nature*, *589*(7842), 363–370.

https://doi.org/10.1038/s41586-020-03128-0

Jiang, L., & Patel, S. N. (2008). Community assembly in the presence of disturbance: A

microcosm experiment. *Ecology*, *89*(7), 1931–1940. https://doi.org/10.1890/07-

1263.1

Johnson, P. T. J. (2006). Amphibian diversity: Decimation by disease. *Proceedings of*

the National Academy of Sciences, *103*(9), 3011–3012.

https://doi.org/10.1073/pnas.0600293103

Johnson, P. T. J., Ostfeld, R. S., & Keesing, F. (2015). Frontiers in research on

biodiversity and disease. *Ecology Letters*, *18*(10), 1119–1133.

https://doi.org/10.1111/ele.12479

Johnson, P. T. J., Preston, D. L., Hoverman, J. T., & LaFonte, B. E. (2013). Host and

parasite diversity jointly control disease risk in complex communities.

Proceedings of the National Academy of Sciences, *110*(42), 16916–16921.

https://doi.org/10.1073/pnas.1310557110

Jones, K. E., Patel, N. G., Levy, M. A., Storeygard, A., Balk, D., Gittleman, J. L., &

Daszak, P. (2008). Global trends in emerging infectious diseases. *Nature*,

451(7181), 990–993. https://doi.org/10.1038/nature06536

Karvonen, A., Terho, P., Seppälä, O., Jokela, J., & Valtonen, E. T. (2006). Ecological

divergence of closely related Diplostomum (Trematoda) parasites. *Parasitology*,

133(02), 229. https://doi.org/10.1017/S0031182006000242

Kim, K. S., Ejima, K., Iwanami, S., Fujita, Y., Ohashi, H., Koizumi, Y., Asai, Y., Nakaoka,

S., Watashi, K., Aihara, K., Thompson, R. N., Ke, R., Perelson, A. S., & Iwami, S.

(2021). A quantitative model used to compare within-host SARS-CoV-2, MERS-

CoV, and SARS-CoV dynamics provides insights into the pathogenesis and

treatment of SARS-CoV-2. *PLOS Biology*, *19*(3), e3001128.

https://doi.org/10.1371/journal.pbio.3001128

Kirk, D., Jones, N., Peacock, S., Phillips, J., Molnár, P. K., Krkošek, M., & Luijckx, P.

(2018). Empirical evidence that metabolic theory describes the temperature

dependency of within-host parasite dynamics. *PLOS Biology*, *16*(2), e2004608.

https://doi.org/10.1371/journal.pbio.2004608

Lafferty, K. D. (2009). The ecology of climate change and infectious diseases. *Ecology*,

90(4), 888–900. https://doi.org/10.1890/08-0079.1

Laprise, F., & Dodson, J. (1994). Environmental variability as a factor controlling spatial

patterns in distribution and species diversity of zooplankton in the St. Lawrence

Estuary. *Marine Ecology Progress Series*, *107*, 67–81.

https://doi.org/10.3354/meps107067

Lawson, C. R., Vindenes, Y., Bailey, L., & van de Pol, M. (2015). Environmental

variation and population responses to global change. *Ecology Letters*, *18*(7),

724–736. https://doi.org/10.1111/ele.12437

Lips, K. R., Diffendorfer, J., Mendelson, J. R., & Sears, M. W. (2008). Riding the Wave:

Reconciling the Roles of Disease and Climate Change in Amphibian Declines.

PLoS Biology, *6*(3), e72. https://doi.org/10.1371/journal.pbio.0060072

Luo, J., Greif, S., Ye, H., Bumrungsri, S., Eitan, O., & Yovel, Y. (2021). Flight rapidly

modulates body temperature in freely behaving bats. *Animal Biotelemetry*, *9*(1),

45. https://doi.org/10.1186/s40317-021-00268-6

Molnár, P. K., Kutz, S. J., Hoar, B. M., & Dobson, A. P. (2013). Metabolic approaches to

understanding climate change impacts on seasonal host-macroparasite

dynamics. *Ecology Letters*, *16*(1), 9–21. https://doi.org/10.1111/ele.12022

Molnár, P. K., Sckrabulis, J. P., Altman, K. A., & Raffel, T. R. (2017). Thermal

Performance Curves and the Metabolic Theory of Ecology—A Practical Guide to

Models and Experiments for Parasitologists. *Journal of Parasitology*, *103*(5), 423.

https://doi.org/10.1645/16-148

Munguia, P., Osman, R., Hamilton, J., Whitlatch, R., & Zajac, R. (2010). Modeling of

priority effects and species dominance in Long Island Sound benthic

communities. *Marine Ecology Progress Series*, *413*, 229–240.

https://doi.org/10.3354/meps08764

Ostfeld, R. S., & Keesing, F. (2012). Effects of Host Diversity on Infectious Disease.

Annual Review of Ecology, Evolution, and Systematics, *43*(1), 157–182.

https://doi.org/10.1146/annurev-ecolsys-102710-145022

Paaijmans, K. P., Blanford, S., Bell, A. S., Blanford, J. I., Read, A. F., & Thomas, M. B.

(2010). Influence of climate on malaria transmission depends on daily

temperature variation. *Proceedings of the National Academy of Sciences*,

107(34), 15135–15139. https://doi.org/10.1073/pnas.1006422107

Paaijmans, K. P., Heinig, R. L., Seliga, R. A., Blanford, J. I., Blanford, S., Murdock, C.

C., & Thomas, M. B. (2013). Temperature variation makes ectotherms more

sensitive to climate change. *Global Change Biology*, *19*(8), 2373–2380.

https://doi.org/10.1111/gcb.12240

Patz, J. A., Daszak, P., Tabor, G. M., Aguirre, A. A., Pearl, M., Epstein, J., Wolfe, N. D.,

Kilpatrick, A. M., Foufopoulos, J., Molyneux, D., Bradley, D. J., & Members of the

Working Group on Land Use Change Disease Emergence. (2004). Unhealthy

Landscapes: Policy Recommendations on Land Use Change and Infectious

Disease Emergence. *Environmental Health Perspectives*, *112*(10), 1092–1098.

https://doi.org/10.1289/ehp.6877

Paull, S. H., LaFonte, B. E., & Johnson, P. T. J. (2012). Temperature-driven shifts in a

host-parasite interaction drive nonlinear changes in disease risk. *Global Change*

Biology, *18*(12), 3558–3567. https://doi.org/10.1111/gcb.12018

Pedersen, A. B., & Fenton, A. (2007). Emphasizing the ecology in parasite community

ecology. *Trends in Ecology & Evolution*, *22*(3), 133–139.

https://doi.org/10.1016/j.tree.2006.11.005

Plowright, R. K., Parrish, C. R., McCallum, H., Hudson, P. J., Ko, A. I., Graham, A. L., &

Lloyd-Smith, J. O. (2017). Pathways to zoonotic spillover. *Nature Reviews*

Microbiology, *15*(8), 502–510. https://doi.org/10.1038/nrmicro.2017.45

Plowright, R., Reaser, J., Locke, H., Woodley, S. J., Patz, J. A., Becker, D., Oppler, G.,

Hudson, P., & Tabor, G. M. (2020). *A call to action: Understanding land use-*

induced zoonotic spillover to protect environmental, animal, and human health

[Preprint]. EcoEvoRxiv. https://doi.org/10.32942/osf.io/cru9w

Pounds, J. A., Fogden, M. P. L., & Campbell, J. H. (1999). Biological response to

climate change on a tropical mountain. *Nature*, *398*(6728), 611–615.

https://doi.org/10.1038/19297

Righetti, D., Vogt, M., Gruber, N., Psomas, A., & Zimmermann, N. E. (2019). Global

pattern of phytoplankton diversity driven by temperature and environmental

variability. *Science Advances*, *5*(5), eaau6253.

https://doi.org/10.1126/sciadv.aau6253

Rohr, J. R., Dobson, A. P., Johnson, P. T. J., Kilpatrick, A. M., Paull, S. H., Raffel, T. R.,

Ruiz-Moreno, D., & Thomas, M. B. (2011). Frontiers in climate change–disease

research. *Trends in Ecology & Evolution*, *26*(6), 270–277.

https://doi.org/10.1016/j.tree.2011.03.002

Rohr, J. R., & Raffel, T. R. (2010). Linking global climate and temperature variability to

widespread amphibian declines putatively caused by disease. *Proceedings of the*

National Academy of Sciences, *107*(18), 8269–8274.

https://doi.org/10.1073/pnas.0912883107

Rohr, J. R., Raffel, T. R., Romansic, J. M., McCallum, H., & Hudson, P. J. (2008).

Evaluating the links between climate, disease spread, and amphibian declines.

Proceedings of the National Academy of Sciences, *105*(45), 17436–17441.

https://doi.org/10.1073/pnas.0806368105

Roxburgh, S. H., Shea, K., & Wilson, J. B. (2004). The intermediate disturbance

hypothesis: Patch dynamics and mechanisms of species coexistence. *Ecology*,

85(2), 359–371. https://doi.org/10.1890/03-0266

Rynkiewicz, E. C., Pedersen, A. B., & Fenton, A. (2015). An ecosystem approach to

understanding and managing within-host parasite community dynamics. *Trends*

in Parasitology, *31*(5), 212–221. https://doi.org/10.1016/j.pt.2015.02.005

Saenz, R. A., Quinlivan, M., Elton, D., MacRae, S., Blunden, A. S., Mumford, J. A.,

Daly, J. M., Digard, P., Cullinane, A., Grenfell, B. T., McCauley, J. W., Wood, J.

L. N., & Gog, J. R. (2010). Dynamics of Influenza Virus Infection and Pathology.

Journal of Virology, *84*(8), 3974–3983. https://doi.org/10.1128/JVI.02078-09

Schoolfield, R. M., Sharpe, P. J. H., & Magnuson, C. E. (1981). Non-linear regression of

biological temperature-dependent rate models based on absolute reaction-rate

theory. *Journal of Theoretical Biology*, *88*(4), 719–731.

https://doi.org/10.1016/0022-5193(81)90246-0

Shurin, J. B., Winder, M., Adrian, R., Keller, W. (Bill), Matthews, B., Paterson, A. M.,

Paterson, M. J., Pinel-Alloul, B., Rusak, J. A., & Yan, N. D. (2010). Environmental

stability and lake zooplankton diversity—Contrasting effects of chemical and

thermal variability. *Ecology Letters*, *13*(4), 453–463.

https://doi.org/10.1111/j.1461-0248.2009.01438.x

Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A.,

Xia, Y., Bex, V., & Midgley, P. M. (Eds.). (2013). *Climate change 2013: The*

physical science basis. In: Contribution of Working Group I to the Fifth

Assessment Report of the Intergovernmental Panel on Climate Change.

Cambridge University Press.

Symons, C. C., & Arnott, S. E. (2014). Timing is everything: Priority effects alter

community invasibility after disturbance. *Ecology and Evolution*, *4*(4), 397–407.

https://doi.org/10.1002/ece3.940

Taylor, L. H., Latham, S. M., & woolhouse, M. E. J. (2001). Risk factors for human

disease emergence. *Philosophical Transactions of the Royal Society of London.*

Series B: Biological Sciences, *356*(1411), 983–989.

https://doi.org/10.1098/rstb.2001.0888

Tracy, A. M., Pielmeier, M. L., Yoshioka, R. M., Heron, S. F., & Harvell, C. D. (2019).

Increases and decreases in marine disease reports in an era of global change.

Proceedings of the Royal Society B: Biological Sciences, *286*(1912), 20191718.

https://doi.org/10.1098/rspb.2019.1718

Vasseur, D. A., DeLong, J. P., Gilbert, B., Greig, H. S., Harley, C. D. G., McCann, K. S.,

Savage, V., Tunney, T. D., & O'Connor, M. I. (2014). Increased temperature

variation poses a greater risk to species than climate warming. *Proceedings of*

the Royal Society B: Biological Sciences, *281*(1779), 20132612.

https://doi.org/10.1098/rspb.2013.2612

Vega-Rodriguez, M., Müller-Karger, F., Hallock, P., Quiles-Perez, G., Eakin, C., Colella,

M., Jones, D., Li, J., Soto, I., Guild, L., Lynds, S., & Ruzicka, R. (2015). Influence

of water-temperature variability on stony coral diversity in Florida Keys patch

reefs. *Marine Ecology Progress Series*, *528*, 173–186.

https://doi.org/10.3354/meps11268

Wright, R. K., & Cooper, E. L. (1981). Temperature effects on ectotherm immune

responses. *Developmental and Comparative Immunology*, *5*, 117–122.

905