1 **Environmental variability can promote parasite diversity within hosts and**
2 **transmission among hosts**
3

4 Madeline Jarvis-Cross (ORCID: 0009-0008-0527-9369)[1]
5 Martin Krkosek (ORCID: 0000-0001-7591 7954)[1]
6 [1]The University of Toronto, Department of Ecology and Evolutionary Biology
7 25 Willcocks Street, Toronto, ON, M5S 3B2
8 Corresponding author: Madeline Jarvis-Cross madeline.jarvis.cross@mail.utoronto.ca.

Abstract

While the mechanisms that govern disease emergence and spread among hosts are increasingly well-
described, the mechanisms that promote parasite diversity within-hosts, affecting host outcomes and
spillover potential, have been comparatively understudied. Furthermore, while attention has been paid to
the effects of increasing temperatures on disease systems, the effects of environmental variability have
been left underexplored, despite rising climatic variability. To investigate the impacts of environmental
variability on parasite diversity within-hosts, we analyzed a model of within-host population dynamics
wherein two parasites indirectly compete through the host's immune response. We simulated the model
under constant, demographically stochastic, environmentally stochastic, and demographically and
environmentally stochastic conditions, and analysed the viability and longevity of non-equilibrium
parasite co-occurrence. We found that environmental stochasticity increased the viability and longevity
of parasite co-occurrence, suggesting that thermal variability arising from climatic change and as a
physiological trait may promote parasite diversity within ectotherms and help explain bats' propensity to
support diverse communities of parasites. Further, we found that under certain conditions, the
transmissibility of co-occurring parasites can surpass the transmissibility of single parasites, suggesting
that thermal variability may increase the transmissibility of co-occurring parasites.

Key Words: within-host; co-infection; environmental variability; co-existence; modelling
28

Acknowledgements

30 This work was supported by The Emerging and Pandemic Infections Consortium Doctoral Award (MJ-
31 C), and Ontario Graduate Scholarship (MJ-C), a Natural Sciences and Engineering Research Council of
32 Canada Discovery Grant (MK), and a Canada Research Chair (MK).

Author Contributions

35 MJC and MK conceived the ideas and designed the methodology; MJC collected the data; MJC analysed
36 the data; MJC led the writing of the manuscript. All authors contributed critically to drafts and gave final
37 approval for publication.
38

Conflicts of Interest

40 The authors have declared that no conflicts of interest exist.
41

Data Availability Statement

43 Time series data available from the Federated Research Data Repository at
44 https://doi.org/10.20383/103.0764. Scripts and replication materials available from GitHub at
45 https://github.com/MadelineJC/EnvironmentalVariability_Repo.
46

Introduction

It has become clear that climate and land-use change can facilitate novel species interactions and increase contact rates between wild populations and humans, resulting in devastating epidemics and enabling spillover and disease emergence events among wild and human hosts alike (Daszak et al., 2000, 2001; Plowright et al., 2017). There is strong theoretical and empirical evidence that within-host parasite diversity influences disease outcomes in individual hosts (Bromenshenk et al., 2010; Johnson & Hoverman, 2012), transmissibility among hosts (Jolles et al., 2008; Susi et al., 2015), and community structure and functioning (Lafferty et al., 2006). However, the mechanisms that promote within-host parasite diversity have been understudied.

Ecological theory predicts a relationship between environmental variability and species diversity. For example, the Intermediate Disturbance Hypothesis (IDH) posits that environmental disturbances to competitive and successional processes will maximize species richness by preventing competitive exclusion (Chesson, 1994; Connell, 1978). In the case of apparent competition, analogized by immune suppression of two infecting parasites (Holt & Bonsall, 2017; Holt & Lawton, 1994), theory indicates that intermediate levels of disturbance will promote coexistence when a consumer has a larger impact on the stronger of two competitors (Wootton, 1998). More recent work suggests that within-host parasite diversity may be maintained by stabilizing and equalizing mechanisms (Bashey, 2015), including niche differentiation (Ramesh & Hall, 2023), competition-colonization trade-offs (T. A. Dallas et al., 2019), and environmental heterogeneity (Greenrod et al., 2024). Song and Spaak (2024) determined that in lower trophic levels, diversity is primarily constrained by fitness differences, which can be minimized via equalizing mechanisms like environmental variation (Song & Spaak, 2024; Spaak et al., 2021).

We considered the relationship between environmental variation and within-host parasite diversity along two axes: magnitude and frequency of stochastic disturbance (Mackey & Currie, 2001; Miller et al., 2011, 2012). First, we considered rising temperature variability as a feature of climate change (Lawson et al., 2015; Stocker et al., 2013; Vasseur et al., 2014). In ectothermic hosts, increasingly erratic

external conditions will affect the internal conditions of the host, thereby also affecting any parasites the

host may harbour (Cizauskas et al., 2017). A relationship between thermal variability and within-host

parasite diversity may be of particular interest when considered alongside the long-debated relationship

between climate change and disease-mediated amphibian declines (Daszak et al., 1999; Johnson, 2006;

Lips et al., 2008; Rohr et al., 2008). While it has been posited that increasing temperatures may drive

these declines, Rohr and Raffel (2010) demonstrated that El Niño events and associated regional

temperature variability increase host susceptibility, and thus contribute to host population declines

(Cohen et al., 2019; Rohr et al., 2008; Rohr & Raffel, 2010). Second, we considered environmental

variability as a host trait. Bats (order *Chiroptera*) have long been acknowledged as prolific reservoirs for

disease, and are unique among mammals in their ability to control body temperature in accordance with

activity level (e.g. torpor, flight) (Brook & Dobson, 2015; Calisher et al., 2006; Irving et al., 2021; Luo

et al., 2021). It has been theorized that thermal variation may contribute to the efficacy of bats' immune

systems, allowing them to carry microparasites without suffering increased morbidity (Fumagalli et al.,

2021; Luo et al., 2021). Similarly, thermal variability may affect parasite coexistence. In combination,

the effects of thermal variability on immune responses and parasite coexistence may help explain bats'

propensity to support large communities of parasites.

We used a model to investigate the impacts of environmental variability on the viability and

maintenance of nonequilibrium parasite co-occurrence within hosts. We began by adapting Antia et al.'s

(1994) model of within-host population dynamics to include two parasites that indirectly compete via the

host's immune system, and thus exhibit apparent competition (Holt, 1977; Råberg et al., 2006;

Williamson, 1957). We analyzed deterministic, demographically stochastic, environmentally stochastic,

and demographically-and-environmentally stochastic implementations of the model. Within the

environmentally stochastic implementations of the model, we draw on the Metabolic Theory of Ecology

to model the parasites' replication rates as temperature-dependent, and vary the size and frequency of

thermal disturbance within the system. Between and among the deterministic and stochastic

implementations of the model, we compared the ability of the secondary parasite to invade a system

containing the primary parasite, and the duration of time for which the primary and secondary parasites

co-occurred. Further, we compared the total transmission of co-occurring parasites with the total

transmission of a single parasite. We found that compared to a constant environment, a variable

environment tended to increase the ability of the secondary parasite to invade the system, facilitated

longer periods of co-occurrence between the primary and secondary parasites, and increased the

transmissibility of co-occurring parasites. These results (1) suggest that climatic changes may have

significant effects on parasite diversity within ectothermic hosts, (2) provide insight on a host's ability to

support large communities of parasites, and propose that variability within-hosts may influence

transmission dynamics among-hosts.

Methods

Model Description

Within-Host Population Dynamics Model

We began with a model of within host population dynamics based on Antia et al., (1994) that

represents the interaction between an infecting parasite and its host's immune system as analogous to the

interaction between a prey item and its predator. Per Antia et al. (1994), the density of the parasite

population, P, is determined by its replication rate, r, and the rate at which it is cleared by the host's

immune system, k. The model assumes that the immune system's replication rate is proportional to

parasite density at low parasite densities, and saturates at high parasite densities, allowing the host's

immune system to either extirpate or succumb to the infection (Antia et al., 1994). We adapted the model

to include a second parasite, P_j, that indirectly competes with the primary parasite, P_i, through the host's

immune response (Eqns. 1-3, Table 1). As such, the model describes an immune-mediated apparent

competition (Holt, 1977; Råberg et al., 2006; Williamson, 1957). Parameters values were not chosen to

describe a specific system, but to produce the dynamical behaviour described above and shown in Figure

1. Given that any number of parameter value combinations may produce this behaviour, we chose to

maintain the initial conditions and select parameter values proposed by Antia et al. (1994). However, we

increased the rate at which parasites are destroyed by the host's immune system (k) and decreased the

maximum replication rates of the primary and secondary parasites (r_i, r_j) to favour extirpation, and

eliminate host death as an infection outcome (Table 1). We also conducted a sensitivity analysis to

determine how variations in (1) the value of k and (2) the shape of the relationship between parasite

replication rate and the temperature of the system would affect our results, and found the model to be

robust (see pages 2, 4-8 of supplement for complete results).

129
130   (1) $P_i(t + 1) = (1 + r_i - (1 - e^{-kI(t)}))P_i(t)$
131   (2) $P_j(t + 1) = (1 + r_j - (1 - e^{-kI(t)}))P_j(t)$
132   (3) $I(t + 1) = (1 + \rho \left(\dfrac{P_i(t)}{P_i(t) + \phi} + \dfrac{P_j(t)}{P_j(t) + \phi} \right))I(t)$

133
134   **Table 1:** Parameter definitions and values.

Symbol	Parameter	Values Stated in Antia et al. (1994)	Values Used
P_i and P_j	Parasite abundance	1*	1*
I	Host immune cell abundance	1*	1*
r_i and r_j	Replication rates of parasites	0.1 – 10.0	0.1 – 2.0[†]
k	Rate at which parasites are destroyed by host's immune system	10^{-3}	10^{-2}
ρ	Maximum replication rate of host's immune system	1.0	1.0
ϕ	Parasite density at which replication rate of host's immune system is half its maximum	10^3	10^3
D	Lethal within-host parasite density	10^9	10^9

135   * Initial conditions
136   [†] Dependent on the temperature of system in environmentally stochastic simulations
137

To compare the impacts of different types of stochasticity on parasite diversity, we wrote deterministic

(D), demographically stochastic (S_D), environmentally stochastic (S_E), and demographically-and-

environmentally stochastic (S_DE) implementations of the model. Within the demographically stochastic

implementation of the model, we wrote the model's replication terms ($r_i \cdot P_i(t)$, $r_j \cdot P_j(t)$, and

$\rho \left(\frac{P_i(t)}{P_i(t)+\phi} + \frac{P_j(t)}{P_j(t)+\phi} \right) I(t))$ as Poisson processes, and mortality term $(1 - e^{-kI(t)})$ as a binomial process

(Eqns. 4-6).

(4) $P_i(t+1) = P_i(t) + B_{P_i}(t) - D_{P_i}(t)$

(5) $P_j(t+1) = P_j(t) + B_{P_j}(t) - D_{P_j}(t)$

(6) $I(t+1) = I(t) + B_I(t)$

In Eqns. (4) and (5), $B_{P_i}(t)$ and $B_{P_j}(t)$ are Poisson random variables such that $B_{P_i}(t) \sim Poisson(\lambda_i)$,

where $\lambda_i = r_i \cdot P_i(t)$, and $B_{P_j}(t) \sim Poisson(\lambda_j)$, where $\lambda_j = r_j \cdot P_j(t)$. In Eqn. (6), $B_I(t)$ is a Poisson

random variable such that $I_B(t) \sim Poisson(\lambda)$, where $\lambda = \rho \left(\frac{P_i(t)}{P_i(t)+\phi} + \frac{P_j(t)}{P_j(t)+\phi} \right) I(t)$. In Eqns. (4) and

(5), $D_{P_i}(t)$ and $D_{P_j}(t)$ are binomial random variables such that $D_{P_i}(t) \sim Binomial(P_i(t), p)$, where $p =$

$1 - e^{-kI(t)}$, and $D_{P_j}(t) \sim Binomial(P_j(t), p)$. To write the environmentally stochastic implementation

of the model, we used randomly generated temperature series to dictate the temperature of the system

(T), and allowed the replication rates of each parasite (r_i, r_j) to vary in response (Eqns. 7-9). We chose to

write r_i and r_j, though not k, as temperature-dependent to isolate the effect of thermal variability on

traits that can be metabolically scaled, and constrain the scope of the work. We did not write ρ or ϕ as

temperature-dependent to constrain the scope of the work.

(7) $P_i(t+1) = (1 + r_i(T(t)) - (1 - e^{-kI(t)}))P_i(t)$

(8) $P_j(t+1) = (1 + r_j(T(t)) - (1 - e^{-kI(t)}))P_j(t)$

(9) $I(t+1) = (1 + \rho \left(\frac{P_i(t)}{P_i(t) + \phi} + \frac{P_j(t)}{P_j(t) + \phi} \right)) I(t)$

We used first-order autocorrelated Gaussian noise to generate temperature series. We initiated the series

with a value of zero, and generated the next value by multiplying the previous value by the coefficient of

autocorrelation (0.25), and summing the resulting value with a value drawn from a normal random

variable with a mean of zero. We varied the standard deviation of the normal random variable to control

the size of disturbance, as described below. We then added the average temperature of the system to each

value in the series. Resulting values outside the thermal range of the system were re-drawn. We then

converted the temperature series into replication rates using various parameterizations of the Sharpe-

Schoolfield sub-model, as described below (Eqns. S1.1, S1.2; Table S2).

Sharpe-Schoolfield Sub-Model

The Metabolic Theory of Ecology (MTE) predicts how metabolic rates control ecological

processes across levels of organization, and explains how metabolic rates vary with body size and

temperature (Brown et al., 2004). Building off the MTE, the Boltzmann-Arrhenius (BA) relation

describes metabolism as a function of temperature. The Sharpe-Schoolfield model characterizes the

performance of a trait as a function of temperature through a thermal performance curve (TPC)

(Schoolfield et al., 1981). When applied to disease systems, the Sharpe-Schoolfield model has been

shown to describe relationships between temperature and species traits, and has proven useful in

investigating the relationship between rising temperatures and disease transmission.

We used various parameterizations of the Sharpe-Schoolfield model to generate paired thermal

performance curves relating the temperature of the system to the replication rates of two parasites (r_i and

r_j) (Fig. S9). We generated four paired thermal performance curves, representing four thermal scenarios:

(1) the identical responses scenario, (2) the offset responses scenario, (3) the P_i-advantage scenario, and

(4) the P_j-advantage scenario (Fig. 1, Table S2). The identical responses scenario describes r_i and r_j as

responding identically to the temperature of the system, and exhibits an average temperature of 19.9°C

(Fig. 1, Table S2, Fig. S9). The remaining scenarios describe a system within which P_i exhibits a higher

replication rate at lower temperatures and P_j exhibits a higher replication rate at higher temperatures, and

are differentiated by the average temperature of the system. The offset responses scenario exhibits an

average temperature of 19.9°C. At this temperature, the replication rate of P_i is equal to that of P_j (Fig. 1,

Table S2, Fig. S9). The P_i-advantage scenario exhibits an average temperature of 18°C, giving P_i a

thermal advantage over P_j (Fig. 1, Table S2, Fig. S9). We considered two versions, (I) and (II), of the P_j-

advantage scenario, each of which give P_j a thermal advantage over P_i. (I) exhibits an average

temperature of 20.9ºC, such that P_j's thermal advantage is comparable in size to that of P_i, as described

above (Fig. 1, Table S2, Fig. S9). (II) replicates the P_i-advantage scenario, but re-assigns the low-

temperature curve to P_j and the high-temperature curve to P_i, to isolate the effects of asymmetry (Fig.

S9). Given that (I) and (II) produced qualitatively comparable results, we will only discuss (I) in the

main text (see supplement for complete results).

Given that the Sharpe-Schoolfield model cannot generate negative values, it cannot be used to

explore cases in which extreme temperatures result in parasite mortality and population decline. To

compensate, we developed an alternative set of paired thermal performance curves, through which

extreme temperatures produce "negative" replication rates, and spur population decline (see "Extended

analysis" in supplement; Fig. S19).

Model Simulations

Within each thermal scenario, we simulated each model (D, S_D, S_E, S_{DE}) for fifty days.

Simulating for fifty days proved sufficient to capture the full course of parasite infection and clearance

across all treatments. We began each set of simulations with the primary parasite and the host's immune

system in the system, at initial abundances of one. We then considered all cases for the introduction of

the secondary parasite from its introduction at the onset of infection, to its introduction on the forty-ninth

day of infection.

For each environmentally stochastic model (S_E, S_{DE}) simulation, we generated a new temperature

series and two corresponding series of replication rates. When determining how new temperatures would

be selected, we considered two axes of thermal disturbance: size and frequency. We controlled the size

of disturbance by changing the standard deviation of the normal random variable in the temperature

selection function, starting from two and increasing by two until the effect was fully characterized at a

value of thirty, resulting in fifteen sizes of disturbance (Fig. 1B). We considered standard deviations

between two and ten to denote small disturbances, between twelve and twenty to denote intermediately

sized disturbances, and between twenty-two and thirty to denote large disturbances (Fig. 1B). To control

the frequency of disturbance, we re-assigned the temperature of the system every ten, nine, eight, seven,

six, five, four, three, and two days, once per day, and two, five, and ten times per day to fully

characterise the effect, resulting in thirteen frequencies of disturbance (Fig. 1B). We considered re-

assignments every ten to seven days to denote infrequent disturbances, every six to two days to denote

intermediately frequent disturbances, and every day to ten times per day to denote frequent disturbances

(Fig. 1B). Considering size and frequency together resulted in 195 unique disturbance regimes. We

conducted 1000 simulations per stochastic model and treatment (Fig. 1A). Two hundred such

simulations are shown in Figures 1A and S10.

Co-Occurrence Analyses

Given successful invasion by the secondary parasite $(P_j(t+1) > P_j(t))$, we defined longevity of

co-occurrence as the amount of time for which both parasites exhibited population abundances equal to

or greater than one.

Figure 1: (A) Shows a schematic detailing simulation experiment. When simulating each of four models (deterministic,

demographically stochastic, environmentally stochastic, and demographically and environmentally stochastic), we considered

four thermal scenarios (identical responses, offset responses, P_i-advantage, and P_j-advantage). When simulating the

environmentally stochastic models (in blue), we considered 15 sizes and 13 frequencies of disturbance, totalling 195 unique

disturbance regimes. We additionally considered 50 days on which the secondary parasite could be introduced. We conducted

1000 simulations per stochastic model and treatment. Examples of such simulations are shown as time series. (B) Shows a

schematic detailing how size and frequency interact to produce disturbance regimes.

We summarised these results as distributions for each stochastic model and treatment. Using the S_D

model as a baseline, we investigated the effects of environmental variability on the viability and

maintenance of parasite diversity by comparing the ability of the secondary parasite to invade a single-

parasite system, and the duration of time for which the primary and secondary parasites co-occurred

across stochastic treatments. Additionally, we analysed (1) the proportion of stochastic simulations per

model and treatment that resulted in longer periods of co-occurrence relative to the deterministic model,

and (2) the competitive outcomes of each treatment to gain insight on underlying mechanisms.

Transmissibility Analyses

To explore the impacts of co-infection on transmission potential, we compared the lifetime

replicative outputs of co-occurring parasites with those of a single parasite in constant and variable

environments. As per Antia et al., (1994), we determined the total transmission (q_i, q_j, q, where q is the

total transmission of a single parasite) of a parasite (P_i, P_j, P, where P is a single parasite) from an

infected host by taking the integral of the rate of transmission (u) of each parasite from the host over the

duration of infection (Eqns. 10.1-10.3). For simplicity, we set the rate of transmission to one.

(10.1) $\quad q_i = \int_0^\infty u P_i(t) dt$

(10.2) $\quad q_j = \int_0^\infty u P_j(t) dt$

(10.3) $\quad q = \int_0^\infty u P(t) dt$

256
To compare constant and variable environments, we analysed multi-parasite and single-parasite versions

of the S_D and S_{DE} models.

Results

Analytically, the secondary parasite could invade ($P_j(t+1) > P_j(t)$) when $I(t) < \frac{-\ln(1-r_j)}{k}$. Our

parameterization of the S_D model allowed the secondary parasite to invade when introduced before the

tenth to twelfth day of infection, resulting in 8.4 to 9.2 days of co-occurrence, depending on thermal

scenario (Fig. S11, Supplementary Datafile 1-1:10).

Co-occurrence Analysis

Simulation outputs show that compared to a deterministic baseline, the addition of demographic

stochasticity has little effect on co-occurrence (Supplementary Datafile 2-1:10), while the addition of

thermal variability increases the ability of the secondary parasite to invade the system and facilitates

longer periods of co-occurrence between parasites. While longevities of co-occurrence are maximised

under the P_j-advantage scenario, the difference between deterministic and environmentally stochastic

implementations is not consistently maximised by any one thermal scenario. Additionally, while larger

and more frequent disturbances tend to promote longer periods of co-occurrence, increasingly large

disturbances produce a less steep effect than do increasingly frequent disturbances. These findings are

common among thermal scenarios.

When both parasites were introduced simultaneously, S_D model simulations resulted in periods of

co-occurrence that were, on average, 4.5% to 3.1% shorter (0.4 to 0.3 days) than deterministically

simulated periods of co-occurrence (Figs. 2, S12). In comparison, simulations of the environmentally

stochastic models (S_E, S_{DE}) resulted in periods of co-occurrence that were up to 46.2% (S_E) and 36.0%

(S_{DE}) longer (3.9 (S_E) and 3.0 (S_{DE}) days) than deterministically simulated periods of co-occurrence

(Figs. 2, S12). Across all 195 disturbance regimes, simulations of the S_E model resulted in mean periods

of co-occurrence that were between 2.7% shorter (0.2 days) and 46.2% longer (3.9 days) than their

deterministic counterparts (Figs. 2, S12). The identical responses scenario promoted the shortest and

longest mean periods of co-occurrence (Figs. 2-3, S12-S13). Simulations of the S_{DE} model resulted in

mean periods of co-occurrence that were between 6.8% shorter (0.6 days) and 36.0% longer (3.0 days)

than deterministically simulated periods of co-occurrence (Figs. 2, S12). Again, the identical responses

scenario promoted the shortest and longest mean periods of co-occurrence (Figs. 2-3, S12-S13). Across

thermal scenarios, mean periods of co-occurrence were shortest when disturbances were large (standard

deviation of the normal random variable in the temperature autocorrelation function = 22 to 30) and

infrequent (once every 7 to 9 days) (Figs. 3, S13). Mean periods of co-occurrence were longest when

disturbances were large (SD = 24 to 30) and frequent (five to ten times per day) (Figs. 3, S13).

More generally, as disturbances increased in size, periods of co-occurrence increased in length

towards a saturation point (standard deviation of the normal random variable in the temperature

autocorrelation function ≈ 24-30, depending on thermal scenario) (Figs. S14, S18, Supplementary

Datafiles 3-1:5, 4-1:5). However, an analysis of simulations within which an alternative set of TPCs

allowed for extreme temperatures to cause population decline revealed a parabolic relationship between

disturbance size and longevity of co-occurrence, that peaked when disturbances were intermediately

sized (standard deviation of the normal random variable in the temperature autocorrelation function ≈ 8-

16) (Fig. S20).

When disturbances were intermediately frequent (once every two to six days), periods of co-

occurrence were similar in length to deterministically simulated periods of co-occurrence. Interestingly,

when disturbances were infrequent (once every six to ten days), mean periods of co-occurrence were

shorter than deterministically simulated periods of co-occurrence (Fig. S14, S21). Per the deterministic

model, P_i and P_j can only co-occur for 8.4 to 9.2 days. Thus, a system subjected to infrequent

disturbances is functionally similar to the deterministic system, making the outcome of infection reliant

on initial conditions. When disturbances were frequent (ten times per day to once per day), mean periods

of co-occurrence were longer than deterministically simulated periods of co-occurrence. Within a given

disturbance regime, longevities of co-occurrence were normally distributed (Figs. S14-S15, S23).

Figure 2: The mean percent difference in longevity of co-occurrence relative to deterministically simulated periods of co-

occurrence, when both parasites are introduced at the same time. Single points above "S_D" labels represent the mean percent

difference in longevity of co-occurrence relative to deterministic thresholds across 1000 demographically stochastic

simulations. Each point above "S_E"/"S_{DE}" labels represents the mean percent difference in longevity of co-occurrence relative

to deterministic thresholds across 1000 environmentally stochastic simulations. As such, there are 195 points above each

"S_E"/"S_{DE}" label, corresponding to 195 environmental disturbance regimes.

Rarity and Competitive Outcomes

When the secondary parasite was introduced before or on the invasion threshold (10th or 12th day

of infection, depending on thermal scenario), 29%-71% (S_E) and 46%-58% (S_{DE}) of environmentally

stochastic simulations resulted in longer periods of co-occurrence relative to deterministic simulations

(Fig. 4). When the secondary parasite was introduced after the invasion threshold, the likelihood that an

environmentally stochastic simulation resulted in some period of co-occurrence sharply declined. When

the secondary parasite was introduced one to twelve days after the invasion threshold, 1%-31% (S_E) and

1%-46% (S_{DE}) of environmentally stochastic simulations resulted in some period of co-occurrence (Fig.

4). Thus, environmental variability extended the invasion threshold by 50%-75% (S_E) and 100%-120%

(S_{DE}), depending on thermal scenario (Fig. 4).

Figure 3: Fully factorial representations of the impacts of varyingly sized and timed disturbances on the longevity of co-

occurrence. The plot shows the mean number of time steps (10 time steps/day) for which P_i and P_j have population

abundances greater than one as per the S_{DE} model. The circled numbers denote the number of time steps for which P_i and P_j

can co-occur, per the deterministic model.

Analyzing competitive outcomes at intermediate levels of disturbance (SD = 10, frequency of

disturbance = once per day) revealed that under the offset responses scenario, when P_i and P_j were

introduced at the same time, P_j outlasted P_i 65.5% (S_E) and 63.4% (S_{DE}) of the time (Fig. S22-S23).

Additionally, under the P_i-advantage scenario, when P_i and P_j were introduced at the same time, P_i and

P_j were equally likely to outlast one another (P_j outlasts P_i 49.9% (S_E) and 51.0% (S_{DE}) of the time), but

under the P_j-advantage, P_j was much more likely to outlast P_i (P_j outlasts P_i 76.6% (S_E) and 70.4% (S_{DE})

of the time) (Fig. S21-S23). Across thermal scenarios, when the secondary parasite, P_j, was introduced in

the middle of the infection period, P_i was more likely to outlast P_j. When P_j was introduced late in the

infection period, P_j was more likely to outlast P_i (Fig. S22-S23). Regardless of when P_j was introduced,

simulations resulting in especially long periods of co-occurrence are equally likely to show either

parasite outlasting the other (Fig. S22-S23).

Figure 4: Proportion of environmentally stochastic simulations (SD = 10, frequency of disturbance = once per day) resulting

in longer periods of co-occurrence relative to deterministic simulations by the day on which the secondary parasite was

introduced. Vertical lines show the first day on which the introduction of the secondary parasite does not result in any co-

occurrence between the two parasites, as per the deterministic model (D).

Transmissibility

Across thermal scenarios, simulations of the multi-parasite S_D model produced lifetime

replicative outputs (LROs) that were one to two orders of magnitude lower than those produced by

simulations of the single-parasite S_D model (Supplementary Datafiles 5-1:16). Similarly, under the first

three thermal scenarios, simulations of the multi-parasite S_{DE} model produced LROs that were one to

two orders of magnitude lower than those produced by simulations of the single-parasite S_{DE} model.

However, under the P_j-advantage scenario and among 62 of 195 (32%) disturbance regimes, the average

transmission potential of the co-occurring secondary parasite, P_j, exceeded that of a single parasite (Fig.

5). Among these disturbance regimes, the transmission potential of P_j exceeded that of a single parasite

by an average of 2% to 135% (Supplementary Datafiles 5-8). The transmission potentials of co-

occurring parasites peaked when disturbances were infrequent (once every ten days) to intermediately

frequent (once every six days) (Fig. S24). While intermediately large and frequent disturbances

maximally increased longevities of co-occurrence, such infections did not result in particularly high

transmission potentials, indicating that especially long periods of co-occurrence may be facilitated by

lower lifetime replicative outputs that do not trigger a large immune response, and thus allow for more

persistent, but less transmissible, infections.

Figure 5: The impacts of environmental variability on transmission potential under the P_j-advantage scenario. The plot shows

a fully factorial representation of the impacts of thermal disturbance on the transmission potential of P_j, relative to the

equivalent single-parasite treatment. The black line indicates disturbance regimes under which the average transmission

potential of P_j was equal to that of a single parasite. Colours warmer than denoted by the black line indicate disturbance

regimes under which the average transmission potential of P_j exceeded that of a single parasite; colours cooler than denoted

by the black line indicate the opposite.

Discussion

We investigated the potential for environmental variability to affect the viability and maintenance

of within-host parasite diversity. Our work provides theoretical evidence that relative to a constant

environment, a variable environment can increase the ability of a secondary parasite to invade a single-

parasite system, promote longer periods of co-occurrence between parasites, and increase the

transmissibility of co-occurring parasites. In more detail, thermal disturbances of various sizes and

frequencies allowed a secondary parasite to invade a single-parasite system up to twelve days after the

deterministic invasion threshold, extending the threshold by 100% to 120% (S$_{DE}$ model, Fig. 4). Large

and frequent thermal disturbances enabled parasites to co-occur for up to 37.8% longer, relative to a

constant environment (S$_{DE}$ model, Figs. 2, S12). Finally, under the P_j-advantage scenario and among

32% disturbance regimes, the average transmission potential of the co-occurring secondary parasite

exceeded that of a single parasite (S$_{DE}$ model, Fig. 5).

Across thermal scenarios, the relationship between disturbance size and longevity of co-

occurrence was initially positive and eventually saturating (Fig. S14, S18). Large disturbances produced

a saturating response rather than the expected parabolic response due to the structure of the Sharpe-

Schoolfield sub-model. Allowing for extreme temperatures to result in parasite mortality and population

decline produced a parabolic response, implying that intermediate sizes of disturbance maximised the

viability and longevity of parasite co-occurrence (Figs. S19-S20). When thermal disturbances were

small, the parasite with the higher initial replication rate was more likely to grow quickly and trigger a

commensurate immune response that would competitively exclude the other. As thermal disturbances

increased in size, it became increasingly likely that when the temperature of the system changed, the

competitive advantage would shift from one parasite to the other, minimizing fitness differences over the

course of an infection and promoting co-occurrence. However, per the extended analysis, large thermal

disturbances caused parasite populations to alternatively explode and crash, triggering a sizable immune

response, increasing the likelihood of extirpation, and precluding co-occurrence.

The relationship between climate change and disease-mediated amphibian declines has long been debated. While some argue that climatic changes may impede immune function or result in performance gaps between parasites and their hosts, making amphibians more susceptible to infection, others argue that insufficient evidence should preclude a causal link (Cohen et al., 2019; Rohr et al., 2008). While much of the literature proposes mean-climate hypotheses, some evidence suggests that intensifying El Niño events are increasing thermal variability, which may in turn affect amphibians' susceptibility to or ability to defend against infection (Rohr & Raffel, 2010). Congruently, our findings provide theoretical evidence that thermal variability promotes the viability and maintenance of within-host parasite diversity, and thus suggest that as the climate continues to change, attention should be paid to shifts in microparasite communities and their impacts on ectothermic hosts. In particular, well-documented climate-driven increases in disease incidence among a wide variety of marine organisms may constitute an important starting point for future work (Bates et al., 2009; Bruno et al., 2007; Tracy et al., 2019).

Across thermal scenarios, low and high frequencies of disturbance impeded and promoted the viability and longevity of parasite co-occurrence, respectively (Figs. 3, S13-S14, S21). Given the transient nature of infection specified by the underlying model, infrequent disturbances (once every seven to ten days) resulted in the temperature of the system being disturbed towards the end of the infection period, when parasite populations were small and especially susceptible to unfavourable conditions. Conversely, frequent disturbances (ten times per day to once per day) resulted in the temperature of the system being disturbed throughout the duration of the infection period, and as such, neither favourable nor unfavorable conditions lasted long enough for one parasite to gain a decisive advantage over and exclude the other, and thus promoted extended periods of co-occurrence. While it has been theorized that internal temperature variation may aid bats' immune responses, allowing them to carry microparasites without suffering outsized morbidity (Fumagalli et al., 2021), these findings suggest that internal temperature variation may also facilitate parasite co-occurrence within bats, and thus may help explain bats' propensity to support diverse communities of parasites.

Under the offset responses scenario, P_j often outlasted P_i, while under the P_i-advantage scenario,

P_i and P_j were equally likely to outlast each other (Fig. S22-S23). Under the P_j-advantage scenario, P_j

was much more likely to outlast P_i (Fig. S22-S23). These outcomes are consequences of the

asymmetrical structure of the thermal performance curves, which made unfavourable conditions more

disadvantageous for P_i than P_j (Sinclair et al., 2016). Further, regardless of when P_j was introduced,

simulations resulting in especially long periods of co-occurrence were equally likely to show either

parasite outlasting the other. Taken together, these results suggest that to predict the competitive

outcome of a multiple infection, one may need to consider the interplay between how well adapted a

parasite is to its host, priority effects, and the responses of biological rates to environmental factors.

While one might expect a specialist or well-adapted parasite to exploit its host more efficiently than a

generalist or novel parasite, the results of this work suggest that environmental variability can even the

scales, and allow a generalist or novel parasite to establish and persist, as predicted by the intermediate

disturbance hypothesis (Chesson, 1994; Connell, 1978; Roxburgh et al., 2004). Similarly, while one

might additionally expect a colonizing parasite to benefit from an initial lack of competition, the results

of this work again suggest that environmental variability and disturbance may contribute to the

structuring of communities by allowing latecomers to establish and persist more often and for longer

than they otherwise would (Jiang & Patel, 2008; Symons & Arnott, 2014).

While the metabolic theory of ecology posits that certain biological rates (e.g. metabolic rate,

replication/growth rate, etc.) will exhibit an asymmetrical and unimodal response to temperature, other

vital rates may exhibit symmetrical or bimodal responses to temperature (Dell et al., 2014; Molnár et al.,

2017; Sinclair et al., 2016). This work suggests that the shape of a thermal performance curve can

meaningfully affect the nature and outcome of an infection. The shape of a thermal performance curve

can also dictate the effects of stochasticity on persistence. Here, thermal variation pushes the system

away from a parasite's optimal temperature, and thus, away from its maximum replication rate. In

accordance with Jensen's inequality, compared to a deterministic system, the introduction of

stochasticity results in a lower average population growth rate (Denny, 2017; Jensen, 1906; Vasseur et

al., 2014). In the absence of thermal dependency, the introduction of stochasticity will also result in a

lower average population growth rate over time (Tuljapurkar, 1990). In this system, lower average

parasite replication rates slow down the infection and subsequently, the immune response, allowing

parasites to co-occur for longer. As anthropogenic changes alter host-parasite interaction networks,

understanding how evolutionary and environmental factors combine to affect interactions between

parasites will be essential in predicting and managing spillover and emergence in wildlife and human

populations (Paull et al., 2012; Pedersen & Fenton, 2007).

Developing an understanding of how the impacts of environmental variability on parasite

diversity within hosts may scale to affect among-host processes and events, including transmission,

spillover, and emergence will be essential to the maintenance of wildlife, human, and ecosystem health

(Allocati et al., 2016; Halpin et al., 2000; Johnson, 2006). While increased within-host parasite diversity

provides more strains or species of a parasite or parasites an opportunity to transmit, within-host

competition may result in decreased parasite loads, thus decreasing the transmission probability of any

one strain or species, and altering expected transmission patterns (Johnson et al., 2013; Pedersen &

Fenton, 2007). Similarly, our findings provide theoretical evidence that when environmental variations

are large and frequent, within-host competition decreases the total transmission of co-occurring parasites.

As such, while large and frequent disturbances may maximally increase the amount of time for which

two parasites co-occur, such infections do not result in particularly high transmission potentials,

indicating that especially long periods of co-occurrence may be facilitated by lower parasite replication

rates that do not trigger a large immune response, and thus allow for more persistent infections.

However, we found that approximately a third of tested disturbance regimes allowed the replicative

outputs of co-occurring parasites to exceed those of single parasites, indicating that the effects of

environmental variability may not be limited to the promotion of within-host parasite diversity, but may

also promote transmission between hosts (Fig. 5, S24). Disturbances of intermediate size and low to

intermediate frequency may constitute a sizable region within which environmental variation can

decrease fitness differences between parasites and promote within-host parasite diversity, while allowing

parasites to achieve sufficiently high parasite loads, resulting in more diverse and transmissible

infections (Fig. 5, S24).

While both Antia et al.'s (1994) model and our modified model should be broadly applicable to

an array of within-host multi-parasite systems, including multi-strain or cross-species, both models are

relatively simple as to remain generalizable, and thus cannot represent chronic infections, systems within

which parasites exhibit more complex interactions with their host or their host's immune system, nor

systems within which parasites directly interact with one another. As such, it may be interesting to

validate the theoretical insights of this work in an empirical system. This work is also limited by the

scope with which thermal dependency was considered. Given the thermal sensitivity of immune function

in ectotherms and endotherms, it may be worthwhile to consider the thermal dependency of immune

processes like response time and replication rate in tandem with the thermal dependency of parasite

replication rates (Butler et al., 2013; Wright & Cooper, 1981). Similarly, using nonlinear averaging to

assess the response of a nonlinear process to variability is contentious. While many theoretical works and

some empirical studies support the use of nonlinear averaging and find evidence for Jensen's inequality,

borne out as disproportionate effects of thermal variability on performance, many empirical studies have

contested an implicit assumption of nonlinear averaging: that traits should respond instantaneously to

change, without acclimation (Bernhardt et al., 2018; Estay et al., 2011; Krichel et al., 2023; Sinclair et

al., 2016). Thermal acclimation through phenotypic plasticity alters thermal performance curves, and as

such, produces a more complex relationship between temperature and biological rates that may not be

well-characterized by nonlinear averaging (Schulte et al., 2011; Sckrabulis et al., 2022). With regards to

host-parasite interactions, the temperature variability hypothesis posits that owing to their small relative

size, parasites should acclimate more quickly than their hosts, though the speed at which various

parasites may acclimate remains unclear (Raffel et al., 2013; Rohr et al., 2013). Interestingly, some

empirical studies of the impacts of environmental variability on host-parasite interactions have

considered thermal acclimation, yielding a range of conclusions including decreased pathogen

prevalence, reduced infection prevalence, and decreased resistance in hosts (T. Dallas & Drake, 2016;

Duncan et al., 2011; Raffel et al., 2013).

Ecological theory has long predicted that environmental variability supports species diversity,

from the IDH (Chesson, 1994; Connell, 1978; Huston, 2014; Roxburgh et al., 2004) to modern

coexistence theory (Barabás et al., 2018; Chesson, 1994, 2000, 2003; Chesson & Huntly, 1997). Work

on trophic interactions, wherein apparent competition is considered analogous to interactions between

the host's immune system and infecting parasites (Holt & Bonsall, 2017; Holt & Lawton, 1994),

indicates that intermediate levels of disturbance can promote coexistence (Wootton 1998) and that within

lower trophic levels (e.g. among competing parasites), diversity is primarily constrained by fitness

differences, which can be minimized via equalizing mechanisms like environmental variation (Song &

Spaak). Here, we provide theoretical evidence that the predicted relationship between environmental

variability and species diversity is borne out in within-host disease systems. Camus wrote that plagues,

despite their commonality, find us unprepared (Camus, Albert, 2008). Improving our understanding of

how environmental variability affects parasite diversity within- and between-hosts will expand our

understanding of the mechanisms that can facilitate disease spread and emergence, and allow us to make

robust predictions, and develop and implement mitigation strategies as we move into an increasingly

uncertain future. In brief, we will be found *less* unprepared.

References

Allocati, N., Petrucci, A. G., Di Giovanni, P., Masulli, M., Di Ilio, C., & De Laurenzi, V. (2016). Bat–man disease transmission: Zoonotic pathogens from wildlife reservoirs to human populations. *Cell Death Discovery*, *2*(1), 16048. https://doi.org/10.1038/cddiscovery.2016.48

Antia, R., Levin, B. R., & May, R. M. (1994). Within-Host Population Dynamics and the Evolution and Maintenance of Microparasite Virulence. *The American Naturalist*, *144*(3), 457–472. https://doi.org/10.1086/285686

Bashey, F. (2015). Within-host competitive interactions as a mechanism for the maintenance of parasite diversity. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *370*(1675), 20140301. https://doi.org/10.1098/rstb.2014.0301

Bates, A., Hilton, B., & Harley, C. (2009). Effects of temperature, season and locality on wasting disease in the keystone predatory sea star Pisaster ochraceus. *Diseases of Aquatic Organisms*, *86*, 245–251. https://doi.org/10.3354/dao02125

Bernhardt, J. R., Sunday, J. M., Thompson, P. L., & O'Connor, M. I. (2018). Nonlinear averaging of thermal experience
predicts population growth rates in a thermally variable environment. *Proceedings of the Royal Society B: Biological*
Sciences, *285*(1886), 20181076. https://doi.org/10.1098/rspb.2018.1076
Brook, C. E., & Dobson, A. P. (2015). Bats as 'special' reservoirs for emerging zoonotic pathogens. *Trends in Microbiology*,
23(3), 172–180. https://doi.org/10.1016/j.tim.2014.12.004
Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., & West, G. B. (2004). Toward a metabolic theory of ecology.
Ecology, *85*(7), 1771–1789. https://doi.org/10.1890/03-9000
Bruno, J. F., Selig, E. R., Casey, K. S., Page, C. A., Willis, B. L., Harvell, C. D., Sweatman, H., & Melendy, A. M. (2007).
Thermal Stress and Coral Cover as Drivers of Coral Disease Outbreaks. *PLoS Biology*, *5*(6), e124.
https://doi.org/10.1371/journal.pbio.0050124
Butler, M. W., Stahlschmidt, Z. R., Ardia, D. R., Davies, S., Davis, J., Guillette, L. J., Johnson, N., McCormick, S. D.,
McGraw, K. J., & DeNardo, D. F. (2013). Thermal Sensitivity of Immune Function: Evidence against a Generalist-
Specialist Trade-Off among Endothermic and Ectothermic Vertebrates. *The American Naturalist*, *181*(6), 761–774.
https://doi.org/10.1086/670191
Calisher, C. H., Childs, J. E., Field, H. E., Holmes, K. V., & Schountz, T. (2006). Bats: Important Reservoir Hosts of
Emerging Viruses. *Clinical Microbiology Reviews*, *19*(3), 531–545. https://doi.org/10.1128/CMR.00017-06
Chesson, P. (1994). Multispecies Competition in Variable Environments. *Theoretical Population Biology*, *45*(3), 227–276.
https://doi.org/10.1006/tpbi.1994.1013
Cizauskas, C. A., Carlson, C. J., Burgio, K. R., Clements, C. F., Dougherty, E. R., Harris, N. C., & Phillips, A. J. (2017).
Parasite vulnerability to climate change: An evidence-based functional trait approach. *Royal Society Open Science*,
4(1), 160535. https://doi.org/10.1098/rsos.160535
Cohen, J. M., Civitello, D. J., Venesky, M. D., McMahon, T. A., & Rohr, J. R. (2019). An interaction between climate change
and infectious disease drove widespread amphibian declines. *Global Change Biology*, *25*(3), 927–937.
https://doi.org/10.1111/gcb.14489
Connell, J. H. (1978). Diversity in Tropical Rain Forests and Coral Reefs: High diversity of trees and corals is maintained
only in a nonequilibrium state. *Science*, *199*(4335), 1302–1310. https://doi.org/10.1126/science.199.4335.1302
Dallas, T. A., Laine, A.-L., & Ovaskainen, O. (2019). Detecting parasite associations within multi-species host and parasite
communities. *Proceedings of the Royal Society B: Biological Sciences*, *286*(1912), 20191109.
https://doi.org/10.1098/rspb.2019.1109
Dallas, T., & Drake, J. M. (2016). Fluctuating temperatures alter environmental pathogen transmission in a *Daphnia* –
pathogen system. *Ecology and Evolution*, *6*(21), 7931–7938. https://doi.org/10.1002/ece3.2539
Daszak, P., Berger, L., Cunningham, A. A., Hyatt, A. D., Green, D. E., & Speare, R. (1999). Emerging Infectious Diseases
and Amphibian Population Declines. *Emerging Infectious Diseases*, *5*(6), 735–748.
https://doi.org/10.3201/eid0506.990601
Daszak, P., Cunningham, A. A., & Hyatt, A. D. (2000). Emerging Infectious Diseases of Wildlife—Threats to Biodiversity
and Human Health. *Science*, *287*(5452), 443–449. https://doi.org/10.1126/science.287.5452.443
Daszak, P., Cunningham, A. A., & Hyatt, A. D. (2001). Anthropogenic environmental change and the emergence of infectious
diseases in wildlife. *Acta Tropica*, *78*(2), 103–116. https://doi.org/10.1016/S0001-706X(00)00179-0
Dell, A. I., Pawar, S., & Savage, V. M. (2014). Temperature dependence of trophic interactions are driven by asymmetry of
species responses and foraging strategy. *Journal of Animal Ecology*, *83*(1), 70–84. https://doi.org/10.1111/1365-
2656.12081
Denny, M. (2017). The fallacy of the average: On the ubiquity, utility and continuing novelty of Jensen's inequality. *Journal*
of Experimental Biology, *220*(2), 139–146. https://doi.org/10.1242/jeb.140368
Duncan, A. B., Fellous, S., & Kaltz, O. (2011). Temporal variation in temperature determines disease spread and maintenance
in *Paramecium* microcosm populations. *Proceedings of the Royal Society B: Biological Sciences*, *278*(1723), 3412–
3420. https://doi.org/10.1098/rspb.2011.0287
Estay, S. A., Clavijo-Baquet, S., Lima, M., & Bozinovic, F. (2011). Beyond average: An experimental test of temperature
variability on the population dynamics of *Tribolium confusum*. *Population Ecology*, *53*(1), 53–58.
https://doi.org/10.1007/s10144-010-0216-7
Fumagalli, M. R., Zapperi, S., & La Porta, C. A. M. (2021). Role of body temperature variations in bat immune response to
viral infections. *Journal of The Royal Society Interface*, *18*(180), 20210211. https://doi.org/10.1098/rsif.2021.0211
Greenrod, S. T. E., Cazares, D., Johnson, S., Hector, T. E., Stevens, E. J., MacLean, R. C., & King, K. C. (2024). Warming
alters life-history traits and competition in a phage community. *Applied and Environmental Microbiology*, *90*(5),
e00286-24. https://doi.org/10.1128/aem.00286-24
Halpin, K., Young, P. L., Field, H. E., & Mackenzie, J. S. (2000). Isolation of Hendra virus from pteropid bats: A natural
reservoir of Hendra virus. *Journal of General Virology*, *81*(8), 1927–1932. https://doi.org/10.1099/0022-1317-81-8-
1927
Holt, R. D. (1977). Predation, apparent competition, and the structure of prey communities. *Theoretical Population Biology*,
12(2), 197–229. https://doi.org/10.1016/0040-5809(77)90042-9

Holt, R. D., & Bonsall, M. B. (2017). Apparent Competition. *Annual Review of Ecology, Evolution, and Systematics*, *48*(1), 447–471. https://doi.org/10.1146/annurev-ecolsys-110316-022628

Holt, R. D., & Lawton, J. H. (1994). THE ECOLOGICAL CONSEQUENCES OF SHARED NATURAL ENEMIES. *Annual Review of Ecology and Systematics*, *25*(1), 495–520. https://doi.org/10.1146/annurev.es.25.110194.002431

Huston, M. A. (2014). Disturbance, productivity, and species diversity: Empiricism vs. logic in ecological theory. *Ecology*, *95*(9), 2382–2396. https://doi.org/10.1890/13-1397.1

Irving, A. T., Ahn, M., Goh, G., Anderson, D. E., & Wang, L.-F. (2021). Lessons from the host defences of bats, a unique viral reservoir. *Nature*, *589*(7842), 363–370. https://doi.org/10.1038/s41586-020-03128-0

Jensen, J. L. W. V. (1906). Sur les fonctions convexes et les inégalités entre les valeurs moyennes. *Acta Mathematica*, *30*(0), 175–193. https://doi.org/10.1007/BF02418571

Jiang, L., & Patel, S. N. (2008). Community assembly in the presence of disturbance: A microcosm experiment. *Ecology*, *89*(7), 1931–1940. https://doi.org/10.1890/07-1263.1

Johnson, P. T. J. (2006). Amphibian diversity: Decimation by disease. *Proceedings of the National Academy of Sciences*, *103*(9), 3011–3012. https://doi.org/10.1073/pnas.0600293103

Johnson, P. T. J., Preston, D. L., Hoverman, J. T., & LaFonte, B. E. (2013). Host and parasite diversity jointly control disease risk in complex communities. *Proceedings of the National Academy of Sciences*, *110*(42), 16916–16921. https://doi.org/10.1073/pnas.1310557110

Jolles, A. E., Ezenwa, V. O., Etienne, R. S., Turner, W. C., & Olff, H. (2008). INTERACTIONS BETWEEN MACROPARASITES AND MICROPARASITES DRIVE INFECTION PATTERNS IN FREE-RANGING AFRICAN BUFFALO. *Ecology*, *89*(8), 2239–2250. https://doi.org/10.1890/07-0995.1

Krichel, L., Kirk, D., Pencer, C., Hönig, M., Wadhawan, K., & Krkošek, M. (2023). Short-term temperature fluctuations increase disease in a Daphnia-parasite infectious disease system. *PLOS Biology*, *21*(9), e3002260. https://doi.org/10.1371/journal.pbio.3002260

Lafferty, K. D., Dobson, A. P., & Kuris, A. M. (2006). Parasites dominate food web links. *Proceedings of the National Academy of Sciences*, *103*(30), 11211–11216. https://doi.org/10.1073/pnas.0604755103

Lawson, C. R., Vindenes, Y., Bailey, L., & van de Pol, M. (2015). Environmental variation and population responses to global change. *Ecology Letters*, *18*(7), 724–736. https://doi.org/10.1111/ele.12437

Lips, K. R., Diffendorfer, J., Mendelson, J. R., & Sears, M. W. (2008). Riding the Wave: Reconciling the Roles of Disease and Climate Change in Amphibian Declines. *PLoS Biology*, *6*(3), e72. https://doi.org/10.1371/journal.pbio.0060072

Luo, J., Greif, S., Ye, H., Bumrungsri, S., Eitan, O., & Yovel, Y. (2021). Flight rapidly modulates body temperature in freely behaving bats. *Animal Biotelemetry*, *9*(1), 45. https://doi.org/10.1186/s40317-021-00268-6

Mackey, R. L., & Currie, D. J. (2001). *The Diversity-Disturbance Relationship: Is It Generally Strong and Peaked?*

Miller, A. D., Roxburgh, S. H., & Shea, K. (2011). How frequency and intensity shape diversity–disturbance relationships. *Proceedings of the National Academy of Sciences*, *108*(14), 5643–5648. https://doi.org/10.1073/pnas.1018594108

Miller, A. D., Roxburgh, S. H., & Shea, K. (2012). Timing of disturbance alters competitive outcomes and mechanisms of coexistence in an annual plant model. *Theoretical Ecology*, *5*(3), 419–432. https://doi.org/10.1007/s12080-011-0133-1

Molnár, P. K., Sckrabulis, J. P., Altman, K. A., & Raffel, T. R. (2017). Thermal Performance Curves and the Metabolic Theory of Ecology—A Practical Guide to Models and Experiments for Parasitologists. *Journal of Parasitology*, *103*(5), 423. https://doi.org/10.1645/16-148

Paull, S. H., LaFonte, B. E., & Johnson, P. T. J. (2012). Temperature-driven shifts in a host-parasite interaction drive nonlinear changes in disease risk. *Global Change Biology*, *18*(12), 3558–3567. https://doi.org/10.1111/gcb.12018

Pedersen, A. B., & Fenton, A. (2007). Emphasizing the ecology in parasite community ecology. *Trends in Ecology & Evolution*, *22*(3), 133–139. https://doi.org/10.1016/j.tree.2006.11.005

Plowright, R. K., Parrish, C. R., McCallum, H., Hudson, P. J., Ko, A. I., Graham, A. L., & Lloyd-Smith, J. O. (2017). Pathways to zoonotic spillover. *Nature Reviews Microbiology*, *15*(8), 502–510. https://doi.org/10.1038/nrmicro.2017.45

Råberg, L., De Roode, J. C., Bell, A. S., Stamou, P., Gray, D., & Read, A. F. (2006). The Role of Immune-Mediated Apparent Competition in Genetically Diverse Malaria Infections. *The American Naturalist*, *168*(1), 41–53. https://doi.org/10.1086/505160

Raffel, T. R., Romansic, J. M., Halstead, N. T., McMahon, T. A., Venesky, M. D., & Rohr, J. R. (2013). Disease and thermal acclimation in a more variable and unpredictable climate. *Nature Climate Change*, *3*(2), 146–151. https://doi.org/10.1038/nclimate1659

Ramesh, A., & Hall, S. R. (2023). Niche theory for within-host parasite dynamics: Analogies to food web modules via feedback loops. *Ecology Letters*, *26*(3), 351–368. https://doi.org/10.1111/ele.14142

Rohr, J. R., & Raffel, T. R. (2010). Linking global climate and temperature variability to widespread amphibian declines putatively caused by disease. *Proceedings of the National Academy of Sciences*, *107*(18), 8269–8274. https://doi.org/10.1073/pnas.0912883107

Rohr, J. R., Raffel, T. R., Blaustein, A. R., Johnson, P. T. J., Paull, S. H., & Young, S. (2013). Using physiology to understand climate-driven changes in disease and their implications for conservation. *Conservation Physiology*, *1*(1), cot022–cot022. https://doi.org/10.1093/conphys/cot022

Rohr, J. R., Raffel, T. R., Romansic, J. M., McCallum, H., & Hudson, P. J. (2008). Evaluating the links between climate, disease spread, and amphibian declines. *Proceedings of the National Academy of Sciences*, *105*(45), 17436–17441. https://doi.org/10.1073/pnas.0806368105

Roxburgh, S. H., Shea, K., & Wilson, J. B. (2004). The intermediate disturbance hypothesis: Patch dynamics and mechanisms of species coexistence. *Ecology*, *85*(2), 359–371. https://doi.org/10.1890/03-0266

Schoolfield, R. M., Sharpe, P. J. H., & Magnuson, C. E. (1981). Non-linear regression of biological temperature-dependent rate models based on absolute reaction-rate theory. *Journal of Theoretical Biology*, *88*(4), 719–731. https://doi.org/10.1016/0022-5193(81)90246-0

Schulte, P. M., Healy, T. M., & Fangue, N. A. (2011). Thermal Performance Curves, Phenotypic Plasticity, and the Time Scales of Temperature Exposure. *Integrative and Comparative Biology*, *51*(5), 691–702. https://doi.org/10.1093/icb/icr097

Sckrabulis, J. P., Altman, K. A., & Raffel, T. R. (2022). Using Metabolic Theory to Describe Temperature and Thermal Acclimation Effects on Parasitic Infection. *The American Naturalist*, *199*(6), 789–803. https://doi.org/10.1086/719409

Sinclair, B. J., Marshall, K. E., Sewell, M. A., Levesque, D. L., Willett, C. S., Slotsbo, S., Dong, Y., Harley, C. D. G., Marshall, D. J., Helmuth, B. S., & Huey, R. B. (2016). Can we predict ectotherm responses to climate change using thermal performance curves and body temperatures? *Ecology Letters*, *19*(11), 1372–1385. https://doi.org/10.1111/ele.12686

Song, C., & Spaak, J. W. (2024). Trophic tug-of-war: Coexistence mechanisms within and across trophic levels. *Ecology Letters*, *27*(4), e14409. https://doi.org/10.1111/ele.14409

Spaak, J. W., Godoy, O., & De Laender, F. (2021). Mapping species niche and fitness differences for communities with multiple interaction types. *Oikos*, *130*(12), 2065–2077. https://doi.org/10.1111/oik.08362

Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., & Midgley, P. M. (Eds.). (2013). *Climate change 2013: The physical science basis. In: Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press.

Susi, H., Barrès, B., Vale, P. F., & Laine, A.-L. (2015). Co-infection alters population dynamics of infectious disease. *Nature Communications*, *6*(1), 5975. https://doi.org/10.1038/ncomms6975

Symons, C. C., & Arnott, S. E. (2014). Timing is everything: Priority effects alter community invasibility after disturbance. *Ecology and Evolution*, *4*(4), 397–407. https://doi.org/10.1002/ece3.940

Tracy, A. M., Pielmeier, M. L., Yoshioka, R. M., Heron, S. F., & Harvell, C. D. (2019). Increases and decreases in marine disease reports in an era of global change. *Proceedings of the Royal Society B: Biological Sciences*, *286*(1912), 20191718. https://doi.org/10.1098/rspb.2019.1718

Tuljapurkar, S. (1990). *Population Dynamics in Variable Environments* (Vol. 85). Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-642-51652-8

Vasseur, D. A., DeLong, J. P., Gilbert, B., Greig, H. S., Harley, C. D. G., McCann, K. S., Savage, V., Tunney, T. D., & O'Connor, M. I. (2014). Increased temperature variation poses a greater risk to species than climate warming. *Proceedings of the Royal Society B: Biological Sciences*, *281*(1779), 20132612. https://doi.org/10.1098/rspb.2013.2612

Williamson, M. H. (1957). An Elementary Theory of Interspecific Competition. *Nature*, *180*(4583), 422–425. https://doi.org/10.1038/180422a0

Wootton, J. T. (1998). Effects of Disturbance on Species Diversity: A Multitrophic Perspective. *The American Naturalist*, *152*(6), 803–825. https://doi.org/10.1086/286210

Wright, R. K., & Cooper, E. L. (1981). Temperature effects on ectotherm immune responses. *Developmental and Comparative Immunology*, *5*, 117–122.