**Environmental variability can promote parasite diversity within hosts and transmission among hosts**

Madeline Jarvis-Cross[1], Martin Krkosek[1]

[1] The University of Toronto, Department of Ecology and Evolutionary Biology

25 Willcocks Street, Toronto, ON, M5S 3B2

Corresponding author: Madeline Jarvis-Cross madeline.jarvis.cross@mail.utoronto.ca

**Acknowledgements**

This work was supported by The Emerging and Pandemic Infections Consortium Doctoral Award (MJ-C), and Ontario Graduate Scholarship (MJ-C), an NSERC Discovery Grant (MK), and a Canada Research Chair (MK).

**Author Contributions**

MJC and MK conceived the ideas and designed methodology; MJC collected the data; MJC analysed the data; MJC led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

**Conflicts of Interest**

The authors have declared that no conflicts of interest exist.

**Data Availability Statement**

Time series data available from the Federated Research Data Repository at https://doi.org/10.20383/103.0764. Scripts and replication materials available from GitHub at https://github.com/MadelineJC/EnvironmentalVariability_Repo.

**Abstract**

1. While the mechanisms that govern disease emergence and spread among hosts are increasingly

well-described, the mechanisms that promote parasite diversity within-hosts, affecting host

outcomes and spillover potential, have been comparatively understudied. Furthermore, while

attention has been paid to the effects of increasing temperatures on disease systems, the effects of

environmental variability have been left underexplored, despite rising climatic variability.

2. To investigate the impacts of environmental variability on parasite diversity within-hosts, we

analyzed a model of within-host population dynamics wherein two parasites indirectly compete

through the host's immune response.

3. We simulated the model under constant, demographically stochastic, environmentally stochastic,

and demographically and environmentally stochastic conditions, and analysed the viability and

longevity of non-equilibrium parasite co-occurrence.

4. We found that environmental stochasticity increased the viability and longevity of parasite co-

occurrence, suggesting that thermal variability arising from climatic change and as a physiological

trait may promote parasite diversity within ectotherms and help explain bats' propensity to support

diverse communities of parasites.

5. Further, we found that under certain conditions, the transmissibility of co-occurring parasites can

surpass the transmissibility of single parasites, suggesting that thermal variability may increase the

transmissibility of co-occurring parasites.

**Introduction**

It has become clear that climate and land-use change can facilitate novel species interactions and

increase contact rates between wild populations and humans, resulting in devastating epidemics and

enabling spillover and disease emergence events among wild and human hosts alike (Daszak et al., 2000,

2001; Plowright et al., 2017). However, despite the consequential interplay between within- and among-

host processes, the mechanisms that promote parasite diversity within hosts and enable events observed

among hosts have been understudied.

An intermediate level of disturbance may promote species diversity by minimizing the competitive

advantages apparent at low levels of disturbance, and forestalling the extinction risks brought on by high

levels of disturbance (Chesson, 1994; Connell, 1978). Roxburgh et al. (2004) posited that because

different species respond to disturbances in different ways, there are myriad ways in which diversity may

be generated (Roxburgh et al., 2004). As such, Roxburgh et al. (2004) conceptualized the intermediate

disturbance hypothesis not as a mechanism of coexistence unto itself, but as a collection of mechanisms

through which disturbances in resource availability or environmental conditions may promote diversity

(Roxburgh et al., 2004). Developing a generalizable understanding of the relationship between

disturbance and diversity has been complicated by the conceptual multidimensionality of "disturbance",

and by a lack of understanding with regards to how various ecological factors affect the shape and

intensity of such relationships (Mackey & Currie, 2001). Further, in multi-trophic systems, Wootton

(1998) demonstrated that the effects of disturbance on species diversity are dependent on the structure of

the system (Wootton, 1998). For example, when competitors have a shared enemy, intermediate levels of

disturbance are more likely to promote coexistence when consumers have a larger impact on the stronger

competitor (Holt & Lawton, 1994; Wootton, 1998).

While niche theory contends that that two species occupying the same niche should not coexist,

such species are often observed to do so (Hardin, 1960; Hutchinson, 1961). It has been proposed that this

contradiction is a consequence of historical disagreement regarding the definition and measurement of

the niche (Sales et al., 2021; Vandermeer, 1972). Modern coexistence theory aims to circumvent the

ambiguity of "the niche", and explain coexistence as a product of either stabilizing, or equalizing

mechanisms (Chesson, 2000). While stabilizing mechanisms (e.g. niche differentiation), which promote

coexistence despite fitness differences, have been employed to explain the maintenance of parasite

diversity within hosts, the contributions of equalizing mechanisms (e.g. relative nonlinearity), which

decrease average fitness differences between competitors, have yet to be examined (Bashey, 2015; Chesson, 1994, 2000, 2003; Chesson & Huntly, 1997). Originally designed to consider two species directly competing for a shared resource within a single trophic level, modern coexistence theory has been expanded to consider more complex communities, including multi-trophic communities (Barabás et al., 2018; Song & Spaak, 2024; Spaak et al., 2021; Spaak & De Laender, 2020). In a recent paper, Song and Spaak (2024) demonstrated that in a multi-trophic system, diversity in lower trophic levels is primarily constrained by fitness differences, while diversity in higher trophic levels is primarily constrained by niche differences (Song & Spaak, 2024). As stated above, fitness differences can be minimized via equalizing mechanisms, like relative nonlinearity, whereby competing species are differentially affected by variation, including environmental variation.

Here, we consider the relationship between environmental variation and within-host parasite diversity along two axes (Mackey & Currie, 2001; Miller et al., 2011, 2012). First, we consider rising temperature variability as a feature of climate change (Stocker et al., 2013). While much attention has been paid to the effects of increasing temperatures on disease systems, far less has been invested in exploring the effects of increasing temperature variability, despite evidence of its outsized impacts on species outcomes (Dobson & Carper, 1992; Lawson et al., 2015; Vasseur et al., 2014). In ectothermic hosts, increasingly erratic external conditions will affect the internal conditions of the host, thereby also affecting any parasites the host may harbour (Cizauskas et al., 2017). A relationship between thermal variability and within-host parasite diversity may be of particular interest when considered alongside the long-debated relationship between climate change and widespread disease-mediated amphibian declines (Daszak et al., 1999; Johnson, 2006; Lips et al., 2008; Rohr et al., 2008). While it's been posited that increasing temperatures may drive these declines, Rohr and Raffel (2010) demonstrated that El Niño events and associated regional temperature variability increase host susceptibility to disease, and thus help drive disease-mediated amphibian declines (Cohen et al., 2019; Rohr et al., 2008; Rohr & Raffel, 2010).

Second, one may consider environmental variability as a physiological trait of hosts. Bats (order

*Chiroptera*) have long been acknowledged as prolific reservoirs for disease and have often been

implicated in zoonotic spillover events (Brook & Dobson, 2015; Calisher et al., 2006; Irving et al.,

2021). Bats are also unique among mammals in that their internal temperature varies widely in

accordance with activity level (e.g. torpor, flight) (Luo et al., 2021). It has been theorized that internal

temperature variability may aid bats' immune responses, allowing them to carry microparasites without

suffering outsized morbidity (Fumagalli et al., 2021). Similarly, internal temperature variability may

impose environmental variability on parasites, and affect their ability to coexist. In combination, the

effects of internal temperature variability on immune responses and parasites coexistence may help

explain bats' propensity to support large communities of parasites.

We used a model to investigate the impacts of environmental variability on the viability and

maintenance of parasite diversity within hosts. We began by adapting Antia et al.'s (1994) model of

within-host population dynamics to include two parasites that indirectly compete via the host's immune

system, and thus exhibit apparent competition. We analyzed deterministic, demographically stochastic,

environmentally stochastic, and demographically-and-environmentally stochastic implementations of the

model. Within the environmentally stochastic implementations of the model, we varied the magnitude

and frequency of thermal disturbance within the system. Between and among the deterministic and

stochastic simulations of the model, we compared the ability of the secondary parasite to invade a system

containing the primary parasite, and the duration of time for which the primary and secondary parasites

co-occurred. Further, we compared the total transmission of co-occurring parasites with the transmission

potential of a single parasite. We found that compared to a constant environment, a variable environment

tended to increase the ability of the secondary parasite to invade the system, facilitated longer periods of

co-occurrence between the primary and secondary parasites, and increased the transmissibility of co-

occurring parasites. These results (1) suggest that climatic changes may have significant effects on

parasite diversity within ectothermic hosts, (2) provide insight on a host's ability to support large

communities of parasites, and propose that variability within-hosts may influence transmission dynamics

among-hosts.

**Methods**

**Model Description**

*Within-Host Population Dynamics Model*

We began with a model of within host population dynamics based on Antia et al., (1994) that

represents the interaction between an infecting parasite and its host's immune system as analogous to the

interaction between a prey item and its predator. Per Antia et al. (1994), the density of the parasite

population, $P$, is determined by its replication rate, $r$, and the rate at which it is cleared by the host's

immune system, $k$. The model assumes that the replication rate of the host's immune system is

proportional to parasite density at low parasite densities, and saturates at high parasite densities, allowing

the host's immune system to either extirpate or succumb to the infection (Antia et al., 1994). We adapted

the model to include a second parasite, $P_j$, that indirectly competes with the primary parasite, $P_i$, through

the host's immune response (Eqns. 1-3, Table 1). As such, the model describes an immune-mediated

apparent competition (Holt, 1977; Råberg et al., 2006; Williamson, 1957). Parameters values were not

chosen to describe a specific system, but to replicate a dynamical behaviour, as described above and

shown in Figure 1. Given that an infinite number of parameter value combinations can produce this

behaviour, we opted to maintain the initial conditions and select parameter values proposed by Antia et

al. (1994). However, we increased the rate at which parasites are destroyed by the host's immune system

($k$) and decreased the maximum replication rates of the primary and secondary parasites ($r_i, r_j$) to favour

extirpation, and eliminate host death as an infection outcome (Table 1). We also conducted a sensitivity

analysis to determine how variations in (1) the value of $k$ and (2) the shape of the relationship between

parasite replication rate and the temperature of the system would affect our results, and found the model

to be robust (see pages 2, 4-8 of supplement for complete results).

(1) $\quad P_i(t+1) = (1 + r_i - (1 - e^{-kI(t)}))P_i(t)$

(2) $\quad P_j(t+1) = (1 + r_j - (1 - e^{-kI(t)}))P_j(t)$

(3) $\quad I(t+1) = (1 + \rho\left(\dfrac{P_i(t)}{P_i(t)+\phi} + \dfrac{P_j(t)}{P_j(t)+\phi}\right))I(t)$

153
154   **Table 1:** Parameter definitions and values.

| Symbol | Parameter | Values Stated in Antia et al. (1994) | Values Used |
|---|---|---|---|
| $P_i$ and $P_j$ | Parasite abundance | 1* | 1* |
| $I$ | Host immune cell abundance | 1* | 1* |
| $r_i$ and $r_j$ | Replication rates of parasites | 0.1 – 10.0 | $0.1 – 2.0^\dagger$ |
| $k$ | Rate at which parasites are destroyed by host's immune system | $10^{-3}$ | $10^{-2}$ |
| $\rho$ | Maximum replication rate of host's immune system | 1.0 | 1.0 |
| $\phi$ | Parasite density at which replication rate of host's immune system is half its maximum | $10^3$ | $10^3$ |
| $D$ | Lethal within-host parasite density | $10^9$ | $10^9$ |

* Initial conditions

$^\dagger$ Dependent on the temperature of system in environmentally stochastic simulations

To compare the impacts of different types of stochasticity on parasite diversity, we wrote deterministic

(D), demographically stochastic ($S_D$), environmentally stochastic ($S_E$), and demographically-and-

environmentally stochastic ($S_{DE}$) implementations of the model. Within the demographically stochastic

implementation of the model, we wrote the model's growth terms as Poisson processes, and mortality

term as a binomial process (Eqns. 4-6).

163
(4) $\quad P_i(t+1) = P_i(t) + B_{P_i}(t) - D_{P_i}(t)$

(5) $\quad P_j(t+1) = P_j(t) + B_{P_j}(t) - D_{P_j}(t)$

(6) $\quad I(t+1) = I(t) + B_I(t)$

In Eqns. (4) and (5), $B_{P_i}(t)$ and $B_{P_j}(t)$ are Poisson random variables such that $B_{P_i}(t) \sim Poisson(1, \lambda_i)$,

where $\lambda_i = r_i \cdot P_i(t)$, and $B_{P_j}(t) \sim Poisson(1, \lambda_j)$, where $\lambda_j = r_j \cdot P_j(t)$. In Eqn. (6), $B_I(t)$ is a Poisson

random variable such that $I_B(t) \sim Poisson(1, \lambda)$, where $\lambda = \rho\left(\dfrac{P_i(t)}{P_i(t)+\phi} + \dfrac{P_j(t)}{P_j(t)+\phi}\right)I(t)$. In Eqns. (4) and

(5), $D_{P_i}(t)$ and $D_{P_j}(t)$ are binomial random variables such that $D_{P_i}(t) \sim Binomial(1, P_i(t), p)$, where

$p = 1 - e^{-kI(t)}$, and $D_{P_j}(t) \sim Binomial(1, P_j(t), p)$. To write the environmentally stochastic

implementation of the model, we generated a stochastic temperature series and allowed the replication

rates of each parasite $(r_i, r_j)$ to vary in response to the temperature of the system (Eqns. 7-9). We chose

to write $r_i$ and $r_j$, though not $k$, as temperature-dependent isolate the effect of thermal variability on

traits that can be metabolically scaled, and constrain the scope of the work.

177
178  (7)   $P_i(t+1) = (1 + r_i(T(t)) - (1 - e^{-kI(t)}))P_i(t)$
179  (8)   $P_j(t+1) = (1 + r_j(T(t)) - (1 - e^{-kI(t)}))P_j(t)$
180  (9)   $I(t+1) = (1 + \rho \left( \dfrac{P_i(t)}{P_i(t) + \phi} + \dfrac{P_j(t)}{P_j(t) + \phi} \right))I(t)$

181
We used first-order autocorrelated Gaussian noise to generate stochastic temperature series. We began

with a value of 0, and generated the next value, $T(t+1)$, by multiplying the previous value by the

coefficient of autocorrelation (AC = 0.25), and summing the resulting value with a value drawn from a

normal random variable with a mean of zero. We varied the standard deviation of the normal random

variable, as described below. We then added the average temperature of the system to each value in the

series. Resulting values outside the thermal range of the system were re-drawn. We then converted the

stochastic temperature series into series of replication rates using various parameterizations of the

Sharpe-Schoolfield sub-model (Eqns. S1.1, S1.2; Table S2).

*Sharpe-Schoolfield Sub-Model*

The Metabolic Theory of Ecology (MTE) predicts how metabolic rates control ecological

processes across levels of organization, and explains how metabolic rates vary with body size and

temperature (Brown et al., 2004). Building off the MTE, the Boltzmann-Arrhenius (BA) relation

describes metabolism as a function of temperature. The Sharpe-Schoolfield model characterizes the

performance of a trait as a function of temperature through a thermal performance curve (TPC)

(Schoolfield et al., 1981). When applied to disease systems, the Sharpe-Schoolfield model has been

shown to describe relationships between temperature and species traits, and has proven useful in

investigating the relationship between rising temperatures and disease transmission.

We used various parameterizations of the Sharpe-Schoolfield model to generate paired thermal

performance curves relating the temperature of the system to the replication rates of two parasites ($r_i$ and

$r_j$) (Fig. S9). We generated four paired thermal performance curves, representing four thermal scenarios:

(1) the identical responses scenario, (2) the offset responses scenario, (3) the $P_i$-advantage scenario, and

(4) the $P_j$-advantage scenario (Table S2). Each thermal scenario describes a different relationship

between the temperature of the system and parasite replication rates. The identical responses scenario

describes $r_i$ and $r_j$ as responding identically to the temperature of the system (Fig. S9). Under the offset

responses scenario, $r_i$ and $r_j$ peak on opposite sides of the average temperature of the system (Fig. S9).

To develop the $P_i$-advantage scenario, we decreased the average temperature of the system to give the

primary parasite, $P_i$, a thermal advantage (Fig. S9). We considered two versions, (I) and (II), of the $P_j$-

advantage scenario, each of which gave the secondary parasite, $P_j$, a thermal advantage. To develop (I),

we increased the average temperature of the system to give the secondary parasite a quantitatively

comparable thermal advantage (Fig. S9). To develop (II), we replicated the $P_i$-advantage scenario, and

assigned the TPC with the lower optimum to $P_j$, to give the secondary parasite the same thermal

advantage given to the primary parasite and isolate the effects of asymmetry (Fig. S9). Given that (I) and

(II) produced qualitatively comparable results, we will only discuss (I) in the main text (see supplement

for complete results). Within each thermal scenario, the average temperature of the system was chosen to

maintain quantitatively comparable (1) parasite replication rates between thermal scenarios and (2)

differences between replication rates at the average temperature of the system.

Given that the Sharpe-Schoolfield model cannot generate negative values, it cannot be used to

explore cases in which extreme temperatures result in parasite mortality and population decline. To

compensate, we developed an alternative set of paired thermal performance curves, though which

extreme temperatures produce "negative" replication rates, and spur population decline (see "Extended

analysis" in supplement; Fig. S19).

**Model Simulations**

224           Within each thermal scenario, we simulated the deterministic model (D), and each of the three

225     stochastic models ($S_D$, $S_E$, $S_{DE}$) for fifty days with a discretization time step of 0.1, which was sufficient

226     to capture the full course of parasite infection and clearance across all simulations. We began each set of

227     simulations with the primary parasite and the host's immune system in the system, at initial abundances

228     of 1 as the initial condition. We then considered all cases for the introduction of the secondary parasite

229     from its introduction at the onset of infection, to its introduction on the forty-ninth day of infection.

230           For each environmentally stochastic model ($S_E$, $S_{DE}$) simulation, we generated a new temperature

series and two corresponding series of replication rates. When determining the ways in which new

temperatures would be selected, we considered two axes of thermal disturbance: magnitude and

frequency. To vary magnitude of disturbance, we varied the standard deviation of the normal random

variable in the temperature autocorrelation function, starting from zero (no thermal disturbance) and

increasing by two until the effect of magnitude on co-occurrence was fully characterized (SD = 30),

resulting in fifteen discrete magnitudes of disturbance. To vary frequency of disturbance, we re-assigned

the temperature of the system every 10, 9, 8, 7, 6, 5, 4, 3, and 2 days, once per day, and 2, 5, and 10

times per day, resulting in thirteen frequencies of disturbance. Considering these magnitudes and

frequencies of disturbance fully-factorially resulted in 195 disturbance regimes. We conducted 1000

simulations per stochastic model and treatment (Fig. 1). Two hundred such simulations are shown in

Figures 1 and S10.

Figure 1: Schematic detailing simulation experiment. We considered four thermal scenarios, and four models (deterministic,

demographically stochastic, environmentally stochastic, and demographically and environmentally stochastic). When

simulating the environmentally stochastic models, we considered 195 disturbance regimes. When simulating each of the four

models, we considered 50 days on which the secondary parasite could be introduced. We conducted 1000 simulations per

stochastic model and treatment.

**Co-Occurrence Analyses**

Given successful invasion by the secondary parasite ($P_j(t + 1) > P_j(t)$), we quantified

longevities of co-occurrence by counting the number of time steps for which primary and secondary

parasite population abundances were equal to or greater than one within each simulated timeseries. We

summarised these results as distributions for each stochastic model and treatment. Using the $S_D$ model as

a baseline, we investigated the effects of environmental variability on the viability and maintenance of

parasite diversity by comparing the ability of the secondary parasite to invade a single-parasite system,

and the duration of time for which the primary and secondary parasites co-occurred across stochastic

treatments. Additionally, we analysed (1) the proportion of stochastic simulations per model and

treatment that resulted in longer periods of co-occurrence relative to the deterministic model to quantify

rarity, and (2) the competitive outcomes of each treatment to gain insight on underlying mechanisms.

**Transmissibility Analyses**

To explore the impacts of co-infection on transmission potential, we compared the lifetime

replicative outputs of co-occurring parasites with those of a single parasite in constant and variable

environments. As per Antia et al., (1994), we determined the total transmission ($q_i, q_j, q$, where $q$ is the

total transmission of the single parasite) of a parasite ($P_i, P_j, P$, where $P$ is the single parasite) from an

infected host by taking the integral of the rate of transmission ($u$) of each parasite from the host over the

duration of infection (Eqns. 10.1-10.3). For simplicity, we set the rate of transmission to one.

(10.1)   $q_i = \displaystyle\int_0^\infty uP_i(t)dt$

(10.2)   $q_j = \displaystyle\int_0^\infty uP_j(t)dt$

(10.3)   $q = \displaystyle\int_0^\infty uP(t)dt$

269
To compare constant and variable environments, we analysed multi-parasite and single-parasite versions

of the $S_D$ and $S_{DE}$ models.

**Results**

Analytically, the secondary parasite could invade ($P_j(t+1) > P_j(t)$) when $I(t) < \frac{-\ln(1-r_j)}{k}$. Our

parameterization of the $S_D$ model allowed the secondary parasite to invade when introduced before the

tenth to twelfth day of infection, resulting in 8.4 to 9.2 days of co-occurrence, depending on thermal

scenario (Fig. S11, Supplementary Datafile 1-1:10).

**Co-occurrence Analysis**

Simulation outputs show that compared to a deterministic baseline, the addition of demographic

stochasticity has little effect on co-occurrence (Supplementary Datafile 2-1:10), while the addition of

thermal variability increases the ability of the secondary parasite to invade the system and facilitates

longer periods of co-occurrence between the primary and secondary parasites. While longevities of co-

occurrence are maximised under the $P_j$-advantage scenario, the difference between deterministic and

environmentally stochastic implementations is not consistently maximised by any one thermal scenario.

Additionally, while larger and more frequent disturbances tend to promote longer periods of co-

occurrence, increasingly large disturbances produce a less steep effect than do increasingly frequent

disturbances. These findings are common among thermal scenarios.

When both parasites were introduced simultaneously, $S_D$ model simulations resulted in periods of

co-occurrence that were, on average, 4.5% to 3.1% shorter (0.4 to 0.3 days) than deterministically

simulated periods of co-occurrence (Figs. 2, S12). Under the same conditions, simulations of the

environmentally stochastic models ($S_E$, $S_{DE}$) resulted in periods of co-occurrence that were up to 46.2%

($S_E$) and 36.0% ($S_{DE}$) longer (3.9 ($S_E$) and 3.0 ($S_{DE}$) days) than deterministically simulated periods of co-

occurrence (Figs. 2, S12). Across all 195 disturbance regimes, simulations of the $S_E$ model resulted in

mean periods of co-occurrence that were between 2.7% shorter (0.2 days) and 46.2% longer (3.9 days)

than their deterministic counterparts (Figs. 2, S12). The identical responses scenario promoted the

shortest and longest mean periods of co-occurrence (Figs. 2-3, S12-S13). Simulations of the $S_{DE}$ model

resulted in mean periods of co-occurrence that were between 6.8% shorter (0.6 days) and 36.0% longer

(3.0 days) than deterministically simulated periods of co-occurrence (Figs. 2, S12). Again, the identical

responses scenario promoted the shortest and longest mean periods of co-occurrence (Figs. 2-3, S12-

S13). Across thermal scenarios, mean periods of co-occurrence were shortest when disturbances were

large (standard deviation of the normal random variable in the temperature autocorrelation function = 22

to 30) and infrequent (once every 7 to 9 days) (Figs. 3, S13). Mean periods of co-occurrence were

longest when disturbances were large (SD = 24 to 30) and frequent (five to ten times per day) (Figs. 3,

S13).

More generally, as disturbances increased in size, periods of co-occurrence increased in length

towards a saturation point (standard deviation of the normal random variable in the temperature

autocorrelation function ≈ 24-30, depending on thermal scenario) (Figs. S14, S18, Supplementary

Datafiles 3-1:5, 4-1:5). An analysis of simulations within which an alternative set of TPCs allowed for

extreme temperatures to cause population decline revealed a parabolic relationship between disturbance

size and longevity of co-occurrence, that peaked when the standard deviation of the normal random

variable in the temperature autocorrelation function $\approx 8$ (Fig. S20).

When disturbances were infrequent (once every seven to ten days), periods of co-occurrence

were similar in length to deterministically simulated periods of co-occurrence. Per the deterministic

model, $P_i$ and $P_j$ can only co-occur for a maximum of 9.2 days. Thus, systems subjected to infrequent

disturbances were functionally similar to systems subjected to a constant environment, making the

outcome of a simulation reliant on the initial conditions of the system. Interestingly, when disturbances

were intermediately frequent (once per day to once every few days), mean periods of co-occurrence were

shorter than deterministically simulated periods of co-occurrence (Fig. S14, S21). When disturbances

were frequent (twice to ten times per day), mean periods of co-occurrence were longer than

deterministically simulated periods of co-occurrence. Finally, within a given disturbance regime,

distributions of longevities of co-occurrence were normally distributed (Figs. S14-S15, S23).

Figure 2: The mean percent difference in longevity of co-occurrence relative to deterministically simulated periods of co-

occurrence, when both parasites are introduced at the same time. Each point represents the mean percent difference in

longevity of co-occurrence relative to deterministically simulated periods of co-occurrence across 1000 stochastic

simulations, as per a given environmental disturbance regime. As such, there are 195 points above each "$S_E$"/"$S_{DE}$" label.

Figure 3: Fully factorial representations of the impacts of varyingly sized and timed disturbances on the longevity of co-

occurrence. The plot shows the mean number of time steps (10 time steps/day) for which $P_i$ and $P_j$ have population

abundances greater than one as per the $S_{DE}$ model. The circled numbers denote the number of time steps for which $P_i$ and $P_j$

can co-occur, per the deterministic model.

**Rarity and Competitive Outcomes**

When the secondary parasite was introduced before or on the invasion threshold (10[th] or 12[th] day

of infection, depending on thermal scenario), 29%-71% ($S_E$) and 46%-58% ($S_{DE}$) of environmentally

stochastic simulations resulted in longer periods of co-occurrence relative to deterministic simulations

(Fig. 4). When the secondary parasite was introduced after the invasion threshold, the likelihood that an

environmentally stochastic simulation resulted in some period of co-occurrence sharply declined. When

the secondary parasite was introduced one to twelve days after the invasion threshold, 1%-31% ($S_E$) and

1%-46% ($S_{DE}$) of environmentally stochastic simulations resulted in some period of co-occurrence (Fig.

4). Thus, environmental variability extended the invasion threshold by 50%-75% ($S_E$) and 100%-120%

($S_{DE}$), depending on thermal scenario (Fig. 4).

Analyzing competitive outcomes at intermediate levels of disturbance (SD = 10, frequency of

disturbance = once per day) revealed that under the offset responses scenario, when $P_i$ and $P_j$ were

introduced at the same time, $P_j$ outlasted $P_i$ 65.5% ($S_E$) and 63.4% ($S_{DE}$) of the time (Fig. S22-S23).

Additionally, under the $P_i$-advantage scenario, when $P_i$ and $P_j$ were introduced at the same time, $P_i$ and

$P_j$ were equally likely to outlast one another ($P_j$ outlasts $P_i$ 49.9% ($S_E$) and 51.0% ($S_{DE}$) of the time), but

under the $P_j$-advantage, $P_j$ was much more likely to outlast $P_i$ ($P_j$ outlasts $P_i$ 76.6% ($S_E$) and 70.4% ($S_{DE}$)

of the time) (Fig. S21-S23). Across thermal scenarios, when the secondary parasite, $P_j$, was introduced in

the middle of the infection period, $P_i$ was more likely to outlast $P_j$. When $P_j$ was introduced late in the

infection period, $P_j$ was more likely to outlast $P_i$ (Fig. S22-S23). Regardless of when $P_j$ was introduced,

simulations resulting in especially long periods of co-occurrence are equally likely to show either

parasite outlasting the other (Fig. S22-S23).

Figure 4: Proportion of environmentally stochastic simulations (SD = 10, frequency of disturbance = once per day) resulting

in longer periods of co-occurrence relative to deterministic simulations by the day on which the secondary parasite was

introduced. Vertical lines show the first day on which the introduction of the secondary parasite does not result in any co-

occurrence between the two parasites, as per the deterministic model (D).

**Transmissibility**

Across thermal scenarios, simulations of the multi-parasite $S_D$ model produced lifetime

replicative outputs (LROs) that were one to two orders of magnitude lower than those produced by

simulations of the single-parasite $S_D$ model (Supplementary Datafiles 5-1:16). Similarly, under the first

three thermal scenarios, simulations of the multi-parasite S$_{DE}$ model produced LROs that were one to

two orders of magnitude lower than those produced by simulations of the single-parasite S$_{DE}$ model.

However, under the $P_j$-advantage scenario and among 62 of 195 (32%) disturbance regimes, the average

transmission potential of the co-occurring secondary parasite, $P_j$, exceeded that of a single parasite (Fig.

5). Among these disturbance regimes, the transmission potential of $P_j$ exceeded that of a single parasite

by an average of 2% to 135% (Supplementary Datafiles 5-8). The transmission potentials of co-

occurring parasites peaked when disturbances were infrequent (once every ten days) to intermediately

frequent (once every six days) (Fig. S24). While large and frequent disturbances maximally increased

longevities of co-occurrence, such infections did not result in particularly high transmission potentials,

indicating that especially long periods of co-occurrence may be facilitated by lower lifetime replicative

outputs that do not trigger a large immune response, and thus allow for more persistent infections.

Figure 5: The impacts of environmental variability on transmission potential under the $P_j$-advantage scenario. The plot shows

a fully factorial representation of the impacts of thermal disturbance on the transmission potential of $P_j$, relative to the

equivalent single-parasite treatment. The black line indicates disturbance regimes under which the average transmission

potential of $P_j$ was equal to that of a single parasite. Colours warmer than denoted by the black line indicate disturbance

regimes under which the average transmission potential of $P_j$ exceeded that of a single parasite; colours cooler than denoted

by the black line indicate the opposite.

**Discussion**

381   We investigated the potential for environmental variability, both as a feature of climate change

382  and as a trait unique to a prolific reservoir for disease to affect the viability and maintenance of within-

383  host parasite diversity. Our work provides theoretical evidence that relative to a constant environment, a

384  variable environment can increase the ability of a secondary parasite to invade a single-parasite system,

385  promote longer periods of co-occurrence between parasites, and increase the transmissibility of co-

386  occurring parasites. In more detail, thermal disturbances of various sizes and frequencies allowed a

387  secondary parasite to invade a single-parasite system up to twelve days after the invasion threshold,

388  extending the invasion threshold by 100% to 120% ($S_{DE}$ model, Fig. 4). Large and frequent thermal

389  disturbances enabled parasites to co-occur for up to 37.8% longer, relative to a constant environment

390  ($S_{DE}$ model, Figs. 2, S12). Finally, under the $P_j$-advantage scenario and among 32% disturbance regimes,

391  the average transmission potential of the co-occurring secondary parasite exceeded that of a single

392  parasite ($S_{DE}$ model, Fig. 5).

393   Across thermal scenarios and disturbance regimes, the relationship between disturbance size and

394  longevity of co-occurrence was initially positive and eventually saturating (Fig. S14, S18). Large

395  disturbances produced a saturating response rather than the expected parabolic response due to the

396  structure of the Sharpe-Schoolfield sub-model. Allowing for extreme temperatures to result in parasite

397  mortality and population decline produced a parabolic response, implying that intermediate magnitudes

398  of disturbance maximised the viability and longevity of parasite co-occurrence (Figs. S19-S20). When

399  thermal disturbances were small, the parasite with the higher initial replication rate was more likely to

400  grow quickly and trigger a commensurate immune response that would competitively exclude the other.

401  As thermal disturbances increased in size, it became increasingly likely that when the temperature of the

402  system changed, the competitive advantage would shift from one parasite to the other, minimizing fitness

403  differences over the course of an infection and promoting co-occurrence. However, per the extended

404  analysis, large thermal disturbances caused parasite populations to alternatively explode and crash,

405  triggering a sizable immune response and increasing the likelihood of extirpation.

The relationship between climate change and disease-mediated amphibian declines has long been

debated. While some argue that climatic changes may impede immune function or result in performance

gaps between parasites and their hosts, making amphibians more susceptible to infection, others argue

that insufficient evidence should preclude a causal link (Cohen et al., 2019; Rohr et al., 2008). While

much of the literature proposes mean-climate hypotheses, some evidence suggests that intensifying El

Niño events are increasing thermal variability, which may in turn affect amphibians' susceptibility to or

ability to defend against infection (Rohr & Raffel, 2010). Congruently, our findings provide theoretical

evidence that thermal variability promotes the viability and maintenance of within-host parasite

diversity, and thus suggest that as the climate continues to change, attention should be paid to shifts in

microparasite communities and their impacts on ectothermic hosts. In particular, well-documented

climate-driven increases in disease incidence among a wide variety of marine organisms may constitute

an important starting point for future work (Bates et al., 2009; Bruno et al., 2007; Tracy et al., 2019).

Across thermal scenarios and disturbance regimes, intermediate and high frequencies of

disturbance impeded and promoted the viability and longevity of parasite co-occurrence, respectively

(Figs. 3, S13-S14, S21). Given the transient nature of infection specified by the underlying model,

intermediate frequencies of disturbance (once every seven days, for example) resulted in the temperature

of the system being disturbed towards the end of the infection period, when parasite populations were

small and especially susceptible to unfavourable conditions. Conversely, high frequencies of disturbance

resulted in the temperature of the system being disturbed throughout the duration of the infection period,

and as such, neither favourable nor unfavorable conditions lasted long enough for one parasite to gain a

decisive advantage over and exclude the other. While it has been theorized that internal temperature

variation may aid bats' immune responses, allowing them to carry microparasites without suffering

outsized morbidity (Fumagalli et al., 2021), these findings suggest that internal temperature variation

may also facilitate the co-occurrence of parasites within bats, and thus may help explain bats' propensity

to support diverse communities of parasites.

Under the offset responses scenario, $P_j$ often outlasted $P_i$, while under the $P_i$-advantage scenario,

$P_i$ and $P_j$ were equally likely to outlast each other (Fig. S22-S23). Under the $P_j$-advantage scenario, $P_j$

was much more likely to outlast $P_i$ (Fig. S22-S23). These outcomes are consequences of the

asymmetrical structure of the thermal performance curves, which made unfavourable conditions more

disadvantageous for $P_i$ than $P_j$ (Sinclair et al., 2016). Further, regardless of when $P_j$ was introduced,

simulations resulting in especially long periods of co-occurrence were equally likely to show either

parasite outlasting the other. Taken together, these results suggest that to predict the competitive

outcome of a multiple infection, one may need to consider the interplay between how well adapted a

parasite is to its host, priority effects, and the responses of biological rates to environmental factors.

While one might expect a specialist or well-adapted parasite to exploit its host more efficiently than a

generalist or novel parasite, the results of this work suggest that environmental variability can even the

scales, and allow a generalist or novel parasite to establish and persist, as predicted by the intermediate

disturbance hypothesis (Chesson, 1994; Connell, 1978; Roxburgh et al., 2004). Similarly, while one

might additionally expect a colonizing parasite to benefit from an initial lack of competition, the results

of this work again suggest that environmental variability and disturbance may contribute to the

structuring of communities by allowing latecomers to establish and persist more often and for longer

than they otherwise would (Jiang & Patel, 2008; Symons & Arnott, 2014).

While the metabolic theory of ecology posits that certain biological rates (e.g. metabolic rate,

replication/growth rate, etc.) will exhibit an asymmetrical and unimodal response to temperature, other

vital rates may exhibit symmetrical or bimodal responses to temperature (Dell et al., 2014; Molnár et al.,

2017; Sinclair et al., 2016). This work suggests that the shape of a thermal performance curve can

meaningfully affect the nature and outcome of an infection. The shape of a thermal performance curve

can also dictate the effects of stochasticity on persistence. Here, thermal variation pushes the system

away from a parasite's optimal temperature, and thus, away from its maximum replication rate. In

accordance with Jensen's inequality, compared to a deterministic system, the introduction of

stochasticity results in a lower average population growth rate (Denny, 2017; Jensen, 1906; Vasseur et

al., 2014). In the absence of thermal dependency, the introduction of stochasticity will also result in a

lower average population growth rate over time (Tuljapurkar, 1990). In this system, lower average

parasite replication rates slow down the infection and subsequently, the immune response, allowing

parasites to co-occur for longer. As anthropogenic changes alter host-parasite interaction networks,

understanding how evolutionary and environmental factors combine to affect interactions between

parasites will be essential in predicting and managing spillover and emergence in wildlife and human

populations (Paull et al., 2012; Pedersen & Fenton, 2007).

Developing an understanding of how the impacts of environmental variability on parasite

diversity within hosts may scale to affect among-host processes and events, including transmission,

spillover, and emergence will be essential to the maintenance of wildlife, human, and ecosystem health

(Allocati et al., 2016; Halpin et al., 2000; Johnson, 2006). While increased within-host parasite diversity

provides more strains or species of a parasite or parasites an opportunity to transmit, within-host

competition may result in decreased parasite loads, thus decreasing the transmission probability of any

one strain or species, and altering expected transmission patterns (Johnson et al., 2013; Pedersen &

Fenton, 2007). Similarly, our findings provide theoretical evidence that when environmental variations

are large and frequent, within-host competition decreases the total transmission of co-occurring parasites.

As such, while large and frequent disturbances may maximally increase the amount of time for which

two parasites co-occur, such infections do not result in particularly high transmission potentials,

indicating that especially long periods of co-occurrence may be facilitated by lower parasite replication

rates that do not trigger a large immune response, and thus allow for more persistent infections.

However, we found that approximately a third of tested disturbance regimes allowed the replicative

outputs of co-occurring parasites to exceed those of single parasites, indicating that the effects of

environmental variability may not be limited to the promotion of within-host parasite diversity, but may

also promote transmission between hosts (Fig. 5, S24). Disturbances of intermediate size and low to

intermediate frequency may constitute a sizable region within which environmental variation can

decrease fitness differences between parasites and promote within-host parasite diversity, while allowing

parasites to achieve sufficiently high parasite loads, resulting in more diverse and transmissible

infections (Fig. 5, S24).

While both Antia et al.'s (1994) model and our modified model should be broadly applicable to

an array of within-host multi-parasite systems, including multi-strain or cross-species, both models are

relatively simple as to remain generalizable, and thus cannot represent chronic infections, systems within

which parasites exhibit more complex interactions with their host or their host's immune system, nor

systems within which parasites directly interact with one another. As such, it may be interesting to

validate the theoretical insights of this work in an empirical system. This work is also limited by the

scope with which thermal dependency was considered. Given the thermal sensitivity of immune function

in ectotherms and endotherms, it may be worthwhile to consider the thermal dependency of immune

processes like response time and replication rate in tandem with the thermal dependency of parasite

replication rates (Butler et al., 2013; Wright & Cooper, 1981). Similarly, using nonlinear averaging to

assess the response of a nonlinear process to variability is contentious. While many theoretical works and

some empirical studies support the use of nonlinear averaging and find evidence for Jensen's inequality,

borne out as disproportionate effects of thermal variability on performance, many empirical studies have

contested an implicit assumption of nonlinear averaging: that traits should respond instantaneously to

change, without acclimation (Bernhardt et al., 2018; Estay et al., 2011; Krichel et al., 2023; Sinclair et

al., 2016). Thermal acclimation through phenotypic plasticity alters thermal performance curves, and as

such, produces a more complex relationship between temperature and biological rates that may not be

well-characterized by nonlinear averaging (Schulte et al., 2011; Sckrabulis et al., 2022). With regards to

host-parasite interactions, the temperature variability hypothesis posits that owing to their small relative

size, parasites should acclimate more quickly than their hosts, though the speed at which various

parasites may acclimate remains unclear (Raffel et al., 2013; Rohr et al., 2013). Interestingly, some

empirical studies of the impacts of environmental variability on host-parasite interactions have

considered thermal acclimation, yielding a range of conclusions including decreased pathogen

prevalence, reduced infection prevalence, and decreased resistance in hosts (Dallas & Drake, 2016;

Duncan et al., 2011; Raffel et al., 2013).

The intermediate disturbance hypothesis predicts that an intermediate level of disturbance in

resource availability or environmental conditions will promote coexistence by minimizing the

competitive advantages apparent at low levels of disturbance, and forestalling the extinction risks

brought on by high levels of disturbance (Chesson, 1994; Connell, 1978; Roxburgh et al., 2004).

Similarly, modern coexistence theory posits that variation can promote coexistence through equalizing

mechanisms, like relative nonlinearity (Barabás et al., 2018; Chesson, 1994, 2000, 2003; Chesson &

Huntly, 1997). Work by Wootton (1998), and more recently, Song and Spaak (2024) argue that the

maintenance of diversity is dependent on the trophic structure of the system (Song & Spaak, 2024;

Wootton, 1998). In particular, Wootton (1998) demonstrates that when competitors have a shared

enemy, intermediate levels of disturbance are more likely to promote coexistence, while Song and Spaak

(2024) argue that diversity within lower trophic levels is primarily constrained by fitness differences

(Song & Spaak, 2024; Wootton, 1998). Both the intermediate disturbance hypothesis and modern

coexistence theory predict a relationship between variability and diversity that is borne out in this work.

Here, we provide theoretical evidence that environmental variability promotes parasite diversity within

hosts and transmission among hosts. Camus wrote that plagues, despite their commonality, find us

unprepared. Improving our understanding of how environmental variability affects parasite diversity

within- and between-hosts will expand our understanding of the mechanisms that can facilitate disease

spread and emergence, and allow us to make robust predictions, and develop and implement mitigation

strategies as we move into an increasingly uncertain future. In brief, we will be found *less* unprepared.

**References**

Allocati, N., Petrucci, A. G., Di Giovanni, P., Masulli, M., Di Ilio, C., & De Laurenzi, V. (2016). Bat–man disease transmission: Zoonotic pathogens from wildlife reservoirs to human populations. *Cell Death Discovery*, *2*(1), 16048. https://doi.org/10.1038/cddiscovery.2016.48

Antia, R., Levin, B. R., & May, R. M. (1994). Within-Host Population Dynamics and the Evolution and Maintenance of Microparasite Virulence. *The American Naturalist*, *144*(3), 457–472. https://doi.org/10.1086/285686

Barabás, G., D'Andrea, R., & Stump, S. M. (2018). Chesson's coexistence theory. *Ecological Monographs*, *88*(3), 277–303. https://doi.org/10.1002/ecm.1302

Bashey, F. (2015). Within-host competitive interactions as a mechanism for the maintenance of parasite diversity. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *370*(1675), 20140301. https://doi.org/10.1098/rstb.2014.0301

Bates, A., Hilton, B., & Harley, C. (2009). Effects of temperature, season and locality on wasting disease in the keystone predatory sea star Pisaster ochraceus. *Diseases of Aquatic Organisms*, *86*, 245–251. https://doi.org/10.3354/dao02125

Bernhardt, J. R., Sunday, J. M., Thompson, P. L., & O'Connor, M. I. (2018). Nonlinear averaging of thermal experience predicts population growth rates in a thermally variable environment. *Proceedings of the Royal Society B: Biological Sciences*, *285*(1886), 20181076. https://doi.org/10.1098/rspb.2018.1076

Brook, C. E., & Dobson, A. P. (2015). Bats as 'special' reservoirs for emerging zoonotic pathogens. *Trends in Microbiology*, *23*(3), 172–180. https://doi.org/10.1016/j.tim.2014.12.004

Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., & West, G. B. (2004). Toward a metabolic theory of ecology. *Ecology*, *85*(7), 1771–1789. https://doi.org/10.1890/03-9000

Bruno, J. F., Selig, E. R., Casey, K. S., Page, C. A., Willis, B. L., Harvell, C. D., Sweatman, H., & Melendy, A. M. (2007). Thermal Stress and Coral Cover as Drivers of Coral Disease Outbreaks. *PLoS Biology*, *5*(6), e124. https://doi.org/10.1371/journal.pbio.0050124

Butler, M. W., Stahlschmidt, Z. R., Ardia, D. R., Davies, S., Davis, J., Guillette, L. J., Johnson, N., McCormick, S. D., McGraw, K. J., & DeNardo, D. F. (2013). Thermal Sensitivity of Immune Function: Evidence against a Generalist-Specialist Trade-Off among Endothermic and Ectothermic Vertebrates. *The American Naturalist*, *181*(6), 761–774. https://doi.org/10.1086/670191

Calisher, C. H., Childs, J. E., Field, H. E., Holmes, K. V., & Schountz, T. (2006). Bats: Important Reservoir Hosts of Emerging Viruses. *Clinical Microbiology Reviews*, *19*(3), 531–545. https://doi.org/10.1128/CMR.00017-06

Chesson, P. (1994). Multispecies Competition in Variable Environments. *Theoretical Population Biology*, *45*(3), 227–276. https://doi.org/10.1006/tpbi.1994.1013

Chesson, P. (2000). Mechanisms of Maintenance of Species Diversity. *Annual Review of Ecology and Systematics*, *31*(1), 343–366. https://doi.org/10.1146/annurev.ecolsys.31.1.343

Chesson, P. (2003). Quantifying and testing coexistence mechanisms arising from recruitment fluctuations. *Theoretical Population Biology*, *64*(3), 345–357. https://doi.org/10.1016/S0040-5809(03)00095-9

Chesson, P., & Huntly, N. (1997). The Roles of Harsh and Fluctuating Conditions in the Dynamics of Ecological Communities. *The American Naturalist*, *150*(5), 519–553. https://doi.org/10.1086/286080

Cizauskas, C. A., Carlson, C. J., Burgio, K. R., Clements, C. F., Dougherty, E. R., Harris, N. C., & Phillips, A. J. (2017). Parasite vulnerability to climate change: An evidence-based functional trait approach. *Royal Society Open Science*, *4*(1), 160535. https://doi.org/10.1098/rsos.160535

Cohen, J. M., Civitello, D. J., Venesky, M. D., McMahon, T. A., & Rohr, J. R. (2019). An interaction between climate change and infectious disease drove widespread amphibian declines. *Global Change Biology*, *25*(3), 927–937. https://doi.org/10.1111/gcb.14489

Connell, J. H. (1978). Diversity in Tropical Rain Forests and Coral Reefs: High diversity of trees and corals is maintained only in a nonequilibrium state. *Science*, *199*(4335), 1302–1310. https://doi.org/10.1126/science.199.4335.1302

Dallas, T., & Drake, J. M. (2016). Fluctuating temperatures alter environmental pathogen transmission in a *Daphnia* – pathogen system. *Ecology and Evolution*, *6*(21), 7931–7938. https://doi.org/10.1002/ece3.2539

Daszak, P., Berger, L., Cunningham, A. A., Hyatt, A. D., Green, D. E., & Speare, R. (1999). Emerging Infectious Diseases and Amphibian Population Declines. *Emerging Infectious Diseases*, *5*(6), 735–748. https://doi.org/10.3201/eid0506.990601

Daszak, P., Cunningham, A. A., & Hyatt, A. D. (2000). Emerging Infectious Diseases of Wildlife—Threats to Biodiversity and Human Health. *Science*, *287*(5452), 443–449. https://doi.org/10.1126/science.287.5452.443

Daszak, P., Cunningham, A. A., & Hyatt, A. D. (2001). Anthropogenic environmental change and the emergence of infectious diseases in wildlife. *Acta Tropica*, *78*(2), 103–116. https://doi.org/10.1016/S0001-706X(00)00179-0

Dell, A. I., Pawar, S., & Savage, V. M. (2014). Temperature dependence of trophic interactions are driven by asymmetry of species responses and foraging strategy. *Journal of Animal Ecology*, *83*(1), 70–84. https://doi.org/10.1111/1365-2656.12081

Denny, M. (2017). The fallacy of the average: On the ubiquity, utility and continuing novelty of Jensen's inequality. *Journal of Experimental Biology*, *220*(2), 139–146. https://doi.org/10.1242/jeb.140368

Dobson, A. P., & Carper, R. E. (1992). Global Warming and Potential Changes in Host-Parasite and Disease-Vector Relationships. In R. L. Peters & T. E. Lovejoy (Eds.), *Global Warming and Biological Diversity* (pp. 201–207). Yale University Press.

Duncan, A. B., Fellous, S., & Kaltz, O. (2011). Temporal variation in temperature determines disease spread and maintenance in *Paramecium* microcosm populations. *Proceedings of the Royal Society B: Biological Sciences*, *278*(1723), 3412–3420. https://doi.org/10.1098/rspb.2011.0287

Estay, S. A., Clavijo-Baquet, S., Lima, M., & Bozinovic, F. (2011). Beyond average: An experimental test of temperature variability on the population dynamics of *Tribolium confusum*. *Population Ecology*, *53*(1), 53–58. https://doi.org/10.1007/s10144-010-0216-7

Fumagalli, M. R., Zapperi, S., & La Porta, C. A. M. (2021). Role of body temperature variations in bat immune response to viral infections. *Journal of The Royal Society Interface*, *18*(180), 20210211. https://doi.org/10.1098/rsif.2021.0211

Halpin, K., Young, P. L., Field, H. E., & Mackenzie, J. S. (2000). Isolation of Hendra virus from pteropid bats: A natural reservoir of Hendra virus. *Journal of General Virology*, *81*(8), 1927–1932. https://doi.org/10.1099/0022-1317-81-8-1927

Hardin, G. (1960). The Competitive Exclusion Principle. *Science*, *131*(3409), 1292–1297. https://doi.org/10.1126/science.131.3409.1292

Holt, R. D. (1977). Predation, apparent competition, and the structure of prey communities. *Theoretical Population Biology*, *12*(2), 197–229. https://doi.org/10.1016/0040-5809(77)90042-9

Holt, R. D., & Lawton, J. H. (1994). THE ECOLOGICAL CONSEQUENCES OF SHARED NATURAL ENEMIES. *Annual Review of Ecology and Systematics*, *25*(1), 495–520. https://doi.org/10.1146/annurev.es.25.110194.002431

Hutchinson, G. E. (1961). The Paradox of the Plankton. *The American Naturalist*, *95*(882), 137–145. https://doi.org/10.1086/282171

Irving, A. T., Ahn, M., Goh, G., Anderson, D. E., & Wang, L.-F. (2021). Lessons from the host defences of bats, a unique viral reservoir. *Nature*, *589*(7842), 363–370. https://doi.org/10.1038/s41586-020-03128-0

Jensen, J. L. W. V. (1906). Sur les fonctions convexes et les inégalités entre les valeurs moyennes. *Acta Mathematica*, *30*(0), 175–193. https://doi.org/10.1007/BF02418571

Jiang, L., & Patel, S. N. (2008). Community assembly in the presence of disturbance: A microcosm experiment. *Ecology*, *89*(7), 1931–1940. https://doi.org/10.1890/07-1263.1

Johnson, P. T. J. (2006). Amphibian diversity: Decimation by disease. *Proceedings of the National Academy of Sciences*, *103*(9), 3011–3012. https://doi.org/10.1073/pnas.0600293103

Johnson, P. T. J., Preston, D. L., Hoverman, J. T., & LaFonte, B. E. (2013). Host and parasite diversity jointly control disease risk in complex communities. *Proceedings of the National Academy of Sciences*, *110*(42), 16916–16921. https://doi.org/10.1073/pnas.1310557110

Krichel, L., Kirk, D., Pencer, C., Hönig, M., Wadhawan, K., & Krkošek, M. (2023). Short-term temperature fluctuations increase disease in a Daphnia-parasite infectious disease system. *PLOS Biology*, *21*(9), e3002260. https://doi.org/10.1371/journal.pbio.3002260

Lawson, C. R., Vindenes, Y., Bailey, L., & van de Pol, M. (2015). Environmental variation and population responses to global change. *Ecology Letters*, *18*(7), 724–736. https://doi.org/10.1111/ele.12437

Lips, K. R., Diffendorfer, J., Mendelson, J. R., & Sears, M. W. (2008). Riding the Wave: Reconciling the Roles of Disease and Climate Change in Amphibian Declines. *PLoS Biology*, *6*(3), e72. https://doi.org/10.1371/journal.pbio.0060072

Luo, J., Greif, S., Ye, H., Bumrungsri, S., Eitan, O., & Yovel, Y. (2021). Flight rapidly modulates body temperature in freely behaving bats. *Animal Biotelemetry*, *9*(1), 45. https://doi.org/10.1186/s40317-021-00268-6

Mackey, R. L., & Currie, D. J. (2001). *The Diversity-Disturbance Relationship: Is It Generally Strong and Peaked?*

Miller, A. D., Roxburgh, S. H., & Shea, K. (2011). How frequency and intensity shape diversity–disturbance relationships. *Proceedings of the National Academy of Sciences*, *108*(14), 5643–5648. https://doi.org/10.1073/pnas.1018594108

Miller, A. D., Roxburgh, S. H., & Shea, K. (2012). Timing of disturbance alters competitive outcomes and mechanisms of coexistence in an annual plant model. *Theoretical Ecology*, *5*(3), 419–432. https://doi.org/10.1007/s12080-011-0133-1

Molnár, P. K., Sckrabulis, J. P., Altman, K. A., & Raffel, T. R. (2017). Thermal Performance Curves and the Metabolic Theory of Ecology—A Practical Guide to Models and Experiments for Parasitologists. *Journal of Parasitology*, *103*(5), 423. https://doi.org/10.1645/16-148

Paull, S. H., LaFonte, B. E., & Johnson, P. T. J. (2012). Temperature-driven shifts in a host-parasite interaction drive nonlinear changes in disease risk. *Global Change Biology*, *18*(12), 3558–3567. https://doi.org/10.1111/gcb.12018

Pedersen, A. B., & Fenton, A. (2007). Emphasizing the ecology in parasite community ecology. *Trends in Ecology & Evolution*, *22*(3), 133–139. https://doi.org/10.1016/j.tree.2006.11.005

Plowright, R. K., Parrish, C. R., McCallum, H., Hudson, P. J., Ko, A. I., Graham, A. L., & Lloyd-Smith, J. O. (2017). Pathways to zoonotic spillover. *Nature Reviews Microbiology*, *15*(8), 502–510. https://doi.org/10.1038/nrmicro.2017.45

Råberg, L., De Roode, J. C., Bell, A. S., Stamou, P., Gray, D., & Read, A. F. (2006). The Role of Immune-Mediated Apparent
Competition in Genetically Diverse Malaria Infections. *The American Naturalist*, *168*(1), 41–53.
https://doi.org/10.1086/505160
Raffel, T. R., Romansic, J. M., Halstead, N. T., McMahon, T. A., Venesky, M. D., & Rohr, J. R. (2013). Disease and thermal
acclimation in a more variable and unpredictable climate. *Nature Climate Change*, *3*(2), 146–151.
https://doi.org/10.1038/nclimate1659
Rohr, J. R., & Raffel, T. R. (2010). Linking global climate and temperature variability to widespread amphibian declines
putatively caused by disease. *Proceedings of the National Academy of Sciences*, *107*(18), 8269–8274.
https://doi.org/10.1073/pnas.0912883107
Rohr, J. R., Raffel, T. R., Blaustein, A. R., Johnson, P. T. J., Paull, S. H., & Young, S. (2013). Using physiology to
understand climate-driven changes in disease and their implications for conservation. *Conservation Physiology*, *1*(1),
cot022–cot022. https://doi.org/10.1093/conphys/cot022
Rohr, J. R., Raffel, T. R., Romansic, J. M., McCallum, H., & Hudson, P. J. (2008). Evaluating the links between climate,
disease spread, and amphibian declines. *Proceedings of the National Academy of Sciences*, *105*(45), 17436–17441.
https://doi.org/10.1073/pnas.0806368105
Roxburgh, S. H., Shea, K., & Wilson, J. B. (2004). The intermediate disturbance hypothesis: Patch dynamics and mechanisms
of species coexistence. *Ecology*, *85*(2), 359–371. https://doi.org/10.1890/03-0266
Sales, L. P., Hayward, M. W., & Loyola, R. (2021). What do you mean by "niche"? Modern ecological theories are not
coherent on rhetoric about the niche concept. *Acta Oecologica*, *110*, 103701.
https://doi.org/10.1016/j.actao.2020.103701
Schoolfield, R. M., Sharpe, P. J. H., & Magnuson, C. E. (1981). Non-linear regression of biological temperature-dependent
rate models based on absolute reaction-rate theory. *Journal of Theoretical Biology*, *88*(4), 719–731.
https://doi.org/10.1016/0022-5193(81)90246-0
Schulte, P. M., Healy, T. M., & Fangue, N. A. (2011). Thermal Performance Curves, Phenotypic Plasticity, and the Time
Scales of Temperature Exposure. *Integrative and Comparative Biology*, *51*(5), 691–702.
https://doi.org/10.1093/icb/icr097
Sckrabulis, J. P., Altman, K. A., & Raffel, T. R. (2022). Using Metabolic Theory to Describe Temperature and Thermal
Acclimation Effects on Parasitic Infection. *The American Naturalist*, *199*(6), 789–803.
https://doi.org/10.1086/719409
Sinclair, B. J., Marshall, K. E., Sewell, M. A., Levesque, D. L., Willett, C. S., Slotsbo, S., Dong, Y., Harley, C. D. G.,
Marshall, D. J., Helmuth, B. S., & Huey, R. B. (2016). Can we predict ectotherm responses to climate change using
thermal performance curves and body temperatures? *Ecology Letters*, *19*(11), 1372–1385.
https://doi.org/10.1111/ele.12686
Song, C., & Spaak, J. W. (2024). Trophic tug-of-war: Coexistence mechanisms within and across trophic levels. *Ecology*
*Letters*, *27*(4), e14409. https://doi.org/10.1111/ele.14409
Spaak, J. W., & De Laender, F. (2020). Intuitive and broadly applicable definitions of niche and fitness differences. *Ecology*
*Letters*, *23*(7), 1117–1128. https://doi.org/10.1111/ele.13511
Spaak, J. W., Godoy, O., & De Laender, F. (2021). Mapping species niche and fitness differences for communities with
multiple interaction types. *Oikos*, *130*(12), 2065–2077. https://doi.org/10.1111/oik.08362
Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., & Midgley, P. M.
(Eds.). (2013). *Climate change 2013: The physical science basis. In: Contribution of Working Group I to the Fifth*
*Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press.
Symons, C. C., & Arnott, S. E. (2014). Timing is everything: Priority effects alter community invasibility after disturbance.
*Ecology and Evolution*, *4*(4), 397–407. https://doi.org/10.1002/ece3.940
Tracy, A. M., Pielmeier, M. L., Yoshioka, R. M., Heron, S. F., & Harvell, C. D. (2019). Increases and decreases in marine
disease reports in an era of global change. *Proceedings of the Royal Society B: Biological Sciences*, *286*(1912),
20191718. https://doi.org/10.1098/rspb.2019.1718
Tuljapurkar, S. (1990). *Population Dynamics in Variable Environments* (Vol. 85). Springer Berlin Heidelberg.
https://doi.org/10.1007/978-3-642-51652-8
Vandermeer, J. H. (1972). Niche Theory. *Annual Review of Ecology, Evolution, and Systematics*, *3*(Volume 3, 1972), 107–
132. https://doi.org/10.1146/annurev.es.03.110172.000543
Vasseur, D. A., DeLong, J. P., Gilbert, B., Greig, H. S., Harley, C. D. G., McCann, K. S., Savage, V., Tunney, T. D., &
O'Connor, M. I. (2014). Increased temperature variation poses a greater risk to species than climate warming.
*Proceedings of the Royal Society B: Biological Sciences*, *281*(1779), 20132612.
https://doi.org/10.1098/rspb.2013.2612
Williamson, M. H. (1957). An Elementary Theory of Interspecific Competition. *Nature*, *180*(4583), 422–425.
https://doi.org/10.1038/180422a0
Wootton, J. T. (1998). Effects of Disturbance on Species Diversity: A Multitrophic Perspective. *The American Naturalist*,
*152*(6), 803–825. https://doi.org/10.1086/286210

Wright, R. K., & Cooper, E. L. (1981). Temperature effects on ectotherm immune responses. *Developmental and*
*Comparative Immunology*, *5*, 117–122.
708