Imidacloprid disrupts larval molting regulation and nutrient energy

metabolism, causing developmental delay in honey bee *Apis mellifera*

Zhi Li [a,b,*,1], Yuedi Wang [a,b,1], Qiqian Qin [a,b], Lanchun Chen [a,b],Xiaoqun Dang [a,b],

Zhengang Ma [a,b], Zeyang Zhou [a,b,c,d]

[a] College of Life Sciences, Chongqing Normal University, Chongqing, China

[b] Key Laboratory of Pollinator Resources Conservation and Utilization of the Upper Yangtze River, Ministry
of Agriculture and Rural Affairs, Chongqing, China

[c] Chongqing Key Laboratory of Microsporidia Infection and Control, China

[d] The State Key Laboratory of Silkworm Genome Biology, Southwest University, Chongqing, China.

* Corresponding author. E-mail address: lizhicqnu@gmail.com (Z. Li).
[1] These authors contributed equally to the study
13

Abstract

Imidacloprid is a global health threat that severely poisons the economically and

ecologically important honeybee pollinator, *Apis mellifera*. However, its effects on

developing bee larvae remain largely unexplored. Our pilot study showed that

imidacloprid causes developmental delay in bee larvae, but the underlying toxicological

mechanisms remain incompletely understood. In this study, we exposed bee larvae to

imidacloprid at environmentally relevant concentrations of 0.7, 1.2, 3.1, and 377 ppb.

There was a marked dose-dependent delay in larval development, characterized by

reductions in body mass, width, and growth index. However, imidacloprid did not affect

on larval survival and food consumption. The primary toxicological effects induced by

elevated concentrations of imidacloprid (377 ppb) included inhibition of neural

transmission gene expression, induction of oxidative stress, gut structural damage, and

apoptosis, inhibition of developmental regulatory hormones and genes, suppression of

gene expression levels involved in proteolysis, amino acid transport, protein synthesis, carbohydrate catabolism, oxidative phosphorylation, and glycolysis energy production. In addition, we found that the larvae may use antioxidant defenses and P450 detoxification mechanisms to mitigate the effects of imidacloprid. Ultimately, this study provides the first evidence that environmentally exposed imidacloprid can affect the growth and development of bee larvae by disrupting molting regulation and limiting the metabolism and utilization of dietary nutrients and energy. These findings have broader implications for studies assessing pesticide hazards in other juvenile animals.

Keywords: Imidacloprid; *Apis mellifera*; Toxicity effects; Developmental retardation; Juvenile animal;

1. Introduction

Hazardous pesticides are used extensively around the world to protect crop production, particularly in agriculture-dependent developing countries. However, exposure to these chemicals has become a global health concern. Imidacloprid, introduced in 1991, is known for its high animal toxicity and has received international attention (Jeschke et al., 2011). The World Health Organization and the U.S. Environmental Protection Agency have listed it as an acute oral and dermal toxicant for animals. The excessive use of imidacloprid can lead to environmental contamination, endangering human and animal health through food chains, poisoning non-target pollinators, and disrupting ecological balances (Tudi et al., 2021).

Imidacloprid is also known for its adverse effects on human health. Adults exposed to high doses of imidacloprid (9.6%) experience dizziness, apathy, movement disorders, labored breathing, and temporary growth retardation (Wu et al., 2001). Imidacloprid also causes adverse congenital, hematological, hepatic, and renal effects, along with degenerative changes in various organs (Khan et al., 2010; Shaw et al., 2014; Yang et al.,

2014). Imidacloprid even at low doses causes severe liver, thyroid, and body weight

problems, reproductive toxicity, developmental retardation, and neurobehavioral

deficits in rats and rabbits (Anatra-Cordone M, 2005). Additionally, imidacloprid affects

the singing ability of birds, making it difficult for them to attract mates and reproduce. It

is a significant contributor to the decline in terrestrial insect populations since 1991,

averaging 9% per decade (Cardoso et al., 2020; Sánchez-Bayo and Wyckhuys, 2019).

Honey bees (*Apis mellifera* L.) are valuable model organisms in insects with significant

economic and ecological importance as crop pollinators and in maintaining ecological

balance (Zheng and Fu-Liang, 2009). However, the widespread use of pesticides,

including imidacloprid, has resulted in colony collapse disorder (CCD), causing severe

impacts on the beekeeping industry and ecological balance (Mahé et al., 2021). Studies

have shown that imidacloprid is applied as a seed coating (Simon-Delso et al., 2015) and

accumulates in the plant (Dively and Kamel, 2012; Rocha et al., 2022), exposing non-

target bees to imidacloprid residues during pollen and nectar collection (Tong et al.,

2018). Even low levels of systemic imidacloprid cause sublethal effects in adult bees,

including neurotoxicity (Belzunces et al., 2012), behavioural changes (Karahan et al.,

2015; Medrzycki et al., 2003), reproductive impairment (Chaimanee et al., 2016),

reduced immunity (Pettis et al., 2013), apoptosis and autophagy (Carneiro et al., 2022),

and shortened lifespan (Anderson and Harmon-Threatt, 2021). A recent finding shows

that imidacloprid induces oxidative stress in honey bees (Balieira et al., 2018). This

oxidative stress is a result of the long-term synergistic effect between imidacloprid and

its target receptors (nAChRs) (Makoto et al., 2020) in bees, which leads to an excessive

influx of Ca^{2+} into the bee brain (Tahirare, 2013), disrupting Ca^{2+} homeostasis and

ultimately causing mitochondrial dysfunction and increased mitochondrial ROS

generation, exacerbating oxidative stress.

The development and health status of juvenile animals play a critical role in

population structure and future growth (Gill et al., 2012). However, they are more

susceptible to the toxic effects of pesticides than adults. Despite this, there is limited

information on the toxic effects of imidacloprid on juvenile animals. Available reports mainly focus on the epistatic effects of toxicity, with limited investigation into the involved biochemical mechanisms. Imidacloprid is known to pose a potential health risk to children through dietary exposure (Anderson and Harmon-Threatt, 2021), and may cause lead to developmental abnormalities in amphibians (Samojeden et al., 2022) and fish (Islam et al., 2019). Furthermore, negative effects on pupal development in insects have been observed (Jia and Li, 2023). For instance, cat fleas (*Ctenocephalides felis*) larvae treated with imidacloprid become immobile, and their intestines show pulsatile movements for up to one hour, eventually dying after two hours (Mehlhorn et al., 1999). Additionally, imidacloprid exposure can affect the average weight gain per day in *Harmonia axyridis* (Coleoptera: Coccinellidae) larvae (Charles et al., 2000). It has also been found that imidacloprid exposure affects various cellular aspects, such as mitochondria, energy, lipids, and the transcriptome in Drosophila (Felipe et al., 2020), and prolongs the larval pupation duration and affects forewing development in the butterfly *Pieris brassicae* (Whitehorn et al., 2018).

In bees, larvae play a crucial role in the growth and development of colonies. However, the broader toxic effects of imidacloprid on bee larvae remain to be investigated. Limited reports suggest that developing bee larvae are exposed to imidacloprid through contact with contaminated hives or ingestion of contaminated food (Bhme et al., 2018). However, their survival, developmental rate, and body weight are not affected (Dai et al., 2019). Nevertheless, other studies have shown that imidacloprid can lead to impaired olfactory learning behavior (Peng and Yang, 2016), and other behavioral abnormalities (Wu et al., 2017). The current understanding is that honey bee larvae are more tolerant to imidacloprid than adults, as they can survive treatment and develop into adults (Dai et al., 2017). However, they may experience developmental delays, such as delayed plumage (Woyciechowski and Moroń, 2009; Wu et al., 2011), and reduced rates of capping, pupation, and eclosion (Yang et al., 2012). A recent study using next-generation sequencing has indicated that sublethal imidacloprid

treatment during the larval stage causes changes in gene expression in larvae, pupae, and adults, suggesting a persistent sublethal impact on honey bee development (Chen et al., 2021). Furthermore, a latest study has demonstrated that imidacloprid induces differential gene expression in bumble bee larvae, resulting in unique expression patterns related to various biological processes such as starvation response, cuticle genes, neural development, and cell growth (Martín-Blázquez et al., 2023).

Although progress has been made in understanding the toxic effects of imidacloprid on honey bee larvae, the molecular basis of this toxicity remains poorly understood. Further research is needed to uncover the mechanisms and pathways involved in the developmental toxicity of imidacloprid in honey bee larvae. In our pilot study, we observed a significant developmental delay in bee larvae exposed to imidacloprid; However, the molecular and biochemical mechanisms underlying this delay have not been systematically investigated. Therefore, this study aims to comprehensively evaluate the toxicity of imidacloprid on honey bee larvae at multiple levels, including histology, neurotoxicity, molting regulation, oxidative stress, detoxification, nutrition, and energy metabolism. Our study provides first-hand evidence of the developmental delay caused by imidacloprid in bee larvae. Because these mechanisms involved are relatively conserved in animals, our findings have broader implications for understanding and assessing the potential risk of continuous exposure to imidacloprid during animal development.

Results

Imidacloprid causes bee larval developmental retardation

The 96-hour observation period showed no significant difference in larval survival between the four groups exposed to imidacloprid and the control group. However, as the exposure concentration and duration increased, here was a significant decrease in larval developmental progress, weight, and width (Fig. 1A-1E). Further analysis was

conducted on the most adverse exposure level of 377 ppb, which is the highest reported
environmental residue concentration detected in bee products, such as beeswax
(Kapoor et al., 2014). It is important to note that while the tested doses of the 377 ppb
was found in bee products, it is likely that the bees themselves were exposed to even
higher doses. Out result demonstrate that exposure to 377 ppb imidacloprid did not
have a significant impact on survival, feeding (Fig. 1C, 1F, and 1G), and the time required
for larvae to enter the pupal stage successfully (Fig. 1H). However, it did lead to slow
developmental progress, a lower initial rate of successful pupation, and lower growth
index. Additionally, imidacloprid consistently resulted in lower larval weight (Fig. 1I) and
width (Fig. 1J) up to the pupal stage (Fig. 1B).

Imdacloprid neurotoxicity in bee larvae

Under imidacloprid environmental stress at 377 ppb, AChE activity, which is closely
related to the disruption of neurotransmission, was significantly inhibited in larvae (Fig.
2A). Neither acetylcholine receptor (nAChR) alpha 1 (*Alph1*) nor acetylcholinesterase 2
(*Ace2*) gene expression was significantly different from the control. In contrast,
acetylcholine receptor alpha 2 (*Alph2*) and acetylcholinesterase 1 (*Ace1*) gene
expression was significantly higher than that in the control group (Fig. 2B).

Imidacloprid disrupts the homeostasis of developmental regulation in larvae

In imidacloprid-exposed larvae, the developmental regulatory gene juvenile hormone
acid methyl transferase (*JHAMT*) and the transcription factor broad complex (*Br-c*) were
downregulated, while vitellogenin (*Vg*) gene expression was upregulated by seven-fold
in comparison to control groups (Fig. 3A). In parallel, imidacloprid exposure decreased
the major regulator of insect metamorphosis 20-hydroxyecdysone (20E) titers but had
no effect on preventing premature metamorphosis juvenile hormone (JH-3) titers (Fig.
3B).

Imidacloprid induces detoxification in larvae

According to the qRT-PCR analysis, the expression levels of cytochrome P450
monooxygenase family members such as CYPq1, CYPq2, CYP450, CYP6AS14, CYP4G11,
and CYP306A1, except for CYPq3, were observed to be significantly upregulated by
several fold in imidacloprid-induced developmental retardation larvae. Among them,
CYPq1 showed an abnormal 94-fold increase (Fig. 4).

Imidacloprid toxicity causes oxidative stress and induces antioxidant defense in larvae

Larvae exposed to imidacloprid exhibited a significant increase in reactive oxygen
species (ROS) levels (Fig. 5A) and upregulation of the antioxidant genes GPX and Trx
expression (Fig. 5C), as well as increased activities of the antioxidants CAT, SOD and GSH
(Fig. 5B), resulting in a significant increase in total antioxidant capacity (T-AOC) (Fig. 5B)
compared to the control group. Under these conditions, MDA levels, used as a marker of
lipid peroxidation, were observed to increase, while protein carbonylation damage
indicator PCO levels remained unchanged (Fig. 5A).

Imidacloprid induces gut apoptosis and tissue damage in larvae

Hematoxylin-eosin staining revealed a significant reduction in the number of cells in
the muscular layer of the larval gut exposed to imidacloprid with clear apoptotic
characteristics compared to the standard feeding control groups (Fig. 5D). This was
evidenced by increased nuclear staining and compacted chromatin, as well as disrupted
cell arrangement in the basal layer of the gut, increased nuclear staining of basal lamina
cells, and apoptotic signals. The integrity of the peritrophic membrane structure of the
gut was better in the control group, while imidacloprid-exposed larvae were still in the
formation stage, resulting in the accumulation of undigested food residues in their guts
(Fig. 5D).

Imidacloprid impairs larval digestion and breakdown of dietary nutrients

Exposure to imidacloprid did not significantly affect the daily food intake of
developmentally delayed larvae compared to controls (Fig. 1F and 1G). However,

imidacloprid toxicity suppressed the expression of nutrient catabolism genes, including

alpha-amylase (*α-Am*) and alpha-glucosidase (*α-Glu*), which are involved in

carbohydrate catabolism (Fig. 6A), carboxypeptidase (*CPs*) and aminopeptidase (*APs*)

involved in proteolysis (Fig. 6B), and glycine tRNA ligase (*Aats-Gly*) and tyrosine tRNA

ligase (*Aats-Tyr*) involved in amino acid transport (Fig. 6C).

Imidacloprid causes restriction of larval protein synthesis and energy metabolic

dysfunction

RT-qPCR analysis revealed that imidacloprid exposure reduced the expression of

critical genes involved in energy metabolism, including cytochrome c oxidase copper

chaperone (*COX17*) and NADH dehydrogenase [ubiquinone] 1β subcomplex subunit 7

(*NDUFB7*), in developmentally delayed larvae (Fig. 6D). Imidacloprid exposure also

inhibited the expression of genes related to glycolysis and energy production, such as

glyceraldehyde-3-phosphate dehydrogenase (*Gapdh*) and glucosamine-6-phosphate

isomerase (*Oscillin*) (Fig. 6E). In addition, the expression of transcription initiation

factors related to protein synthesis, including translation initiation factor 3 subunit M

(*Tango7*), translation initiation factor 3 subunit B (*elF3-S9*), and translation initiation

factor 3 subunit A (*elF3-S10*), was significantly suppressed by imidacloprid toxicity (Fig.

6F). Finally, the total levels of ATP, glycogen and protein, which are closely related to

energy production and reserves, were significantly lower in developmentally delayed

larvae than in controls (Fig. 6G-6I).

Correlation between developmental phenotypic changes and molecular features

caused by imidacloprid toxicity

To better understand the mechanism of imidacloprid-induced larval developmental

retardation, we performed a comprehensive correlation analysis between epigenetic

growth traits and molecular characteristics (Fig. 7). The results showed that in larvae

with imidacloprid-induced developmental retardation, body weight, width, growth

index, and developmental rate were positively correlated with energy metabolism and

reserves, nutrient catabolism, protein synthesis, and developmental regulation.
Conversely, negative correlations were observed between these growth traits and
parameters associated with detoxification and antioxidant defense processes that
require additional energy expenditure. Moreover, oxidative stress and damage,
detoxification, and antioxidant defense all exhibited significant positive correlations (Fig.
7).

Discussion

Toxic effects and the molecular basis of environmental concentrations of imidacloprid on bee larvae

Previous research has shown that bee larvae are more susceptible to the adverse
effects of pesticides than adult bees (Tomé et al., 2020). In this study, larvae were
exposed to environmental concentrations of imidacloprid, resulting in sublethal effects
such as decreased body weight, width, developmental rate, and growth index (Fig. 1).
Our results support the notion that pesticides can affect larval development (Dai et al.,
2018; Zhu et al., 2014). However, current research has primarily focused on
characterizing the effects of imidacloprid on larval phenotype and behavior, and the
molecular mechanisms underlying its ecotoxicological effects remain poorly understood.

To obtain molecular evidence of imidacloprid toxicity to bee larvae, we first examined
the expression of acetylcholine receptors and AChE activity. The results showed that
imidacloprid exposure increased the expression of the *Alph2* gene and inhibited AChE
activity (Fig. 2). Normally, acetylcholine stimulates nerve impulses by binding to the Alph
receptor, and AChE degrades acetylcholine to terminate nerve conduction. However,
imidacloprid competitively binds to AChE receptors, inhibits AChE activity, and causes
continuous excitation of nerve conduction, ultimately leading to paralysis and death
(Katić et al., 2021; Shan et al., 2020). Therefore, our results strongly suggest that
imidacloprid is neurotoxic to bee larvae.

The growth and development of juvenile animals depends on the digestion, absorption and utilization of food nutrients. Therefore, we next investigated whether imidacloprid affects the digestion and utilization of dietary by larvae. The results indicate that exposure to imidacloprid suppressed the hydrolysis genes *CPs* and *APs* and the amino acid transport pathway genes *Aats-Gly* and *Aats-Tyr* in larvae (Fig. 6B and 6C). Since these genes are responsible for cleaving dietary proteins into absorbable peptides and amino acids and transporting them for metabolism (2019), inhibiting the expression of these genes would severely affect larval digestion, resulting in protein deficiency and impaired metabolic activity. In addition, imidacloprid also affected the expression of *α-Am* and *α-Glu* genes (Fig. 6A), which are involved in carbohydrate digestion, indicating that imidacloprid also affects the digestion of dietary carbohydrates. Indeed, histological observations also confirmed that a large amount of undigested food remained in the intestine of larvae exposed to imidacloprid (Fig. 5D). In conclusion, imidacloprid inhibits the digestion and utilization of dietary proteins and carbohydrates by larvae, which is extremely detrimental to their growth and development.

In addition to nutrition, energy metabolism is critical for the maintenance of all life activities. In this study, imidacloprid exposure inhibited the expression of genes related to mitochondrial oxidative phosphorylation (*COX17*, *NDUFB7*) and its alternative glycolytic pathway (*Gapdh*, *Oscillin*). Mitochondrial oxidative phosphorylation is the central pathway for ATP generation during energy metabolism (Tian et al., 2020a), while glycolysis is an alternative mechanism for ATP synthesis (Jiang et al., 2020). Since both biochemical processes are essential for energy production, their inhibition resulted in reduced ATP levels in bee larvae (Fig. 6G). These results strongly suggest that imidacloprid induces energy metabolism dysfunction leading to insufficient energy production in larvae. Since larval energy production is insufficient, this is likely to lead to a depletion of energy reserves. Indeed, further analysis revealed a significant decrease in glycogen and protein energy reserves in larvae exposed to imidacloprid (Fig. 6H and 6I). These findings are consistent with previous reports that exposure to pesticides such

as endosulfan, chlorpyrifos, insecticides, and fipronil can lead to a reduction in energy reserves in organisms (Bouayad et al., 2012a; Dutra et al., 2009; Radwan et al., 2008; Rambabu and Rao, 1994; Ribeiro et al., 2001). In conclusion, our results suggest that imidacloprid induces energy metabolism dysfunction and affects energy production and reserves in bee larvae.

Insects rely primarily on cytochrome P450 detoxification to resist toxic compounds (Li et al., 2007), and the level of CYP450 expression indicates the severity of pesticide toxicity. While adult worker bees have shown a positive response to imidacloprid (Mao et al., 2009), it is uncertain whether bee larvae can respond to pesticide toxicity. In this study, we found that seven cytochrome P450 family transcripts, including CYPq1, CYPq2, CYP450, CYP6AS14, CYP4G11, and CYP306A1, were significantly upregulated in bee larvae, with CYP9q1 being 94-fold higher than the control group (Fig. 4). These results are consistent with previous findings that the CYP450 family genes CYP6AS3, CYP6AS4, and CYP9S1 were significantly upregulated in adult worker bees exposed to quercetin, coumaphos, and fluvalinate (Mao et al., 2011) and that CYP9Q1, CYP9Q2, and CYP9Q3 help bees detoxify fluvalinate and coumaphos (Gregorc et al., 2018). This suggests that the P450 detoxification system is already established in bee larvae and they respond positively by activating the detoxification function of CYP450 upon exposure to imidacloprid.

Oxidative stress is the excessive production of ROS in aerobic organisms (Felton, 1995; Ozcan and Ogun, 2015). Pesticides, including coumaphos (Gregorc et al., 2018), fipronil (Paris et al., 2017), paraquat (Li-Byarlay et al., 2016), organophosphorus, pyrethroid, organochlorinated (Chakrabarti et al., 2015), and chlorpyrifos (Shafiq-ur-Rehman et al., 2012), have been shown to cause oxidative stress in bees. In this study, exposure to imidacloprid resulted in increased levels of ROS and MDA in larvae (Fig. 5A), indicating severe oxidative stress and lipid damage. Organisms have evolved antioxidant defense mechanisms to scavenge excessive ROS and resist oxidative stress, including increasing

levels of antioxidants such as CAT, SOD, GSH, TrxR, and GSH (Dong et al., 2013; Tian et

al., 2020b). In this regard, we have recently confirmed this in worker bees (Li et al.,

2022), which is also well reflected in the present study. We observed a significant

increase in the activities of antioxidant enzymes SOD and CAT, gene expression of Trx

and Gpx, and GSH levels in larvae after exposure to imidacloprid (Fig. 5B and 5C). These

results show that the antioxidant system in developing larvae is established and used to

resist oxidative stress caused by imidacloprid, similar to that in adult worker bees.

Notably, compared to our recent study in adult bees (Li et al., 2022), the larvae in this

study exhibited more robust antioxidant activity when exposed to higher concentrations

of imidacloprid toxicity than worker bees. However, the underlying mechanism requires

further investigation. The underlying mechanisms involved require further study.

Notably, in addition to routine antioxidants, imidacloprid induces a 7-fold increase in

Vg expression (Fig. 3A). Vg is known to provide nutrition to developing embryos

(Jorgensen et al., 2009) and to help animals cope with pollution-induced stress (Matozzo

et al., 2008). In bees, Vg plays an important antioxidant role by protecting other

hemolymph molecules from ROS and increasing cellular oxidative stress tolerance

(Corona et al., 2007). Previous studies have shown a correlation between high Vg

expression in adult worker bees and resistance to oxidative stress resulting from

exposure to the pesticide paraquat (Seehuus et al., 2006). In this study, our results

support this observation and suggest that larvae may also resist oxidative stress caused

by imidacloprid through increased Vg production.

Mechanisms of imidacloprid-induced developmental retardation in bee larvae

Impaired juvenile development can have a significant impact on the growth and

sustainability of entire populations. Deviations in juvenile development due to

environmental stress have recently received considerable attention. *Drosophila*

melanogaster exposed to heat stress exhibit developmental delays due to the failure of

early embryos to fully inhibit the synthesis of non-heat shock proteins (Bergh and Arking,

1984). Similarly, exposure to nutrient and heavy metal stress can cause delayed or

arrested development in *Caenorhabditis elegans* due to altered gene expression and

disruption of developmental signaling pathways (Carranza-García and Navarro, 2020;

Rashid et al., 2021). In the present study, we also observed typical developmental delays

in honeybee larvae exposed to imidacloprid, but the underlying mechanisms remain

unclear.

Insect development is regulated by the hormones JH and 20E, which control

morphological remodeling events during molting. 20E promotes pupation (Yuan et al.,

2020), while JH maintains juvenile characteristics and prevents metamorphosis by

antagonizing the effects of 20E (Luo et al., 2021). To elucidate the mechanism

underlying imidacloprid-induced larval developmental delay, we first focused on the

hormonal regulation of molting development and found that imidacloprid inhibits the

expression of JHAMT (Fig. 3A), a positive regulator of JH synthesis (Liu et al., 2018), but

did not affect JH titer levels (Fig. 3B). This suggests that the antagonistic effect of JH still

restricts the developmental process, thereby maintaining larval morphology. In parallel,

imidacloprid toxicity also resulted in reduced ecdysteroid 20E titer and Br-c expression

(Fig. 3). The role of 20E is to induce larval molting and pupation by triggering

downstream response factors, such as Br-c (Deng et al., 2012). Therefore, our results

suggest that larval developmental delay may be caused by imidacloprid reducing 20E

titer and inhibiting *Br-c* expression, thereby blocking molting. Notably, we also observed

that a decrease in 20E levels was accompanied by a decrease in AChE activity (Fig. 2A),

which is known to cause delayed pupation (He et al., 2012) and developmental arrest

(Desneux et al., 2007) in insect larvae. Therefore, our results suggest that imidacloprid

neurotoxicity may cause developmental delay in bee larvae by inhibiting the molting

hormone 20E. This conclusion is supported by a correlation analysis showing that

decreases in 20E titer, *Br-c* expression and AChE activity were positively correlated with

decreased developmental rates and growth index, weight and width growth (Fig. 7). In

conclusion, imidacloprid neurotoxicity inhibits the 20E and *Br-c* genes, thereby blocking
molting, which may be an important cause of delayed larval development in honey bees.

Animals employ selective foraging as a survival strategy to avoid ingesting toxic foods
that may cause adverse physiological effects (Berenbaum and Johnson, 2015). Although
bees cannot actively reject toxins innately, postingestive malaise allows them to learn to
avoid ingesting toxic-containing foods with adverse physiological effects (Hurst et al.,
2014). Given this, we speculate that it is likely that the honey bee larvae in this study
reduced their intake of toxic imidacloprid-containing foods, making it difficult to meet
their nutritional and energy needs, which may have resulted in delayed development.
However, monitoring of daily food intake did not show a significant difference between
larvae exposed to imidacloprid and the control group (Fig. 1F and 1G), indicating that
larval food intake was not reduced and larvae did not initiate avoidance behavior
toward toxic food. Therefore, food intake was not a critical factor in the delayed larval
development caused by imidacloprid.

Nutrition and energy are critical for the growth and development of juvenile animals.
Exposure to heavy metals and neonicotinoid pesticides has been shown to affect
digestion, energy reserves, production, and utilization in animals (Jiang et al., 2020; Li et
al., 2021). Given this, since the lack of the necessary correlation between delayed larval
development and food intake in this study prompts us to consider potential problems in
larval food and energy utilization that led to delayed development. Consequently, our
focus shifted to the investigation of larval nutrition and energy metabolism.
Interestingly, we found severe pathological changes in the gut tissues of
developmentally delayed larvae exposed to imidacloprid (Fig. 5D). Imidacloprid caused a
significant reduction in the tightness of the cell arrangement in the basal layer of the gut,
and a drastic decrease in the number of cells. The peritrophic membrane was
incompletely formed, and undigested food residues were present in the gut. qRT-PCR
analysis revealed a significant decrease in larval digestion and catabolism of dietary

proteins and carbohydrates, amino acid transport, oxidative phosphorylation, and

glycolysis (Fig. 6D-6F). Biochemical assays showed that imidacloprid stress also

decreased ATP levels and total glycogen and protein content, resulting in a lack of

energy reserves and increased energy expenditure (Fig. 6G-6I). Correlation analysis

indicated a positive relationship between impaired catabolism and dietary nutrient

utilization caused by imidacloprid and larval developmental delay (Fig. 7). These

evidence strongly suggest that imidacloprid exposure affects the digestion and

utilization of nutrients and energy in larvae, resulting in insufficient nutrients and energy

for growth and development, which may be another important factor contributing to

imidacloprid-induced larval developmental delay.

The Dynamic Energy Budget (DEB) theory suggests that environmental stress can

significantly affect the energy balance of organisms, leading to increased energy

expenditure (Kooijman, 2009). To cope with stress, animals may deplete their energy

reserves, such as glycogen and protein (Bouayad et al., 2012b; Matsukura et al., 2008).

In this study, we found that larvae with delayed development caused by imidacloprid

had lower energy reserves (ATP, total protein, and total glycogen), and these energy

reserves were negatively correlated with larval antioxidant and detoxification (Fig. 7),

suggesting that the reduction in energy reserves caused by imidacloprid is related to the

energy-consuming activities of CYP450 detoxification and antioxidant defense. To meet

the high energy cost demands of environmental stress, animals often have to allocate

most of their food intake to defense (Guedes et al., 2006), and this redistribution of

energy resources will inevitably lead to an insufficient energy supply for growth and

development (Beyers et al., 1999). Therefore, the P450 detoxification and antioxidant

defense used by the larvae in this study to resist imidacloprid toxicity will inevitably

force them to consume more food resources that should have been used for growth and

development, resulting in a severe lack of energy needed for development. This may

also be another important reason why imidacloprid causes delayed development in bee

larvae.

It is worth noting that our investigation into the causes of imidacloprid-induced

developmental delay in honey bee larvae has provided limited answers. Future

comparative analyses of different tissues and durations of exposure, using genetic

manipulation and histochemical methods, would improve our understanding of the

effects of sublethal doses of imidacloprid on larval development, growth and survival.

Conclusion

Imidacloprid toxicity caused growth and developmental delay in *A. mellifera* larvae.

The toxic mechanism may include imidacloprid disrupting the regulatory balance of molt

development, resulting in restricted development. In addition, the gut damage caused

by imidacloprid restricts the metabolism and utilization of food nutrients and energy in

the larvae. Third, the additional energy consumed by larval P450 detoxification and

antioxidant defenses further reduces the supply of nutrients and energy for growth and

development (Fig. 8). This study is the first multi-level investigation of the toxic effects

of imidacloprid on *A. mellifera* larvae. These findings have broader implications for

understanding and assessing the threat of harmful pesticides to animal larvae.

Materials and Methods

Larval rearing

The Animal Bioethics Committee at Chongqing Normal University, China approved all

experimental protocols. The chosen colonies were healthy and not exposed to

pathogens or pesticides. Two-day-old larvae from the same frames of the same hive

were individually transferred to sterile 24-well cell culture plates. The plates contained

standard food, which included royal jelly, glucose, fructose, water, and yeast extract

(Aupinel et al., 2005). The larvae were cultured in dark conditions at 35°C and 96% RH,

with daily feeding volumes adjusted according to their age (20 µL for three-day-old

larvae, 30 µL for four-day-old larvae, 40 µL for five-day-old larvae, and 50 µL for six-day-old larvae). Any remaining food was recorded and removed.

Experimental design

This study employed four different concentrations of imidacloprid, which were residual concentrations found in honey (0.7 ppb) (Chauzat et al., 2009), bees (1.2 ppb) (Chauzat et al., 2011), pollen (3.1 ppb) (Mullin et al., 2010), and beeswax (377 ppb) (Pareja et al., 2011). The experiment was conducted with three treatment groups: control group (CK), solvent control group (CKac), and imidacloprid-treated group (IMI), each with three replicates. Each replicate contained 32 bees that were distributed in standard 24-well cell culture plates (1 bee/well). The CK group consisted of larval bees fed acetone-royal jelly glucose-fructose-yeast extract-distilled water (0.00377%-50%-6%-6%-1%-36.99623%, v/w/w/w/w/v). The IMI group consisted of larvae-fed control solution containing imidacloprid (99% active ingredient, Hubei Norna Technology Co. Ltd, Hubei, China) at 0.7 ppb, 1.2 ppb, 3.1 ppb, and 377 ppb. The CKac group served as a solvent control and contained only acetone at a concentration of 0.00377%. A total of 312 larvae were used for survival statistics (96 bees), body weight and width (96 bees), and gene expression, enzyme activity, histochemical content, and histopathologic analysis. Food was administered daily according to age (20 µL for three-day-old larvae, 30 µL for four-day-old larvae, 40 µL for five-day-old larvae, and 50 µL for six-day-old larvae). Dead individuals and food remain were recorded daily, and incubation conditions were 35°C and 96% RH. Surviving larvae were collected 72 h after consumption of the test solution and stored at −80°C for further analysis, including evaluations of gene expression, enzyme activity, and chemical content evaluations. Furthermore, HE-stained sections were used to evaluate any imidacloprid-induced tissue damage and apoptosis.

Developmental phenotypic traits

Larval individuals were randomly selected daily from the CK and IMI groups to

measure body weight and width. Food residues, larval mortality, developmental

progression, and the amount of time spent in the larval developmental stage (from the

first imidacloprid exposure to before pupa) were recorded daily to calculate food

consumption, survival rates, developmental rates, and development times. The growth

index (GI) was calculated for larvae exposed to imidacloprid from the three-day-old to

the six-day-old using the equations described by Zhang (Zhang et al., 1993). The

developmental rate was measured as the number of larvae successfully reaching the

pupal stage as a percentage of the total initial sample size.

RNA extraction, cDNA synthesis, and real-time quantitative PCR analysis

Total RNA was extracted from larvae using TRIzol reagent (Thermo Fisher Scientific)

and RNAex Pro Reagent (Accurate Biology, China) as instructed by the manufacturer.

The RNA quality was evaluated by gel electrophoresis and absorbance determination

using a NanoDrop 2000 spectrophotometer (Thermo Scientific, New York, USA).

Genomic DNA was removed, and cDNA was synthesized using a reverse transcription kit

(Evo M-MLV RT Mix Kit with gDNA Clean for qPCR, Accurate Biology, China) according to

the manufacturer's instructions. Briefly, 1 µg of RNA reacted with the 5X gDNA Clean

Reaction Mix provided in the kit at 42°C for 5 min to remove residual genomic DNA to

ensure the accuracy of the quantitative results. After the reaction, 4 µL of 5X Evo M-

MLV RT Reaction Mix and 4 µL of RNase-free water were added to the reaction tube

for 20 µL. The mixture was incubated at 37°C for 15 min and 85°C for 5 sec to generate

cDNA. Then, quantitative RT–PCR was performed with a Bio-Rad CFX96™ Real-time PCR

Detection System (Bio-Rad, Hercules, CA, USA) according to standard protocols and

cycling conditions. RP49 (ribosomal protein 49, Accession no. AF441189) (Lourenco et al.,

2008) was used as the endogenous control. The primers for target genes and RP49 are

listed in Table S1. qPCR system (20 µL): EvaGreen Express 2X qPCR MasterMix (10 µL),

each forward and reverse primer (0.4 µL, 0.2 µmol/L), cDNA (1 µL, 100 ng/µL), ddH$_2$O

(8.2 μL). The thermal cycling program was as follows: 95°C for 30 sec, followed by 40

cycles of 95°C for 5 s and 60°C for 30 s. The qPCR products were sequenced by Shanghai

Biological Engineering Technology Services Co., Ltd. to confirm whether the amplified

fragments were the target sequences. Relative mRNA expression levels of target genes

were calculated using the $2^{-\Delta\Delta Ct}$ method.

Histochemical measurement

Juvenile hormone 3 (JH-III), 20-hydroxyecdysone (20E) titers, and reactive oxygen

species (ROS) levels in larvae were determined using ELISA enzyme-linked

immunosorbent assay kits (Jining, Shanghai, China) according to the manufacturer's

instructions. Briefly, the larva was homogenized on ice in 5 volumes (M: V) of

phosphate-buffered saline (PBS) buffer, and the supernatant was collected after

centrifugation at 12,000 x g for 20 min. Subsequently, 50 μL of reference standard

solution or 10 μL of sample supernatant plus 40 μL of dilution were added to the

microtiter wells and incubated at 37°C for 60 min with 100 μL of HRP-labeled detection

antibody. Each well was washed five times with 400 μL of PBS buffer containing 0.05%

Tween 20, pat dried, and left to stand at room temperature for 5 min. TMB chromogen

solution A (50 μL) and solution B (50 μL) were added to each well, gently mixed, and

incubated at 37°C for 15 min in the dark. Then, 50 μL of H_2SO_4 (4 mol/L) was added to

each well to terminate the reaction. Finally, the OD value was measured at 450 nm with

a NanoDrop 2000 spectrophotometer (Thermo Scientific, New York, USA), and the

content of the detected substance was calculated from a calibration curve constructed

by reference standards.

To prepare samples for analysis of protein carbonylation (PCO), acetylcholinesterase

(AchE), catalase (CAT), malondialdehyde (MDA), total antioxidant capacity (T-AOC),

superoxide dismutase (SOD), and glutathione (GSH), larvae was weighed and

homogenized on ice in 5 volumes (M: V) of PBS buffer, then centrifuged at 12,000 x g for

15 min to collect the supernatant. Subsequently, PCO content was measured using the

2,4-dinitrophenyl-hydrazine (DNPH) method with slight modifications. The above-prepared supernatant was mixed with streptomycin sulfate at a 9:1 volume (V: M), kept at room temperature for 10 min, and then centrifuged at 12,000 x g for 15 min. Then, 80 µL of supernatant was mixed with 160 µL of 10 mM DNPH in 2 M HCl and reacted at 37°C for 30 min. Sample controls were also prepared by adding an equal volume (500 µL) of 2 M HCl. Then, 200 µL of 20% TCA was added, mixed thoroughly, and centrifuged at 4°C for 10 min at 12,000 x g, and the precipitate was collected. Add 400 µL of ethanol: ethyl acetate mixture (1:1), vortex and mix, centrifuge at 4°C 12,000 x g for 10 min, and collect the precipitate to remove unbound DNPH. This DNPH removal step was repeated once. The pellet was mixed with 400 µL of 6 M guanidine hydrochloride, kept at room temperature for 15 min, and then centrifuged at 4°C 12,000 x g for 10 min to collect the supernatant. The carbonyl and protein contents were determined by reading the absorbance at $\lambda = 370$ nm and 280 nm with a SpectraMax 190 spectrophotometer (Molecular Devices, California, USA) for each sample against an appropriate blank. Finally, the carbonyl content per unit of protein was calculated based on the protein content.

AChE activity was measured by the Ellman method as described by Orhan et al. (Orhan et al., 2007) with slight modifications. Briefly, 20 µL of the prepared sample supernatant was mixed with 160 µL of 0.1 mM PBS buffer (pH 8.0) and 10 µL of an AChE (0.28 U mL^{-1}) and incubated for 15 min at room temperature. Then, 10 µL of ATChI (acetylthiocholine iodide, 4 mg mL^{-1}) was added together with 20 µL of DTNB (5,5-dithiobis-2-nitrobenzoic acid, 1.2 mg mL^{-1}) and immediately mixed, and the absorbance was read at 412 nm. The results were expressed as activity percentages relative to the sample solvent negative control.

The CAT activity was determined using a kit (Beyotime, Shanghai, China) according to the manufacturer's instructions with slight modifications. Briefly, 10 µL of the prepared sample supernatant was mixed with 30 µL of Tris-HCl buffer (pH 7.0, 50 mmol L^{-1}), 10 µL of 250 mM H_2O_2 was added, and the mixture was immediately mixed and kept at 25°C

for 5 min. Then, 450 µL of 1.8 mol L-1 H_2SO_4 was added, and the mixture was vortexed

to terminate the reaction. Subsequently, 10 µL of this termination reaction solution was

mixed with 40 µL of Tris-HCl buffer (pH 7.0, 50 mmol L^{-1}). Then, 10 µL of this mixture

was added to a 96-well plate mixed with 200 µL of color development working solution

containing peroxidase and incubated for 15 min at 25°C. Finally, the absorbance was

measured at 520 nm, and the CAT activity was calculated using a calibration curve of 5

mM H_2O_2.

T-AOC was measured using the total antioxidant capacity assay kit with the ABTS

(2,2'-azino-bis(3-ethylbenzthiazoline-6-sulfonic acid) method based on a previous report

(Chen et al., 2019). Add 10 µL of the prepared sample supernatant to 200 µL of ABTS

working solution. After 6 min of reaction, the absorbance was measured at 734 nm.

Finally, the antioxidant activity was displayed by Trolox equivalents antioxidant capacity

as mmol Trolox equivalents/g extract. The ABTS working solution was prepared as

follows: 400 µL of potassium persulfate was added to 400 µL of ABTS to generate an

ABTS stock solution, which was then incubated for 16 h at room temperature in the dark.

Before use, the stock solution was diluted with phosphate buffer, and the absorbance at

734 nm was adjusted to 0.70 ± 0.05.

Additionally, MDA, SOD, and GSH levels were detected using ELISA enzyme-linked

immunosorbent assay kits (Beyotime, Shanghai, China) according to our recent study (Li

et al., 2022).

Histopathological analysis

Microsectioning and hematoxylin-eosin (HE) staining was performed according to

standard histological protocols (Khan et al., 2010) with some modifications. Briefly,

entire fresh larvae were fixed in 10% neutral-buffered formalin solution for 12 h at 4°C

after 72 h of imidacloprid exposure. After isopropyl alcohol dehydration, samples were

embedded in paraffin and sectioned. Three-micron-thick sections were stained with a

Hematoxylin and Eosin Staining Kit (Beyotime, Shanghai, China), observed with an

Olympus BX51 microscope, and photographed with an Olympus digital camera.

Statistical analysis

Data were obtained from three independent experiments performed in triplicate and are presented as the mean ± SEM. SPSS statistical package was used for statistical analysis using Student's t-test and one-way ANOVA followed by Tukey's multiple comparison test (*$P < 0.05$, **$P < 0.01$, and ***$P < 0.001$ were considered statistically significant). GraphPad software was used for data normalization, Pearson correlation coefficient calculation was analyzed using SPSS, and a heat map was generated.

Acknowledgments

The authors would like to thank the other colleagues in our laboratory who contributed to this study. This study received financial support from the grants from the earmarked fund for China Agriculture Research System (No. CARS-44); the Science and Technology Project of the Chongqing Municipal Education Commission of China (No. KJZD-K202100502); the Natural Science Foundation Project of Chongqing of China (No. cstc2021jcyj-msxmX0422); and the Natural Science Foundation Project of the State Key Laboratory of Silkworm Genome Biology of China (No. SKLSGB-ORP202103).

Additional information

Author contributions

Zhi Li: Supervision, Designed research, Conceptualization, Methodology, Funding acquisition, Writing-original draft, Writing-review & editing. **Yuedi wang:** Investigation, Methodology, Validation, Formal analysis, Project administration. **Lanchun Chen:** Investigation, Data curation. **Qiqian Qing:** Investigation, Data curation. **Xiaoqun Dang:** Data curation. **Zhengang Ma:** Data curation. **Zeyang Zhou:** Conceived the research, Research guidance. All the authors read, corrected, and approved the manuscript.

Ethics

Animal experimentation: This study was performed in strict accordance with the guidelines of the Animal Bioethics Committee at Chongqing Normal University, China approved all experimental protocols.

Data Availability

All the data needed to evaluate the conclusions in this paper are present in the paper and/or Supplementary Materials.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Additional files

Supplementary files：Table S1. Primers used in real-time quantitative PCR.

References

(2019) **A Review on Microbial Alkaline Protease: An Essential Tool for Various Industrial Approaches** *Industrial Biotechnology* **15:**69-78

Anatra-Cordone M D. P (2005) **Imidacloprid. Human health assessment and ecological risk assessment Final report**. **Syracuse Environmental Research Associates Inc, New York** *SERA TR*:05-43.

Anderson N. L., Harmon-Threatt, A. N (2021) **Chronic contact with imidacloprid during development may decrease female solitary bee foraging ability and increase male competitive ability for mates** *Chemosphere* **283**

Aupinel P., et al. (2005) **Improvement of artificial feeding in a standard in vitro method for rearing Apis mellifera larvae** *Bulletin of Insectology* **58:**107-111

Belzunces L. P., et al. (2012) **Neural effects of insecticides in the honey bee**. **Springer-Verlag**

Berenbaum M. R., Johnson R. M. (2015) **Xenobiotic detoxification pathways in honey bees** *Curr Opin Insect Sci* **10:**51-58

Bergh S., Arking R. (1984) **Development profile of the heat shock response in early embryos of Drosophila** *J Exp Zool* **231:**379-91

Beyers D. W., et al. (1999) **Estimating physiological cost of chemical exposure: integrating energetics and stress to quantify toxic effects in fish** *Canadian Journal of Fisheries and Aquatic Sciences* **56:**814-822

Bhme F., et al. (2018) **From field to food—will pesticide-contaminated pollen diet lead to a contamination of royal jelly? Springer Paris**

Bouayad N., et al. (2012a) **Dietary effects of harmine, a beta-carboline alkaloid, on development, energy reserves and alpha-amylase activity of Plodia interpunctella Hubner (Lepidoptera: Pyralidae)** *Saudi J Biol Sci* **19:**73-80

Bouayad N., et al. (2012b) **Dietary effects of harmine, a β-carboline alkaloid, on development, energy reserves and α-amylase activity of Plodia interpunctella Hübner (Lepidoptera: Pyralidae)** *Saudi J Biol Sci* **19**:73-80

Cardoso P., et al. (2020) **Scientists' warning to humanity on insect extinctions** *Biological Conservation* **242:**108426

Carneiro L. S., et al. (2022) **Acute oral exposure to imidacloprid induces apoptosis and autophagy in the midgut of honey bee Apis mellifera workers** *Sci Total Environ* **815:**152847

Carranza-García E., Navarro R. E. (2020) **Insights Into the Hypometabolic Stage Caused by Prolonged Starvation in L4-Adult Caenorhabditis elegans Hermaphrodites** *Front Cell Dev Biol* **8:**124

Chaimanee V., et al. (2016) **Sperm viability and gene expression in honey bee queens (Apis mellifera) following exposure to the neonicotinoid insecticide imidacloprid and the organophosphate acaricide coumaphos** *J Insect Physiol* **89:**1-8

Chakrabarti P., et al. (2015) **Pesticide-induced oxidative stress in laboratory and field populations of native honey bees along intensive agricultural landscapes in two Eastern Indian states** Apidologie **46**:107-129

Charles V., et al. (2000) **Effects of imidacloprid on Harmonia axyridis (Coleoptera: Coccinellidae) larval biology and locomotory behavior** *Eur J Entomol* **97**:501-506

Chauzat M. P., et al. (2009) **Influence of pesticide residues on honey bee (Hymenoptera: Apidae) colony health in France** *Environ Entomol* **38:**514-23

Chauzat M. P., et al. (2011) **An assessment of honeybee colony matrices, Apis mellifera (Hymenoptera: Apidae) to monitor pesticide presence in continental France** *Environ Toxicol Chem* **30:**103-11

Chen J., et al. (2019) **Cytoprotective Effect of Ligustrum robustum Polyphenol Extract against Hydrogen Peroxide-Induced Oxidative Stress via Nrf2 Signaling Pathway in Caco-2 Cells** *Evid Based Complement Alternat Med* **2019**:5026458.

Chen Y. R., et al. (2021) **Chronic Effects of Imidacloprid on Honey Bee Worker Development— Molecular Pathway Perspectives** *Int J Mol Sci* **22**:11835

Corona M., et al. (2007) **Vitellogenin, juvenile hormone, insulin signaling, and queen honey bee longevity** *Proc Natl Acad Sci* **104**, 7128-33

Dai P., et al. (2018) **The Herbicide Glyphosate Negatively Affects Midgut Bacterial Communities and Survival of Honey Bee during Larvae Reared in Vitro** *J Agric Food Chem* **66**:7786-7793

Dai P., et al. (2019) **Chronic toxicity of clothianidin, imidacloprid, chlorpyrifos, and dimethoate to Apis mellifera L. larvae reared in vitro** *Pest Manag Sci* **75**:29-36

Deng H., et al. (2012) **Homeodomain POU and Abd-A proteins regulate the transcription of pupal genes during metamorphosis of the silkworm, Bombyx mori** *Proc Natl Acad Sci* **109**:12598-12603.

Desneux N., et al. (2007) **The sublethal effects of pesticides on beneficial arthropods** *Annu Rev Entomol* **52**:81-106

Dively G. P., Kamel A. (2012) **Insecticide residues in pollen and nectar of a cucurbit crop and**
their potential exposure to pollinators *J Agric Food Chem* **60**:4449-56
Dong M., et al. (2013) **Toxic effects of 1-decyl-3-methylimidazolium bromide ionic liquid on the**
antioxidant enzyme system and DNA in zebrafish (Danio rerio) livers *Chemosphere*
91:1107-1112
Dutra B. K., et al. (2009) **Carbofuran-induced alterations in the energy metabolism and**
reproductive behaviors of Hyalella castroi (Crustacea, Amphipoda) *Comp Biochem*
Physiol C Toxicol Pharmacol **149**:640-646
Felipe M., et al. (2000) **Low doses of the neonicotinoid insecticide imidacloprid induce ROS**
triggering neurological and metabolic impairments in Drosophila *Proc Natl Acad Sci*
117:25840-25850
Gill R. J., et al. (2012) **Combined pesticide exposure severely affects individual- and colony-level**
traits in bees *Nature* **491**:105-108
Gregorc A., et al. (2018) **Effects of coumaphos and imidacloprid on honey bee (Hymenoptera:**
Apidae) lifespan and antioxidant gene regulations in laboratory experiments *Sci Rep*
8:15003
Guedes R. N. C., et al. (2006) **Cost and mitigation of insecticide resistance in the maize weevil,**
Sitophilus zeamais *Physiological Entomology* **31**:30-38
He G., et al. (2012) **RNA interference of two acetylcholinesterase genes in Plutella xylostella**
reveals their different functions *Arch Insect Biochem Physiol* **79**:75-86
Hurst V., et al. (2014) **Toxins induce 'malaise' behaviour in the honeybee (Apis mellifera)** *J Comp*
Physiol A Neuroethol Sens Neural Behav Physiol **200:**881-890
Islam M. A., et al. (2019) **Acute Toxicity of Imidacloprid on the Developmental Stages of**
Common Carp Cyprinus carpio *Toxicol Environ Health Sci* **11**:244-251
Jeschke P., et al. (2011) **Overview of the status and global strategy for neonicotinoids** *J Agric*
Food Chem **59**:2897-908
Jia Q., Li, S. (2023) **Mmp-induced fat body cell dissociation promotes pupal development and**
moderately averts pupal diapause by activating lipid metabolism *Proc Natl Acad Sci*
120:e2215214120.
Jiang D., et al. (2020) **Cd exposure-induced growth retardation involves in energy metabolism**
disorder of midgut tissues in the gypsy moth larvae *Environ Pollut* **266**:115173
Jorgensen P., et al. (2009) **The mechanism and pattern of yolk consumption provide insight into**
embryonic nutrition in Xenopus *Development* **136**:1539-1548
Kapoor U., et al. (2014) **Disposition and acute toxicity of imidacloprid in female rats after single**
exposure *Food Chem Toxicol* **68**:190-195
Karahan A., et al. (2015) **Sublethal imidacloprid effects on honey bee flower choices when**
foraging *Ecotoxicology*.**24**:2017-2025
Katić A., et al. (2021) **Effects of low-level imidacloprid oral exposure on cholinesterase activity,**
oxidative stress responses, and primary DNA damage in the blood and brain of male
Wistar rats *Chemico-Biological Interactions* **338**:109287
Khan D. A., et al. (2010) **Monitoring health implications of pesticide exposure in factory**
workers in Pakistan *Environ Monit Assess* **168**:231-240
Kooijman B., (2009) **Dynamic Energy Budget Theory for Metabolic Organisation** *Cambridge*
University Press **Cambridge**

Li-Byarlay H., et al. (2016) **Honey bee (Apis mellifera) drones survive oxidative stress due to**
increased tolerance instead of avoidance or repair of oxidative damage *Exp Gerontol*
83:15-21
Li X., et al. (2007) **Molecular mechanisms of metabolic resistance to synthetic and natural**
xenobiotics *Annu Rev Entomol* **52**:231-253
Li X., et al. (2021) **New insights into crosstalk between apoptosis and necroptosis co-induced by**
chlorothalonil and imidacloprid in Ctenopharyngodon idellus kidney cells *Sci Total*
Environ **780**:146591
Li Z., et al. (2022) **Melatonin enhances the antioxidant capacity to rescue the honey bee Apis**
mellifera from the ecotoxicological effects caused by environmental imidacloprid
Ecotoxicol Environ Saf **239**:113622
Liu D., et al. (2018) **Toxicity and sublethal effects of fluralaner on Spodoptera litura Fabricius**
(Lepidoptera: Noctuidae) *Pestic Biochem Physiol* **152**:8-16
Lourenco A. P., et al. (2008) **Validation of reference genes for gene expression studies in the**
honey bee, Apis mellifera, by quantitative real-time RT-PCR *Apidologie* **39**:372-385
Luo W., et al. (2021) **Juvenile hormone signaling promotes ovulation and maintains egg shape**
by inducing expression of extracellular matrix genes *Proc Natl Acad Sci*
118:e2104461118
Mahé C., et al. (2021) **The countryside or the city: Which environment is better for the**
honeybee? *Environ Res* **195**:110784
Mao W., et al. (2009) **Quercetin-metabolizing CYP6AS enzymes of the pollinator Apis mellifera**
(Hymenoptera: Apidae) *Comp Biochem Physiol B Biochem Mol Biol* **154**:427-434
Mao W., et al. (2011) **CYP9Q-mediated detoxification of acaricides in the honey bee (Apis**
mellifera) *Proc Natl Acad Sci* **108**:12657-12662
Martín-Blázquez, R., et al. (2023) **Gene expression in bumble bee larvae differs qualitatively**
between high and low concentration imidacloprid exposure levels *Sci Rep* **13**:9415.
Makoto I., et al. (2020) **Cofactor-enabled functional expression of fruit fly, honeybee, and**
bumblebee nicotinic receptors reveals picomolar neonicotinoid actions *Proc Natl Acad*
Sci **117**:16283-16291
Matozzo V., et al. (2008) **Vitellogenin as a biomarker of exposure to estrogenic compounds in**
aquatic invertebrates: a review *Environ Int* **34**:531-545
Matsukura K., et al. (2008) **Changes in chemical components in the freshwater apple snail,**
Pomacea canaliculata (Gastropoda: Ampullariidae), in relation to the development of
its cold hardiness *Cryobiology* **56**:131-137
Mehlhorn H., et al. (1999) **Effects of imidacloprid on adult and larval stages of the flea**
Ctenocephalides felis after in vivo and in vitro application: a light- and electron-
microscopy study *Parasitol Res* **85**:625-637
Medrzycki P., et al. (2003) **Effects of imidacloprid administered in sub-lethal doses on honey**
bee behaviourr. Laboratory tests *Bulletin of Insectology* **56**:59-62
Mullin C. A., et al. (2010) **High levels of miticides and agrochemicals in North American apiaries:**
implications for honey bee health *PLoS One* **5**:e9754
Orhan I., et al. (2007) **Antioxidant and anticholinesterase evaluation of selected Turkish Salvia**
species *Food Chemistry* **103**:1247-1254
Paris L., et al. (2017) **Disruption of oxidative balance in the gut of the western honeybee Apis**
mellifera exposed to the intracellular parasite Nosema ceranae and to the insecticide
fipronil *Microb Biotechnol* **10**:1702-1717

Peng Y. C., Yang, E. C. (2016) **Sublethal Dosage of Imidacloprid Reduces the Microglomerular**
Density of Honey Bee Mushroom Bodies *Sci Rep* **6**:19298
Pettis J. S., et al. (2013) **Crop pollination exposes honey bees to pesticides which alters their**
susceptibility to the gut pathogen Nosema ceranae *PLoS One* **8**:e70182
Radwan M. A., et al. (2008) **Biochemical and histochemical studies on the digestive gland of**
Eobania vermiculata snails treated with carbamate pesticides *Pesticide Biochemistry*
and Physiology **90**:154-167
Rambabu J. P., Rao M. B. (1994) **Effect of an organochlorine and three organophosphate**
pesticides on glucose, glycogen, lipid, and protein contents in tissues of the freshwater
snail Bellamya dissimilis (Muller) *Bull Environ Contam Toxicol* **53**:142-148
Rashid S., et al. (2021) **Developmental plasticity and the response to nutrient stress in**
Caenorhabditis elegans *Dev Biol* **475**:265-276
Ribeiro S., et al. (2001) **Effect of endosulfan and parathion on energy reserves and physiological**
parameters of the terrestrial isopod Porcellio dilatatus *Ecotoxicol Environ Saf* **49**:131-
138
Rocha C. M., et al. (2022) **Environmental photochemical fate of pesticides ametryn and**
imidacloprid in surface water (Paranapanema River, Sao Paulo, Brazil) *Environ Sci Pollut*
Res Int **29**:42290-42304
Samojeden C. G., et al. (2022) **Toxicity and genotoxicity of imidacloprid in the tadpoles of**
Leptodactylus luctator and Physalaemus cuvieri (Anura: Leptodactylidae) *Sci Rep*
12:11926.
Sánchez-Bayo F., Wyckhuys, K. A. G. (2019) **Worldwide decline of the entomofauna: A review of**
its drivers *Biological Conservation* **232**:8-27
Seehuus S.-C., et al. (2006) **Reproductive protein protects functionally sterile honey bee**
workers from oxidative stress. *Pro Nat Aca Sci* **103**:962-967
Shafiq-ur-Rehman, et al. (2012) **Chlorpyrifos-induced neuro-oxidative damage in bee** *Toxico &*
Enviro Health Sci. **4**:30-36
Shan Y., et al. (2020) **Effect of imidacloprid on the behavior, antioxidant system,**
multixenobiotic resistance, and histopathology of Asian freshwater clams (Corbicula
fluminea) *Aquat Toxicol* **218**:105333
Shaw G. M., et al. (2014) **Early pregnancy agricultural pesticide exposures and risk of**
gastroschisis among offspring in the San Joaquin Valley of California *Birth Defects Res A*
Clin Mol Teratol **100**:686-694
Simon-Delso N., et al. (2015) **Systemic insecticides (neonicotinoids and fipronil): trends, uses,**
mode of action and metabolites *Environ Sci Pollut Res Int* **22**:5-34
Tahira F. (2013) **A potential link among biogenic amines-based pesticides, learning and memory,**
and colony collapse disorder: A unique hypothesis *Neurochem Int* **62**:122-36.
Tian F., et al. (2020a) **Effects of Thymoquinone on Small-Molecule Metabolites in a Rat Model**
of Cerebral Ischemia Reperfusion Injury Assessed using MALDI-MSI *Metabolites* **10**:27
Tian X., et al. (2020b) **Neonicotinoids caused oxidative stress and DNA damage in juvenile**
Chinese rare minnows (Gobiocypris rarus) *Ecotoxicol Environ Saf* **197**:110566
Tomé H. V. V., et al. (2020) **Frequently encountered pesticides can cause multiple disorders in**
developing worker honey bees *Environ Pollut* **256**:113420
Tong Z., et al. (2018) **A survey of multiple pesticide residues in pollen and beebread collected in**
China *Sci Total Environ* **640-641**:1578-1586

Tudi M., et al. (2021) **Agriculture Development, Pesticide Application and Its Impact on the**
Environment *Int J Environ Res Public Health* **18**:1112
Woyciechowski M.; Moroń, D (2009) L**ife expectancy and onset of foraging in the honeybee**
(Apis mellifera) *Insectes Soc* **56**:193–201.
Whitehorn P.R., et al. (2018) **Larval exposure to the neonicotinoid imidacloprid impacts adult**
size in the farmland butterfly Pieris brassicae *PeerJ* **6**:e4772
Wu I. W., et al. (2001) **Acute poisoning with the neonicotinoid insecticide imidacloprid in N-**
methyl pyrrolidone *J Toxicol Clin Toxicol* **39**:617-621
Wu J.Y. et al. (2011) **Sub-lethal effects of pesticide residues in brood comb on worker honey**
bee (Apis mellifera) development and longevity *PLoS One* **6**:e14720
Wu M. C., et al. (2017) **Gene expression changes in honey bees induced by sublethal**
imidacloprid exposure during the larval stage *Insect Biochem Mol Biol* **88**:12-20
Yang E. C., et al. (2012) **Impaired olfactory associative behavior of honeybee workers due to**
contamination of imidacloprid in the larval stage *PLoS One* **7**:e49472
Yang W., et al. (2014) **Residential agricultural pesticide exposures and risk of neural tube**
defects and orofacial clefts among offspring in the San Joaquin Valley of California *Am*
J Epidemiol **179**:740-748
Yuan D., et al. (2020) **The AMPK-PP2A axis in insect fat body is activated by 20-**
hydroxyecdysone to antagonize insulin/IGF signaling and restrict growth rate *Pro Nat*
Aca Sci **117**:9292-9301
Zhang M., et al. (1993) **Quantification of insect growth and its use in screening of naturally**
occurring insect control agents *J Chem Ecol* **19**:1109-1118
Zheng H. Q., Fu-Liang H. U. (2009) **Honeybee:a newly emerged model organism** *Acta*
Entomologica Sinica 210-215
Zhu W., et al. (2014) **Four common pesticides, their mixtures and a formulation solvent in the**
hive environment have high oral toxicity to honey bee larvae *PLoS One* **9**:e77547
843

 Figure legends

Figure 1. Imidacloprid causes larval developmental retardation in *Apis mellifera*. Larvae

were orally administered imidacloprid (IMI) or artificial food (CK) beginning at three-

day-old, followed by developmental monitoring for 96 hours. (A) Larval developmental

phenotypes at 0.7, 1.2, 3.1, and 377 ppb environmental concentrations of imidacloprid

exposure; (B) Larval developmental phenotype at 377 ppb imidacloprid exposure; (C)

Effect of 0.7, 1.2, 3.1, and 377 ppb imidacloprid on larval survival; (D) and (E) body width

and weight of larvae exposed to 0.7, 1.2, 3.1, and 377 ppb imidacloprid for 96 hours; (F)

and (G) statistics for daily (F) and three-day cumulative feeding (G) for larvae exposed to

377 ppb imidacloprid; (H) larval development time from three-day-old to pupal stage; (I)

The number of larvae that successfully reached the pupal stage as a percentage of the

total initial sample size; (J) The growth index of larvae. Statistical significance was set at

$*P < 0.05$ and $**P < 0.01$.

Figure 2. Imidacloprid neurotoxicity in *Apis mellifera* larvae. (A) AChE activity and (B)

nerve conduction-related gene expression analysis in larvae exposed to 377 ppb

imidacloprid for 72 hours or a control group. IMI is the larvae exposed to imidacloprid

for 72 hours. CK is the artificial food control group. Statistical significance was set at $*P <$

0.05 and $**P < 0.01$.

Figure 3. Effects of imidacloprid on the homeostasis of developmental regulation in *Apis*

mellifera larvae. (A) Relative gene expression and (B) hormone levels of developmental

regulatory-related genes in larvae exposed to 377 ppb imidacloprid for 72 h or a control

group. Statistical significance was set at $*P < 0.05$ and $**P < 0.01$.

Figure 4. Imidacloprid induced a detoxification response in *Apis mellifera* larvae. Relative

expression levels of cytochrome P450 monooxygenase family genes in larvae exposed to

imidacloprid for 72 hours. IMI is the larvae exposed to imidacloprid for 72 hours. CK is

the artificial food control group. Statistical significance was set at *$P < 0.05$, **$P < 0.01$,

and ***$P < 0.001$.

Figure 5. Imidacloprid exposure causes oxidative stress, gut apoptosis, and tissue

structural damage while inducing antioxidant defenses in *Apis mellifera* larvae. (a)

Analysis of oxidative stress and damage in larvae exposed to imidacloprid for 72 hours.

ROS levels indicate the degree of oxidative stress. MDA is a marker for lipid peroxidation.

PCO levels reflect the extent of protein carbonylation damage; (b) Activity of

antioxidants in larvae exposed to imidacloprid for 72 hours; (c) The relative expression

levels of antioxidant genes in larvae exposed to imidacloprid for 72 hours; (d)

Histological sections of the larval gut stained with H&E. The circled inserts in the figure

show magnified views of the muscle layer. The blue arrow indicates the cell nucleus of

the muscle layer. The boxed inset in the figure is a magnified view of the basal layer

(black arrow), peritrophic membrane (red arrow), and food residues (green arrow). IMI is

the larvae exposed to imidacloprid for 72 hours. CK is the artificial food control group.

Statistical significance was set at *$P < 0.05$, **$P < 0.01$, and ***$P < 0.001$.

Figure 6. Imidacloprid toxicity inhibited the expression of genes involved in nutrient

catabolism, protein synthesis, and energy metabolism and reserves. (A) Carbohydrate

catabolism; (B) Proteolysis; (C) Amino acid transport; (D) Mitochondrial oxidative

phosphorylation; (E) Glycolysis; (F) Protein synthesis. (G) The total contents of ATP,

glycogen (H), and protein (I). IMI is the larvae exposed to imidacloprid for 72 hours. CK is

the artificial food control group. Statistical significance was set at *$P < 0.05$, **$P < 0.01$,

and ***$P < 0.001$.

Figure 7. Correlation between phenotypes and molecular characteristics in imidacloprid-

exposed larvae with developmental retardation. Data were normalized using GraphPad

software, followed by a Pearson correlation coefficient calculation using SPSS, and a

heatmap was generated.

Figure 8. A model interpreting the ecotoxicological effects of imidacloprid on *Apis*

mellifera larvae and the mechanisms of imidacloprid-induced developmental retardation.

Imidacloprid toxicity caused growth and developmental delay in *A. mellifera* larvae. It

disrupts the regulatory balance of molt development, resulting in restricted

development. In addition, the damage caused by imidacloprid restricts the metabolism

and utilization of food nutrients and energy in larvae. The additional energy consumed

by larval P450 detoxification and antioxidant defenses further reduces the supply of

nutrients and energy supply for growth and development. The arrows indicate induction

or promotion, the straight lines indicate inhibition, and the green dotted lines indicate

speculative inhibition.

Imidacloprid toxicity and the underlying mechanism causing larval development delay

Imidacloprid

3rd instar

4th instar

6th instar

7th instar

72 hours Exposure

Developmental retardation

Toxic effects

Epigenetic traits

Energy reserves

Nutrient catabolism

Energy metabolism

Protein synthesis

Developmental hormone regulation

Nerve conduction

Detoxification

Oxidative stress and damage

Antioxidant

Gut tissue damage

Muscle layer

Basal layer, Peritrophic membrane, Food

Correlation Analysis

Epigenetic growth traits

Energy reserves

Nutrient catabolism and transport

Energy metabolism

Protein synthesis

Developmental regulation

Nerve conduction

Detoxification

Oxidative stress and damage

Antioxidant