October 31, 2023

The *Toxoplasma* monocarboxylate transporters are involved in the metabolism within the apicoplast and are linked to parasite survival

Hui Dong[a,b,1], Jiong Yang[c,1], Kai He[a,b,1], Wen-Bin Zheng[d], De-Hua Lai[c], Jing Liu[a,b], Hui-Yong Ding[a,b], Rui-Bin Wu[a,b], Kevin M. Brown[e], Geoff Hide[f], Zhao-Rong Lun[c], Xing-Quan Zhu[d], Shaojun Long[a,b,#]

[a]National Key Laboratory of Veterinary Public Health Safety, and College of Veterinary Medicine, China Agricultural University, Beijing 100193, China

[b]National Animal Protozoa Laboratory and School of Veterinary Medicine, China Agricultural University, Beijing 100193, China

[c]MOE Key Laboratory of Gene Function and Regulation, State Key Laboratory of Biocontrol, School of Life Sciences, Sun Yat-Sen University, Guangzhou 510275, China

[d]College of Veterinary Medicine, Shanxi Agricultural University, Taigu 030801, Shanxi Province, China

[e]Department of Microbiology and Immunology, University of Oklahoma Health Sciences Center, Oklahoma City, Oklahoma, USA

[f]Biomedical Research and Innovation Centre and Environmental Research and Innovation Centre, School of Science, Engineering and Environment, University of Salford, Salford, M5 4WT, UK

[1]These authors contributed equally to the work

[#]To whom correspondence may be addressed:

Shaojun Long, LongS2018@163.com

Abstract

The apicoplast is a four-membrane plastid found in the apicomplexans, which harbors

biosynthesis and organelle housekeeping activities in the matrix. However, the

mechanism driving the flux of metabolites, in and out, remains unknown. Here we

used TurboID and genome engineering to identify apicoplast transporters in

Toxoplasma gondii. Among the many novel transporters, we show that one pair of

apicomplexan monocarboxylate transporters (AMTs) appears to have evolved from a

putative host cell that engulfed a red alga. Protein depletion showed that AMT1 and

AMT2 are critical for parasite growth. Metabolite analyses supported the notion that

AMT1 and AMT2 are associated with biosynthesis of isoprenoids and fatty acids.

However, stronger phenotypic defects were observed for AMT2, including in the

inability to establish *T. gondii* parasite virulence in mice. This study clarifies,

significantly, the mystery of apicoplast transporter composition and reveals the

importance of the pair of AMTs in maintaining the apicoplast activity in apicomplexans.

Introduction

The phylum Apicomplexa encompasses thousands of intracellular parasites, including

Plasmodium. spp., the notorious agent of malaria, and *Toxoplasma gondii*, the highly

successful parasite that infects almost all warm-blooded animals and humans

(Plattner and Soldati-Favre, 2008). It is believed that apicomplexan evolution included

two key independent, and sequential, endosymbiotic events, finally leading to the

common ancestor of the chromalveolates (e.g. alveolates, heterokonts, oomycetes

and Cryptomonads) (Cavalier-Smith, 1999; Gould et al., 2008; Janouskovec et al.,

2010). This ancestor eventually gave rise to the apicomplexans, most of which still

contain a secondary non-photosynthetic plastid, the apicoplast (McFadden and Yeh,

2017; Obornik et al., 2009; Salomaki and Kolisko, 2019). Though endosymbiont

origins of the apicoplast were proposed (Köhler et al., 1997; McFadden et al., 1996;

Wilson et al., 1996), it is believed that the apicoplast originates from a red alga (Foth

and McFadden, 2003; McFadden and van Dooren, 2004). Therefore, the apicoplast

membranes have been proposed to evolve from the red algal plastid, the algal plasma

membrane and the host endomembrane when counting from the innermost to

outermost membranes of the apicoplast (Gould et al., 2008; McFadden and Yeh,

2017). This heavily compartmentalized feature poses great challenges for metabolic

integration of the organelle within these parasites and, in particular, the transport of

metabolites across these membrane systems.

The apicoplast houses essential biosynthetic pathways. These pathways include the

fatty acid synthesis type II (FASII) pathway, methylerythritol phosphate (MEP)

pathway, Fe-S cluster biosynthesis pathway, (partial) heme biosynthesis pathway and

lysophosphatidic acid synthesis pathway (using the FASII products) (Amiar et al.,

2016; Fata and Rabuzzi, 1988; Jomaa et al., 1999; Kloehn et al., 2021; Krishnan et al.,

2020b; Ralph et al., 2004; Tjhin et al., 2020; Waller et al., 1998). They are able to

make critical components of fatty acids and isoprenoid precursors for the parasites, in

T. gondii and *Plasmodium falciparum* (McFadden and Yeh, 2017; Ramakrishnan et al.,

2012; Striepen, 2011; Vaughan et al., 2009; Yeh and DeRisi, 2011). The apicoplast

FASII pathway in *T. gondii* is key to understand the metabolic flexibility, and can

become essential under metabolic adaptation conditions (Amiar et al., 2020; Botté et

al., 2013; Krishnan et al., 2020a; Primo et al., 2021). Moreover, the apicoplast

maintains housekeeping activities for genome replication, transcription and translation

(Low et al., 2018; McFadden and Roos, 1999). Accordingly, it requires adequate

supplies of carbon molecules, energy, reducing equivalents, cofactors, amino acids,

nucleotides as well as inorganic ions, to feed the biosynthetic pathways and plastid

maintenance activities in the matrix (Kloehn et al., 2021; McFadden and Yeh, 2017;

Ralph et al., 2004). However, compared to plant plastids, the nature of the channels

and transporters in the apicoplast remains obscure.

The apicoplast biosynthetic pathways were characterized through a

homology/bipartite signal-based approach (Foth et al., 2003; Ralph et al., 2004). A

similar approach was utilized to identify phosphate translocators in *T. gondii* (i.e.

*Tg*APT1) and *P. falciparum* (i.e. *Pf*iTPT and *Pf*oTPT), and a two-pore channel TPC in

T. gondii. These transporters are capable of supplying phosphorylated carbons,

(Brooks et al., 2010; Fleige et al., 2007; Karnataki et al., 2007; Lim et al., 2010; Mullin

et al., 2006), and calcium to the apicoplast (Li et al., 2021). It was proposed that an

early step in the endosymbiotic event was to integrate transporters that exchange

metabolites across the membranes (Cavalier-Smith, 2000; Weber et al., 2006).

Consistent with that, studies demonstrated that 57-58% of the chloroplast transporters

have an origin in the putative host cell in the first endosymbiosis while 20-30% are

from bacteria (Karkar et al., 2015; Tyra et al., 2007). It is plausible that a complex

transporting system was produced in the evolution of the secondary plastid (i.e. the

apicoplast).

Recent studies suggested that we still do not know the major source of pyruvate

required for the biosynthesis pathways in the apicoplast of *T. gondii* and *P. falciparum*

(Swift et al., 2020; Xia et al., 2019). Therefore, identifying the apicoplast transporter

composition and their functions is critical for an understanding of how these

transporters evolved and how the organelle maintains its metabolic and housekeeping

activities. In this study, we utilized the apicomplexan model organism *T. gondii* to

screen apicoplast membrane proteins, identified from available computational and

proteomic information (Barylyuk et al., 2020; Boucher et al., 2018) and from our

APT1-TurboID proximity proteomic candidates, and discovered several novel

apicoplast transporters. Among the many transporter candidates, one pair of

monocarboxylate transporters (AMTs) appeared to have evolved from the putative

host cell involved in the evolution of the secondary plastid. Detailed analyses support

association of the AMTs with the apicoplast biosynthesis pathways likely by supplying

the pathways with substrates. This study opens up the possibilities of tackling difficult

questions of substrate transport and especially for the pair of monocarboxylate

transporters in *T. gondii* and in other members of this phylum of deadly parasites.

Results

Discovery of novel transporters in the apicoplast of *T. gondii*

To identify apicoplast transporter proteins in *T. gondii*, we first analyzed candidates

identified in recent studies using combined proteomic and computational analyses in *T.*

gondii and *P. falciparum* (Barylyuk et al., 2020; Boucher et al., 2018). The hyperplexed

localization of organelle proteins (hyperLOPIT) assigned 129 proteins to the

apicoplast (Barylyuk et al., 2020), of which three proteins were likely transporters, i.e.

APT1 (Transmembrane Domain, TMD=6), one major facilitator superfamily (MFS)

transporter (TGME49_309580) (HAP1, hypothetical apicoplast protein 1) and one

hypothetical protein (TGME49_267690) (HAP2) (Table S1). We also mined an

apicoplast proteomic dataset obtained by the BioID fusion at acyl carrier protein (ACP)

in *P. falciparum* (Boucher et al., 2018), which reported 16 transmembrane domains

(TMDs) ≥ 6 protein candidates, and two additional candidates annotated as ABC

transporters (TMDs<6). We identified the orthologues in *T. gondii* using the *P.*

falciparum candidates (Boucher et al., 2018; Sayers et al., 2018), resulting in 8

candidates in *T. gondii*. They were temporarily named as HAP3-9 (Table S1). Taken

together, these datasets provided only limited numbers of candidate transporters.

To better define the repertoire of apicoplast transporters, we employed a biotin

proximity labeling strategy that is useful for defining organelle proteomes. A

promiscuous biotin ligase tag called TurboID was endogenously fused with APT1,

using a CRISPR approach (Table S2-4) (Branon et al., 2018; Brooks et al., 2010;

Long et al., 2018). The apicoplast Acetyl-CoA carboxylase 1 (ACC1) is a strongly

biotinylated protein (Zuther et al., 1999), which might provide a clear and strong

streptavidin labeling at the apicoplast (Figure 1A). Though the streptavidin signal and

the protein itself were slightly diffused (Karnataki et al., 2007), APT1 was reported to

mainly localize to four of the apicoplast membranes by an Immuno-Electron

Microscopy approach for HA epitopes tagged at the C-terminus of APT1 (Karnataki et

al., 2007). This feature of APT1 thus provided good biotinylation labeling at potential

proteins localized to membranes of the organelle. Predictably, many additional bands

present on the APT1-TurboID lane were likely to be apicoplast proteins (Figure 1B).

The resultant mass-spectrometry datasets (N=3 for each) (Dataset S1) were analyzed

by comparing the APT1-fusion to the parental line (Figure 1C and Table S5). This

analysis showed that APT1 itself has the highest *p*-value associated with its

enrichment in the TurboID line, as would be expected for a bait protein. The TMD

candidates (TMD≥6 and 2≤TMD<6) were evenly distributed on the volcano map

plotted by fold changes against *p*-values in a similar pattern to the distribution of

known apicoplast proteins (Figure 1C and Table S6). These features suggested that

APT1-TurboID had enriched apicoplast proteins. Thereafter, we compiled a list of the

hits in order of TMD numbers, and selected all transporter candidates (including

carriers) and hypothetical proteins (TMD≥6) as candidates (Table S7). These

candidates were merged with the previously described ones (HAP1-9) and

temporarily named as HAP10-50 (Table S8).

After the above candidate selections, as described in our workflow (Figure1D), we

compiled a list of 50 candidates (HAP1-50) for endogenous tagging and IFA analysis

in *T. gondii*. We successfully generated endogenous HAP-6Ty fusion lines (46 HAPs)

using a CRISPR-Cas9 tagging approach (Figure S1A and Tables S2-4), as described

in our previous study (Long et al., 2017b). Indirect fluorescence assay (IFA) showed

that many HAP-6Ty fusions were localized to foci inside the parasites, resembling the

position of the apicoplast location (Figure S1A), whereas some candidates were likely

to be in the cytosolic vesicles (e.g. HAP13, HAP30, HAP35), the apical region (e.g.

HAP19) or other structures (e.g. HAP22) (Figure S1B). Further analysis was

performed by co-localization of the fusions with an apicoplast marker ACP, which

showed that many of the foci had only partial or no co-localization with ACP (Figure

S2A). However, strict analyses clearly identified 15 HAP fusions that were almost

perfectly co-localized with ACP (Figure 1E).

Of the 15 novel apicoplast membrane proteins, we identified in *T. gondii* that 12

proteins are annotated as putative transporters or carriers in TOXODB, and 3 are

annotated as hypothetical proteins (TMD≥6) (Figure 1F and Table S9). These novel

proteins included 1 from hyperLOPIT, 3 from ACP-BirA and 14 from our APT1-TurboID

(Figure 1G), among which three of these proteins were shared between the

APT1-TurboID and the combinations of hyperLOPIT and ACP-BirA (Figure 1H).

Collectively, TurboID proved to be a reasonably robust approach for the identification

of putative apicoplast transporters.

Novel transporters are mostly conserved in the Apicomplexa

To determine if the apicoplast transporters we identified in *T. gondii* are conserved in

the Apicomplexa, we searched for orthologues of these novel proteins in different

species, as shown in Fig. 2A-B. The chromalveolate hypothesis proposed a plastid

origin in a common ancestor of a single red alga, which unites the Alveolates with the

chromists (Cavalier-Smith, 1999). We identified orthologues in species belonging to

the taxa using a powerful Hidden Markov Model (HMM) profiling strategy. The best

hits (the cutoff E-value: 1×10^{-7}) remained on the list of resultant orthologues in these

species, which were then used for conservation analyses in the ComplexHeatmap

package in R (v4.2.0) (Gu et al., 2016) (Table S10, contains the hits with E-values).

The conservation heatmap showed that proteins with conserved transport domains

(HAP1, 3, 4, 9, 10, 28, 33, 34, 41 and 50) are mostly conserved in the Apicomplexa

(Figure 2A), including the Gregarina and *Cryptosporidium* that have lost the apicoplast.

Further analysis revealed that most orthologues with the exception of HAP3 and

HAP4, can be identified in green algae, red algae and plants (Figure 2B). As the red

alga is considered to have given rise to the ancestor of the secondary plastid (Gould

et al., 2008; Janouskovec et al., 2010), the results thus indicated that those

transporters were possibly derived from the red alga. However, further analysis would

be required to clarify their origins, as shown for chloroplastid transporters (Karkar et

al., 2015).

AMT1 and AMT2 might have critical roles in metabolic integration of the

apicoplast

Several transporters, such as HAP1, 3, 4, and 34 and 50, are predicted to be essential

in *T. gondii* and in *P. falciparum* (Figure 2A-B, and Table S9 contains the *P. falciparum*

orthologues with fitness scores) (Sidik et al., 2016; Zhang et al., 2018). The analyses

with InterProScan showed that three of these transporters (HAP1, 3 and 4) belong to

the major facilitator superfamily (MFS) (Figure S2B). The MFS is a large family of

transporters capable of transporting small solutes in response to chemiosmotic ion

gradients (Pao et al., 1998; Walmsley et al., 1998). These MFS transporters contain a

pair of monocarboxylate transporter (MCT) with 11-12 TMDs (Figure S2C). The MCTs

play essential roles in the transport of nutrients, such as pyruvate, lactate, amino

acids and so on, in the plasma membrane of human cells (Felmlee et al., 2020).

These features attracted our intense attention, and thus named the MCTs as

Apicoplast **M**onocarboxylate **T**ransporters (AMTs, AMT1 for HAP3, TGGT1_233540;

AMT2 for HAP4, TGGT1_297245).

We noticed that these two transporters are absent in red algae, green algae and

plants (Figure 2B). This observation is in contrast to APT1, which was identifiable in

these primary plastid-containing taxa (e.g. plants) but not in cyanobacteria (Figure 2B).

Evolution and phylogenetic analyses suggested that the plant orthologues of APT1

are derived from a putative host cell that previously engulfed a cyanobacteria (Knappe

et al., 2003; Weber et al., 2006). The pair of AMTs appears similar to the APT1

situation, indicating that they were possibly derived from the heterotrophic protist – the

putative host cell - that previously engulfed a red alga. The AMTs were identifiable in

the photosynthetic relative (e.g. *Chromera velia*) of the apicomplexans, and a distant

relative taxa of Oomycetes (Figure 2A), indicating a broader conservation in these

species. We further analyzed orthologues in selected species of apicomplexans and

found that most species contain at least two identifiable orthologues, including

Crytosporidium spp. (Table S11). Collectively, the pair of AMTs is likely to have critical

roles in metabolic integration of the endosymbiont in the parasite cytosol during

evolution.

AMT1 and AMT2 are essential for the parasites

The confocal imaging showed that AMT1-6Ty was almost perfectly co-localized with

AMT2-6HA (Figure S3A), suggesting that the putative transporters are highly likely to

be localized at the apicoplast membranes. In support of this model, immuno-Electron

Microscopy (immuno-EM) provided a further piece of evidence, supporting the

membrane localization of AMT1 and AMT2 (Figure S3B). Due to the complexity of the

apicoplast membranes, and heterogeneity of immunolabeling, the precise topology of

AMT1 and AMT2 within the apicoplast membranes merits future investigation. To

analyze the physiological functions of the AMTs, we generated the conditional

knockdown lines using an IAA-inducible degron (AID) system in *T. gondii* (Brown et al.,

2017; Long et al., 2017b). The CRISPR-Cas9 genome editing approach was utilized

to specifically fuse the AID-6Ty at the 3'-terminus of the genes in the IAA receptor

(TIR1) expressing line RH TIR1 (Figure S4A). To conditionally study both proteins

together, we fused AMT2 with AID-3HA at the 3'-terminus in the background of

AMT1-AID-6Ty, creating a double knockdown line AMT1-AID-6Ty/AMT2-AID-3HA

(dKD). Correct integration of the AID fragments was confirmed by diagnostic PCR as

demonstrated in the schematic diagram (Figure S4A-B). As expected, the AID fusions

were efficiently and stably depleted in IAA treatments for 3-36 hours, as shown on

Western blots (Figure 3A). It was noteworthy that Western blots detected two bands in

the single and double AID fusion lines, which might be the precursor and mature

forms of AMT1 and AMT2 (Figure 3A). So far, several membrane proteins have been

identified in the apicoplast of *T. gondii*, four of which were observed to be processed

(FtsH1, TiC20, ATrx1 and TPC) (DeRocher et al., 2008; Karnataki et al., 2007;

Karnataki et al., 2009; van Dooren et al., 2008). Yet, at least ATrx1 and FtsH1 lack

canonical signal peptides. However, in *P. falciparum,* PfiTPT is processed and cleaved,

while PfoTPT is unprocessed due to its absence of a signal peptide (Mullin et al.,

2006). The processing and topology of AMTs is complex and unpredictable in *T. gondii,*

thus requiring further work to clarify the origin of either the processing hypothesis,

alternative splicing or alternative initiator codons in the parasite. Further analysis by

IFA showed that the AID fusions were correctly localized to the apicoplast and were

depleted within 24 hours of IAA treatment (Figure 3B and Figure S4C); however, the

parasite bodies were morphologically normal. To assess lytic growth of the AID lines,

we assayed plaque formation of the AID parasites on host cell monolayers for 7 days.

We observed that the AID lines grown in IAA did not form discernible plaques (Figure

3C), which was supported by further analysis of the plaque numbers and areas

(Figure 3D-E). In summary, the AMTs are potentially essential for *in vitro* parasite

growth.

We wondered if parasite replication contributed to the plaque defects. The parasites

were grown in IAA for 24 hours, followed by scraping, purification and re-infection for

the assay in the 2nd 24 hours and 3rd 24 hours. Using IFA analysis of the parasites, we

observed that parasite replication was significantly affected in the first 24 hours of

growth and became worse in the following cycles of 24 hours, especially the 3rd 24

hour period (Figure 3F). These results suggested that depletion of the transporters

caused strong growth defects and delayed death of the parasites. We wondered if

depletion of the proteins could cause loss or breakdown of the apicoplast, as reported

previously for many apicoplast proteins (Kennedy et al., 2019b). ACP, an apicoplast

marker, was used to test the apicoplast status (Waller et al., 1998). We observed that

ACP partially diffused in the cytosol of intracellular parasites grown in IAA for 24 hours

(Figure S5A), whereas it completely diffused in the cytosol of extracellular parasites

that had grown in IAA for 24 hours (Figure S5B). This observation suggested swelling

or breakdown of the apicoplast in the intracellular or extracellular parasites depleted

with the AMTs, respectively. Detailed analysis of the phenotypes showed that the ACP

diffusion gradually increased while comparing the three cycles for the same AID line in

both the intracellular and extracellular parasites (Figure 3G-H). However, it was as low

as ~30% in AMT1-AID, compared to those (60-70%) in AMT2-AID and dKD (Figure

3G-H). These results suggested that depletion of AMT2-AID resulted in much more

severe defects (swelling or breakdown) of the apicoplast than that of AMT1-AID, and

further depletion of AMT1-AID had no further effect.

Depletion of the proteins resulted in defects in metabolic pathways

Based on the evolutionary predictions and features of monocarboxylate transporters,

we suspected that the AMTs might be involved in transport of substrates, such as

pyruvate, lactate, amino acids, as reported for human MCTs (Felmlee et al., 2020). As

such, depletion of AMT1 and AMT2 would result in reduced levels of the metabolites

isopentenyl pyrophosphate (isoprenoid precursors, IPP) and fatty acids in *T. gondii*.

We first designed experiments to examine the possible defect of IPP synthesis in the

AID lines upon induction, by assaying the capabilities of endocytic trafficking of GFP

vesicles to the vacuolar compartment (VAC) (Dou et al., 2014). Our recent data

showed that GFP transport to the VAC following its uptake from host cell cytosol is

dependent on protein prenylation of Rab1B and YKT6.1 (Wang et al., 2023). These

proteins are modified at the C-terminal motif by isoprenoids, the synthesis of which

could be specifically inhibited by atorvastatin at the host cells and by zoledronate in

the parasite (Li et al., 2013), as demonstrated in a schematic and in positive

experiments (TIR1 with two drugs at IC50 concentrations) with reduced GFP transport

(Figure 4A-B). Therefore, measuring the endocytic trafficking of GFP vesicles in

parasites in the absence/presence of atorvastatin could reflect levels of protein

prenylation and isoprenoids (IPP). The substrates modified at the proteins were

mainly derived from the apicoplast biosynthesis pathway - the MEP pathway. We then

assayed the TIR1 and AID lines by growing them in GFP expressing HFF monolayers,

where experimental groups were controlled by absence/presence of atorvastatin and

IAA (Figure 4B), and where LHVS was added to avoid degradation of GFP in the VAC.

Following endocytosis of GFP vesicles by the micropore (manuscript in press), GFP

vesicles were successfully transported to and accumulated in the VAC of TIR without

IAA, with a high percentage (>20%) of parasites containing GFP foci (Figure 4B). In

contrast, the GFP vesicles diffused through the cytosol, and were unable to

accumulate at the VAC in the AID parasites grown in IAA with the absence/presence

of atorvastatin, as shown by significantly reduced percentages (~8%) of the parasites

in the AID lines induced for 12 hours (Figure 4B). We further tested Rab1B stability in

the AMT-AID lines grown in IAA for 12 hours, which showed a strong defect of

reduced protein abundance of Rab1B by scoring of parasites with normal or abnormal

level of Rab1B using an IFA approach (Figure S5C). This phenotype was observed in

parasites with a reduced supply of isoprenoids by drug inhibition or knockdown of

prenyl-transferase GGT-2 (unpublished data). These results altogether thus

suggested that the IPP synthesis is significantly decreased in the AID lines depleted

with the transporters. However, it was noteworthy that parasites depleted with

AMT1-AID and AMT2-AID had similar levels of reduced GFP transport and parasites

with reduced protein abundance of Rab1B at the 12 hours of IAA induction,

suggesting that depletions of AMT1 and AMT2 resulted in similar levels of IPP

reduction in parasites.

Our results in *T. gondii* reported a physiological effect that was observed in *P.*

falciparum, where inhibition of the MEP pathway by fosmidomycin caused disruption

of hemoglobin transport (Howe et al., 2013; Kennedy et al., 2019a). Herein, our

results underscored the importance of the transporters for supporting the MEP

pathway in the apicoplast. This was further manifested by a reduced level of

mitochondrial membrane potential detected by mitotracker red in the AMT1-AID or

AMT2-AID lines grown in IAA for 24 hours (Figure S5D). IPP is the precursor for

synthesis of ubiquinone, an essential electron carrier required for electron transport in

the mitochondria (Kennedy et al., 2019b; Sleda et al., 2022). The reduced

mitochondrial potential was also observed in the parasites depleted with APT1

(Brooks et al., 2010). Thus, the reduced mitochondrial membrane potential provided

another piece of evidence supporting the notion of defective IPP synthesis in

parasites depleted with the transporters.

We further examined activity of the FASII pathway in the TIR1 and AID lines grown in

IAA for 18 hours, by testing metabolites (e.g. FA) in the parasites using untargeted

metabolomics (GCMS and LCMS). The metabolomics detected 66 metabolites in total

in these lines (Dataset S2). After analyses using Dunnette's multiple comparison

methods, 15, 23, and 34 differential metabolites were identified in the parasite groups

depleted with either AMT1-AID, AMT2-AID or both (dKD) (Table S12), in comparison

with TIR1. Interestingly, three AID lines shared 13 differential metabolites (Figure 4C).

The shared metabolites included fatty acids and lipids, such as myristic acid (C14 fatty

acid), palmitoleic acid (C16 fatty acid), trans-oleic acid (C18 fatty acid) and

phospholipids containing C14 and C18 fatty acids (Figure 4D-E). These results were

consistent with previous studies where inactivated FASII proteins led to reduced

synthesis of fatty acids up to 18 carbons long, and exogenous additions of the C14

and C16 fatty acids could complement growth defects (Liang et al., 2020;

Ramakrishnan et al., 2012). Studies suggested that lipids can be partially acquired

from the host cells (Krishnan et al., 2020a), which includes those in the endocytic

vesicles (Koreny et al., 2023; Wan et al., 2023). Here we observed that the endocytic

vesicles containing GFP were diffused in the parasite cytosol, indicating that the host

cell-derived lipids were not affected in the protein-depleted parasites. We then

concluded that   depletion of the transporters significantly impaired the *de novo*

biosynthesis of fatty acids by the FASII activity in the parasite. In contrast to the same

level of reduced endocytic trafficking of GFP vesicles in the AID lines, an overall trend

toward lower levels of myristic acid (C14 fatty acid), palmitoleic acid (C16 fatty acid)

and trans-oleic acid (C18 fatty acid) was observed in AMT2-AID and tKD parasites in

comparison with the AMT1-AID parasites (Figure 4E. These results supported the

notion of a stronger impairment of the FASII pathway in parasites depleted with

AMT2-AID.

Intriguingly, the metabolomics also identified metabolites related to the TCA cycle,

such as gamma-aminobutyric acid (GABA), glutamic acid, and succinic acid (Figure

4E), which were shared among three of the AID lines. This would likely result from

defective membrane potential caused by a reduced supply of IPP for synthesis of

ubiquinone, as described above, or synthesis of heme that requires co-operation

between the mitochondrion and the apicoplast (Harding et al., 2020). Another two

metabolites, creatine and lactic acid, were likely associated with reduced metabolic

levels in the parasites (Figure 4E). The untargeted metabolomics tends to identify

differential metabolites by comparing the AID to TIR1 lines. Pyruvate was not

identified, as it is actually a product of glycolysis in the cytosol. Collectively, the

transporters are expectedly associated with the MEP and FASII pathways, to support

synthesis of IPP and FA in the apicoplast.

Depletion of AMT2 affects stability of ACC1

Depletion of AMT2 resulted in stronger defects on ACP diffusion and synthesis of fatty

acids, suggesting a more important role of AMT2 in the apicoplast. To find out the

reason for this, we first examined the protein level of one AMT (fused with 6HA) in

another AMT's AID-Ty line, thus assuring no reciprocal effect on protein level. After

IAA induction for 36 hours, we did not see obvious changes to protein levels in the

6HA fusions in the AID-Ty lines on western blots and IFA (Figure 5A-B). Considering a

MCT could transport biotin in mammalian cells (Daberkow et al., 2003), we wondered

if AMT2 was able to transport biotin for activation of acetyl-CoA carboxylase ACC1,

which is a key enzyme in FASII (Zuther et al., 1999). ACC1 was endogenously fused

with 6HA in the AID lines by a CRISPR approach. Upon protein depletion in IAA for 36

hours, ACC1 was found to be gone only in the AMT2-AID parasites, but not in the

AMT1-AID parasites, as demonstrated by Western blots (Figure 5C). Further analysis

showed that ACC1 was gone in some parasites within a single vacuole in the

AMT2-AID-depleted parasites, where ACP was clearly stained with a slight diffusion

(Figure 5D). Based on the presence or absence of ACC1 in parasites of the vacuoles,

we can classify the vacuoles into three types - all parasites without ACC1 (type 1),

one parasite (type 2) or two parasites (type 3) with ACC1 (Figure 5E). The

quantification assay showed that only ~40% of the vacuoles contained one or two

parasites with ACC1 signal (type 2 and type 3) (Figure 5F). However, ACC1 in those

parasites was still biotinylated, as shown in streptavidin stains (Figure 5G).

Collectively, AMT2 is associated with stability of ACC1 in a way, as yet, unknown, but

unrelated with biotin availability in the apicoplast.

AMT2 is essential to parasite virulence in mice

We next examined the *in vivo* growth of the AID parasites in mice administered with

IAA. Prior to the mice assay, we attempted to gain further information about the

ultra-structures of the apicoplast upon depletion of the AMTs. The transmission

electron microscopy (TEM) demonstrated that the four membranes surrounding the

apicoplast remained visible, but were loosened, and the interior of the organelle

appeared less dense and even electron lucent in the AID parasites, which had been

induced for 24 hours in comparison with the TIR1 parasites (Figure 6A). We further

quantified the apicoplasts with a reduced electron density in the AID parasites induced

for 12, 24 and 36 hours vs the TIR1 in IAA for 36 hours, by examining 300 parasites in

each time point. This examination showed that the apicoplast numbers identified by

TEM dropped in the AMT1-AID parasites, when induced for 24 hours and 36 hours,

and the numbers dropped by much greater levels in the AMT2-AID parasites,

following 12 hours of IAA induction (Figure 6B). Notably, the apicoplast was identified

only 3 times in the AMT2-AID parasites at 36 hours induction. Accordingly, the

percentage of the abnormal apicoplast increased for both of the AID lines. In contrast,

depletion of AMT2-AID resulted in much higher levels of abnormal apicoplasts at the

three induction time points, compared to those in parasites depleted with AMT1-AID

(Figure 6B). These results suggested that depletion of AMT1 caused modest loss of

the apicoplast, yet the AMT2 depletion appeared to have a greater effect on the

apicoplast loss.

The parental line TIR1 and its derivative lines (AMT1-AID, AMT2-AID and dKD) were

then used to intraperitoneally infect mice, at 100 parasites per mouse, followed by

administering the mice orally and intraperitoneally with IAA in a solution on a daily

basis. Those mice infected with TIR1 succumbed to lethal toxoplasmosis by day 10

post infection (Figure 6C). Conversely, the IAA treatment ultimately prevented acute

virulence caused by the parasite lines with the AMT2 depletion (i.e. AMT2-AID and

dKD), and it postponed mouse death by the AMT1-AID depletion. We next tested sera

extracted from the surviving mice (which were infected by AMT2-AID) by performing

IFA on *T. gondii* parasites. The IFA showed that the mouse sera clearly recognized

proteins of the parasitophorous vacuoles of the parasites (Figure S6A). We also

further analyzed the efficacy of protein degradation of the AMT1-AID fusions in mice,

by treatment with or without IAA for 24 hours after 5 days of growth, as illustrated in

Figure S7B. It was clear that AMT1-AID was efficiently degraded in mice by the

treatment of IAA in only 24 hours (Figure S6B). These results demonstrated that the

mice infection assays were robust. However, the strong growth impairment conferred

by AMT1 depletion in parasites in cell culture surprisingly resulted in only reduced

parasite virulence in mice. We assume that this discrepancy is related to the relatively

modest defects, such as the ACP diffusion/apicoplast abnormality (revealed by

TEM)/FASII activity, as observed in the AMT1-depleted parasites. Thus, the remaining

apicoplast is still functionally generating an adequate amount of metabolites, allowing

the parasites to survive in mice. However, the discrepancy is possibly associated with

different transport capabilities of substrates by AMT1 and AMT2. Though this does not

perfectly explain the discrepancy, weaker phenotypes in AMT1-depleted parasites

argue for this possibility. Collectively, AMT2 was obviously essential for establishing *T.*

gondii infection and virulence in mice.

Discussion

Most of the apicomplexan parasites, such as *T. gondii* and *Plasmodium spp.*, harbor a

vestigial plastid, which undergoes several fitness-conferring or essential metabolic

pathways in the matrix (Kloehn et al., 2021; Lim and McFadden, 2010). However, the

transporters that are responsible for exchanging metabolites between the apicoplast

and the parasite remain elusive, due to difficulties in identification of those

transporters in the multiple membranes that surround the organelle. Here we fully

exploited the currently available data, and the high-efficiency of proximity biotin

labeling and CRISPR-Cas9 tagging approaches in the model apicomplexan *T. gondii*,

and discovered several novel transporters in the apicoplast. Among those putative

transporters, most of them appear to be derived from the endosymbiont (i.e. a red

alga) in the evolution of the secondary plastid, and only the pair of monocarboxylate

transporters AMT1 and AMT2 is likely to be derived from the putative host cell that

originally engulfed the endosymbiont. Protein depletion of the AMT1 and AMT2 using

a highly efficient technique, the TIR1-AID system, resulted in defects of plaque

formation *in vitro*, ACP diffusion/apicoplast loss and biosynthetic pathways of the MEP

and FASII. However, depletion of AMT2 clearly caused stronger defects including

ACP diffusion/apicoplast loss, FASII produced fatty acids, ACC1 stability and removal

of mouse virulence. These results indicate that complex transporting functions are

likely to exist in the apicoplast involving AMT1 and AMT2.

Recent studies have highlighted the difficulty in identifying apicoplast transporters,

because of strong backgrounds in the proteomics resulting from high activity and

nonspecific labeling of the biotin ligase (BioID) fused with apicoplast proteins

(Boucher et al., 2018; Sayers et al., 2018). This remained true for our proximity

proteomics from APT1-TurboID. Fortunately, our focused screening leveraged high

efficiency by genomic editing in *T. gondii,* yielding positive results for 12 putative

transporters. These novel transporters provided an important resource for evolution

and functional analyses. Their conservation, including in plastid-deficient taxa, i.e.

Gregarina and Cryptosporidia, indicates that these proteins were relics of the former

plastid in evolution via endosymbiont gene transfers (EGT) (Gould et al., 2008).

Studies suggested that the metabolic integration of the primary plastid (e.g.

chloroplast) was mainly driven by targeting transporters from a heterotrophic protist -

the host cell, but not from the endosymbiont - cyanobacteria (Karkar et al., 2015; Tyra

et al., 2007). However, our preliminary study in evolution indicated that most novel

transporters were likely to be derived from the endosymbiont. In sharp contrast, the

pair of monocarboxylate transporters (AMT1 and AMT2) may have an origin in a

heterotrophic protist – the putative host cell. This observation therefore provides a

perspective that the pair of AMTs may play critical roles in the metabolic integration of

the algal endosymbiont into the parasites.

Our immuno-EM and confocal imaging provided a strong support for the membrane

localization of the pair of transporters AMT1 and AMT2 in the apicoplast, indicating

that the AMTs are likely to transport substrates in the membranes. We observed that

the AMTs appeared to have two bands on Western blot, which might be processed

versions, or produced by alternative splicing or produced by an alternative initiator

codon. Concerning the processing and targeting, apicoplast membrane proteins in *T.*

gondii appear to be more complex than those in *P. falciparum*, by comparison to

PfiTPT and PfoTPT (Mullin et al., 2006). Our phenotypic analysis showed that

depletion of the transporters AMT1 and AMT2 caused strong defects on plaque

formation and parasite replication, suggesting that both transporters are critical on the

apicoplast membranes. However, we observed disparate degrees of phenotypic

disruption in the AMT1-depleted and AMT2-depleted parasites. Depletion of AMT2

always caused stronger defects in ACP diffusion, FASII fatty acids, apicoplast loss

and mouse virulence. AMT2 is additionally associated with loss of ACC1, a key

enzyme for FASII, found in the majority of parasites in a vacuole, likely providing

another piece of evidence to explain the greater phenotypic defects. Several attempts

to delete ACC1 failed in our study, supporting a critical role of ACC1 in the FASII

process. Genetic studies suggest that depletion of FASII enzymes led to loss of

apicoplast (Krishnan et al., 2020b; Liang et al., 2020). Thus, the observation of ACC1

loss supports stronger phenotypes in AMT2-AID depleted parasites. In addition,

depletion of the AMTs caused an initially gradual defect, but then a much more

focused sharper defect in parasite replication in the 3rd 24 hours and showed a

delayed death phenotype. Previous studies showed that parasite replication exhibited

gradual deceases upon depletion of TPC and APT1 by utilization of a low efficient

knockdown strategy - the tetracyclin-inducible system (TATi) (Brooks et al., 2010; Li et

al., 2021). It is unknown whether the TATi system would mask the growth phenotype,

yet, the AMT depletion appeared to have such a sharper change in the third cycle (24

hours in a growth cycle). Collectively, the AMTs are critical to parasites, yet AMT2 is

more important to the apicoplast membrane function.

Based on knowledge of MCTs (Felmlee et al., 2020), the pair of AMTs is likely to

transport pyruvate, lactate, amino acids and other substrates. We thus suspect that

AMT1 and AMT2 might be associated with transporting pyruvate and amino acids,

since these are essential for the apicoplast metabolism. Pyruvate is the central

metabolite that connects many metabolic pathways in living organisms. It must be

imported into organelles for specific metabolic pathways, as suggested with pyruvate

import by the NHD1/Bass2-protein coupling function in the chloroplast of C4 plants,

and by the mitochondrial pyruvate carrier MPC (Bricker et al., 2012; Furumoto, 2016;

Furumoto et al., 2011; Gasperotti et al., 2020; Herzig et al., 2012). In *T. gondii*, and *P.*

falciparum, studies have suggested that the major pyruvate source has still not been

established (Niu et al., 2022; Swift et al., 2020; Xia et al., 2019). Pyruvate is one of the

key carbon sources for the condensation with triose phosphates to initiate the

biosynthesis pathway of MEP in the parasite. Meanwhile, pyruvate is the only carbon

source in the FASII pathway for biosynthesis of fatty acids (Brooks et al., 2010). The

possible role on pyruvate import by the AMTs could provide a good explanation for the

defects on MEP and FASII pathways, specifically, the phenotypes reflecting the

reduced levels of IPP synthesis in the AMT-depleted parasites. We attempted to

define this possible function of the AMTs by a pyruvate sensor PyronicSF in the

parasite (Arce-Molina et al., 2020), and by a complementation of an *Escherichia coli*

knockout strain that is deficient in pyruvate import (a generous gift from Prof. Kirsten

Jung) (Gasperotti et al., 2020).Hhowever, several attempts with these approaches

were not successful in generating the strain required for detailed examination of

pyruvate import capability either in the parasite or in the *E. coli* strain.

Our work successfully identifies several novel apicoplast transporters, thus greatly

unravelling the mystery of the apicoplast membrane composition and opening up

possibilities to tackle difficult questions about metabolite transport in the

apicomplexans. The most intriguingly findings in this study are the identification of the

pair of monocarboxylate transporters AMT1 and AMT2, which are likely derived from

the putative host cell that engulfed a red algal during the evolution of this group of

parasites. Metabolic analysis together with other phenotypic analyses supported the

notion that the pair of AMTs is associated with the MEP and FASII biosynthesis

pathways. However, though both transporters belong to the MCTs, the phenotypic

defects differ greatly, supporting a complex function for AMT2 in the apicoplast,

however, the substrates transported by the AMTs still await further clarification.

Collectively, the monocarboxylate transporters are likely to control similar essential

processes in related protists, especially in other members of this phylum of deadly

parasites.

Limitations of This Study

This study has begun to look into the possibly evolutional origins of newly identified

transporters although, as yet, observations are preliminary and await further analysis

to clarify specific questions, as demonstrated in the detailed evolutionary analysis of

chloroplast transporters (Karkar et al., 2015). Our further analysis on the phenotypes

of two of these transporters, AMT1 and AMT2, using parasites depleted with these

activities, demonstrates differing degrees of defects at critical points in the parasite's

physiology and metabolism. We also observed similar phenotypes reflecting

decreasedlevels of the MEP biosynthesis activity upon depletion of AMT1 and AMT2,

indicating that phenotypic outcomes did not result from apicoplast abnormality, since

other phenotypic outcomes were more severe in the AMT2-depleted parasites. We

predict, therefore, that the metabolic defects possibly result from a reduced supply of

substrates by each of the transporter. It is noteworthy that other studies provided us

with comparable knowledge in phenotypes in TPC-depleted and APT1-depleted

parasites. In these parasites, metabolic defects in the apicoplast pathways were

observed by indirect measurements as well (Brooks et al., 2010; Li et al., 2021).

However, these phenotypes were possibly caused by pleiotropic effects of apicoplast

loss, since they were analyzed by a comparably slow inactivation system to deplete

the transporters of TPC and APT1 (with an induction of at least 3-4 days).

Nevertheless, in our case with a highly efficient inactivation system, we still cannot

exclude the effect of apicoplast loss on phenotypic results caused by depletion of the

AMTs. Therefore, a more precise understanding of the functions of the AMTs will

require further identification of the topology, protein interactions and substrate

transport capability in the complex configuration of the apicoplast membranes. Our

work, as yet, unsuccessful, in attempting to use a pyruvate sensor and/or a

heterologous expression the AMTs in *E. coli*, is the first attempt to examine the

transport capability of pyruvate by AMT1 and AMT2. Thereafter, the application of

more sophisticated approaches, such as *Xenopus* oocytes, an improved *E. coli*

system, or a yeast expression system, may enable us to tackle the important question

of substrate transport by AMT1 and AMT2. Future progress in this direction will thus

shed critical insights into the discrepancies in parasite growth that we observe when

comparing cell culture versus infections in mice.

REFERENCES

Amiar, S., Katris, N.J., Berry, L., Dass, S., Duley, S., Arnold, C.S., Shears, M.J., Brunet,
C., Touquet, B., McFadden, G.I.*, et al.* (2020). Division and Adaptation to Host
Environment of Apicomplexan Parasites Depend on Apicoplast Lipid Metabolic
Plasticity and Host Organelle Remodeling. Cell Rep *30*, 3778-3792.e3779.
Amiar, S., MacRae, J.I., Callahan, D.L., Dubois, D., van Dooren, G.G., Shears, M.J.,
Cesbron-Delauw, M.F., Maréchal, E., McConville, M.J., McFadden, G.I.*, et al.* (2016).
Apicoplast-Localized Lysophosphatidic Acid Precursor Assembly Is Required for Bulk
Phospholipid Synthesis in Toxoplasma gondii and Relies on an Algal/Plant-Like
Glycerol 3-Phosphate Acyltransferase. PLoS Pathog *12*, e1005765.
Arce-Molina, R., Cortes-Molina, F., Sandoval, P.Y., Galaz, A., Alegria, K., Schirmeier, S.,
Barros, L.F., and San Martin, A. (2020). A highly responsive pyruvate sensor reveals
pathway-regulatory role of the mitochondrial pyruvate carrier MPC. Elife *9*.
Barylyuk, K., Koreny, L., Ke, H., Butterworth, S., Crook, O.M., Lassadi, I., Gupta, V.,
Tromer, E., Mourier, T., Stevens, T.J.*, et al.* (2020). A Comprehensive Subcellular

Atlas of the Toxoplasma Proteome via hyperLOPIT Provides Spatial Context for
Protein Functions. Cell Host Microbe *28*, 752-766 e759.
Bastin, P., Bagherzadeh, Z., Matthews, K.R., and Gull, K. (1996). A novel epitope tag
system to study protein targeting and organelle biogenesis in *Trypanosoma brucei*.
Molec Biochem Parasitol *77*, 235-239.
Botté, C.Y., Yamaryo-Botté, Y., Rupasinghe, T.W., Mullin, K.A., MacRae, J.I., Spurck,
T.P., Kalanon, M., Shears, M.J., Coppel, R.L., Crellin, P.K.*, et al.* (2013). Atypical lipid
composition in the purified relict plastid (apicoplast) of malaria parasites. Proc Natl
Acad Sci U S A *110*, 7506-7511.
Boucher, M.J., Ghosh, S., Zhang, L., Lal, A., Jang, S.W., Ju, A., Zhang, S., Wang, X.,
Ralph, S.A., Zou, J.*, et al.* (2018). Integrative proteomics and bioinformatic prediction
enable a high-confidence apicoplast proteome in malaria parasites. PLoS Biol *16*,
e2005895.
Branon, T.C., Bosch, J.A., Sanchez, A.D., Udeshi, N.D., Svinkina, T., Carr, S.A.,
Feldman, J.L., Perrimon, N., and Ting, A.Y. (2018). Efficient proximity labeling in living
cells and organisms with TurboID. Nat Biotechnol *36*, 880-887.
Bricker, D.K., Taylor, E.B., Schell, J.C., Orsak, T., Boutron, A., Chen, Y.C., Cox, J.E.,
Cardon, C.M., Van Vranken, J.G., Dephoure, N.*, et al.* (2012). A mitochondrial
pyruvate carrier required for pyruvate uptake in yeast, Drosophila, and humans.
Science *337*, 96-100.
Brooks, C.F., Johnsen, H., van Dooren, G.G., Muthalagi, M., Lin, S.S., Bohne, W.,
Fischer, K., and Striepen, B. (2010). The toxoplasma apicoplast phosphate
translocator links cytosolic and apicoplast metabolism and is essential for parasite
survival. Cell Host Microbe *7*, 62-73.
Brown, K.M., Long, S., and Sibley, L.D. (2017). Plasma Membrane Association by
N-Acylation Governs PKG Function in Toxoplasma gondii. mBio *8*.
Brown, K.M., Long, S., and Sibley, L.D. (2018). Conditional Knockdown of Proteins
Using Auxin-inducible Degron (AID) Fusions in Toxoplasma gondii. Bio Protoc *8*.
Brown, K.M., and Sibley, L.D. (2018). Essential cGMP Signaling in Toxoplasma Is
Initiated by a Hybrid P-Type ATPase-Guanylate Cyclase. Cell Host Microbe *24*,
804-816 e806.
Cavalier-Smith, T. (1999). Principles of protein and lipid targeting in secondary
symbiogenesis: euglenoid, dinoflagellate, and sporozoan plastid origins and the
eukaryote family tree. J Eukaryot Microbiol *46*, 347-366.
Cavalier-Smith, T. (2000). Membrane heredity and early chloroplast evolution. Trends
Plant Sci *5*, 174-182.
Daberkow, R.L., White, B.R., Cederberg, R.A., Griffin, J.B., and Zempleni, J. (2003).
Monocarboxylate transporter 1 mediates biotin uptake in human peripheral blood
mononuclear cells. J Nutr *133*, 2703-2706.
DeRocher, A.E., Coppens, I., Karnataki, A., Gilbert, L.A., Rome, M.E., Feagin, J.E.,
Bradley, P.J., and Parsons, M. (2008). A thioredoxin family protein of the apicoplast
periphery identifies abundant candidate transport vesicles in Toxoplasma gondii.
Eukaryot Cell *7*, 1518-1529.

Dou, Z., McGovern, O.L., Di Cristina, M., and Carruthers, V.B. (2014). Toxoplasma
gondii ingests and digests host cytosolic proteins. mBio *5*, e01188-01114.
Fata, J.J., and Rabuzzi, D.D. (1988). Aggressive juvenile fibromatosis presenting as a
parotid mass. Ear Nose Throat J *67*, 678, 681-672, 684 passim.
Felmlee, M.A., Jones, R.S., Rodriguez-Cruz, V., Follman, K.E., and Morris, M.E. (2020).
Monocarboxylate Transporters (SLC16): Function, Regulation, and Role in Health and
Disease. Pharmacol Rev *72*, 466-485.
Fleige, T., Fischer, K., Ferguson, D.J., Gross, U., and Bohne, W. (2007). Carbohydrate
metabolism in the Toxoplasma gondii apicoplast: localization of three glycolytic
isoenzymes, the single pyruvate dehydrogenase complex, and a plastid phosphate
translocator. Eukaryot Cell *6*, 984-996.
Foth, B.J., and McFadden, G.I. (2003). The apicoplast: a plastid in Plasmodium
falciparum and other Apicomplexan parasites. Int Rev Cytol *224*, 57-110.
Foth, B.J., Ralph, S.A., Tonkin, C.J., Struck, N.S., Fraunholz, M., Roos, D.S., Cowman,
A.F., and McFadden, G.I. (2003). Dissecting apicoplast targeting in the malaria
parasite Plasmodium falciparum. Science *299*, 705-708.
Furumoto, T. (2016). Pyruvate transport systems in organelles: future directions in C4
biology research. Curr Opin Plant Biol *31*, 143-148.
Furumoto, T., Yamaguchi, T., Ohshima-Ichie, Y., Nakamura, M., Tsuchida-Iwata, Y.,
Shimamura, M., Ohnishi, J., Hata, S., Gowik, U., Westhoff, P.*, et al.* (2011). A plastidial
sodium-dependent pyruvate transporter. Nature *476*, 472-475.
Gao, X., Pujos-Guillot, E., and Sebedio, J.L. (2010). Development of a quantitative
metabolomic approach to study clinical human fecal water metabolome based on
trimethylsilylation derivatization and GC/MS analysis. Anal Chem *82*, 6447-6456.
Gasperotti, A., Going, S., Fajardo-Ruiz, E., Forne, I., and Jung, K. (2020). Function and
Regulation of the Pyruvate Transporter CstA in Escherichia coli. Int J Mol Sci *21*.
Gould, S.B., Waller, R.F., and McFadden, G.I. (2008). Plastid evolution. Annu Rev Plant
Biol *59*, 491-517.
Gu, Z., Eils, R., and Schlesner, M. (2016). Complex heatmaps reveal patterns and
correlations in multidimensional genomic data. Bioinformatics *32*, 2847-2849.
Harding, C.R., Sidik, S.M., Petrova, B., Gnadig, N.F., Okombo, J., Herneisen, A.L.,
Ward, K.E., Markus, B.M., Boydston, E.A., Fidock, D.A.*, et al.* (2020). Genetic screens
reveal a central role for heme metabolism in artemisinin susceptibility. Nat Commun
11, 4813.
Heaslip, A.T., Dzierszinski, F., Stein, B., and Hu, K. (2010). TgMORN1 is a key
organizer for the basal complex of Toxoplasma gondii. PLoS Pathog *6*, e1000754.
Herzig, S., Raemy, E., Montessuit, S., Veuthey, J.L., Zamboni, N., Westermann, B.,
Kunji, E.R., and Martinou, J.C. (2012). Identification and functional expression of the
mitochondrial pyruvate carrier. Science *337*, 93-96.
Howe, R., Kelly, M., Jimah, J., Hodge, D., and Odom, A.R. (2013). Isoprenoid
biosynthesis inhibition disrupts Rab5 localization and food vacuolar integrity in
Plasmodium falciparum. Eukaryot Cell *12*, 215-223.
Huynh, M.H., and Carruthers, V.B. (2009). Tagging of endogenous genes in a

Toxoplasma gondii strain lacking Ku80. Eukaryot Cell *8*, 530-539.

Janouskovec, J., Horak, A., Obornik, M., Lukes, J., and Keeling, P.J. (2010). A common
red algal origin of the apicomplexan, dinoflagellate, and heterokont plastids. Proc Natl
Acad Sci U S A *107*, 10949-10954.

Jomaa, H., Wiesner, J., Sanderbrand, S., Altincicek, B., Weidemeyer, C., Hintz, M.,
Türbachova, I., Eberl, M., Zeidler, J., Lichtenthaler, H.K.*, et al.* (1999). Inhibitors of the
nonmevalonate pathway of isoprenoid biosynthesis as antimalarial drugs. Science
285, 1573-1576.

Karkar, S., Facchinelli, F., Price, D.C., Weber, A.P., and Bhattacharya, D. (2015).
Metabolic connectivity as a driver of host and endosymbiont integration. Proc Natl
Acad Sci U S A *112*, 10208-10215.

Karnataki, A., Derocher, A., Coppens, I., Nash, C., Feagin, J.E., and Parsons, M. (2007).
Cell cycle-regulated vesicular trafficking of Toxoplasma APT1, a protein localized to
multiple apicoplast membranes. Mol Microbiol *63*, 1653-1668.

Karnataki, A., DeRocher, A.E., Feagin, J.E., and Parsons, M. (2009). Sequential
processing of the Toxoplasma apicoplast membrane protein FtsH1 in topologically
distinct domains during intracellular trafficking. Mol Biochem Parasitol *166*, 126-133.

Kennedy, K., Cobbold, S.A., Hanssen, E., Birnbaum, J., Spillman, N.J., McHugh, E.,
Brown, H., Tilley, L., Spielmann, T., McConville, M.J.*, et al.* (2019a). Delayed death in
the malaria parasite Plasmodium falciparum is caused by disruption of
prenylation-dependent intracellular trafficking. PLoS Biol *17*, e3000376.

Kennedy, K., Crisafulli, E.M., and Ralph, S.A. (2019b). Delayed Death by Plastid
Inhibition in Apicomplexan Parasites. Trends Parasitol *35*, 747-759.

Kloehn, J., Lacour, C.E., and Soldati-Favre, D. (2021). The metabolic pathways and
transporters of the plastid organelle in Apicomplexa. Curr Opin Microbiol *63*, 250-258.

Knappe, S., Flugge, U.I., and Fischer, K. (2003). Analysis of the plastidic phosphate
translocator gene family in Arabidopsis and identification of new phosphate
translocator-homologous transporters, classified by their putative substrate-binding
site. Plant Physiol *131*, 1178-1190.

Köhler, S., Delwiche, C.F., Denny, P.W., Tilney, L.G., Webster, P., Wilson, R.J., Palmer,
J.D., and Roos, D.S. (1997). A plastid of probable green algal origin in Apicomplexan
parasites. Science *275*, 1485-1489.

Koreny, L., Mercado-Saavedra, B.N., Klinger, C.M., Barylyuk, K., Butterworth, S., Hirst,
J., Rivera-Cuevas, Y., Zaccai, N.R., Holzer, V.J.C., Klingl, A.*, et al.* (2023). Stable
endocytic structures navigate the complex pellicle of apicomplexan parasites. Nat
Commun *14*, 2167.

Krishnan, A., Kloehn, J., Lunghi, M., Chiappino-Pepe, A., Waldman, B.S., Nicolas, D.,
Varesio, E., Hehl, A., Lourido, S., Hatzimanikatis, V.*, et al.* (2020a). Functional and
Computational Genomics Reveal Unprecedented Flexibility in Stage-Specific
Toxoplasma Metabolism. Cell Host Microbe *27*, 290-306.e211.

Krishnan, A., Kloehn, J., Lunghi, M., Chiappino-Pepe, A., Waldman, B.S., Nicolas, D.,
Varesio, E., Hehl, A., Lourido, S., Hatzimanikatis, V.*, et al.* (2020b). Functional and
Computational Genomics Reveal Unprecedented Flexibility in Stage-Specific

Toxoplasma Metabolism. Cell Host Microbe *27*, 290-306 e211.

Li, Z.H., King, T.P., Ayong, L., Asady, B., Cai, X., Rahman, T., Vella, S.A., Coppens, I.,
Patel, S., and Moreno, S.N.J. (2021). A plastid two-pore channel essential for
inter-organelle communication and growth of Toxoplasma gondii. Nat Commun *12*,
5802.

Li, Z.H., Ramakrishnan, S., Striepen, B., and Moreno, S.N. (2013). Toxoplasma gondii
relies on both host and parasite isoprenoids and can be rendered sensitive to
atorvastatin. PLoS Pathog *9*, e1003665.

Liang, X., Cui, J., Yang, X., Xia, N., Li, Y., Zhao, J., Gupta, N., and Shen, B. (2020).
Acquisition of exogenous fatty acids renders apicoplast-based biosynthesis
dispensable in tachyzoites of Toxoplasma. J Biol Chem *295*, 7743-7752.

Lim, L., Linka, M., Mullin, K.A., Weber, A.P., and McFadden, G.I. (2010). The carbon
and energy sources of the non-photosynthetic plastid in the malaria parasite. FEBS
Lett *584*, 549-554.

Lim, L., and McFadden, G.I. (2010). The evolution, metabolism and functions of the
apicoplast. Philos Trans R Soc Lond B Biol Sci *365*, 749-763.

Long, S., Anthony, B., Drewry, L.L., and Sibley, L.D. (2017a). A conserved ankyrin
repeat-containing protein regulates conoid stability, motility and cell invasion in
Toxoplasma gondii. Nat Commun *8*, 2236.

Long, S., Brown, K.M., Drewry, L.L., Anthony, B., Phan, I.Q.H., and Sibley, L.D. (2017b).
Calmodulin-like proteins localized to the conoid regulate motility and cell invasion by
Toxoplasma gondii. PLoS Pathog *13*, e1006379.

Long, S., Brown, K.M., and Sibley, L.D. (2018). CRISPR-mediated Tagging with BirA
Allows Proximity Labeling in Toxoplasma gondii. Bio Protoc *8*.

Long, S., Wang, Q., and Sibley, L.D. (2016). Analysis of Noncanonical
Calcium-Dependent Protein Kinases in Toxoplasma gondii by Targeted Gene Deletion
Using CRISPR/Cas9. Infect Immun *84*, 1262-1273.

Low, L.M., Stanisic, D.I., and Good, M.F. (2018). Exploiting the apicoplast:
apicoplast-targeting drugs and malaria vaccine development. Microbes Infect *20*,
477-483.

McFadden, G.I., Reith, M.E., Munholland, J., and Lang-Unnasch, N. (1996). Plastid in
human parasites. Nature *381*, 482.

McFadden, G.I., and Roos, D.S. (1999). Apicomplexan plastids as drug targets. Trends
Microbiol *7*, 328-333.

McFadden, G.I., and van Dooren, G.G. (2004). Evolution: red algal genome affirms a
common origin of all plastids. Curr Biol *14*, R514-516.

McFadden, G.I., and Yeh, E. (2017). The apicoplast: now you see it, now you don't. Int J
Parasitol *47*, 137-144.

Mullin, K.A., Lim, L., Ralph, S.A., Spurck, T.P., Handman, E., and McFadden, G.I.
(2006). Membrane transporters in the relict plastid of malaria parasites. Proc Natl
Acad Sci U S A *103*, 9572-9577.

Niu, Z., Ye, S., Liu, J., Lyu, M., Xue, L., Li, M., Lyu, C., Zhao, J., and Shen, B. (2022).
Two apicoplast dwelling glycolytic enzymes provide key substrates for metabolic

pathways in the apicoplast and are critical for Toxoplasma growth. PLoS Pathog *18*,
e1011009.
Obornik, M., Janouskovec, J., Chrudimsky, T., and Lukes, J. (2009). Evolution of the
apicoplast and its hosts: from heterotrophy to autotrophy and back again. Int J
Parasitol *39*, 1-12.
Pao, S.S., Paulsen, I.T., and Saier, M.H., Jr. (1998). Major facilitator superfamily.
Microbiol Mol Biol Rev *62*, 1-34.
Plattner, F., and Soldati-Favre, D. (2008). Hijacking of host cellular functions by the
Apicomplexa. Annu Rev Microbiol *62*, 471-487.
Primo, V.A., Jr., Rezvani, Y., Farrell, A., Murphy, C.Q., Lou, J., Vajdi, A., Marth, G.T.,
Zarringhalam, K., and Gubbels, M.J. (2021). The Extracellular Milieu of Toxoplasma's
Lytic Cycle Drives Lab Adaptation, Primarily by Transcriptional Reprogramming.
mSystems *6*, e0119621.
Ralph, S.A., van Dooren, G.G., Waller, R.F., Crawford, M.J., Fraunholz, M.J., Foth, B.J.,
Tonkin, C.J., Roos, D.S., and McFadden, G.I. (2004). Tropical infectious diseases:
metabolic maps and functions of the Plasmodium falciparum apicoplast. Nat Rev
Microbiol *2*, 203-216.
Ramakrishnan, S., Docampo, M.D., Macrae, J.I., Pujol, F.M., Brooks, C.F., van Dooren,
G.G., Hiltunen, J.K., Kastaniotis, A.J., McConville, M.J., and Striepen, B. (2012).
Apicoplast and endoplasmic reticulum cooperate in fatty acid biosynthesis in
apicomplexan parasite Toxoplasma gondii. J Biol Chem *287*, 4957-4971.
Salomaki, E.D., and Kolisko, M. (2019). There Is Treasure Everywhere: Reductive
Plastid Evolution in Apicomplexa in Light of Their Close Relatives. Biomolecules *9*.
Sayers, C.P., Mollard, V., Buchanan, H.D., McFadden, G.I., and Goodman, C.D. (2018).
A genetic screen in rodent malaria parasites identifies five new apicoplast putative
membrane transporters, one of which is essential in human malaria parasites. Cell
Microbiol *20*.
Sidik, S.M., Huet, D., Ganesan, S.M., Huynh, M.H., Wang, T., Nasamu, A.S., Thiru, P.,
Saeij, J.P.J., Carruthers, V.B., Niles, J.C.*, et al.* (2016). A Genome-wide CRISPR
Screen in Toxoplasma Identifies Essential Apicomplexan Genes. Cell *166*, 1423-1435
e1412.
Sleda, M.A., Li, Z.H., Behera, R., Baierna, B., Li, C., Jumpathong, J., Malwal, S.R.,
Kawamukai, M., Oldfield, E., and Moreno, S.N.J. (2022). The Heptaprenyl
Diphosphate Synthase (Coq1) Is the Target of a Lipophilic Bisphosphonate That
Protects Mice against Toxoplasma gondii Infection. mBio *13*, e0196622.
Soldati, D., and Boothroyd, J.C. (1993). Transient transfection and expression in the
obligate intracellular parasite Toxoplasma gondii. Science *260*, 349-352.
Striepen, B. (2011). The apicoplast: a red alga in human parasites. Essays Biochem *51*,
111-125.
Swift, R.P., Rajaram, K., Keutcha, C., Liu, H.B., Kwan, B., Dziedzic, A., Jedlicka, A.E.,
and Prigge, S.T. (2020). The NTP generating activity of pyruvate kinase II is critical for
apicoplast maintenance in Plasmodium falciparum. Elife *9*.
Tjhin, E.T., Hayward, J.A., McFadden, G.I., and van Dooren, G.G. (2020).

Characterization of the apicoplast-localized enzyme TgUroD in Toxoplasma gondii

reveals a key role of the apicoplast in heme biosynthesis. J Biol Chem *295*,

1539-1550.

Tyra, H.M., Linka, M., Weber, A.P., and Bhattacharya, D. (2007). Host origin of plastid

solute transporters in the first photosynthetic eukaryotes. Genome Biol *8*, R212.

van Dooren, G.G., Tomova, C., Agrawal, S., Humbel, B.M., and Striepen, B. (2008).

Toxoplasma gondii Tic20 is essential for apicoplast protein import. Proc Natl Acad Sci

U S A *105*, 13574-13579.

Vaughan, A.M., O'Neill, M.T., Tarun, A.S., Camargo, N., Phuong, T.M., Aly, A.S.,

Cowman, A.F., and Kappe, S.H. (2009). Type II fatty acid synthesis is essential only

for malaria parasite late liver stage development. Cell Microbiol *11*, 506-520.

Waller, R.F., Keeling, P.J., Donald, R.G., Striepen, B., Handman, E., Lang-Unnasch, N.,

Cowman, A.F., Besra, G.S., Roos, D.S., and McFadden, G.I. (1998). Nuclear-encoded

proteins target to the plastid in Toxoplasma gondii and Plasmodium falciparum. Proc

Natl Acad Sci U S A *95*, 12352-12357.

Walmsley, A.R., Barrett, M.P., Bringaud, F., and Gould, G.W. (1998). Sugar transporters

from bacteria, parasites and mammals: structure-activity relationships. Trends

Biochem Sci *23*, 476-481.

Wan, W., Dong, H., Lai, D.H., Yang, J., He, K., Tang, X., Liu, Q., Hide, G., Zhu, X.Q.,

Sibley, L.D.*, et al.* (2023). The Toxoplasma micropore mediates endocytosis for

selective nutrient salvage from host cell compartments. Nat Commun *14*, 977.

Wang, Q.Q., Sun, M., Tang, T., Lai, D.H., Liu, J., Maity, S., He, K., Wu, X.T., Yang, J., Li,

Y.B.*, et al.* (2023). Functional screening reveals Toxoplasma prenylated proteins

required for endocytic trafficking and rhoptry protein sorting. mBio *14*, e0130923.

Weber, A.P., Linka, M., and Bhattacharya, D. (2006). Single, ancient origin of a plastid

metabolite translocator family in Plantae from an endomembrane-derived ancestor.

Eukaryot Cell *5*, 609-612.

Wilson, R.J., Denny, P.W., Preiser, P.R., Rangachari, K., Roberts, K., Roy, A., Whyte, A.,

Strath, M., Moore, D.J., Moore, P.W.*, et al.* (1996). Complete gene map of the

plastid-like DNA of the malaria parasite Plasmodium falciparum. J Mol Biol *261*,

155-172.

Xia, N., Ye, S., Liang, X., Chen, P., Zhou, Y., Fang, R., Zhao, J., Gupta, N., Yang, S.,

Yuan, J.*, et al.* (2019). Pyruvate Homeostasis as a Determinant of Parasite Growth

and Metabolic Plasticity in Toxoplasma gondii. mBio *10*.

Yeh, E., and DeRisi, J.L. (2011). Chemical rescue of malaria parasites lacking an

apicoplast defines organelle function in blood-stage Plasmodium falciparum. PLoS

Biol *9*, e1001138.

Yesbolatova, A., Saito, Y., Kitamoto, N., Makino-Itou, H., Ajima, R., Nakano, R.,

Nakaoka, H., Fukui, K., Gamo, K., Tominari, Y.*, et al.* (2020). The auxin-inducible

degron 2 technology provides sharp degradation control in yeast, mammalian cells,

and mice. Nat Commun *11*, 5701.

Zhang, M., Wang, C., Otto, T.D., Oberstaller, J., Liao, X., Adapa, S.R., Udenze, K.,

Bronner, I.F., Casandra, D., Mayho, M.*, et al.* (2018). Uncovering the essential genes

of the human malaria parasite Plasmodium falciparum by saturation mutagenesis.
Science *360*.

Zuther, E., Johnson, J.J., Haselkorn, R., McLeod, R., and Gornicki, P. (1999). Growth of
Toxoplasma gondii is inhibited by aryloxyphenoxypropionate herbicides targeting
acetyl-CoA carboxylase. Proc Natl Acad Sci U S A *96*, 13387-13392.

ACKNOWLEDGMENTS

We are grateful to Dr. Sophie Alvarez and Dr. Michael Naldrett at the Proteomics and

Metabolomics Facility, Center for Biotechnology at the University of Nebraska-Lincoln

for the proteomics analysis, Dr. Wen-Chao Wang at the Shanghai Profleader Biotech

Co., Ltd for untargeted metabolomics analysis, and the imaging center at the School

of Life Sciences, Sun Yat-Sun University for assistance with TEM and immuno-EM.

We thank Prof. Vern Carruthers for the kind gift of RH*Δku80Δhxgprt* line, and Prof.

Philippe Bastin for monoclonal antibodies against the epitope tag Ty (BB2).

This research was supported by the National Key Research and Development

Program of China (Grant Nos. 2021YFC2300800 and 2021YFC2300802) to X.Q. Z,

by the National Natural Science Foundation of China (31873009 to S.L. and

31772445 to D.H.L), and by the University Startup Package to S.L..

Author Contributions

Conceptualization, S.L.; Methodology, H.D.; J.Y.; K.H.; W.B.Z.; J.L.; Investigation,

H.D.; J.Y.; K.H.; W.B.Z.; H.Y.D.; R.B.W.; Formal Analysis, S.L.; H.D.; K.H.; J.Y.; D.H.L.

Writing – Original Draft, S.L.; Writing – Review & Editing, S.L.; G.H.; K.M.B.; X.Q.Z.;

D.H.L.;Z.R.L.; Funding Acquisition, X.Q.Z.; S.L.; D.H.L.; Resources, S.L.; D.H.L.;

X.Q.Z.; Supervision, S.L..

DECLARATION OF INTERESTS

The authors declare no competing interests.

REAGENT OR RESOURCE	SOURCE	IDENTIFIER

908   **Materials and Methods**

909   Key resource table listed below

antibodies

Mouse anti-Ty	In house hybridoma (Bastin et al., 1996)	N/A
Rabbit polyclonal anti-HA	ThermoFisher Scientific	Cat#71-5500
Mouse monoclonal anti-HA	BioLegend	Cat#901501
Rabbit polyclonal anti-GAP45	Pmid: 36813769	Polyclonal
Rabbit polyclonal anti-actin	Pmid: 36813769	Polyclonal
Rabbit polyclonal anti-IMC1	Pmid: 36813769	Polyclonal
Rabbit polyclonal anti-ACP	Pmid: 36813769	Polyclonal
Mouse monoclonal anti-GFP	ThermoFisher Scientific	Cat#MA5-15349
Streptavidin Alexa Fluor-488 conjugate	LICOR	Cat#926-3230
Alexa Fluor 488 Goat anti-mouse IgG(H+L)	ThermoFisher Scientific	Cat#A-11029
Alexa Fluor 568 Goat anti-mouse IgG(H+L)	ThermoFisher Scientific	Cat#A-11031
Alexa Fluor 568 Goat anti-rabbit IgG(H+L)	ThermoFisher Scientific	Cat#A-11036
Alexa Fluor 488 Goat anti-rabbit IgG(H+L)	ThermoFisher Scientific	Cat#A-11034
IRDye 800CW Goat anti-mouse IgG (H+L)	LI-COR	Cat#926-32210
IRDye 800CW Goat anti-rabbit IgG (H+L)	LI-COR	Cat#926-32211
IRDye 680CW Goat anti-rabbit IgG (H+L)	LI-COR	Cat#926-68071
IRDye 680CW Goat anti-mouse IgG (H+L)	LI-COR	Cat#926-68070
IRDye 800CW Streptavidin	LI-COR	Cat#926-32230

Chemical, Peptides, and Recombinant Proteins

Pyrimethamine	Sigma-Aldrich	Cat#46706
Mycophenolic acid	Sigma-Aldrich	Cat#M5255
Xanthine	Sigma-Aldrich	Cat#X4002
3-Indoleacetic Acid (auxin)	Sigma-Aldrich	Cat#I2886
D-biotin	Sigma-Aldrich	Cat#B4639
Penicillin-Streptomycin	Solarbio	Cat#P1400
4% paraformaldehyde solution	Solarbio	Car#P1110

Critical Commercial Assays `

pEASY-Basic Seamless Cloning and Assembly Kit	Transgene	CU201-02
Pierce Streptavidin Magnetic Beads	ThermoFisher Scientific	Cat#88816
ProLong Gold Antifade Mountant without DAPI	ThermoFisher Scientific	Cat#36930
ProLong Gold Antifade Mountant with DAPI	ThermoFisher Scientific	Cat#36931

Experimental Models: Cell lines

Human Foreskin Fibroblasts	ATCC	Cat#SCRC-1041
Fetal Bovine Serum	Transgene	FS201-02
Dulbecco's Modified Eagle Medium (DMEM)	ThermoFisher Scientific	Cat#12800-082
Penicillin-streptomycin	Solarbio	P1400
Glutamine	Solarbio	G0200

Experimental Models: Organisms/Strains		
Parasites lines in Table S2	Parental lines: _T. gondii_ RHΔ_hxgprt_Δ_ku80;_ RHΔ_hxgprt_Δ_ku80_/TIR1	
BALB/C female mice (six week old)	Division of Experimental Animals, Beijing University	
Bacterial and Virus Strains		
E. coli Competent Cells (DH10b)	New England Biolab	Cat#C3019I
Recombinant Plasmids		
Plasmids listed in Table S3		
Oligonucleotides		
Primers for PCR listed in Table S4		
Software and Algorithms		
PRISM, version 8.0	GraphPAD	http://www.graphpad.com
InterPro:protein sequence analysis & classification	InterPro	https://www.ebi.ac.uk/interpro/
Toxoplasma Genomics Resource	ToxoDB	http://toxodb.org/toxo/
Eukaryotic Pathogen Genomics Resources	VEuPathDB	https://veupathdb.org/veupathdb
MetaboAnalyst 5.0	MetaboAnalyst	https://www.metaboanalyst.ca
jackhmmer	HMMER v3.3.2	http://hmmer.org/
trimAL v1.2.rev57		https://vicfero.github.io/trimal/
TBtools		https://github.com/CJ-Chen/TBtools
R v4.2.0		https://www.r-project.org/
ComplexHeatmap	R	https://bioconductor.org/packages/release/bioc/html/ComplexHeatmap.html
ChiPlot		https://www.chiplot.online/
Mafft v7.490		https://mafft.cbrc.jp/alignment/software/

Parasite and cell culture

T. gondii RHΔ_hxgprt_Δ_ku80_ (Huynh and Carruthers, 2009), RHΔ_hxgprt_Δ_ku80_/TIR1

(Brown et al., 2017) and their derivative lines are listed in Table S2. The tachyzoite

lines were grown in human foreskin fibroblast (HFF-1 from ATCC® SCRC-1041™) in

D5 media prepared by DMEM supplemented with 5% inactivated fetal bovine serum

(FBS) (Transgene, FS201-02), 10 mM glutamine and penicillin-streptomycin. The

lines and HFF-1 were maintained mycoplasma negative as described previously

(Long et al., 2016). The TIR1 and AID lines were cultured in HFF-1 with 500 µM auxin

(+IAA) or 0.1% ethanol alone (-IAA) for phenotypic assays, as previously described

(Brown et al., 2017; Long et al., 2017b). Parasites were harvested by passing through

22g needles and by filtration through 3.0 micron polycarbonate membranes, from

which extracellular parasites were used for experimental assays.

Mice

T. gondii parasite growth *in vivo* was tested in six-week old BALB/C mice. Mice were

maintained under specific pathogen-free (SPF) conditions in filter-top cages with

provision of sterile water and food. Mice were randomly assigned to experimental

groups (n=5 per group) and challenged by intraperitoneal (i.p.) injection of 100 *T.*

gondii tachyzoites, followed by monitoring on a daily basis to record their health by

checking their appearance, bodyweight and responsiveness. The mouse experiments

were conducted according to the guidelines and regulations issued by the Veterinary

Office of the China Agricultural University (Issue No. AW11402202-2-1).

Generation of Plasmids and *T. gondii* Lines

A CRISPR/Cas9 sgRNA 3' plasmid (Table S3) targeting a region close to a stop codon

at a specific gene can efficiently produce Cas9 and sgRNA to create DNA double

strand breaks (DSB) in parasites, which facilitates the integration of a tagging

amplicon at the 3' terminus of a specific gene. The sgRNA specific sequences were

selected, as described in our previous protocol (Brown et al., 2018). The sequences

were integrated into the pCas9 plasmid using a basic seamless cloning and assembly

kit (CU201-02, TransGen Biotech) with primers listed in Table S4, as described in our

recent study (Wan et al., 2023). The amplicon for a specific integration at the DSB was

generated from a generic tagging plasmid (Table S3) using a pair of primers L and T

(Table S4), as described in our previous study (Long et al., 2017b), which was

illustrated in Fig. S4A. These generic plasmids included pL-6Ty-HXGPRT (Brown and

Sibley, 2018), pL-6HA-DHFR (Long et al., 2017a), pL-6HA-HXGPRT (addgene

#86552) (Long et al., 2017b), pL-TurboID-4Ty-DHFR (Wan et al., 2023),

pL-AID-6Ty-DHFR (Wan et al., 2023) and pL-AID-3HA-HXGPRT (addgene #86553)

(Long et al., 2017b). One additional generic plasmid pL-AID-GFP-HXGPRT was

generated using pL-AID-3HA-HXGPRT as the PCR template of plasmid backbone,

followed by cloning the GFP fragment into the backbone with primers listed in Table

S4. The amplicon generated by primers L and T from the generic plasmids contains

short homologs (HR1 and HR2, 41 bp) for targeting the upstream (HR1) of the stop

codon and the downstream (HR2) of the Cas9 cleavage site (DSB), resulting in

integration of epitope tags or AID fragments in frame with the encoding sequences of

specific genes. The drug-selected lines were sub-cloned, followed with screening with

diagnostic PCR, IFA and western blots. The resistance markers containing LoxP sites

can be excised by transfection with pmini-Cre, as previously described (Heaslip et al.,

2010).

Transfection and Selection

Parasites were harvested and resuspended in Cytomix (10 mM KPO_4, 120 mM KCl, 5

mM $MgCl_2$, 25 mM HEPES, 2 mM EDTA) $1x10^8$ parasites/ml, then combined with

10-20 µg pCas9-sgRNA 3' and 1-5 µg amplicons with 2 mM ATP, 2 mM GSH and 150

µM $CaCl_2$ to a final volume of 250 µl, followed by electroporation using a BTX ECM

830 electroporator (Harvard Apparatus), as described previously (Soldati and

Boothroyd, 1993). Parasites were selected on the second day with addition of

appropriate drugs mycophenolic acid (MPA) (25 µg/ml) and 6-xanthine (6Xa) (50

µg/ml), or pyrimethamine (Pyr) (3 µM). Negative selection was applied with

6-thio-xanthine (200 µg/ml) when parasites were transfected with a pCre plasmid to

remove the resistant marker HXGPRT. Positive selection pools of parasites were

subcloned in 96-well plates at 3 parasites/well and allowed to grow for 6-7 days before

selection of wells containing a single plaque. Pure clones were diagnosed by PCR

using appropriate primers (Table S4) and IFA analysis using appropriate antibodies if

applicable.

Indirect Fluorescence Assay (IFA)

Parasites were fixed by 4% paraformaldehyde in PBS, followed by blocking and

permeabilization using PBS containing 2.5% BSA and 0.25% Triton X100. Antibody

incubation was performed with indicated combinations of primary antibodies for 30

min, followed by incubation with secondary antibodies conjugated with fluorescent

dyes (Thermofisher Scientific) for 30 min. The above procedures were applied to the

ACP diffusion analysis for parasites as stated in the results and legends. The blind

counting was conducted for at least 150 vacuoles (intracellular parasites) or single

parasites (extracellular parasites) in each replicate. ProLong Gold Antifade Mountant

with or without DAPI were used to mount the coverslips on glass slides. Parasites

were then imaged using a Nikon Ni-E microscope C2+ equipped with a DS-Ri2

Microscope Camera. The co-localization analyses were performed using the Ni-E

software system NIS Elements AR. All the IFA analyses were carefully repeated at

least three times, and similar results were obtained for the same experiments before

finalizing the analysis.

Western blotting

Parasites were resuspended in PBS, followed by addition of 5x Laemmli sample

buffer supplemented with 1 mM DTT for a further incubation at 98 °C for 10 min.

Protein lysis was then resolved in SDS-PAGE, blotted by a BioRad wet-blotting

system, subsequently incubated with different combinations of primary and secondary

antibodies, followed by secondary antibodies conjugated with LI-COR reagents, or

streptavidin LI-COR 800CW. The blot membranes were then visualized using a

Bio-RAD ChemiDOC MP system.

TurboID and Biotinylated Proteins

The APT1-TurboID line was grown on HFF monolayers for 24 hours, subsequently

incubated with 500 μM D-biotin for 90 min. The parasites were fixed with 4%

paraformaldehyde solution for IFA or harvested for western blots. IFA and western

blots were performed with appropriate streptavidin reagents to visualize biotinylated

proteins in parasites. The parental line and APT1-TurboID line were grown in HFF

cells for 36 hours, followed with labeling of proteins by 500 μM D-biotin for 90 min, and

harvesting of the parasites for purification of biotinylated proteins. In parallel, the

APT1-TurboID parasites after 36 hours growing were harvested, and extracellular

parasites were incubated with 500 μM D-biotin for 90 min, followed with by collection

of parasites for purification of biotinylated proteins. The parasites were lysed in a

buffer containing 1% Triton X-100, 0.2% SDS and 0.5% deoxycholate, and sonicated

with a microtip in 550 sonic dismembrator (ThermoFisher Scientific). Biotinylated

proteins in cleared supernatant were purified using streptavidin magnetic beads,

exactly following the protocol described elsewhere (Long et al., 2018).

Proteomic analysis

The purified proteins were separated on SDS-PAGE and stained by Coomassie blue

R250, 45% methanol and 10% glacial acetic acid, subsequently dried on SDS-PAGE

slices by vacuum. The dried gels were rehydrated, alkylated and washed to remove

the stain and SDS, and subjected to steps of mass-spectrometry analysis, following

the protocol described in the Bio-Protocol (Long et al., 2018). The parental line and

TurboID fusion line were analyzed in parallel with three technical and biological

replicates. Resulting spectra were searched against a combined database of *T. gondii*

ME49 (http://ToxoDB.org, release 29) and human proteins and a decoy database,

using Mascot and Scaffold. The current views in Scaffold were exported into an excel

spreadsheet using the following settings of peptide 2, protein threshold 99% and

peptide threshold 95%. Hits were analyzed using embedded tools in ToxoDB

(https://toxodb.org/toxo/app/) to predict the transmembrane domain. Peptide numbers

of the hits in the TurboID were compared to those in the parental line (0 peptide was

considered as 1), and fold changes were used to plot a volcano map against $\mathrm{Log10}^{p}$

^{value} in ggplot2 package in R.

Conservation and phylogenetic analysis

The apicoplast proteins identified in this study and known apicoplast proteins as listed

in the results (Table S10) were used as queries for searching orthologues in

organisms in Eukaryote, including species belonging to the chromalveolates, plants,

metazoan and fungi, and Bacteria (Table S9). These proteins were simultaneously

searched with the profile Hidden Markov models (HMMs) search engine

(jackhammer). The E-value cutoff was set at 1e-7 and hits with an E-value < 1e-7

were considered as orthologues. The best hits from the representative species in

above taxa were merged and resultant list of orthologues (Table S10) were used for

conservation analysis with ComplexHeatmap package in R (v4.2.0) (Gu et al., 2016).

This analysis pulled up the protein conservation heatmap, with indication by a heat

scaled with E-value for each individual orthologue. The tree dendrogram without

meaningful branch lengths across these species, and across the proteins were

clustered according to the Euclidean distance using UPGMA method.

Representatives from the known apicoplast proteins were analyzed in parallel as the

control for comparison.

Plaque Formation

Parasites (150 parasites) were inoculated on confluent HFF monolayers in one 6-well

in plates, followed with addition of either 500 µM IAA or ethanol alone (0.1%) and

growth in D5 media at 37°C for 7 days. The parasites and monolayers were then fixed

with 70% ethanol for 15 min, stained with 0.5% crystal violet for 5 min, subsequently

washed with dH2O and dried at room temperature. The plaque monolayers were

recorded by scanning using a HP Scanjet G4050, and plaque numbers and plaque

sizes were measured in ImageJ.

Parasite Replication

The TIR1 and its derived AID lines were grown in 500 μM IAA for 24 hours on

monolayers in 24-well plates with coverslips. The parasites were fixed with 4%

paraformaldehyde for 10 min, followed with blocking and permeabilization with 0.25%

Triton X-100 in PBS containing 2.5% BSA. The parasites were then incubated with

GAP45 antibodies and secondary anti-rabbit antibodies conjugated with Alexa Fluor

568. The coverslips were mounted and visualized under a Nikon Ni-E microscope C+.

Vacuoles containing different numbers of parasites were scored on the same images

with blindness, and at least 150 vacuoles were counted in each replicate of three

independent experiments. Ratios of vacuoles containing different numbers of

parasites were plotted against the total vacuoles examined.

GFP acquisition assay

HFF cells expressing cytosolic GFP were prepared in our previous study (Wan et al.,

2023). In brief, pLVX-EF1a-GFP-IRES-puromycin was combined with packaging

plasmids psPAX2 (addgene #12260) and pMD2.G (addgene #12259) in Opti-MEM I

(31985062, Gibco), and mixed with Fugene HD (E2311, Promega), and transfected

into 293T cells. Viruses were collected and mixed with polybrene in a 12-well plate

well containing 30-50% confluent HFF-1 cell monolayers. After 24 hours of

post-transduction, the cell monolayers were re-seeded with incubation of puromycin

(2.5 μg/ml) for 3 days. The cell monolayers were then maintained and expanded in D5

media. The TIR1 and AID lines were grown in the cell monolayers with IAA, 10 μM

LHVS and 12.5 μM atorvastatin for 12, 24 or 36 hours. LHVS inhibits cysteine

protease CPL to accumulate GFP derived from host cell cytosol, as reported

previously (Dou et al., 2014), Atorvastatin can inhibit synthesis of FPP and GGPP in

host cells, to decrease the supply of FPP and GGPP from the host cells to the

parasites. In parallel, TIR1 was grown in the host cells with 10 μM LHVS, 0.3 μM

zoledronic acid and 12.5 μM atorvastatin, serving as a positive control to observe

defects in GFP transport by decreasing protein prenylation in parasites. The addition

of atorvastatin and zoledronic acid could specifically inhibit the FPP and GGPP supply

both in host cells and parasites (Li et al., 2013), thus inhibiting GFP transport to the

VAC in parasites, as demonstrated in Fig. 5A-B. Purified parasites were adhered onto

poly-lysine-coated coverslips, followed with fixation with 4% paraformaldehyde for 10

min. The parasites were permeabilized by 0.25% Triton X-100 in PBS, subsequently

mounted in ProLong Gold Antifade Mountant with DAPI for visualization under a Nikon

Ni-E microscope C+. Parasites were judged by observations of parasites with either

GFP foci, GFP diffusion or GFP absence, as demonstrated in Figure 4B.

Fields/images were selected blind, and all parasites in the same images were counted

for scoring of parasites with or without GFP signal. The counting was performed in

replicate for at least 150 parasites. Percentages of parasites containing GFP foci were

calculated and expressed as percentages in the populations.

Immuno-staining for Electron Microscopy (Immuno-EM)

Parasites were grown on HFF monolayers for 24 hours, subsequently digested with

trypsin for collecting samples. Samples were fixed in a 0.1 M sodium cacodylate

buffer (pH7.4) containing 3% (w/v) paraformaldehyde, 0.1% glutaraldehyde (v/v), 4%

sucrose (w/v) at 4°C overnight. The samples were then washed four times by a 0.1 M

sodium cacodylate buffer containing 4% (w/v) sucrose and neutralized on ice by a 0.1

M sodium cacodylate buffer containing 4% (w/v) sucrose, 0.1 M glycine, and

dehydrated in ethanol. The samples were immersed in LR White resin (Ted Pella, Inc.)

at 4°C overnight, subsequently polymerised at light intensity 1.2×10^5 μJ/cm^2 on ice for

5 hours. The LR White imbedded samples were sectioned with a Leica

ultramicrotome (EM UC7) at 100 nm and were picked onto 100-mesh

formvar-carbon-coated nickel grids. The loaded nickel grids were then blocked and

incubated with mouse anti-Ty antibodies (1:5) at RT for 2 hours and overnight at 4°C,

followed with washing by PBS. The samples on grids were incubated with Alexa

FluorTM 488 goat anti-mouse IgG 10 nm colloidal gold particle (1:100) (15nm or 10

nm in size) for 1 hours, followed with washes in PBS and double-distilled water. The

samples on grids were stained by 3% phosphotungstic acid and observed under a

JEM1400FLASH transmission electron microscope operation at 120 KV.

Transmission Electron Microscopy (TEM)

Parasites grown on HFF cell monolayers were collected by scraping off, fixed by a

fixative mixture of 2% paraformaldehyde and 3% glutaraldehyde at 4 °C overnight.

The samples were washed, fixed in 1% osmium tetroxide in 50 mM phosphate buffer

at 4 °C for 1 hour, washed three times and dehydrated by sequential concentration of

ethanol, and by acetone and Spurr for 1 hour. The samples were then imbedded in

Spurr and sectioned to 100 nm slices by a Leica ultramicrotome (EM UC7). The

ultrathin-sections were collected on 100 mesh grids. The grids were then viewed by

JEOL transmission electron microscopy (JEM1400FLASH).

Untargeted Metabolomics

Parasites were grown in IAA for 18 hours and purified in chilled PBS, followed by

immediate quenching by addition of -40°C methanol and sterile ddH2O (400 µl

methanol + 100 µl H_2O). The samples were processed for the analysis of metabolites

by GC-MS and LC-MS, exactly following the protocol reported in our recent study

(Wan et al., 2023). Briefly, the GC-MS analysis was performed with An OPTIMA 5 MS

Accent fused-silica capillary column on Agilent7890A/5975C GC-MS system (Agilent

Technologies Inc., CA, USA), while the LC-MS analysis was performed on a

ThermoFisher Ultimate 3000 UHPLC system with a Waters ACQUITY UPLC BEH

C18 column (2.1mm × 100 mm, 1.7µm) using standard positive and negative modes.

For GC-MS, the peak picking, alignment, deconvolution and further processing of

resulting raw data were carried out according to the previously published protocols

(Gao et al., 2010). The final data was exported as a peak table file, including

observations, variables (rt_mz), and peak areas. The data was normalized against the

total peak value of total peaks, and differential metabolites were identified by

statistical analysis performed in R platform, where parametric and nonparametric

tests were done using One-way ANOVA with Dunnett's multiple comparison, to

identify differential metabolites with $p<0.05$ and fold change >1.2 for further analysis.

For LC-MS, The raw data was transformed to mzXML format by ProteoWizard and

then processed by XCMS and CAMERA packages in R software platform. The final

data was exported as a peak table file, including sample names, variables (rt_mz) and

peak areas. The peak areas data was normalized to internal standards before the

analysis of statistics. The in-house database was referenced to identify the compound

by using m/z (MS1), mass spectra (MS2) and retention times (RT). Normalization for

relative parasite amounts was based on the total integrated peak area values of

metabolites within an experimental batch. Metabolites identified by the GCMS and

LCMS were merged together for further analysis (Dataset S2). Enrichment of

Metabolic Pathways (Table S13) were analyzed using the differential metabolites

identified in three of the AID lines versus the TIR1 line. MetaboAnalyst 5.0 server was

used to enrich the metabolite sets with KEGG, in which the p-value below 0.05 was

considered to be significant.

Parasite growth in mice

Mice were randomly assigned to groups and were intraperitoneally injected with 100

parasites per mouse. Subsequently, mice (n=5 mice for each line) were orally

administered with IAA (3-indoleacetic acid, auxin, Sigma) and by an intraperitoneal

injection on a daily basis (Brown and Sibley, 2018; Yesbolatova et al., 2020). The

drinking water contained sterile water with 1 mg/ml IAA, 3 mM NaOH, 5% sucrose

(w/v), flavored with 2 mg/ml TANG (Mondelēz International), pH8.0. Meanwhile, mice

were intraperitoneally injected with a daily 0.2 ml of sterile solution containing 15

mg/ml IAA, 1 M NaOH, pH7.8. The mice infected with the TIR1 line served as the

control, which received the drinking water and injection of IAA-containing solution in

the same way to the AID lines. The mice were weighed and their health conditions

were monitored on a daily basis until day 20.

Quantification and statistical analysis

Experiments were conducted blind for at least two-three biological replicates and

statistical analyses were conducted in Graphpad v8.0. One-way or two-way ANOVA

with Tukey's multiple comparison was conducted for data that fit normal a distribution

or the Mann-Whitney test for non-parametric comparison, while one-way ANOVA with

Dunnett's multiple comparison was tested for data with small sizes or abnormal

distributions. $p<0.05$ was considered significant. Experiment-specific statistical

information is provided in the figure legends or associated method details including

replicates (n), trials (N), and statistical tests performed.

Data availability

The proteomic data reported in this paper have been deposited in the OMIX, China

National Center for Bioinformation/Beijing Institute of Genomics, Chinese Academy of

Sciences (https://ngdc.cncb.ac.cn/omix) with accession number: OMIX002059. The

metabolomics raw data have been deposited in the MetaboLights

(https://www.ebi.ac.uk/metabolights/) with accession number: MTBLS7060. Minimally

processed data of the proteomics and metabolomics data are available in Dataset S1

and Dataset S2, respectively. *Toxoplasma gondii* genome information can be found in

ToxoDB release 53 (http://toxodb.org) and Eukaryotic pathogen, Vector & Host

Information Resources can be found in VEupathDB (http://veupathdb.org).

1181

1182

1183

1184

1185

1186

1187 **Figure Legends**

1188

Figure 1. Discovery of apicoplast membrane proteins

(A-B) Activity test of biotinylation of proximal proteins by APT1-TurboID-4Ty. Parasites

were grown in 500 µM D-biotin for 90 min, followed by detection of biotinylated

proteins (green) by streptavidin reagents on IFA (A), and on Western blots (B). Actin

served as the loading control. Scale = 5 µm.

(C) Volcano plot analysis comparing the TurboID fusion to the parental line. Three

replicate mass-spectrometry experiments were analyzed by the student t-test. Known

apicoplast proteins (red) and candidates with different numbers of TMDs (deep and

light blue were indicated, and novel apicoplast proteins were pointed out. See Dataset

S1.

(D) Workflow of the discovery of novel apicoplast membrane proteins. Candidates

from hyperLOPIT (Barylyuk et al., 2020), ACP-BirA (Boucher et al., 2018) and

APT1-TurboID (this study) were screened by epitopte tagging and IFA.

(E) Confocal co-localization of HAP-6Ty with the apicoplast marker ACP, showing

confocal imaging of HAP-6Ty (red) and ACP (green), and Pearson Corelation

Coefficiency (PCC). The PCC values over the merged fluoresent foci for candidate

proteins and ACP were analyzed by a co-localization and ROI intensity analysis

module in the NIS Elements AR software and shown with mean ± SEM (N=6 for each).

Scale = 5 µm.

(F-H) Summary and comparison of the screening from three datasets. The putative

transporters are grouped into diverse types (F) and identified from three different

datasets (G), which shared three novel proteins, as shown by the venn diagram (H).

Three independent experiments were performed with similar outcomes (A, B, E).

1212

Figure 2. Conservation analysis of putative transporters

(A-B) Heatmap indicating conservation of known apicoplast proteins and newly

identified proteins among chromalveolates (A), red alga, green alga, plants, fungi,

metazoa and bacteria (B). The conservation was scaled by a heat bar (red) with

E-values for putative orthologues of the listed proteins (see Dataset S2). ToxoDB

protein numbers (left) and protein (temporary) names (right) are shown. Their

sequences were searched for orthologues using the Hidden Markov Model (HMM) (E

value<E-7) and the best orthologue was used for the heatmap analysis in

ComplexHeatmap package in R. Parasite fitness scores were shown on the right with

the green heat bar for *T. gondii* (ToxoDB)(Sidik et al., 2016) and the blue bar for *P.*

falciparum (PlasmoDB)(Zhang et al., 2018). The tree dendrogram across the genes

were clustered according to the euclidean distance using UPGMA method. The

species tree (top) shows phylogenetic relationships and major clades: *Theil.*, *Theileria*;

Babes, *Babesia*; *Crypt.*, *Cryptosporidium*; *Proteo*, *Proteobacteria*.

1227

Figure 3. AMT1 and AMT2 are localized at the apicoplast membrane and essential for

parasite growth *in vitro*

(A) Western blot detection of protein depletion in the AID lines. The lines were induced

by auxin (IAA) for different hours (hrs) as indicated. Western blots detected immatured

(p) and matured (m) forms of the protein fusions in the non-induced lanes. Actin

served as the control.

(B) IFA detection of protein depletion in the AID lines. Parasites were grown in IAA for

24 hours, followed by IFA with antibodies against Ty or HA (red) and IMC1 (green).

Scale = 5 μm.

(C-E) Plaque formation by the TIR1 and AID lines on HFF monolayers in ± IAA for 7

days. Numbers (D) and sizes (E) of the plaques were measured by ImageJ. Scale =

0.5 cm. Two independent experiments with triplicates were performed. Data are

shown as a mean ± SEM.

(F) Parasite replication of the TIR1 and AID lines grown in IAA. The parasites were

grown in IAA for 24 hours, followed by scraping, harvesting and infection for the 2nd

and 3rd rounds of parasite growth in IAA. The parasites stained with GFP45 were

counted in vacuoles (at least 150 vacuoles in each replicate). In comparing to TIR1

(1st, 2nd and 3rd), $p<0.0001$ for the AID lines with 2 and 8 parasites/vacuole in the 1st

round, and for the AID lines with 1-8 parasites/vacuole in the 2nd round, $p<0.0001$ for

the AID lines with 1 and 4 parasites/vacuoles in the 3rd round.

(G-H) ACP diffusion in the TIR1 and AID lines grown in IAA. Parasites were grown in

IAA for the 1st, 2nd and 3rd lytic cycles, as described in the parasite replication assay

(intracellular parasites). Those parasites in the 1st and 2nd round of growth were forced

to egress for the same analysis (extracellular parasites). Vacuoles or single parasites

were scored (n>150 for each replicate).

Fields/images were selected blind and all parasites/vacuoles were scored on the

same fields/images (F-H). Three independent experiments were performed (A, B, F, G

and H), data are shown as a mean ± SEM with two-way ANOVA with Tukey's multiple

comparisons (compared with the TIR1). ***, $p<0.0001$.

1257

Figure 4. Depletion of AMT1 and AMT2 caused defects of MEP and FASII pathways

(A-C) The MEP activity was strongly affected in the AID lines. GFP transport to the

VAC was assayed by growning parasties in GFP expressing HFF cells (HFF-GFP).

LHVS (10 µM) was added to inhibit GFP digestion in the VAC (Dou et al., 2014). The

isoprenoid synthesis in host cells and parasites was inhibited by IC50 concentrations

of atorvastatin (10 µM) and zoledronate (0.3 µM)(Li et al., 2013), respectively (A). The

inhibition over the TIR1 line using two of the drugs was used as a positive control (B).

The AID parasites were grown in HFF-GFP cells in the presence of LHVS for 36 hours

in total, with the addition of atorvastatin, zoledronate or IAA as stated for different

experiments in the figure in the last 12 hours. Parasites were immediately harvested,

adhered and scored, as shown in the example images in B. Percentages of parasites

with GFP foci were plotted (C). Fields/images were selected blind and all

parasites/vacuoles were scored on the same fields/images (C). Three independent

experiments with triplicates were performed. Data are shown as a mean ± SEM with

one-way ANOVA with Tukey's multiple comparison. ***, $p<0.0001$, scale = 5 µm.

(D-F) Fatty acid synthesis and TCA metabolites were significantly reduced in IAA

induced AID parasites. Parasites were grown in IAA for 18 hours and harvested for

untargeted metabolomics with GCMS and LCMS (N=3), and metabolites detected

were analyzed by Dunnette's multiple comparison by comparing the AID parasites to

the TIR1. Three AID lines shared 13 differential metabolites (D). These shared

metabolites included fatty acids and lipids, such as myristic acid, palmitoleic acid,

oleic acid, and phosphatidylcholine (PC) (E), and metabolites related to the TCA cycle

(F). *, $p<0.05$; **, $p<0.001$; ***, $p<0.0001$.

1281

Figure 5. Depletion of AMT2 resulted in ACC1 instability.

(A-B) No reciprocal effect on protein levels by depletion of AMT1 or AMT2. AMT1 and

AMT2 were reciprocally tagged at C-terminus with 6HA in the AID lines, and the HA

fusions were detected by Western blots in parasites in ±IAA for 36 hours (A), and by

IFA in parasites in ±IAA for 24 hours (B). IFA was performed with IMC1 (green) and

Ty/HA (red).

(C-D) ACC1 was lost upon depletion of AMT2-AID. ACC1 was endogenously tagged

with 6HA at its C-terminus in the AMT1-AID and AMT2-AID lines, and parasites were

grown in IAA for 36 hours for Western blots (C) and IFA (D). IFA analyses were

performed with anti-ACP and anti-HA antibodies, while Western blots were done with

anti-HA and streptavidin Li-COR 800CW. Actin served as the control.

(E-F) Representative images of ACC1 on IFA in parasites of AMT2-AID induced in IAA

for 24 hours. ACC1 (red) was completely lost in parasites of a vacuole (type 1),

remained only in one parasite of a single vacuole (type 2), or in two parasites of a

single vacuole (type 3) (E). The ACC1 types were scored in AMT2-AID induced in IAA

for 24 hours, and expressed as percentages of different types (F). At least 150

parasites were scored in each replicate. Three experiments with triplicates were

performed, and data are shown as a mean ± SEM and analyzed by two-way ANOVA

with multiple comparison, ***, $p<0.0001$.

(G) The remaining ACC1 was still biotinylated. Parasites were grown in ±IAA for 24

hours for IFA analysis with antibodies against HA and streptavidin Alexa Fluor 488.

Parasites containing both ACC1-6HA (red) and biotinylated ACC1 (green) was

observed.

Three independent experiments were performed with similar outcomes, and

representative images were shown. Scale = 5 µm.

1307

Figure 6. Examination of the apicoplast ultra-structures, and mouse virulence of the

parasites upon depletion of the AID fusions.

(A) Transmission electron microscopy (TEM) observed less electron dense or electron

lucent lumen and loosened membranes in the apicoplast in the AID parasites induced

in IAA for 24 hours. Representative images were shown. Scale = 200 nm.

(B) Quantification of the apicoplast with the lumen less dense or lucent in parasites

induced for 12, 24 and 36 hours and TIR in IAA for 36 hours. The TEM examined 300

parasites for three lines (TIR1, AMT1-AID and AMT2-AID) at each time point, and

numbers of abnormal apicoplast (red numbers) and apicoplast identified (green

numbers) were shown by the columns. The columns express the percentages of

abnormal apicoplast in all the identified apicoplasts.

(C) Survival curve of BALB/C mice infected with 100 parasites intraperitoneally and

treated with IAA for 20 days. The Gehan-Breslow-Wilcoxon test was used to compare

differences between the survival curves (the AID line versus TIR1), ***, $p<0.0001$. The

experiment of mouse infection was performed with 5 mice in each group, and results

from three independent experiments (1st, 2nd and 3rd) were shown together with

symbols for the mouse groups.

A

TIR1　　　　AMT1-AID　　　　AMT2-AID

B

abnormal api (%)

TIR1	2/42
AMT1-AID 12	10/43
24	10/20
36	10/20
AM2-AID 12	6/11
24	8/12
36 hr	3/3

C

- TIR1 +AA
- AMT1-AID
- AMT2-AID
- AMT1-AID/AMT2-AID

Percentage of Survival

1st
0.0116
***
***

2nd
0.0069
***
***

3rd
0.0019
***
***

Days Post Infection