1   **Article**
2
3   **Intraspecific genetic variation is critical to robust toxicological predictions of aquatic**
4   **contaminants**
5

6   René S. Shahmohamadloo*[a,b], Seth M. Rudman*[a,b], Catherine I. Clare[a], Judy A. Westrick[c],
7   Xueqi Wang[b], Luc De Meester[d], John M. Fryxell[b]
8
[a] School of Biological Sciences, Washington State University, Vancouver, Washington, 98686,
United States
[b] Department of Integrative Biology, University of Guelph, Guelph, Ontario, N1G 2W1, Canada
[c] Department of Chemistry, Wayne State University, Detroit, Michigan, 48202, United States
[d] Laboratory of Aquatic Ecology, Evolution, and Conservation, University of Leuven, Charles
Deberiotstraat 32, 3000 Leuven, Belgium
15
16   *These authors contributed equally to this work.
17
Address co-correspondence to:
René S. Shahmohamadloo, PhD
Liber Ero Postdoctoral Fellow, NSERC Postdoctoral Fellow
School of Biological Sciences | Washington State University
14204 NE Salmon Creek Avenue | Vancouver | Washington | 98686 | United States
Email: rene.shahmohamadloo@wsu.edu
24
Address co-correspondence to:
Seth M. Rudman, PhD
Assistant Professor
School of Biological Sciences | Washington State University
14204 NE Salmon Creek Avenue | Vancouver | Washington | 98686 | United States
Email: seth.rudman@wsu.edu
31

32   **Author Contributions:** R.S.S., S.M.R., L.D.M., and J.M.F. designed the research. R.S.S. and
33   S.M.R. conducted the experiments. J.A.W. and X.W. conducted supporting data collections.
34   S.M.R., R.S.S., and C.I.C. analyzed the data. R.S.S. and S.M.R. wrote the paper with insights
35   from L.D.M., C.I.C., and J.M.F. All authors read, amended, and approved the final manuscript.
36
37   **Competing Interest Statement:** The authors declare no competing interests.
38
39   **Data Availability Statement:** The primary dataset that support the findings of this study are
40   included in the Supporting Information.
41
42   **Keywords:** *Risk assessment*, *Genetic variation*, *Toxicology, Daphnia, Contaminants monitoring*
43

44   **This file includes:** Main Text, Figure 1 to 3
45

Abstract

Environmental risk assessment is a critical tool for protecting aquatic life and its effectiveness is predicated on predicting how natural populations respond to contaminants. Yet, routine toxicity testing typically examines only one genotype, which may render risk assessments inaccurate as populations are most often composed of genetically distinct individuals. To determine the importance of intraspecific variation in the translation of toxicity testing to populations, we quantified the magnitude of genetic variation within 20 *Daphnia magna* clones derived from one lake using whole genome sequencing and phenotypic assays. We repeated these assays across two exposure levels of microcystins, a cosmopolitan and lethal aquatic contaminant produced by harmful algal blooms. We found considerable intraspecific genetic variation in survival, growth, and reproduction, which was amplified by microcystins exposure. Finally, using simulations we demonstrate that the common practice of employing a single genotype to calculate toxicity tolerance failed to produce an estimate within the 95% confidence interval over half of the time. These results illuminate the importance of incorporating intraspecific genetic variation into toxicity testing to reliably predict how natural populations will respond to aquatic contaminants.

Main Text

Reliable translation of aquatic toxicity testing into predictions of population-level consequences in nature is a cornerstone of environmental risk assessments [1–6]. Such predictions are classically obtained by quantifying individual phenotypic responses under constant conditions in a laboratory [7]. Research has historically favored single-species toxicity testing that generates point estimates such as the lethal concentration at which a contaminant will kill 50% of a test population (LC50) [8]. For more than fifty years, regulatory agencies have relied on such lethality data to inform environmental risk assessments, mandated to safeguard biodiversity and associated ecosystem services by protecting rare or economically significant species [6]. Yet, reliably translating laboratory toxicity studies into population-level predictions requires building on traditional methods to incorporate aspects that are key to ecological realism [8].

Genetic variation within species is recognized as a critical driver of both ecological and evolutionary processes [9–11]. In ecological contexts, genetic variation begets variation in functional traits, which can have profound impacts on populations, communities, and ecosystems [9–12]. Genetic variation within species is central to evolution, as the process of natural selection shaping genetic variation is what produces adaptation [10,13]. Among the most visible and societally challenging examples of rapid adaptation are cases where populations adapt to agrochemicals [14] or contaminants [15]. Despite an appreciation of the role of adaptation in resistance and some demonstrations of the amount of standing variation in tolerance [4,5,16–18], recognition of the importance of intraspecific genetic variation continues to be ignored by regulators within the discipline of toxicology and, more broadly, the environmental risk assessment process.

Investigating the magnitude of intraspecific variation within an aquatic population represents a crucial and understudied element in toxicology and more broadly environmental monitoring programs. Little is known about the magnitude and ecological importance of intraspecific genetic variation in toxicological responses within natural populations. Choosing a genotype based on availability (e.g., standard lab strain) may lead to inferences that are unrepresentative of imperiled populations. More broadly, conducting measurements on any single genotype is using a single point to estimate a distribution that likely has substantial variance [19,20]. Detailed study of specific populations that includes both measures of the extent of genetic variation within focal

97 populations and classical toxicity testing are needed to infer the potential for this variation to
98 impact population-level responses under toxic insults [21].
99
Daphnia magna (water fleas) are a powerful eco-genomic model system because they are
simple to culture, have a fully-sequenced genome, and they play an important and well-
documented ecological role in many freshwater ecosystems [22,23]. With short generation times
and considerable genetic diversity derived from diapausing eggs in sediments, *Daphnia* can
undergo rapid adaptation in response to environmental variation [22–26]. These eco-evolutionary
insights have not translated to change in routine laboratory testing of *Daphnia*, which is a model
system used in toxicology to test chemical compounds before they are produced, released or
sold [23]. Toxicologists previously advocated that toxicological studies should account for genetic
variation in *Daphnia* based on discrepancies between several laboratory strains that were
exposed to the same reference toxicants [2,3,7,27]. Yet, determining the loss in accuracy associated
with choosing a single clone from a natural and genetically heterogenous population for
environmental risks assessments is crucial to demonstrating the overall importance of
accounting for genetic variation.
113
Determining the magnitude of variation within a population is crucial both for predicting
immediate toxicological responses and as an indicator of the potential for rapid adaptation.
Doing so requires accounting for the scale and scope of genetic diversity within a population
and conducting toxicity testing on a heterogeneous and realistic population of *Daphnia*.
Genomic data can be useful in meeting these objectives as genome-wide data provide
information on the scale and scope of genetic diversity being studied and can be translated to
provide population-level insight. For investigations on the relationship between genetic diversity
and toxicological responses whole genome sequencing data yields an ability to identify
populations or clones and compare the extent to which they are genetically divergent, to
determine whether the degree of genomic differentiation between populations is correlated with
toxicological responses, and to test for relationships between genomic diversity and tolerance.
Depending on the underlying genetic architecture of variation in toxin tolerance it may be
possible to identify the genetic basis of resistance [28], though finding a 'simple' genetic basis may
not be common [29]. The use of candidate loci, identified through prior mapping efforts or through
changes in gene expression, provides an opportunity to test hypotheses about the relationship
between genetic variation in potentially important genes and toxicological responses that could
link genetic and phenotypic variation [30–32]. At a minimum, incorporating population genetic
information can help standardized toxicity tests and the information generated also has the
potential to better predict the risk of toxicant exposure on populations, and increase
predictability on the associated community and ecosystem functions.
134
To determine the importance of intraspecific variation in toxicity projections key to the
environmental risk assessment process, we sequenced the genomes of 20 *Daphnia magna*
clones collected from a single Belgian lake and quantified the chronic effects of the
cyanobacterium *Microcystis* on their life-history traits (survival, growth, reproduction, broods).
Harmful algal blooms (HABs) of *Microcystis* are a prominent and toxic ecological disturbance [33]
that strongly impact a wide-range of aquatic taxa due to the production of microcystin toxins [34–
39]. As a keystone zooplankton grazer, *Daphnia* shows substantial intraspecific diversity and
rapid adaptation in response to HABs [24,40,41], including some evidence that particular genotypes
can effectively feed on and reduce the abundance of toxic *Microcystis* [42,43]. Determining whether
intraspecific variation in toxin tolerance influences vital demographic rates and population
dynamics by *Daphnia* clarifies the role of intraspecific genetic variation in toxicology, with
implications for regulatory agencies tasked with maintaining biodiversity under toxic bloom
conditions.

148
In this study, 30 individuals per genotype were divided and exposed to three treatments (i.e.,
common gardens): 100% green algae (*Chlorella vulgaris*, non-toxic diet), 2:1
chlorella:microcystis (a moderately toxic diet), and 1:1 chlorella:microcystis (a severely toxic
diet). Common garden rearing keeps environmental conditions constant allowing for the
estimation of the contribution of genetic variation to phenotypic variation. We chose 2:1 and 1:1
toxic treatments that resulted in microcystin concentrations of 3.3 ± 0.001 μg L^{-1} and 5.1 ± 0.001
μg L^{-1} (Dataset S1), respectively, which have been considered sublethal [44] to lethal [45,46] in
Daphnia laboratory assays and are common ratios in freshwater ecosystems with HABs [36]. We
sought to address the following four questions: 1) How much intraspecific genomic and
phenotypic variation occurs within a single population of *D. magna*? 2) Does exposure to toxic
cyanobacteria reduce the magnitude of intraspecific phenotypic variation? 3) Is there an
interaction between clonal growth and reproduction under favorable (green algae) and toxic
(cyanobacteria) conditions? 4) Using data from these trials, how would sampling design
influence the interpretation of reproductive output with and without contaminant exposure? We
predicted that *D. magna* clones would exhibit considerable genetic and phenotypic intraspecific
variation in toxicological responses, as *Daphnia* populations have been shown to be variable.
Additionally, we predicted that phenotypic variation would be reduced when exposed to a toxic
insult due to a decrease in trait values in high performing genotypes. Moreover, we predicted a
trade-off between growth and reproduction under favorable and toxic conditions, with clones
having the greatest reproductive output varying across cyanobacterial concentrations. Finally,
we predicted that sampling a sufficient number of genotypes would be critical to producing an
accurate estimate of reproduction in both favorable and toxic conditions.
171
172   *Intraspecific genomic and phenotypic variation*
173
Genetic variation among *D. magna* clones was considerable in both phenotype and genotype
(Figure 1). When reared on a *Chlorella*-only non-toxic diet, phenotypic variation among clones
was significant for survival ($X^2 = 4.52$, $P = 0.033$), growth ($X^2 = 67.40$, $P < 0.0001$), and
reproduction ($X^2 = 40.22$, $P < 0.0001$). The magnitude of genome-wide divergence between
pairwise sets of clones varied (Figure 1a), with the most similar clones having ~1.2M SNP
differences and the most divergent ~1.6M.
180
We tested for an association between the degree of genomic divergence between clones and
the magnitude of their phenotypic differentiation. We calculated a phenotypic distance between
each pair of clones based on results of phenotypic assays for survival, growth, time to
reproduction, and total reproduction when reared on non-toxic, moderately toxic, and severely
toxic diets. We found no significant association between the degree of genomic divergence
genome-wide and the magnitude of overall phenotypic divergence between clones (Figure 1)
(mantel $r = 0.033$, $P = 0.781$). We also examined the relationship between genetic diversity
found within clones (i.e., total heterozygosity) and clonal survival and reproduction (Supporting
Information, Dataset S1). We found no significant association between the amount of genetic
diversity within clones and their ability to survive or reproduce across diet conditions.
191
We used a similar analysis approach to assess the association between phenotypic
differentiation between clones and genetic variation in 8 gene regions previously implicated in
Daphnia transcriptomic responses to microcystins [47,48]. A PCA of the all variant sites with these
8 gene regions revealed that clone 4, a phenotypically distinct clone due to its overall low
survival and growth rates across conditions (Figure 1B), is largely separated from all other
clones by PC1. We found no significant association between the amount of divergence at these
putatively functional gene regions and the magnitude of overall phenotypic divergence between

clones (Figure 1) (mantel $r = 0.054$, $P = 0.584$). We also found no association between the
amount of genetic diversity at these putatively functional gene regions and their ability to survive
or reproduce at high (1:1) *Microcystis* exposure.
202

203
204 **Figure 1.** Multivariate a) whole genome genomic and b) phenotypic divergence across the all
205 treatments between 20 *Daphnia magna* clones visualized using principal component analysis
206 (PCA). Panel c) shows a PCA constructed from genetic variation at putative candidate loci for
207 variation in tolerance to *Microcystis* in *Daphnia*.
208
209 *Magnitude of phenotypic variation from toxic exposure*
210
Exposure to cyanobacteria reduced the overall fitness of *D. magna* genotypes following a
monotonic dose response in the moderately toxic (2:1 *Chlorella* : *M. aeruginosa*) and severely
toxic (1:1 *Chlorella* : *M. aeruginosa*) treatments (Figure 2). For example, mean survival across
all clones for the 14-day trial declined from 94% in the non-toxic diet, to 86.5% in the moderately
toxic diet, and to 53% in the severely toxic diet. We tested for differences in the magnitude of
clonal variation with increasing exposure to *M. aeruginosa* for each of the three fitness-

associated phenotypes (survival, growth, reproduction). Intraspecific variation in survival did not increase from the control to 2:1 ($F^* = 3.60$, $P = 0.066$) but did from 2:1 to 1:1 ($F^* = 22.59$, $P < 0.001$). Intraspecific variation in growth rate increased from the control to 2:1 ($F^* = 10.28$, $p = 0.003$) but decreased from 2:1 to 1:1 ($F^* = 11.82$, $P = 0.002$). Neonate production showed the most consistent pattern, with intraspecific variation between clones declining with increasing *M. aeruginosa* exposure (control to 2:1 ($F^* = 18.54$, $P = 0.0001$); 2:1 to 1:1 ($F^* = 7.44$, $P = 0.009$)).

Figure 2. Phenotypic variation among the 20 *Daphnia magna* clones at the end of the 14-d chronic toxicity study. Phenotypes measured in parents were a) survival, b) growth from day 0 to 14, c) total number of neonates produced, and d) total number of broods produced. Cell concentration was standardized across three algal food treatments: non-toxic - "C" , moderately-toxic - "2:1 C:M", and severely-toxic - "1:1 C:M". Plots show means and standard errors for each clone when reared on each diet.

Interactions under favorable and toxic conditions

We tested for interactions associated with performance across variations in diet toxicity. On average, survival declined with increasing toxicity by 8% in the moderately-toxic diet and 44% in the severely toxic diet relative to controls. However, we found evidence of an interaction between clonal identity and a *Microcystis* diet ($X^2 = 59.68$, $P = 0.014$). Some of these interactive effects were striking. Relative to rearing on the *Chlorella* diet, clones 5 and 15 showed an 80% decline in mean survival when reared on the severely toxic diet while clone 19 exhibited a 20% increase in survival. Similarly, we found a significant interaction associated with growth across treatments ($X^2 = 244.17$, $P < 0.0001$). Finally, neonate production showed a similar pattern, with a significant interaction between clonal identity and a *Microcystis* diet ($X^2 = 264.42$, $P < 0.0001$).

Of particular interest was that certain genotypes performed better in one of the toxic treatments
compared to the *Chlorella* diet (e.g., genotypes 6 in 2:1, and 13 in both 2:1 and 1:1, across all
life-history traits), while other genotypes performed better in the 1:1 versus 2:1 toxic treatments
(e.g., genotypes 10, 14, 18, 19, 20). These findings indicate that relative performance of clones
is dependent on the magnitude of toxicant exposure. reinforce the shifts in relative performance
across clones associated with changing *M. aeruginosa* conditions.
249
250 *Influence of genetic diversity on toxicological inferences*
251
We used a bootstrapping approach to determine how variation in the number of clones in the
experimental sampling design could influence the robustness of the data relative to including 20
clones. Our results demonstrate that sampling individuals from a larger number of genotypes is
critical to generating a robust estimate of reproduction in all 3 rearing conditions (Figure 3a). To
quantify the loss of robustness and accuracy that would result from assaying a reduced amount
of genetic diversity we assessed how frequently samples at five reduced diversity levels would
yield survival estimates within the 95% confidence interval (CI) obtained using all 20 clones
(Figure 3b). While including 15 clones always produced a survival estimate with the 95% CI
assays using only a single clone failed to produce an estimate within the 95% CI over half the
time in all three toxicity treatments. Overall, this reinforces the importance of accounting for
genetic diversity in toxicity testing, as failing to do so means estimates of reproduction and
survival are likely to be highly variable and inaccurate.
264

265
Figure 3. Panel A shows simulations of mean reproductive outputs for experimental designs
with varying amounts of genetic diversity of *Daphnia magna* across three treatments of chlorella
"C", 2:1 chlorella:microcystis "2:1 C:M", and 1:1 chlorella:microcystis "1:1 C:M". Symbol and
error bars represent the mean and standard deviation. A 95% confidence interval was
calculated for survival using all 20 clones. Panel B shows simulations of how often estimates of
survival fell within this 95% confidence interval when the data were subsetted to include a
reduced number of genotypes.
273
274 **Discussion**
275
Genetic variation can meaningfully impact population dynamics [49], community composition [50],
ecosystem function [51], and underlie rapid adaptation [10]. Incorporating natural genetic variation in
toxicology, which directly informs environmental regulations and policy, is critical to increasing
the reliability of benchtop toxicity assays to natural populations. Our data demonstrate that
genetic variation, even within a population from a single locality, has a profound impact on
toxicological responses in a widely used model system. For example, the moderately toxic diet
used in this study could be interpreted as an LC50 (i.e., clone 4) or an LC0 (clone 18 and 19)

depending on the clone selected for the assay. When reared on a severely toxic diet, mortality
ranged from 0% to 90%. Toxicity studies that do not include intraspecific genetic variation
necessarily cannot estimate this variation or provide context on whether the genotype or strain
used in the assay is in any way representative of the species or genus. This can lead to
considerable uncertainty when extrapolating results from laboratory to field, and have
implications for the accuracy and efficacy of risk management strategies [52]. The ramifications of
this oversight are likely to be exacerbated in natural populations where natural selection may
strongly shape genetic variation and toxicological responses [53,54].
291
The variation we observed came from a single population of *D. magna* and is doubtless an
underestimate of the magnitude of variation that would be found across populations that
experience dramatically different exposures to HABs. For example, if the species of focus spans
a wide range of ecological conditions, particularly with regards to prior toxicant exposure that
may have driven adaptation, there is likely to be substantially more intraspecific variation than
we observed within the single population we studied here. Hence, we advocate for the inclusion
of multiple genotypes in clonal test organisms or individuals from multiple populations in sexual
organisms when conducting toxicity studies. Ideally, studies should survey across a sufficiently
wide array of genetic diversity that is similar to populations that could be impacted by the
toxicant to allow 'clone' or 'population origin' to be treated as a random effect in linear mixed
models and hence generate reasonable estimates of the variation in key response variables
across the populations of interest [55].
304
In practice, we recognize that such robust sampling designs may be challenging and would
require shifting effort and focus in many toxicology studies. As a preliminary step, it must be
acknowledged by toxicology regulatory agencies that LC50s derived from real, genetically
heterogeneous populations will differ at local, regional, or global scale. Each geographic scale
presents its own sources of variation and shapes the requirements in terms of the sampling
required to achieve the stated objective of a meaningful safety limit recommendation based on
an LC50. Therefore, testing multiple genotypes within a single population (local/regional scale)
or multiple populations of genotypes (international scale) and reporting a CI of LC50s calculated
across each would be a strong step forward. Doing so would provide both a more accurate and
robust measure of the LC50, as it would be calculated from multiple genotypes or populations
and an indication of the uncertainty. The uncertainty of this estimate would not arise from
experimental noise but from real differences in biology. Reporting LC50s would then include
both the estimate and the variance around this estimate stemming from genetic variation. This
would be useful in discussions about risk and the connection between laboratory assays and
toxicological outcomes in natural settings, a key area of growth in translating benchtop
experiments to wild populations.
321
The study here demonstrates that considerable variation is found in toxicological responses
within a population. However, it is important to consider the overall scale and scope of variation
in toxicological responses. Current methods often examine focal taxa across broad phylogenetic
categories (e.g., use of a single surrogate species for a family or order), which is important. But
meaningful biological variation can be found across levels of organization. Future work
examining variation across biological scales —within populations, across populations, between
congeners, and across genera— is critical for parsing the levels of biodiversity most relevant for
risk assessment and management action. Identifying the biological scales at which the largest
fraction of functional variation is found has been transformative in ecology [56] and a similar
approach could also lead to a major advance in toxicology.
332

The level of genomic variation observed across clones in our study demonstrates the large
magnitude of genetic variation found within a single population of *Daphnia*. From a toxicological
perspective this variation is important both as it provides phenotypic variation, as demonstrated
here, and this variation is the raw material on which natural selection can act to drive
adaptation. Previous work has suggested that rapid adaptation could be a critical mechanism of
suppressing *Microcystis* blooms as field populations of *Daphnia* can rapidly adapt to tolerate
Microcystis and can suppress the biomass of HABs that contain lethal concentrations of
microcystins (3 to 6.5 µg L^{-1}) [43]. These data underscore the need for toxicological theory to
integrate genetic diversity, and potentially rapid adaptation in routine testing dose-responses
between chronically exposed [24,42,43,57] and unexposed [3,58] populations might well differ due to
genomic variation.

The pace and ecological consequences of resistance evolution has been appreciated for
decades [59], but a growing body of research is seeking to understand the molecular genetic
underpinnings of variation in resistance to toxins [15,60,61]. We conducted whole genome
sequencing of all 20 *D. magna* clones assayed in the study. These data provide a holistic view
of the genetic diversity found within a population but do not provide the power required to
discover the genetic basis of clonal differences in resistance to *M. aeruginosa*. This is
underscored by the lack of associations both between phenotypic variation and genomic
differentiation and genomic variation within clones and phenotypic responses. In short, it
appears variation in responses to cyanobacteria are not the primary force shaping genetic
variation and the raw amount of genetic variation itself is not a key determinant in performance
when exposed to toxicants. These results largely fit with the prevailing population genetic theory
for genome wide patterns [62].

The utility of genomic data in toxicology has been touted as potentially transformative - with a
gene-level understanding of resistance presented as a major goal [63]. It is possible to conduct
quantitative trait locus (QTL) mapping or genome wide association studies (GWAS) to find the
underlying genomic basis of trait variation. However, generating the resolution needed to
identify a causal genomic region, particularly specific genes, would require a herculean effort, a
relatively simple genetic basis, and likely some recombinational luck - efforts most likely to yield
payoff only in model systems. There is considerable interest in understanding the genetic basis
of microcystin resistance in *Daphnia* [47,48,64]. We investigated the link between variation in genes
putatively involved in resistance and a clones' ability to survive and reproduce when exposed to
Microcystis and found no significant relationship. Clone 4, which was distinct in both phenotype
and genotype for putatively functional loci, does have a private haplotype containing 17
singleton SNPs over a 5 kb window of the tyrosine-protein kinase ABL1. However, this clone
had >22,000 unique singleton SNPs making any inference about the function of specific genetic
variants exceeding tenuous. Given the complex and environmentally variable genetic
architecture of trait variation for many complex traits, we advocate that toxicologists focus
genetic studies on assessments of clonal, family, or population level variation. Explicitly
including this information on origin in study designs and statistical models will yield powerful
insights without requiring major changes in the methods most often employed in the discipline.

Conclusions

Studies of genetic variation have improved our understanding of ecological and evolutionary
processes [60]. Arguments in favor of incorporating genetic variation, and to an extent evolution,
in the environmental risk assessment process are growing stronger as the recognition that
meaningful levels of intraspecific genetic variation exist in key toxicological traits [21,23,52,65–67].
Ignoring genetic variation can lead to inaccurate reporting of LC50 toxicity thresholds [23], which

may obscure safety measures adhered by companies invested in the commercial production of
chemicals and other categories of stakeholders more concerned with toxicants threatening
aquatic lifebiodiversity. Here, we show that genotypic and phenotypic differences in *D. magna*
tolerance to the aquatic contaminant *M. aeruginosa* were substantial across multiple life-history
traits. Significant interactions between growth and reproduction were additionally observed
among genotypes in green algae versus cyanobacteria treatments. These results suggest that
genetic diversity can strongly influence population dynamics with and without toxicant exposure.
Using simulations and bootstrapping approaches we further demonstrate that the common
toxicological practice of using a single genotype to calculate toxicity tolerance leads to an
estimate outside of the 95% confidence interval >50% of the time. We suggest that integrating
intraspecific genetic variation into routine laboratory toxicity studies would improve the reliability
and biological realism of LC50 calculations and improve both the environmental risk
assessment process and our understanding of the transgenerational impacts of aquatic
contaminants on natural populations.
398
399 **Methods**
400
401 *Daphnia magna field collection and culturing*
402
Twenty genotypes of *D. magna* were collected from 'Langerodevijver' (LRV; 50° 49' 42.08", 04°
38' 20.60"), a small lake (surface area = 140,000 m^2, max depth = 1 m) within the nature
reserve of Doode Bemde, Vlaams-Brabant, Belgium [68], in late spring and early summer. LRV
has a single basin and has seasonal *Microcystis* HABs typical to many lakes globally and has a
large resident population of *D. magna*. Parthenogenetic lines of each genotype were maintained
for over five years (approximately 300 generations) in continuous cultures at 20 ºC in UV-filtered
dechlorinated municipal tap water containing 2 mg C L^{-1} of the green alga *C. vulgaris* (strain
CPCC 90; Canadian Phycological Culture Centre, Waterloo, ON, Canada). *C. vulgaris* was
grown in COMBO medium [69].
412
To prepare for this study, we isolated one adult female *D. magna* per genotype in separate 50-
mL glass tubes inoculated with COMBO medium and *C. vulgaris* at 2 mg C L^{-1}, and monitored
them daily for reproduction. *D. magna* neonates born within 24 hr were collected from their
respective genotypes and individually separated into 50-mL glass tubes as previously
described, totalling 10 replicates per genotype and 200 tubes total. These 200 *D. magna*
representing 20 genotypes were the founding mothers of this study. All *D. magna* were
incubated under constant conditions (temperature of 21 ± 1 ºC, cool-white fluorescent light of
600 ± 15 lx, with a photoperiod of 16:8 h light:dark).
421
422 *Microcystis aeruginosa culturing*
423
Following our previously described method [70], *M. aeruginosa* (strain CPCC 300; Canadian
Phycological Culture Centre, Waterloo, ON, Canada) was cultured in BG-11 media and kept in a
growth chamber under axenic conditions with a fixed temperature of 21 ± 1 ºC, cool-white
fluorescent light of 600 ± 15 lx, with a photoperiod of 16:8 h light:dark. The culture was grown
for a minimum of one month before preparation for the chronic study. *M. aeruginosa* CPCC 300
produces microcystins-LR (CAS: 101043-37-2, $C_{49}H_{74}N_{10}O_{12}$) and its desmethylated form [D-
Asp³]-microcystin-LR (CAS: 120011-66-7, $C_{48}H_{72}N_{10}O_{12}$), which occur widely in freshwater
ecosystems [34,36] and are toxic to zooplankton.
432
433 To prepare *M. aeruginosa* for testing on *D. magna*, an aliquot of the stock was inoculated in
434 100% COMBO medium for two weeks prior to test initiation and cultured to a cell concentration

435   of $1.25 \pm 0.02 \times 10^7$ cells mL^{-1}. This medium was chosen because it supports the growth of
436   algae and cyanobacteria and is non-toxic to zooplankton [69].
437
438   *Chronic toxicity study*
439
A 14-d chronic toxicity study was performed in accordance with the test guideline of the
Organization for Economic Cooperation and Development to assess the effects of contaminants
on the survival, growth, and reproductive output of *D. magna* [71]. The OECD test guidelines are
widely used in toxicity laboratory studies. Three treatments were included in this study: 100%
chlorella (non-toxic diet), 2:1 chlorella:microcystis (moderately toxic diet), and 1:1
chlorella:microcystis (severely toxic diet). Following these ratios all *D. magna* were fed 2 mg C
L^{-1}, corresponding to 3×10^6 cells total, which corroborates with previous literature exposing
daphnids to dietary combinations of green algae and cyanobacteria [41,44,72].
448
Following the OECD test guidelines (Test No. 211: Daphnia magna Reproduction Test [71]), *D.*
magna neonates born within 24 hr were collected from their respective genotype, and 10
neonates were individually placed in 50-mL glass tubes containing 50 mL of UV-filtered water
and its respective treatment. To ensure there were no systematic differences in ages between
D. magna neonates that could drive variation in responses within and across clones, a stringent
protocol was implemented to remove neonates twice, daily in the morning and at night in
preparation for the toxicity test. The UV-filtered water met the validity criteria for the test
following an accredited method for *D. magna* culturing and toxicity testing [73]. A total of 600
animals were used for this study (i.e., 10 replicates per genotype, 200 animals per treatment).
Since this was a semi-static test, solutions were renewed 3 × wk on Mondays, Wednesdays,
and Fridays by transferring *D. magna* from old to new glass tubes, followed by supplying each
D. magna with 3×10^6 cells of food (e.g., the 2:1 treatment received 2×10^6 *C. vulgaris* cells
and 1×10^6 *M. aeruginosa* cells, corresponding with 2 mg C L^{-1}). During this time, survival,
growth, reproductive output, and number of broods were recorded to assess for interactions
between genotype and treatment effects. Growth (mm) for each replicate was also measured on
days 0, 7, and 14. The study was incubated under 400–800 lx cool-white fluorescent light at 20
$\pm$ 1 °C with a 16:8 light:dark cycle.
466
467   *Genomics sequencing and analysis*
468
Pools of individuals were collected from each clone and DNA was extracted using the QIAGEN
MagAttract HMW DNA kit. Extractions were checked for DNA concentration using a Qubit 2.0
fluorometer and libraries for each clone were prepared using the KAPA Hyper Prep kit. The
mean DNA concentration and mean total quantity of DNA used for library construction was 6.0 ±
0.4 ng µL^{-1} and 304.3 ± 15.6 ng, respectively. Sequencing of these libraries was generated on a
NovaSeq 6000 generating 150bp paired-end reads. Sequences were then trimmed using
trimmomatic/0.39 using a quality threshold of 20 and a min length of 50 and subsequently
mapped to the *D. magna* reference genome (GCA_020631705.2) using *bwa/0.7.17* (default
parameters). Duplicates were marked using *Picard tools/2.21.2* and then the bam files were
input into *freebayes/1.2.0* to generate a VCF of all variant positions among all individuals on the
10 *D. magna* chromosomes (not scaffolds). *Vcftools* was then used to trim to a minimum quality
score of 30. This produced a whole genome dataset with an average genome sequencing
coverage of 29.5 (min=26.6, max=35.5, SE=0.34) for each of the 20 clones. The total number of
variant sites and the number of heterozygous sites for each clone was extracted using
VCFtools. Genetic distance was calculated using *VCF2dis* (https://github.com/BGI-
shenzhen/VCF2Dis). Genotype likelihood (beagle) files were generated using *angsd* (-GL 1 -out

D23Genolike -doMaf 2 -SNP_pval 1e-6 -doMajorMinor 1 -doGlf 2), and *pcangsd* was used to
generate principal component analysis.
487
488 *Water sampling for analysis*
489
490 Water samples were collected from each treatment at test initiation, solution changes, and
491 termination to quantify the cell concentration, fluorescence, and standard water parameters and
492 were quick-frozen at -80 °C before analysis for microcystins and oligopeptides (Supporting
493 Information, Dataset S1).
494
495 *Chemicals and Reagents for water analysis*
496
Water, acetonitrile, methanol, acetic acid, and formic acid were all Optima LC/MS grade
solvents and purchased from Fisher Scientific (Tewksbury, MA, USA). Microcystin standards D-
Asp^3-RR, MC-RR, YR, HtyR, LR, D-Asp^3-LR, HilR, WR, LA, LY, LW, and LF, as well as
nodularin and cylindrospermopsin were purchased from Enzo Life Sciences (Farmingdale, NY,
USA). Microcystin Dha^7-LR was purchased from National Research Council Canada (Ottawa,
ON, Canada). Microcystin Leu^1-LR was purchased from GreenWater Laboratories (Palatka, FL,
USA). D-Asp^3-E-Dhb^7-RR was purchased from Sigma Aldrich. Anabaenopeptin A and B,
oscillamide Y, oscillaginin A, microginin 690 methyl ester, aeruginosamide B and C,
cyanopeptolin 1040 MB and 1007, and micropeptin 1106 were purchased from MARBIONC
(Wilmington, NC, USA). Anabaenopeptin F was synthetically created by the Stockdill laboratory
at Wayne State University (Detroit, MI, USA). ^{15}N MC-LR was purchased from Cambridge
Isotope Laboratories, INC. (Tewksbury, MA, USA).
509
510 *Microcystin and cyanopeptide water analysis*
511
Microcystins were measured using a method based on the US EPA method 544 [74]. Quality
assurance and quality control met the criteria set by the US EPA method 544. Dataset S1
depicts the standard curve range, retention time, and linearity, as well as the quantitative and
qualitative ions that were used for further identification of the microcystins. Surrogate recovery
for the samples were within the US EPA method 544 limit of 60-130%, and the control
standards were within the limits of 70-130% set by this same method. All duplicates relative %
differences were calculated to be < 30%.
519
Microcystin and cyanopeptide standards and samples were analyzed using a Thermo Scientific
TSQ Altis™ triple quadrupole mass spectrometer (Waltham, MA, USA) with a TriPlus™ RSH
EQuan 850 system. Both online concentration analysis and direct inject analysis were
completed [74]. Quantitative results shown use the direct inject analysis method. Detection limits
for the online concentration method were between 0.5-5 ng L^{-1}, and the standard curve range
was between 0.5-500 ng L^{-1}.
526
For direct injection analysis, 10 µL injections of standards and samples were loaded onto a
Thermo Hypersil GOLD™ C18 2.1 x 50 mm, 1.9 µm column using a TriPlus RSH autosampler
equipped with a 100 µL sampling syringe. The analytical separation gradient consisted of 0.1%
formic acid in water (mobile phase A) and 0.1% formic acid in acetonitrile (mobile phase B). At a
flow rate of 0.5 mL/min, the gradient started at 24% B and was held for 0.7 min. It was then
ramped to 26% B from 0.7–1.7 min, from 26–50% B from 1.7–5.5 min. The column was washed
at 98% B from 5.51–6.5 min at a flow rate 1.0 mL/min, then brought back down to original
conditions to equilibrate. The total run time was 8.5 min. The column oven temperature was

held at 35°C. The standard curve was prepped in LC-MS water and the range measured
between 0.05-50 µg L^{-1}.
537
538 Mass spectrometry analysis was performed using electrospray ionization source (ESI). The MS
539 source settings were as follows: spray voltage 3500 V, ion transfer tube temperature at 325 °C,
540 vaporizer temperature at 350°C, sheath gas 35 (arbitrary units), auxiliary gas 10 (arbitrary
541 units), sweep gas 0 (arbitrary units), collision gas at 1.5 mTorr, and Q1/Q3 resolution set to 0.7
542 Da.
543
544 Control samples and standards were performed every 20 samples during each run. Control
545 standards included negative control (blank) and positive control (10 ppb standard) as well as
546 duplicate and fortified samples. For each standard, a quantitative and qualitative ion transition
547 were chosen for use with selected reaction monitoring (SRM).
548
549 *Selective ion mode analysis*
550
For selective ion mode (SIM) analysis, the following masses in Supporting Information, Dataset
S1 were investigated because they are known to be produced by *M. aeruginosa* CPCC 300
(Racine et al. 2019). The gradient for the method consisted of 0.1% formic acid in water (A) and
0.1% formic acid in acetonitrile (B). At a flow rate of 0.35 mL min^{-1}, the gradient started at 0% B
and was held for 1.0 min. It was then ramped to 60% B from 1.0-8.5 min. The column was
washed at 98% B from 8.51–10.5 min at a flow rate 0.6 mL min^{-1}, then brought back down to
original conditions to equilibrate. The total run time was 14 min. The injection volume was 10 µL
and the column oven temperature was held at 35 °C. Mass spectrometer source settings were
the same as the quantitative method above. The SIM method was accomplished in Q3. The
area under the m/z peak was calculated and reported.
561
562 *Testing criteria and toxin analysis*
563
Measured microcystin concentrations were 3.3 ± 0.001 µg L^{-1} and 5.1 ± 0.001 µg L^{-1} in the 2:1
and 1:1 toxic treatments, respectively (Dataset S1). Water parameters before and after solution
renewals were also within the test criteria (Dataset S1). There were no signs of hypoxia stress
on *D. magna* behavior. Following the criteria for survival and reproduction of controls, all test
acceptability criteria were met. Mortality and immobilization were first observed within 48 h after
microcystin exposure in either treatment, and continued onward throughout the test. This was
expected as the microcystin concentrations chosen can be sublethal [44] to lethal [45,46] in *Daphnia*
laboratory assays and are commonly measured in freshwater ecosystems with HABs [36].
572
573 *Statistical analysis*
574
575 *For survival, growth, and neonate production*
576
To test the magnitude of clonal variation, we used a Linear Mixed Effects (LME) model (or a
Generalized Linear Model (GLM) for binomial survival data) for each response with clone
treated as a random effect and tested for significant differences in fit between models with and
without clone included as random effect [75]. Second, to determine whether the concentration of
Microcystis influenced the magnitude of clonal variation we conducted Brown-Forsythe tests for
homogeneity of variances after generating means for each clone for each response variable.
These Brown-Forsythe tests were done sequentially - meaning comparisons were between
Chlorella and 2:1, then 2:1 and 1:1. Finally, to test for interactions between response variables
across *Microcystis* concentrations we used linear models with concentration and clone as fixed

effects (*glm* with binomial for survival, *lm* for growth, and *glm* with a gamma distribution for
neonate production).
588
589 *Association between clonal genomic diversity, genomic divergence, and phenotypic distance*
590
We tested for an association between the amount of genetic diversity found within clones and
clonal survival and reproduction across each of the three diet conditions. To do so we regressed
the total number of heterozygous sites and each response variable for each rearing condition.
To test for associations between genomic diversity and phenotypic distance, we first generated
distance matrices from mean values for all 20 clones. The genomic distance matrix was
constructed from all polymorphic sites. The phenotypic distance matrix was constructed using
neonate production, time to first brood, total broods, week one growth (day 0-7), week two
growth (day 7-14), week one survival, and week two survival for each clone for each diet
condition. A two-tailed Mantel test was then conducted to determine whether there was
significant correlation between matrices that differed from 0.
601
We conducted a similar set of analyses exclusively using genes previously identified as involved
in microcystin tolerance. These two unique *Daphnia* studies identified potential candidate genes
for microcystin tolerance in *Daphnia* through a combination of differential expression analysis
(RNAseq) and employing functional annotations [47,48]. We search the gene names provided by
the authors in the NCBI database. If only a single gene region matched the name we included it
in our downstream analysis. For gene names that had multiple matches we selected from the
most exact. Finally, when authors provided primer regions we used ePCR to identify the gene
region that would be amplified in the reference. Using these methods we were able to locate 8
potential candidate loci for genetic variation in microcystin tolerance. We then identified all
variant sites across the 20 *Daphnia* clones within these eight candidate gene regions, including
1,000 bp up and downstream of the demarcated gene, for each of 20 clones. We constructed a
PCA using this data, conducted a two-tailed Mantel test to determine whether there was
significant correlation between differences at these loci and phenotypic variation, and tested for
associations between the number of variants in these eight genes and survival and reproduction
in high (1:1) *Microcystis* conditions.
617
618 *Examining the effects of assays with reduced diversity*
619
We sought to determine how the choice to conduct lower diversity assays might influence the
estimate of the effects of the toxicant on reproduction and survival. For reproductive output we
calculated means and standard errors for all 20 clones and for 15, 10, 5, 3, and 1 clonal
diversity groups created by subsetting the full dataset 100 times at each diversity level. For
survival, we calculated the 95% CI of survival probability using the means of all 20 clones. We
then created subsamples containing 15, 10, 5, 3, and 1 clones and determined how frequently
they would produce survival estimates within the 95% CI obtained from the full dataset.
627
628 All analyses were completed in R version 4.2.2 [76].
629
630 **Acknowledgements**
631

632 We thank the Rudman and Fryxell labs for helpful comments and discussions. This work was
633 supported by the Canada First Research Excellence Fund through the Food from Thought
634 Program (JMF), an NSERC Discovery Grant (JMF), the National Institute of General Medicine of
635 the National Institutes of Health (Award #1R35GM147264; SMR), a Banting Postdoctoral

639 **Supporting Information**
640
641 Dataset S1: Tables S1-S8, Figure S1
642
643 **References**
644
1.   Levin, S. A., Harwell, M. A., Kelly, J. R. & Kimball, K. D. Ecotoxicology: Problems and
Approaches. in *Ecotoxicology: Problems and Approaches* (eds. Levin, S. A., Kelly, J. R.,
Harwell, M. A. & Kimball, K. D.) 3–7 (Springer New York, 1989).
2.   Soares, A. M. V. M., Baird, D. J. & Calow, P. Interclonal variation in the performance
ofDaphnia magnastraus in chronic bioassays. *Environ. Toxicol. Chem.* **11**, 1477–1483
(1992).
3.   Barata, C., Baird, D. J. & Soares, A. M. V. M. Determining genetic variability in the
distribution of sensitivities to toxic stress among and within field populations of Daphnia
magna. *Environ. Sci. Technol.* **36**, 3045–3049 (2002).
4.   Clements, W. H. & Rohr, J. R. Community responses to contaminants: using basic
ecological principles to predict ecotoxicological effects. *Environ. Toxicol. Chem.* **28**, 1789–
1800 (2009).
5.   Newman, M. C. *Fundamentals of ecotoxicology*. (CRC press, 2009).
6.   Stark, J. D., Vargas, R. I. & Banks, J. E. Incorporating variability in point estimates in risk
assessment: Bridging the gap between LC50 and population endpoints. *Environ. Toxicol.
Chem.* **34**, 1683–1688 (2015).
7.   Barata, C., Baird, D. J. & Markich, S. J. Influence of genetic and environmental factors on
the tolerance of Daphnia magna Straus to essential and non-essential metals. *Aquat.
Toxicol.* **42**, 115–137 (1998).
8.   Relyea, R. & Hoverman, J. Assessing the ecology in ecotoxicology: a review and synthesis
in freshwater systems. *Ecol. Lett.* **9**, 1157–1171 (2006).
9.   Crutsinger, G. M. *et al.* Plant genotypic diversity predicts community structure and governs
an ecosystem process. *Science* **313**, 966–968 (2006).
10. Barrett, R. D. H. & Schluter, D. Adaptation from standing genetic variation. *Trends Ecol.
Evol.* **23**, 38–44 (2008).
11. Des Roches, S. *et al.* The ecological importance of intraspecific variation. *Nat Ecol Evol* **2**,
57–64 (2018).
12. Whitham, T. G., Allan, G. J., Cooper, H. F. & Shuster, S. M. Intraspecific Genetic Variation
and Species Interactions Contribute to Community Evolution. *Annu. Rev. Ecol. Evol. Syst.*
51, 587–612 (2020).
13. Rudman, S. M. *et al.* Direct observation of adaptive tracking on ecological time scales in
Drosophila. Science **375**, eabj7484 (2022).
14. Weedall, G. D. *et al.* An Africa-wide genomic evolution of insecticide resistance in the
malaria vector Anopheles funestus involves selective sweeps, copy number variations,
gene conversion and transposons. *PLoS Genet.* **16**, e1008822 (2020).
15. Reid, N. M. *et al.* The genomic landscape of rapid repeated evolutionary adaptation to toxic
pollution in wild fish. *Science* **354**, 1305–1308 (2016).
16. Lemaire, V. *et al.* Genotype × genotype interactions between the toxic cyanobacterium
Microcystis and its grazer, the waterflea Daphnia. *Evol. Appl.* **5**, 168–182 (2012).
17. Hochmuth, J. D., De Meester, L., Pereira, C. M. S., Janssen, C. R. & De Schamphelaere,
K. A. C. Rapid Adaptation of a Daphnia magna Population to Metal Stress Is Associated
with Heterozygote Excess. *Environ. Sci. Technol.* **49**, 9298–9307 (2015).

18. Jansen, M. *et al.* Experimental evolution reveals high insecticide tolerance in Daphnia
inhabiting farmland ponds. *Evol. Appl.* **8**, 442–453 (2015).

19. Albert, C. H. *et al.* A multi-trait approach reveals the structure and the relative importance of
intra- vs. interspecific variability in plant traits. *Funct. Ecol.* **24**, 1192–1201 (2010).

20. Lemmen, K. D., Butler, O. M., Koffel, T., Rudman, S. M. & Symons, C. C. Stoichiometric
Traits Vary Widely Within Species: A Meta-Analysis of Common Garden Experiments.
Frontiers in Ecology and Evolution **7**, (2019).

21. Bickham, J. W. The four cornerstones of Evolutionary Toxicology. *Ecotoxicology* **20**, 497–
502 (2011).

22. Miner, B. E., De Meester, L., Pfrender, M. E., Lampert, W. & Hairston, N. G., Jr. Linking
genes to communities and ecosystems: Daphnia as an ecogenomic model. *Proc. Biol. Sci.*
279, 1873–1882 (2012).

23. Ebert, D. Daphnia as a versatile model system in ecology and evolution. *Evodevo* **13**, 16
(2022).

24. Hairston, N. G., Jr *et al.* Natural selection for grazer resistance to toxic cyanobacteria:
evolution of phenotypic plasticity? *Evolution* **55**, 2203–2214 (2001).

25. Govaert, L., De Meester, L., Spaak, P. & Hairston, N. G. Eco-Evolutionary Dynamics in
Freshwater Systems. in *Reference Module in Earth Systems and Environmental Sciences*
(Elsevier, 2021).

26. Lynch, M., Wei, W., Ye, Z. & Pfrender, M. The Genome-wide Signature of Short-term
Temporal Selection. *bioRxiv* 2023.04.28.538790 (2023) doi:10.1101/2023.04.28.538790.

27. Baird, D. J., Barber, I., Bradley, M., Soares, A. M. & Calow, P. A comparative study of
genotype sensitivity to acute toxic stress using clones of Daphnia magna straus. *Ecotoxicol.*
Environ. Saf. **21**, 257–265 (1991).

28. Rudman, S. M. *et al.* What genomic data can reveal about eco-evolutionary dynamics. *Nat*
Ecol Evol **2**, 9–15 (2018).

29. Boyle, E. A., Li, Y. I. & Pritchard, J. K. An Expanded View of Complex Traits: From
Polygenic to Omnigenic. *Cell* **169**, 1177–1186 (2017).

30. van Straalen, N. M. & Feder, M. E. Ecological and Evolutionary Functional Genomics—How
Can It Contribute to the Risk Assessment of Chemicals? *Environ. Sci. Technol.* **46**, 3–9
(2012).

31. Rennison, D. J., Rudman, S. M. & Schluter, D. Genetics of adaptation: Experimental test of
a biotic mechanism driving divergence in traits and genes. *Evol Lett* **3**, 513–520 (2019).

32. Betancourt, N. J. *et al.* Allelic polymorphism at foxo contributes to local adaptation in
Drosophila melanogaster. *Mol. Ecol.* **30**, 2817–2830 (2021).

33. Merder, J. *et al.* Geographic redistribution of microcystin hotspots in response to climate
warming. *Nature Water* **1**, 844–854 (2023).

34. Harke, M. J. *et al.* A review of the global ecology, genomics, and biogeography of the toxic
cyanobacterium, Microcystis spp. *Harmful Algae* **54**, 4–20 (2016).

35. Reid, A. J. *et al.* Emerging threats and persistent conservation challenges for freshwater
biodiversity. *Biol. Rev. Camb. Philos. Soc.* **94**, 849–873 (2019).

36. Chorus, I. & Welker, M. *Toxic Cyanobacteria in Water:A Guide to Their Public Health*
Consequences, Monitoring and Management. (CRC Press, 2021).

37. Shahmohamadloo, R. S. *et al.* Lake Erie fish safe to eat yet afflicted by algal hepatotoxins.
Sci. Total Environ. **861**, 160474 (2023).

38. Shahmohamadloo, R. S. *et al.* Diseases and Disorders in Fish due to Harmful Algal
Blooms. in *Climate Change on Diseases And Disorders Of Finfish In Cage Culture, 3rd*
Edition (eds. Woo, P. T. K. & Subasinghe, R. P.) 387–429 (CAB International, 2023).

39. Shahmohamadloo, R. S. *et al.* Cyanotoxins accumulate in Lake St. Clair fish yet their fillets
are safe to eat. *Sci. Total Environ.* **874**, 162381 (2023).

40. Hairston, N. G., Jr *et al.* Lake ecosystems: Rapid evolution revealed by dormant eggs.
Nature **401**, 446–446 (1999).
41. Isanta-Navarro, J. *et al.* Reversed evolution of grazer resistance to cyanobacteria. *Nat.*
Commun. **12**, 1945 (2021).
42. Sarnelle, O. & Wilson, A. E. Local adaptation ofDaphnia pulicariato toxic cyanobacteria.
Limnol. Oceanogr. **50**, 1565–1570 (2005).
43. Chislock, M. F., Sarnelle, O., Jernigan, L. M. & Wilson, A. E. Do high concentrations of
microcystin prevent Daphnia control of phytoplankton? *Water Res.* **47**, 1961–1970 (2013).
44. Shahmohamadloo, R. S., Poirier, D. G., Ortiz Almirall, X., Bhavsar, S. P. & Sibley, P. K.
Assessing the toxicity of cell-bound microcystins on freshwater pelagic and benthic
invertebrates. *Ecotoxicol. Environ. Saf.* **188**, 109945 (2020).
45. Ferrão-Filho, A. S., Azevedo, S. M. F. O. & DeMott, W. R. Effects of toxic and non-toxic
cyanobacteria on the life history of tropical and temperate cladocerans. *Freshw. Biol.* **45**, 1–
19 (2000).
46. Lürling, M. & van der Grinten, E. Life-history characteristics ofDaphniaexposed to dissolved
microcystin-LR and to the cyanobacteriumMicrocystis aeruginosawith and without
microcystins. *Environ. Toxicol. Chem.* **22**, 1281–1287 (2003).
47. Schwarzenberger, A. *et al.* Deciphering the genetic basis of microcystin tolerance. *BMC*
Genomics **15**, 776 (2014).
48. Lyu, K. *et al.* Transcriptomic analysis dissects the mechanistic insight into the Daphnia
clonal variation in tolerance to toxic Microcystis. *Limnol. Oceanogr.* **64**, 272–283 (2019).
49. Yoshida, T., Jones, L. E., Ellner, S. P., Fussmann, G. F. & Hairston, N. G., Jr. Rapid
evolution drives ecological dynamics in a predator-prey system. *Nature* **424**, 303–306
(2003).
50. Farkas, T. E., Mononen, T., Comeault, A. A., Hanski, I. & Nosil, P. Evolution of camouflage
drives rapid ecological change in an insect community. *Curr. Biol.* **23**, 1835–1843 (2013).
51. Rudman, S. M. *et al.* Adaptive genetic variation mediates bottom-up and top-down control
in an aquatic ecosystem. *Proc. Biol. Sci.* **282**, 20151234 (2015).
52. Coutellec, M.-A. & Barata, C. An introduction to evolutionary processes in ecotoxicology.
Ecotoxicology **20**, 493–496 (2011).
53. Hawkins, N. J., Bass, C., Dixon, A. & Neve, P. The evolutionary origins of pesticide
resistance. *Biol. Rev. Camb. Philos. Soc.* **94**, 135–155 (2019).
54. Weston, D. P. *et al.* Multiple origins of pyrethroid insecticide resistance across the species
complex of a nontarget aquatic crustacean, Hyalella azteca. *Proc. Natl. Acad. Sci. U. S. A.*
110, 16532–16537 (2013).
55. Bolker, B. M. *Ecological Models and Data in R*. (Princeton University Press, 2008).
56. McGill, B. J., Enquist, B. J., Weiher, E. & Westoby, M. Rebuilding community ecology from
functional traits. *Trends Ecol. Evol.* **21**, 178–185 (2006).
57. Gustafsson, S. & Hansson, L.-A. Development of tolerance against toxic cyanobacteria in
Daphnia. *Aquat. Ecol.* **38**, 37–44 (2004).
58. Forbes, V. E. Genetics and ecotoxicology—insights from the interface. in *Genetics and*
Ecotoxicology 1–8 (CRC Press, 1998).
59. Mallet, J. The evolution of insecticide resistance: Have the insects won? *Trends Ecol. Evol.*
4, 336–340 (1989).
60. Oziolor, E. M., Bickham, J. W. & Matson, C. W. Evolutionary toxicology in an omics world.
Evol. Appl. **10**, 752–761 (2017).
61. Oziolor, E. M. *et al.* Adaptive introgression enables evolutionary rescue from extreme
environmental pollution. *Science* **364**, 455–457 (2019).
62. Kimura, M. *The Neutral Theory of Molecular Evolution*. (Cambridge University Press, 1983).
63. Gaytán, B. D. & Vulpe, C. D. Functional toxicology: tools to advance the future of toxicity
testing. *Front. Genet.* **5**, 110 (2014).

64. Schwarzenberger, A., Kuster, C. J. & Von Elert, E. Molecular mechanisms of tolerance to
cyanobacterial protease inhibitors revealed by clonal differences in Daphnia magna. *Mol.*
Ecol. **21**, 4898–4911 (2012).
65. Breitholtz, M., Rudén, C., Hansson, S. O. & Bengtsson, B.-E. Ten challenges for improved
ecotoxicological testing in environmental risk assessment. *Ecotoxicol. Environ. Saf.* **63**,
324–335 (2006).
66. Oziolor, E. M., DeSchamphelaere, K., Lyon, D., Nacci, D. & Poynton, H. Evolutionary
Toxicology-An Informational Tool for Chemical Regulation? *Environ. Toxicol. Chem.* **39**,
257–268 (2020).
67. Loria, A., Cristescu, M. E. & Gonzalez, A. Genotype diversity promotes the persistence of
Daphnia populations exposed to severe copper stress. *J. Evol. Biol.* **35**, 265–277 (2022).
68. Orsini, L., Spanier, K. I. & DE Meester, L. Genomic signature of natural and anthropogenic
stress in wild populations of the waterflea Daphnia magna: validation in space, time and
experimental evolution. *Mol. Ecol.* **21**, 2160–2175 (2012).
69. Kilham, S. S., Kreeger, D. A., Lynn, S. G., Goulden, C. E. & Herrera, L. COMBO: a defined
freshwater culture medium for algae and zooplankton. *Hydrobiologia* **377**, 147–159 (1998).
70. Shahmohamadloo, R. S. *et al.* An efficient and affordable laboratory method to produce and
sustain high concentrations of microcystins by Microcystis aeruginosa. *MethodsX* **6**, 2521–
2535 (2019).
71. OECD. Test No. 211: Daphnia magna Reproduction Test, OECD Guidelines for the Testing
of Chemicals, Section 2. in (OECD Publishing, 2012).
72. Rohrlack, T. *et al.* Ingestion of microcystins byDaphnia: Intestinal uptake and toxic effects.
Limnol. Oceanogr. **50**, 440–448 (2005).
73. Ontario Ministry of the Environment and Climate Change Aquatic Toxicology Unit. *Standard*
Operating Procedure: Daphnia Magna Culturing, SOP DM1, Volume 9. (2014).
74. Birbeck, J. A., Westrick, J. A., O'Neill, G. M., Spies, B. & Szlag, D. C. Comparative Analysis
of Microcystin Prevalence in Michigan Lakes by Online Concentration LC/MS/MS and
ELISA. *Toxins* **11**, (2019).
75. Bolker, B. M. *et al.* Generalized linear mixed models: a practical guide for ecology and
evolution. *Trends Ecol. Evol.* **24**, 127–135 (2009).
76. R Core Team. R: A Language and Environment for Statistical Computing. Preprint at
(2022).