1

Re-annotation of SARS-CoV-2 proteins using an HHpred-based approach opens new opportunities for a better understanding of this virus

Pierre Brézellec[*1,2]

[1] Institut Systématique Evolution Biodiversité (ISYEB UMR 7205), Sorbonne Université, MNHN, CNRS, EPHE, UA, Paris, France.

[2] Université de Versailles Saint Quentin, 45 avenue des Etats Unis, 78000 Versailles, France.

[*]Corresponding author

Correspondence: pierre.brezellec@uvsq.fr

ABSTRACT

Since the publication of the genome of SARS-CoV-2 – the causative agent of COVID-19 – in January 2020, many bioinformatic tools have been applied to annotate its proteins. Although effcient methods have been used, such as the identification of protein domains stored in Pfam, most of the proteins of this virus have no detectable homologous protein domains outside the viral taxa. As it is now well established that some viral proteins share similarities with proteins of their hosts, we decided to explore the hypothesis that this lack of homologies could be, at least in part, the result of the documented loss of sensitivity of Pfam Hidden Markov Models (HMMs) when searching for domains in "divergent organisms". In order to improve the annotation of SARS-CoV-2 proteins, we used the HHpred protein annotation tool. To avoid "false positive predictions" as much as possible, we designed a robustness procedure to evaluate the HHpred results. In total, 6 robust similarities involving 6 distinct SARS-CoV-2 proteins were detected. Of these 6 similarities, 3 are already known and well documented, and

one is in agreement with recent crystallographic results. We then examined carefully the two
similarities that have not yet been reported in the literature. We first show that the C-terminal
part of Spike S (the protein that binds the virion to the cell membrane by interacting with the
host receptor, triggering infection) has similarities with the human prominin-1/CD133; after
reviewing what is known about prominin-1/CD133, we suggest that the C-terminal part of Spike
S could both improve the docking of Spike S to ACE2 (the main cell entry receptor for SARS-
CoV-2) and be involved in the delivery of virions to regions where ACE2 is located in cells.
Secondly, we show that the SARS-CoV-2 ORF3a protein shares similarities with human G
protein-coupled receptors (GPCRs) belonging mainly to the "Rhodopsin family"; on the basis of
the literature, we then show that specific G protein-coupled receptors (GPCRs) of this family
are known to form ion channels; we emphasize this is consistent with a recent Cryo-EM
structure of SARS-CoV-2 ORF3a suggesting that it can form a non-selective Ca^{2+}-permeable
cation channel; furthermore, we highlight that some of the GPCRs identified as sharing
similarities with ORF3a are targeted by antibodies in patients with COVID-19 and Long-COVID,
suggesting that these similarities may trigger some of the observed autoimmune responses. We
conclude that the approach described here (or similar approaches) opens up new avenues of
research to better understand SARS-CoV-2 and could be used to complement virus annotations,
particularly for less-studied viruses.

Keywords: Pfam Domains, HHpred, Hidden Markov Models (HMMs), Bioinformatics, Protein annotation,
SARS-CoV-2.

Introduction

A significant fraction of the proteins expressed by viruses often lack homologs. These proteins are termed "orphan" to emphasise that no homologs are detected, or "taxonomically restricted" to indicate that they have no detectable homologs outside a given taxon (Kuchibhatla *et al.*, 2014). SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2), the causative agent of COVID-19, is no exception. According to UniProt (UniProt Consortium, 2021), this virus expresses 17 proteins (see Supplemental file 1 for more details). If we consider the Pfam annotations (Mistry *et al.*, 2021, http://pfam-legacy.xfam.org/) of the proteins expressed by this virus, we observe that *i/* 4 of these 17 proteins are not Pfam annotated, *ii/* the other 13 proteins are annotated by a set of 40 domains, 39 of which are strictly associated with viruses (the Macro domain being an exception to the rule). This clearly shows that SARS-CoV-2 domains are mostly similar to viral domains (97.5% ((39/40)*100) which are generally poorly annotated.

These results can be interpreted in two different (but complementary) ways:

1./ This virus, like many viruses, essentially contains virus-like proteins that are only present in viruses and not elsewhere,

2./ As it has been established that *i/* some viral proteins show similarities to some proteins of their host and that *ii/* this "molecular mimicry" is increasingly recognised (Elde & Malik, 2009), this lack of homologies outside of viral taxa can also be seen, at least in part, as a consequence of weaknesses in annotation methods.

It has been shown that HMMs stored in Pfam can lack sensitivity when searching for domains in "divergent organisms" (where the relevant signals become too weak to be identified (Terrapon *et al.*, 2012)). We thus decided here to explore the second way. We naturally turned to HHpred which is known to be an efficient tool for remote protein homology detection and can be easily used via a fast server (Gabler *et al.*, 2020). HHpred offers many possibilities such as searching for homologs among all proteins in an organism. HHpred is based on HHsearch and HHblits, which perform pairwise comparison of HMM profiles. Given their proven efficiency, HHsearch and HHblits have been used for some years to annotate viruses, and in particular Coronaviruses (Forni *et al.*, 2022). They have obviously also been used to annotate proteins expressed by SARS-CoV-2 (O'Donoghue *et al.*, 2021). However, the two previous works limited the homology search to viral proteins. For our part, we focused on searching for homologs in human.

To avoid "false positive predictions" as much as possible, we designed a procedure, mainly based on two ideas suggested in (Gabler *et al.*, 2020) but not implemented, to assess the robustness of HHpred results. Using HHpred and this procedure, we detected 6 robust similarities.

Materials and Methods

SARS-CoV-2 protein sequences

The 17 proteins studied in this article were extracted from UniProt (https://www.uniprot.org/, UniProt Consortium, 2021). UniProt provides polyproteins 1a (pp1a) and 1ab (pp1ab) as two separate entries. The pp1ab polyprotein is cleaved to form 15 shorter proteins; the first 10 proteins, *i.e.*, NSPs 1-10, are also cleaved from pp1a; NSPs 12-16 are unique to pp1ab. The list of proteins is given below. For each protein, we give its "Recommended Name", its "Short Name", its "AC - Uniprot ID", and its length:

Replicase polyprotein 1a / pp1a / P0DTC1 - R1A_SARS2 / Length 4,405

Replicase polyprotein 1ab / pp1ab / P0DTD1 - R1AB_SARS2 / Length 7,096

Envelope small membrane protein / E; sM protein / P0DTC4 - VEMP_SARS2 / Length 75

Membrane protein / M / P0DTC5 - VME1_SARS2 / Length 222

Nucleoprotein / N / P0DTC9 - NCAP_SARS2 / Length 419

Spike glycoprotein/ S glycoprotein / P0DTC2 - SPIKE_SARS2 / Length 1,273
ORF3a protein/ ORF3a / P0DTC3 - AP3A_SARS2 / Length 275
ORF3c protein / ORF3c / P0DTG1 - ORF3C_SARS2 / Length 41
ORF6 protein / ORF6 / P0DTC6 - NS6_SARS2 / Length 61
ORF7a protein / ORF7a / P0DTC7 - NS7A_SARS2 / Length 121
ORF7b protein / ORF7b / P0DTD8 - NS7B_SARS2 / Length 43
ORF8 protein / ORF8 / P0DTC8 - NS8_SARS2 / Length 121
ORF9b protein / ORF9b / P0DTD2 - ORF9B_SARS2 / Length 97
Putative ORF3b protein/ ORF3b / P0DTF1 - ORF3B_SARS2 / Length 22
Putative ORF3d protein/ _ / P0DTG0 - ORF3D_SARS2 / Length 57
Putative ORF9c protein / ORF9c / P0DTD3 - ORF9C_SARS2 / Length 73
Putative ORF10 protein / ORF10 / A0A663DJA2 - ORF10_SARS2 / Length 38

Sequence similarity searches
For remote homology detection, we used HHpred (Gabler *et al.*, 2020). First, starting from single
sequences or multiple sequence alignments (MSAs), it transforms them into a query HMM; using this HMM,
it then searches the Uniclust database30 and adds significantly similar sequences found to the query MSA for
the next search iteration. This strategy is very effective in detecting remotely homologous sequences but, as
the user guide points out (https://github.com/soedinglab/hh-suite/wiki), "the higher the number of search
iterations, the greater the risk of non-homologous sequences or sequence segments entering the MSA and
recruiting other sequences of the same type in subsequent iterations". To avoid this problem, we set the
number of iterations to 0, *i.e.* the parameter "MSA generation iterations" was set to 0. The default settings
were used for the other parameters. Note that we also briefly present in the Results section the HHpred
results obtained using the default setting for "MSA generation iterations", *i.e.* 3 (iterations).
Finally, it is important to note here that we use HHpred to look for similarities independently of the
mechanisms underlying these similarities, *i.e.* homologies, horizontal transfers (e.g. obtained by
"recombination" between SARS-CoV-2 and its current host, between ancestors of SARS-CoV-2 and their
hosts, between SARS-CoV-2 and another virus, *etc*.), convergent evolutions, *etc*.

Procedure for assessing the robustness of HHpred results
According to (Gabler *et al.*, 2020), when the reported probability value for a hit is greater than 95%,
homology is highly probable. Since viral and human proteins are being compared here, it can be assumed
that the 95% threshold is too high to detect similarities. In order to be more sensitive, while controlling
specificity (*i.e.* avoiding "false positive predictions"), we have devised a procedure that we describe below. Its
purpose is to assess the robustness of the results provided by HHpred. It is based on two ideas described in
(Gabler *et al.*, 2020) (section "Understanding Results") but not taken into account in HHpred.
This procedure is divided into 4 steps. From an algorithmic point of view, this procedure can be described
as a "gready search algorithm". It is performed for each protein expressed by the SARS-CoV-2 virus (see
Figure) (note that this was done by hand, as there were few data to process):
1./ For a given SARS-CoV-2 protein, hereafter referred to as "query", HHpred is run (using the default
parameters, except for the "MSA generation iterations" parameter which we set to 0, see section above) on
the *Homo sapiens* proteome of HHpred.
2.1/ The examination of the results provided by HHpred starts with the probability threshold of 0.95. Hits
with a probability greater than or equal to 0.95 are selected. If no hits meet this constraint, the threshold is
successively lowered to 0.9, 0.85 and finally to 0.80. As soon as a threshold satisfies the constraint (*i.e.* there
is at least one hit with a probability greater than or equal to the threshold), all hits above the threshold are

selected. If no threshold satisfies the constraint, we consider that no similarity between the query and the
human proteins can be detected.

2.2./ All previously selected hits are collected in a list and ranked from highest to lowest probability. The
best hit is then used as a seed to build a family of hits as follows: hits located at the same position as this best
hit on the query sequence and of similar size to it (+-5 amino acids for a best hit of length < 150, and +-15 for
a best hit of length > 150) feed the family under construction and are removed from the list ; hits that
overlap the seed are also removed from the list. The highest hit in the updated list is used as the "new" seed
and the process continues until the list is empty. As it is possible for a protein to have only one homolog in
human, families of singletons are not excluded.

3.1/ The query is then run on four HHpred proteomes, called "test" proteomes, corresponding to the
following four species: *Arabidopsis thaliana*, *Drosophila melanogaster*, *Escherichia coli* and *Haloferax volcanii*
(an archaea).

3.2/ For each species, the following step is performed:

First, the hits whose probability is greater than or equal to the previously selected threshold (see 2.1) are
selected. Then, depending on their size and location on the query sequence, they are assigned, if possible, to
a previously built family (see 2.2).

At the end of step 3, a family is thus made up of hits belonging at least to *Homo sapiens* and possibly to
Arabidopsis thaliana, *Drosophila melanogaster*, *Escherichia coli* or *Haloferax volcanii*. If a family includes only
human proteins, the robustness assumption can neither be rejected nor established. In this case, the
threshold is lowered and step 2 is performed again.

4./ For each family, InterPro annotations (Blum *et al.*, 2020) of proteins associated with hits are collected
and inspected manually (in particular the part of these proteins that corresponds to the hits). If the
annotations of the human proteins are similar to each other and to all proteins from at least one other
organism, this family/similarity is considered "robust"; these annotations are then associated with the
corresponding part of the viral protein (the query) ; if not, no similarities can be identified, and we consider
that no similarity between the query and the human proteins can be detected.

It should be noted that when the threshold of 0.8 is reached and it is not possible to reject or establish
the robustness hypothesis, an in-depth examination of the results is carried out by relaxing the constraints *i/*
on the probability threshold, which is then set to 0.5 (in accordance with the HHpred documentation which
states that "typically, a match should be seriously considered if it has a probability value >50%")), and *ii/* on
the size and location of hits ; the annotations of the proteins found by relaxing the constraints are then
examined; if at least 90% of human proteins are similarly annotated and these are also similarly annotated to
100% of the proteins of at least one other organism, this family/similarity is considered "robust"; these
annotations are then associated with the corresponding part of the viral protein (the query).

The similarities identified at the 0.95 and 0.9 probability levels will be labeled by "highly robust"; the
similarities identified at the 0.85 and 0.8 probability levels will be labeled by "very robust"; finally, the
similarities identified during the relaxation stage of constraints will be labeled by "quite robust".

Note: only proteins beginning with the prefix NP are considered in the analysis. XP records (proteins) are
not curated and are therefore not considered here; furthermore, proteins identified by HHpred that do not
have a match in "UniProtKB reviewed (Swiss-Prot)" (name and size in amino acids) were not considered
either.

181   **Results**

In the following sections, we present the main results of our study, *i.e.*, the list of the 6 robust similarities
we have identified. For reasons of clarity, for each family/similarity considered here, only the best hit in each
organism is provided. All results can be found in Supplemental file 2 (this file contains a condensed version of

the results produced by HHpred which are enriched by the InterPro annotations). The raw HHpred results are
stored in a separate gzip file called Supplemental file 4.

Note that the Pfam annotations of the proteins come from the InterPro or Pfam legacy (http://pfam-
legacy.xfam.org/) websites; the two sites generally give similar predictions; however, the domain boundaries
may sometimes differ very slightly.

NSP2 harbors a "Casein kinase II regulatory subunit" domain (very robust similarity)

NSP2 is derived from polyprotein 1a (181-818). The length of this protein is 638 A.A.

At the 0.85 probability threshold, only one human protein shares similarity with NSP2. Specifically, the
151-195 part of NSP2 is similar to the 101-142 part of the human protein "Casein kinase II subunit beta"
(CSK2B_HUMAN/NP_001311, length = 215 A.A.). The 109-140 part of the human "Casein kinase II subunit
beta" protein is annotated with the PROSITE "Casein kinase II regulatory subunit signature" motif ; in
addition, parts 105-126 and 127-148 of this protein are annotated with the PRINTS motif "CASNKINASEII".
This suggests that the 151-195 part of NSP2 could also be a "casein kinase II regulatory subunit signature".
Note that when the "MSA generation iterations" parameter is set to 3 (default setting), no significant results
are obtained (the probability of the best hit is 0.32).

For the given threshold of 0.85, this part of NSP2 is also similar to a part of an *Arabidopsis thaliana*
protein. Specifically, the 151-194 part of NSP2 is similar to the 182-222 part of the *Arabidopsis thaliana*
protein "Casein kinase II subunit beta" (CSK2D_ARATH/NP_191584.1, length = 276 AA). The 190-221 part of
the latter is annotated with the PROSITE motif "Casein kinase II regulatory subunit signature" ; in addition,
parts 186-207 and 208-229 of this protein are annotated with the PRINTS motif "CASNKINASEII".

This strongly suggests that NSP2 carries a "regulatory subunit signature of casein kinase II".

NSP3 harbors a Macro domain (highly robust similarity)

NSP3 is derived from polyprotein 1a (819-2763). The length of this protein is 1945 A.A.

For the 0.95 probability threshold, 7 human proteins share similarity with NSP3. The best match is the
human "Core histone macro-H2A.2" protein (H2AW_HUMAN/NP_061119, length = 372) whose 187-371 part
is similar to the 210-377 part of NSP3, *i.e.* the 1029-1197 part of polyprotein 1a. The 187-371 region of this
human protein contains the Pfam Macro domain (216-329). This suggests that the 210-377 part of NSP13
also shares similarity with the Macro domain. Note that when the "MSA generation iterations" parameter is
set to 3 (default setting), similar results are obtained.

For the probability threshold considered (*i.e.*, 0.95), one *Escherichia coli* protein shares similarity with
NSP3: the "O-acetyl-ADP-ribose deacetylase" protein (YMDB_ECOLI/NP_415563, length = 177) whose 3-166
part is similar to the 218-367 part of NSP3; the 218-367 region of this bacterial protein contains the Pfam
Macro domain (21-137).

The above strongly suggests that NSP3 hosts a Macro domain.

NSP13 harbors AAA domains (highly robust similarity)

NSP13 is derived from polyprotein 1ab (5325-5925). The length of this protein is 601 A.A.

At the 0.95 probability level, many human proteins share similarity with NSP13. The best match is the
human "DNA-binding protein SMUBP-2" (SMBP2_HUMAN/NP_002171, length = 993) whose 207-618 part is
similar to the 275-582 part of NSP13. The 207-618 part of this human protein is involved in two Pfam
domains, namely AAA_11/191-411 and AAA_12/418-615, which are both members of the P-loop NTPase clan
(CL0023). This suggests that the 275-582 part of NSP13, *i.e.* the 5600-5907 part of polyprotein 1ab, harbours
AAA domains. Note that when the "MSA generation iterations" parameter is set to 3 (default setting), similar
results are obtained.

The previously considered part of NSP13 is similar to three *Arabidopsis thaliana* proteins. The best match
is the *Arabidopsis thaliana* "probable helicase" protein (MAA3_ARATH/NP_001329005, length = 818) whose
273-734 part is similar to the 275-581 part of NSP13. The 273-734 part of this plant protein is involved in
three Pfam domains, namely AAA_11/257-436 + AAA_11/451-526 + AAA_12 /534-731, which are members
of the P-loop NTPase clan (CL0023). This result is in agreement with what has been found in human.
Our results strongly suggest that the 275-582 part of NSP13 hosts AAA domains.

234   NSP16 is a methyltransferase (highly robust similarity)
NSP16 is derived from polyprotein 1ab (6799-7096). The length of this protein is 298 A.A.
At the 0.95 probability level, two human proteins share similarity with NSP13. The best match is the "pre-
rRNA 2'-O-ribose RNA methyltransferase FTSJ3" protein (SPB1_HUMAN/NP_060117, length = 847). Its 31-217
part is similar to the 46-230 part of NSP16, *i.e.* the 6845-7029 part of polyprotein 1ab. The 31-217 part of this
human protein corresponds quite well to the Pfam "FtsJ-like methyltransferase" domain, FtsJ/21-207. Note
that when the "MSA generation iterations" parameter is set to 3 (default setting), similar results are
obtained.
The part of NSP16 considered above is similar to two *Drosophila melanogaster* proteins. The best match
is the fly protein "Putative tRNA (cytidine(32)/guanosine(34)-2'-O)-methyltransferase 1"
(TRM71_DROME/NP_650590, length = 302) whose 28-211 part is similar to the 46-215 part of NSP16. The
28-211 part of this fly protein corresponds quite well to the Pfam "FtsJ-like methyltransferase" domain,
FtsJ/21-207. This result is in agreement with what was found in human.
This strongly suggests that NSP16 is a methyltransferase.

248   Spike S harbors a part of a "Prominin domain" (highly robust similarity)
The length of this protein is 1273 A.A.
At the 0.90 probability level, 2 human proteins share similarity with Spike S (prominin-1 and prominin-2
proteins). The best match is human prominin-1 (PROM1_HUMAN/NP_006008, length = 865). Its 186-482 part
is similar to the 908-1254 part of Spike S; the 186-482 part of this human protein is included in the Pfam
"Prominin" domain, Prominin/19-820. Note that when the "MSA generation iterations" parameter is set to 3
(default setting), similar results are obtained.
For the given threshold of 0.90, one fly protein annotated with the Prominin domain of Pfam shares
similarities with Spike S: the fly protein "Prominin-like protein" (PROML_DROME/NP_001261351.1, length =
1013) whose 235-534 part is similar to the 911-1254 part of Spike S; the 235-534 part of this fly protein is
included in the "Prominin" domain of Pfam, Prominin/76-881.
This strongly suggests that Spike S hosts part of the "Prominin domain".

260   ORF3a has similarities with some "G Protein-Coupled Receptors" (quite robust similarity)
The length of this protein is 275 A.A.
At the 0.80 probability level, a human protein shares similarity with ORF3a, the human "lutein-
choriogonadotropic hormone receptor" (LSHR_HUMAN/NP_000224, length = 699). Its 537-693 part is similar
to the 41-183 part of ORF3a. A large part of this 537-693 region is included in the "7 transmembrane
receptor (rhodopsin family)" Pfam domain, *i.e.* 7tm_1/376-623. Note that when the "MSA generation
iterations" parameter is set to 3 (default setting), no significant results are obtained (the probability of the
best hit is 0.66).
For the given threshold of 0.80, no similarity is detected with proteins belonging to the 4 "test"
proteomes. However, a number of factors support this similarity when certain constraints are relaxed (see
Materials and methods):

Looking at the list of hits found by HHpred between ORF3a and the human proteome (see Supplemental
file 2), it is immediately obvious that the vast majority of human proteins found are G Protein-Coupled
Receptors (GPCRs). Indeed, it appears that out of 28 hits, 26 concern GPCRs (26/28 = 0.928), while the other
two correspond to transmembrane segments of proteins that are not linked to GPCRs.
Considering the fly proteome and applying the same methodology as previously used in human, it
appears that out of 3 hits, 3 concern GPCRs (see Supplemental file 2).
Overall (see Materials and Methods), this suggests that the similarity found is quite robust and that
ORF3a shares similarities with human GPCRs.

279   **Discussion**

The documented loss of sensitivity of Pfam HMMs when searching for domains in "divergent organisms"
(Terrapon *et al.*, 2012) prompted us to use HHpred (Gabler *et al.*, 2020) to annotate SARS-CoV-2 proteins.
Given a query sequence, this annotation tool offers the possibility to search for homologs among all proteins
in an organism. Each protein in the organism is represented by an HMM built according to a different
strategy than that used by Pfam (for more details, see the section "Creating custom databases" in the user
guide (https://github.com/soedinglab/hh-suite/wiki)). We speculated that this difference might give HHpred
the ability to discover similarities not detectable by Pfam (it should be noted that a theoretical comparison
between the Pfam and HHpred HMMs, as well as a full empirical comparison, is beyond the scope of this
paper).
To avoid as much as possible false predictions when using HHpred, we decided to disable its first step
which is based on an iterative search strategy. Indeed, the greater the number of search iterations, the
greater the risk of recruiting non-homologous sequences in the following iterations (see Materials and
Methods). Furthermore, in addition to the probability assigned by HHpred to each hit, we decided to
evaluate the robustness of these latter. Our evaluation procedure is based on two unimplemented ideas
described in (Gabler *et al.*, 2020) and can be summarized as follows (see Materials and Methods for more
details ; see also Figure):
A probability threshold is set; the starting value is 0.95 (according to (Gabler *et al.*, 2020), when the
probability of a hit is greater than 95%, homology is highly probable). Each viral protein ("query" sequence) is
compared to the human proteome using HHpred; all hits with a probability above the chosen threshold are
selected (if no hit meets this criterion, the threshold is successively lowered to 0.9, 0.85 and 0.80) ; if all hits
of similar size located at the same position on the query sequence (*i.e.*, a family of homologous hits) are
annotated with the same InterPro domain (Blum *et al.*, 2020), their probability of actually being homologous
to the query is very high ("Check relationships among top hits", first idea from (Gabler et al., 2020)); the
query is then run on a set of "test" proteomes to check whether similarly annotated homologous hits are
returned ("Check if you can reproduce the results with other parameters", second idea of (Gabler *et al.*,
2020)); if so, a family of homologous hits defined a "robust similarity"; if not, we consider that no similarities
can be identified. Note that when a family includes only human proteins, the robustness assumption can
neither be rejected nor established; in this case, the threshold is lowered and the study is carried out again. It
should be also noted that when the threshold of 0.8 is reached and it is not possible to reject or establish the
robustness hypothesis, a thorough examination of the results is carried out by relaxing the constraints
(mainly on the size, location and/or probability associated with the hits, see Materials and Methods for more
details). Similarities identified at the 0.95 and 0.9 probability levels are labeled "highly robust"; similarities
identified at the 0.85 and 0.8 probability levels are labeled "very robust"; finally, the similarities identified
when certain constraints are relaxed are described as "quite robust".
The organisms used to evaluate the HHpred results are *Arabidopsis thaliana*, *Drosophila melanogaster*,
Escherichia coli and *Haloferax volcanii* (an archaea). Note that, in order to potentially increase the identified

similarities, we would have liked to include proteomes from organisms closer to humans in our study.
Unfortunately, the online server currently does not offer the option to use such proteomes. To successfully
accomplish this task, it is necessary to perform the local installation of the free HH-suite software and build
these proteomes using this software. This work needs to be done (future works).
Below we present a summary of our results.
We subjected the 17 proteins of the SARS-CoV-2 proteome (see Materials & Methods and Results
sections) to our annotation procedure. UniProt considers polyproteins 1a (pp1a) and 1ab (pp1ab) as two
separate entries; polyprotein pp1ab is proteolytically cleaved to form 15 shorter proteins; the first 10
proteins (NSP1, ..., NSP10) are also cleaved from pp1a; NSP12, ..., NSP16 are unique to pp1ab. We therefore
subjected 30 proteins to our evaluation procedure.

No "robust" similarities were found for the following 24 proteins
NSP1, NSP4-10, NSP12, NSP14-15, Nucleoprotein, Envelope small membrane, Membrane Protein M,
ORF3B, ORF3C, ORF3D, ORF6, ORF7a, ORF7b, ORF8, ORF9b, ORF9C, ORF10.

A "highly robust" or "very robust" similarity, already documented in literature, was detected on the
following 4 proteins
NSP3 harbors a Macro domain; NSP13 harbors AAA domains; NSP16 is a methyltransferase. As these
similarities are well documented and widely discussed, the interested reader is invited to consult the InterPro
annotations.
In a more interesting manner, we have shown that part 151-195 of NSP2, *i.e.* part 332-376 of polyprotein
1a, contains a "signature of the beta subunit of casein kinase II". According to PROSITE, such a domain could
be involved in the binding of a metal such as zinc. Interestingly, the structure of the N-terminal part of NSP2
was recently solved (Ma *et al.*, 2021). It shows that NSP2 has three zinc fingers: Zn1, Zn2 and Zn3. Two Zn2
(resp. Zn3) binding sites are located at positions 161 and 164 (resp. at positions 190 and 193). Our prediction
is therefore in agreement with this structure of the N-terminal domain of SARS-CoV-2 NSP2.

A previously unknown "highly robust" similarity was detected on Spike S protein
The Spike S protein (1273 A.A.) is composed of two subunits: the S1 subunit (14-685 residues), and the S2
subunit (686-1273 residues), which are responsible for receptor binding and membrane fusion respectively
(Huang *et al.*, 2020). We have shown that the 908-1254 part of the Spike S protein is similar to the 186-482
part of human prominin-1 (length = 865). This similarity encompasses the heptapeptide repeat 1 sequence,
i.e. HR1 (912-984 residues), HR2 (1163-1213 residues), the TM domain (1213-1237) and part of the
cytoplasmic domain (1237-1273) of the S2 subunit; however, it excludes the fusion peptide (FP) (788-806) of
S2 which plays an essential role in mediating membrane fusion. HR1 and HR2, which are part of the
similarity, have been shown to form a six-helix bundle that is essential for the fusion and viral entry function
of the S2 subunit (Xia *et al.*, 2020).
Recently, in searching for proteins involved in SARS-CoV-2 entry into host cells, (Kotani *et al.*, 2022) found
that the glycoprotein CD133, the other name for prominin-1, colocalises with ACE2 – the main cell entry
receptor for SARS-CoV-2 – bound to the Spike S protein in Caco-2 cells. They demonstrated that the SARS-
CoV-2 Spike protein exhibited increased binding capacity in cells co-expressing ACE2 and CD133, compared
to cells expressing ACE2 alone. In addition, they experimentally infected HEK293T cells with a SARS-CoV-2
pseudovirus and showed that infectivity was twice as high in HEK293T cells co-expressing CD133-ACE2 than
in HEK293T cells expressing ACE2 alone. They concluded that CD133, although not a primary receptor for the
SARS-CoV-2 Spike protein, is a cofactor (a co-receptor) that partially contributes to infection in the expressing
cells. All these results suggest that the C-terminal part of Spike S, which has similarities with prominin-1, may

be involved in the docking of Spike S to ACE2 (insofar as CD133 enhances the ability of Spike S to bind to
ACE2). This obviously remains to be demonstrated but is clearly an interesting avenue of research.

While considerable work has been done to characterise the cellular receptors and pathways mediating
virus internalisation, little is known about the onset of the infection process, which begins when the virus
comes into contact with the host cell surface; some studies have shown that viruses "diffuse" onto the
surface of host cells after "landing" on them; this process ranges from a random walk to a constrained
diffusion where the virus particles appear to be confined to a specific microdomain of the cell membrane
(Boulant *et al.*, 2015). From this point of view, it is interesting to note that it was recently shown by (Rouaud
et al., 2022) that *i/* ACE2 concentrates at epithelial apical cell junctions in cultured epithelial cell lines, and
that *ii/* (Pinto *et al.*, 2022*)* showed that ACE2 and TMPRSS2 (which is used by SARS-CoV-2 for Spike S-protein
priming (Hoffmann *et al.*, 2020)) were localised at the plasma membrane, including the microvilli, in human
airway epithelium. Interestingly, about 25 years ago, prominin was shown to be localised to the apical
surface of various epithelial cells, where it is selectively associated with microvilli and microvillus-related
structures (Weigmann *et al.*, 1997). Furthermore, Weigmann and colleagues showed that prominin
expressed ectopically in non-epithelial cells was also selectively found in microvillus-like protrusions of the
plasma membrane. Two years later, (Corbeil *et al.*, 1999) showed that prominin contains dual targeting
information, for direct delivery to the apical domain of the plasma membrane and for enrichment in the
microvilli subdomain. Furthermore, they showed that this dual targeting does not require the cytoplasmic C-
terminal tail of prominin (*i.e.*, part 814-865 of CD133). From the above results, it is tempting to assume that
the prominin-like part of Spike S is involved in the delivery of the virus to the apical domain of the plasma
membrane where the ACE2 proteins are located. This hypothesis is all the more tempting as the similarity
between Spike S and prominin does not concern the C-terminal part of prominin, which, as we have pointed
out above, is not necessary for prominin targeting (recall that we have shown that the 186-482 part of
human prominin-1 is similar to the 908-1254 part of Spike S). Unfortunately, to date, the molecular nature of
the prominin apical sorting signal is unknown. It has been suggested in (Weigmann *et al.*, 1997) that
prominin may interact with the actin cytoskeleton, or that plasma membrane protrusions may have a specific
lipid composition/organisation for which prominins may have a preference.

Finally, it should be noted that the "SARS-CoV(-1)" glycoprotein Spike, which, like SARS-CoV-2 Spike,
binds to human ACE2 (Li *et al.*, 2003), is also similar to human prominin-1. Specifically, using HHpred, we
showed that the 177-473 part of the latter is similar to the 890-1236 part of Spike (with an associated
probability of 0.95 – see Supplemental file 4, raw HHpred data). In contrast, the MERS-CoV Spike
glycoprotein (like SARS-CoV and SARS-CoV-2, MERS-CoV is a betacoronavirus), which uses human DPP4 as an
entry receptor (Raj *et al.*, 2013), is similar to human mucin-1: the 292-421 part of mucin-1 is similar – with an
associated probability of 0.89 – to the 1230-1344 part of MERS-CoV Spike (see Supplemental file 4, raw
HHpred data). It is also interesting to note that (Kotani *et al.*, 2022) showed that the DPP4 protein also
colocalises with ACE2 and CD133 in Caco-2 cells. This suggests that it is likely that *i/* different coronaviruses
compete at the same positions on the cell, but *ii/* use different entry receptors and therefore different types
of spike proteins to reach these sites and fuse with the cells.

A previously unknown "quite robust" similarity was detected on ORF3a protein
The 41-183 part of ORF3a (275 A.A.) shows similarities to human G Protein-Coupled Receptors (GPCRs)
(which are cell surface receptor proteins that detect molecules from outside the cell and trigger cellular
responses (Lagerström & Schiöth, 2008)) and in particular to the GPCRs annotated with the Pfam domain "7
transmembrane receptor (rhodopsin family)/7tm_1" (see Results section and Supplemental file 2). According
to Pfam, this family contains, among other GPCRs, members of the opsin family, which are considered typical
members of the rhodopsin superfamily.

The ORF3a protein of "SARS-CoV(-1)" has been shown to form an ion channel (Lu *et al.*, 2006). Recently,
(Kern *et al.*, 2021) presented Cryo-EM determined structures of SARS-CoV-2 ORF3a at a resolution of 2.1Å.
The authors provide evidence suggesting that ORF3a forms a large polar cavity in the inner half of the
transmembrane region (TM) that could form ionic conduction paths (TM1 (43-61), TM2 (68-99) and TM3
(103-133)). Interestingly, the similarity we detected on ORF3a (41-183) encompasses the transmembrane
portion of ORF3a (43-133) which could form ionic permeation pathways. As mentioned earlier, we have
shown that this part of ORF3a resembles many GPCRs which belong to the Rhodopsin family (22 of 28 human
proteins sharing similarities with ORF3a, see Supplemental file 2 for more details). It is interesting to note
that some GPCRs, called "Rhodopsin channels", directly form ion channels (see (Nagel *et al.*, 2002) and
(Nagel *et al.*, 2003)). From this point of view, our prediction is therefore in line with the work of (Kern *et al.*,
2021).

However, it should be noted that if our method is applied to the ORF3a of SARS-CoV(-1), no similarities
are identified. More precisely, none of the similarities found by HHpred are significant, *i.e.* the probability of
the best hit is 0.72, which is below our threshold of 0.8; moreover, this best hit does not correspond to a
GPCR (see Supplemental file 4). This result may suggest a lack of sensitivity of HHpred. That said, although
HHpred is a fairly effective tool for detecting very distant homologies, not all similarities are detectable.
Furthermore, although the ORF3a of SARS-CoV(-1) and SARS-CoV-2 share 72% sequence identity and are
similar in the arrangement of the TM domains, the differences observed in the ion channel properties
between these two proteins suggest a different mode of action between them (Zhang *et al.*, 2022).

Autoantibodies targeting GPCRs have been found in patients with COVID-19 and Long-COVID-19. It is
therefore tempting to speculate that the similarity between ORF3a and certain human GPCRs could be the
cause of the autoimmune reactions observed. There is some evidence to support this hypothesis:

- Autoantibodies targeting GPCRs (and RAS-related molecules) have been shown to be associated with
the severity of COVID-19 (Cabral-Marques *et al.*, 2022). Among the anti-GPCR autoantibodies, the authors of
the latter paper identified the chemokine receptor CXCR3 and the RAS-related molecule AGTR1 as antibody
targets with the strongest association with disease severity. Strikingly, of the 26 GPCRs we identified as
sharing similarity with ORF3a (see Supplemental file 2), 5 are chemokine receptors, including the chemokine
receptor CXCR3,

- Functional autoantibodies against G protein-coupled receptors have been found in patients with
persistent symptoms of Long-COVID-19 (Wallukat *et al.*, 2021). In particular, the authors of the latter paper
identified functional autoantibodies against the M2 muscarinic receptor in the blood of Long-COVID patients.
Strinkgly, of the 26 GPCRs we identified as sharing similarity with ORF3a (see Supplemental file 2), 3 are
muscarinic receptors, including the muscarinic acetylcholine M2 receptor. In the same study, functional
autoantibodies against the alpha 1-adrenoceptor and the beta 2-adrenoceptor were also identified.
Interestingly, of the 26 above mentioned GPCRs, 3 are adrenoreceptors, namely alpha-1D, alpha-2A and
alpha-2C (see Supplemetal file 2).

To conclude this section, we would like to emphasize that our main goal is to identify similarities between
SARS-CoV-2 proteins and human proteins in order to gain a better understanding of the functions of SARS-
CoV-2 proteins, rather than seeking mimics that could trigger autoimmune processes. This problem is usually
solved by searching for n-mers, which is obviously not done here. (Khavinson *et al.*, 2021), for example,
specifically addresses this problem and concludes that ORF3a does not appear to be involved in triggering an
autoimmune response. Furthermore, based solely on the similarity between ORF3a and certain human
GPCRs targeted by autoantibodies in patients with COVID-19 and Long-COVID-19, it is difficult to state that
this similarity is the cause of the autoimmune phenomena observed. As Cabral-Marques *et al.* (2022) point
out, the mechanisms by which SARS-CoV-2 infection triggers the production of autoantibodies remain
unknown to this day; according to these authors, molecular mimicry between SARS-CoV-2 and certain human
proteins is obviously not the only hypothesis to explain these phenomena: a hyperinflammatory response

triggered by the virus could cause tissue damage, leading to systemic autoimmune reactions. However, our
results suggest that further studies should be conducted.

Comparison of our results with those of "Pfam clans"

As indicated in the introduction to this article (see also Supplemental file 1), of the 40 Pfam domains that
annotate SARS-CoV-2 proteins, only one domain is not confined to viruses, the Macro domain that annotates
NSP3. This observation can be modulated at the level of Pfam clans which are collections of related domains.
At this level, 12 domains belong to clans whose domains are not strictly viral (see Supplemental file 1). These
clans allow the annotation of the following 9 proteins (more generally, of only part of each protein): NSP3,
NSP5, NSP13, NSP14, NSP15, NSP16, ORF7a, ORF8, and Spike S. 4 of these proteins are annotated by both
Pfam and our approach: NSP3, NSP13, NSP16 and Spike S. In the case of NSP3, NSP13 and NSP16, the
annotations are similar (note however that for NSP3, Pfam detects two domains related to the MACRO clan;
only one Macro domain is detected by our approach) whereas in the case of Spike S, our annotations refer to
a different part of the protein than that annotated by Pfam. We also identified similarities, not restricted to
viruses unlike Pfam, for ORF3a and NSP2.

Evaluation of our results in light of the known weaknesses of HHpred

As reported in (Gabler *et al.*, 2020) and (Kuchibhatla *et al.*, 2014), some false positive HHpred hits may
have high scores because they have coiled-coil, transmembrane or low complexity segments. Of our 6
"robust similarities", 2 have transmembrane segments and/or disordered areas (according to InterPro
annotations).

ORF3a

As previously indicated, ORF3a shares similarity with G Protein-Coupled Receptors (GPCRs) annotated
with the Pfam domains "7 transmembrane receptor (rhodopsin family)/7tm_1" or "7 transmembrane
receptor (secretin-like) 7tm_2" (see Results or Supplemental file 2).
Since transmembrane proteins are a large family of proteins – according to UniProt, out of 80581
proteins expressed by humans, 13876 are transmembrane proteins – it is legitimate to ask whether the
(observed) distribution of transmembrane proteins found by HHpred – out of 28 proteins found by HHpred,
28 are transmembrane proteins – is the same as the (expected) distribution of transmembrane proteins in
UniProt. Using a Fisher's exact test, we conclude (see Supplemental file 3 for proof) that the results found by
HHpred are not randomly drawn from the UniProt human proteome (p-value = 6.2059249716913E-11).
Similarly, as transmembrane proteins can be grouped into many different classes (the Pfam clan "Family
A G protein-coupled receptor-like superfamily", to which 7tm_1 and 7tm_2 belong, alone contains 53
different domains), it can also be argued that the similarities found by HHpred are due to chance. Of the 28
transmembrane proteins found by HHpred, 26 belong to the 7tm_1 or 7tm_2 classes. Knowing that the
number of human proteins belonging to the 7tm_1 or 7tm_2 classes is – according to UniProt – 540, we
show (see Supplemental file 3 for proof) using a Fisher's exact test that the results obtained by HHpred do
not arise from random selection within the different classes of the transmembrane protein family (p-value =
2.8739559680731E-12).

Spike glycoprotein

As shown previously, the 908-1254 part of the Spike S protein of SARS-CoV-2 is similar to the 186-482
part of human prominin-1. The 179-432 part of this prominin is annotated as
"NON_CYTOPLASMIC_DOMAIN" (*i.e.* non-cytoplasmic loops of a TM protein) by Phobius (for completeness,
note that the 253-283 part is annotated as a coil by COILS).

In contrast to the case of ORF3a, no reliable statistical test can be performed here (the number of human prominins, *i.e.* proteins annotated by Pfam as "prominin" (Pfam PF05478), is 5). However, such a calculation seems unnecessary here. HHpred identified a similarity between Spike S and human and fly prominins (see Results section). Human and fly belonging to lineages that were separated over 700 million years ago (median time of divergence 694 MYA (see http://timetree.org/, (Kumar et al., 2017)), this similarity is clearly not a coincidence (unless one imagines a recent horizontal transfer).

Conclusion

We used HHpred to search for similarities between SARS-Cov-2 and human proteins. To avoid false predictions, the robustness of each similarity was assessed using a procedure based on "test sets/proteomes". We found six robust similarities in six different proteins, of which three are already documented, one is in agreement with recent crystallographic results, and two are not reported in the literature. We focused on these last two similarities and showed how they open new avenues of research to better understand this virus. Obviously, our work is limited to making predictions that need to be validated experimentally. Furthermore, the origin of the similarities (evolutionary convergence, horizontal transfer, etc.) has not been addressed in this work. Nevertheless, we believe that our approach (or one similar to it) can be profitably used to open up lines of research and to improve the annotation of any virus, especially "orphan viruses", *i.e.* viruses which, for various reasons, are far much less studied than SARS-CoV-2.

Acknowledgements

We would like to thank all those who initiated Pfam, InterPro and HH-suit and all those who maintain and improve these databases and tools for the benefit of our community. Without them, the work presented here would not have been possible. We also warmly thank Joël Pothier (ISYEB, MNHN) whose comments have clearly contributed to improve the quality of this paper.

Supplementary information availability

Supplemental files 1, 2, 3 and 4 are available online (*cf*. bioRxiv, DOI of this document).

Conflict of interest disclosure

The author declares that he complies with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

References

Boulant S., Stanifer M., Lozach P-Y. Dynamics of virus-receptor interactions in virus binding, signaling, and endocytosis. Viruses. 2015 Jun 2;7(6):2794-815. https://doi.org/10.3390/v7062747

Blum M., Chang H., Chuguransky S., Grego T., Kandasaamy S., Mitchell A., Nuka G., Paysan-Lafosse T., Qureshi M., Raj S., Richardson L., Salazar G. A., Williams L., Bork P., Bridge A., Gough J., Haft D. H., Letunic I., Marchler-Bauer A., Mi H., Natale D. A., Necci M., Orengo C. A., Pandurangan A. P., Rivoire C., Sigrist C. J. A., Sillitoe I., Thanki N., Thomas P. D., Tosatto S. C. E., Wu C. H., Bateman A. and Finn R. D. The InterPro protein families and domains database: 20 years on. Nucleic Acids Research, Nov 2020. https://doi.org/10.1093/nar/gkaa977

Cabral-Marques O., Halpert G., Schimke L. F., Ostrinski Y., Vojdani A., Baiocchi G. C., Freire P. P., Filgueiras I. S., Zyskind I., Lattin M. T., Tran F., Schreiber S., Marques A. H. C., Plaça D. R., Fonseca D. L. M., Humrich J. Y., Müller A., Giil L. M., Graßhoff H., Schumann A., Hackel A., Junker J., Meyer C., Ochs H. D., Lavi Y. B., Scheibenbogen C., Dechend R., Jurisica I., Schulze-Forster K., Silverberg J. I., Amital H., Zimmerman J., Heidecke H., Rosenberg A. Z., Riemekasten G., and Shoenfeld Y. Autoantibodies targeting GPCRs and RAS-related molecules associate with COVID-19 severity. Nature Communications. 2022 Mar 9;13(1):1220. https://doi.org/10.1038/s41467-022-28905-5

Corbeil D., Röper K., Hannah M. J., Hellwig A., Huttner W. B. Selective localization of the polytopic membrane protein prominin in microvilli of epithelial cells - a combination of apical sorting and retention in plasma membrane protrusions. *J Cell Sci* (1999) 112 (7): 1023–1033. https://doi.org/10.1242/jcs.112.7.1023

Elde N. C., Malik H. S. The evolutionary conundrum of pathogen mimicry. Nature Reviews Microbiology. 2009 Nov;7(11):787-97. https://doi.org/10.1038/nrmicro2222

Forni D., Cagliani R., Molteni C., Arrigoni F., Mozzi A., Clerici M., De Gioia L., Sironi M. Homology-based classification of accessory proteins in coronavirus genomes uncovers extremely dynamic evolution of gene content. Molecular Ecology. 2022 Jul;31(13):3672-3692. https://doi.org/10.1111/mec.16531

Gabler F., Nam S.-Z., Till S., Mirdita M., Steinegger M., Söding J., Lupas A. N., Alva V. Protein Sequence Analysis Using the MPI Bioinformatics Toolkit. Current Protocols in Bioinformatics. 2020 Dec;72(1). https://doi.org/10.1093/nar/gkl217

Hoffmann M., Kleine-Weber H., Schroeder S., Krüger N., Herrler T., Erichsen S., Schiergens T. S., Herrler G., Wu N.-H., Nitsche A., Müller M. A., Drosten C., Pöhlmann S. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. Cell. 2020 Apr 16;181(2):271-280. https://doi.org/10.1016/j.cell.2020.02.052

Huang Y., Yang C., Xu X.-F. , Xu W., Liu S.-W. Structural and functional properties of SARS-CoV-2 spike protein: potential antivirus drug development for COVID-19. Acta Pharmalogica Sinica. 2020 Sep;41(9):1141-1149. https://doi.org/10.1038/s41401-020-0485-4

Kern D. M., Sorum B.,Mali S. S., Hoel C. M., Sridharan S., Remis J. P., Toso D. B., Kotecha A., Bautista D. M., Brohawn S. G. Cryo-EM structure of SARS-CoV-2 ORF3a in lipid nanodiscs. Nat Struct Mol Bio. 2021 Jul;28(7):573-582. https://doi.org/10.1038/s41594-021-00619-0

Khavinson V., Terekhov A., Kormilets D., Maryanovich A. Homology between SARS CoV-2 and human proteins. Sci Rep. 2021 Aug 25;11(1):17199. https://doi.org/10.1038/s41598-021-96233-7

Kotani N., Nakano T., Kuwahara R. Host cell membrane proteins located near SARS-CoV-2 spike protein attachment sites are identified using proximity labeling and proteomic analysis. J Biol Chem. 2022 Nov;298(11):102500. https://doi.org/10.1016/j.jbc.2022.102500

Kuchibhatla D. B., Sherman W. A., Chung B. Y. W., Cook S., Schneider G., Eisenhaber B., Karlin D. G. Powerful sequence similarity search methods and in-depth manual analyses can identify remote homologs in many apparently "orphan" viral proteins. Journal of Virology, 2014 Jan; 88(1): 10–20. https://doi.org/10.1128/JVI.02595-13

Kumar S., Stecher G., Suleski M., Hedges S. B. TimeTree: A Resource for Timelines, Timetrees, and Divergence Times. Mol Biol Evol. 2017 Jul 1;34(7):1812-1819. https://doi.org/10.1093/molbev/msx116

Lagerström M. C., Schiöth H. B. Structural diversity of G protein-coupled receptors and significance for drug discovery. Nature Reviews Drug Discovery. 2008 Apr;7(4):339-57. https://doi.org/10.1038/nrd2518

Li W., Moore M. J., Vasilieva N., Sui J., Wong S. K., Berne M. A., Somasundaran M., Sullivan J., Luzuriaga K., Greenough T. C., Choe H., Farzan M. Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus. Nature. 2003 Nov 27;426(6965):450-4. https://doi.org/10.1038/nature02145

Lu W., Zheng B.J., Xu K., Schwarz W., Du L., Wong C.K., Chen J., Duan S., Deubel V., Sun B. Severe acute respiratory syndrome-associated coronavirus 3a protein forms an ion channel and modulates virus release. Proc. Natl. Acad. Sci. USA. 2006;103:12540–12545. https://doi.org/10.1073/pnas.0605402103

Ma J., Chen Y., Wu W., Chen Z. Structure and Function of N-Terminal Zinc Finger Domain of SARS-CoV-2
NSP2. Virologica sinica. 2021 Oct;36(5):1104-1112. https://doi.org/10.1007/s12250-021-00431-6

Mistry J., Chuguransky S., Williams L., Qureshi M., Salazar G. A., Sonnhammer E. L. L., Tosatto S. C. E., Paladin
L., Raj S., Richardson L. J., Finn R. D., Bateman A. Pfam: The protein families database in 2021. Nucleic
Acids Research. 2021 8 January Volume 49, Issue D1, Pages D412–D419.
https://doi.org/10.1093/nar/gkaa913

Nagel G., Ollig D., Fuhrmann M., Kateriya S.,Musti A. M., Bamberg E.,Hegemann P. Channelrhodopsin-1: a
light-gated proton channel in green algae. Science. 2002 Jun 28;296(5577):2395-8.
https://doi.org/10.1126/science.1072068

Nagel G., Szellas T., Huhn W., Kateriya S., Adeishvili N., Berthold P., Ollig D., Hegemann P., and Bamberg E.
Channelrhodopsin-2, a directly light-gated cation-selective membrane channel. Proc Natl Acad Sci U S A.
2003 Nov 25; 100(24): 13940–13945. https://doi.org/10.1073/pnas.1936192100

O'Donoghue S., Schafferhans A., Sikta N., Stolte C., Kaur S., Ho B. K., Anderson S., Procter J. B., Dallago C.,
Bordin N., Adcock M., Rost B. SARS-CoV-2 structural coverage map reveals viral protein assembly,
mimicry, and hijacking mechanisms. Molecular System Biology. 2021 Sep;17(9).
https://doi.org/10.15252/msb.202010079

Pinto A. L., Rai R. K., Brown J. C., Griffin P., Edgar J. R., Shah A., Singanayagam A., Hogg C., Barclay W. S.,
Futter C. E., Burgoyne T. Ultrastructural insight into SARS-CoV-2 entry and budding in human airway
epithelium. Nat Comm. 2022 Mar 25;13(1):1609. https://doi.org/10.1038/s41467-022-29255-y

Raj V. S., Mou H., Smits S. L., Dekkers D. H. W., Müller M. A., Dijkman R., Muth D., Demmers J. A. A., Zaki A.,
Fouchier Ron A. M., Thiel V., Drosten C., Rottier P. J. M., Osterhaus A. D. M. E., Bosch B. J., Haagmans B. L.
Dipeptidyl peptidase 4 is a functional receptor for the emerging human coronavirus-EMC. Nature.
2013;495:251–254. https://doi.org/10.1038/nature12005

Rouaud F., Méan I., Citi S. The ACE2 Receptor for Coronavirus Entry Is Localized at Apical Cell-Cell Junctions of
Epithelial Cells. Cells. 2022 Feb 11;11(4):627. https://doi.org/10.3390/cells11040627

Terrapon N. , Gascuel O., Maréchal E., Bréhélin L. Fitting hidden Markov models of protein domains to a
target species: application to *Plasmodium falciparum*. BMC Bioinformatics. 2012 May 1;13:67.
https://doi.org/10.1186/1471-2105-13-67

UniProt Consortium. UniProt: the universal protein knowledgebase in 2021. Nucleic Acids Research. 2021 Jan
8;49(D1):D480-D489. https://doi.org/10.1093/nar/gkaa1100

Wallukat G., Hohberger B., Wenzel K., Fürst J., Schulze-Rothe S., Wallukat A., Hönicke A.-S., Müller J.
Functional autoantibodies against G-protein coupled receptors in patients with persistent Long-COVID-19
symptoms. Journal of Translational Autoimmunity. 2021;4:100100.
https://doi.org/10.1016/j.jtauto.2021.100100

Weigmann A., Corbeil D., Hellwig A., Huttner W. B. Prominin, a novel microvilli-specific polytopic membrane
protein of the apical surface of epithelial cells, is targeted to plasmalemmal protrusions of non-epithelial
cells. PNAS. 1997 Nov 11;94(23):12425-30. https://doi.org/10.1073/pnas.94.23.12425

Xia S., Zhu Y., Liu M., Lan Q., Xu W., Wu Y., Ying T., Liu S., Shi Z., Jiang S., Lu L. Fusion mechanism of 2019-
nCoV and fusion inhibitors targeting HR1 domain in spike protein. Cell Mol Immunol. 2020 Jul;17(7):765-
767. https://doi.org/10.1038/s41423-020-0374-2

Zhang J., Ejikemeuwa A., Gerzanich V., Nasr M., Tang Q., Simard J.M., Zhao R.Y. Understanding the Role of
SARS-CoV-2 ORF3a in Viral Pathogenesis and COVID-19. Front Microbiol. 2022 Mar 9;13:854567.
https://doi.org/10.3389/fmicb.2022.854567.

Figure - Using HHpred, a viral protein is compared to the human proteome and to a set of other proteomes, called "test" proteomes, which include the fly proteome. The probability threshold was set at 0.9, so only hits with a probability value of 0.90 or greater are considered relevant here. 4 homologous hits (*i.e.*, hits of similar sizes and located at a similar position on the query sequence) exceeding the given threshold were found by HHpred (black boxes): 3 are found in humans and one in flies; the InterPro annotation of all the "black box" hits are the same (red oval); as the annotations of all these homologous hits are identical and at least one of these hits belongs to a test proteome, the corresponding family of homologous hits is considered to be a "robust/similar family"; this similarity will be used to annotate the corresponding hit on the viral protein.

664