1   **Using a *P. falciparum* genetic cross to dissect the relative contributions of *pfcrt* and**
2   ***plasmepsin II/III* to piperaquine response-related traits**
3

4   John Kane[1], Xue Li[2], Sudhir Kumar[3], Katrina A. Button-Simons[1], Katelyn Vendrely Brenneman[4],
5   Haley Dahlhoff[1], Mackenzie A.C. Sievert[1], Lisa A. Checkley[1], Douglas A. Shoue[1], Puspendra P.
6   Singh[1], Meseret T. Haile[3], [2]Ann Reyes, Ian H. Cheeseman[5], Ashley M. Vaughan[3,6], Timothy J.
7   C. Anderson[2], & Michael T. Ferdig[1]*.
8
[1]Eck Institute for Global Health, Department of Biological Sciences, University of Notre Dame,
Notre Dame, IN, USA. [2]Disease Intervention and Prevention Program, Texas Biomedical
Research Institute, San Antonio, TX, USA. [3]Center for Global Infectious Disease Research,
Seattle Children's Research Institute, Seattle, WA, USA. [4]Harvard T.H. Chan School of Public
Health, Boston, MA, USA. [5]Host Pathogen Interactions Program, Texas Biomedical Research
Institute, San Antonio, TX, USA. [6]Department of Pediatrics, University of Washington, Seattle,
WA, USA.
16
17   * Corresponding Author

18
<u>Abstract</u>

Piperaquine (PPQ) is widely used in combination with dihydroartemisinin (DHA) as a first-line

treatment against malaria parasites. Resistance to this artemisinin combination therapy (ACT)

has emerged in Cambodia and threatens to spread to other malaria-endemic regions. Multiple

drivers of PPQ resistance have been reported, including mutations in the *Plasmodium*

falciparum chloroquine resistance transporter (*pfcrt*) and increased copies of *plasmepsin II/III*

(*pm*). Experimental genetic crosses allow for the precise quantification of the relative contributions

of specific loci to a phenotype and have the distinct advantage of assaying the effects of naturally

evolved genetic variation. We generated a cross between a Cambodia-derived multi-drug

resistant KEL1/PLA1 lineage parasite isolate (KH004) and a drug susceptible parasite isolated in

Malawi (Mal31). Mal31 harbors a wild-type (3D7-like*)* *pfcrt* allele and a single copy of *pm,* while

KH004 has a chloroquine-resistant (Dd2-like*)* *pfcrt* allele with an additional G367C substitution

and four copies of *pm*. We cloned 101 unique recombinant progeny from this cross, and from this

pool, we identified a targeted set of progeny representing all possible combinations of variants *at*

pfcrt and pm for detailed analysis of a range of PPQ response and competitive fitness phenotypes.

Fortuitously, progeny clones that inherited the KH004 *pm* allele displayed a range of different

copies, allowing us to directly test the role *of pm* copy number on a range of resistance

phenotypes, including IC50, area under the curve (AUC), and PSA as well as competitive fitness.

We find that all progeny inheriting both *pfcrt* and *pm* alleles from KH004 are PPQ resistant by

(PSA and AUC) and that inheritance of KH004 *pfcrt* is required for resistance (including progeny

that inherited the single copy *pm* from Mal31). Our findings clearly display how progeny clones

from experimental crosses can be used to parse apart the contributions *of pfcrt, pm,* and parasite

genetic background*,* to a range of PPQ-related traits and confirm the critical role of the PfCRT

G367C substitution on PPQ resistance.

<u>Introduction</u>

Malaria is a life-threatening parasitic disease that nearly half of the world's population is

at risk of contracting (1). The World Health Organization estimates that in 2020 there were 247

million malaria cases and 619,000 deaths reported across 84 countries, with 76% of those deaths

occurring in children under the age of 5 (1). The continued evolution of drug-resistant *Plasmodium*

falciparum asexual blood-stage parasites has been one of the major obstacles to global malaria

elimination and eradication efforts (1). Current frontline treatments for uncomplicated *P.*

falciparum infection are artemisinin-based combination therapies (ACT), which are composed of

a fast-acting artemisinin (ART) derivative paired with one or more longer-lasting partner drugs (2).

ACTs began to replace historically efficient antimalarials, such as chloroquine (CQ) and

sulfadoxine-pyrimethamine (SP), as first-line therapies in the early 2000s after resistance to these

two therapies swept throughout Africa (3–5). The WHO has recommended ACTs for

uncomplicated malaria since 2006 due to their greater efficacy and decreased resistance

compared to other single-drug therapeutics, with the most common ACTs used in Southeast Asia

(SEA) being dihydroartemisinin/piperaquine (DHA+PPQ), artemether/lumefantrine (AL), and

artesunate/mefloquine (AS+MQ). The ACT combination DHA-PPQ has been most widely used

throughout the Greater Mekong Subregion (GMS) of SEA which has historically been a hotbed

for antimalarial drug resistance (6–8).

The spread of resistance to DHA+PPQ is largely attributed to the spread of the rapidly

expanding, multidrug-resistant KEL1/PLA1 lineage of parasites (9–11). This lineage of "superbug"

parasites emerged out of western Cambodia in 2008 and has since spread to Vietnam, Laos, and

northeastern Thailand in subsequent years (10). KEL1/PLA1 parasites are characterized by

mutations in *kelch13* (KEL1), specifically the C580Y substitution which has been strongly

associated with ART resistance, as well as a copy number (CN) amplification of the two aspartic

protease genes *plasmepsin II/III* (PLA1), which has been linked with decreased sensitivity to PPQ

(9, 10, 12, 13). The decreased sensitivity to ART derivatives results in a greater remaining parasite

load which has to be cleared by the partner drug, leading to an increased chance of ACT failure

(9, 14–17).

After the original detection of PPQ resistance in Cambodia in 2015, independent genome-

wide association studies identified that variation in *plasmepsin II/III* (*pm*) was associated with

PPQ-R parasites (16, 18–20). The proposed mechanism supporting the association between *pm*

amplification and PPQ-R is that plasmepsins reduce concentrations of reactive heme in the

parasite digestive vacuole (DV), which counters the inhibitory mechanism of PPQ (20). Increased

parasite survival at higher PPQ concentrations amongst Cambodian field isolates was shown to

be associated with amplification of *pm*, further connecting plasmepsins with a PPQ-R phenotype

(13). Further *in vitro* studies of *pm* amplification have found that in a PPQ-S 3D7 background,

inactivation of *pm* results in a mild increase in PPQ susceptibility, whereas overexpression of *pm*

alone did not impact parasite sensitivity to PPQ, artesunate (AS), or CQ (21, 22). While *pm* has

clearly been shown to impact parasite response to PPQ, variations of *in vitro* and isolate-based

genotype-phenotype associations suggest that while CN is associated with decreased sensitivity

to PPQ, additional loci contribute to resistance (13, 23).

Additional to the role of *pm*, decreased copies of *pfmdr1* and single nucleotide

polymorphisms (SNP) in several other genes, such as *exonuclease* (PF3D7_1362500) and the

Plasmodium falciparum chloroquine resistance transporter (*pfcrt*) have been shown to contribute

to PPQ-R (16, 18, 24–26). Novel haplotypes containing amino acid substitutions in PfCRT were

originally identified due to their increasing prevalence in areas utilizing DHA+PPQ, and *in vitro*

studies have found that edited isogenic parasites with these substitutions alone can impact

parasite uptake of PPQ and drug sensitivity independent of *pm* copy number (25, 27–30). Field

studies from within the GMS have noted that some of these *pfcrt* haplotypes have continued to

expand within the last 5 years and are often found in parasites that also have multiple copies of

pm, indicating no detrimental fitness cost associated with this allelic combination (31, 32).

Additionally, the main PfCRT substitutions associated with PPQ-R, such as H97Y, F145I, M343L,

G350R, and G353V, all arose from either a Dd2 or 7G8-like background (33, 34). This further

connects genetic background to the original evolution of these resistance-conferring SNPs.

Compared to other antimalarials, assessing *in vitro* PPQ-R has been a challenge due to

the shape of the dose-response curve in PPQ-R parasites. These resistant parasites show

incomplete clearance at the higher concentrations used during phenotyping, resulting in a bimodal

dose-response curve (13). To address this non-traditional curve shape, area under the curve

(AUC) has been used as a proxy for traditional IC50 assays, with a greater AUC representing

decreased sensitivity (13). Alternatively, the piperaquine survival assay (PSA) was developed to

measure parasite survival rate after exposure to a pharmacologically relevant (200nM) dose of

PPQ (18). While both measurements can correlate with overall PPQ-R, a multi-phenotype

approach can not only validate the strongest factors driving resistance, which should be observed

in all phenotypes, but can potentially identify different secondary loci due to slight differences in

the measured phenotype.

To further understand the complexity and contributions of several loci to PPQ-R, we used

progeny from a genetic cross between a multidrug-resistant Cambodian parasite of KEL1/PLA1

lineage (KH004) and a drug-sensitive Malawi isolate (Mal31). KH004 carries a C508Y Kelch13, a

Dd2-like PfCRT (74I, 75E, 76T, 220S, 271E, 326S, 356T, and 371I) with an addition G367C

substitution, and 4 copies of *pm*. Mal31 has a wild-type genotype at both *kelch13* and *pfcrt* and

carries a single copy of *pm* (Figure 1A). The G367C substitution in PfCRT carried by KH004 has

previously been identified as a novel haplotype present in SEA, however, this is the first study

that directly connects it to *in vitro* PPQ resistance (25, 33, 35).

The genetic background of *P. falciparum* has continuously been shown to play a central

role in the emergence of drug resistance as well as supporting parasite fitness. While several

genetic factors have been implicated in PPQ resistance, understanding the epistatic interactions

between these loci is crucial for identifying parasites that can easily develop resistance or lineages

which have the potential to expand. A genetic cross-based approach to studying PPQ-R allows

for recombination across the entire genome and can identify the contributions of all loci, both

known and unidentified, when paired with linkage analysis such as quantitative trait loci (QTL)

mapping. In this study, we use three different phenotypes of resistance (limited-point IC50, AUS,

and PSA) to identify the contributions of a novel *pfcrt* haplotype and amplification of *pm* to PPQ

resistance as well as detect additional loci which further contribute to resistance and modulate

parasite fitness.

Results

Novel allelic combinations of pfcrt and plasmepsin II/III

From our genetic cross between KH004 and Mal31, previously described in Brenneman

et al. (36), we recovered 101 unique recombinant progeny with various combinations of

inheritance of *pfcrt, plasmepsin II/III,* alleles and *pm* copy number, ranging from single copy to

five copies (Figure 1B). Genome-wide, we see two large skews in inheritance favoring Mal31

alleles on Chr 7 and 14 (Supplementary Figure 1A). Both of these skews towards the African

drug-sensitive parent have been observed in previous crosses, and indicate a potential fitness

cost associated with the inheritance of these alleles (37). Additionally, we see no evidence for

pairwise linkage disequilibrium or co-inheritance among these genes or other loci across the

genome, including when accounting for *pm* CN (Supplementary Figure 1B**)**.

PPQ dose response phenotype

Traditional IC50 assays have not been useful in phenotyping PPQ resistance due to the

well-documented biphasic dose-response curve observed in PPQ-R parasites. Based on the

dose-response phenotypes of the progeny from our KH004 × Mal31 genetic cross, we find that

the combination of mutations in *pfcrt* plus multiple copies of *pm* is responsible for this biphasic

phenotype. Parasites with single copies of *pm*, as well as those which inherited the *pfcrt* allele

from Mal31, all display a traditional sigmoidal dose-response curve. Conversely, parasites that

carry the KH004 *pfcrt* allele while also expressing multiple copies of *pm* produce a biphasic curve.

PfCRT is the main driver of PPQ resistance

One of the major challenges to phenotyping PPQ drug response is the biphasic dose-

response curve, indicating incomplete parasite killing at higher drug concentrations observed in

PPQ-R parasites. To study the response of parasites with novel combinations of *pfcrt* and *pm* to

PPQ independent of dose curve shape, the piperaquine survival assay (PSA) was performed on

48 progeny and both parents from the KH004 × Mal31 cross and encompassed all available

combinations of *pfcrt* and *pm* alleles and *pm* CN. Our results show a significant difference in PPQ

sensitivity based on the *pfcrt* genotype (Figure 2A). Within the PPQ-R population inheriting the

pfcrt allele from the KH004 parent, we further see an increase in resistance associated with an

increase in *pm* CN. Additionally, QTL mapping of the PSA phenotype revealed a highly significant

(1% threshold) 144 kb region on chromosome 7 (LOD score of 10.615), which encompasses *pfcrt*

(Figure 2B,). Additional measures of variance identify that this strong peak containing *pfcrt* is

responsible for 61.9**%** of the observed phenotypic variance in the PSA phenotype. Due to this

large and potentially masking effect of *pfcrt*, secondary scans of residual variation were used after

statistically removing the effect from the chromosome 7 locus. These residual scans identified

another highly significant (1% threshold, LOD score of 4.175) marker on chromosome 14 (Figure

2C). Using a 95% confidence interval (CI) from the highest LOD score, this secondary peak

contains the *plasmepsin II* gene. We further support the association between *pm* CN amplification

and PPQ resistance by using a secondary measure of resistance, AUC (Figure 2D-F). QTL

mapping based on the AUC phenotype identifies a strong peak on chromosome 7 (LOD 6.94),

similar to the one identified using PSA, which accounts for 48.4% of phenotypic variance (Figure

2E). Residual scans of this phenotype also identify a significant (5% threshold, LOD of 4.199)

marker on chromosome 10, which contains 61 genes (Supplementary Table 1).

Plasmepsin II/III plays an additive role in PPQ resistance

Based on our data from PSA and AUC measurements, we identified that copy number

amplification of *pm* contributes to a decreased sensitivity to PPQ (Figure 2A, C, D). However, this

effect of copy number is often masked by *pfcrt* (Figure 2B). The main reason why IC50 is not

traditionally used for phenotyping PPQ is due to the biphasic drug response curve associated with

resistance. By using a limited point IC50 (LP-IC50), we can remove the biphasic portion of the

curve and therefore calculate an IC50 for PPQ. These results show an increased IC50 due to the

inheritance of the *pfcrt* allele as well as *pm* amplification in the Mal31 *pfcrt* population (Figure 3A).

However, this decreased sensitivity to PPQ reaches an IC50 of 6nm, which is still comparable to

other PPQ-S parasites such as 3D7 (21). Once again, QTL mapping of this phenotype identified

a strong association between IC50 and *pfcrt* (Figure 3B). While this IC50 phenotype was sufficient

to map the impact of *pfcrt*, it did not provide the same resolution in differentiating PPQ-R and

PPQ-S parasites as the other phenotypes did.

PfCRT G367C confers PPQ resistance

Using progeny from our KH004×Mal31 genetic cross, we have further validated the results

of previous studies, which indicate that *pfcrt* is the main driver of PPQ resistance. From this

progeny set, inheritance of the *pfcrt* allele from KH004 was the major factor shared by PPQ-R

parasites (Figure 2,3). This parasite carries a Dd2-like *pfcrt* with the addition of a novel mutation

G367C. Most drug resistance-associated mutations in PfCRT are found in one of the 10

transmembrane (TM) domains, and specifically regions that make up the central cavity of the

transporter (33, 38). The G36C substitution differs from other PPQ-R-associated mutations in that

while it is found within TM9 with other known mutations, it is found in the digestive vacuole (DV)

exposed portion of this protein and does not part of the core structure of the transporter's central

cavity (Supplementary Figure 2). While the Dd2 *pfcrt* haplotype confers to resistance to other

antimalarials, such as CQ, the haplotype does not confer PPQ resistance, indicating the direct

influence of the G367C mutation. To validate this, we used PSA to measure PPQ sensitivity

between Dd2, KH004, a selfed KH004 parasite from our cross, which carries only a single copy

of *plasmepsin II/II, and* NHP4026 which is from SEA and also expresses a Dd2 *pfcrt*. PSA reveals

that both parasites containing the G367C SNP are significantly less sensitive to PPQ compared

to the Dd2 parasite, which lacks that mutation (Supplementary Figure 3).

This novel G367C mutation has previously been mentioned by Ross et al (25) for its

presence in SEA but was not globally prevalent at the time of analysis. Since the KEL1/PLA1

lineage has continued to spread throughout SEA and PPQ resistance has continued to emerge,

we analyzed the current prevalence of this novel mutation using the MalariaGen Pf7 dataset. Out

of 16,203 global *pfcrt* haplotypes, 29 samples carried the G367C substitution (Figure 4). This

substitution was originally identified in 2012 and was restricted to Cambodia but has emerged in

neighboring countries Vietnam and Laos since 2017 (Supplementary Table 2). Additionally, 93%

(27/29) of these filed samples also carry multiple copies of *pm*.

To further identify selective pressures on *pfcrt* in Cambodia, Laos, and Vietnam, we used

Tajima's D and nucleotide diversity to characterize positive selective pressure on this gene

(Figure 4). A Tajima's D value <2 results from a significant abundance in rare alleles that deviates

from expected mutation-drift equilibrium, indicative of positive selection on the gene. While *pfcrt*

remains in equilibrium with no selective pressures in Cambodia and Vietnam, we observed

significant positive selection for this gene in Laos (D= -2.718 p = 0.007) (Figure 4). Measurements

of Tajima's D over time also indicate that this positive selection event is recent, with a significant

negative Tajima's D score being observed only in 2018 (Supplementary Figure 4).

232
Impacts of pfcrt and pm inheritance on parasite fitness

Our data across multiple resistance phenotypes support that *pfcrt* is the primary driver of PPQ-

R, with an additive increase in resistance resulting from increased copies of *pm.* It has also been

well-documented that mutations in major resistance-associated genes can be associated with a

loss in parasite fitness (30, 39, 40). For these novel resistance-conferring genotypes to increase

in global prevalence, parasites need to preserve fitness either through low-cost resistance

mutations or the acquisition of secondary compensatory mutations that preserve or improve

fitness. As a measurement of blood-stage fitness, we performed competitive growth assays

between progeny of the KH004 × Mal31 cross and both parental parasites (Supplementary Figure

5). These co-cultures were maintained for a maximum of 40 days or until one parasite reached

fixation (>90% of culture), with samples being collected every 7 days. Progeny vs. parent

competitions revealed that the majority of progeny had a fitness level between the two parents,

and four parasites were outside of the fitness range of the parents (two more fit than KH004 and

two less fit than Mal31), with no significant clustering in fitness based on inherited *pfcrt* and *pm*

genotypes (Supplementary Figure 5).

Based on the similar fitness phenotypes generated from competitions against the parental

parasites, a select subset of 18 progeny and both parents were set up in overlapping competitions

against common competitors, resulting in 212 competitive outcomes (win, lose, or draw). The

outcomes of these pairwise competitions were used to generate an Elo Rating for each parasite

using the *EloRating* package in R (41). This Elo phenotype has previously been used in ranking

dominance rankings but has not been adapted for ranking *P. falciparum* fitness hierarchies. Elo

ratings provide a weighted ranking of parasite fitness based on competitive growth outcome as

well as the fitness ranking of the competitor, with a higher Elo Rating representing a greater

competitive fitness level. Elo Ratings for these 20 parasites provide a much finer resolution of

fitness differences between parasites of seemingly similar fitness. Similar to competitions against

both parents, Elo Rating does not cluster based on *pfcrt* and *plasmepsin* genotype (Figure 5A).

However, QTL analysis using Elo phenotypes reveals a highly significant (1% threshold) on

chromosome 9 (LOD score of 4.489), which encompasses a 35kb region containing 15 genes

(Figure 5B). Parasites with the greatest competitive fitness were associated with inheritance of

this 35kb region from the KH004 parent Sequence comparison between KH004 and Mal31 for

these 15 genes identified 4 coding SNPs in two genes: PF3D7_0934100 and PF3D7_0934700

(Supplementary Table 3). In order to isolate the role of *pm* CN variation with respect to parasite

fitness, we compared fitness levels of two sets of isogenic progeny as well as parental parasites,

which have been selfed and varied in copy number in our competitive growth assays and

generated Elo ratings (Supplementary Figure 6). Based on these Elo scores, we observe no major

influence of *pm* CN alone on parasite fitness.

<u>Discussion</u>

In this study, we used a genetic cross between a multidrug-resistant Cambodian parasite

and a drug-sensitive isolate from Malawi to characterize the genetic contributors to PPQ-R.

Previous work on this subject has either utilized naturally resistant patient isolates or conducted

in vitro manipulations to confer PPQ-R (13, 16, 20, 25). By using a genetic cross, we can track

the inheritance of alleles that have previously been associated with resistance, such as SNPs in

PfCRT or amplification of *pm*, while accounting for variation in the parasite's genetic background.

Within our progeny set, we have identified that the inheritance of the KH004 PfCRT (Dd2 +

G367C) is the main driver of PPQ-R, with the amplification of *pm* serving an additive role amongst

resistant parasites.

The KH004 parent of this cross expresses a novel G367C substitution in PfCRT, which

has previously been identified in SEA but has never been associated with *in vitro* resistance to

PPQ. Through comparisons between KH004, Dd2, and other Southeast Asian parasites, we

observed a > 30-fold change in PSA between KH004 and parasites not expressing a Dd2-like

PfCRT, indicating a direct connection between this substitution and PPQ-R. While the G367C

substitution has not previously been connected to PPQ-R, this residue is within the binding site

of the structurally similar chloroquine derivative perfluorophenylazido biotinylated chloroquine

(AzBCQ) (42). Through the association of this residue with PPQ-R, we have contributed to a

further understanding of structural regions of PfCRT that influence resistance to PPQ. While we

have shown that the addition of the G367C substitution confers PPQ-R, the overall *pfcrt* haplotype

is likely what facilitates the evolution of these resistance-conferring SNPs. The major *pfcrt*

haplotypes associated with PPQ-R have all come out of a 7G8 or Dd2 background, and

transfection of these mutations into different *pfcrt* backgrounds has led to slight decreases in

resistance compared to those in a Dd2 background (28, 33).

Previously studied PPQ-R-associated PfCRT substitutions such as T93S, H97Y, C101F,

F145I, M343L, and G353V were originally identified based on their growing prevalence in areas

using the ACT DHA+PPQ (25, 33). The novel G367C substitution expressed by KH004 arose in

western Cambodia in 2012 and was found at a low abundance in the population, which is why it

had been overlooked compared to other mutations. Since its initial emergence, the Dd2+G367C

haplotype has continued to persist at a consistently low abundance but has spread to neighboring

countries, Laos and Vietnam, with the latter accounting for the majority of identifications in recent

years. As most of SEA has transitioned from DHA+PPQ to AL as the frontline ACT, Vietnam

continues to use PPQ, which might explain the recent increase in abundance of this novel

haplotype within the country (1). On the other hand, the introduction of this haplotype into Laos

can potentially be accounted for by a recent introduction of PPQ-R parasites into the country from

Cambodia (43).

Due to the difficult nature of phenotyping PPQ-R, we used 3 previously utilized resistance

phenotypes: AUC, limited point IC50, and PSA. By using these different assays, we can break

the resistance phenotype into three components, survival under high single dose PPQ (PSA), low

dose growth inhibition (IC50), and shape of the dose-response curve under high PPQ

concentrations (AUC), to identify the genetic factors driving variation within each of these

phenotypes (13, 18, 21, 22). All three of these phenotypes were able to identify PfCRT as the

main underlying contributor to resistance but identified different secondary loci for each

phenotype. QTL scans based on PSA phenotype after removing the large effect of *pfcrt* further

identify a region on chromosome 14 containing *plasmepsin II* as a secondary contributor to

resistance. This association supports the additive effect of *pm* copy number amplification

observed in our PPQ-R progeny (Figure 2). Using the AUC phenotype supported the strong

impact of *pfcrt*, but also identified a secondary association between the phenotype and a locus

on chromosome 10. One of the genes with the strongest association within this region is

autophagy-related gene 18 (*atg18*). This protein, which localizes to the parasite DV, has

previously been associated with decreased sensitivity to several antimalarials such as DHA,

artemether (AM), and PPQ, while also being connected to increased survival under nutrient

deprivation (44–46). Due to its previous association with PPQ-R and the association only with the

AUC phenotype, *atg18* may potentially be one of the underlying factors for the non-traditional

dose-response curve encountered when studying *in vitro* PPQ-R

Previous studies into the impact of *pm* amplification have hypothesized that instead of

directly modulating PPQ-R, expressing multiple copies of this gene might serve a compensatory

role in parasite fitness (22). This hypothesis, paired with extensive previous findings on the fitness

costs of *pfcrt* resistance mutations, led us to assess the competitive fitness levels of our progeny

with various combinations of *pfcrt* and *pm* (29, 39, 47). We found that compared to the KH004

and Mal31 parents of this cross, there was no significant clustering of *pfcrt/pm* genotype

combinations that were associated with parasite fitness. Additionally, we have utilized Elo ratings

as a new way to assess parasite fitness within our set of progeny. Elo ratings are an effective

method for efficiently comparing head-to-head fitness levels across a large progeny set and

provide a more refined phenotype, which is necessary for effective QTL mapping. Using this Elo

phenotype, we have been able to identify a highly significant association between parasite fitness

and a narrow 35kb region on chromosome 9. This region is mainly composed of genes involved

in development, transcriptional regulation, and metabolism and contains two genes with coding

region SNP differences between KH004 and Mal31. Neither of these genes have previously been

directly connected to parasite fitness and therefore require future work to further examine the role

that they play in parasite fitness.

Understanding the genetic determinants of PPQ resistance is essential for future control

and prevention measures. Through our genetic cross, we have confirmed the major drivers of

resistance, including a novel *pfcrt* mutation, as well as identified secondary genetic factors that

contribute to parasite resistance and fitness. Additionally, we have been able to incorporate

several previously used PPQ-R phenotypes into our analysis and identify unique genetic regions

that are associated with changes in each phenotype. By combining this novel genetic cross with

rigorous *in vitro* resistance and fitness phenotyping, we have identified the genetic architecture

which promotes decreased sensitivity to PPQ while remaining competitively fit. While the

DHA+PPQ ACT continues to be used as a frontline ACT, it is crucial to understand the genetic

underpinnings of resistance. Furthermore, this genetic cross also highlights the recombination

events that would be required to confer PPQ-R if one of these multidrug resistant parasite lineages

were to expand to Africa.

References

1. World Health Organization. 2022. World malaria report 2022. Geneva.
2. White NJ, Pukrittayakamee S, Hien TT, Faiz MA, Mokuolu OA, Dondorp AM. 2014. Malaria. The Lancet 383:723–735.
3. Blasco B, Leroy D, Fidock DA. 2017. Antimalarial drug resistance: linking Plasmodium falciparum parasite biology to the clinic. Nat Med 23:917–928.
4. Payne D. 1987. Spread of chloroquine resistance in Plasmodium falciparum. Parasitol Today 3:241–246.
5. Mita T, Tanabe K, Kita K. 2009. Spread and evolution of Plasmodium falciparum drug resistance. Parasitol Int 58:201–209.
6. Imwong M, Suwannasin K, Kunasol C, Sutawong K, Mayxay M, Rekol H, Smithuis FM, Hlaing TM, Tun KM, van der Pluijm RW, Tripura R, Miotto O, Menard D, Dhorda M, Day NPJ, White NJ, Dondorp AM. 2017. The spread of artemisinin-resistant Plasmodium falciparum in the Greater Mekong subregion: a molecular epidemiology observational study. Lancet Infect Dis 17:491–497.
7. Imwong M, Dhorda M, Myo Tun K, Thu AM, Phyo AP, Proux S, Suwannasin K, Kunasol C, Srisutham S, Duanguppama J, Vongpromek R, Promnarate C, Saejeng A, Khantikul N, Sugaram R, Thanapongpichat S, Sawangjaroen N, Sutawong K, Han KT, Htut Y, Linn K, Win AA, Hlaing TM, van der Pluijm RW, Mayxay M, Pongvongsa T, Phommasone K, Tripura R, Peto TJ, von Seidlein L, Nguon C, Lek D, Chan XHS, Rekol H, Leang R, Huch C, Kwiatkowski DP, Miotto O, Ashley EA, Kyaw MP, Pukrittayakamee S, Day NPJ, Dondorp AM, Smithuis FM, Nosten FH, White NJ. 2020. Molecular epidemiology of resistance to antimalarial drugs in the Greater Mekong subregion: an observational study. Lancet Infect Dis 20:1470–1480.
8. Ecker A, Lehane AM, Clain J, Fidock DA. 2012. PfCRT and its role in antimalarial drug resistance. Trends in Parasitology 28:504–514.
9. Amato R, Pearson RD, Almagro-Garcia J, Amaratunga C, Lim P, Suon S, Sreng S, Drury E, Stalker J, Miotto O, Fairhurst RM, Kwiatkowski DP. 2018. Origins of the current outbreak of multidrug-resistant malaria in southeast Asia: a retrospective genetic study. Lancet Infect Dis 18:337–345.
10. Hamilton WL, Amato R, van der Pluijm RW, Jacob CG, Quang HH, Thuy-Nhien NT, Hien TT, Hongvanthong B, Chindavongsa K, Mayxay M, Huy R, Leang R, Huch C, Dysoley L, Amaratunga C, Suon S, Fairhurst RM, Tripura R, Peto TJ, Sovann Y, Jittamala P, Hanboonkunupakarn B, Pukrittayakamee S, Chau NH, Imwong M, Dhorda M, Vongpromek R, Chan XHS, Maude RJ, Pearson RD, Nguyen T, Rockett K, Drury E, Gonçalves S, White NJ, Day NP, Kwiatkowski DP, Dondorp AM, Miotto O. 2019. Evolution and expansion of multidrug-resistant malaria in southeast Asia: a genomic epidemiology study. Lancet Infect Dis 19:943–951.
11. MalariaGEN Plasmodium falciparum Community Project. 2016. Genomic epidemiology of artemisinin resistant malaria. Elife 5:e08714.
12. MalariaGEN, Abdel Hamid MM, Abdelraheem MH, Acheampong DO, Ahouidi A, Ali M, Almagro-Garcia J, Amambua-Ngwa A, Amaratunga C, Amenga-Etego L, Andagalu B, Anderson T, Andrianaranjaka V, Aniebo I, Aninagyei E, Ansah F, Ansah PO, Apinjoh T, Arnaldo P, Ashley E, Auburn S, Awandare GA, Ba H, Baraka V, Barry A, Bejon P, Bertin GI, Boni MF, Borrmann S, Bousema T, Bouyou-Akotet M, Branch O, Bull PC, Cheah H, Chindavongsa K, Chookajorn T, Chotivanich K, Claessens A, Conway DJ, Corredor V, Courtier E, Craig A, D'Alessandro U, Dama S, Day N, Denis B, Dhorda M, Diakite M, Djimde A, Dolecek C, Dondorp A, Doumbia S, Drakeley C, Drury E, Duffy P, Echeverry DF, Egwang TG, Enosse SMM, Erko B, Fairhurst RM, Faiz A, Fanello CA, Fleharty M, Forbes M, Fukuda M, Gamboa D, Ghansah A, Golassa L, Goncalves S, Harrison GLA, Healy SA, Hendry JA, Hernandez-Koutoucheva A, Hien TT, Hill CA, Hombhanje F, Hott A, Htut Y, Hussein M, Imwong M, Ishengoma D, Jackson SA, Jacob CG, Jeans J, Johnson KJ, Kamaliddin C, Kamau E, Keatley J, Kochakarn T, Konate DS, Konaté A, Kone A, Kwiatkowski DP, Kyaw MP, Kyle D, Lawniczak M, Lee SK, Lemnge M, Lim P, Lon C, Loua KM, Mandara CI, Marfurt J, Marsh K, Maude RJ, Mayxay M, Maïga-Ascofaré O, Miotto O, Mita T, Mobegi V,

Mohamed AO, Mokuolu OA, Montgomery J, Morang'a CM, Mueller I, Murie K, Newton PN, Ngo Duc T, Nguyen T, Nguyen T-N, Nguyen Thi Kim T, Nguyen Van H, Noedl H, Nosten F, Noviyanti R, Ntui VN-N, Nzila A, Ochola-Oyier LI, Ocholla H, Oduro A, Omedo I, Onyamboko MA, Ouedraogo J-B, Oyebola K, Oyibo WA, Pearson R, Peshu N, Phyo AP, Plowe CV, Price RN, Pukrittayakamee S, Quang HH, Randrianarivelojosia M, Rayner JC, Ringwald P, Rosanas-Urgell A, Rovira-Vallbona E, Ruano-Rubio V, Ruiz L, Saunders D, Shayo A, Siba P, Simpson VJ, Sissoko MS, Smith C, Su X, Sutherland C, Takala-Harrison S, Talman A, Tavul L, Thanh NV, Thathy V, Thu AM, Toure M, Tshefu A, Verra F, Vinetz J, Wellems TE, Wendler J, White NJ, Whitton G, Yavo W, Van Der Pluijm RW. 2023. Pf7: an open dataset of Plasmodium falciparum genome variation in 20,000 worldwide samples. Wellcome Open Res 8:22.

13. Bopp S, Magistrado P, Wong W, Schaffner SF, Mukherjee A, Lim P, Dhorda M, Amaratunga C, Woodrow CJ, Ashley EA, White NJ, Dondorp AM, Fairhurst RM, Ariey F, Menard D, Wirth DF, Volkman SK. 2018. Plasmepsin II–III copy number accounts for bimodal piperaquine resistance among Cambodian Plasmodium falciparum. Nat Commun 9:1769.

14. Anderson TJC, Nair S, McDew-White M, Cheeseman IH, Nkhoma S, Bilgic F, McGready R, Ashley E, Pyae Phyo A, White NJ, Nosten F. 2017. Population Parameters Underlying an Ongoing Soft Sweep in Southeast Asian Malaria Parasites. Mol Biol Evol 34:131–144.

15. Ariey F, Witkowski B, Amaratunga C, Beghain J, Langlois A-C, Khim N, Kim S, Duru V, Bouchier C, Ma L, Lim P, Leang R, Duong S, Sreng S, Suon S, Chuor CM, Bout DM, Ménard S, Rogers WO, Genton B, Fandeur T, Miotto O, Ringwald P, Le Bras J, Berry A, Barale J-C, Fairhurst RM, Benoit-Vical F, Mercereau-Puijalon O, Ménard D. 2014. A molecular marker of artemisinin-resistant Plasmodium falciparum malaria. Nature 505:50–55.

16. Amato R, Lim P, Miotto O, Amaratunga C, Dek D, Pearson RD, Almagro-Garcia J, Neal AT, Sreng S, Suon S, Drury E, Jyothi D, Stalker J, Kwiatkowski DP, Fairhurst RM. 2017. Genetic markers associated with dihydroartemisinin–piperaquine failure in Plasmodium falciparum malaria in Cambodia: a genotype–phenotype association study. The Lancet Infectious Diseases 17:164–173.

17. Tumwebaze PK, Conrad MD, Okitwi M, Orena S, Byaruhanga O, Katairo T, Legac J, Garg S, Giesbrecht D, Smith SR, Ceja FG, Nsobya SL, Bailey JA, Cooper RA, Rosenthal PJ. 2022. Decreased susceptibility of Plasmodium falciparum to both dihydroartemisinin and lumefantrine in northern Uganda. Nat Commun 13:6353.

18. Duru V, Khim N, Leang R, Kim S, Domergue A, Kloeung N, Ke S, Chy S, Eam R, Khean C, Loch K, Ken M, Lek D, Beghain J, Ariey F, Guerin PJ, Huy R, Mercereau-Puijalon O, Witkowski B, Menard D. 2015. Plasmodium falciparum dihydroartemisinin-piperaquine failures in Cambodia are associated with mutant K13 parasites presenting high survival rates in novel piperaquine in vitro assays: retrospective and prospective investigations. BMC Med 13:305.

19. Amaratunga C, Lim P, Suon S, Sreng S, Mao S, Sopha C, Sam B, Dek D, Try V, Amato R, Blessborn D, Song L, Tullo GS, Fay MP, Anderson JM, Tarning J, Fairhurst RM. 2016. Dihydroartemisinin-piperaquine resistance in Plasmodium falciparum malaria in Cambodia: a multisite prospective cohort study. Lancet Infect Dis 16:357–365.

20. Witkowski B, Duru V, Khim N, Ross LS, Saintpierre B, Beghain J, Chy S, Kim S, Ke S, Kloeung N, Eam R, Khean C, Ken M, Loch K, Bouillon A, Domergue A, Ma L, Bouchier C, Leang R, Huy R, Nuel G, Barale J-C, Legrand E, Ringwald P, Fidock DA, Mercereau-Puijalon O, Ariey F, Ménard D. 2017. A surrogate marker of piperaquine-resistant Plasmodium falciparum malaria: a phenotype–genotype association study. The Lancet Infectious Diseases 17:174–183.

21. Mukherjee A, Gagnon D, Wirth DF, Richard D. 2018. Inactivation of Plasmepsins 2 and 3 Sensitizes Plasmodium falciparum to the Antimalarial Drug Piperaquine. Antimicrob Agents Chemother 62.

22. Loesbanluechai D, Kotanan N, de Cozar C, Kochakarn T, Ansbro MR, Chotivanich K, White NJ, Wilairat P, Lee MCS, Gamo FJ, Sanz LM, Chookajorn T, Kümpornsin K. 2019. Overexpression of plasmepsin II and plasmepsin III does not directly cause reduction in Plasmodium falciparum sensitivity to artesunate, chloroquine and piperaquine. Int J Parasitol Drugs Drug Resist 9:16–22.

23.   Parobek CM, Parr JB, Brazeau NF, Lon C, Chaorattanakawee S, Gosi P, Barnett EJ, Norris LD,
Meshnick SR, Spring MD, Lanteri CA, Bailey JA, Saunders DL, Lin JT, Juliano JJ. 2017. Partner-
Drug Resistance and Population Substructuring of Artemisinin-Resistant Plasmodium falciparum in
Cambodia. Genome Biol Evol 9:1673–1686.
24.   Boonyalai N, Vesely BA, Thamnurak C, Praditpol C, Fagnark W, Kirativanich K, Saingam P,
Chaisatit C, Lertsethtakarn P, Gosi P, Kuntawunginn W, Vanachayangkul P, Spring MD, Fukuda
MM, Lon C, Smith PL, Waters NC, Saunders DL, Wojnarski M. 2020. Piperaquine resistant
Cambodian Plasmodium falciparum clinical isolates: in vitro genotypic and phenotypic
characterization. Malar J 19:269.
25.   Ross LS, Dhingra SK, Mok S, Yeo T, Wicht KJ, Kümpornsin K, Takala-Harrison S, Witkowski B,
Fairhurst RM, Ariey F, Menard D, Fidock DA. 2018. Emerging Southeast Asian PfCRT mutations
confer Plasmodium falciparum resistance to the first-line antimalarial piperaquine. Nat Commun
9:3314.
26.   Dondorp AM. 2017. New genetic marker for piperaquine resistance in Plasmodium falciparum. The
Lancet Infectious Diseases 17:119–121.
27.   Okombo J, Mok S, Qahash T, Yeo T, Bath J, Orchard LM, Owens E, Koo I, Albert I, Llinás M,
Fidock DA. 2022. Piperaquine-resistant PfCRT mutations differentially impact drug transport,
hemoglobin catabolism and parasite physiology in Plasmodium falciparum asexual blood stages.
PLoS Pathog 18:e1010926.
28.   Wicht KJ, Small-Saunders JL, Hagenah LM, Mok S, Fidock DA. 2022. Mutant PfCRT Can Mediate
Piperaquine Resistance in African Plasmodium falciparum With Reduced Fitness and Increased
Susceptibility to Other Antimalarials. J Infect Dis 226:2021–2029.
29.   Small-Saunders JL, Hagenah LM, Wicht KJ, Dhingra SK, Deni I, Kim J, Vendome J, Gil-Iturbe E,
Roepe PD, Mehta M, Mancia F, Quick M, Eppstein MJ, Fidock DA. 2022. Evidence for the early
emergence of piperaquine-resistant Plasmodium falciparum malaria and modeling strategies to
mitigate resistance. PLoS Pathog 18:e1010278.
30.   Dhingra SK, Gabryszewski SJ, Small-Saunders JL, Yeo T, Henrich PP, Mok S, Fidock DA. 2019.
Global Spread of Mutant PfCRT and Its Pleiotropic Impact on Plasmodium falciparum Multidrug
Resistance and Fitness. mBio 10:e02731-18.
31.   van der Pluijm RW, Imwong M, Chau NH, Hoa NT, Thuy-Nhien NT, Thanh NV, Jittamala P,
Hanboonkunupakarn B, Chutasmit K, Saelow C, Runjarern R, Kaewmok W, Tripura R, Peto TJ,
Yok S, Suon S, Sreng S, Mao S, Oun S, Yen S, Amaratunga C, Lek D, Huy R, Dhorda M,
Chotivanich K, Ashley EA, Mukaka M, Waithira N, Cheah PY, Maude RJ, Amato R, Pearson RD,
Gonçalves S, Jacob CG, Hamilton WL, Fairhurst RM, Tarning J, Winterberg M, Kwiatkowski DP,
Pukrittayakamee S, Hien TT, Day NP, Miotto O, White NJ, Dondorp AM. 2019. Determinants of
dihydroartemisinin-piperaquine treatment failure in Plasmodium falciparum malaria in Cambodia,
Thailand, and Vietnam: a prospective clinical, pharmacological, and genetic study. Lancet Infect
Dis 19:952–961.
32.   Dhingra SK, Small-Saunders JL, Ménard D, Fidock DA. 2019. Plasmodium falciparum resistance to
piperaquine driven by PfCRT. The Lancet Infectious Diseases 19:1168–1169.
33.   Kim J, Tan YZ, Wicht KJ, Erramilli SK, Dhingra SK, Okombo J, Vendome J, Hagenah LM,
Giacometti SI, Warren AL, Nosol K, Roepe PD, Potter CS, Carragher B, Kossiakoff AA, Quick M,
Fidock DA, Mancia F. 2019. Structure and drug resistance of the Plasmodium falciparum
transporter PfCRT. Nature 576:315–320.
34.   Pelleau S, Moss EL, Dhingra SK, Volney B, Casteras J, Gabryszewski SJ, Volkman SK, Wirth DF,
Legrand E, Fidock DA, Neafsey DE, Musset L. 2015. Adaptive evolution of malaria parasites in
French Guiana: Reversal of chloroquine resistance by acquisition of a mutation in *pfcrt*. Proc Natl
Acad Sci USA 112:11672–11677.
35.   Mairet-Khedim M, Leang R, Marmai C, Khim N, Kim S, Ke S, Kauy C, Kloeung N, Eam R, Chy S,
Izac B, Mey Bouth D, Dorina Bustos M, Ringwald P, Ariey F, Witkowski B. 2021. Clinical and In

Vitro Resistance of Plasmodium falciparum to Artesunate-Amodiaquine in Cambodia. Clin Infect
Dis 73:406–413.

36. Brenneman KV, Li X, Kumar S, Delgado E, Checkley LA, Shoue DA, Reyes A, Abatiyow BA,
Haile MT, Tripura R, Peto T, Lek D, Button-Simons KA, Kappe SHI, Dhorda M, Nosten F,
Nkhoma SC, Cheeseman IH, Vaughan AM, Ferdig MT, Anderson TJC. 2022. Optimizing bulk
segregant analysis of drug resistance using Plasmodium falciparum genetic crosses conducted in
humanized mice. iScience 25:104095.

37. Button-Simons KA, Kumar S, Carmago N, Haile MT, Jett C, Checkley LA, Kennedy SY, Pinapati
RS, Shoue DA, McDew-White M, Li X, Nosten FH, Kappe SH, Anderson TJC, Romero-Severson
J, Ferdig MT, Emrich SJ, Vaughan AM, Cheeseman IH. 2021. The power and promise of genetic
mapping from Plasmodium falciparum crosses utilizing human liver-chimeric mice. Commun Biol
4:734.

38. Willems A, Kalaw A, Ecer A, Kotwal A, Roepe LD, Roepe PD. 2023. Structures of Plasmodium
falciparum Chloroquine Resistance Transporter (PfCRT) Isoforms and Their Interactions with
Chloroquine. Biochemistry 62:1093–1110.

39. Osman ME, Mockenhaupt FP, Bienzle U, Elbashir MI, Giha HA. 2007. Field-based evidence for
linkage of mutations associated with chloroquine (pfcrt/pfmdr1) and sulfadoxine–pyrimethamine
(pfdhfr/pfdhps) resistance and for the fitness cost of multiple mutations in P. falciparum. Infection,
Genetics and Evolution 7:52–59.

40. Nair S, Li X, Arya GA, McDew-White M, Ferrari M, Nosten F, Anderson TJC. 2018. Fitness Costs
and the Rapid Spread of kelch13-C580Y Substitutions Conferring Artemisinin Resistance.
Antimicrob Agents Chemother 62:e00605-18.

41. Neumann C, Kulik L. 2014. Package"EloRating" (0.43).

42. Lekostaj JK, Natarajan JK, Paguio MF, Wolf C, Roepe PD. 2008. Photoaffinity labeling of the
Plasmodium falciparum chloroquine resistance transporter with a novel perfluorophenylazido
chloroquine. Biochemistry 47:10394–10406.

43. Wasakul V, Disratthakit A, Mayxay M, Chindavongsa K, Sengsavath V, Thuy-Nhien N, Pearson
RD, Phalivong S, Xayvanghang S, Maude RJ, Gonçalves S, Day NP, Newton PN, Ashley EA,
Kwiatkowski DP, Dondorp AM, Miotto O. 2022. Malaria outbreak in Laos driven by a selective
sweep for Plasmodium falciparum kelch13 R539T mutants: a genetic epidemiology analysis. Lancet
Infect Dis S1473-3099(22)00697–1.

44. Sudhakar R, Das D, Thanumalayan S, Gorde S, Sijwali PS. 2021. Plasmodium falciparum Atg18
localizes to the food vacuole via interaction with the multi-drug resistance protein 1 and
phosphatidylinositol 3-phosphate. Biochem J 478:1705–1732.

45. Wang Z, Cabrera M, Yang J, Yuan L, Gupta B, Liang X, Kemirembe K, Shrestha S, Brashear A, Li
X, Porcella SF, Miao J, Yang Z, Su X-Z, Cui L. 2016. Genome-wide association analysis identifies
genetic loci associated with resistance to multiple antimalarials in Plasmodium falciparum from
China-Myanmar border. Sci Rep 6:33891.

46. Breglio KF, Amato R, Eastman R, Lim P, Sa JM, Guha R, Ganesan S, Dorward DW, Klumpp-
Thomas C, McKnight C, Fairhurst RM, Roberts D, Thomas C, Simon AK. 2018. A single
nucleotide polymorphism in the Plasmodium falciparum atg18 gene associates with artemisinin
resistance and confers enhanced parasite survival under nutrient deprivation. Malar J 17:391.

47. Dhingra SK, Redhi D, Combrinck JM, Yeo T, Okombo J, Henrich PP, Cowell AN, Gupta P,
Stegman ML, Hoke JM, Cooper RA, Winzeler E, Mok S, Egan TJ, Fidock DA. 2017. A Variant
PfCRT Isoform Can Contribute to *Plasmodium falciparum* Resistance to the First-Line Partner
Drug Piperaquine. mBio 8:e00303-17.

48. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R,
Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-Paredes
B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger M,
Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW, Kavukcuoglu K,

Kohli P, Hassabis D. 2021. Highly accurate protein structure prediction with AlphaFold. Nature
596:583–589.
49. Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, Yuan D, Stroe O, Wood G,
Laydon A, Žídek A, Green T, Tunyasuvunakool K, Petersen S, Jumper J, Clancy E, Green R, Vora
A, Lutfi M, Figurnov M, Cowie A, Hobbs N, Kohli P, Kleywegt G, Birney E, Hassabis D, Velankar
S. 2022. AlphaFold Protein Structure Database: massively expanding the structural coverage of
protein-sequence space with high-accuracy models. Nucleic Acids Res 50:D439–D444.
50. Davis SZ, Singh PP, Vendrely KM, Shoue DA, Checkley LA, McDew-White M, Button-Simons
KA, Cassady Z, Sievert MAC, Foster GJ, Nosten FH, Anderson TJC, Ferdig MT. 2020. The
extended recovery ring-stage survival assay provides a superior association with patient clearance
half-life and increases throughput. Malar J 19:54.
51. Tirrell AR, Vendrely KM, Checkley LA, Davis SZ, McDew-White M, Cheeseman IH, Vaughan
AM, Nosten FH, Anderson TJC, Ferdig MT. 2019. Pairwise growth competitions identify relative
fitness relationships among artemisinin resistant Plasmodium falciparum field isolates. Malaria
Journal 18.
52. Amambua-Ngwa A, Button-Simons KA, Li X, Kumar S, Brenneman KV, Ferrari M, Checkley LA,
Haile MT, Shoue DA, McDew-White M, Tindall SM, Reyes A, Delgado E, Dalhoff H, Larbalestier
JK, Amato R, Pearson RD, Taylor AB, Nosten FH, D'Alessandro U, Kwiatkowski D, Cheeseman
IH, Kappe SHI, Avery SV, Conway DJ, Vaughan AM, Ferdig MT, Anderson TJC. 2023.
Chloroquine resistance evolution in Plasmodium falciparum is mediated by the putative amino acid
transporter AAT1. Nat Microbiol https://doi.org/10.1038/s41564-023-01377-z.
53. Neumann C, Duboscq J, Dubuc C, Ginting A, Irwan AM, Agil M, Widdig A, Engelhardt A. 2011.
Assessing dominance hierarchies: validation and advantages of progressive evaluation with Elo-
rating. Animal Behaviour 82:911–921.
54. Sen S, Churchill GA. 2001. A statistical framework for quantitative trait mapping. Genetics
159:371–387.
55. Broman KW, Wu H, Sen S, Churchill GA. 2003. R/qtl: QTL mapping in experimental crosses.
Bioinformatics 19:889–890.
56. Churchill GA, Doerge RW. 1994. Empirical threshold values for quantitative trait mapping.
Genetics 138:963–971.
57. Paradis E. 2010. pegas: an R package for population genetics with an integrated–modular approach.
Bioinformatics 26:419–420.
595

A.

Parasite name	KH004	Mal31
Origin	2016, Western Cambodia	2016, Malawi
Kelch13	C580Y	Wild-type
PfCRT	M74I, N75E, K76T, A220S, Q271E, N326S, I356T, R371I, **G367C**	Wild-type
Plasmepsin II/III Copy #	~4 copies	1 copy
PfAAT1	S258L, F313S	S258L
PfMDR1	Y184F	Y184F
PfMDR2	T484I, F423Y, G229D, S208N	I492V, S208N
Competitive fitness	Mild	Unfit

B.

PfCRT/PM Copy #/Plasmepsin	# of Available Progeny
M1M	47
M1K	10
M2K	13
M3K	7
M4K	3
M5K	4
K1M	8
K1K	3
K2K	7
K3K	1

Figure 1. Description of cross parents and progeny demographics. A) KH004 is a multidrug Cambodian oarasite of the KEL1/PLA1 lineage and was isolated in 2016. It contains several traditional markers of drug resistance such as C580Y in Kelch13, a Dd2-like PfCRT with an additional G367C substitution, and multiple copies of plasmepsin II/III. Mal31 is a Malawi drug-sensitive parasite which has a wild-type genetic background at major resistance markers and expresses a single copy of plasmepsin II/III. B) Through our genetic cross, we have recovered 101 unique recombinant progeny which have various allelic combinations at pfcrt and pm. Progeny were grouped into 10 categories based on the allele inheritance at pfcrt and pm as well as pm copy number. K = allele inherited from KH004. M = allele inherited from Mal31

Figure 2: PfCRT is the major driver of PPQ-R. A) Measuring PPQ sensitivity through PSA shows that inheritance of pfcrt allele is the major factor differentiating PPQ-R vs PPQ-S parasites. Additionally, pm copy number amplification further promotes resistance in a PPQ-R parasite expressing the KH004 pfcrt allele. PSA phenotype is represented in fold change between parasites exposed to 200nM PPQ and the no drug control. B) QTL mapping further supports the strong influence of pfcrt (chr 7 peak, LOD = 10.615) on PSA phenotype. C) Removal of the influence of pfcrt on this phenotype identifies a secondary region on chr 14, which encompasses plasmepsin II, influencing PPQ sensitivity through PSA. D) Measuring AUC phenotype reveals additive contributions of pm in parasites with either KH004 or Mal31 pfcrt alleles. E) Similar to QTL from PSA, we identify pfcrt as the major factor contributing to AUC phenotype, and residual scans F) identify a region within chr 10 that is associate with the AUC phenotype. This peak is centered around the gene atg18, which has previously been connected to PPQ-R. By using different phenotypes, we are still able to see the major role of pfcrt in PPQ-R parasites while also using these different phenotypes to identify novel secondary alleles associated with each phenotype.

A.

B.

Figure 3: Additive role of *pm* copy number within PPQ-S population identified through LP-IC50. Using a limited point dose response curve allows us to calculate an IC50 independent of the biphasic nature of the curve. A) By using this modified IC50 phenotype, we see a significant difference in IC50 between *pfcrt* alleles as well as a decreased sensitivity associated with increased *pm* CN in a Mal31-*pfcrt* background. B) QTL mapping based on LP-IC50 solely associates *pfcrt* with this phenotype.

Country	% of PPQ-R samples	# of G367C samples	Nucleotide diversity (π)	Tajima's D
Cambodia (n = 1040)	37.5%	13	0.002600824	-0.8810487
Vietnam (n= 1325)	40.2%	14[a]	0.0033925	-0.8555029
Laos (n= 1061)	17.4%	3	0.005063148	-2.717986 **
Global population (n = 16203)		24		

Figure 4. Prevalence of G367C mutation in SEA. Based on available data from MalariaGen Pf7, this novel polymorphism has been observed throughout SEA, although at low levels, since 2012. Apart from two samples, this mutation is found in PPQ-R samples and is also accompanied by increased copies of *plasmepsin II/III*. Measurements of neutrality (Tajima's D) have identified an abundance of rare alleles which significantly deviates from mutation-drift equilibrium.

[a] Determination of resistance was based on *plasmepsin II/III* copy number amplification. Two samples from Vietnam carried a *single plasmepsin copy* and was deemed sensitive independent of *pfcrt* polymorphisms.

* p < 0.05 ** p < 0.01

Figure 5. QTL mapping of fitness-associated loci using Elo rating phenotype. A) *Mean Elo rating of 18 progeny + 2 parents of the KH004 × Mal31 genetic cross based on 50 permutations of the order of competitions. Elo rating was calculated using the R package EloRating and was based on the each parasite's competitive growth assay outcome. Phenotype distribution is ordered from lowest Elo (least fit) on the left to highest Elo (most fit) on the right.* **B)** *QTL scan based on Elo phenotype reveals a highly significant peak (1% threshold) on chromosome 9. This peak encompasses a 35,264bp region which contains 15 genes. Genes identified from this QTL scan are proposed to directly contribute to parasite fitness.*

601
602

Methods

Parasites used in this study

This study utilizes parents and progeny of the KH004 × Mal31 genetic cross generated by utilizing human liver-chimeric mice and has previously been described in Brenneman et al (36). From this cross, one hundred and one unique recombinant progeny were recovered through limited dilution cloning at 0.3 cells per well. Individual wells containing parasites were identified through qPCR and analysis of progeny relatedness and allele inheritance were based on the methods outlined in Button-Simons et al. (37). Methods for sequencing of progeny, construction of the genetic map, and generation of visual recombination map are all based on Button-Simons et al. (37). Analysis of allelic co-inheritance utilized the entire progeny set of one hundred and one parasites and phenotyping assays utilized fifty unique parasites, forty eight progeny and the two parental parasites.

Parasite culture

Cryopreserved *P. falciparum* stocks of KH004 × Mal31 progeny and KH004-020-019-H9 and Mal31-9040-C11 parents were thawed and cultured in complete media (CM) consisting of 0.5% Albumax II (Gibco, Life Technologies) supplemented RPMI 1640 with L-glutimate (Gibco, Life Technologies) with additional 50 mg/L hypoxanthine (Calbiochem, Sigma-Aldrich), 25 mM HEPES (Corning, VWR), 10ug/ml gentamycin (Gibco, Life Technologies), and 0.225% sodium bicarbonate (Corning, VWR). Parasite cultures were maintained at 5% hematocrit in O+ red blood cells (RBC) (Interstate Blood Bank, Memphis, TN) in separate flasks and maintained consistent temperature (37ºC) and atmosphere (5% CO_2/5% O_2/90% N_2). Parasitemia of cultures was kept below 2% and media changes were performed every 48 hours, corresponding with one intraerythrocytic development cycle.

PfCRT Structure Analysis

Three-dimensional homology model of PfCRT was predicted using AlphaFold (48, 49)

and accessed through the PDB database (6UKJ) (33). Additionally, the F′(ab) fragment used in

7G8 PfCRT cryo-EM structure elucidation (33) was removed from the structure. Protein

mutagenesis and visualization of resistance-associated mutations were conducted using PyMol

software (v2.5.4; Schrödinger, LLC).

Drug susceptibility assays

Prior to susceptibility assay setup, parasites were synchronized using a single layer of

70% Percoll (Sigma Aldrich) in 1X RPMI with 13.3% sorbitol in phosphate buffered saline.

400μL of packed erythrocytes infected by a majority of schizont stage parasites were

resuspended in 2mL of incomplete media, layered over the percoll layer, and centrifuged (1561

x g for 10 in, no brake). The top layer was removed to isolate late-stage schizonts, washed

twice with incomplete media, resuspended at 5% hematocrit in complete media, and placed on

a shaker for 4 hours at 37ºC. Following the 4 hour incubation, parasitemia and stage were

determined through flow cytometry by staining with SYBR Green I and SYTO 61 and analyzed

on a Guava easyCyte HT (Luminex) after 50,000 events were counted. If cultures were >70%

ring-stage, they were diluted to 0.15% parasitemia at 2% hematocrit and set up in a 96-well

plate at a volume of 150μL per well. Within each assay plate, two technical replicates of each

parasite were exposed to 10 concentrations of a 2-fold dilution series of PPQ as well as both

untreated parasite and uninfected RBC controls. Parasites were exposed to PPQ for 72 hours.

Parasite density was determined using SYBR Green and dose-response curves were generated

by plotting percent survival against the log of drug concentration. Analysis of IC50 and AUC was

performed using Prism 9.0 (GraphPad). A nonlinear log inhibitor vs response 4 parameter dose-

response curve was generated based on percent survival. To generate the limited point curve,

data points which made up the biphasic portion of the plot were excluded to allow for a

sigmoidal shaped curve, which was then used for IC50 calculations.

PSA
658
659   PSA was set up as originally described (18) with synchronized early ring stage parasites

being expose to a single 200nM dose of PPQ for 48 hours. This protocol was modified to

resemble the extended recovery RSA (eRRSA) phenotype (50), in which samples are analyzed

at 120 hours post-exposure instead of 72 hours. This extended timepoint has been shown to

provide superior differentiation between resistant and sensitive parasites and allows for finer

resolution between PPQ-S parasites. Parasite prevalence was measured through qPCR and

survival was calculated as fold change between untreated and treated samples for three

technical replicates (3 replicates within the plate as well as 3 independent qPCR reads) and at

least 3 biological replicates per sample. Additionally, to ensure consistency between our

resistance phenotypes, parasites from the same synchronization were used for both PSA and

IC50/AUC susceptibility assays.

670
671   ***Competitive growth assays***

672   To determine relative fitness levels between parasites, we used competitive growth

assays between progeny of the KH004 × Mal31 cross and both parental parasites as a proxy for

in vitro fitness. Parasitemia and stage of parasites was quantified using flow cytometry 15 hours

after synchronization with a 70% Percoll gradient. Synchronized parasites were each adjusted

to 0.5% parasitemia at 5% hematocrit and set up in a 1:1 ratio (1% total parasitemia) in 96-well

plates as previously described in (51, 52). Parasitemia was assessed (Giemsa-stained slides

and microscopy) every two days, in accordance with one intraerythrocytic life cycle, adjusted to

1% parasitemia and supplemented with fresh blood and media. Samples from each well were

collected at regular intervals upon sample dilution and stored at -80ºC for genotyping. These

competitive growth assays were maintained for a maximum of 40 days, or until one parasite

reached fixation (>95% of the co-culture). Parasites were deemed to have equivalent fitness

levels if by 40 days neither parasite reached fixation.

To analyze relative parasite densities in each competition, six microsatellite markers

(MS) were used for genotyping (Supplementary Table 4) to differentiate parasites in co-culture.

PCR amplification was performed using the Phusion Blood Direct PCR Kit (ThermoFisher, cat

#F547L) for 30 cycles as previously described in (51). Amplified microsatellites were analyzed

using the Applied Biosystems 3730xl DNA Analyzer (ThermoFisher). Raw fragment analysis

data was uploaded to the Thermo Fisher Connect platform and analyzed using the microsatellite

analysis tool. The proportion of the two competing parasites in co-culture was determined by

taking the ratio of fluorescent peak height associated with the corresponding PCR product size.

Due to the large number of competitive growth assays required to directly compete all

parasites against each other, all parasites were competed against both parents. Additionally, a

genetically diverse subset of parasites were competed against each other. Based on the

outcomes of the progeny v progeny competitions, an Elo rating system, which has previously

been used for animal dominance hierarchies, was utilized to assign a quantitative score to each

parasite based on their win/loss record while also taking into account the record of the parasite

competed against (53). The package "EloRating" (41) in R Studio (2023.03.0+386). was used

for assigning scores. In order to reduce the effect of competition order on ranking, the

competitive outcomes file was randomized 1000 times to generate 1000 Elo scores which were

then averaged. The distribution of Elo ratings figure was created using Prism 9.0 (GraphPad).

QTL Analysis

708   Statistical analysis of the QTL data for this cross was performed using the computational

methods previously described (54). Blood stages of *P. falciparum* are haploid, therefore, only

two genetic classes are present for each locus. QTL mapping was performed using the "R/qtl"

package (55) in R Studio. One thousand permutations of the trait values determined the 37%,

5%, and 1% genome-wide significance thresholds and the strength of each linkage was

expressed as a logarithm of the odds (LOD) score (56). The main QTL and corresponding mean

trait values from these scans were used to obtain estimates of residual empirical thresholds to

identify secondary loci that contribute to the phenotype of interest (56)

Global haplotype analysis

718   Analysis of global *pfcrt* haplotypes was performed using publicly available sequences

719   through the Pf7 dataset (MalariaGen) (12). PPQ resistance status for this dataset was inferred

720   based on *plasmepsin II/III* amplification, therefore, all samples labeled as PPQ-R are assumed

721   to express more than one copy of these genes. Tajima's D and nucleotide diversity were used

722   as metrics of genetic diversity within and were calculated using the R package "pegas" (57).

728   **Additional Information**

729   Additional figures and tables are available for this paper.

731   Correspondence and requests for materials should be addressed to Michael T. Ferdig (ferdig.1@nd.edu)

733   **Funding**

734   National Institutes of Health grant P01 12072248 (MTF)

736