A *Plasmodium falciparum* genetic cross reveals the contributions of *pfcrt* and

***plasmepsin II/III* to piperaquine drug resistance**

John Kane[a], Xue Li[b], Sudhir Kumar[c], Katrina A. Button-Simons[a], Katelyn M. Vendrely

Brenneman[a*], Haley Dahlhoff[a], Mackenzie A.C. Sievert[a], Lisa A. Checkley[a], Douglas A. Shoue[a],

Puspendra P. Singh[a*], Biley A. Abatiyow[c], Meseret T. Haile[c], Shalini Nair[b], Ann Reyes[b], Rupam

Tripura[d,e], Tom Peto[d,e], Dysoley Lek[f,g], Stefan H.I. Kappe[c,h], Mehul Dhorda[d,e], Standwell C

Nkhoma[i], Ian H. Cheeseman[j], Ashley M. Vaughan[c,h], Timothy J. C. Anderson[b], & Michael T.

Ferdig[a#].

[a]Eck Institute for Global Health, Department of Biological Sciences, University of Notre Dame,

Notre Dame, IN, USA.

[b]Disease Intervention and Prevention Program, Texas Biomedical Research Institute, San

Antonio, TX, USA.

[c]Center for Global Infectious Disease Research, Seattle Children's Research Institute, Seattle,

WA, USA.

[d]Mahidol-Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol

University, Bangkok, Thailand.

[e]Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine Research

Building, University of Oxford Old Road Campus, Oxford, UK.

[f]National Center for Parasitology, Entomology and Malaria Control, Phnom Penh, Cambodia.

[g]School of Public Health, National Institute of Public Health, Phnom Penh, Cambodia.

[h]Department of Pediatrics, University of Washington, Seattle, WA, USA.

[i]BEI Resources, American Type Culture Collection (ATCC), Manassas, VA, USA

[j]Host Pathogen Interactions Program, Texas Biomedical Research Institute, San Antonio, TX,

USA.

Running Title: Genetic determinants of piperaquine resistance

Address correspondence to Michael T. Ferdig, ferdig.1@nd.edu

* Present addresses: Katelyn Vendrely Brenneman, Harvard T.H. Chan School of Public Health,

Boston, MA, USA; Puspendra P. Singh, ECD Division, Indian Council of Medical Research, New

Delhi, India.

Word Count: 231 (abstract), xxx (text)

Abstract

Piperaquine (PPQ) is widely used in combination with dihydroartemisinin (DHA) as a first-line treatment against malaria parasites. Multiple genetic drivers of PPQ resistance have been reported, including mutations in the *Plasmodium falciparum chloroquine resistance transporter* (*pfcrt*) and increased copies of *plasmepsin II/III* (*pm2/3*). We generated a cross between a Cambodia-derived multi-drug resistant KEL1/PLA1 lineage isolate (KH004) and a drug susceptible parasite isolated in Malawi (Mal31). Mal31 harbors a wild-type (3D7-like*) pfcrt* allele and a single copy of *pm2/3,* while KH004 has a chloroquine-resistant (Dd2-like*) pfcrt* allele with an additional G367C substitution and four copies of *pm2/3*. We recovered 104 unique recombinant progeny and examined a targeted set of progeny representing all possible combinations of variants at *pfcrt and pm2/3* for detailed analysis of competitive fitness and a range of PPQ susceptibility phenotypes, including PPQ survival assay (PSA), area under the dose-response curve (AUC), and a limited point IC$_{50}$ (LP-IC$_{50}$). We find that inheritance of the KH004 *pfcrt* allele is required for PPQ resistance, whereas copy number variation in *pm2/3* further enhances resistance but does not confer resistance in the absence of PPQ-R-associated mutations in *pfcrt*. Deeper investigation of genotype-phenotype relationships demonstrates that progeny clones from experimental crosses can be used to understand the relative contributions *of pfcrt, pm2/3,* and parasite genetic background*,* to a range of PPQ-related traits and confirm the critical role of the PfCRT G367C substitution in PPQ resistance.

Importance

Resistance to PPQ used in combination with DHA has emerged in Cambodia and threatens to spread to other malaria-endemic regions. Understanding the causal mutations of drug resistance and their impact on parasite fitness is critical for surveillance and intervention, and can also reveal new avenues to limiting the evolution and spread of drug resistance. An experimental genetic cross is a powerful tool for pinpointing the genetic determinants of key

drug resistance and fitness phenotypes and have the distinct advantage of assaying the effects

of naturally evolved genetic variation. Our study was significantly strengthened because the full

a range of copies of KH004 *pm2/3* was inherited among the progeny clones, allowing us to

directly test the role *of pm2/3* copy number on resistance-related phenotypes in the context of a

unique *pfcrt* allele. Our multi-gene model suggests an important role for both loci in the evolution

of this ACT resistant parasite lineage.

68
69 **Introduction**
70
Malaria is a life-threatening parasitic disease that puts nearly half of the world's

population at risk (1). The World Health Organization estimates that in 2020 there were 247

million malaria cases and 619,000 deaths reported across 84 countries, with 76% of those

deaths occurring in children under the age of 5 (1). The continued evolution of drug-resistant

Plasmodium falciparum asexual blood-stage parasites has been one of the major obstacles to

global malaria elimination and eradication efforts (1). Current frontline treatments for

uncomplicated *P. falciparum* infection are artemisinin-based combination therapies (ACT),

which are composed of a fast-acting artemisinin (ART) derivative paired with one or more

longer-lasting partner drugs (2). ACTs began to replace historically efficient antimalarials, such

as chloroquine (CQ) and sulfadoxine-pyrimethamine (SP), as first-line therapies in the early

2000s after resistance to these previous therapies swept throughout Africa (3–5). The WHO has

recommended ACTs for uncomplicated malaria since 2006 due to their greater efficacy and

decreased resistance compared to other single-drug therapeutics, with the most common ACTs

used in Southeast Asia (SEA) being dihydroartemisinin/piperaquine (DHA+PPQ),

artemether/lumefantrine (AL), and artesunate/mefloquine (AS+MQ). The ACT combination

DHA-PPQ has been widely used throughout the Greater Mekong Subregion (GMS) of SEA,

which has historically been a hotbed for the evolution of antimalarial drug resistance (6–8).

The spread of resistance to DHA+PPQ has been attributed to the rapidly expanding,

multidrug-resistant KEL1/PLA1 lineage of parasites (9–11). These parasites originally emerged

out of western Cambodia in 2008 and have since spread to Vietnam, Laos, and northeastern

Thailand (10). KEL1/PLA1 parasites are characterized by ART-resistant mutations in *kelch13*

(KEL1), specifically the C580Y substitution, as well as a copy number (CN) amplification of the

two aspartic protease genes *plasmepsin II/III* (PLA1) associated with decreased susceptibility to

PPQ (9, 10, 12, 13). Reduced susceptibility to ART component of and ACT results in a greater

remaining parasite load that has to be cleared by the partner drug, increasing the selective

pressure for ACT failure (9, 14–17).

After the original detection of PPQ resistance in Cambodia in 2015, independent

genome-wide association studies identified an association between *plasmepsin II/III* (*pm2/3*)

variants and PPQ-R (16, 18–20). The mechanism proposed to support this association is an

amplification of *pm2/3* that reduces concentrations of reactive heme in the parasite digestive

vacuole (DV) to counter the inhibitory action of PPQ (20). Increased parasite survival at higher

PPQ concentrations amongst Cambodian field isolates was associated with the amplification of

pm2/3 (13). Further *in vitro* studies of *pm2/3* amplification found that in the PPQ-S 3D7

background, inactivation of *pm2/3* results in a mild increase in PPQ susceptibility, whereas

overexpression of *pm2/3* alone did not impact the degree of parasite susceptibility to PPQ,

artesunate (AS), or CQ (21, 22). Nevertheless, while *pm2/3* was initially heralded as a defining

feature of PPQ-R, variations of *in vitro* and isolate-based genotype-phenotype associations

suggest that additional loci are needed to fully account for resistance (13, 23).

Beyond the role of *pm2/3*, decreased copies of *pfmdr1* and single nucleotide

polymorphisms (SNP) in several other genes, including *exonuclease* (PF3D7_1362500) and the

P. falciparum chloroquine resistance transporter (*pfcrt*) have been associated with PPQ-R

parasites (16, 18, 24–26). Novel haplotypes containing amino acid substitutions in PfCRT were

initially identified due to their increasing prevalence in areas utilizing DHA+PPQ, and *in vitro*

studies showed that isogenic parasites edited with these substitutions alone can impact parasite

uptake of PPQ and drug susceptibility independent of *pm2/3* copy number (25, 27–30). Field

studies in the GMS report a continued expansion of some of these *pfcrt* haplotypes in the last

five years that often co-occur in parasites carrying multiple copies of *pm2/3*, suggesting there is

no fitness cost associated with this allelic combination (31, 32). Well studied PfCRT

substitutions linked to PPQ-R, including H97Y, F145I, M343L, G350R, and G353V, arose in

either a Dd2- or 7G8-like CQ-resistant (CQ-R) genetic background (33, 34).

Compared to other antimalarial drugs, assessing *in vitro* PPQ-R has been challenging

due to the unusual dose-response effect in PPQ-R parasites. These resistant parasites show

incomplete killing at the higher concentrations, resulting in an unusual bimodal dose-response

curve (13). To account for this non-traditional drug concentration-biological effect relationships,

various modifications to standard IC_{50} values have been used, including total area under the

curve (AUC) (13). Alternatively, the piperaquine survival assay (PSA) was developed to

measure the parasite survival rate after exposure to a single pharmacologically relevant (200

nM) dose of PPQ (18). While these different readouts will correlate in their characterization of

resistance, they also likely capture distinct aspects of the drug-parasite interaction. We predict

that genetic loci will contribute differentially to these different response readouts, perhaps

pointing to distinct underlying biological in progeny inheriting unique allele combinations. .

The genetic background of *P. falciparum* has been shown to play a central role in the

emergence of drug resistance and its impact on parasite fitness (35–39). While several

individual genes have been implicated in PPQ-R, the understanding how these genes interact is

lacking any may provide insights about how parasites develop resistance and how some

lineages have strong potential to expand. To further dissect the contributions of different loci to

PPQ-R, here we examined progeny from a genetic cross between a multidrug-resistant

Cambodian parasite of KEL1/PLA1 lineage (KH004) and a drug-sensitive Malawian isolate

(Mal31). KH004 carries a C508Y Kelch13 substitution associated with ART-R, a Dd2-like PfCRT

(74I, 75E, 76T, 220S, 271E, 326S, 356T, and 371I), associated with CQ-R, with an additional

G367C substitution, and 4 copies of *pm2/3*. Mal31 has a wild-type genotype at *kelch13* and *pfcrt*

and carries a single copy of *pm2/3* (Table S1). The G367C substitution in PfCRT carried by

KH004 has previously been identified as a novel haplotype present in SEA. However, it's

association with *in* vitro PPQ resistance has not been determined (25, 33, 40).

Pairing a novel genetic cross with quantitative trait loci (QTL) mapping can be a powerful

tool for quantifying the contributions of all loci to phenotypes of interest. We take a multi-

phenotype linkage approach to validate and characterize known resistance determinants as well

as identify novel secondary loci associated with each unique resistance-related phenotype,

underscoring slight differences in biological features of these measured traits. We find that

increasing levels of resistance to PPQ prominently impacted by *pfcrt,* specifically a novel G367C

substitution, with additional contributions from *pm2/3* amplification and other novel loci.

Results

Experimental genetic crosses

We generated two biological replicate genetic crosses between Mal31 and KH004 using

Anopheles stephensi mosquitoes and human liver-chimeric FRG huHep mice, as described in

Vaughan *et al.* (41). There are 14,455 core-genome SNPs (defined in Miles et al. (42))

distinguishing these two parental parasites (43) and both parental parasites were cloned by

limiting dilution. We generated the two recombinant pools for this cross using independent

groups of 250 mosquitoes and intravenously injected salivary gland sporozoites from these two

pools into separate FRG huHep mice. The oocyst prevalence from the two pools were 21% and

33% respectively and the average number of oocysts, 0.5 and 0.6. We were able to isolate

approximately 225,000 salivary gland sporozoites from each pool, approximately 900

sporozoites per mosquito. Results from the parental feeds alone also showed a low oocyst

prevalence (Mal31, 41% and KH004, 33%, with total sporozoite numbers per mosquito

averaging 800 for Mal31 and 480 for KH004. The total number of potential recombinants from

each pool (assuming 50% of oocysts are recombinant and 50% are parental), was 0.5 oocysts ×

4 recombinants per oocyst x 250 mosquitoes x 0.5 = 250 for pool 1 and 0.6 oocysts x 4

recombinants × 250 mosquitoes x 0.5 = 300 for pool 2. The initial allele frequencies of Mal31 of

the two recombinant pools directly from mice were 0.70 and 0.66, suggesting the existence of

Mal31 selfed progeny.

We cloned and sequenced 567 progeny from the two recombinant pools (286 from pool

1 and 281 from pool 2) (Table S2). Initial analysis led to the removal of 153 non-clonal cultures

(F_{WS} <0.9, or non-clonal based on allele frequency plot). Of the 414 clonal progeny cultures, 92

were selfed Mal31, 3 were selfed KH004, and 319 were recombinant. We identified a total of

104 unique recombinants (identical by descent [IBD] clusters), of which 65 were from pool 1 and

39 were from pool 2. There were 44 IBD clusters with more than 2 clones, and 60 were

singletons. The results show a skewing towards selfing of Mal31, as predicted and overall,

319/414 = 77% of the cloned progeny were recombinant.

Determination of plasmepsin II/III copy numbers in parental parasites and progeny

We measured *pm2/3* CN for parental parasites with both nanopore long reads and

Illumina short reads (Figure 1A). Mal31 harbors one copy of *pm2/3*, while KH004 harbors

multiple copies of *pm2/3.* We were able to extract four long reads from KH004 nanopore

sequences that cover the *pm2/3* genes alongside >10kb flanking regions (both upstream and

downstream) and two of the reads had 5 copies, one had 4, and one had 3 copies of *pm2/3.*

We further used Illumina short reads to analyze *pm2/3* CN in progeny (Figure 1B and C)

which revealed that *pm2/3* CN ranges from 1 to 5 in progeny; all selfed KH004 contained only

one copy of *pm2/3*, in contrast to the parental KH004 that had multiple copy of *pm2/3*; copy

number variations (CNV) of *pm2/3* were detected inside IBD clusters, i.e., C18 and C21 (Figure

1C), suggesting rapid CNV evolution during *in vitro* blood culture and/or meiosis.

Assessment of novel allelic* combinations *of pfcrt and pm2/3

From our set of 104 unique recombinant progeny, parasites were grouped into 10 unique

clusters based on the inheritance of *pfcrt* and *pm2/3* alleles as well as *pm2/3* CN (Table S3).

Progeny carrying the Mal31 *pfcrt* and KH004 *pm2/3* alleles inherited all potential copies of

pm2/3, from 1 copy to 5 copies, whereas no progeny carry KH004 *pfcrt* with more than 3 copies

of *pm2/3.* Regardless of *pfcrt* inheritance, only 28% (13/46) of parasites which inherited the

KH004 *pm2/3* allele have >2 copies, indicating a potential fitness cost to carrying an excess

number of copies.

Genome-wide, we observed two large skews towards Mal31 alleles, one on

chromosome (chr) 7, and the second on chr 14 (Figure S1). The skewed region on chr 7 is

centered with *pfcrt*, which is known to carry high fitness cost with CQ resistance alleles (8). The

apicoplast ribosomal protein S10 (ARPS10, PF3D7_1460900) is located at the peak of chr 14

skewed region. Selection against the same *ARPS10* allele (Val127Met and Asp128His) has

been previously detected in two other independent crosses between Asian and African

parasites (38, 44), indicating a strong fitness cost carried by this allele from a Southeast Asia

parent (45).

We detected no skewed inheritance at *pm2/3*. We also saw no evidence for pairwise linkage

disequilibrium or co-inheritance between *pfcrt, kelch13*, or *pm2/3* alleles. However, we did

observe a significant (p = 0.006) association between inheritance of the KH004 ART-R *kelch13*

allele and multiple copies of *pm2/3* (Table S4).

PfCRT mutations are the major contributor to PPQ resistance

A major challenge to quantifying response to PPQ is the biphasic dose-response curve

which suggests incomplete parasite killing at higher drug concentrations as has been previously

reported in PPQ-R parasites (13). To comprehensively assess the impact of novel combinations

of *pfcrt* and *pm2/3* on parasite response to PPQ , we measured three unique phenotypes - PSA,

AUC, and a modified IC_{50} based on a limited-point dose response curve (LP-IC_{50}) in both

parents from the KH004 × Mal31 cross as well as 48 select progeny chosen to encompass all

available combinations of *pfcrt* and *pm2/3* CN.

IC_{50} is not traditionally used for phenotyping PPQ is due to the atypical biphasic drug

response curve generated by resistant parasites (13).We find that progeny parasites that

inherited the KH004 *pfcrt* allele and also multiple copies of *pm2/3* generate a biphasic dose

response curve, whereas progeny with only one copy of *pm2/3* and carrying the Mal31 *pfcrt*

allele produce a standard sigmoidal dose response (Figure S2A). This observed range of dose-

response curve shapes within our progeny set affirms our use of multiple trait measurements,

including a modification of standard IC_{50} values that rely on only the linear portion of the

sigmoidal dose-response (Figure S2B) and the biphasic portion of the curve to calculate AUCs.

We detected significant differences in PPQ susceptibilities strongly impacted by *pfcrt*

genotype for all three traits (Figure 2), with *p* values equal to 2.2e-13 for PSA, 1.0e-08 for AUC

and 1.0e-06 for LP-IC_{50} (Figure 2A). Additionally, QTL mapping of the PSA phenotype revealed

a highly significant 144 kb region on chr 7 (LOD score of 10.3) that centered on *pfcrt* (Figure

3A). This strong peak is responsible for 62% of the observed inherited phenotypic variance in

the PSA phenotype across the progeny set, whereas inheritance of this allele is responsible for

48% of the variance in the distribution of the AUC phenotype. QTL mapping of AUC and LP-IC_{50}

identify single QTLs on this chr 7 locus (LOD scores of 6.9 and 5.2 respectively) (Figure 3C,D).

pm2/3 CNV contributes epistatically to increased PPQ resistance

On top of the major segregation of PPQ responsiveness based on *pfcrt* inheritance, we

further detected that the trait distributions within *pfcrt* genotype sub-clusters associate with

pm2/3 CNV (Figure 2). However, unlike the progeny distribution shaped by the *pfcrt* allele

across all three phenotypes, the effect of *pm2/3* amplification varied depending on the *pfcrt*

allele and by phenotype measurement. For example, for PSA, we see an increase in the degree

of PPQ resistance associated with increased *pm2/3* CN only in parasites carrying the KH004

pfcrt allele (Figure 2A).

To examine the role of *pm2/3* CN independent of other loci, we utilized two sets of

parasites: isogenic (IBD) progeny and selfed parental parasites, are identical genotypes except

for differences in CN. These parasites are PPQ-S because they inherited the Mal31 *pfcrt* allele.

Notably, we did not recover isogenic progeny with the KH004 *pfcrt* allele but displaying different

CN. For both PSA and AUC, amplification from single copy *pm2/3* to 4 copies in a PPQ-R

(KH004 *pfcrt*) background significantly impacts both phenotypes, however, variation of the

number of multiple copies did not significantly impact either phenotype in individuals expressing

the Mal31 *pfcrt* allele (Figure 4).

Identification of additional loci contributing to PPQ response using QTL mapping

Due to the large and potentially masking effect of *pfcrt*, secondary scans of residual

variation were examined for all traits after statistically removing the effect from that chr 7 locus.

While no secondary effects were identified for either the PSA or LP-IC$_{50}$ phenotypes, residual

scans of the AUC phenotype identify a significant (LOD = 4.0) QTL on chr 10, which contains 61

genes (95% CI) (Table S5). Additionally, we used a two-dimensional, two-QTL genome scan to

identify and characterize significant interactions between loci that contribute to the various

resistance phenotypes (Table 1). Using a two-locus model, we identified a significant additive

interaction between loci on chr 7 and chr 10 (LOD additive: 10.7; 5% threshold) for the AUC

phenotype. Conversely, no QTL interactions were identified using the PSA phenotype.

PfCRT G367C likely confers PPQ resistance

Comprehensive analysis of KH004 × Mal31progeny points to *pfcrt* as the major

determinant of PPQ resistance, along with modulatory effects from other loci. Inheritance of the

pfcrt allele from KH004 was the major factor shared by PPQ-R parasites (Figures 2 and 4).

KH004 carries a Dd2-like *pfcrt*, differing from Dd2 by only a G367C substitution. While the Dd2

pfcrt haplotype confers resistance to CQ and other antimalarials, it is not associated with PPQ

resistance (25, 27–29). Thus, we suspect the G367C substitution is the major determinant of

PPQ-R in KH004. We further explored PPQ susceptibility among the parental strain, KH004, by

studying a selfed KH004 parasite (FG0305) that carries only one copy of *pm2/3），* Dd2*,* and

NHP4026 (recently isolated from parent at Thailand-Myanmar border (45) that carries a Dd2

pfcrt allele and one copy of *pm* 2/3). PSA shows clearly that both parasites (parental KH004 and

selfed KH004) with G367C are significantly PPQ-R, while those without G367C (*Dd*2 and

NHP4026) have PPQ-S PSA values similar to the African parasites NF54 and Mal31, which

have a 3D7-like *pfcrt* (Figure S4).

Reported drug resistance-associated mutations in PfCRT generally reside in one of the

10 transmembrane (TM) domains and, specifically, in regions that make up the central cavity of

the transporter (33, 46). The G367C substitution is distinct from experimentally validated PPQ-

R-associated mutations in that it resides in the digestive vacuole (DV)-exposed portion of TM9

and is not part of the core structure of the transporter's central cavity (27, 28, 33, 47, 48) (Figure

S3).

This novel G367C mutation was noted by Ross et al (25) to be present in SEA at low

prevalence in a small geographic focus. Because the KEL1/PLA1 lineage has continued to

spread throughout SEA coinciding with the continuing emergence of PPQ-R, we analyzed the

prevalence over time of this novel mutation using the MalariaGEN Pf7 dataset. Of 16,203 global

pfcrt haplotypes, 29 samples carried G367C (Table 3), first identified in 2012 and restricted to

Cambodia but has more recently emerged in neighboring countries Vietnam and Laos (Table

S6). Furthermore, 93% of these 29 clinical samples also carry multiple copies of *pm2/3*. We also

note the timeline of the emergence of these novel PPQ-R conferring *pfcrt* mutations: Kelch13-

mediated ART-R mutations emerged first while the rise of novel PPQ-R-associated *pfcrt*

mutations quickly followed the increase of samples expressing multiple copies of *pm2/3* (Figure

S5).

Impacts of pfcrt and pm2/3 inheritance on parasite blood-stage fitness

Our data across multiple resistance phenotypes support that *pfcrt* is the primary driver of

PPQ-R, with a boost to the degree of resistance resulting from increased copies of *pm2/3.* It has

also been well-documented that mutations in major resistance-associated genes can be

associated with a loss in parasite fitness (30, 49, 50). Deleterious fitness effects of the KH004

pfcrt allele are clearly evident from the reduced representation of this allele in the progeny

recovered from the genetic cross (Figure S1). For these novel resistance-associated genotypes

to increase in global prevalence in the absence of selection from widespread drug treatment,

parasites must preserve fitness either through low-cost resistance mutations or the acquisition

of secondary compensatory mutations that preserve or improve fitness. We conducted

competitive growth assays between progeny of the KH004 × Mal31 cross and both parental

parasites to assess blood-stage fitness differentials (Figure S6). Co-cultures were maintained

for a maximum of 40 d or until one parasite reached fixation (>90% of culture), with samples

being collected every 2 d. Progeny vs. parent competitions revealed that most progeny had a

relative fitness situated between two parents, while four progeny parasites were outside of the

parental fitness range (two more fit than KH004 and two less fit than Mal31), with no significant

clustering in fitness based on inherited *pfcrt* and *pm2/3* genotypes (Figure S6).

Based on the similar fitness phenotypes generated from competitions against the

parental parasites, a select subset of 18 progeny and both parents were set up in overlapping

competitions against common competitors, resulting in 212 competitive outcomes (win, lose, or

draw). The outcomes of these pairwise competitions were used to generate an Elo Rating for

each parasite using the *EloRating* package in R (51). Elo ratings provide a weighted ranking of

parasite fitness based on competitive growth outcome as well as the fitness ranking of the

competitor, with a higher Elo Rating representing a greater competitive fitness level. Elo Ratings

for these 20 parasites (parents plus progeny) provide a much finer resolution of fitness

differences between parasites of seemingly similar fitness. Unlike the competitions against both

parents, Elo Ratings provide a more continuous phenotype, but still do not cluster based on

pfcrt and *pm2/3* genotype (Figure 5A). However, QTL analysis using Elo phenotypes reveals a

highly significant (1% threshold) on chromosome 9 (LOD score of 4.5), which encompasses a

35 kb region containing 15 genes (Figure 5B). Parasites with the greatest competitive fitness

were associated with inheritance of this 35 kb region from the KH004 parent. Sequence

comparison between KH004 and Mal31 for these 15 genes identified 4 non-synonymous coding

SNPs in two genes: PF3D7_0934100 and PF3D7_0934700 (Table S7).

Discussion

In this study, we used a genetic cross between a multidrug-resistant Cambodian parasite

and a drug-sensitive isolate from Malawi to characterize the genetic contributors to PPQ-R.

Previous work on this subject has relied on naturally resistant parasites isolated from patience

or *in vitro* experimental manipulations to test specific mutations' ability to confer PPQ-R (13, 16,

20, 25). Use of a genetic cross allows us to track the inheritance and impact on phenotypes of

alleles previously associated with resistance, such as SNPs in *pfcrt* or amplification of *pm2/3*,

while also searching for additional trait-impacting loci across the genome . This classical genetic

approach demonstrates that the inheritance of the KH004 PfCRT (Dd2 + G367C) is the main

driver of PPQ-R, with the amplification of *pm2/3* serving an additive role amongst resistant

parasites.

The KH004 parent of this cross expresses a novel G367C substitution in PfCRT,

previously noted in SEA but not associated with PPQ-R. Through comparisons between KH004,

Dd2, and other SEA parasites, we observed a > 30-fold change in PSA between KH004 and

parasites expressing a standard Dd2-like PfCRT, indicating a direct connection between this

substitution and PPQ-R. While the G367C substitution has not previously been connected to

PPQ-R, this residue is within the binding site of the structurally similar chloroquine derivative

perfluorophenylazido biotinylated chloroquine (AzBCQ) (52). This finding of an association of

this residue with PPQ-R contributes to a broader understanding of PfCRT structure that

influences a wide range of drug susceptibilities, i.e. the range of *pfcrt* haplotypes shaped by

evolution to drug pressure. The major *pfcrt* haplotypes associated with PPQ-R have all evolved

on a 7G8 or Dd2 background, and transfection of these mutations into different *pfcrt*

backgrounds generates slight decreases in PPQ-R levels compared to those in a Dd2

background (28, 33).

Previously studied PPQ-R-associated PfCRT substitutions including T93S, H97Y,

C101F, F145I, M343L, and G353V were suspected due to their increasing prevalence in sites

using the DHA+PPQ (25, 33). The novel G367C substitution expressed by KH004 arose in

western Cambodia in 2012 and was found at a low abundance in the population. Since its initial

emergence, the Dd2+G367C haplotype has continued to persist at a consistently low

abundance but has spread to neighboring countries, Laos, and Vietnam, with the latter

accounting for the majority of identifications in recent years. As most of SEA has transitioned

from DHA+PPQ to alternative combination therapies as frontline treatments, Vietnam continues

to use PPQ, which might explain the recent increase in abundance of this novel haplotype within

the country (1, 53, 54). On the other hand, the introduction of this haplotype into Laos can

potentially be accounted for by a recent introduction of PPQ-R parasites into the country from

Cambodia (55).

Due to unusual PPQ-parasite interactions that are not effectively captured by standard

dose-response (IC_{50}) curves, we utilize three unique phenotypes to illuminate different aspects

of PPQ-R: AUC, limited point IC_{50}, and PSA. By using these different assays, we deconstruct

the resistance phenotype into three related but distinct biological readouts, survival under high

single dose PPQ (PSA), low dose growth inhibition (LP-IC_{50}), and shape of the dose-response

curve under high PPQ concentrations (AUC), to refine our search for the genetic factors

controlling these phenotypes (13, 18, 21, 22). QTL Mapping for each trait identified mutant *pfcrt*

as the main underlying contributor to resistance. Additionally, statistically removing the effect of

the *pfcrt* locus allowed for the identification of secondary contributors to the corresponding

phenotype. While the PSA and LP-IC_{50} residual scans did not identify any secondary loci, an

additional locus on chr 10 had a highly significant association with the AUC phenotype in the

absence of *pfcrt* (Figure 2). One of the genes with the strongest association within this region is

autophagy-related gene 18 (*atg18*). KH004 contains a single non-synonymous SNP in *atg18*,

leading to a Thr38Ile substitution, whereas Mal31 carries the WT 3D7-like variant of this gene,

highlighting a candidate mutation for further investigation. This protein localizes to the parasite

DV and has been associated with decreased susceptibility to several antimalarials such as

DHA, artemether (AM), and PPQ, while also being connected to increased survival under

nutrient deprivation (56–58). Due to its previous association with PPQ-R and the association

only with the AUC phenotype, *atg18* may contribute to the non-traditional dose-response curve

encountered when studying *in vitro* PPQ-R. Due to the fact that inheritance of the KH004 *pm2/3*

allele does not directly correspond to a set number of copies, the physical map used for QTL

analysis only associates allelic inheritance, not copy number, with each phenotype. Given that

pm2/3 CN is what impacts PPQ susceptibility, we do not observe any QTL encompassing this

locus on either or main or residual scans.

Previous studies hypothesized that instead of directly modulating PPQ-R, expressing

multiple copies of *pm2/3* amplification may play a compensatory role in parasite fitness (22).

This hypothesis, paired with extensive previous findings on the fitness costs of *pfcrt* resistance

mutations, led us to assess the competitive fitness levels of our progeny with various

combinations of *pfcrt* and *pm2/3* (29, 47, 49). By comparing progeny fitness to KH004 and

Mal31 parents, there found no significant clustering of *pfcrt/pm2/3* genotype combinations that

suggest a simple genetic determination of parent-derived fitness. Consequently, we leveraged

all-on-all competitions to generate Our novel implementation of Elo rankings can efficiently

compare head-to-head fitness levels across a large progeny set to generate a refined

phenotype for QTL mapping. Using this approach, we identified a highly significant association

between parasite fitness and a narrow 35 kb region on chr 9. This region is includes genes

involved in development, transcriptional regulation, and metabolism including two prioritized

candidate genes with coding variations between KH004 and Mal31. Neither gene has been

linked previously to parasite fitness.

Deeper knowledge of the genetic determinants of PPQ resistance can point to novel

control and prevention strategies. Our classical genetic approach informed our understanding

of the major drivers of resistance, including a novel *pfcrt* mutation, as well as identified

secondary genetic factors that contribute to parasite resistance and fitness. Incorporation of a

range of PPQ-R phenotype measures into our analysis revealed unique genetic regions and

their interactions associated with changes in each phenotype. By combining this novel genetic

cross with rigorous *in vitro* resistance and fitness phenotyping, we have identified the genetic

architecture which underlies decreased susceptibility to PPQ. While the DHA+PPQ ACT

continues to be used as a frontline ACT, it is crucial to understand the genetic underpinnings of

resistance.

References

1. World Health Organization. 2022. World malaria report 2022. Geneva.

2. White NJ, Pukrittayakamee S, Hien TT, Faiz MA, Mokuolu OA, Dondorp AM. 2014. Malaria. The
Lancet 383:723–735.

3. Blasco B, Leroy D, Fidock DA. 2017. Antimalarial drug resistance: linking Plasmodium falciparum
parasite biology to the clinic. Nat Med 23:917–928.

4. Payne D. 1987. Spread of chloroquine resistance in Plasmodium falciparum. Parasitol Today 3:241–
246.

5. Mita T, Tanabe K, Kita K. 2009. Spread and evolution of Plasmodium falciparum drug resistance.
Parasitol Int 58:201–209.

6. Imwong M, Suwannasin K, Kunasol C, Sutawong K, Mayxay M, Rekol H, Smithuis FM, Hlaing
TM, Tun KM, van der Pluijm RW, Tripura R, Miotto O, Menard D, Dhorda M, Day NPJ, White NJ,
Dondorp AM. 2017. The spread of artemisinin-resistant Plasmodium falciparum in the Greater
Mekong subregion: a molecular epidemiology observational study. Lancet Infect Dis 17:491–497.

7. Imwong M, Dhorda M, Myo Tun K, Thu AM, Phyo AP, Proux S, Suwannasin K, Kunasol C,
Srisutham S, Duanguppama J, Vongpromek R, Promnarate C, Saejeng A, Khantikul N, Sugaram R,
Thanapongpichat S, Sawangjaroen N, Sutawong K, Han KT, Htut Y, Linn K, Win AA, Hlaing TM,
van der Pluijm RW, Mayxay M, Pongvongsa T, Phommasone K, Tripura R, Peto TJ, von Seidlein
L, Nguon C, Lek D, Chan XHS, Rekol H, Leang R, Huch C, Kwiatkowski DP, Miotto O, Ashley
EA, Kyaw MP, Pukrittayakamee S, Day NPJ, Dondorp AM, Smithuis FM, Nosten FH, White NJ.
2020. Molecular epidemiology of resistance to antimalarial drugs in the Greater Mekong subregion:
an observational study. Lancet Infect Dis 20:1470–1480.

8.   Ecker A, Lehane AM, Clain J, Fidock DA. 2012. PfCRT and its role in antimalarial drug resistance.

Trends in Parasitology 28:504–514.

9.   Amato R, Pearson RD, Almagro-Garcia J, Amaratunga C, Lim P, Suon S, Sreng S, Drury E, Stalker

J, Miotto O, Fairhurst RM, Kwiatkowski DP. 2018. Origins of the current outbreak of multidrug-

resistant malaria in southeast Asia: a retrospective genetic study. Lancet Infect Dis 18:337–345.

10.   Hamilton WL, Amato R, van der Pluijm RW, Jacob CG, Quang HH, Thuy-Nhien NT, Hien TT,

Hongvanthong B, Chindavongsa K, Mayxay M, Huy R, Leang R, Huch C, Dysoley L, Amaratunga

C, Suon S, Fairhurst RM, Tripura R, Peto TJ, Sovann Y, Jittamala P, Hanboonkunupakarn B,

Pukrittayakamee S, Chau NH, Imwong M, Dhorda M, Vongpromek R, Chan XHS, Maude RJ,

Pearson RD, Nguyen T, Rockett K, Drury E, Gonçalves S, White NJ, Day NP, Kwiatkowski DP,

Dondorp AM, Miotto O. 2019. Evolution and expansion of multidrug-resistant malaria in southeast

Asia: a genomic epidemiology study. Lancet Infect Dis 19:943–951.

11.   MalariaGEN Plasmodium falciparum Community Project. 2016. Genomic epidemiology of

artemisinin resistant malaria. Elife 5:e08714.

12.   MalariaGEN, Abdel Hamid MM, Abdelraheem MH, Acheampong DO, Ahouidi A, Ali M,

Almagro-Garcia J, Amambua-Ngwa A, Amaratunga C, Amenga-Etego L, Andagalu B, Anderson T,

Andrianaranjaka V, Aniebo I, Aninagyei E, Ansah F, Ansah PO, Apinjoh T, Arnaldo P, Ashley E,

Auburn S, Awandare GA, Ba H, Baraka V, Barry A, Bejon P, Bertin GI, Boni MF, Borrmann S,

Bousema T, Bouyou-Akotet M, Branch O, Bull PC, Cheah H, Chindavongsa K, Chookajorn T,

Chotivanich K, Claessens A, Conway DJ, Corredor V, Courtier E, Craig A, D'Alessandro U, Dama

S, Day N, Denis B, Dhorda M, Diakite M, Djimde A, Dolecek C, Dondorp A, Doumbia S, Drakeley

C, Drury E, Duffy P, Echeverry DF, Egwang TG, Enosse SMM, Erko B, Fairhurst RM, Faiz A,

Fanello CA, Fleharty M, Forbes M, Fukuda M, Gamboa D, Ghansah A, Golassa L, Goncalves S,

Harrison GLA, Healy SA, Hendry JA, Hernandez-Koutoucheva A, Hien TT, Hill CA, Hombhanje

F, Hott A, Htut Y, Hussein M, Imwong M, Ishengoma D, Jackson SA, Jacob CG, Jeans J, Johnson KJ, Kamaliddin C, Kamau E, Keatley J, Kochakarn T, Konate DS, Konaté A, Kone A, Kwiatkowski DP, Kyaw MP, Kyle D, Lawniczak M, Lee SK, Lemnge M, Lim P, Lon C, Loua KM, Mandara CI, Marfurt J, Marsh K, Maude RJ, Mayxay M, Maïga-Ascofaré O, Miotto O, Mita T, Mobegi V, Mohamed AO, Mokuolu OA, Montgomery J, Morang'a CM, Mueller I, Murie K, Newton PN, Ngo Duc T, Nguyen T, Nguyen T-N, Nguyen Thi Kim T, Nguyen Van H, Noedl H, Nosten F, Noviyanti R, Ntui VN-N, Nzila A, Ochola-Oyier LI, Ocholla H, Oduro A, Omedo I, Onyamboko MA, Ouedraogo J-B, Oyebola K, Oyibo WA, Pearson R, Peshu N, Phyo AP, Plowe CV, Price RN, Pukrittayakamee S, Quang HH, Randrianarivelojosia M, Rayner JC, Ringwald P, Rosanas-Urgell A, Rovira-Vallbona E, Ruano-Rubio V, Ruiz L, Saunders D, Shayo A, Siba P, Simpson VJ, Sissoko MS, Smith C, Su X, Sutherland C, Takala-Harrison S, Talman A, Tavul L, Thanh NV, Thathy V, Thu AM, Toure M, Tshefu A, Verra F, Vinetz J, Wellems TE, Wendler J, White NJ, Whitton G, Yavo W, Van Der Pluijm RW. 2023. Pf7: an open dataset of Plasmodium falciparum genome variation in 20,000 worldwide samples. Wellcome Open Res 8:22.

13. Bopp S, Magistrado P, Wong W, Schaffner SF, Mukherjee A, Lim P, Dhorda M, Amaratunga C, Woodrow CJ, Ashley EA, White NJ, Dondorp AM, Fairhurst RM, Ariey F, Menard D, Wirth DF, Volkman SK. 2018. Plasmepsin II–III copy number accounts for bimodal piperaquine resistance among Cambodian Plasmodium falciparum. Nat Commun 9:1769.

14. Anderson TJC, Nair S, McDew-White M, Cheeseman IH, Nkhoma S, Bilgic F, McGready R, Ashley E, Pyae Phyo A, White NJ, Nosten F. 2017. Population Parameters Underlying an Ongoing Soft Sweep in Southeast Asian Malaria Parasites. Mol Biol Evol 34:131–144.

15. Ariey F, Witkowski B, Amaratunga C, Beghain J, Langlois A-C, Khim N, Kim S, Duru V, Bouchier C, Ma L, Lim P, Leang R, Duong S, Sreng S, Suon S, Chuor CM, Bout DM, Ménard S, Rogers WO, Genton B, Fandeur T, Miotto O, Ringwald P, Le Bras J, Berry A, Barale J-C, Fairhurst RM,

Benoit-Vical F, Mercereau-Puijalon O, Ménard D. 2014. A molecular marker of artemisinin-

resistant Plasmodium falciparum malaria. Nature 505:50–55.

16. Amato R, Lim P, Miotto O, Amaratunga C, Dek D, Pearson RD, Almagro-Garcia J, Neal AT, Sreng

S, Suon S, Drury E, Jyothi D, Stalker J, Kwiatkowski DP, Fairhurst RM. 2017. Genetic markers

associated with dihydroartemisinin–piperaquine failure in Plasmodium falciparum malaria in

Cambodia: a genotype–phenotype association study. The Lancet Infectious Diseases 17:164–173.

17. Tumwebaze PK, Conrad MD, Okitwi M, Orena S, Byaruhanga O, Katairo T, Legac J, Garg S,

Giesbrecht D, Smith SR, Ceja FG, Nsobya SL, Bailey JA, Cooper RA, Rosenthal PJ. 2022.

Decreased susceptibility of Plasmodium falciparum to both dihydroartemisinin and lumefantrine in

northern Uganda. Nat Commun 13:6353.

18. Duru V, Khim N, Leang R, Kim S, Domergue A, Kloeung N, Ke S, Chy S, Eam R, Khean C, Loch

K, Ken M, Lek D, Beghain J, Ariey F, Guerin PJ, Huy R, Mercereau-Puijalon O, Witkowski B,

Menard D. 2015. Plasmodium falciparum dihydroartemisinin-piperaquine failures in Cambodia are

associated with mutant K13 parasites presenting high survival rates in novel piperaquine in vitro

assays: retrospective and prospective investigations. BMC Med 13:305.

19. Amaratunga C, Lim P, Suon S, Sreng S, Mao S, Sopha C, Sam B, Dek D, Try V, Amato R,

Blessborn D, Song L, Tullo GS, Fay MP, Anderson JM, Tarning J, Fairhurst RM. 2016.

Dihydroartemisinin-piperaquine resistance in Plasmodium falciparum malaria in Cambodia: a

multisite prospective cohort study. Lancet Infect Dis 16:357–365.

20. Witkowski B, Duru V, Khim N, Ross LS, Saintpierre B, Beghain J, Chy S, Kim S, Ke S, Kloeung

N, Eam R, Khean C, Ken M, Loch K, Bouillon A, Domergue A, Ma L, Bouchier C, Leang R, Huy

R, Nuel G, Barale J-C, Legrand E, Ringwald P, Fidock DA, Mercereau-Puijalon O, Ariey F,

Ménard D. 2017. A surrogate marker of piperaquine-resistant Plasmodium falciparum malaria: a

phenotype–genotype association study. The Lancet Infectious Diseases 17:174–183.

21. Mukherjee A, Gagnon D, Wirth DF, Richard D. 2018. Inactivation of Plasmepsins 2 and 3

Sensitizes Plasmodium falciparum to the Antimalarial Drug Piperaquine. Antimicrob Agents

Chemother 62.

22. Loesbanluechai D, Kotanan N, de Cozar C, Kochakarn T, Ansbro MR, Chotivanich K, White NJ,

Wilairat P, Lee MCS, Gamo FJ, Sanz LM, Chookajorn T, Kümpornsin K. 2019. Overexpression of

plasmepsin II and plasmepsin III does not directly cause reduction in Plasmodium falciparum

sensitivity to artesunate, chloroquine and piperaquine. Int J Parasitol Drugs Drug Resist 9:16–22.

23. Parobek CM, Parr JB, Brazeau NF, Lon C, Chaorattanakawee S, Gosi P, Barnett EJ, Norris LD,

Meshnick SR, Spring MD, Lanteri CA, Bailey JA, Saunders DL, Lin JT, Juliano JJ. 2017. Partner-

Drug Resistance and Population Substructuring of Artemisinin-Resistant Plasmodium falciparum in

Cambodia. Genome Biol Evol 9:1673–1686.

24. Boonyalai N, Vesely BA, Thamnurak C, Praditpol C, Fagnark W, Kirativanich K, Saingam P,

Chaisatit C, Lertsethtakarn P, Gosi P, Kuntawunginn W, Vanachayangkul P, Spring MD, Fukuda

MM, Lon C, Smith PL, Waters NC, Saunders DL, Wojnarski M. 2020. Piperaquine resistant

Cambodian Plasmodium falciparum clinical isolates: in vitro genotypic and phenotypic

characterization. Malar J 19:269.

25. Ross LS, Dhingra SK, Mok S, Yeo T, Wicht KJ, Kümpornsin K, Takala-Harrison S, Witkowski B,

Fairhurst RM, Ariey F, Menard D, Fidock DA. 2018. Emerging Southeast Asian PfCRT mutations

confer Plasmodium falciparum resistance to the first-line antimalarial piperaquine. Nat Commun

9:3314.

26.   Dondorp AM. 2017. New genetic marker for piperaquine resistance in Plasmodium falciparum. The

544   Lancet Infectious Diseases 17:119–121.

545   27.   Okombo J, Mok S, Qahash T, Yeo T, Bath J, Orchard LM, Owens E, Koo I, Albert I, Llinás M,

546   Fidock DA. 2022. Piperaquine-resistant PfCRT mutations differentially impact drug transport,

547   hemoglobin catabolism and parasite physiology in Plasmodium falciparum asexual blood stages.

548   PLoS Pathog 18:e1010926.

549   28.   Wicht KJ, Small-Saunders JL, Hagenah LM, Mok S, Fidock DA. 2022. Mutant PfCRT Can Mediate

550   Piperaquine Resistance in African Plasmodium falciparum With Reduced Fitness and Increased

551   Susceptibility to Other Antimalarials. J Infect Dis 226:2021–2029.

552   29.   Small-Saunders JL, Hagenah LM, Wicht KJ, Dhingra SK, Deni I, Kim J, Vendome J, Gil-Iturbe E,

553   Roepe PD, Mehta M, Mancia F, Quick M, Eppstein MJ, Fidock DA. 2022. Evidence for the early

554   emergence of piperaquine-resistant Plasmodium falciparum malaria and modeling strategies to

555   mitigate resistance. PLoS Pathog 18:e1010278.

556   30.   Dhingra SK, Gabryszewski SJ, Small-Saunders JL, Yeo T, Henrich PP, Mok S, Fidock DA. 2019.

557   Global Spread of Mutant PfCRT and Its Pleiotropic Impact on Plasmodium falciparum Multidrug

558   Resistance and Fitness. mBio 10:e02731-18.

559   31.   van der Pluijm RW, Imwong M, Chau NH, Hoa NT, Thuy-Nhien NT, Thanh NV, Jittamala P,

560   Hanboonkunupakarn B, Chutasmit K, Saelow C, Runjarern R, Kaewmok W, Tripura R, Peto TJ,

561   Yok S, Suon S, Sreng S, Mao S, Oun S, Yen S, Amaratunga C, Lek D, Huy R, Dhorda M,

562   Chotivanich K, Ashley EA, Mukaka M, Waithira N, Cheah PY, Maude RJ, Amato R, Pearson RD,

563   Gonçalves S, Jacob CG, Hamilton WL, Fairhurst RM, Tarning J, Winterberg M, Kwiatkowski DP,

564   Pukrittayakamee S, Hien TT, Day NP, Miotto O, White NJ, Dondorp AM. 2019. Determinants of

565   dihydroartemisinin-piperaquine treatment failure in Plasmodium falciparum malaria in Cambodia,

Thailand, and Vietnam: a prospective clinical, pharmacological, and genetic study. Lancet Infect
Dis 19:952–961.

32. Dhingra SK, Small-Saunders JL, Ménard D, Fidock DA. 2019. Plasmodium falciparum resistance to
piperaquine driven by PfCRT. The Lancet Infectious Diseases 19:1168–1169.

33. Kim J, Tan YZ, Wicht KJ, Erramilli SK, Dhingra SK, Okombo J, Vendome J, Hagenah LM,
Giacometti SI, Warren AL, Nosol K, Roepe PD, Potter CS, Carragher B, Kossiakoff AA, Quick M,
Fidock DA, Mancia F. 2019. Structure and drug resistance of the Plasmodium falciparum
transporter PfCRT. Nature 576:315–320.

34. Pelleau S, Moss EL, Dhingra SK, Volney B, Casteras J, Gabryszewski SJ, Volkman SK, Wirth DF,
Legrand E, Fidock DA, Neafsey DE, Musset L. 2015. Adaptive evolution of malaria parasites in
French Guiana: Reversal of chloroquine resistance by acquisition of a mutation in *pfcrt*. Proc Natl
Acad Sci USA 112:11672–11677.

35. Miotto O, Amato R, Ashley EA, MacInnis B, Almagro-Garcia J, Amaratunga C, Lim P, Mead D,
Oyola SO, Dhorda M, Imwong M, Woodrow C, Manske M, Stalker J, Drury E, Campino S,
Amenga-Etego L, Thanh T-NN, Tran HT, Ringwald P, Bethell D, Nosten F, Phyo AP,
Pukrittayakamee S, Chotivanich K, Chuor CM, Nguon C, Suon S, Sreng S, Newton PN, Mayxay M,
Khanthavong M, Hongvanthong B, Htut Y, Han KT, Kyaw MP, Faiz MA, Fanello CI, Onyamboko
M, Mokuolu OA, Jacob CG, Takala-Harrison S, Plowe CV, Day NP, Dondorp AM, Spencer CCA,
McVean G, Fairhurst RM, White NJ, Kwiatkowski DP. 2015. Genetic architecture of artemisinin-
resistant Plasmodium falciparum. Nat Genet 47:226–234.

36. Wootton JC, Feng X, Ferdig MT, Cooper RA, Mu J, Baruch DI, Magill AJ, Su X-Z. 2002. Genetic
diversity and chloroquine selective sweeps in Plasmodium falciparum. Nature 418:320–323.

37.   Tirrell AR, Vendrely KM, Checkley LA, Davis SZ, McDew-White M, Cheeseman IH, Vaughan

589   AM, Nosten FH, Anderson TJC, Ferdig MT. 2019. Pairwise growth competitions identify relative

590   fitness relationships among artemisinin resistant Plasmodium falciparum field isolates. Malaria

591   Journal 18.

592   38.   Li X, Kumar S, McDew-White M, Haile M, Cheeseman IH, Emrich S, Button-Simons K, Nosten F,

593   Kappe SHI, Ferdig MT, Anderson TJC, Vaughan AM. 2019. Genetic mapping of fitness

594   determinants across the malaria parasite Plasmodium falciparum life cycle. PLoS Genet

595   15:e1008453.

596   39.   Amambua-Ngwa A, Button-Simons KA, Li X, Kumar S, Brenneman KV, Ferrari M, Checkley LA,

597   Haile MT, Shoue DA, McDew-White M, Tindall SM, Reyes A, Delgado E, Dalhoff H, Larbalestier

598   JK, Amato R, Pearson RD, Taylor AB, Nosten FH, D'Alessandro U, Kwiatkowski D, Cheeseman

599   IH, Kappe SHI, Avery SV, Conway DJ, Vaughan AM, Ferdig MT, Anderson TJC. 2023.

600   Chloroquine resistance evolution in Plasmodium falciparum is mediated by the putative amino acid

601   transporter AAT1. Nat Microbiol https://doi.org/10.1038/s41564-023-01377-z.

602   40.   Mairet-Khedim M, Leang R, Marmai C, Khim N, Kim S, Ke S, Kauy C, Kloeung N, Eam R, Chy S,

603   Izac B, Mey Bouth D, Dorina Bustos M, Ringwald P, Ariey F, Witkowski B. 2021. Clinical and In

604   Vitro Resistance of Plasmodium falciparum to Artesunate-Amodiaquine in Cambodia. Clin Infect

605   Dis 73:406–413.

606   41.   Vaughan AM, Pinapati RS, Cheeseman IH, Camargo N, Fishbaugher M, Checkley LA, Nair S,

607   Hutyra CA, Nosten FH, Anderson TJC, Ferdig MT, Kappe SHI. 2015. Plasmodium falciparum

608   genetic crosses in a humanized mouse model. Nat Methods 12:631–633.

609   42.   Miles A, Iqbal Z, Vauterin P, Pearson R, Campino S, Theron M, Gould K, Mead D, Drury E,

610   O'Brien J, Ruano Rubio V, MacInnis B, Mwangi J, Samarakoon U, Ranford-Cartwright L, Ferdig

M, Hayton K, Su X, Wellems T, Rayner J, McVean G, Kwiatkowski D. 2016. Indels, structural

variation, and recombination drive genomic diversity in *Plasmodium falciparum*. Genome Res

26:1288–1299.

43. Brenneman KV, Li X, Kumar S, Delgado E, Checkley LA, Shoue DA, Reyes A, Abatiyow BA,

Haile MT, Tripura R, Peto T, Lek D, Button-Simons KA, Kappe SHI, Dhorda M, Nosten F,

Nkhoma SC, Cheeseman IH, Vaughan AM, Ferdig MT, Anderson TJC. 2022. Optimizing bulk

segregant analysis of drug resistance using Plasmodium falciparum genetic crosses conducted in

humanized mice. iScience 25:104095.

44. Kumar S, Li X, McDew-White M, Reyes A, Delgado E, Sayeed A, Haile MT, Abatiyow BA,

Kennedy SY, Camargo N, Checkley LA, Brenneman KV, Button-Simons KA, Duraisingh MT,

Cheeseman IH, Kappe SHI, Nosten F, Ferdig MT, Vaughan AM, Anderson TJC. 2022. A Malaria

Parasite Cross Reveals Genetic Determinants of Plasmodium falciparum Growth in Different

Culture Media. Front Cell Infect Microbiol 12:878496.

45. Button-Simons KA, Kumar S, Carmago N, Haile MT, Jett C, Checkley LA, Kennedy SY, Pinapati

RS, Shoue DA, McDew-White M, Li X, Nosten FH, Kappe SH, Anderson TJC, Romero-Severson

J, Ferdig MT, Emrich SJ, Vaughan AM, Cheeseman IH. 2021. The power and promise of genetic

mapping from Plasmodium falciparum crosses utilizing human liver-chimeric mice. Commun Biol

4:734.

46. Willems A, Kalaw A, Ecer A, Kotwal A, Roepe LD, Roepe PD. 2023. Structures of Plasmodium

falciparum Chloroquine Resistance Transporter (PfCRT) Isoforms and Their Interactions with

Chloroquine. Biochemistry 62:1093–1110.

47. Dhingra SK, Redhi D, Combrinck JM, Yeo T, Okombo J, Henrich PP, Cowell AN, Gupta P,

Stegman ML, Hoke JM, Cooper RA, Winzeler E, Mok S, Egan TJ, Fidock DA. 2017. A Variant

PfCRT Isoform Can Contribute to *Plasmodium falciparum* Resistance to the First-Line Partner

Drug Piperaquine. mBio 8:e00303-17.

48. Gomez GM, D'Arrigo G, Sanchez CP, Berger F, Wade RC, Lanzer M. 2023. PfCRT mutations

conferring piperaquine resistance in falciparum malaria shape the kinetics of quinoline drug binding

and transport. PLoS Pathog 19:e1011436.

49. Osman ME, Mockenhaupt FP, Bienzle U, Elbashir MI, Giha HA. 2007. Field-based evidence for

linkage of mutations associated with chloroquine (pfcrt/pfmdr1) and sulfadoxine–pyrimethamine

(pfdhfr/pfdhps) resistance and for the fitness cost of multiple mutations in P. falciparum. Infection,

Genetics and Evolution 7:52–59.

50. Nair S, Li X, Arya GA, McDew-White M, Ferrari M, Nosten F, Anderson TJC. 2018. Fitness Costs

and the Rapid Spread of kelch13-C580Y Substitutions Conferring Artemisinin Resistance.

Antimicrob Agents Chemother 62:e00605-18.

51. Neumann C, Kulik L. 2014. Package"EloRating" (0.43).

52. Lekostaj JK, Natarajan JK, Paguio MF, Wolf C, Roepe PD. 2008. Photoaffinity labeling of the

Plasmodium falciparum chloroquine resistance transporter with a novel perfluorophenylazido

chloroquine. Biochemistry 47:10394–10406.

53. Hassett MR, Roepe PD. 2019. Origin and Spread of Evolving Artemisinin-Resistant Plasmodium

falciparum Malarial Parasites in Southeast Asia. The American Journal of Tropical Medicine and

Hygiene 101:1204–1211.

54. Sattabongkot J, Cui L, Bantuchai S, Chotirat S, Kaewkungwal J, Khamsiriwatchara A, Kiattibutr K,

Kyaw MP, Lawpoolsri S, Linn NYY, Menezes L, Miao J, Nguitragool W, Parker D, Prikchoo P,

Roobsoong W, Sa-Angchai P, Samung Y, Sirichaisinthop J, Sriwichai P, Suk-Uam K, Thammapalo

S, Wang B, Zhong D. 2022. Malaria Research for Tailored Control and Elimination Strategies in the

Greater Mekong Subregion. Am J Trop Med Hyg 107:152–159.

55.   Wasakul V, Disratthakit A, Mayxay M, Chindavongsa K, Sengsavath V, Thuy-Nhien N, Pearson

RD, Phalivong S, Xayvanghang S, Maude RJ, Gonçalves S, Day NP, Newton PN, Ashley EA,

Kwiatkowski DP, Dondorp AM, Miotto O. 2022. Malaria outbreak in Laos driven by a selective

sweep for Plasmodium falciparum kelch13 R539T mutants: a genetic epidemiology analysis. Lancet

Infect Dis S1473-3099(22)00697–1.

56.   Sudhakar R, Das D, Thanumalayan S, Gorde S, Sijwali PS. 2021. Plasmodium falciparum Atg18

localizes to the food vacuole via interaction with the multi-drug resistance protein 1 and

phosphatidylinositol 3-phosphate. Biochem J 478:1705–1732.

57.   Wang Z, Cabrera M, Yang J, Yuan L, Gupta B, Liang X, Kemirembe K, Shrestha S, Brashear A, Li

X, Porcella SF, Miao J, Yang Z, Su X-Z, Cui L. 2016. Genome-wide association analysis identifies

genetic loci associated with resistance to multiple antimalarials in Plasmodium falciparum from

China-Myanmar border. Sci Rep 6:33891.

58.   Breglio KF, Amato R, Eastman R, Lim P, Sa JM, Guha R, Ganesan S, Dorward DW, Klumpp-

Thomas C, McKnight C, Fairhurst RM, Roberts D, Thomas C, Simon AK. 2018. A single

nucleotide polymorphism in the Plasmodium falciparum atg18 gene associates with artemisinin

resistance and confers enhanced parasite survival under nutrient deprivation. Malar J 17:391.

59.   Manske M, Miotto O, Campino S, Auburn S, Almagro-Garcia J, Maslen G, O'Brien J, Djimde A,

Doumbo O, Zongo I, Ouedraogo J-B, Michon P, Mueller I, Siba P, Nzila A, Borrmann S, Kiara SM,

Marsh K, Jiang H, Su X-Z, Amaratunga C, Fairhurst R, Socheat D, Nosten F, Imwong M, White

NJ, Sanders M, Anastasi E, Alcock D, Drury E, Oyola S, Quail MA, Turner DJ, Ruano-Rubio V,

Jyothi D, Amenga-Etego L, Hubbart C, Jeffreys A, Rowlands K, Sutherland C, Roper C, Mangano

679 V, Modiano D, Tan JC, Ferdig MT, Amambua-Ngwa A, Conway DJ, Takala-Harrison S, Plowe

680 CV, Rayner JC, Rockett KA, Clark TG, Newbold CI, Berriman M, MacInnis B, Kwiatkowski DP.

681 2012. Analysis of Plasmodium falciparum diversity in natural infections by deep sequencing.

682 Nature 487:375–379.

683 60. Schaffner SF, Taylor AR, Wong W, Wirth DF, Neafsey DE. 2018. hmmIBD: software to infer

684 pairwise identity by descent between haploid genotypes. Malar J 17:196.

685 61. Vendrely KM, Kumar S, Li X, Vaughan AM. 2020. Humanized Mice and the Rebirth of Malaria

686 Genetic Crosses. Trends Parasitol 36:850–863.

687 62. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R,

688 Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-Paredes

689 B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger M,

690 Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW, Kavukcuoglu K,

691 Kohli P, Hassabis D. 2021. Highly accurate protein structure prediction with AlphaFold. Nature

692 596:583–589.

693 63. Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, Yuan D, Stroe O, Wood G,

694 Laydon A, Žídek A, Green T, Tunyasuvunakool K, Petersen S, Jumper J, Clancy E, Green R, Vora

695 A, Lutfi M, Figurnov M, Cowie A, Hobbs N, Kohli P, Kleywegt G, Birney E, Hassabis D, Velankar

696 S. 2022. AlphaFold Protein Structure Database: massively expanding the structural coverage of

697 protein-sequence space with high-accuracy models. Nucleic Acids Res 50:D439–D444.

698 64. Davis SZ, Singh PP, Vendrely KM, Shoue DA, Checkley LA, McDew-White M, Button-Simons

699 KA, Cassady Z, Sievert MAC, Foster GJ, Nosten FH, Anderson TJC, Ferdig MT. 2020. The

700 extended recovery ring-stage survival assay provides a superior association with patient clearance

701 half-life and increases throughput. Malar J 19:54.

702   65.   Neumann C, Duboscq J, Dubuc C, Ginting A, Irwan AM, Agil M, Widdig A, Engelhardt A. 2011.

703   Assessing dominance hierarchies: validation and advantages of progressive evaluation with Elo-

704   rating. Animal Behaviour 82:911–921.

705   66.   Sen Ś, Churchill GA. 2001. A statistical framework for quantitative trait mapping. Genetics

706   159:371–387.

707   67.   Broman KW, Wu H, Sen S, Churchill GA. 2003. R/qtl: QTL mapping in experimental crosses.

708   Bioinformatics 19:889–890.

709   68.   Churchill GA, Doerge RW. 1994. Empirical threshold values for quantitative trait mapping.

710   Genetics 138:963–971.

Figure 1: Plasmepsin II/III (pm 2/3) copy number variations in parents and progeny. *A) IGV plot showing copy*
number variations in parental parasites at pm gene region with the pm repeat unit labeled with a red box. B) pm 2/3
copy number of parental parasites and recombinant and selfed progeny. Mal31.IBD indicates recombinant progeny
with pm allele inherited from Mal31; KH004.IBD represents progeny with pm from KH004. Parental parasites Mal31
and KH004 are labeled with red diamonds. C) rapid evolution of pm 2/3 copy numbers in identical-by-descent (IBD)
progeny. C18 and C21 are two IBD progeny groups. Of which, C18 contains 5 clones, with 4 of them containing 5
copies and 1 of them containing 3 copies of pm; while C21 contains 4 clones, with 3 of them contain 4 copies, and 1
contains 3 copies of pm. D) Nanopore long reads that cover the pm 2/3 CNV and flanking regions. The boundaries of
the repeat unit are labeled with [red square brackets].

Figure 2: PfCRT mutations determine the majority of the variation in PPQ response. *Progeny of the KH004 × Mal31 cross were grouped based on inheritance of parental allele at pfcrt (KH004 or Mal31) and pm2/3 copy number and were phenotyped using 3 different measurements of resistance: PSA (A), AUC (B), and LP-IC$_{50}$ (C). We find that with all 3 phenotypes, inheritance of the KH004 pfcrt allele leads to significantly decreased susceptibility to PPQ. Additionally, we observed that increased copies of pm2/3 led to additional decreases in susceptibility, however the impact on CNV with respect to pfcrt varied by phenotype.*

Figure 3: QTL mapping identifies major and secondary loci associated with PPQ-R. *We performed QTL mapping based on parasite response to PPQ using our 3 metrics of resistance: PSA, AUC, and LP-IC$_{50}$. For each of the 3 phenotypes, a region of chr 7 centered around pfcrt was most strongly associated with the variation associated with each phenotype. This further supports that pfcrt is the major factor associated with decreased susceptibility. To eliminate the potentially masking effect of the chr 7 QTL, residual scans of variation were performed. These scans identified a secondary QTL on chr 10 which was associated with AUC. Pairing QTL mapping with resistance phenotyping further supports the strong influence on pfcrt as well as identifies secondary contributors to PPQ-R.*

Phenotype	Pairs of Loci	Position locus 1	Position locus 2	LOD interaction (epistasis)	LOD additive
AUC	Chr 7: Chr 10	357253	485200	1.39	**10.7**

Table 1: Additive interactions influencing PPQ-R phenotypes. Two-dimensional genome scan identifies interacting loci on chromosomes 7 and 10 contribute to the AUC phenotype . We have identified that these two loci interact additively in their contributions to AUC. We identified no interacting loci associated with PSA phenotype. Significant LOD scores are bolded and have surpassed the 5% threshold.

Figure 4: Effect of plasmepsin II/III copy number on PPQ-R is dependent on genetic background. Four groups of isogenic parasites which differ only in pm CN were assayed using A) PSA and B) AUC to identify the isolated role of pm CNV independent of genetic background. Of which, FG0175 is a selfed Mal31 progeny; FG0098 and FG0191 were IBD; FG0132 and FG0288 were IBD; FG0305 is a selfed KH004 progeny. We found that CNV in a Mal31-pfcrt background does not significantly impact either phenotype.

Country	% of multicopy *pm2/3* samples	# of G367C samples	# of documented PPQ-R mutations[*]
Cambodia (n = 1267)	42.1%	12	327
Vietnam (n= 1404)	39.0%	14[a]	676
Laos (n= 991)	16.4%	3	169
Global population (n = 16203)	7.90%	29	1204

Table 2: Prevalence of G367C mutation in SEA. *Based on available data from MalariaGEN Pf7, this novel polymorphism has been observed throughout SEA, although at low levels, since 2012. Apart from two samples, this mutation is found in PPQ-R samples and is also accompanied by increased copies of pm2/3.*

[a] Two samples from Vietnam carried a single pm2/3 copy and was deemed sensitive independent of pfcrt polymorphisms.

[*] PPQ-R-associated PfCRT substitutions include: T93S, H97Y, F145I, I218F, M343L, G353V. The C350R substitution was omitted as it arose independently out of a 7G8 background and has not been identified on top of a Dd2 background.

Figure 5: QTL mapping of fitness-associated loci using Elo rating phenotype. A) *Mean Elo rating of 18 progeny + 2 parents of the KH004 ✕ Mal31 genetic cross based on 50 permutations of the order of competitions. Elo rating was calculated using the R package EloRating and was based on the each parasite's competitive growth assay outcome. Phenotype distribution is ordered from lowest Elo (least fit) on the left to highest Elo (most fit) on the right. **B)** QTL scan based on Elo phenotype reveals a highly significant peak (1% threshold) on chromosome 9. This peak encompasses a 35,264 bp region which contains 15 genes. Genes identified from this QTL scan are proposed to directly contribute to parasite fitness.*

Figure S1: Inheritance breakdown of KH004xMal31 genetic cross. *A) From this cross, we see two strong skews in inheritance on chromosomes 7 and 14, both favoring the Mal31 allele. This selective inheritance has previously been observed in other genetic crosses between Southeast Asian and African parasites. The chromosome 7 peak is centered around pfcrt and therefore potentially represents a fitness cost associated with the inheritance of the KH004 allele at these positions. B) Aside from the selective inheritance on chromosomes 7 and 14, we observe no significant co-inheritance patterns between pfcrt, pm2/3, or kelch13. Inheritance of the KH004 allele is denoted in blue and the Mal31 allele in orange.*

Figure S2: Shape of dose-response curve changes based on pfcrt and pm2/3 genotype. Traditional IC_{50} measurements have been a challenge with PPQ due to the biphasic dose-response curve observed in PPQ-R parasites. A) We have found inheritance of the KH004 pfcrt allele and multiple copies of pm2/3 are required for producing a biphasic curve. Progeny inheriting the Mal31 pfcrt allele produce do not produce a biphasic curve, regardless of CNV. Additionally, all parasites expressing a single copy of pm2/3 produce a sigmoidal curve. To account for these various dose-response curve shapes, AUC is used as a measure of resistance, with a greater AUC indicating decreased PPQ susceptibility. B) By using a limited number of points in our dose response curve (LP-IC_{50}), we can eliminate the secondary peak in the curve and measure an IC50 independent of original curve shape.

Figure S3: Structural position of G367C Substitution: A) Alphafold model of Dd2 PfCRT with previously documented in vitro PPQ-R associated mutations identified (yellow). Additional highlighted AA substitutions and the novel G367C substitution carried by KH004 (red).

pm CN	1	1	2	1	4	1
pfcrt allele		3D7		Dd2	Dd2+ G367C	

Figure S4: PfCRT G367C directly confers PPQ-resistance
PSA between KH004, FG0305 (selfed KH004 with single copy pm) and two other southeast Asian parasites (Dd2 and NH4026) which carry a Dd2-like PfCRT allele, and two African parasites carrying a WT PfCRT identifies that a Dd2 PfCRT background alone is not sufficient for PPQ-R. This further supports the conclusion that PfCRT G367C directly confers PPQ resistance.

Figure S5: Rise of pm2/3 amplification and novel pfcrt mutations associated with PPQ-R in the GMS. Using available data from MalariaGEN Pf7 release (n = 3,359), samples from Cambodia, Vietnam, and Laos were analyzed for the association between pm2/3 amplification and novel PPQ-R conferring pfcrt mutations. The rise in novel pfcrt PPQ-R mutations immediately following amplification of pm2/3 further confirms the important roles of both pm2/3 and pfcrt in the evolution of PPQ-R.

Figure S6: Competitive growth assessment between progeny and parents. *Head-to-head competitive growth assays were set up between progeny and parents and were ranked based on competitive outcome. The fitness level of most progeny fell in between the two parents and no apparent clustering was observed between relative fitness and pfcrt or pm2/3 genotype.*

Methods

Ethics approval and consent to participate

The study was performed in strict accordance with the recommendations in the Guide for the

Care and Use of Laboratory Animals of the National Institutes of Health (NIH), USA. To this end,

the Seattle Children's Research Institute (SCRI) has an Assurance from the Public Health

Service (PHS) through the Office of Laboratory Animal Welfare (OLAW) for work approved by

its Institutional Animal Care and Use Committee (IACUC). All of the work carried out in this

study was specifically reviewed and approved by the SCRI IACUC.

Preparation of the genetic crosses

We generated the crosses using FRG NOD huHep mice with human chimeric livers and *A. stephensi* mosquitoes as described by Vaughan *et al. (41).* Two individual recombinant pools

were generated for each cross, by using different cages of infected mosquitoes. To start each

cross, gametocytes from both parental parasite strains were diluted to 0.5% gametocytemia in a

human serum erythrocyte mix, to generate infectious blood meals (IBMs). IBMs from each

parent were mixed at equal proportions and fed to three cages of mosquitos (150 per cage).

We examined the mosquito infection rate and oocyst number per infected mosquito 7-10 days

post-feeding. Fifteen mosquitoes were randomly picked from each cage and dissected under

microscopy. Sporozoites were isolated from infected mosquito salivary glands and 2-4 million

sporozoites from each cage of mosquitoes were injected into three FRG huHep mice (one cage

per mouse), intravenously. To allow the liver stage-to-blood stage transition, mice are infused

with human erythrocytes six and seven days after sporozoite injection. Four hours after the

second infusion, the mice are euthanized and exsanguinated to isolate the circulating ring stage

P. falciparum-infected human erythrocytes. The parasites from each mouse constitute the initial

recombinant pools of recombinant progeny for genetic mapping experiments. We maintained

the initial pools in AlbuMAX supplemented RPMI media; we genome sequenced aliquots from

each pool to check allele frequencies from both parents.

Library preparation and sequencing

We used Qiagen DNA mini kit to extract and purify the genomic DNA, and Quant-iT™

PicoGreen® Assay (Invitrogen) to quantify the amount of DNA. For samples with less than 50ng

DNA obtained, whole genome amplification (WGA) was performed before NGS library

preparation. WGA reactions were performed following Nair et al (Nair et al., 2014). Each 25 µl

reaction contained at least 5ng of *Plasmodium* DNA, 1× BSA (New England Biolabs), 1 mM

dNTPs (New England Biolabs), 3.5 µM of Phi29 Random Hexamer Primer, 1× Phi29 reaction

buffer (New England Biolabs), and 15 units of Phi29 polymerase (New England Biolabs). We

used a PCR machine (SimpliAmp, Applied Biosystems) programmed to run a "stepdown"

protocol: 35 °C for 10 min, 34 °C for 10 min, 33 °C for 10 min, 32 °C for 10 min, 31 °C for 10

min, 30 °C for 6 h then heating at 65 °C for 10 min to inactivate the enzymes prior to cooling to 4

°C. Samples were cleaned with AMPure XP Beads (Beckman Coulter) at a 1:1 ratio. We

constructed next generation sequencing (NGS) libraries using 50-100 ng DNA or WGA product

following the KAPA HyperPlus Kit protocol with 3-cycle of PCR. All libraries were sequenced at

150bp pair-end using Illumina Novaseq S4 or Hiseq X sequencers. We sequenced all bulk

samples to a minimum coverage of 100×.

Mapping and genotyping

We individually mapped whole-genome sequencing reads for each library against the *P.*

falciparum 3D7 reference genome (PlasmoDB, release32) using the alignment algorithm BWA

mem (http://bio-bwa.sourceforge.net/) under the default parameters. The resulting alignments

were then converted to SAM format, sorted to BAM format, and deduplicated using picard tools

v2.0.1 (http://broadinstitute.github.io/picard/). We used Genome Analysis Toolkit GATK v3.7

(https://software.broadinstitute.org/gatk/) to recalibrate the base quality score based on a set of

verified known variants (42).

After alignment, we excluded the highly variable genome regions (subtelomeric repeats,

hypervariable regions and centromeres) and only performed genotype calling in the 21 Mb core

genome (defined in (42)). We called variants for each sample using HaplotypeCaller, and calls

from every 100 samples were merged using CombineGVCFs with default parameters. Variants

were further called at all sample-level using GenotypeGVCFs, with parameters: --

max_alternate_alleles 6 --variant_index_type LINEAR --variant_index_parameter 128000 --

sample_ploidy 2 -nt 20. We further filtered the variants calls by calculating the recalibrated

variant quality scores (VQSR) of genotypes from parental parasites. Loci with VQSR less than 1

or not distinguishable between two parents were removed from further analysis. The variants in

VCF format were annotated for predicted functional effect on genes and proteins using snpEff

v4.3 (https://pcingola.github.io/SnpEff/) with 3D7 (PlasmoDB, release32) as the reference.

Cloned progeny analysis

To identify unsuccessfully cloned progeny, we measured the multiplicity of progeny samples

with F_{WS} (59), which is configured in moimix (https://github.com/bahlolab/moimix). Samples with

F_{WS} < 0.9 were assumed to be non-clonal and were removed from further analysis. Allele

frequencies across the genome were also plotted and manual inspected to detect further

possible mix infections.

We calculated Identity-by-descent (IBD) between clonal progeny and parents with hmmIBD

(60) under default parameters. The proportions of shared IBD were used to determine

relatedness among parental and progeny parasites and to identify genome regions inherited

from each parent: 1) progeny with >90% shared IBD with either of the parents were assumed to

result from selfing; 2) progeny with <90% shared IBD with both parents were defined as

recombinants; 3) recombinant progeny with >90% IBD against each other were defined as non-

unique clones (61).

Nanopore sequencing of parental parasites

We used an optimized a Phenol:Chloroform DNA isolation protocol that generates 500 ng to

3 µg large molecular weight genomic DNA from 40 mL of in vitro blood cultures. In brief, (i)

short, fragmented DNA (<60kb) is removed using Small Read Eliminator Kit (Nanopore); (ii)

after DNA clean-up, we prepared sequencing libraries with Ligation Sequencing Kit (Nanopore),

which adds barcode to each sample. We used the Nanopore MinION Mk1C sequencer to

generate long reads. We obtained 3.5 Gb (150× genome coverage) and 2.5 Gb (105×) of long-

read sequencing data for Mal31 and KH004. Fifty percent of the data obtained comprise reads

with length >62Kb (N50 = 62kb) with the longest read of 485 kb.

We mapped the nanopore long reads to *plasmepsin* gene sequences using *blastn*

(https://blast.ncbi.nlm.nih.gov/Blast.cgi). Reads that cover all three *plasmepsin* genes and

>10kb flanking regions on both sides were then extracted and compared to 3D7 reference

genome to identify the repeat unit at *plasmepsin* genes.

Plasmepsin II/III copy number identification

We analyzed mapped reads on the 67 kb extended loci of the *plasmepsin* genes (from

260,000 to 327,000 at chromosome 14), which including 28 kb upstream *plasmepsin* I and 28

kb downstream *plasmepsin* III. The number of reads mapped onto each position (coverage)

were determined from the deduplicated BAM file using *bedtools*

(https://bedtools.readthedocs.io/en/latest/). We visualized reads mapping to the 67 kb region

using the Integrative Genomics Viewer (IGV) (Fig S1A). The repeat unit of at *plasmepsin* genes

was confirmed by both IGV plot and nanopore long-read sequences. We calculated the

Plasmepsin copy number using mean[coverage at repeat unit]/mean[coverage at extension

regions].

Linkage Analysis

Linkage between the *pfcrt*, *pm2/3* and *pfkelch13* loci was assessed by calculating D and D′ and utilizing Fisher's exact test to determine significance. Association between *pfcrt* or *pfkelch13* alleles and *pm2/3* CN were assessed by binning multicopy *pm2/3* CN and conducting Fisher's exact test.

Parasites used in this study

This study utilizes parents and progeny of the KH004 × Mal31 genetic cross generated by utilizing human liver-chimeric. From this cross, one hundred and four unique recombinant progeny were recovered through limited dilution cloning at 0.3 cells per well. Individual wells containing parasites were identified through qPCR and analysis of progeny relatedness and allele inheritance were based on the methods outlined in Button-Simons et al. (45). Methods for sequencing of progeny, construction of the physical map, and generation of visual recombination map are all based on Button-Simons et al. (45). Analysis of allelic co-inheritance utilized the entire progeny set of one hundred and four parasites and phenotyping assays utilized fifty unique parasites, forty-eight progeny and the two parental parasites.

Parasite culture

Cryopreserved *P. falciparum* stocks of KH004 × Mal31 progeny and KH004-020-019-H9 and Mal31-9040-C11 parents were thawed and cultured in complete media (CM) consisting of 0.5% Albumax II (Gibco, Life Technologies) supplemented RPMI 1640 with L-glutamine (Gibco, Life Technologies) with additional 50 mg/L hypoxanthine (Calbiochem, Sigma-Aldrich), 25 mM HEPES (Corning, VWR), 10ug/ml gentamycin (Gibco, Life Technologies), and 0.225% sodium bicarbonate (Corning, VWR). Parasite cultures were maintained at 5% hematocrit in O+ red blood cells (RBC) (Interstate Blood Bank, Memphis, TN) in separate flasks and maintained

consistent temperature (37ºC) and atmosphere (5% CO_2/5% O_2/90% N_2). Parasitemia of

cultures was kept below 2% and media changes were performed every 48 hours, corresponding

with one intraerythrocytic development cycle.

PfCRT Structure Analysis

Three-dimensional homology model of PfCRT was predicted using AlphaFold (62, 63)

and accessed through the PDB database (6UKJ) (33). Additionally, the F'(ab) fragment used in

7G8 PfCRT cryo-EM structure elucidation (33) was removed from the structure. Protein

mutagenesis and visualization of resistance-associated mutations were conducted using PyMol

software (v2.5.4; Schrödinger, LLC).

Drug susceptibility assays

Prior to susceptibility assay setup, parasites were synchronized using a single layer of

70% Percoll (Sigma Aldrich) in 1X RPMI with 13.3% sorbitol in phosphate buffered saline.

400µL of packed erythrocytes infected by a majority of schizont stage parasites were

resuspended in 2mL of incomplete media, layered over the Percoll layer, and centrifuged (1561

x g for 10 in, no brake). The top layer was removed to isolate late-stage schizonts, washed

twice with incomplete media, resuspended at 5% hematocrit in complete media, and placed on

a shaker for 4 hours at 37ºC. Following the 4 hour incubation, parasitemia and stage were

determined through flow cytometry by staining with SYBR Green I and SYTO 61 and analyzed

on a Guava easyCyte HT (Luminex) after 50,000 events were counted. If cultures were >70%

ring-stage, they were diluted to 0.15% parasitemia at 2% hematocrit and set up in a 96-well

plate at a volume of 150µL per well. Within each assay plate, two technical replicates of each

parasite were exposed to 10 concentrations of a 2-fold dilution series of PPQ as well as both

untreated parasite and uninfected RBC controls. Parasites were exposed to PPQ for 72 hours.

Parasite density was determined using SYBR Green and dose-response curves were generated

by plotting percent survival against the log of drug concentration. Analysis of IC_{50} and AUC was

performed using Prism 9.0 (GraphPad). A nonlinear log inhibitor vs response 4 parameter dose-

response curve was generated based on percent survival. To generate the limited point curve,

data points which made up the biphasic portion of the plot were excluded to allow for a

sigmoidal shaped curve, which was then used for IC_{50} calculations.

PSA
899

PSA was set up as originally described (18) with synchronized early ring stage parasites

being expose to a single 200nM dose of PPQ for 48 hours. This protocol was modified to

resemble the extended recovery RSA (eRRSA) phenotype (64), in which samples are analyzed

at 120 hours post-exposure instead of 72 hours. This extended timepoint has been shown to

provide superior differentiation between resistant and sensitive parasites and allows for finer

resolution between PPQ-S parasites. Parasite prevalence was measured through qPCR and

survival was calculated as fold change between untreated and treated samples for three

technical replicates (3 replicates within the plate as well as 3 independent qPCR reads) and at

least 3 biological replicates per sample. Additionally, to ensure consistency between our

resistance phenotypes, parasites from the same synchronization were used for both PSA and

IC_{50}/AUC susceptibility assays.

Competitive growth assays

To determine relative fitness levels between parasites, we used competitive growth

assays between progeny of the KH004 **×** Mal31 cross and both parental parasites as a proxy for

in vitro fitness. Parasitemia and stage of parasites was quantified using flow cytometry 15 hours

after synchronization with a 70% Percoll gradient. Synchronized parasites were each adjusted

to 0.5% parasitemia at 5% hematocrit and set up in a 1:1 ratio (1% total parasitemia) in 96-well

plates as previously described in (37, 39). Parasitemia was assessed (Giemsa-stained slides

and microscopy) every two days, in accordance with one intraerythrocytic life cycle, adjusted to

1% parasitemia and supplemented with fresh blood and media. Samples from each well were

collected at regular intervals upon sample dilution and stored at -80ºC for genotyping. These

competitive growth assays were maintained for a maximum of 40 days, or until one parasite

reached fixation (>95% of the co-culture). Parasites were deemed to have equivalent fitness

levels if by 40 days neither parasite reached fixation.

To analyze relative parasite densities in each competition, six microsatellite markers

(MS) were used for genotyping (Table S8) to differentiate parasites in co-culture. PCR

amplification was performed using the Phusion Blood Direct PCR Kit (ThermoFisher, cat

#F547L) for 30 cycles as previously described in (37). Amplified microsatellites were analyzed

using the Applied Biosystems 3730xI DNA Analyzer (ThermoFisher). Raw fragment analysis

data was uploaded to the Thermo Fisher Connect platform and analyzed using the microsatellite

analysis tool. The proportion of the two competing parasites in co-culture was determined by

taking the ratio of fluorescent peak height associated with the corresponding PCR product size.

Due to the large number of competitive growth assays required to directly compete all

parasites against each other, all parasites were competed against both parents. Additionally, a

genetically diverse subset of parasites were competed against each other. Based on the

outcomes of the progeny v progeny competitions, an Elo rating system, which has previously

been used for animal dominance hierarchies, was utilized to assign a quantitative score to each

parasite based on their win/loss record while also taking into account the record of the parasite

competed against (65). The package "EloRating" (51) in R Studio (2023.03.0+386). was used

for assigning scores. In order to reduce the effect of competition order on ranking, the

competitive outcomes file was randomized 1000 times to generate 1000 Elo scores which were

then averaged. The distribution of Elo ratings figure was created using Prism 9.0 (GraphPad).

QTL Analysis

Statistical analysis of the QTL data for this cross was performed using the computational

methods previously described (66). Blood stages of *P. falciparum* are haploid, therefore, only

two genetic classes are present for each locus. QTL mapping was performed using the "R/qtl"

package (67) in R Studio. One thousand permutations of the trait values determined the 37%,

5%, and 1% genome-wide significance thresholds, and the strength of each linkage was

expressed as a logarithm of the odds (LOD) score (68). The main QTL and corresponding mean

trait values from these scans were used to obtain estimates of residual empirical thresholds to

identify secondary loci that contribute to the phenotype of interest (68). After identifying QTLs

associated with each phenotype, QTL interaction was tested using the scantwo function of the

"R/qtl" package and was based on 1,000 permutations.

Global haplotype analysis

959   Analysis of global *pfcrt* haplotypes was performed using publicly available sequences

960 through the Pf7 dataset (MalariaGEN) (12). PPQ resistance status for this dataset was inferred

961 based on *plasmepsin II/III* amplification, therefore, all samples labeled as PPQ-R are assumed

962 to express more than one copy of these genes.

Data Availability

965   All data needed to evaluate the conclusions in the paper are present in the paper and/or the

966 Supplementary Materials. All raw sequencing data have been submitted to the NABI Sequence

967 Read Archive (SRA, https://www.ncbi.nlm.nih.gov/sra) under the project number of

968 PRJNA524855. Additional data related to this paper may be requested from the authors.

Code Availability

The code used in analysis and data analysed are available at GitHub through the following

links: https://github.com/kbuttons/MalariaGeneticCrosses-MAL31xKH004

Acknowledgments

This work was supported by National Institutes of Health (NIH) program project grant P01

AI127338 (to MF) and by NIH grant R37 AI048071 (to TJCA). Nanopore sequencing was

supported by Texas Biomedical Research Institute Forum Grant 20-04866 (to XL). Work at

Texas Biomedical Research Institute was conducted in facilities constructed with support from

Research Facilities Improvement Program grant C06 RR013556 from the National Center for

Research Resources. The parental line, Mal31, used in the *Mal31×KH004* cross was sampled

from a Malawian patient in 2016 as part of a cross-sectional study funded by the Wellcome

Trust of Great Britain (Grant no. 099992/Z/12/Z to SCN). We thank the patients who provided

parasites used in this work.

984
985
986
987   **Additional Information**

988   Additional figures and tables are available for this paper.

990