Sex-specific mechanisms of cerebral microvascular BK$_{Ca}$ dysfunction in a mouse model of Alzheimer's disease.

Josiane F. Silva[1], Felipe D. Polk[1,*], Paige E. Martin[1,*], Stephenie H. Thai[1], Andrea Savu[1], Matthew Gonzales[1], Allison M. Kath[1], Michael T. Gee[1], Paulo W. Pires[1,2,3,#]

[1]Department of Physiology, [2]Sarver Heart Center, [3]Bio5 Institute, University of Arizona College of Medicine, Tucson, AZ, 85721, USA.

[*]Equal contribution

[#]Corresponding Author

Running title: Sex-dependent neurovascular dysfunction in *5x-FAD*

Address for Correspondence: Paulo W. Pires, Ph.D.

1656 E Mabel St, Room 426

Medical Research Building

Tucson, AZ 85721

PO Box 245051

Phone: +1 (520) 626-8632

Email: ppires@arizona.edu

Keywords: cerebral pial arteries, BK$_{Ca}$ channels, nitrosylation, cerebral functional hyperemia, Ca^{+2} sparks, sex as a biological variable

Abstract

Background: Cerebral microvascular dysfunction and nitro-oxidative stress are present in patients with Alzheimer's disease (AD) and may contribute to disease progression and severity. Large conductance Ca^{2+}-activated K^+ channels (BK_{Ca}) play an essential role in vasodilatory responses and maintenance of myogenic tone in resistance arteries. BK_{Ca} impairment can lead to microvascular dysfunction and hemodynamic deficits in the brain. We hypothesized that reduced BK_{Ca} function in cerebral arteries mediates microvascular and neurovascular responses in the *5x-FAD* model of AD. *Methods:* BK_{Ca} activity in the cerebral microcirculation was assessed by patch clamp electrophysiology and pressure myography, in situ Ca^{2+} sparks by spinning disk confocal microscopy, hemodynamics by laser speckle contrast imaging. Molecular and biochemical analyses were conducted by affinity-purification assays, qPCR, Western blots and immunofluorescence. *Results:* We observed that pial arteries from 5-6 months-old male and female *5x-FAD* mice exhibited a hyper-contractile phenotype than wild-type (WT) littermates, which was linked to lower vascular BK_{Ca} activity and reduced open probability. In males, BK_{Ca} dysfunction is likely a consequence of an observed lower expression of the pore-forming subunit BK_α and blunted frequency of Ca^{2+} sparks, which are required for BK_{Ca} activity. However, in females, impaired BK_{Ca} function is, in part, a consequence of reversible nitro-oxidative changes in the BK_α subunit, which reduces its open probability and regulation of vascular tone. We further show that BK_{Ca} function is involved in neurovascular coupling in mice, and its dysfunction is linked to neurovascular dysfunction in the model. *Conclusion:* These data highlight the central role played by BK_{Ca} in cerebral microvascular and neurovascular regulation, as well as sex-dependent mechanisms underlying its dysfunction in a mouse model of AD.

Introduction

Alzheimer's disease (AD) is a neurodegenerative disorder associated with global cognitive decline that affects more than 6 million people in the United States alone[1,2], and approximately 2/3 of all AD cases are women[1,3]. Alarmingly, it is projected that more than 130 million people will live with AD worldwide by 2050[1,3]. One of the hallmarks of AD is cerebral amyloid angiopathy (CAA), characterized by perivascular accumulation of amyloid-β, which is present in more than 80% of all AD patients[4]. This accumulation is proposed to induce microvascular fragility, contributing to the development of intracerebral hemorrhages[5], loss of blood-brain-barrier integrity [6], and blood flow deficits[7]. These deficits may be a consequence of microvascular dysfunction, which impairs cerebrovascular reserve and neurovascular coupling[8], a vital process that matches local hyperemia to increased neuronal activity[9]. The ability of the cerebral microcirculation to accommodate these second-to-second changes in local perfusion depends on two main phenomena: the extent of myogenic tone (the intrinsic contractile property of vascular smooth muscle cells that stipulates the vasodilatory reserve) and mechanisms underlying vasorelaxation. Both mechanisms are dependent on the membrane potential of smooth muscle cells: depolarization favors constriction, whereas hyperpolarization favors dilation. Thus, regulation of ion flux is paramount for microvascular function, particularly modulation of K^+ permeability, of which the large-conductance Ca^{2+}-activated K^+ channel (BK_{Ca}) is central.

BK_{Ca} are widely expressed channels that play a critical role in regulating arterial tone by conducting hyperpolarizing currents in vascular smooth cells to oppose vasoconstriction[10,11]. Opening of BK_{Ca} channels occurs downstream from localized intracellular Ca^{2+} release from ryanodine receptors located in the sarcoplasmic reticulum (Ca^{2+} sparks), which activates BK_{Ca} channels, leading to hyperpolarization and dilation[10,12]. BK_{Ca} channels are composed of two subunits, a selective pore (BK_{α}) and regulatory ($BK_{\beta 1,}$ BK_{γ}) subunits[13]. Further, BK_{Ca} undergoes extensive pre- and post-translational modifications (PTM), including phosphorylation[14-16], oxidation and nitrosylation, all affecting channel open probability (Po) and conductance[16]. In some tissues, oxidation increases channel activity[17,18], whereas, in others, it markedly inhibits BK_{Ca}[19-21]. Similarly, nitrosylation of thiol moieties due to an electrophilic attack by peroxynitrite, formed by a reaction between nitric oxide (NO) and superoxide (O_2^-) in a pro-oxidative environment[22-24], as seen in AD[25-27], inhibits BK_{Ca} currents[19]. Despite recent evidence of BK_{Ca} impairment in male APP23 mice[28], it remains unknown if the reduction in cerebral microvascular BK_{Ca} occurs in females *via* similar mechanisms, and whether neurovascular dysfunction accompanies loss of BK_{Ca} activity. Using a combination of molecular and biochemical tools, ex vivo patch clamp

electrophysiology and microvascular pressure myography, and *in vivo* hemodynamic
assessments, we show that BK_{Ca} dysfunction mediates cerebral microvascular and neurovascular
dysfunction in in both males and females *5x-FAD*. Remarkably, the mechanisms underlying this
dysfunction show sex-specificity: In males, we observed lower BK_{α} expression and reduced Ca^{2+}
sparks frequency; however, in females, there is an exacerbated and partially reversible nitro-
oxidative modification of BK_{α}.

Methods

All animal procedures in this study were approved by the Institutional Animal Care and
Use Committee of the University of Arizona College of Medicine (IACUC protocol 18-473) and are
in accordance with the National Institutes of Health's Guide for the Care and Use of Laboratory
Animals, 8th edition. All animal experiments are reported in compliance with ARRIVE guidelines.
The *5x-FAD* mice (5-6 months-old) used in this study are a transgenic model of early-onset AD
that combines five different mutations in two genes associated with Familial AD. Three of the
mutations are located in the gene that codes human amyloid precursor protein: Swedish (K670N,
M671L), Florida (I716V), and London (V717I). The other two mutations are inserted in the gene
of human presenilin, M146L and L286V. Transgene expression is controlled by the neuronal-
specific *Thy1* promoter[29]. All mice were bred on a C57bl/6J background, and wild-type (WT)
littermates were used as controls. Following euthanasia, the brain was collected and the pial
arteries were dissected for patch clamp electrophysiology, pressure myography and quantification
of vascular smooth muscle cells Ca^{2+} sparks activity by spinning disk confocal microscopy. The
brain was either fixed in 4% paraformaldehyde for immunofluorescence, or flash-frozen for
molecular biology and biochemical experiments (oxidized glutathione assay, affinity-purification
of nitrosylated proteins followed by Western blot and qPCR). Baseline cortical perfusion and
functional hyperemia following somatosensory stimulation were measured by laser speckle
contrasting imaging.

Post-mortem brain tissues from CAA / AD and age-matched controls (no CAA or AD
diagnosed) were used for assessment of BK_{α} S-NO by affinity-purification followed by Western
blot. All human samples were kindly donated by the Brain and Body Donation Program from
Banner Health, and are in compliance with all guidelines and requirements for patient de-
identification. The complete and detailed methodology can be found in the *Supplemental*
Information - Methods.

Results

Amyloid-β accumulation is higher in female 5x-FAD mice

One of the hallmarks of Alzheimer's disease is the accumulation of amyloid-β peptides in the brain and cerebral microvasculature[30]. In this study, *5x-FAD* female mice showed a trend towards higher density of amyloid-β plaques in the cortex and hippocampus when compared with age-matched male *5x-FAD* (Figures S1A and S1B). No significant perivascular amyloid-β accumulation was observed in the parenchymal microcirculation of male or female *5x-FAD* mice (Figures S1C and S1D). We previously showed that pial arteries in this model exhibit extensive amyloid-β accumulation[31]. However, in our previous publication, we did not separate the incidence of pial vessel CAA by sex. Thus, we reanalyzed those data and observed no sex differences in pial vessel CAA in *5x-FAD* mice (Figures S1E, and S1F). Given the lack of microvascular intraparenchymal CAA in *5x-FAD*, we focused our vascular reactivity studies on the pial microcirculation.

Increased contractility of pial arteries in 5x-FAD mice.

Along with amyloid-β plaque accumulation, microvascular dysfunction is implicated in AD pathophysiology[30]. We observed an exacerbated contractile response in pial arteries of both female and male *5x-FAD* when compared to wild-type (WT) littermates, shown as higher spontaneous myogenic tone at a purported physiological intraluminal pressure compared to the respectively WT (50 mmHg, Figures 1A-B and 1E-F). Similarly, contractile response to a receptor-dependent stimulus (endothelin-1 [ET-1], 30 nM, Figures 1C and 1G) was higher in females and males *5x-FAD* mice than WT. No differences were observed in contractile responses to a receptor-independent depolarizing stimulus (60 mM $[K^+]_E$, Figures 1D and 1H). Thus, we conclude that contractile responses to a steady intraluminal pressure, as well as receptor-induced, are exacerbated in pial arteries of *5x-FAD* mice.

We then characterized autoregulatory responses (myogenic reactivity) of pial arteries in *5x-FAD* and WT littermates by step-wise increases in intraluminal pressure (from 5 mmHg to 160 mmHg). Under our experimental conditions, myogenic reactivity was not altered in *5x-FAD*, both females and males (Figure S2). Similarly, pial arteries from female or male *5x-FAD* do not undergo significant structural remodeling (Figure S3) or alterations in biomechanical properties (Figure

152 S4)**.** Thus, the enhanced contractility of pial arteries observed is independent of microvascular
153 remodeling.

Reduced BK_{Ca} channel function in pial arteries from 5x-FAD

BK_{Ca} channel is an important mechanism that provides negative feedback to myogenic
constriction[32]. Therefore, to test whether the increased contractility observed in *5x-FAD* was due
to BK_{Ca} dysfunction, we incubated pial arteries with the BK_{Ca} channel blocker iberiotoxin (IbTox,
30 nM). IbTox-induced vasoconstriction was significantly smaller in pial arteries isolated from *5x-*
FAD when compared to WT littermates, for both females and males (Figures 2A-B and 2E-F),
suggesting impaired BK_{Ca} activity in *5x-FAD* mice.

We next corroborated the pressure myography findings by recording pial artery smooth
muscle cell single channel BK_{Ca} currents in excised, inside-out patch-clamp electrophysiology
experiments. We observed an approximately 50% lower BK_{Ca} open probability (Po) at +60 mV in
patches from *5x-FAD* female than those from WT littermates (Figures 2C and 2D). However, in
males, BK_{Ca} Po was not significantly different between WT and *5x-FAD* (Figures 2G and 2H).

Interestingly, when comparing IbTox constriction between WT females and males, we
observed a significantly smaller constriction in pial arteries of males than in females
(Vasoconstriction (%): 10.6 ± 1.2 vs 6.6 ± 1.3, WT females vs WT males, p = 0.0494, two-tailed
Student's t-test). This finding was again corroborated by patch-clamp electrophysiology: BK_{Ca} Po
was significantly lower in WT males when compared to females (BK_{Ca} Po at +60 mV: 0.118 ±
0.025 vs 0.040 ± 0.013, WT females vs WT males, p = 0.0125, two-tailed *Student*'s t-test). These
data suggest that physiological BK_{Ca} activity is lower in pial artery smooth muscle cells from males
than in female WT mice.

5x-FAD males, but not females, have lower BK_α expression in pial arteries

To investigate if the reduced constriction to IbTox in *5x-FAD* mice was due to lower
expression of BK_{Ca}, we assessed mRNA expression of BK_α and $BK_{\beta 1}$ subunits in lysates of pial
arteries from female and male *5x-FAD* and WT littermates by qPCR. Pial artery mRNA expression
of both subunits was not significantly different between *5x-FAD* and WT in females (Figures 3A
and S5A) or males (Figures 3C and S5B). Using Western blot, we then measured expression of
the BK_α protein in pial artery lysates, and observed that, in females, expression of BK_α was similar

between WT and *5x-FAD* (Figure 3B). However, expression of BK_α protein was significantly lower

in male *5x-FAD* when compared to WT littermates (Figure 3D). An image of the membrane used

for the inserts can be found in Figure S6.

In brain lysates, we observed a downregulation of BK_α mRNA in female *5x-FAD* when

compared to WT littermates (Figure S5C), without changes in mRNA for $BK_{\beta 1}$ (Figure S5D). No

significant differences were observed in mRNA for BK_α and $BK_{\beta 1}$ in brain lysates from male *5x-*

FAD when compared to WT (Figures S5E and S5F).

Lower frequency of Ca²⁺ sparks in pial arteries from male 5x-FAD mice, but not in females

Lower frequency of Ca^{2+} sparks in pial arteries from male 5x-FAD mice, but not in females

We next investigate whether the observed BK_{Ca} dysfunction was downstream from

alterations in intracellular Ca^{2+} sparks, as shown previously in male APP23 mice[28]. Our data show

that Ca^{2+} sparks frequency is significantly higher in pial artery smooth muscle cells of female *5x-*

FAD mice than WT littermates (Figure 3I, Movies S1 and S2), without changes in the number of

Ca^{2+} sparks sites per cell (Figures 3E, 3F and 3J). In contrast, Ca^{2+} spark frequency was

significantly lower in pial artery smooth muscle cells of male *5x-FAD* than in WT (Figure 3K,

Movies S3 and S4). This reduction was not due to a lower number of active Ca^{2+} sparks site per

cell, as these were not different between males *5x-FAD* and WT (Figures 3G, 3H and 3L).

Altogether, our findings show that BK_{Ca} dysfunction in *5x-FAD* male is due to lower expression of

BK_α and reduced Ca^{2+} sparks frequency, whereas in females we observed an increase in Ca^{2+}

sparks frequency without changes in BK_α expression.

Oxidative stress impairs BK_Ca in female 5x-FAD

Oxidative stress impairs BK$_{Ca}$ in female 5x-FAD

AD is associated with oxidative stress and inflammation that can contribute to vascular

and neuronal damage[26,33]. Thus, we investigated if oxidative stress was present in the brain of

5x-FAD using an oxidized glutathione assay. We observed that, in females, total glutathione (Total

GSH) levels were similar between WT and *5x-FAD* (Figure 4A). However, brain levels of oxidized

glutathione (GSSG, Figure 4B) and the ratio GSSG / Total GSH (Figure 4C) were significantly

higher in female *5x-FAD* than WT littermates, suggesting the presence of an oxidative

environment. None of the assessed parameters were altered in male *5x-FAD* when compared to

WT littermates (Figure S7A-S7C).

213   We then hypothesized that this oxidant environment leads to reversible PTM in pial artery

214 BK_{Ca} in female *5x-FAD*, thus causing its impairment. We tested this hypothesis by incubating *ex*

215 *vivo* pial arteries and excised inside-out membrane patches with the broad-spectrum reducing

216 agent dithiothreitol (DTT, 10 µM). We observed that DTT partially reversed pial artery sensitivity

217 to IbTox in females, observed as a larger vasoconstriction response after incubation with DTT

218 (Figures 4D and 4E). These findings were corroborated in patch-clamp electrophysiology

219 experiments, as incubation of patches with DTT partially restored BK_{Ca} Po (Figures 4F and 4G).

220 DTT did not affect BK_{Ca} Po in pial artery smooth muscle cells isolated from male *5x-FAD* (Figure

221 S7D-S7E). Altogether, these data suggest that the oxidative environment in female, but not males,

222 *5x-FAD* leads to oxidation of BK_{Ca}, consequently reducing its function in pial artery smooth muscle

223 cells.

Exacerbated BK_{Ca} S-nitrosylation in female 5x-FAD and CAA / AD patients

Exacerbated BK_{Ca} S-nitrosylation in female 5x-FAD and CAA / AD patients

226   Next, we sought to identify the likely type of PTM responsible for the observed BK_{Ca}

227 impairment. The inducible nitric oxide synthase (iNOS) isoform is a pro-inflammatory enzyme that

228 is increased in the AD brain[34,35]. This isoform produces a high amount of NO, which can lead to

229 nitro-oxidative damage of cellular structures[36,37], including excessive nitrosylation of target

230 proteins. Thus, we investigated if iNOS expression is increased in the pial arteries of *5x-FAD*

231 females. Using qPCR, we observed that the expression of iNOS mRNA was increased in the pial

232 arteries from female *5x-FAD* mice compared to WT (Figure 5A). No significant differences were

233 observed for endothelial (eNOS, Figure S8A) or neuronal NOS (nNOS, Figure S8B) in pial

234 arteries. To corroborate the mRNA findings, we performed a semi-quantitative assessment of

235 fluorescence intensity of iNOS immunolabeling in pial arteries, which was significantly higher in

236 pial vessels from female *5x-FAD* than in WT mice (Figures 5B and 5C). These results suggest

237 that iNOS is upregulated on the pial vessels from female, but not male, *5x-FAD* mice. Lastly,

238 inhibition of NOS with L-NAME (200 µM) induced a greater pial artery constriction in *5x-FAD* than

239 in WT littermates (Figure S8C).

240   Excessive NO production in the presence of an oxidative environment can lead to

241 excessive protein S-NO[38], thus we investigated if this particular PTM was involved in the BK_{Ca}

242 dysfunction observed in female *5x-FAD*. Our findings show that total S-NO proteins were

243 significantly higher in brain lysates of female *5x-FAD* when compared to WT littermates (Figures

244 5D-5E), but not in male *5x-FAD*, where they were actually significantly reduced (Figure S9). We

then performed an affinity-purification of S-NO proteins from brain lysates of female *5x-FAD* and
WT littermates, followed by Western blot against the BK_α subunit. We observed a significant
increase in the amount of BK_α S-NO (Figures 5F-G, whole blot of inserts in Figure S10). We
corroborated these findings by performing a co-immuno-localization assay to investigate
vascular-specific BK_α S- NO (Figure 5H). Our data shows that there is a significant increase in
co-localization between the BK_α subunit and S-NO proteins, as evidenced by the M1 / M2 –
Mander's coefficients (Figures 5I-5J) and the Pearson's coefficient (Figure 5K). Together, these
findings support our hypothesis that the pro-nitro-oxidative environment in *5x-FAD* females leads
to excessive BK_α S-NO.

We then investigated if the increase in BK_α S-NO observed in *5x-FAD* was also present in
humans with CAA or AD. We obtained post-mortem brain samples from the Banner Brain and
Body Donation Program, from age-matched women and men, and separated them in 3 groups:
healthy (non-CAA / non-ADA, n=6), CAA / non-AD (CAA, n=6) and AD + CAA (n=12). Healthy
patients were defined no presence of dementia or symptoms of parkinsonism, and no presence
of CAA or senile plaques in histopathology. CAA / non-AD were patients without dementia or
parkinsonism, but with evident perivascular amyloid plaques and no presence of senile plaques.
AD + CAA were patients with Alzheimer's disease with a minimum of intermediate or high NIA-
Reagan criteria[39], and presence of CAA and / or senile plaques. Using Western blot, we observed
that there was a trend towards an increase in BK_α subunit expression in brain from AD + CAA
patients (p=0.0976, Brown-Forsythe and Welch's ANOVA), without evident sex differences
(Figures 6A-6B). When performing an affinity purification of S-NO proteins followed by Western
blot, we again observed a trend towards an increase in BK_α S-NO in AD + CAA patients
(p=0.0772, Brown-Forsythe and Welch's ANOVA), without sex differences (Figures 6C-6D).
Images of the whole blots shown in the cropped insets (Figures 6A and 6C) can be found in the
Figure S11. These findings suggest that BK_α S-NO is also present in AD patients, although in
both women and men.

Cerebral perfusion deficits in female 5x-FAD

Next, we investigated if there are changes in basal cortical perfusion in *5x-FAD* females
and males compared to sex-matched WT littermates. Using laser speckle contrast imaging, we
observed that the baseline perfusion of the entire superficial cortex was not different between WT
and *5x-FAD* females (Figures S12B-S12C, WC: whole cortex**)** or males (Figures S12D-S12E**,**

WC: whole cortex). However, when analyzing discrete perfusion of specific vascular territories

(diagram in Figures S12A)[40], we observed that *5x-FAD* females showed lower basal perfusion to

the frontal cortex (Figures S12B-S12C), but not in other regions. No significant differences were

observed in cortical perfusion of male *5x-FAD* mice (Figures S12D-S12E).

Impaired functional neurovascular responses in females and males 5x-FAD

We next tested the hypothesis that impaired vascular BK_{Ca} would result in neurovascular

dysfunction in *5x-FAD*. First, we showed that acutely inhibiting BK_{Ca}, achieved by applying IbTox

(30 nM) atop the thinned-skull cranial window, significantly blunted hyperemic responses on the

somatosensory cortex during mechanical stimulation of contralateral whiskers[41] (Figures 7A-7C),

suggesting that BK_{Ca} play a role in neurovascular responses. We then assessed neurovascular

responses in *5x-FAD* mice and compared to WT littermates. We observed that both females

(Figures 7D-7F) and males (Figures 7G-7I) *5x-FAD* had significantly lower hyperemic responses

than WT littermates. Together, these findings suggest that the cerebral vascular impairment

observed in *5x-FAD* mediates neurovascular dysfunction in the model.

Discussion

Our findings show that cerebral BK_{Ca} microvascular dysfunction is present in both female

and male *5x-FAD*, likely mediating neurovascular impairments in the model. A major novelty of

our study lies in our findings elucidating the distinct mechanisms underlying BK_{Ca} dysfunction in

females and males. In males, we observed reduced expression of the BK_{α} subunit and lower Ca^{2+}

sparks frequency. On the other hand, in females, we show that BK_{α} expression is unchanged and

Ca^{2+} sparks frequency is higher, yet BK_{Ca} activity is likewise impaired. Upon further investigation,

we uncovered that the BK_{α} subunit in female *5x-FAD* undergoes oxidative modification,

particularly excessive S-NO, and this partially underlies cerebral microvascular BK_{Ca} dysfunction.

The exacerbation in nitrosylation is downstream from increased vascular iNOS expression and

general oxidative stress in the brain environment, none of which were present in males (Figure

S13). Lastly, we describe two other novel findings in our study: basal cerebral microvascular BK_{Ca}

activity seems lower in male than in female C57Bl/6J mice, and that BK_{Ca} function is involved in

physiological neurovascular responses, as its blockade blunts somatosensory-induced functional

hyperemia. Altogether, our data highlight the complexities of microvascular regulation in the

context of AD, and the differential effects of biological sex that may hinder putative therapies

ineffective as a "one size fits all" approach.

AD is a progressive neurodegenerative disease characterized by accumulation of amyloid-

β and neurofibrillary tangles, particularly in the temporal lobe and neocortical structures[42], which

lead to cognitive deficits[30]. Beyond misfolded protein pathology, cerebral vascular dysfunction

contributes to AD progression, and much attention has been placed on loss of endothelial function

and blood-brain-barrier breakdown[43,44]. However, less is known on how cerebral microvascular

smooth muscle cell function is affected in AD, as well as its participation in AD's development and

progression. A recent report by Taylor *et al.* described that acute exposure of cerebral arteries

isolated from male C57Bl/6 mice to amyloid-$\beta_{(1-40)}$ increases myogenic tone due to impaired BK_{Ca}

activity[28]. Our findings expand on this report, and we show that not only spontaneous myogenic

tone is higher in *5x-FAD*, a mouse model of AD with accumulation of both amyloid-$\beta_{(1-40)}$ and

amyloid-$\beta_{(1-42)}$ (predominant), but so is contractility to ET-1. These data suggest the presence of

a general hyper-contractile state in the cerebral microcirculation in both female and male arteries

during AD progression. Our data suggests that BK_{Ca} impairment likely underlies this hyper-

contractile status in *5x-FAD*, as its function is impaired in *ex vivo* pressurized arteries.

Paradoxically, we did not observe any significant changes in autoregulation. Although loss of

autoregulation is observed in cerebral small vessel disease[45] and it was shown to occur in some

rodent models of AD[46], human studies have also failed to observe cerebral autoregulation

impairments in AD patients[47]. The mechanistic explanation underlying these differences in

autoregulation between AD models and humans remain unclear and require further exploration.

Impairment in BK_{Ca} function is associated with vascular dysfunction in various disease

models, including diabetes[48,49], hypertension[50,51], menopause[41,52], cigarette smoking[53] and AD[28].

One limitation of the aforementioned studies is that they were all performed in a single sex, and

the vast majority in males (with the exception of studies focused on menopause). Our findings

expand on these previous observations and show a sex-dependent mechanism of BK_{Ca}

impairment in *5x-FAD*. We show that BK_{Ca} activity is reduced in pial arteries independently of

biological sex, observed as blunted constriction to iberiotoxin. However, the cellular and molecular

mechanisms responsible for the lower $BK_{C}a$ activity diverge between females and males. We

observed that, in male *5x-FAD*, single channel BK_{Ca} Po was not lower than that recorded from

WT littermates, indicating that BK_{Ca} dysfunction was likely upstream from intracellular signals

regulating BK_{Ca} activity. Upon further investigation, we describe that expression of the pore-

forming subunit BK_{α} is lower in male *5x-FAD* than in WT littermates, which could account for lower

sensitivity to Iberiotoxin in pressure myography experiments. In addition, frequency of Ca^{2+} sparks

was significantly reduced in male *5x-FAD*, similar to data published by Taylor *et al.* in male APP23

mice[28].   On the other hand, female *5x-FAD* showed reduced single channel BK_{Ca} Po despite

similar levels of BK_α expression and an increase in Ca^{2+} spark frequency, suggesting that, in

females, BK_{Ca} dysfunction was likely due to molecular alterations in the BK_α subunit itself, such

as PTMs resulting from nitro-oxidative stress.

In support of this hypothesis, we observed that brains of females *5x-FAD*, but not males,

show an increase in markers of oxidative and nitrosative stress, observed as higher levels of

oxidized glutathione (GSSG) and higher levels of global S-NO proteins. Together, this nitro-

oxidative environment can result in exacerbated PTMs within the BK_α subunit. PTMs are the result

of binding of different moieties to proteins, such as phosphorylation, acetylation, lipidation,

oxidation and nitrosylation[24,54]. PTMs play a vital role in the known heterogeneity and tissue

specificity of the biophysical properties of BK_{Ca}, as they have profound effects on gating, voltage

and Ca^{2+} sensitivity, and membrane availability of the channel[16]. However, how S-NO affects BK_{Ca}

remains controversial: some studies show that NO can, directly or indirectly, activate BK_{Ca}[55,56],

whereas others show that NO itself, or peroxynitrite, do not affect or can blunt BK_{Ca} activity. For

instance, a NO-donor failed to activate the BK_{Ca} channel in a heterologous expression system[57].

Moreover, Liu *et al.* showed that peroxynitrite reduces the open-state probability of BK_{Ca} channels

in isolated vascular smooth muscle cells[58]. We show that, in female *5x-FAD*, the BK_α subunit

undergoes excessive S-NO, and this nitro-oxidative PTM likely mediates channel dysfunction in

a reversible manner. This assertion is supported by our data that the broad-spectrum reducing

agent DTT partially recover single BK_{Ca} Po and arterial responsiveness to iberiotoxin. Although

the molecular mechanisms of how S-NO impair BK_{Ca} remain undetermined, we speculate that

they may reduce Ca^{2+} sensitivity of the BK_α subunit, which may explain the increase in Ca^{2+} sparks

frequency observed in female *5x-FAD*, likely to compensate for such lower sensitivity. This,

however, remains untested.

We further made efforts to elucidate the cellular source responsible for excessive protein

S-NO, and observed an increase in cerebral vascular iNOS expression. Inducible NOS (NOS2)

is commonly upregulated under stress and inflammatory conditions[34], such as is observed in AD

and other dementias[33,59]. Distinctly from the other NOS isoforms (eNOS and nNOS), iNOS is

effectively locked in an active position due to the tight binding of calmodulin[34,60], leading to

sustained basal NO production. Together with oxidative stress, the higher NO production can

result in excessive protein S-NO, as we observed in female *5x-FAD*. Further, we observed a trend

towards an increase in BK_α S-NO in *post-mortem* brain samples from patients with AD + CAA

when compared to age-matched non-dementia controls (healthy), without any evident changes in

BK_α expression. Interestingly, in humans, our data does not indicate the presence of sex

differences, as both women and men show BK_α S-NO. The discrepancies between the human

and *5x-FAD* findings are not immediately apparent, but we speculate that other co-morbidities in

males may increase BK_α S-NO in men, including cigarette smoking or cardiovascular disease.

We currently do not have the medical records of the patients included in our samples; therefore,

this possibility remains a speculation.

Ultimately, our observed microvascular BK_{Ca} dysfunction may underlie the functional

perfusion deficits observed in *5x-FAD*. We show that acute application of iberiotoxin atop the

cranial window blunts neurovascular responses in both female and male C57Bl/6J mice, showing

the role of BK_{Ca} in these functional hemodynamic responses, as shown by others using an *ex vivo*

brain slice preparations[61]. Further, both females and males *5x-FAD* mice showed a reduction in

neurovascular responses during somatosensory stimulation. A similar neurovascular impairment

was recently reported by us, in which female mice that underwent chemically-induced menopause

showed reduced functional hyperemia responses associated with cerebral microvascular BK_{Ca}

impairment[41]. However, it is important to note that these studies, and our data presented herein,

are associative, and further experiments are needed to definitively show the role of cerebral

microvascular BK_{Ca} in neurovascular coupling responses. Regardless, our previous and current

findings, in conjunction with findings from others[28,51,62], reiterate the central role played by BK_{Ca} in

basal physiological hemodynamic regulation in the brain, as well as its importance in

cerebrovascular diseases.

Lastly, the molecular basis of the sex differences observed remains unclear. Sex

differences are commonly reported in mouse models of AD[63,64], as well as in humans, and sex

hormones seem to play a role. Female *5x-FAD* and AD patients have lower expression of

aromatase in the hippocampus, an important enzyme that produces 17β-estradiol[65]. Since

estradiol can play a role in vascular function[66,67], neuroplasticity[68], synaptic stability[68], as well as

attenuation of oxidative stress[69], it is possible that reduction in its concentration or bioavailability

may result underlie the heightened AD pathology in females. Further, in *5x-FAD* mice harboring

human ApoE3 or ApoE4, ovariectomy worsens cognitive function, and β-estradiol partially

recovers cognition in *5x-FAD* x ApoE3, with little effect in *5x-FAD* x ApoE4. Testes levels of

testosterone are higher in male *5x-FAD* than in wild-type, without changes in sperm viability,

testosterone levels are higher[70]. Currently, there are no published studies showing the effects of
castration on *5x-FAD* males.

In conclusion, our results show that the BK_{Ca}-dependent microvascular and neurovascular
dysfunction occurs in both male and female *5x-FAD*, although the mechanisms underlying BK_{Ca}
impairment differ based on biological sex. Further, we show that one of such mechanisms, namely
excessive $BK_α$ S-NO, may also be present in humans with AD and CAA, suggesting that
manipulation of this basic pathophysiological mechanism could have therapeutic potential.

Novelty and Significance

Cerebrovascular dysfunction is a hallmark of Alzheimer's disease and related dementias.
A property of the resistance vasculature is to constrict when pressurized (myogenic tone);
however, over-constriction can be detrimental. Feedback mechanisms, including the opening of
BK_{Ca}, exist to keep the system in balance. Here, using a combination of molecular biology with *ex*
vivo and *in vivo* assessments, we show a novel mechanism associated with BK_{Ca} dysfunction in
the cerebral microvasculature of female *5x-FAD* mice. We report increased BK_{Ca} S-nitrosylation
linked to higher myogenic tone and reduced channel activity. These changes were associated
with lower perfusion of the frontal cortex and impaired neurovascular reactivity, suggesting that
nitro-oxidative stress is an important mechanism of vascular dysfunction in Alzheimer's disease
in mice and human.

Acknowledgments

We thank Patricia Jansma, co-manager of the Office Research, Innovation & Impact's
Imaging Cores - Optical Facility at the University of Arizona, for providing training service at the
microscope and helping with the microscopy image acquisition. We would further like to thank Dr.
Anne M. Dorrance (Michigan State University) for her critical review of our manuscript.

We are grateful to the Banner Sun Health Research Institute Brain and Body Donation
Program of Sun City, Arizona for the provision of human biological materials. The Brain and Body
Donation Program has been supported by the National Institute of Neurological Disorders and
Stroke (U24 NS072026 National Brain and Tissue Resource for Parkinson's Disease and Related
Disorders), the National Institute on Aging (P30 AG19610 and P30AG072980, Arizona
Alzheimer's Disease Center) the Arizona Department of Health Services (contract 211002,

Arizona Alzheimer's Research Center), the Arizona Biomedical Research Commission (contracts 4001, 0011, 05-901, and 1001 to the Arizona Parkinson-s Disease Consortium) and the Michael J. Fox Foundation for Parkinson's Research.

The mouse strain used for this research project, B6SJL-Tg(APPSwFlLon, PSEN1*M146L *L286V)6799Vas/Mmjax, RRID:MMRRC_034840-JAX, was obtained from the Mutant Mouse Resource and Research Center (MMRRC) at The Jackson Laboratory, an NIH-funded strain repository, and was donated to the MMRRC by Robert Vassar, Ph.D., Northwestern University.

Funding Support

This work was supported by the National Institutes of Health (R00 HL140106, R01 AG073230 to PWP), the Alzheimer's Association (AARGD-21-850835 to PWP), and the Arizona Alzheimer's Consortium, Arizona Department of Health Services.

Conflict of interest

The authors declare no conflicts of interest.

Supplemental Material

A detailed description of the materials and methods is provided in the *Supplemental Material.*

Author Contribution

JFS – conceived, designed, performed the experiments, analyzed data, and wrote the initial draft of the manuscript;

FDP – performed experiments and analyzed data;

PM – performed experiments and analyzed data;

SHT – performed experiments and analyzed data;

ECL – performed experiments and analyzed data;

AS – performed the experiments and analyzed data;

AMK – performed experiments and analyzed data;

MTG – performed the experiments and analyzed data;

PWP – conceived, designed, performed experiments, analyzed data and revised the draft of the
manuscript.

All authors declare that they reviewed and edited the manuscript, and agree to its submission.

References

1.   2021 Alzheimer's disease facts and figures. *Alzheimers Dement*. 2021;17:327-406. doi:
10.1002/alz.12328

2.   Girard SD, Baranger K, Gauthier C, Jacquet M, Bernard A, Escoffier G, Marchetti E,
Khrestchatisky M, Rivera S, Roman FS. Evidence for early cognitive impairment related
to frontal cortex in the 5XFAD mouse model of Alzheimer's disease. *J Alzheimers Dis*.
2013;33:781-796. doi: 10.3233/JAD-2012-120982

3.   Collaborators GBDDF. Estimation of the global prevalence of dementia in 2019 and
forecasted prevalence in 2050: an analysis for the Global Burden of Disease Study 2019.
Lancet Public Health. 2022;7:e105-e125. doi: 10.1016/S2468-2667(21)00249-8

4.   Jellinger KA. Alzheimer disease and cerebrovascular pathology: an update. *J Neural*
Transm (Vienna). 2002;109:813-836. doi: 10.1007/s007020200068

5.   Mehndiratta P, Manjila S, Ostergard T, Eisele S, Cohen ML, Sila C, Selman WR. Cerebral
amyloid angiopathy-associated intracerebral hemorrhage: pathology and management.
Neurosurg Focus. 2012;32:E7. doi: 10.3171/2012.1.FOCUS11370

6.   Gireud-Goss M, Mack AF, McCullough LD, Urayama A. Cerebral Amyloid Angiopathy and
Blood-Brain   Barrier   Dysfunction.   *Neuroscientist*.   2021;27:668-684.   doi:
10.1177/1073858420954811

7.   Greenberg SM, Gurol ME, Rosand J, Smith EE. Amyloid angiopathy-related vascular
cognitive   impairment.   *Stroke*.   2004;35:2616-2619.   doi:
10.1161/01.STR.0000143224.36527.44

8.   Shin HK, Jones PB, Garcia-Alloza M, Borrelli L, Greenberg SM, Bacskai BJ, Frosch MP,
Hyman BT, Moskowitz MA, Ayata C. Age-dependent cerebrovascular dysfunction in a
transgenic mouse model of cerebral amyloid angiopathy. *Brain*. 2007;130:2310-2319. doi:
10.1093/brain/awm156

9.   Iadecola C. The Neurovascular Unit Coming of Age: A Journey through Neurovascular
Coupling   in   Health   and   Disease.   *Neuron*.   2017;96:17-42.   doi:
10.1016/j.neuron.2017.07.030

10.   Nelson MT, Quayle JM. Physiological roles and properties of potassium channels in
arterial   smooth   muscle.   *Am   J   Physiol*.   1995;268:C799-822.   doi:
10.1152/ajpcell.1995.268.4.C799

11.   Brenner R, Perez GJ, Bonev AD, Eckman DM, Kosek JC, Wiler SW, Patterson AJ, Nelson
MT, Aldrich RW. Vasoregulation by the beta1 subunit of the calcium-activated potassium
channel. *Nature*. 2000;407:870-876. doi: 10.1038/35038011

12.   Jaggar JH, Porter VA, Lederer WJ, Nelson MT. Calcium sparks in smooth muscle. *Am J*
506   *Physiol Cell Physiol*. 2000;278:C235-256. doi: 10.1152/ajpcell.2000.278.2.C235

507   13.   Krishnamoorthy-Natarajan G, Koide M. BK Channels in the Vascular System. *Int Rev*
508   *Neurobiol*. 2016;128:401-438. doi: 10.1016/bs.irn.2016.03.017

509   14.   Tian L, Jeffries O, McClafferty H, Molyvdas A, Rowe IC, Saleem F, Chen L, Greaves J,
510   Chamberlain LH, Knaus HG, et al. Palmitoylation gates phosphorylation-dependent
511   regulation of BK potassium channels. *Proc Natl Acad Sci U S A*. 2008;105:21006-21011.
512   doi: 10.1073/pnas.0806700106

513   15.   Schubert R, Nelson MT. Protein kinases: tuners of the BKCa channel in smooth muscle.
514   *Trends Pharmacol Sci*. 2001;22:505-512. doi: 10.1016/s0165-6147(00)01775-2

515   16.   Kyle BD, Braun AP. The regulation of BK channel activity by pre- and post-translational
516   modifications. *Front Physiol*. 2014;5:316. doi: 10.3389/fphys.2014.00316

517   17.   Thuringer D, Findlay I. Contrasting effects of intracellular redox couples on the regulation
518   of maxi-K channels in isolated myocytes from rabbit pulmonary artery. *J Physiol*. 1997;500
519   (Pt 3):583-592. doi: 10.1113/jphysiol.1997.sp022044

520   18.   Lang RJ, Harvey JR, McPhee GJ, Klemm MF. Nitric oxide and thiol reagent modulation of
521   Ca2+-activated K+ (BKCa) channels in myocytes of the guinea-pig taenia caeci. *J Physiol*.
522   2000;525 Pt 2:363-376. doi: 10.1111/j.1469-7793.2000.00363.x

523   19.   Brzezinska AK, Gebremedhin D, Chilian WM, Kalyanaraman B, Elliott SJ. Peroxynitrite
524   reversibly inhibits Ca(2+)-activated K(+) channels in rat cerebral artery smooth muscle
525   cells. *Am J Physiol Heart Circ Physiol*. 2000;278:H1883-1890. doi:
526   10.1152/ajpheart.2000.278.6.H1883

527   20.   Wang ZW, Nara M, Wang YX, Kotlikoff MI. Redox regulation of large conductance Ca(2+)-
528   activated K+ channels in smooth muscle cells. *J Gen Physiol*. 1997;110:35-44. doi:
529   10.1085/jgp.110.1.35

530   21.   Soto MA, Gonzalez C, Lissi E, Vergara C, Latorre R. Ca(2+)-activated K+ channel
531   inhibition by reactive oxygen species. *Am J Physiol Cell Physiol*. 2002;282:C461-471. doi:
532   10.1152/ajpcell.00167.2001

533   22.   Furuta S. Basal S-Nitrosylation Is the Guardian of Tissue Homeostasis. *Trends Cancer*.
534   2017;3:744-748. doi: 10.1016/j.trecan.2017.09.003

535   23.   Butterfield DA, Boyd-Kimball D. Mitochondrial Oxidative and Nitrosative Stress and
536   Alzheimer Disease. *Antioxidants (Basel)*. 2020;9. doi: 10.3390/antiox9090818

24. Handy DE, Loscalzo J. Nitric oxide and posttranslational modification of the vascular
proteome: S-nitrosation of reactive thiols. *Arterioscler Thromb Vasc Biol*. 2006;26:1207-
1214. doi: 10.1161/01.ATV.0000217632.98717.a0

25. Su B, Wang X, Nunomura A, Moreira PI, Lee HG, Perry G, Smith MA, Zhu X. Oxidative
stress signaling in Alzheimer's disease. *Curr Alzheimer Res*. 2008;5:525-532. doi:
10.2174/156720508786898451

26. Asiimwe N, Yeo SG, Kim MS, Jung J, Jeong NY. Nitric Oxide: Exploring the Contextual
Link with Alzheimer's Disease. *Oxid Med Cell Longev*. 2016;2016:7205747. doi:
10.1155/2016/7205747

27. Girouard H, Park L, Anrather J, Zhou P, Iadecola C. Cerebrovascular nitrosative stress
mediates neurovascular and endothelial dysfunction induced by angiotensin II.
Arterioscler   Thromb   Vasc   Biol.   2007;27:303-309.   doi:
10.1161/01.ATV.0000253885.41509.25

28. Taylor JL, Pritchard HAT, Walsh KR, Strangward P, White C, Hill-Eubanks D, Alakrawi M,
Hennig GW, Allan SM, Nelson MT, Greenstein AS. Functionally linked potassium channel
activity in cerebral endothelial and smooth muscle cells is compromised in Alzheimer's
disease.   *Proc   Natl   Acad   Sci   U   S   A*.   2022;119:e2204581119.   doi:
10.1073/pnas.2204581119

29. Oakley H, Cole SL, Logan S, Maus E, Shao P, Craft J, Guillozet-Bongaarts A, Ohno M,
Disterhoft J, Van Eldik L, et al. Intraneuronal beta-amyloid aggregates,
neurodegeneration, and neuron loss in transgenic mice with five familial Alzheimer's
disease mutations: potential factors in amyloid plaque formation. *J Neurosci*.
2006;26:10129-10140. doi: 10.1523/JNEUROSCI.1202-06.2006

30. 2022 Alzheimer's disease facts and figures. *Alzheimers Dement*. 2022;18:700-789. doi:
10.1002/alz.12638

31. Peters EC, Gee MT, Pawlowski LN, Kath AM, Polk FD, Vance CJ, Sacoman JL, Pires PW.
Amyloid-beta disrupts unitary calcium entry through endothelial NMDA receptors in mouse
cerebral arteries. *Journal of cerebral blood flow and metabolism : official journal of the
International Society of Cerebral Blood Flow and Metabolism*. 2022;42:145-161. doi:
10.1177/0271678X211039592

32. Jackson WF. Myogenic Tone in Peripheral Resistance Arteries and Arterioles: The
Pressure Is On! *Front Physiol*. 2021;12:699517. doi: 10.3389/fphys.2021.699517

33. Newcombe EA, Camats-Perna J, Silva ML, Valmas N, Huat TJ, Medeiros R. Inflammation:
the link between comorbidities, genetics, and Alzheimer's disease. *J Neuroinflammation*.
2018;15:276. doi: 10.1186/s12974-018-1313-3

34. Cinelli MA, Do HT, Miley GP, Silverman RB. Inducible nitric oxide synthase: Regulation,
structure, and inhibition. *Med Res Rev*. 2020;40:158-189. doi: 10.1002/med.21599

35. Kleinert H, Schwarz PM, Forstermann U. Regulation of the expression of inducible nitric
oxide synthase. *Biol Chem*. 2003;384:1343-1364. doi: 10.1515/BC.2003.152

36. Forstermann U, Xia N, Li H. Roles of Vascular Oxidative Stress and Nitric Oxide in the
Pathogenesis   of   Atherosclerosis.   *Circ   Res*.   2017;120:713-735.   doi:
10.1161/CIRCRESAHA.116.309326

37. Weidinger A, Mullebner A, Paier-Pourani J, Banerjee A, Miller I, Lauterbock L, Duvigneau
JC, Skulachev VP, Redl H, Kozlov AV. Vicious inducible nitric oxide synthase-
mitochondrial reactive oxygen species cycle accelerates inflammatory response and
causes   liver   injury   in   rats.   *Antioxid   Redox   Signal*.   2015;22:572-586.   doi:
10.1089/ars.2014.5996

38. Viner RI, Williams TD, Schoneich C. Peroxynitrite modification of protein thiols: oxidation,
nitrosylation, and S-glutathiolation of functionally important cysteine residue(s) in the
sarcoplasmic   reticulum   Ca-ATPase.   *Biochemistry*.   1999;38:12408-12415.   doi:
10.1021/bi9909445

39. Newell KL, Hyman BT, Growdon JH, Hedley-Whyte ET. Application of the National
Institute on Aging (NIA)-Reagan Institute criteria for the neuropathological diagnosis of
Alzheimer   disease.   *J   Neuropathol   Exp   Neurol*.   1999;58:1147-1155.   doi:
10.1097/00005072-199911000-00004

40. Polycarpou A, Hricisak L, Iring A, Safar D, Ruisanchez E, Horvath B, Sandor P, Benyo Z.
Adaptation of the cerebrocortical circulation to carotid artery occlusion involves blood flow
redistribution between cortical regions and is independent of eNOS. *Am J Physiol Heart
Circ Physiol*. 2016;311:H972-H980. doi: 10.1152/ajpheart.00197.2016

41. Blackwell JA, Silva JF, Louis EM, Savu A, Largent-Milnes TM, Brooks HL, Pires PW.
Cerebral arteriolar and neurovascular dysfunction after chemically induced menopause in
mice.   *Am   J   Physiol   Heart   Circ   Physiol*.   2022;323:H845-H860.   doi:
10.1152/ajpheart.00276.2022

42. Breijyeh Z, Karaman R. Comprehensive Review on Alzheimer's Disease: Causes and
Treatment. *Molecules*. 2020;25. doi: 10.3390/molecules25245789

43.   Bryant A, Li Z, Jayakumar R, Serrano-Pozo A, Woost B, Hu M, Woodbury ME, Wachter
603   A, Lin G, Kwon T, et al. Endothelial Cells Are Heterogeneous in Different Brain Regions
604   and Are Dramatically Altered in Alzheimer's Disease. *J Neurosci*. 2023;43:4541-4557. doi:
605   10.1523/JNEUROSCI.0237-23.2023

606   44.   Andjelkovic AV, Situ M, Citalan-Madrid AF, Stamatovic SM, Xiang J, Keep RF. Blood-
607   Brain Barrier Dysfunction in Normal Aging and Neurodegeneration: Mechanisms, Impact,
608   and Treatments. *Stroke*. 2023;54:661-672. doi: 10.1161/STROKEAHA.122.040578

609   45.   Scheuermann BC, Parr SK, Schulze KM, Kunkel ON, Turpin VG, Liang J, Ade CJ.
610   Associations of Cerebrovascular Regulation and Arterial Stiffness With Cerebral Small
611   Vessel Disease: A Systematic Review and Meta-Analysis. *J Am Heart Assoc*.
612   2023;12:e032616. doi: 10.1161/JAHA.123.032616

613   46.   Niwa K, Kazama K, Younkin L, Younkin SG, Carlson GA, Iadecola C. Cerebrovascular
614   autoregulation is profoundly impaired in mice overexpressing amyloid precursor protein.
615   *Am J Physiol Heart Circ Physiol*. 2002;283:H315-323. doi: 10.1152/ajpheart.00022.2002

616   47.   Heutz R, Claassen J, Feiner S, Davies A, Gurung D, Panerai RB, Heus R, Beishon LC.
617   Dynamic cerebral autoregulation in Alzheimer's disease and mild cognitive impairment: A
618   systematic review. *Journal of cerebral blood flow and metabolism : official journal of the
619   International Society of Cerebral Blood Flow and Metabolism*. 2023;43:1223-1236. doi:
620   10.1177/0271678X231173449

621   48.   Nieves-Cintron M, Syed AU, Buonarati OR, Rigor RR, Nystoriak MA, Ghosh D, Sasse KC,
622   Ward SM, Santana LF, Hell JW, Navedo MF. Impaired BK(Ca) channel function in native
623   vascular smooth muscle from humans with type 2 diabetes. *Sci Rep*. 2017;7:14058. doi:
624   10.1038/s41598-017-14565-9

625   49.   Lu T, Chai Q, Jiao G, Wang XL, Sun X, Furuseth JD, Stulak JM, Daly RC, Greason KL,
626   Cha YM, Lee HC. Downregulation of BK channel function and protein expression in
627   coronary arteriolar smooth muscle cells of type 2 diabetic patients. *Cardiovasc Res*.
628   2019;115:145-153. doi: 10.1093/cvr/cvy137

629   50.   Grimm PR, Sansom SC. BK channels and a new form of hypertension. *Kidney Int*.
630   2010;78:956-962. doi: 10.1038/ki.2010.272

631   51.   Taylor JL, Walsh KR, Mosneag IE, Danby TGE, Luka N, Chanda B, Schiessl I, Dunne RA,
632   Hill-Eubanks D, Hennig GW, et al. Uncoupling of Ca(2+) sparks from BK channels in
633   cerebral arteries underlies hypoperfusion in hypertension-induced vascular dementia.
634   *Proc Natl Acad Sci U S A*. 2023;120:e2307513120. doi: 10.1073/pnas.2307513120

52.	Reed JT, Pareek T, Sriramula S, Pabbidi MR. Aging influences cerebrovascular myogenic reactivity and BK channel function in a sex-specific manner. *Cardiovasc Res*. 2020;116:1372-1385. doi: 10.1093/cvr/cvz314

53.	Le T, Martin-Aragon Baudel M, Syed A, Singhrao N, Pan S, Flores-Tamez VA, Burns AE, Man KNM, Karey E, Hong J, et al. Secondhand Smoke Exposure Impairs Ion Channel Function and Contractility of Mesenteric Arteries. *Function (Oxf)*. 2021;2:zqab041. doi: 10.1093/function/zqab041

54.	Liu YP, Zhang TN, Wen R, Liu CF, Yang N. Role of Posttranslational Modifications of Proteins in Cardiovascular Disease. *Oxid Med Cell Longev*. 2022;2022:3137329. doi: 10.1155/2022/3137329

55.	Bolotina VM, Najibi S, Palacino JJ, Pagano PJ, Cohen RA. Nitric oxide directly activates calcium-dependent potassium channels in vascular smooth muscle. *Nature*. 1994;368:850-853. doi: 10.1038/368850a0

56.	Hou S, Heinemann SH, Hoshi T. Modulation of BKCa channel gating by endogenous signaling molecules. *Physiology (Bethesda)*. 2009;24:26-35. doi: 10.1152/physiol.00032.2008

57.	Hou S, Xu R, Heinemann SH, Hoshi T. The RCK1 high-affinity Ca2+ sensor confers carbon monoxide sensitivity to Slo1 BK channels. *Proc Natl Acad Sci U S A*. 2008;105:4039-4043. doi: 10.1073/pnas.0800304105

58.	Liu Y, Terata K, Chai Q, Li H, Kleinman LH, Gutterman DD. Peroxynitrite inhibits Ca2+-activated K+ channel activity in smooth muscle of human coronary arterioles. *Circ Res*. 2002;91:1070-1076. doi: 10.1161/01.res.0000046003.14031.98

59.	Kinney JW, Bemiller SM, Murtishaw AS, Leisgang AM, Salazar AM, Lamb BT. Inflammation as a central mechanism in Alzheimer's disease. *Alzheimers Dement (N Y)*. 2018;4:575-590. doi: 10.1016/j.trci.2018.06.014

60.	Venema RC, Sayegh HS, Kent JD, Harrison DG. Identification, characterization, and comparison of the calmodulin-binding domains of the endothelial and inducible nitric oxide synthases. *J Biol Chem*. 1996;271:6435-6440. doi: 10.1074/jbc.271.11.6435

61.	Filosa JA, Bonev AD, Straub SV, Meredith AL, Wilkerson MK, Aldrich RW, Nelson MT. Local potassium signaling couples neuronal activity to vasodilation in the brain. *Nat Neurosci*. 2006;9:1397-1403. doi: 10.1038/nn1779

62.	Pritchard HAT, Pires PW, Yamasaki E, Thakore P, Earley S. Nanoscale remodeling of ryanodine receptor cluster size underlies cerebral microvascular dysfunction in Duchenne muscular dystrophy. *Proc Natl Acad Sci U S A*. 2018. doi: 10.1073/pnas.1804593115

63. Dennison JL, Ricciardi NR, Lohse I, Volmar CH, Wahlestedt C. Sexual Dimorphism in the
3xTg-AD Mouse Model and Its Impact on Pre-Clinical Research. *J Alzheimers Dis*.
2021;80:41-52. doi: 10.3233/JAD-201014

64. Toro CA, Zhang L, Cao J, Cai D. Sex differences in Alzheimer's disease: Understanding
the molecular impact. *Brain Res*. 2019;1719:194-207. doi: 10.1016/j.brainres.2019.05.031

65. Prange-Kiel J, Dudzinski DA, Prols F, Glatzel M, Matschke J, Rune GM. Aromatase
Expression in the Hippocampus of AD Patients and 5xFAD Mice. *Neural Plast*.
2016;2016:9802086. doi: 10.1155/2016/9802086

66. Arias-Loza PA, Muehlfelder M, Pelzer T. Estrogen and estrogen receptors in
cardiovascular oxidative stress. *Pflugers Arch*. 2013;465:739-746. doi: 10.1007/s00424-
013-1247-7

67. Tran QK. Reciprocality Between Estrogen Biology and Calcium Signaling in the
Cardiovascular System. *Front Endocrinol (Lausanne)*. 2020;11:568203. doi:
10.3389/fendo.2020.568203

68. Cheng YJ, Lin CH, Lane HY. From Menopause to Neurodegeneration-Molecular Basis
and Potential Therapy. *Int J Mol Sci*. 2021;22. doi: 10.3390/ijms22168654

69. Razmara A, Duckles SP, Krause DN, Procaccio V. Estrogen suppresses brain
mitochondrial oxidative stress in female and male rats. *Brain Res*. 2007;1176:71-81. doi:
10.1016/j.brainres.2007.08.036

70. Gadomsky L, Dos Santos Guilherme M, Winkler J, van der Kooij MA, Hartmann T, Grimm
M, Endres K. Elevated Testosterone Level and Urine Scent Marking in Male 5xFAD
Alzheimer Model Mice. *Curr Alzheimer Res*. 2020;17:80-92. doi:
10.2174/1567205017666200217105537

Figure Legends

Figure 1. *Cerebral microvascular hyper-contractility in 5x-FAD*. **A,E)** Representative traces of the

lumen diameter of pial arteries isolated from female (**A**) and male (**E**) WT (black, top) or *5x-FAD*

(lower, red) during the generation of spontaneous myogenic tone. Arrows indicate the moment

when intraluminal pressure was raised from 15 to 50 mmHg. **B, F)** Summary data showing higher

spontaneous myogenic tone in pial arteries from female (**B**) and male (**F**) *5x-FAD*. **C, G)** Pial

artery contractility to endothelin-1 (ET-1) was significantly increased in female (**C**) and male (**G**)

5x-FAD when compared to WT littermates. **D, H)** Receptor-independent contraction to a

depolarizing stimulus (60 mM KCl) was unchanged in female (**D**) and male (**H**). All data are means

± SEM. Each data point in graphs represents 1 pial artery from 1 mouse. Statistical analysis:

unpaired two-tailed Mann-Whitney (**B, F**) or unpaired two-tailed *Student*'s t-test (**C, D, G, H**).

Figure 2. *BK_{Ca} impairment in pial arteries from 5x-FAD.* **A, E)** Representative traces of the lumen

diameter of pial arteries isolated from female (**A**) and male (**E**) WT (black, top) or *5x-FAD* (lower,

red) showing constriction to the BK_{Ca} blocker iberiotoxin (IbTox, 30 nM). Note the blunted

constriction in *5x-FAD* mice. Orange boxes: regions where data were obtained**. B, F)** Summary

bar graph showing blunted vasoconstriction to IbTox in female (**B**) and male (**F**) *5x-FAD*. Each

data point in the graph represents 1 pial artery from 1 mouse. Statistical test: unpaired two-tailed

Mann-Whitney test. **C, G)** Representative traces of an inside-out patch clamp from freshly-isolated

pial artery smooth muscle cells showing single channel BK_{Ca} currents at a holding potential of +60

mV. Note the higher frequency of openings (downward deflections) in the patch from female (**C**)

WT (top trace, black) compared to *5x-FAD* (lower trace, red). Single channel BK_{Ca} openings were

not different between males WT and *5x-FAD* (**G**). C: closed; 1: 1 channel; 2: 2 channels. **D, H)**

Summary graph showing BK_{Ca} open probability (Po) over a range of holding potentials (+20, +40,

+60 mV). Po was significantly lower in inside-out patches from pial artery smooth muscle cells

isolated from female *5x-FAD* mice (**D**), whereas BK_{Ca} Po was not different between males *5x-*

FAD and WT littermates (**H**). Number of female replicates: WT n = 10 patches from 5 different

mice; *5x-FAD* n = 7 patches from 4 different mice. Number of male replicates: WT n = 9 patches

from 5 different mice; *5x-FAD* n = 9 patches from 6 different mice. All data are means ± SEM,

two-way ANOVA with a Sidak *post-hoc* test.

Figure 3. *BK_{Ca} impairment in males is a consequence of lower $BK_α$ expression and reduced Ca^{2+}*

sparks activity. **A-B)** Expression levels of $BK_α$ mRNA (**A**) and protein (**B**) are not different between

female *5x-FAD* and WT littermates. WB: Western blot. Each data point in the graph is a pial artery

lysate from 1 mouse. Statistical analysis: unpaired two-tailed *Student*'s t-test. **C-D)** Expression

levels of BK$_\alpha$ mRNA are similar between male *5x-FAD* and WT littermates (**C**); however BK$_\alpha$

protein expression is significantly lower in male *5x-FAD* when compared to WT littermates (**D**).

Each data point in the graph is a pial artery lysate from 1 mouse. Statistical analysis: unpaired

two-tailed *Student*'s t-test. **E-J)** Representative greyscale (**Ei, Fi**) and pseudocolored (**Eii, Fii**)

images of fields of view of pial arteries from female WT (**E**) and *5x-FAD* (**F**) prepared *reverse en*

face for recording of Ca^{2+} sparks (arrows). Ca^{2+} sparks frequency (**I**) was significantly higher in

pial arteries from female *5x-FAD* than in WT littermates, without differences in the number of

active sites per cell (**J**). Each data point in the graph is 1 field of view from a pial artery, for a total

of 5 fields of view per artery from 3 different mice of each genotype. Statistical analysis: unpaired

two-tailed *Student*'s t-test. **G-L)** Representative greyscale (**Gi, Hi**) and pseudocolored (**Gii, Hii**)

images of fields of view of pial arteries from male WT (**G**) and *5x-FAD* (**H**) prepared *reverse en*

face for recording of Ca^{2+} sparks (arrows). Ca^{2+} sparks frequency (**K**) was significantly lower in

pial arteries from male *5x-FAD* than in WT littermates, without differences in the number of active

sites per cell (**L**). Each data point in the graph is 1 field of view from a pial artery, for a total of 5

fields of view per artery from 3 different mice of each genotype. Statistical analysis: unpaired two-

tailed *Student*'s t-test. All data are means ± SEM.

Figure 4. *Oxidative-dependent BK$_{Ca}$ impairment in pial arteries from female 5x-FAD.* **A-C)**

Glutathione assay from brain lysates of female *5x-FAD* showing the presence of an oxidative

environment, observed as a significant increase in oxidized glutathione (GSSG, **B**) and the ratio

between GSSG and reduced glutathione (GSH, **C**). Each data point represents a brain lysate

from the cortex of an individual mouse. Statistical analysis: unpaired two-tailed Student's t-test.

D-E) Incubation of pial arteries from isolated female *5x-FAD* with the broad-spectrum reducing

agent dithiothreitol (DTT, 10 µM) significantly recovers sensitivity to iberiotoxin (30 nM), as

observed by the representative trace (**D**) and summary data (**E**). Orange boxes indicate the

regions where data were extracted. Each data point in the graph represents 1 pial artery from an

individual mouse. Statistical analysis: paired two-tailed *Student*'s t-test. **F-G)** Similarly, acute

incubation with DTT partially recovers BK$_{Ca}$ Po in freshly-isolated pial artery smooth muscle cell

from female *5x-FAD*, as observed in the representative traces (**F**) and summary data (**G**). N = 16

patches from 6 different mice. Statistical analysis: Matching Mixed-Model Anova with a Sidak

post-hoc correction for multiple comparisons. All data are means ± SEM.

Figure 5. *Exacerbated BK$_{Ca}$ S-NO in female 5x-FAD.* **A)** Higher iNOS mRNA expression in

lysates of pial arteries isolated from female *5x-FAD* than in WT littermates. Each data point

represents 1 pial artery lysate from an individual mouse. Statistical analysis: unpaired two-tailed

Student's t-test. B) Immunofluorescence labeling of iNOS (red) in pial arteries (collagen IV – Col

IV, green). Note the apparent higher fluorescence intensity in iNOS labeling in pial arteries from

female *5x-FAD* when compared to WT littermates. Bar = 20 µm. **C)** Blinded, randomized semi-

quantitative analysis of iNOS fluorescence intensity in pial arteries of female WT and *5x-FAD*.

Each data point represents the average fluorescence intensity of 5 pial arteries randomly imaged

from an individual mouse. Statistical analysis: unpaired two-tailed *Student*'s t-test. **D-E)**

Quantification of global S-NO proteins in cortical lysates of *5x-FAD* and WT littermates. Note the

higher expression of S-NO protein in lysates from *5x-FAD* cortex when compared to WT in the

representative blot (**D**) and summary data (**E**). Each data point represents a lysate from an

individual mouse. Statistical analysis: unpaired two-tailed Mann-Whitney test. **F-G)** Affinity-

purification (AP) of S-NO proteins in brain lysates followed by Western blot against BK$_\alpha$ show a

significantly higher expression of BK$_\alpha$ S-NO in female *5x-FAD* when compared to WT littermates,

as evidenced by the representative blot (**F**) and summary data (**G**). Each data point represents a

lysate from an individual mouse. Statistical analysis: unpaired two-tailed *Student*'s test. **H-K)** Co-

localization analysis using Mander's M1 (**I**) and M2 (**J**) coefficients, as well as the Pearson's

coefficient (**K**) show a significantly higher co-localization between of BK$_\alpha$ (**H**, green) and global S-

NO proteins (**H**, red) in pial arteries from female *5x-FAD* (fluorogram in **H**) Bar = 20 µm. Violin

plots represent the distribution of co-localized pixels in each image. Imaging of pial arteries was

performed in a blinded and randomized manner, a total of 3 arteries were imaged per mouse for

a total of 5 mice per group. Statistical analysis: unpaired two-tailed Student's t-test. Data in violin

plots are median ± 95% confidence intervals, all other data are means ± SEM.

Figure 6. *BK$_{Ca}$ S-NO in post-mortem human brain samples.* **A-B)** Western blot (WB) quantification

of BK$_\alpha$ expression in *post-mortem* brain lysates of patients without CAA / AD (Non-AD), CAA or

AD + CAA. Representative blots are shown in **A** and summary data in **B**. Each data point

represents a brain lysate from an individual age-matched woman (♀) or man (♂). Statistical

analysis: Brown-Forsythe and Welch ANOVA with a Dunnett T3 correction for multiple

comparisons. **C-D)** Affinity-purification (AP) of S-NO proteins in brain lysates followed by WB

against the BK$_\alpha$ subunit show a trend towards an increase in BK$_\alpha$ S-NO in AD + CAA patients

when compared to non-AD, as evidenced by the representative blots in **C** and summary data in

D. Each data point represents a brain lysate from an individual age-matched woman (♀) or man

($\male$). Statistical analysis: Brown-Forsythe and Welch ANOVA with a Dunnett T3 correction for

multiple comparisons. All data are means ± SEM.

Figure 7. *Impaired neurovascular coupling after acute BK$_{Ca}$ inhibition and in 5x-FAD.* **A-C)** Acute

application of the BK$_{Ca}$ blocker iberiotoxin (IbTox, 30 nM) atop the thinned-skull cranial window

blunted neurovascular responses following somatosensory stimulation in WT mice, as evidenced

by the pseudocolored perfusion maps (**A**), representative traces (**B**) of perfusion within the region

of interest (black rectangle) and summary data (**C**). Each data point represents an individual

mouse, females in square (□) and males in circles (○), after application of vehicle (□, ○) or IbTox

(■,●). Statistical analysis: paired two-tailed *Student*'s t-test. **D-I)** Neurovascular coupling

responses were significantly impaired in female (**D-F**), and showed a trend towards decrease in

male (**G-I**), *5x-FAD* when compared to WT littermates, as evidenced by the perfusion maps (**D,**

G), representative perfusion traces (**E, H**) and summary data (**F, I**). Each data point represents

an individual mouse. Statistical analysis: paired two-tailed *Student*'s t-test. PU = perfusion units.

All data are means ± SEM.

Figure 1. *Cerebral microvascular hypercontractility in 5x-FAD*

Figure 2. *BK_{Ca} impairment in pial arteries from 5x-FAD*

Figure 3. *BK_{Ca} impairment in males is a consequence of lower BK_α expression and reduced Ca^{2+} sparks activity.*

Figure 4. *Oxidative-dependent BK$_{Ca}$ impairment in pial arteries from female 5x-FAD*

Figure 5. *Exacerbated BK$_{Ca}$ S-NO in female 5x-FAD*

Figure 6. *BK_{Ca} S-NO in post-mortem human brain samples*

Figure 7. *Impaired neurovascular coupling after acute BK$_{Ca}$ inhibition and in 5x-FAD*