DIETS: a simple and sensitive assay to measure and control the

intake of complex solid foods, like high-fat diets, in *Drosophila*

Running title: measuring and controlling complex high calorie diet intake in flies

Manikrao R. Thakare[1, 2, ¶], Prerana Choudhary[1, 2,¶], Bhavna Pydah[1, 2], Suhas Sunke[1],

R Sai Prathap Yadav[3], Pavan Agrawal[3] and Gaurav Das[1*]

[1]*Brain and Feeding Behavior Laboratory, National Centre for Cell Science, S.P.*

Pune University Campus, Ganeshkhind, Pune 411007, India

[2]*Savitribai Phule Pune University, Ganeshkhind, Pune 411007, India*

[3]*Centre for Molecular Neurosciences, Kasturba Medical College, Manipal Academy*

of Higher Education, Udupi 576104, India

[*]Correspondence - gauravdas@nccs.res.in

[¶]These authors contributed equally to this work

Abstract

The fruit fly *Drosophila melanogaster* offers a powerful model to study how diet affects the body and brain. However, existing methods for measuring their food intake often rely on dyes or tags mixed with food, which can be inaccurate due to how the flies absorb and eliminate them. Capillary-based assays like CAFE directly measure consumption, but only work with liquids and shorten fly lifespan. Additionally, capillary assays are incompatible with delivering viscous foods like high-fat diets. Even solidified high-fat diets tend to be sticky death traps for flies. Another longstanding challenge for fly researchers is that dietary restriction in flies involves diluting food, leading to compensatory feeding. To address these shortcomings, we have developed DIETS, a sensitive feeding assay that can be implemented even in low-resource settings. DIETS eliminates the need for labels and directly weighs the solid food consumed by small groups of flies over extended periods of hours to weeks. It allows us to deliver precise amounts of food to flies and implement accurate dietary restrictions. Importantly, DIETS is compatible with studying energy-dense high-fat diets. Using DIETS, we observed that, unlike a high-sugar diet, an isocaloric high-fat diet did not improve the flies' ability to withstand starvation, even though they consumed more calories and had higher fat deposition.

36 **Keywords**

37 *Feeding assay, behavior, Drosophila, dietary restriction, (DR), food choice, high-fat*

38 *diets, high-sugar diets, HFD, HSD, starvation resistance, SR*

Introduction

The quality, quantity, and timing of food intake have a profound impact on the overall health, fecundity, cognition, and longevity of an organism. Food intake is influenced by internal states that reflect a lack of specific nutrients (Bjordal et al. 2014; Ribeiro and Dickson 2010; Toshima and Tanimura 2012; Prasad and Hens 2018; Steck et al. 2018; Musso, Junca, and Gordon 2021; Stafford et al. 2012; Malita et al. 2022; Morton, Meek, and Schwartz 2014), or by internal states that result from stress and pathogenesis (Crompton 1984; Yau and Potenza 2013). Food intake is also influenced by innate and learned sensory inputs, that signal the presence of nutrients (Shiu et al. 2022; Weaver et al. 2023; Das, Lin, and Waddell 2016; Oh et al. 2021; Fantino 1984). Humans and fruit flies share deep evolutionary conservation in physiology, metabolism, and nervous system function (Ugur, Chen, and Bellen 2016; Jennings 2011; Yamaguchi 2018; Diop and Bodmer 2012; Kornberg and Krasnow 2000). Precise genetic tools in flies allow the investigation of the neuro-physiological mechanisms that underlie appetite control and the myriad effects that follow the ingestion of specific foods (Lin, Senapati, and Tsao 2019; Guo, Pan, and Gong 2019; Yoshihara and Ito 2012; Owald, Lin, and Waddell 2015; Pool and Scott 2014).

For all such studies, the ability to measure, or control feeding precisely is critical. A deep understanding of the underlying biology remains incomplete without knowing how much food is consumed. Currently, there are multiple assays for estimating feeding in flies. However, they all suffer from problems that prevent their use across the board for different dietary compositions. Specifically, such shortcomings include relying on proxy measurements of feeding using dyes and other tags, using liquid-

only diets, incompatibility with longitudinal studies, and the need to sacrifice flies for

measurement.

As flies consume only ~ 0.5 - 2.5 microlitres of food in a day (William W. Ja et al.

2007; Wu et al. 2020; Deshpande et al. 2014; Carvalho, Kapahi, and Benzer 2005),

many fly-feeding assays rely on food tagged with dyes, radiolabeled molecules, or

short oligonucleotides (Shell et al. 2018, 2021; Park, Tran, and Atkinson 2018; Wu et

al. 2020; Geer, Olander, and Sharp 1970). The amount of 'tag' ingested, and

eventually excreted acts as a proxy for consumption. Each tag must be carefully

screened to determine whether their taste, absorption, incorporation, and excretion

influence measurement, especially when physiology changes with age or genotype

(Marx 2015; Deshpande et al. 2014; Shell et al. 2018, 2021). Quantification of an

ingested tag requires sacrificing flies. Only the EX-Q (excreta quantification) assay

avoids sacrificing flies by chasing out all ingested dye with an extended feeding

period on unlabeled food. Food intake was then estimated by dye quantification from

the excreta, leaving the flies alive for longitudinal experiments (Wu et al. 2020).

Ethologically, flies likely feed on liquid, solid, and semi-solid food sources. However

multiple tagged and direct assays have been developed for flies solely for measuring

liquid diet intake (Sellier, Reeb, and Marion-Poll 2011; Qi et al. 2015; Murphy et al.

2017; Yapici et al. 2016). Their designs are based on the CApillary FEeder Assay

(CAFE), where liquid food consumed from a microcapillary is directly quantified by

the drop in meniscus (William W. Ja et al. 2007). Recent reports show that feeding

liquid diets over days from capillaries could nutrient-deprive flies and affect both

lifespan and fecundity (Park, Tran, and Atkinson 2018; Lee et al. 2008; Vigne and

Frelin 2010; Deshpande et al. 2014).

Drosophila is being widely and effectively used to gauge the effect of high-fat or

high-sugar diets (HFD and HSD respectively) on the physiology, metabolism,

nervous system function and behavior (Birse et al. 2010; Laura Palanker Musselman

et al. 2011; L. P. Musselman, Fink, and Baranski 2019; Van Dam et al. 2021; Vaziri

et al. 2020; May et al. 2019; Yang et al. 2023; Liao, Amcoff, and Nässel 2020).

Capillary-based liquid assays are likely incompatible with viscous, multi-ingredient

laboratory diets, especially liquid high-fat diets. Even when agarified, solid high-fat

diets tend to become sticky and trap flies on them (Diop, Birse, and Bodmer 2017),

especially with rising temperatures. Such problems could be the reason why very

few fly studies with high-fat diets have attempted to measure actual food intake,

especially over long periods (Shi et al. 2021; Eickelberg et al. 2022).

Here, we describe a simple, and versatile feeding assay that directly measures the

actual amount of solid food consumed by groups of flies. Dubbed DIETS (Direct

Intake Estimation and Tracking of Solid food consumption), the assay is easily

adaptable and inexpensive to establish even in low-resource settings. DIETS can

measure feeding over both short and long periods, typically from 30 minutes to 24 h.

Further, by periodically switching flies to fresh DIETS vials, longitudinal feeding

measurement can be performed over weeks. By placing an additional food cup in a

DIETS vial, we could also monitor short- and long- term feeding preferences in flies

between two distinct foods. Crucially, the small surface area of the food make DIETS

perfectly suited for complex high-fat diets. Flies do not stick to the food and do not

die even at higher temperatures. We have also modified DIETS to a flat arena setup

which can be combined with videography to determine the position of flies during an

experiment and potentially record its contact with food.

DIETS addresses a significant drawback of fly dietary restriction (DR) assays. In

flies, DR is mainly achieved by diluting food or nutrients (Hodge et al. 2022;

Partridge, Piper, and Mair 2005; Carvalho, Kapahi, and Benzer 2005; Piper and

Partridge 2007). However, food dilution leads to compensatory feeding and could

complicate the interpretation of the results (Carvalho, Kapahi, and Benzer 2005;

Vigne and Frelin 2010). With DIETS, we demonstrate that flies could be restricted to

feeding an exact amount of complex foods by offering a predetermined amount of

food in the cups. Hence, DIETS would be extremely useful for DR and Time

Restricted Feeding (TRF) protocols with complex diets.

The mechanisms by which food intake may alter fitness are not all well understood.

We employed DIETS to assess how two distinct calorie-dense, isocaloric food

sources, one rich in sugar and the other rich in saturated fats, could affect

food/calorie intake and contribute towards starvation resistance (SR). Surprisingly,

we found that while flies ingested considerably higher calories on a high-fat diet than

on a high-sugar diet, starvation resistance was only enhanced with the latter. The

size of bodily fat deposits has been strongly correlated with starvation resistance.

Our findings indicate that energy intake and body fat reserves do not necessarily

correlate positively with the flies' response to starvation.

Results

Measurement of feeding by directly weighing food consumed

The DIETS assay quantifies solid food consumption in flies. Consumption is directly measured by weighing the amount of food offered in 4 mm diameter 'cups' to a group of flies at the start of an experiment and finally weighing the amount of food left over at the end of the experimental period (Fig. 1A, Fig. S1A, and Movie S1). A 0.75 % agar bed at the bottom of the vial minimizes evaporation and provides a separate water source. A mean food weight loss from a control diet (CD; see Table ST1) due to evaporation was calculated from evaporation control vials without flies and was subtracted from concurrent intake measurements (Fig. 1A, Fig. S1B). We typically estimated consumption in sated CS-Q flies over 2 to 24 h periods, after which experiments were either terminated, or flies were transferred to fresh DIETS vials for further measurements (Fig. 1A). All experiments were carried out at 25 $^{\circ}$C and 65 - 70 % relative humidity unless otherwise mentioned.

We empirically determined the size of the cups and the distance from the agar bed at which loss of food by evaporation from CD was minimal. Evaporation from a 4 mm diameter cup was lower than that from 6 mm or 8 mm diameter cups, and all were placed 1 mm away from the agar bed (Fig. S1C). Evaporation loss from a 4 mm diameter cup was minimal when placed 1 mm away from the agar bed and went up considerably when the distance of the cup from the agar bed was increased up to 5 mm at both 18 and 25 $^{\circ}$C. As expected, evaporation loss was higher at 25 $^{\circ}$C than at 18 $^{\circ}$C (Fig. S1D).

Intuitively, food consumption should not change significantly with small changes in

food cup position. We observed that the calculated CD consumption dropped if we

increased the distance of food cups from the agar bed from 1 mm to 3 mm, or 5 mm

when subtracting mean evaporation loss (Fig. S1E; left panel). However, the mean

consumption values remained unchanged when the mean evaporation loss was not

accounted for (Fig. S1E; right panel). This suggested that we were perhaps

overestimating evaporation loss using the evaporation control vials, especially when

cups were placed a little further away from the agar bed. As this overestimation of

evaporation loss was the minimum at 1 mm (Fig. S1D), we decided to keep cups at

this distance from the agar bed, unless otherwise required.

DIETS is a sensitive and precise feeding assay

To characterize the resolving power and consistency of the DIETS assay, we

measured feeding on CD after varying sex, group size, temperature, and food

dilutions. Such conditions have been reported to have affected consumption (Wu et

al. 2020; Klepsatel, Wildridge, and Gáliková 2019; William W. Ja et al. 2007;

Deshpande et al. 2014). Firstly, males and females were segregated into groups of

10 to 60 flies to measure food consumption over 24 h at 25 ^{o}C. Like previous reports

(Deshpande et al. 2014; Wu et al. 2020; Carvalho et al. 2006), the mean

consumption of mated females was between 40 % to 125 % more than males from

the same group size (Fig. 1B; see effect sizes or Δ consumption between two

groups). We also observed that the consumption rate per fly increased with the

group size and eventually plateaued with a group size of 50 for males and 25 for

females (Fig. 1B, Table ST2). Therefore, as reported earlier, the alteration of social

interactions with group size seemed to influence individual feeding behavior (Wong

et al. 2009). As expected, the total food consumed by a group increased with the

group size for both sexes and was clearly discernible with DIETS (Fig. S2A, S2B,

and Table ST2). In experiments with mated females, we observed extensive egg

deposition, mostly on the agar bed (Fig. S2C). Only ~ 2 - 4 eggs were found on a

food cup on average, with the number going up to ~ 8 eggs for 60 females (Fig.

S2D). As the mean egg weight was ~ 32   g (data not shown), egg weight could be

ignored, especially if the ones on the outside of the cup were cleared before final

weighing. In case larvae are seen on food, we suggest switching food more

frequently and performing the experiments at a lower temperature if possible.

A recent study in flies has shown that an increase in temperature enhances

metabolic rate and feeding (Klepsatel, Wildridge, and Gáliková 2019). We also saw a

clear increase in CD consumption by male flies over 24 h when we raised the

temperature from 18 ^{o}C to 22 ^{o}C and then from 22 ^{o}C to 25 ^{o}C (Fig. 1C; see effect

sizes). Consumption on CD was even higher at 32 ^{o}C (Fig. S6C). We could detect

such changes in temperature-dependent consumption with a sample size of 12 vials

per condition.

Existing solid food feeding assays can detect compensatory increase or decrease in

feeding in flies upon diluting or concentrating food (Deshpande et al. 2014; Wu et al.

2020). With DIETS, we were able to discern clear differences in feeding between 1X

CD and 0.5X CD (~ 21 % increase) or between 1X and 2X CD (~ 56 % decrease)

(Fig. 1D).

To directly compare the sensitivity of DIETS with the previously reported CAFE and

EX-Q data (Wu et al. 2020), we measured compensatory feeding from 1X to 2X and

to 4X concentrations of 2.5 % sucrose - 2.5 % yeast extract (2.5 % SY: 1X) diet

(n=10, to match with Wu et al. 2020). We compared the standardized effect sizes in

Hedge's g along with its MOE (half the magnitude of 95 % CIs of the Hedge's g: see

Materials and Methods for further details) across assays (Fig. 1E). Between 1X and

2X diets, the analysis showed that DIETS and CAFE report effect sizes that are

close to each other (-2.59 for DIETS, and -3.12 for CAFE). However, a larger effect

size is seen with EX-Q (-5.51), mainly because 1X consumption measured by EX-Q

was elevated (Fig. 1E). Between 2X and 4X diets, all three assays report quite

similar effect sizes (-2.87 for DIETS, -2.39 for CAFE, and -2.98 with EX-Q). Keeping

in mind the small size of the data sets (n=10), the MOE was also generally

comparable (Fig. 1E). We concluded that DIETS was as sensitive as CAFE and EX-

Q and likely equally precise (compare MOEs of all DIETS data reported in Table

ST2).

To test whether DIETS can be used to assess longitudinal feeding, we transferred

groups of 50 male flies into fresh assay vials every 24 h for 4 weeks. Actual

consumption was assessed over the 1^{st}, 4^{th}, 7^{th}, 14^{th}, 21^{st}, and 28^{th} day (schema;

Fig. 1F). Feeding was reliably measured over the whole period, with very little

change in feeding when compared to the 1st-day measurement, except on the 28^{th}

day when a considerable feeding decline was observed. We also observed that with

a large group size of 50, flies started to die around the 14^{th} day, and by the 28^{th} day,

almost 50 % of the flies had perished (Fig. 1F).

Next, to test whether flies were starving in DIETS due to a smaller food source and

hence eventually dying early, we compared the starvation resistance of male flies

(10, 25, and 50 flies/group) after being in DIETS vials or normal food vials for 1 day

and 7 days. We found that flies showed almost similar starvation resistance across

group sizes and modes of food delivery (Fig. S3A). Together, our results suggest

that no discernible change in survival is observed in DIETS up to 7 days. We also

measured feeding on CD in two other strains available in the lab, *CS-BZ,* and w^{1118},

and observed that *CS-BZ* had elevated feeding as compared to the other two strains

(Fig. S3B).

To demonstrate the suitability of DIETS with thermogenetic manipulations at 32 ^{o}C,

we activated Piezo-GAL4 labeled neurons using a temperature-sensitive cation

channel, dTRPA1 (Hamada et al. 2008) during a 30-minute feeding period at 32 °C.

Piezo-GAL4 labeled mechanosensory neurons densely innervate the crop and

anterior midgut and regulate ingestion volume by controlling the crop size (S. Min et

al. 2021). As reported earlier (S. Min et al. 2021), we too observed a defect in

feeding, in fact, a more severe one, upon artificial activation of Piezo-GAL4 neurons

compared to parental controls (Fig. S3C; left panel). Feeding reduction in Piezo-

GAL4>UAS-dTRPA1 flies was not observed at 21 °C (Fig. S3C; right panel).

Activation of serotonergic neurons labeled by R50H05-GAL4 induces hunger and

feeding in sated flies (Albin et al. 2015). We replicated these results with DIETS, with

increased feeding observed in R50H05-GAL4>UAS-dTRPA1 flies at 32 °C (Fig.

S3C; left panel), but not at 21 °C (Fig. S3C; right panel).

Short-term feeding measurements with DIETS

Next, we wanted to determine whether DIETS could be used for short-term feeding

measurements. For this, we starved groups of 50 male flies on 0.75 % agar for 6 h,

12 h, or 24 h. We then measured the consumption of these differentially starved

groups for 30 minutes in DIETS vials with Brilliant Blue FCF labeled CD (Fig. 2A,

schema). Parallelly, flies were also processed after 30 minutes to measure

consumption colorimetrically. Both techniques could detect the linear increase in

consumption with increasing starvation and also detect sizable effect size differences

with each step of starvation. For the sake of comparison across assay types, here

too, we plotted standardized effect sizes in the form of Hegde's g (Fig. 2A). With

DIETS, we had larger Hedge's g values (Fig. 2A) than with blue dye incorporation

across all comparisons (feeding after 6 -vs.- 0, 12 -vs.- 6 and 24 vs. 12 h starvation

times). The MOEs of these effect sizes are also comparable across the two assays

(Fig. 2A; see values represented as effect sizes, MOEs; also in Table ST2). These

results suggested that for short-term measurements, DIETS was more sensitive in

picking differences than blue dye incorporation but equally precise.

Next, to demonstrate whether we can use DIETS for repeated short-term feeding

measurements, we measured feeding every 2 h in groups of 50 male and female

flies for 2 days continuously (Fig. 2B, schema). Previous studies had reported two

peaks of feeding around the morning and evening light-dark transitions (May et al.

2019; Niu et al. 2021; Ro, Harvanek, and Pletcher 2014; Li et al. 2021; Xu, Zheng,

and Sehgal 2008). However, we observed only a single evening feeding peak

around 'lights-off' (Fig. 2B). Combining consumption measurements of the two 'peak'

groups spanning the 'lights off' time and comparing them with similarly merged 'day'

and 'night' groups on either side, we could see the effect of day to night transition on

feeding (Fig. S4A). We also observed that otherwise, feeding levels were quite

similar both during the day and night. From the above experiments, we concluded

that DIETS is also sensitive and precise for both discrete and repeated short-term

food intake measurements in sated flies.

Short-term and long-term two-choice food preference with DIETS

We reasoned that dietary choice between two foods could also be measured if flies

were given access to two cups of distinct foods. To validate food preference, we

arranged choices between CD and CD laced with increasing concentrations (2.5 mM

to 15 mM) of bitter-tasting quinine (QUI), for groups of 50 males, for 24 h (Fig. 2C,

schema). Cups were placed 1 mm from the agar bed and diametrically opposite to

each other. Fully assembled DIETS choice vials were covered with white paper to

act as a diffuser and prevent light bias (Fig. S4B; images). In control DIETS vials,

flies ate equally on average from both the cups with CD. As expected, flies

preferentially fed from the CD cup over the CD + QUI cup, with increasing

concentrations of quinine (Fig. 2C). However, combined consumption of food from

both cups remained unchanged across quinine concentrations (Fig. S4C). Similarly,

we measured short-term preference over 2 h in starved flies. We observed a strong

preference to feed from the CD cup over the CD + 15 mM QUI food cup (Fig. 2D).

Total consumption stayed constant in this case, as well (Fig. S4D). Food preference

in DIETS can also be represented as a consumption preference index (CPI) (Fig. 2C;

see inset). Corresponding CPI values are shown on top of the consumption plots

(Fig. 2C and 2D).

Implementing precise DR and TRF regimens with DIETS

Dietary restriction (DR) in flies is usually carried out by diluting food or protein

components (Partridge, Piper, and Mair 2005; Bass et al. 2007; Piper and Partridge

2007; W. W. Ja et al. 2009; K.-J. Min and Tatar 2006; Hodge et al. 2022; Katewa et

al. 2012). Since we could control the amount of food we offer to flies, we decided to

ascertain whether we could use DIETS to achieve precise DR.

In the 'no DR' control groups, 56 mg of CD was offered to groups of male flies on

average (Fig. 3A). We had determined that this amount of food was non-limiting to a

group of 50 male flies (see Material and Methods for details). To implement 90 %

DR, the mean food offered to a group of flies was adjusted to be ~90-91% of 56 mg

(51 mg), on average to groups of males (Fig. 3A; labeled 'a'). We similarly aimed for

various degrees of DR up to 60% [Fig. 3A; (a / 56) * 100. Ultimately, achieved DR

was calculated based on average actual consumption in each group (Fig. 3A;

labeled 'b'), in proportion to the mean consumption (43 mg) in the 'no DR' group [Fig.

3A; (b / 43) * 100]. We found that the amount fed by groups of flies at all the DR

levels clustered narrowly. This was also reflected in the extremely narrow MOEs of

each group (Fig. 3A, see error distribution of effect sizes. Also see Table ST2). We

concluded that the DIETS setup can be used to restrict the amount of food

consumed by a group of flies within a few percentage points.

Time-restricted feeding (TRF) protocols have been imposed on flies to gauge the

effect on the overall health and longevity of flies (Cabrera, Young, and Axelrod 2020;

Gill et al. 2015; Ulgherait et al. 2021). To investigate the impact of a daily pattern of

feeding and fasting on food intake with DIETS, we measured CD consumption in

groups of 20 male flies for 24 h under three different conditions: (a) the TRF group

where feeding was restricted to 'lights-on' phase, ZT = 0 - 12, (b) An *ad libitum*

feeding "ALF (24)" group with food cup changed every 24 h at ZT = 0, and (c) A

second a*d libitum* feeding group, "ALF (12 + 12)" with food cup changed every 12 h

at ZT = 0 and 12. All groups were habituated as described above for 3 days to let

them adjust to their respective conditions. Consumption was measured for the next

24 h period (Fig. 3B; schema). In agreement with previous reports (Cabrera, Young,

and Axelrod 2020; Gill et al. 2015), we found that flies under TRF consumed similar

amount of food during the daytime as the ALF (24) and ALF (12 + 12) groups

managed to eat over a whole day (Fig. 3B). Next, upon comparing daytime feeding

'a' with nighttime feeding 'b' within the ALF (12+12) group, our data revealed that

flies fed equally during the day and night (Fig. 3B). We concluded that DIETS can be

used to implement TRF protocols with solid foods and simultaneously measure

feeding too.

Combining consumption measurement with video tracking: DIETS-

arena

DIETS vials are not ideal for videographic analysis because of their dimensions and

also due to the location of the cups next to the agar bed. As a proof of concept that

video analysis of fly location can be combined with DIETS style measurement of

food consumption, DIETS-arenas were designed (Fig. S5A). First, to determine the

efficacy of the flat DIETS-arenas, we measured 2 h CD consumption in sated and

starved male flies with both DIETS vials and DIETS-arena side by side (Fig. 3C;

schema). We observed a similar increase in feeding in the starved groups compared

to the sated group with both vials and arenas (Fig. 3C, left panel). The feeding

baseline was higher in the DIETS-arenas compared to DIETS vials for both sated

and starved flies. Further investigation will be needed to determine the cause.

Next, to analyze the temporal dynamics of feeding with the same set of sated and

starved flies, we defined a Food$_{ROI}$ around the food cup in the center of the arena

(see Material and Methods). We found that the percentage of starved flies that

occupied the Food$_{ROI}$, went from 74 % to ~ 40 % in the first 20 minutes. Thereafter

starved flies maintained an average of ~ 44 % Food$_{ROI}$ occupancy. In contrast, fed

flies maintain a low and steady ~ 11 % occupancy of the Food$_{ROI}$ over 90 minutes

(Fig. 3C, right panel). This pattern is also reflected in temporal snapshots from two

representative sets, each of sated and starved flies (Fig. S5B). In conclusion, with

the help of video analysis from DIETS-arenas, we could visualize dynamic changes

in food occupancy that could be correlated with the increased feeding observed in

starved flies.

Longitudinal measurements of high-fat diet (HFD) intake with DIETS

High-fat diets (HFDs) for flies have been typically made by adding coconut oil as a

source of saturated fat to fly food (Birse et al. 2010; Diop, Birse, and Bodmer 2017).

HFDs are difficult to work with as the fat melts even at ~ 25 °C, rendering the food

sticky and lethal for flies unless care is taken (Diop, Birse, and Bodmer 2017). The

sticky consistency of fatty food could potentially affect feeding too. While a previous

study claims the use of CAFE for delivering high-fat food (Shi et al. 2021) , we

sought to independently verify the suitability of CAFE with 10 % sHFD made in the

lab. We re-confirmed earlier observations in the lab that such complex food tends to

separate within the CAFE capillaries, leading to gaps in the continuous supply of

food (Fig. S6A). We reasoned that DIETS, with its low food surface area, may

resolve such issues. We prepared HFDs with added 5 % to 20 % coconut oil at the

expense of water in the CD. We recalibrated evaporation control vials for HFDs and

found that HFD food cups were best placed 2 mm from the agar bed. At lesser

distances (1 mm), the HFD tended to accumulate moisture, leading to misleading

changes in weight (data not shown). Consumption was measured on the 1^{st}, $4^{th,}$ and

7^{th} day (Fig. 4A, schema). We found that consumption of diets with 5 % and 10 %

added coconut oil was comparable with CD throughout (Fig. 4A). As the added lipids

made the food more energy-rich, energy intake was increased at both these

conditions compared to CD (Fig. S6B). We also saw that consumption of the diet

with 20 % added oil was considerably reduced from all the other groups (Fig. 4A; see

effect sizes HFD). As a result, energy intake in the 20 % HFD group was virtually

similar or slightly increased to the 10 % HFD group (Fig. S6B). To determine whether

decreased food intake on HFDs reflects an energy limit sensing mechanism or

simply because higher fat concentrations are taste aversive to flies requires further

investigation. DIETS was also suitable for consumption measurement on HFD even

at 32 ^{o}C (Fig. S6C), enabling thermogenetic experiments with HFDs. Overall, we

concluded that DIETS was suitable for feeding measurements on HFDs over a wide

range of temperatures.

As reported earlier (Birse et al. 2010), upon lipophilic Oil-Red-O staining, we

detected increasing mid-gut fat deposition after 7 days with an increasing percentage

of oil in the flies' diet (Fig. S6D). This showed that HFD feeding in DIETS yielded

expected physiological changes. Next, to ascertain whether feeding from HFDs in

DIETS influenced gene expression changes in key genes, as reported earlier by

RNA-Seq (Stobdan et al. 2019), we performed q-PCR analysis. We extracted RNA

from the heads of male flies kept on either 20 % HFD or CD for 1, 4, and 7 days and

probed gene expression using q-PCR (see Materials and Methods). We re-analyzed

the raw RNA-Seq data generated previously (Stobdan et al. 2019) using DESeq2

(Love, Huber, and Anders 2014) and used p-adjusted < 0.05 as our threshold (Table

ST3). From this analysis, we selected four of the most differentially expressed genes

(DEGs) to analyze from our collected head RNA samples. Out of these, three were

upregulated, namely *takeout* (*to*), *Obp83a*, and *Cyp4e3*, and one down-regulated

gene, *TotA* (Fig. 4B). Our experiment revealed dynamic changes in the expression of

the selected genes due to HFD feeding. On day 1, the expression of these genes

showed variability as compared to the earlier study (Stobdan et al. 2019). However,

by day 4 and then on day 7, their expression was consistent with Stobdan et al.,

2019 (Fig. 4B). We concluded that DIETS enables us to measure feeding in a group

of flies and subsequently use the same animals for post-feeding analyses like gene

expression changes.

A saturated high-fat diet fails to enhance starvation resistance

Chronic consumption of diets rich in sugar improves starvation resistance (SR) in

female flies, likely due to increased bodily fat deposits (van Dam et al. 2020;

Chandegra et al. 2017; Lee and Jang 2014; Strilbytska et al. 2022; Baldal,

Brakefield, and Zwaan 2006; Ballard, Melvin, and Simpson 2008). However,

although sHFDs increase body fat deposits (Fig. S6D), a solitary study has studied

and reported a mild increase in SR upon sHFD feeding (Heinrichsen and Haddad

2012). As DIETS improve how sHFD is delivered to flies and likely decreases

aversion to food due to sticky surface texture, we wanted to re-examine the role of

sHFDs on SR. Hence, we provided OreR flies with an isocaloric 30 % high sugar diet

(HSD: ~21 % sucrose + other sugars) and 10 % sHFD (coconut oil) for 7 days.

Feeding was assessed on days 1, 4, and 7. Subsequently, flies were subjected to

starvation stress in 0.75 % agar vials and survival was ascertained every 24 h

(schema; Fig. 4C). We found that flies on 30 % HSD, consumed significantly less

food, but maintained a total energy intake comparable to flies on CD control food

(Fig. S6E and 4C). In contrast, flies on 10 % sHFD, consumed amounts

intermediated between those of flies on CD and 30 % HSD (Fig. S6E). However,

sHFD being energy dense, the flies consumed significantly more energy than the

control flies (Fig. 4C). Subsequently, we found that a 30 % HSD diet promotes

increased SR in female flies as reported earlier. However, 10 % sHFD-fed flies

showed near identical SR compared to control flies (Fig. 4D). We concluded that a

10% sHFD diet is unable to enhance SR in flies.

Discussion

DIETS is a direct measurement of food consumed by flies with no added dyes or tracers (Fig. 1). Therefore, physiological changes associated with genotype or age, including changes in the gut, do not influence measurement. While DIETS is compatible with group sizes varying from ~ 10 - 60 flies, we recommend that the group size should be optimized for specific experiments. When we measured food consumption in groups of 50 males, there was significant death after 2 weeks of feeding in DIETS vials (Fig. 1E). This could be due to increased stress and struggle for a small food source in larger groups, and likely to be mitigated by using a smaller group size for longitudinal measurements. Indeed, previous longevity studies have used much smaller group sizes of 10 - 15 flies (Lee et al. 2008; Metaxakis and Partridge 2013; Zanco et al. 2021; Krittika and Yadav 2020). We have shown that DIETS could be also utilized for measuring food preferences (Fig. 2). With DIETS we can easily monitor potential shifts in food preference between two, perhaps even three food sources, over days and weeks. The assay is also precise, and with proper humidity control, yields measurements with low variability (see MOEs in Table. ST2).

With regards to versatility, ease, and cost of use, only CAFE and EX-Q assays are comparable with DIETS (Fig. 1E). We show here that DIETS is as sensitive as CAFE (William W. Ja et al. 2007; Diegelmann et al. 2017) and EX-Q (Wu et al. 2020) to discern biologically relevant differences in food consumed (Fig. 1E; see effect sizes).

While CAFE directly measures liquid food intake, it has major drawbacks. Feeding long-term in CAFE significantly reduces both egg-laying and lifespan (Wong et al. 2009; Deshpande et al. 2014). Additionally, while studies have reported using CAFE

for HFDs (Shi et al. 2021), we show that complex diets like HFDs separate inside

capillaries, making it impossible to measure consumption (Fig. S6A). EX-Q is well

suited for longitudinal studies, for use with complex diets, for estimating food

preference with two compatible dyes, and possibly also for dietary restriction

protocols (Wu et al. 2020; Shell et al. 2021). However, any protocol requiring

measurement for short consecutive periods will not be possible with EX-Q because

of the ~ 3 h 'chase' phase on non-dyed food required to excrete out all the dye in

flies. Therefore, longitudinal measurements with EX-Q require an extra fly transfer

step and estimate daily consumption for only a 21 h period. Flies can independently

regulate food and water intake (Van Dam et al. 2021; Fanson, Yap, and Taylor 2012;

W. W. Ja et al. 2009). In DIETS vials, the agar bed provides a separate water source

to allow flies to meet their water needs independently from their food drive (Fig. 1A).

No separate water source is provided in the EX-Q assay (Wu et al. 2020). Ultimately,

EX-Q is still an indirect method of feeding quantification, and any change in dye

absorption or excretion with age or genotype would affect the measurement.

The DIETS assay is versatile and measures feeding over different time scales for

diverse foods, including high-fat diets that are difficult to work with (Diop, Birse, and

Bodmer 2017). This ability of DIETS is an especially significant feature as it allows

long-term experiments with HFDs with accurate fatty food intake measurement. We

are not aware of such studies in the current literature. One significant shortcoming of

DIETS is that it cannot monitor feeding individuality in single flies. Capacitance-

based single fly feeding assays like FLIC, flyPAD, optoPAD, or dFRAME are highly

sensitive and outperform DIETS in determining feeding microstructure, including

feeding duration and frequency of individual feeding bouts (Ro, Harvanek, and

Pletcher 2014; Itskov et al. 2014; Moreira et al. 2019; Niu et al. 2021; Weaver et al.

2023). However, these assays do not measure feeding directly and only quantify the

duration and frequency of fly proboscis contact with food. The contact of the

proboscis with food may not always correlate directly with the amount ingested and

may vary with age or genotype. Ultimately, these assays have much higher setup

and repair costs than DIETS and are not affordable for many. Here, we demonstrate

that direct food intake measurements can be carried out from flat and transparent

DIETS-arenas and combined with video-based temporal analysis of a fly's position.

Similar to previous studies (Niu et al. 2021; Jaime et al. 2023; Weaver et al. 2023),

with the right camera, lighting, and number of flies, the feeding microstructure of a

single 'tagged' fly in a group can also be recorded from DIETS-arenas.

The dilution of food or its protein components leads to compensatory feeding in flies

(Partridge, Piper, and Mair 2005; Chapman and Partridge 1996; Bass et al. 2007;

Mair et al. 2003). To the best of our knowledge, DIETS is the only assay that can

implement precise DR in flies without worrying about compensatory consumption

(Fig. 4A). Additionally, like CAFE, time-restricted feeding (TRF) protocols (Cabrera,

Young, and Axelrod 2020; Gill et al. 2015) are easily carried out with DIETS, with the

added advantage of being able to work with high-fat and high-sugar foods. This

ability to control and measure food intake will also make it possible to calibrate and

deliver precise amounts of drugs or infectious agents to a group of flies.

Multiple lines of evidence indicate that increased fat deposits strongly correlate with

increased SR in flies. Flies selected over generations for increased SR show

enhanced body fat deposits (Schwasinger-Schmidt, Kachman, and Harshman 2012;

Hardy et al. 2017; Chippindale, Chu, and Rose 1996; Harshman, Hoffmann, and

Clark 1999). Foods with higher carbohydrate content (or lower P: C ratio) show

increased bodily lipid content and increased SR (Lee and Jang 2014). As mentioned

above, other studies on HSD and subsequent SR support the same idea (van Dam

et al. 2020; Chandegra et al. 2017; Lee and Jang 2014; Strilbytska et al. 2022).

We tested SR in OreR flies after a week-long exposure to isocaloric HSD and sHFD.

We found that while HSD treatment expectedly improved SR over control, sHFD did

not (Fig. 4D). This was despite higher energy consumption on sHFD (Fig. 4C) and

likely increased fat deposit, similar to our observation for CS flies (Fig. S6D). We

therefore speculate that feeding on HSD versus sHFD, leads to the formation of

bodily fat deposits that could differ in aspects like composition, lipid droplet structure,

and the rate of utilization during starvation. Ultimately, one or more of these factors

could alter SR, and further investigations are needed to understand the exact

mechanisms.

Our DIETS platform will be ideal for similar studies, where precise delivery of

complex diets and accurate measurements of their consumption will be necessary.

We anticipate that the ability of DIETS to precisely estimate and control food intake,

its versatility, and the low cost of deployment will lead to its wide adoption.

Materials and Methods

A. Fly lines and rearing

We used the following fly strains for all experiments: Wild type Canton-S (CS-Q), Oregon-R (OreR), CS-BZ, and w^{1118}, Piezo-GAL4 (BDSC # 59266) (S. Min et al. 2021), R50H05-GAL4 (BDSC # 38764) (Albin et al. 2015), and UAS-dTRPA1 (Hamada et al. 2008). All flies were reared on modified Bloomington's semi-defined food (Laura Palanker Musselman et al. 2011) at 25 and 65 % humidity, with a 12 h:12 h light: dark cycle inside incubators. This food is referred to as the control diet (CD: see Table. ST1 for ingredients). Temperature-sensitive UAS-dTRPA1 flies were raised at 18 °C and transferred to 32 °C 30 minutes before the start of experiments to induce transgene expression.

B. Chemicals and other materials

Food ingredients, namely brewer's yeast (Prime brand, instant dry yeast), table sugar (Madhur brand, refined sugar) and cold-pressed coconut oil (Nutiva), Brilliant blue FCF food grade (Vidhi Manufacturers 2490) were procured from local vendors. Other components, like agar (Himedia: GRM666-500G), peptone (Himedia: GRM001-500G), yeast extract powder (Himedia: RM027-500G), sucrose (MP biomedicals: 194018), propionic acid (Sigma-Aldrich: P5561-1L), methyl-4-hydroxybenzoate (Himedia: GRM1899-500G), $MgSO_4$ X $7H_2O$ (Sigma-Aldrich: M1880), $CaCl_2$ X $2H_2O$ (Sigma-Aldrich: 223506), absolute ethanol (Merck: 64-17-5), Oil-red-O (Sigma-Aldrich: 00625-100G), quinine hydrochloride dihydrate (Sigma-Aldrich: Q1125-5G), Triton X-100 (MP biomedical: SKU:02194854-CF) and paraformaldehyde (Sigma-Aldrich: P6148-500G), 0.22 m syringe filter (MERK: Z260479) were procured from authorized vendors. Light filters used for uniform red

light are 182: light red, 158: deep orange, and 810: zoom diffusion 1 (Bright Light

International). The following chemicals were used for RT-PCR; Trizol (Invitrogen:

15596026), Zymo Direct Zol kit (Zymo Research: R2051), Zymo RNA Clean and

Concentrator kit (Zymo Research: R1013), High-Capacity cDNA Reverse

Transcription kit (Thermo- Fisher Scientific: 4368814), KAPA SYBR FAST Universal

(Sigma-Aldrich: KK4602), QuantStudio™ 6 flex Real-Time PCR System (Applied

Biosystems: 4485691).

C. Diet preparation

To prepare the Control Diet (CD), ingredients (Table ST1) were weighed and mixed

in 60 mL reverse osmosis (RO) filtered water. The mixture was then cooked in a

microwave oven, with 2-3 intermittent stirring for ~ 2 minutes. The food mix was then

allowed to cool to 60   before adding methylparaben, propionic acid, $CaCl_2$, and

$MgSO_4$. The volume was adjusted to 100 mL by adding RO water. This mixture was

then thoroughly stirred with a whisk. To prepare diets with 0.25X to 2X

concentrations of CD (Fig.1D), we separately weighed all the ingredients for each

concentration. Agar concentration was maintained at 1 % for all the dilutions. To

prepare CD spiked with quinine (Fig. 3C), the required amounts of quinine were

added to the molten CD, below 60   and mixed thoroughly by shaking or vortexing.

2.5 % sucrose - 2.5 % yeast diet (1X SY diet), 5 % sucrose - 5 % yeast diet (2X SY

diet), and 10 % Sucrose - 10 % yeast diet (4X SY diet) were prepared as previously

described (Deshpande et al. 2014; Wu et al. 2020). To prepare saturated high-fat

diets (5 %, 10 %, and 20 % HFDs, w/v), the required amount of molten coconut oil

was added to cooked food and then mixed with a whisk. The added coconut oil

replaced an equal volume of water in the recipe. The food mix was allowed to cool to

60   and methylparaben, propionic acid, CaCl$_2$, and MgSO$_4$ were added. The food

was mixed thoroughly again with a whisk, and the volume was adjusted to 100 mL

by adding RO water. A 10 % HFD was isocaloric to 30 % high-sugar diet (HSD). 9%

sucrose is present in the CD and to prepare 30 % w/v HSD, an extra ~ 21 % sucrose

was added to replace the water in the CD. High-fat diets (HFDs) and high-sugar

diets (HSDs) exhibited hygroscopic properties, leading to the accumulation of

moisture on the food surface. We empirically determined the distance at which the

food cups had to be fixed from the agar bed to prevent this. Typically it was between

~ 1- 5 mm.

D. Feeding quantification by DIETS assay

i) DIETS vial preparation

Standard fly vials (inner diameter: 23 mm, outer diameter: 25 mm, height: 90 mm)

and 200 µL microcentrifuge tube caps as food cups were assembled as follows. 7

mL 0.75 % agar solution was added to each vial to form an agar bed. After the agar

solidified, it was allowed to dry for 12 to 18 h at room temperature. If used

immediately, the agar layer leaks water, in which flies get trapped. An agar bed

reduces the weight loss from food cups due to evaporation and provides a separate

water source for flies to drink water. Solid food was provided in 200 µL

microcentrifuge tube (PCR-02-C; Axygen) caps (food cups) of 4 mm diameter. For

most experiments, ~ 70 µL of molten food from stored aliquots were pipetted into

food cups (Supplementary Fig. 1A). Once the food solidified, we noted the initial

weight of the cup + food. The initial weight of the food cups (cup + food) was kept

between a narrow range for consistency, typically ~ 125 - 135 mg for CD. Freshly

prepared food cups were fixed on the inside of the agar vials, normally at a distance

of 1 mm from the agar, with a small square of double-sided tape. Evaporation control

vials were prepared identically.

ii) Food consumption and calorie intake measurement

Flies were collected over the first day after eclosion, transferred to new food bottles,

and allowed to age for one more day. Lines that carried transgenic constructs were

aged for ~ 8 - 10 days after initial collection. On the 3rd day (~ 9th - 11th day for

transgenic lines), flies were segregated by sex under CO_2 anesthesia. Male or

female groups of 50 flies were placed in prepared DIETS vials and habituated for two

further days before actual consumption measurement. Flies were transferred to fresh

DIETS vials once after 24 h during the habituation process (Fig. 1A).

Fly groups were then transferred to DIETS vials with pre-weighed food cups for

feeding measurements. Evaporation control vials with no flies was maintained

alongside (Fig. 1A and Supplementary Fig. 1A). Viscosity of a specific food

influences the rate of evaporation. We also observed that certain viscous foods, like

foods with high fat and high sugar, in cups placed at 1 mm from the agar bed tended

to absorb moisture. Conversely, in diluted, less viscous foods, evaporation loss

increased. Therefore, with each new diet, we empirically determined the position of

the food cups from the agar bed to prevent moisture accumulation or loss by

evaporation in evaporation control vials below 10 % of the total food consumed. The

humidity level of the incubator could also be adjusted to keep evaporation below the

desired level. Most experiments were carried out at 25   and 65 % humidity.

Flies were transferred to fresh DIETS vials at the end of the period of feeding

measurements. The reduction in food cup weight (food consumed + evaporation

loss) was determined. The mean value of evaporation-mediated loss in weight (from

4 - 6 control vials) was deducted from the reduction in food cup weight to derive the

total food consumed by a group of flies. This value was usually plotted as an

average consumption per fly over time (e.g., mg/fly/day) or sometimes as total

consumption for a group of flies (mg/day).

Energy intake (in Fig. 4C and S6B) was calculated from the amount of food

consumed on CD and HFD, using the calorie values given in (Laura Palanker

Musselman et al. 2011) and represented as calorie/fly/day. The caloric value of the

Nutiva coconut oil was taken from the product label.

iii) Feeding measurement in starved flies with DIETS and blue dye uptake

Food intake in starved flies was measured by DIETS assay, and intake was parallelly

validated using a spectrophotometric estimation of Brilliant blue FCF dye. To

measure feeding in starved flies, 2 - 3 day old male flies were habituated in DIETS

vials for 2 days. For starvation, flies were kept in 0.75 % agar vials for specified

times. Flies were then transferred to DIETS vials with pre-weighed 1 % Brilliant blue

FCF 1 dye-labeled CD and allowed to feed for 30 minutes. Flies were homogenized

in 500 µl of PBST (1x PBS + 1 % Triton X-100) using a hand-held homogenizer

(BenchTop Lab Systems: BT-MT-13K). Homogenates were centrifuged at 4,000 g

for 5 minutes, and 100 µl of cleared supernatant was loaded into a 96-well plate.

Sample absorbance was measured at 630 nm on a multi-mode plate reader

(SPECTRAmax 250, Molecular Devices). Absorbance from a control set of flies fed

with food without dye was used to determine the baseline absorbance of fly lysates

(blank).

iv) Round-the-clock circadian feeding measurements

2 - 3 day old male and female flies were habituated in DIETS vials for two days. Feeding measurement was initiated at 12 noon [Zeitgeber time (ZT) = 3.5; our light-on time, or ZT = 0, was 8:30 AM] in fresh vials. Flies were then transferred to fresh DIETS vials every 2 h for the next 50 h, and continuous measurements were taken. The first 2 h feeding measurement was discarded, as it showed spiked feeding, likely due to the fact that they had been transferred to fresh food after 24 h. Fly transfers during the 12 h dark phase were carried out in red light.

v) Two food choice feeding measurements with DIETS

To measure food preference over varying time scales, two cups with distinct foods were placed 1 mm from the agar bed, diametrically opposite each other, inside DIETS vials. During the experiment, the two cups were kept parallel to the tray, facing each other. To avoid a bias due to uneven lighting, individual vials were wrapped in an 8.5 * 8.5 cm white printing paper sheet, acting as a light diffuser (Fig. S3A). Before measuring feeding from two food sources, flies were habituated in DIETS choice set up, with two CD cups for 48 h. Flies were starved, if the experiment demanded, after the habituation step. A consumption preference index (CPI) was also calculated as $CPI_{Food\ A}$ = [(feeding from Cup A - feeding from Cup B) / total consumption from both the food cups].

vi) DR protocol with DIETS

We have achieved DR in flies by offering a fixed amount of diets in food cups to a group of flies for 24 h, after 2 days of habituation on the CD diet. As a 'NO DR' control (*ad-libitum* feeding), an average of 56 mg of CD was offered to groups of 50

male flies. This value was derived from summing the mean consumption of 50 male

flies (~ 39 mg: Fig. S2A), mean evaporation from food cup (~ 4 mg: Fig. S1C), and

mean leftover food (~ 10-13 mg: data not shown). To implement increasing degrees

of DR (~ 90 % to 60 %), we gradually reduced the mean food offered in cups by ~ 3

to 5 mg (labeled 'a'; Fig. 4A). Targeted DR % was calculated with respect to 56 mg

offered in the 'no DR' group. Based on actual food consumed (labeled 'b'; Fig. 4A),

the achieved DR % was calculated with respect to 43 mg, the mean food consumed

by flies in the 'no DR' group (Fig. 4A).

vii) TRF feeding protocol with DIETS

Groups of 20 male flies each were split into three groups. Two *ad libitum* feeding

groups, where flies were switched to fresh food vials either every 24 h [ALF (24h)] or

every 12 h [ALF (12+12)] were set. A third time-restricted feeding (TRF) group was

offered food only during the 'lights on' phase (ZT = 0 - 12). TRF flies were switched

on DIETS vials with no cups during the 'lights off' phase (ZT = 12 - 24). All groups

were habituated similarly to their experimental conditions (schema; Fig. 4B). Food

consumption was measured whenever flies were switched to fresh vials as indicated

(schema, Fig. 4B).

E. Starvation resistance assay

To compare the starvation resistance of flies, we allowed flies in varying group sizes

(10, 25 and 50 males) to feed from DIETS vial and regular food vial for 1 day and 7

days, following two days of usual habituation. For 7 days of feeding in DIETS vials,

flies were transferred to new DIETS vials every 24 h, while normal vials were

changed on alternative days. Flies were then transferred to 0.75 - 1 % agar

starvation vials to measure starvation resistance. Dead flies were counted every 24

h. The survival time of individual flies was recorded after week-long feeding on HSD

or sHFD. The starvation resistance of flies was computed by the Kaplan Meier plot

using graphpad prism.

F. Gut lipid Oil-Red-O staining

For midgut-deposited fat droplet staining, the fly's abdomen was dissected in 1X

PBS with fine forceps (55-INOX, FST) to remove the gut. Gut tissues were fixed in 4

% paraformaldehyde (prepared from pre-dissolved solution) (Sigma-Aldrich: P6148-

500G) for 20 minutes at room temperature on a rotary shaker. Dissected guts are

then rinsed twice for 10 minutes each with 1X PBS to remove the fixative. Fixed guts

were then incubated in Oil-Red-O stain for 30 minutes. Oil-Red-O stains were

prepared by dissolving 10 mg Oil-Red-O (powder) in 10 mL isopropanol (0.1 % Oil-

Red-O). A 6:4 ratio of 0.1 % Oil-Red-O in isopropanol to 4 mL distilled water was

prepared and passed through a 0.22 m syringe filter (Merck: Z260479) before use.

The stained gut tissues were rinsed twice in distilled water for 10 minutes each.

Stained guts were then mounted on a glass slide, and images were captured using

an upright Nikon fluorescent microscope (Nikon Eclipse 80i) with a 20X objective

(Plan Fluor 20X, 0.50 NA Ph1 DLL ∞/0.17 WD 2.1).

G. RNA extraction and qPCR

After habituation on CD for 2 days, male flies were transferred on either 20 % HFD

or CD for 1, 4, and 7 days in DIETS vials and then flash-frozen and stored at −80 °C.

For RNA extraction, 30 flies were vortex-decapitated, and heads were collected on

dry ice, lysed in Trizol (Invitrogen: 15596026), and total RNA was extracted using a

Zymo Direct Zol kit (Cat No: R2051), and RNA was purified using a Zymo RNA

Clean and Concentrator kit (Cat No: R1013) as per the manufacturer's instructions.

After RNA extraction, cDNA was prepared using a High-Capacity cDNA Reverse

Transcription kit (Thermo-Fisher Scientific, 4368814). qPCR was performed using

KAPA SYBR FAST Universal (Sigma Aldrich, KK4602) on a QuantStudio™ 6 Flex

Real-Time PCR System (Applied Biosystems, 4485691). Fold change for each gene

was determined using the $2^{-\Delta\Delta Ct}$ method (Schmittgen and Livak 2008).

To obtain a comprehensive list of DEGs upon exposure to sHFD, we re-analysed the

raw data generated by a previous study (Stobdan et al. 2019) using DESeq2 (Love,

Huber, and Anders 2014) and used p-adjusted value of <0.05 as cut-off. Full analysis

results are included in Table ST3. From this analysis, we selected top three of the

most upregulated genes- *takeout*, *Obp83a*, and *Cyp4e3*, and one downregulated

gene-*TotA*. The primers used were as follows: For RPL32 (endogenous control:

FBgn0002626): RPL32-F: CGCCACCAGTCGGATCGATAT and RPL32-R:

CATGTGGCGGGTGCGCTTGTT. For TotA (FBgn0028396), TotA-F:

CTCTTATGTGCTTTGCACTGCTG and TotA-R: TTTTGGAGTCATCGTCCTGGG.

For Cyp4e3 (FBgn0015035), Cyp4e3-F: CCCGATGTCCAGAGGAAATTGT and

Cyp4e3-R: GCGCTGCGCCTCCTTAATA. For takeout (FBgn0039298), to-F:

CAGCGGAACTGGACAGAGTAAC and to-R: GCGCCTTGTCCCCATTGAACA. For

Obp83a (FBgn0011281), Obp83a-F: CGAGGACGAGAAGCTCAAATGC and

Obp83a-R: CGGATGGACGCAGCCCTTGG.

H. DIETS-arena and video analysis

We developed a flat arena setup, DIETS-arena, with an easily replaceable 3D printed food cup (transparent PLA material) in the center to enable temporal and spatial analysis of flies along with food consumption measurement. Each arena measured 6.5* 6.5 * 0.4 cm, and the whole setup comprised of six arenas (Fig. S6A; left panel). This design was adapted from a previous study (Charroux and Royet 2020). 0 - 4 day old male flies were sorted under CO_2 anesthesia in groups of 15 - 20 and habituated in DIETS vial with CD for 2 days. Arenas were cleaned with warm water and wiped clean thoroughly with a microfibre cloth. 500 L of 1 % agar was pipetted down in a ring form at a distance of 1 cm away from the edge of the central food cup (Fig. S6A; left panel). The agar was left to dry for 1 h, and subsequently, the arenas were covered with a transparent acrylic sheet. The initial weight of the food cup with CD was weighed and was immediately placed into the slot provided at the center of the arena to prevent evaporation. To transfer flies, the whole setup was inverted. Food cups from individual arenas were taken out one at a time, flies were transferred from a vial to the arena via a custom-designed funnel, and food cups were placed back immediately to prevent escape. The arena was then placed in the image capture setup (Fig. S6A; right panel). Arenas without flies were maintained alongside as evaporation controls. This experiment was initiated at ZT = 1. Images were captured via Arducam HQ IR-CUT camera with a 6 mm CS lens (Cat. 1251599, from robu.in), connected to a Raspberry Pi 4B. The Libcamera package was used to capture time-lapse images at 0.2 fps. The captured images were then undistorted using the Computer Vision Toolbox in MATLAB R2022b. These images were later compressed into videos of 10 fps using ImageJ (version 1.53t). A previously

published MATLAB code (Charroux and Royet 2020) was modified to quantify the

number of flies in a defined area.

To determine the number of flies on or near the food cup, we defined a circular zone

around the cup (Food$_{ROI}$), at a distance of 90 pixels from the center of the food cup.

This figure was a summation of the longest male body length (40 pixels: data not

shown) and the food cup radius (50 pixels; see Fig. S6A). As it sometimes proved

difficult to tell flies apart in the Food$_{ROI}$, especially when many were present, we

applied an inverted logic to determine the number of flies in each captured frame

outside the Food$_{ROI}$ in each arena. This ultimately allowed us to calculate the %

occupancy of Food$_{ROI}$ per frame as :

% occupancy of Food$_{ROI}$ = {(Total number of flies in an arena - number of flies

*detected outside Food$_{ROI}$)/Total number of flies in an arena} * 100*

I. Data plotting and statistics

GraphPad Prism version 8.3.1 was used for most main data plotting. Almost all data

are represented as scatter plots with error bars representing the mean with a 95 %

confidence interval (CI). Starvation resistance was plotted as Kaplan-Meier graph.

Estimation statistics, including effect sizes, the margin of error (MOE), and relevant

comparisons between groups performed for each experiment are reported in

Supplementary Table 2 (ST2). For all figures, unless mentioned, the effect size is

calculated as the mean difference between the two groups. This represents the

magnitude of difference between the two groups. The effect size is represented as

the dot in the spread. The two black lines that extend from the effect size dot denote

the 95% CI and are a measure of the precision of the data. To generate the CI,

around 5000 bias-corrected and accelerated (BCa) bootstrap resampling was done

and the effect size of each resample was plotted to generate the CI arms. The

magnitude of one arm of the confidence interval width represents the MOE. The

effect size for Fig. 1E and 2A was measured by the standardized mean difference,

Hedge's g. Hedge's g is the difference in mean of two groups, standardized over the

pooled standard deviation of the two groups, with a correction applied for small (~

less than 20) group sizes (Turner and Bernard 2006). All estimation/delta plots were

generated using an online website: www.estimationstats.com (Ho et al. 2019). We

extracted data for EX-Q and CAFE from (Wu et al. 2020) , Fig. 3C, using

WebPlotDigitizer (automeris.io/WebPlotDigitizer).

Acknowledgment

We thank all members of the brain and feeding behavior laboratory, NCCS, for their valuable insights in the synthesis of the manuscript. We would like to thank Radhika Mohandasan for early reviewing and aiding in the submission of the manuscript and Asmita Dogra for estimating the weight of fly eggs and Mohd. Tanzeel Aalam for helping with the video. We thank Deevitha Balasubramanian for testing the earliest versions of the DIETS assay. We thank the rest of our labs for their feedback on the manuscript. We thank our lab technician Rajkumar Pawar for managing lab purchases and preparing fly food. We thank Umesh Chachala and Jagtap for lab maintenance. MRT was funded by a Department of Biotechnology (DBT) doctoral scholarship. PC was funded by the University Grants Commission (UGC) and supported by a Ramanujan Fellowship award to GD (SB/S2/RJN-048/2017). SS was hired as a project JRF from a Core Research Grant (CRG/2019/005587) from SERB, DST. GoI. BP is funded by the Council of Scientific and Industrial Research (CSIR) doctoral scholarship. RSPY is supported by a TMA Pai fellowship from MAHE. PA lab is funded by the Ramalingaswami Re-entry Fellowship (BT/RLF, Re-entry/34/2018), and DBT, Research grant (BT/PR36166/BRB/10/1859/2020) by the Department of Biotechnology (DBT), Ministry of Science and Technology, Government of India. GD was a Ramanujan Fellow (SB/S2/RJN-048/2017) awarded by SERB. The GD lab is also funded by a Core Research Grant (CRG/2019/005587) from SERB and generous intramural funding support from NCCS.

Contributions

G.D. and M.R.T. and P.C. conceived the design of the DIETS assay. G.D. conceived the DIETS-arenas assay. S.S. built the DIETS-arenas and modified the code for DIETS-arena from work cited. Initial trials and design refinement was done by G.D, M.R.T, P.C, S.S, and B.P. Experiments on starvation resistance following HSD or sHFD were carried out by B.P. qPCR experiments were done by R.S.P.Y performed, and supervised by P.A. Both R.S.P.Y and P.A analyzed the data. All experiments, data analysis, and plotting in the manuscript were performed by G.D, P.A, M.R.T, P.C, B.P., S.S, and R.S.P.Y. Statistical analysis was performed by B.P and S.S. G.D directed the research and wrote the manuscript with inputs from all.

References

Albin, Stephanie D., Karla R. Kaun, Jon-Michael Knapp, Phuong Chung, Ulrike Heberlein, and Julie H. Simpson. 2015. "A Subset of Serotonergic Neurons Evokes Hunger in Adult *Drosophila*." *Current Biology: CB* 25 (18): 2435–40.

Baldal, Egon A., Paul M. Brakefield, and Bas J. Zwaan. 2006. "Multitrait Evolution in Lines of *Drosophila melanogaster* Selected for Increased Starvation Resistance: The Role of Metabolic Rate and Implications for the Evolution of Longevity." *Evolution; International Journal of Organic Evolution* 60 (7): 1435–44.

Ballard, J. William O., R. G. Melvin, and S. J. Simpson. 2008. "Starvation Resistance Is Positively Correlated with Body Lipid Proportion in Five Wild Caught *Drosophila simulans* Populations." *Journal of Insect Physiology* 54 (9): 1371–76.

Bass, Timothy M., Richard C. Grandison, Richard Wong, Pedro Martinez, Linda Partridge, and Matthew D. W. Piper. 2007. "Optimization of Dietary Restriction Protocols in *Drosophila*." *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences* 62 (10): 1071–81.

Birse, Ryan T., Joan Choi, Kathryn Reardon, Jessica Rodriguez, Suzanne Graham, Soda Diop, Karen Ocorr, Rolf Bodmer, and Sean Oldham. 2010. "High-Fat-Diet-Induced Obesity and Heart Dysfunction Are Regulated by the TOR Pathway in *Drosophila*." *Cell Metabolism* 12 (5): 533–44.

Bjordal, Marianne, Nathalie Arquier, Julie Kniazeff, Jean Philippe Pin, and Pierre Léopold. 2014. "Sensing of Amino Acids in a Dopaminergic Circuitry Promotes Rejection of an Incomplete Diet in *Drosophila*." *Cell* 156 (3): 510–21.

Cabrera, Daniel, Michael W. Young, and Sofia Axelrod. 2020. "Time-Restricted Feeding Prolongs Lifespan in *Drosophila* in a Peripheral Clock-Dependent Manner." *bioRxiv*.

Carvalho, Gil B., Pankaj Kapahi, David J. Anderson, and Seymour Benzer. 2006. "Allocrine Modulation of Feeding Behavior by the Sex Peptide of *Drosophila*." *Current Biology: CB* 16 (7): 692–96.

Carvalho, Gil B., Pankaj Kapahi, and Seymour Benzer. 2005. "Compensatory Ingestion upon Dietary Restriction in *Drosophila melanogaster*." *Nature Methods* 2 (11): 813–15.

Chandegra, Bhakti, Jocelyn Lok Yee Tang, Haoyu Chi, and Nazif Alic. 2017. "Sexually Dimorphic Effects of Dietary Sugar on Lifespan, Feeding and Starvation Resistance in *Drosophila*." *Aging* 9 (12): 2521–28.

Chapman, T., and L. Partridge. 1996. "Female Fitness in *Drosophila melanogaster*: An Interaction between the Effect of Nutrition and of Encounter Rate with Males." *Proceedings. Biological Sciences / The Royal Society* 263 (1371): 755–59.

Charroux, Bernard, and Julien Royet. 2020. "Protocol for a Binary Choice Feeding Assay Using Adult, Axenic *Drosophila*." *STAR Protocols* 1 (3): 100117.

891 Chippindale, Adam K., Terence J. F. Chu, and Michael R. Rose. 1996. "Complex
892   Trade-offs and the Evolution of Starvation Resistance in *Drosophila*
893   *melanogaster*." *Evolution; International Journal of Organic Evolution* 50 (2):
894   753–66.

895 Crompton, D. W. 1984. "Influence of Parasitic Infection on Food Intake." *Federation*
896   *Proceedings* 43 (2): 239–45.

897 Dam, Esther van, Lucie A. G. van Leeuwen, Eliano Dos Santos, Joel James, Lena
898   Best, Claudia Lennicke, Alec J. Vincent, et al. 2020. "Sugar-Induced Obesity
899   and Insulin Resistance Are Uncoupled from Shortened Survival in *Drosophila*."
900   *Cell Metabolism* 31 (4): 710–25.e7.

901 Das, Gaurav, Suewei Lin, and Scott Waddell. 2016. "Remembering Components of
902   Food in *Drosophila*." *Frontiers in Integrative Neuroscience* 10 (February): 4.

903 Deshpande, Sonali A., Gil B. Carvalho, Ariadna Amador, Angela M. Phillips, Sany
904   Hoxha, Keith J. Lizotte, and William W. Ja. 2014. "Quantifying *Drosophila* Food
905   Intake: Comparative Analysis of Current Methodology." *Nature Methods* 11 (5):
906   535–40.

907 Diegelmann, Soeren, Annika Jansen, Shreyas Jois, Katharina Kastenholz, Laura
908   Velo Escarcena, Nicole Strudthoff, and Henrike Scholz. 2017. "The CApillary
909   FEeder Assay Measures Food Intake in *Drosophila melanogaster*." *Journal of*
910   *Visualized Experiments: JoVE*, no. 121 (March).

911 Diop, Soda Balla, Ryan T. Birse, and Rolf Bodmer. 2017. "High Fat Diet Feeding and
912   High Throughput Triacylglyceride Assay in *Drosophila melanogaster*." *Journal of*
913   *Visualized Experiments: JoVE*, no. 127 (September).

914 Diop, Soda Balla, and Rolf Bodmer. 2012. "*Drosophila* as a Model to Study the
915   Genetic Mechanisms of Obesity-Associated Heart Dysfunction." *Journal of*
916   *Cellular and Molecular Medicine* 16 (5): 966–71.

917 Eickelberg, Virginia, Gerald Rimbach, Yvonne Seidler, Mario Hasler, Stefanie Staats,
918   and Kai Lüersen. 2022. "Fat Quality Impacts the Effect of a High-Fat Diet on the
919   Fatty Acid Profile, Life History Traits and Gene Expression in *Drosophila*
920   *melanogaster*." *Cells* 11 (24).

921 Fanson, Benjamin G., Sarsha Yap, and Phillip W. Taylor. 2012. "Geometry of
922   Compensatory Feeding and Water Consumption in *Drosophila melanogaster*."
923   *The Journal of Experimental Biology* 215 (Pt 5): 766–73.

924 Fantino, M. 1984. "Role of Sensory Input in the Control of Food Intake." *Journal of*
925   *the Autonomic Nervous System* 10 (3-4): 347–58.

926 Geer, B. W., R. M. Olander, and P. L. Sharp. 1970. "Quantification of Dietary Choline
927   Utilization in Adult *Drosophila melanogaster* by Radioisotope Methods." *Journal*
928   *of Insect Physiology* 16 (1): 33–43.

929 Gill, Shubhroz, Hiep D. Le, Girish C. Melkani, and Satchidananda Panda. 2015.
930   "Time-Restricted Feeding Attenuates Age-Related Cardiac Decline in
931   *Drosophila*." *Science* 347 (6227): 1265–69.

Guo, Chao, Yufeng Pan, and Zhefeng Gong. 2019. "Recent Advances in the Genetic
Dissection of Neural Circuits in *Drosophila*." *Neuroscience Bulletin* 35 (6): 1058–
72.

Hamada, Fumika N., Mark Rosenzweig, Kyeongjin Kang, Stefan R. Pulver, Alfredo
Ghezzi, Timothy J. Jegla, and Paul A. Garrity. 2008. "An Internal Thermal
Sensor Controlling Temperature Preference in *Drosophila*." *Nature* 454 (7201):
217–20.

Hardy, Christopher M., Molly K. Burke, Logan J. Everett, Mira V. Han, Kathryn M.
Lantz, and Allen G. Gibbs. 2017. "Genome-Wide Analysis of Starvation-
Selected *Drosophila melanogaster*—A Genetic Model of Obesity." *Molecular*
Biology and Evolution 35 (1): 50–65.

Harshman, Hoffmann, and Clark. 1999. "Selection for Starvation Resistance in
Drosophila melanogaster : Physiological Correlates, Enzyme Activities and
Multiple Stress Responses." *Journal of Evolutionary Biology* 12 (2): 370–79.

Heinrichsen, Erilynn T., and Gabriel G. Haddad. 2012. "Role of High-Fat Diet in
Stress Response of *Drosophila*." *PloS One* 7 (8): e42587.

Hodge, Brian A., Geoffrey T. Meyerhof, Subhash D. Katewa, Ting Lian, Charles Lau,
Sudipta Bar, Nicole Y. Leung, et al. 2022. "Dietary Restriction and the
Transcription Factor Clock Delay Eye Aging to Extend Lifespan in *Drosophila*
melanogaster." *Nature Communications* 13 (1): 3156.

Ho, Joses, Tayfun Tumkaya, Sameer Aryal, Hyungwon Choi, and Adam Claridge-
Chang. 2019. "Moving beyond P Values: Data Analysis with Estimation
Graphics." *Nature Methods* 16 (7): 565–66.

Itskov, Pavel M., José-Maria Moreira, Ekaterina Vinnik, Gonçalo Lopes, Steve
Safarik, Michael H. Dickinson, and Carlos Ribeiro. 2014. "Automated Monitoring
and Quantitative Analysis of Feeding Behaviour in *Drosophila*." *Nature*
Communications 5 (August): 4560.

Jaime, Maria D. L. A., Ghadi H. Salem, Daniel J. Martinez, Sean Karott, Alejandra
Flores, Cameron D. Palmer, Sharvani Mahadevaraju, et al. 2023. "Whole Animal
Feeding FLat (WAFFL): A Complete and Comprehensive Validation of a Novel,
High-Throughput Fly Experimentation System." *G3* 13 (3).

Ja, William W., Gil B. Carvalho, Elizabeth M. Mak, Noelle N. de la Rosa, Annie Y.
Fang, Jonathan C. Liong, Ted Brummel, and Seymour Benzer. 2007.
"Prandiology of *Drosophila* and the CAFE Assay." *Proceedings of the National*
Academy of Sciences of the United States of America 104 (20): 8253–56.

Ja, W. W., G. B. Carvalho, B. M. Zid, E. M. Mak, T. Brummel, and S. Benzer. 2009.
"Water- and Nutrient-Dependent Effects of Dietary Restriction on *Drosophila*
Lifespan." *Proceedings of the National Academy of Sciences*.

Jennings, Barbara H. 2011. "*Drosophila* – a Versatile Model in Biology & Medicine."
Materials Today 14 (5): 190–95.

Katewa, Subhash D., Fabio Demontis, Marysia Kolipinski, Alan Hubbard, Matthew S. Gill, Norbert Perrimon, Simon Melov, and Pankaj Kapahi. 2012. "Intramyocellular Fatty-Acid Metabolism Plays a Critical Role in Mediating Responses to Dietary Restriction in *Drosophila melanogaster*." *Cell Metabolism* 16 (1): 97–103.

Klepsatel, Peter, David Wildridge, and Martina Gáliková. 2019. "Temperature Induces Changes in *Drosophila* Energy Stores." *Scientific Reports* 9 (1): 5239.

Kornberg, T. B., and M. A. Krasnow. 2000. "The *Drosophila* Genome Sequence: Implications for Biology and Medicine." *Science* 287 (5461): 2218–20.

Krittika, Sudhakar, and Pankaj Yadav. 2020. "Dietary Protein Restriction Deciphers New Relationships between Lifespan, Fecundity and Activity Levels in Fruit Flies *Drosophila melanogaster*." *Scientific Reports* 10 (1): 10019.

Lee, Kwang Pum, and Taehwan Jang. 2014. "Exploring the Nutritional Basis of Starvation Resistance in *Drosophila melanogaster*." *Functional Ecology* 28 (5): 1144–55.

Lee, Kwang Pum, Stephen J. Simpson, Fiona J. Clissold, Robert Brooks, J. William O. Ballard, Phil W. Taylor, Nazaneen Soran, and David Raubenheimer. 2008. "Lifespan and Reproduction in *Drosophila*: New Insights from Nutritional Geometry." *Proceedings of the National Academy of Sciences of the United States of America* 105 (7): 2498–2503.

Liao, Sifang, Mirjam Amcoff, and Dick R. Nässel. 2020. "Impact of High-Fat Diet on Lifespan, Metabolism, Fecundity and Behavioral Senescence in *Drosophila*." *Insect Biochemistry and Molecular Biology*, November, 103495.

Lin, Suewei, Bhagyashree Senapati, and Chang-Hui Tsao. 2019. "Neural Basis of Hunger-Driven Behaviour in *Drosophila*." *Open Biology* 9 (3): 180259.

Li, Wanhe, Zikun Wang, Sheyum Syed, Cheng Lyu, Samantha Lincoln, Jenna O'Neil, Andrew D. Nguyen, Irena Feng, and Michael W. Young. 2021. "Chronic Social Isolation Signals Starvation and Reduces Sleep in *Drosophila*." *Nature* 597 (7875): 239–44.

Love, Michael I., Wolfgang Huber, and Simon Anders. 2014. "Moderated Estimation of Fold Change and Dispersion for RNA-Seq Data with DESeq2." *Genome Biology* 15 (12): 550.

Mair, William, Patrick Goymer, Scott D. Pletcher, and Linda Partridge. 2003. "Demography of Dietary Restriction and Death in *Drosophila*." *Science* 301 (5640): 1731–33.

Malita, Alina, Olga Kubrak, Takashi Koyama, Nadja Ahrentløv, Michael J. Texada, Stanislav Nagy, Kenneth V. Halberg, and Kim Rewitz. 2022. "A Gut-Derived Hormone Suppresses Sugar Appetite and Regulates Food Choice in *Drosophila*." *Nature Metabolism* 4 (11): 1532–50.

Marx, Vivien. 2015. "Metabolism: Feeding Fruit Flies." *Nature Methods* 12 (7): 609–12.

May, Christina E., Anoumid Vaziri, Yong Qi Lin, Olga Grushko, Morteza Khabiri,
    Qiao-Ping Wang, Kristina J. Holme, et al. 2019. "High Dietary Sugar Reshapes
    Sweet Taste to Promote Feeding Behavior in *Drosophila melanogaster*." *Cell
    Reports* 27 (6): 1675–85.e7.

Metaxakis, Athanasios, and Linda Partridge. 2013. "Dietary Restriction Extends
    Lifespan in Wild-Derived Populations of *Drosophila melanogaster*." *PloS One* 8
    (9): e74681.

Min, Kyung-Jin, and Marc Tatar. 2006. "*Drosophila* Diet Restriction in Practice: Do
    Flies Consume Fewer Nutrients?" *Mechanisms of Ageing and Development* 127
    (1): 93–96.

Min, Soohong, Yangkyun Oh, Pushpa Verma, Samuel C. Whitehead, Nilay Yapici,
    David Van Vactor, Greg Sb Suh, and Stephen Liberles. 2021. "Control of
    Feeding by Piezo-Mediated Gut Mechanosensation in *Drosophila*." *eLife* 10
    (February).

Moreira, José-Maria, Pavel M. Itskov, Dennis Goldschmidt, Celia Baltazar, Kathrin
    Steck, Ibrahim Tastekin, Samuel J. Walker, and Carlos Ribeiro. 2019.
    "optoPAD, a Closed-Loop Optogenetics System to Study the Circuit Basis of
    Feeding Behaviors." *eLife* 8 (June).

Morton, Gregory J., Thomas H. Meek, and Michael W. Schwartz. 2014.
    "Neurobiology of Food Intake in Health and Disease." *Nature Reviews.
    Neuroscience* 15 (6): 367–78.

Murphy, Keith R., Jin Hong Park, Robert Huber, and William W. Ja. 2017.
    "Simultaneous Measurement of Sleep and Feeding in Individual *Drosophila*."
    Nature Protocols 12 (11): 2355–66.

Musselman, Laura Palanker, Jill L. Fink, Kirk Narzinski, Prasanna Venkatesh
    Ramachandran, Sumitha Sukumar Hathiramani, Ross L. Cagan, and Thomas J.
    Baranski. 2011. "A High-Sugar Diet Produces Obesity and Insulin Resistance in
    Wild-Type *Drosophila*." *Disease Models & Mechanisms* 4 (6): 842–49.

Musselman, L. P., J. L. Fink, and T. J. Baranski. 2019. "Similar Effects of High-
    Fructose and High-Glucose Feeding in a *Drosophila* Model of Obesity and
    Diabetes." *PloS One*. 14 (5):e0217096.

Musso, Pierre-Yves, Pierre Junca, and Michael D. Gordon. 2021. "A Neural Circuit
    Linking Two Sugar Sensors Regulates Satiety-Dependent Fructose Drive in
    Drosophila." *Science Advances* 7 (49): eabj0186.

Niu, Mengxia, Xiaohang Zhang, Weihan Li, Jianxun Wang, and Yan Li. 2021.
    "dFRAME: A Video Recording-Based Analytical Method for Studying Feeding
    Rhythm in *Drosophila*." *Frontiers in Genetics* 12 (October): 763200.

Oh, Soo Min, Kyunghwa Jeong, Jeong Taeg Seo, and Seok Jun Moon. 2021.
    "Multisensory Interactions Regulate Feeding Behavior in *Drosophila*."
    Proceedings of the National Academy of Sciences of the United States of

America 118 (7).

Owald, David, Suewei Lin, and Scott Waddell. 2015. "Light, Heat, Action: Neural
Control of Fruit Fly Behaviour." *Philosophical Transactions of the Royal Society*
of London. Series B, Biological Sciences 370 (1677): 20140211.

Park, Annie, Tracy Tran, and Nigel S. Atkinson. 2018. "Monitoring Food Preference
in *Drosophila* by Oligonucleotide Tagging." *Proceedings of the National*
Academy of Sciences of the United States of America 115 (36): 9020–25.

Partridge, Linda, Matthew D. W. Piper, and William Mair. 2005. "Dietary Restriction
in *Drosophila*." *Mechanisms of Ageing and Development* 126 (9): 938–50.

Piper, Matthew D. W., and Linda Partridge. 2007. "Dietary Restriction in *Drosophila*:
Delayed Aging or Experimental Artefact?" *PLoS Genetics* 3 (4): e57.

Pool, A. H., and K. Scott. 2014. "Feeding Regulation in *Drosophila*." *Current Opinion*
in Neurobiology.

Prasad, Naveen, and Korneel Hens. 2018. "Sugar Promotes Feeding in Flies via the
Serine Protease Homolog Scarface." *Cell Reports* 24 (12): 3194–3206.e4.

Qi, Wei, Zhe Yang, Ziao Lin, Jin-Yong Park, Greg S. B. Suh, and Liming Wang.
2015. "A Quantitative Feeding Assay in Adult *Drosophila* Reveals Rapid
Modulation of Food Ingestion by Its Nutritional Value." *Molecular Brain* 8
(December): 87.

Ribeiro, Carlos, and Barry J. Dickson. 2010. "Sex Peptide Receptor and Neuronal
TOR/S6K Signaling Modulate Nutrient Balancing in *Drosophila*." *Current*
Biology: CB 20 (11): 1000–1005.

Ro, Jennifer, Zachary M. Harvanek, and Scott D. Pletcher. 2014. "FLIC: High-
Throughput, Continuous Analysis of Feeding Behaviors in *Drosophila*." *PloS*
One 9 (6): e101107.

Schmittgen, Thomas D., and Kenneth J. Livak. 2008. "Analyzing Real-Time PCR
Data by the Comparative C(T) Method." *Nature Protocols* 3 (6): 1101–8.

Schwasinger-Schmidt, T. E., S. D. Kachman, and L. G. Harshman. 2012. "Evolution
of Starvation Resistance in *Drosophila melanogaster*: Measurement of Direct
and Correlated Responses to Artificial Selection." *Journal of Evolutionary*
Biology 25 (2): 378–87.

Sellier, Marie-Jeanne, Pablo Reeb, and Frédéric Marion-Poll. 2011. "Consumption of
Bitter Alkaloids in *Drosophila melanogaster* in Multiple-Choice Test Conditions."
Chemical Senses 36 (4): 323–34.

Shell, Brandon C., Yuan Luo, Scott Pletcher, and Mike Grotewiel. 2021. "Expansion
and Application of Dye Tracers for Measuring Solid Food Intake and Food
Preference in *Drosophila*." *Scientific Reports* 11 (1): 20044.

Shell, Brandon C., Rebecca E. Schmitt, Kristen M. Lee, Jacob C. Johnson, Brian Y.
Chung, Scott D. Pletcher, and Mike Grotewiel. 2018. "Measurement of Solid
Food Intake in *Drosophila* via Consumption-Excretion of a Dye Tracer."

Scientific Reports 8 (1): 11536.

Shi, Dan, Tianshu Han, Xia Chu, Huimin Lu, Xue Yang, Tianqi Zi, Yanhe Zhao, et al. 2021. "An Isocaloric Moderately High-Fat Diet Extends Lifespan in Male Rats and *Drosophila*." *Cell Metabolism* 33 (3): 581–97.e9.

Shiu, Philip K., Gabriella R. Sterne, Stefanie Engert, Barry J. Dickson, and Kristin Scott. 2022. "Taste Quality and Hunger Interactions in a Feeding Sensorimotor Circuit." *eLife* 11 (July).

Stafford, Jeffrey W., Kaylea M. Lynd, Aera Y. Jung, and Michael D. Gordon. 2012. "Integration of Taste and Calorie Sensing in *Drosophila*." *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 32 (42): 14767–74.

Steck, Kathrin, Samuel J. Walker, Pavel M. Itskov, Célia Baltazar, José-Maria Moreira, and Carlos Ribeiro. 2018. "Internal Amino Acid State Modulates Yeast Taste Neurons to Support Protein Homeostasis in *Drosophila*." *eLife* 7 (February).

Stobdan, Tsering, Debashis Sahoo, Priti Azad, Iain Hartley, Erilynn Heinrichsen, Dan Zhou, and Gabriel G. Haddad. 2019. "High Fat Diet Induces Sex-Specific Differential Gene Expression in *Drosophila melanogaster*." *PloS One* 14 (3): e0213474.

Strilbytska, Olha, Tetiana Strutynska, Uliana Semaniuk, Nadia Burdyliyk, Volodymyr Bubalo, and Oleh Lushchak. 2022. "Dietary Sucrose Determines Stress Resistance, Oxidative Damages, and Antioxidant Defense System in *Drosophila*." *Scientifica* 2022 (May): 7262342.

Toshima, Naoko, and Teiichi Tanimura. 2012. "Taste Preference for Amino Acids Is Dependent on Internal Nutritional State in *Drosophila melanogaster*." *The Journal of Experimental Biology* 215 (Pt 16): 2827–32.

Turner, Herbert M. , Iii, and Robert M. Bernard. 2006. "Calculating and Synthesizing Effect Sizes." *Contemporary Issues in Communication Science and Disorders: CICSD* 33 (Spring): 42–55.

Ugur, Berrak, Kuchuan Chen, and Hugo J. Bellen. 2016. "*Drosophila* Tools and Assays for the Study of Human Diseases." *Disease Models & Mechanisms* 9 (3): 235–44.

Ulgherait, Matt, Adil M. Midoun, Scarlet J. Park, Jared A. Gatto, Samantha J. Tener, Julia Siewert, Naomi Klickstein, Julie C. Canman, William W. Ja, and Mimi Shirasu-Hiza. 2021. "Circadian Autophagy Drives iTRF-Mediated Longevity." *Nature* 598 (7880): 353–58.

Van Dam, E., L. A. G. van Leeuwen, E. Dos Santos, J. James, L. Best, C. Lennicke, A. J. Vincent, et al. 2021. "Sugar-Induced Obesity and Insulin Resistance Are Uncoupled from Shortened Survival in *Drosophila*." *Yearbook of Paediatric Endocrinology*.

Vaziri, Anoumid, Morteza Khabiri, Brendan T. Genaw, Christina E. May, Peter L.

Freddolino, and Monica Dus. 2020. "Persistent Epigenetic Reprogramming of
Sweet Taste by Diet." *Science Advances* 6 (46).

Vigne, Paul, and Christian Frelin. 2010. "Food Presentation Modifies Longevity and
the Beneficial Action of Dietary Restriction in *Drosophila*." *Experimental*
Gerontology 45 (2): 113–18.

Weaver, K. J., S. Raju, R. A. Rucker, T. S. Chakraborty, R. A. Holt, and S. D.
Pletcher. 2023. "Behavioral Dissection of Hunger States in *Drosophila*." *eLife* 12
(June): RP84537.

Wong, Richard, Matthew D. W. Piper, Bregje Wertheim, and Linda Partridge. 2009.
"Quantification of Food Intake in *Drosophila*." *PloS One* 4 (6): e6063.

Wu, Qi, Guixiang Yu, Scarlet J. Park, Yue Gao, William W. Ja, and Mingyao Yang.
2020. "Excreta Quantification (EX-Q) for Longitudinal Measurements of Food
Intake in *Drosophila*." *iScience* 23 (1): 100776.

Xu, Kanyan, Xiangzhong Zheng, and Amita Sehgal. 2008. "Regulation of Feeding
and Metabolism by Neuronal and Peripheral Clocks in *Drosophila*." *Cell*
Metabolism 8 (4): 289–300.

Yamaguchi, Masamitsu. 2018. *Drosophila Models for Human Diseases*. Springer.

Yang, Jie, Ruijun Tang, Shiye Chen, Yinan Chen, Kai Yuan, Rui Huang, and Liming
Wang. 2023. "Exposure to High-Sugar Diet Induces Transgenerational Changes
in Sweet Sensitivity and Feeding Behavior via H3K27me3 Reprogramming."
eLife 12 (September).

Yapici, Nilay, Raphael Cohn, Christian Schusterreiter, Vanessa Ruta, and Leslie B.
Vosshall. 2016. "A Taste Circuit That Regulates Ingestion by Integrating Food
and Hunger Signals." *Cell* 165 (3): 715–29.

Yau, Y. H. C., and M. N. Potenza. 2013. "Stress and Eating Behaviors." *Minerva*
Endocrinologica 38 (3): 255–67.

Yoshihara, Moto, and Kei Ito. 2012. "Acute Genetic Manipulation of Neuronal Activity
for the Functional Dissection of Neural Circuits-a Dream Come True for the
Pioneers of Behavioral Genetics." *Journal of Neurogenetics* 26 (1): 43–52.

Zanco, Brooke, Christen K. Mirth, Carla M. Sgrò, and Matthew Dw Piper. 2021. "A
Dietary Sterol Trade-off Determines Lifespan Responses to Dietary Restriction
in *Drosophila melanogaster* Females." *eLife* 10 (January).

Figure 1.

Consumption (mg/fly) = [Δ weight (mg) - mean evaporation (mg)] / number of flies

Figure 1. DIETS (Direct Intake Estimation and longitudinal Tracking of Solid

food consumption) is a sensitive assay that can measure food consumption in

flies over days.

(A) Schematic outline of the DIETS assay. **(B)** Comparison of 24 h control diet (CD)

consumption in males and females of increasing group sizes. Females consistently

eat a greater amount of food than males, regardless of the group size. Mean

consumption by both sexes increased with group size. **(C)** CD consumption by

groups of 50 male flies was measured over 24 h at different temperatures. An

increase in the amount of food consumed was observed with an increase in

temperature. **(D)** Feeding by groups of 50 male flies on varying concentrations of CD

was measured. Flies show compensatory feeding upon diluting or concentrating CD.

(E) Comparing the sensitivity of DIETS, CAFE, and EX-Q assays in resolving food

intake difference between 1X, 2X, and 4X concentrations of 2.5 % sucrose - 2.5 %

yeast extract (2.5 % SY) diet. DIETS show comparable resolution to CAFE and EX-

Q. **(F)** Longitudinal measurement of CD consumption over days 1, 4, 7, 14, 21, and

28 for 50 male flies. Consumption mostly remains unaltered, except for a dip in

feeding by the end of 28 days, possibly due to the mortality observed. Flies start

dying between the 7[th] and the 14[th] day. Scatter plots of raw data with mean ± 95 %

CI are shown. Δ 'effect size' plots below each raw data graph depict mean

differences (black dots; values labeled) between relevant groups or standardized

mean difference, hedge's g for Fig. 1E. The 95 % CI (black lines) and distribution of

the mean differences (curve), generated are also shown. See Materials and Methods

for further details.

Figure 2.

Figure 2. DIETS for short-term feeding measurements and food choice.

(A) Comparing the sensitivity of DIETS with blue dye uptake in measuring short-term

feeding. 30 minutes of feeding on CD supplemented with Brilliant blue FCF in

differentially starved male flies was measured using DIETS. Flies showed an

increase in feeding with increased starvation (left panel). A simultaneous dye

accumulation measurement was conducted by crushing the flies after feeding (right

panel). Hedge's g is effect size, in this case, the difference in mean of two groups,

standardized over the pooled standard deviation of the two groups. DIETS assay

yields larger 'g' values compared to blue dye estimation for comparison of feeding

between the same groups. **(B)** Continuous 2 h feeding measurements in sated male

and female flies measured over 2 days. Both sexes show a distinct evening feeding

peak just around the lights-off time at ZT = 11.5-12.5. **(C)** 24 h, food preferences

between CD and CD mixed with quinine. Flies show an increasing dose-dependent

aversion towards CD adulterated with quinine. **(D)** 2 h food preferences between CD

and CD adulterated with 15 mM quinine in 24 h starved flies. Flies showed a strong

aversion towards quinine-laced food. Scatter plots of raw data with mean ± 95 % CI

are shown. Δ 'effect size' plots below each raw data graph depict mean differences

(black dots; values labeled), between the two relevant groups being compared. The

95 % CI (black lines) and distribution of the mean differences (curve) generated by

bootstrap resampling of the data are also shown. In (A) standardized effect size,

Hedge's g has been plotted, and MOEs have been reported. See Materials and

Methods for further details.

Figure 3.

A

| 56 | 51 | 47 | 42 | 39 | 34 | (mg) avg. food offered (**a**) |
| **43** | 38 | 36 | 32 | 29 | 26 | (mg) avg. actual feeding (**b**) |

0	91	84	75	70	61	Targeted (**a**/56)*100
0	88	84	74	67	60	Achieved (**b**/43)*100

B

ALF = ad-libitum feeding
TRF = time restricted feeding

C

Figure 4. DIETS is well suited to study longitudinal feeding on high-fat diets

and associated transcriptional and physiological changes.

(A) In a longitudinal feeding study, flies were fed 5%, 10% and 20% saturated high-

fat diets made with coconut oil (sHFD), and feeding measurements were conducted

on days 1, 4, and 7. While consumption remained mostly constant up to 10 % extra

coconut oil in the diet, feeding decreased noticeably with the addition of 20 %

coconut oil. **(B)** Differential expression of *takeout* (*to*)*, Cytochrome P450-4e3*

(*Cyp4e3*), *Odorant-binding protein 83a* (*Obp83a*), and *Turandot A* (*TotA*) was

examined by qPCR from heads of flies that have been fed with 20 % sHFD for 1, 4,

and 7 days, with respect to control flies fed CD. These genes were earlier found to

be some of the most differentially expressed genes from an earlier study with a

similar design. The direction and magnitude of differential expression of the above

candidate genes on day 4 and 7 were comparable to that in day 7 samples from the

previous study (Stobdan et al. 2019), as shown. **(C)** Energy intake was determined

over 24 h periods on days 1, 4, and 7 in groups of flies fed 30 % HSD or isocaloric

10 % sHFD. The corresponding mean feeding values are represented above inside

pink boxes. Flies fed on 10 % sHFD showed increased energy intake compared to

the CD and HSD fed groups on all days. **(D)** The same flies were transferred to 0.75

% agar vials after 7 days of dietary treatment and the number of dead flies recorded

every 6 - 12 hours. Survival time (left panel), and percent survival under starvation

(right panel) of flies are shown. While HSD treatment increases starvation

resistance, sHFD treatment does not. Scatter plots of raw data with mean ± 95 % CI

are shown. For Figure 4C starvation resistance was plotted as Kaplan-Meier graph.

Δ 'effect size' plots below each raw data graph depict mean differences (black dots;

values labeled), between the two relevant groups being compared. The 95 % CI

Figure 4.

Figure 4. DIETS is well suited to study longitudinal feeding on high-fat diets

and associated transcriptional and physiological changes.

(A) In a longitudinal feeding study, flies were fed 5%, 10% and 20% saturated high-

fat diets made with coconut oil (sHFD), and feeding measurements were conducted

on days 1, 4, and 7. While consumption remained mostly constant up to 10 % extra

coconut oil in the diet, feeding decreased noticeably with the addition of 20 %

coconut oil. **(B)** Differential expression of *takeout* (*to*)*, Cytochrome P450-4e3*

(*Cyp4e3*), *Odorant-binding protein 83a* (*Obp83a*), and *Turandot A* (*TotA*) was

examined by qPCR from heads of flies that have been fed with 20 % sHFD for 1, 4,

and 7 days, with respect to control flies fed CD. These genes were earlier found to

be some of the most differentially expressed genes from an earlier study with a

similar design. The direction and magnitude of differential expression of the above

candidate genes on day 4 and 7 were comparable to that in day 7 samples from the

previous study (Stobdan et al. 2019), as shown. **(C)** Energy intake was determined

over 24 h periods on days 1, 4, and 7 in groups of flies fed 30 % HSD or isocaloric

10 % sHFD. The corresponding mean feeding values are represented above inside

pink boxes. Flies fed on 10 % sHFD showed increased energy intake compared to

the CD and HSD fed groups on all days. **(D)** The same flies were transferred to 0.75

% agar vials after 7 days of dietary treatment and the number of dead flies recorded

every 6 - 12 hours. Survival time (left panel), and percent survival under starvation

(right panel) of flies are shown. While HSD treatment increases starvation

resistance, sHFD treatment does not. Scatter plots of raw data with mean ± 95 % CI

are shown. For Figure 4C starvation resistance was plotted as Kaplan-Meier graph.

Δ 'effect size' plots below each raw data graph depict mean differences (black dots;

values labeled), between the two relevant groups being compared. The 95 % CI

(black lines) and distribution of the mean differences (curve) generated by bootstrap

resampling of the data are also shown. See Materials and Methods for further

details.