Title:

Trophic ecology of the African riverine elephant fishes (Mormyridae)

Authors:

Gina Maria Sommer[1,*], Samuel Didier Njom[2], Adrian Indermaur[3], Arnold Roger Bitja Nyom[2,4], Petra
Horká[5], Jaroslav Kukla[5], Zuzana Musilova[1,*]

Affiliations:

[1]Department of Zoology, Faculty of Science, Charles University, Viničná 7, 12844 Prague,
Czech Republic

[2]Department of Biological Sciences, University of Ngaoundéré, Ngaoundéré P.O. Box 454
Cameroon

[3]Zoological Institute, University of Basel, Vesalgasse 1, Basel, Switzerland

[4]Department of Management of Fisheries and Aquatic Ecosystems, University of Douala, Douala
P.O. Box 7236, Cameroon

15
16 [5]Institute for Environmental Studies, Faculty of Science, Charles University, Benátská 2, 12844
17 Prague, Czech Republic

18
19
20 * corresponding authors; gina.sommer44@googlemail.com, zuzana.musilova@natur.cuni.cz

Abstract:

Multiple species of the elephant fishes (Mormyridae) commonly coexist in sympatry in most African tropical rivers and lakes. In this study, we investigated the trophic ecology and potential trophic niche partitioning of eleven mormyrid fish species from the Sanaga River system (Cameroon) using the stable isotopes of carbon and nitrogen of muscles and of trophic prey samples. Albeit mormyrids mainly feed on invertebrates, we found differences in isotope signals and the trophic niche partitioning in the studied species. We further show that species with elongated snout tend to show higher carbon and nitrogen isotope signals, suggesting a potential role of snout shape in their trophic preferences. Furthermore, we found significant differences in isotopic signatures within the *Mormyrus* genus, highlighting ecological niche diversification among three closely related species. We also report on different isotopic signals between seasons of the year in four species, possibly caused by species migration and/or anthropogenic agricultural activities. Overall, our research presents robust evidence of the trophic niche partitioning within the entire mormyrid species community, shedding light on the enigmatic evolutionary history of these fascinating African fishes.

Keywords: mormyrids, stable isotopes, trophic ecology, trophic position, trophic resource portioning

1. Introduction

Elephant fishes, Mormyridae, exclusively occur in freshwater habitats of Africa, and they are well known for their additional electric sensory system. The produced electric pulses serve for electrolocation in turbid waters, as well as for conspecific communication (SULLIVAN et al., 2002; CARLSON et al., 2011) and evolution of species-specific signals is often associated with mormyrid diversification (currently 228 described species; FRICKE et al., 2023). Apart from their sensory systems, little is known about the general life history of mormyrids. Seasonal migrations have been reported in some species in several river systems (OKEDI, 1965; BLAKE, 1977, VAN DER WAAL, 1996), and seasonal changes are possibly impacting food availability and in some cases even shifts of diets and consequently trophic position of aquatic animals (e.g., NASIMA et al., 2020; MCMEANS et al., 2019). Studies on ecological aspects like movement patterns, reproductive behavior and feeding strategies are scarce in mormyrids revealing mostly complex behavioral traits such as pack hunting in *Mormyrops anguilloides* in Lake Malawi (ARNEGARD & CARLSON, 2005) or the sound production of *Pollimyrus isidori* mediating their courtship (CRAWFORD, JACOB & BENÉCH, 1997). Feulner et al. (2007) investigated diversification of the genus *Campylomormyrus* and found that shape of the snout protrusions may be associated with different trophic specializations, although this needs further confirmation. Studies on trophic ecology show that mormyrids are mainly invertivorous (WINEMILLER & ADITE, 1997; ADJIBADE et al., 2019; PEEL et al., 2019) or rarely piscivorous (*Mormyrops anguilloides*; BAILEY, 1994; SKELTON, 2001). Differences in trophic preferences have previously been described in co-existent species from lake Ayame, Ivory coast (KOUAMÉLAN et al. (2006) or Sokoto-Rima river basin, Nigeria (HYSLOP, 1986) based on the stomach content analyses showing that different species may specialize on different groups of insect larvae.

The method of stable isotope analysis is a powerful tool to illuminate trophic preferences. Heavier stable isotopes of carbon (^{13}C), nitrogen (^{15}N) or sulfur (commonly ^{34}S) can be detected by stable isotope mass spectrometry and accumulate differently in the organisms compared to the common isotope: the lighter isotopes form weaker bonds than the heavier ones as well as tend to react faster (JARDINE et al., 2003).

Reports based on stable isotope analyses in mormyrids are scarce (SOTO et al., 2019) finding just subtle

differences among species compared to the effect of sexual selection (ARNEGARD et al., 2010) while

being more commonly used in other African fishes (e.g. cichlids, MUSCHICK et al., 2012).

In this study, we focus on the stable isotope analysis in eleven species of mormyrid fishes from the

Sanaga River (Cameroon) to test for differences in trophic resources. We aim to provide the isotopic

analysis throughout the entire mormyrid species community and shed light on the Cameroonian

freshwater fish diversity in one of the less regulated rivers in Africa (until recently). We further focus on

species collected at different seasons of the year (i.e., the wet and the dry season and a transition phase)

and include fish samples from multiple localities on the main Sanaga River around the Nachtigal falls, as

well as in some nearby tributaries.

2. Material and Methods

2.1. Sampling

The study site for this work was the Sanaga River in Cameroon, the country's largest river of over 918

kilometers long (DUBREUIL et al., 1975). There are three large contiguous segments, Upper Sanaga,

Middle Sanaga and Lower Sanaga (BITJA NYOM et al., 2020) and this study took place in the middle part

of the Sanaga River. Sampling localities were located along the main stream in the proximity of the

cascades of Nachtigal complemented by few inflow stream locations (tributaries) (see Map in Fig. 1).

There exist different seasons in this Guinean equatorial climate zone (OLIVRY, 1986; AMOUGOU et al.,

2015), namely a main wet season (from August until November) and a main dry season (from December

to March) plus a transition between the dry and the wet season (from April until July), sometimes divided

as a shorter wet season lasting from April until June and a shorter dry season (from July until early

August) (BITJA NYOM & PARISELLE, 2015).The fish were caught using gillnets of different mesh

sizes. After euthanizing the fish with a lethal amount of MS-222, photographs and measurements were

taken, and fin clips collected for DNA analyses. Afterwards, a 1 to 2 cm piece of lateral muscle was

dissected from an area between the ventral and the caudal fin in proximity to the lateral line. Eleven

species (*Campylomormyrus phantasticus* (PELLEGRIN, 1927), *Hippopotamyrus castor* PAPPENHEIM,

1906, *Marcusenius mento* (BOULENGER, 1890), *Marcusenius sanagaensis* BODEN, TEUGELS & HOPKINS,

1997, *Mormyrops anguilloides* (LINNAEUS, 1758), *Mormyrops caballus* PELLEGRIN, 1927, *Mormyrus*

tapirus PAPPENHEIM, 1905, *Mormyrus* sp. "short snout", *Mormyrus* sp. "long snout", *Paramormyrops*

batesii (BOULENGER, 1906) and *Petrocephalus christyi* BOULENGER, 1920) were sampled in 2017 and

2018. The number of individuals per species vary and the representation of each season is not always

complete and at times unequally distributed (Table 1). All fishes were collected with appropriate research

permits   namely   N°   007,   048,   049/MINRESI/B00/C00/C10/C12   and   N°

2376/PRBS/MINFOF/SETAT/SG/DAFP/SDVEF/SC/ENJ granted by the Ministry of Scientific Research

and Innovation and the Ministry of Forestry and Wildlife in Cameroon.

In total we collected samples from 288 individuals of 11 species. All samples were transfered in 96%

ethanol and stored in a freezer at -20 degrees. We further complemented our muscle sampling by adding

trophic samples consisting of the individual items from the stomachs of nine individuals (two *M.*

caballus, one *M. anguilloides*, two *H. castor*, two *M. tapirus* and one *M.* sp. "short snout"), as well as of

benthic samples, i.e. plant material and invertebrates collected from the river bed area, the riverbank and

the water column of the Sanaga River and a tributary, namely from Nachtigal falls and the Wala river.

These samples target potential prey and a serve as a reference isotope background. The organisms from

these environmental samples were identified taxonomically under a binocular microscope to the highest

level possible, from sometimes only class (Gastropoda), mostly the order (e.g. Ephemeroptera) and

seldom to a family level (e.g. Hydrophilidae, Elmidae) with the help of pictures taken from stomach

content material and invertebrate identification guides (MOISAN J. *et al.*, 2010; MARY, N., 2017). The

water plant material was not further identified, and the various plants were mixed to achieve an amount of

over 2 mg and labelled accordingly as the group called "plant". The isotope values generated by both, the

stomach content material and the benthic material, are jointly represented when later referring to the

values of the "trophic samples", unless otherwise specified.

2.2. Sample processing

The muscle tissue samples were transferred to 2ml Eppendorf tubes and were dried for 48 hours at 60 °C

or at 45 °C for 72 hours over the weekend in a drying oven. If bone structures were present in the sample,

they were carefully removed with a clean forceps. Then samples were ground with two beads at the

frequency 25/s for 20 seconds or at 30/s for 30 seconds in case of a bigger piece using a ball mill

(MM400, Retsch, Germany). After taking the beads out, one millilitre of ethanol was added, and sample

has been shaken until homogenous. The samples were then centrifuged for 30 seconds at 6000 rpm.

About 600 microliters of ethanol were taken out with a pipette and a second drying step was followed at

24 up to 48 hours for 60 °C. In case the sample was not completely transformed into the powder yet, an

additional ball grounding was performed at 25/s frequency for 10 seconds and then centrifuged for 30

seconds at 6000 rpm and final drying for 15 minutes and 60 °C. Subsequently, the samples were weighted

in tin capsules to achieve 0.5 milligrams ideally while allowing a range of 0.45 to 0.55 mg and folded into

tiny cubes using a thin and thick tweezer and put in labelled sample boxes. The trophic samples were

processed in the same manner except grounding happened partly manually and sometimes with only one

silver ball. Plant material had to achieve a weight of 3 mg. Some trophic samples were too light and had

to be excluded from further analysis (17 samples out of 64). Total carbon and nitrogen amounts as well as

the isotopic ratios $\delta^{13}C$ and $\delta^{15}N$ were measured with a Delta V Advantage mass spectrometer coupled to

Conflo IV and elemental analyzer Flash 2000 using a Thermal Conductivity detector (all instrumentation

by Thermo Fisher Scientific, Bremen, Germany). The carbon and nitrogen isotope ratios are expressed

following the delta notation, which is the following: $\delta X = 1000*(R_{sample}/R_{standard} - 1)$, where X stands

for ^{13}C or ^{15}N, respectively, and R is the relative amount of carbon or nitrogen isotopes ($R = {}^{13}C/{}^{12}C$ or

$^{15}N/^{14}N$). International standards were applied to normalise the isotope ratios: The Vienna Pee Dee

Belemnite (VPDB) scale (COPLEN, 1996) and atmospheric N2 was used for the carbon and the nitrogen

isotope ratios, respectively. In addition to repeated measurements of a series of international standards

(IAEA-CH3, IAEA-CH6, IAEA-600, IAEA-N1, IAEA-N2, IAEA-NO3), a glycine standard was included

after every 10th sample to provide calibration for elemental composition as well as serve as a quality

control for isotopic measurement. Analytical precision resulted within $\pm$ 0.2 ‰ for both $\delta^{13}C$ and $\delta^{15}N$.

2.3. Data analysis

Delta 13C ($\delta^{13}C$) and delta 15N ($\delta^{15}N$) values were analysed in R Studio Statistical Software (version

1.4.1717; RStudio Team (2021). RStudio: Integrated Development Environment for R. RStudio, PBC,

Boston, MA URL http://www.rstudio.com/.). The package "tRophicPosition" (QUEZADA-ROMEGIALLI C.

et al., 2018) was used furthermore to access the trophic position of the investigated species using one

baseline of all benthic sourced samples from the trophic samples. The used trophic discrimination factors

(TDF) were McCutchan (MCCUTCHAN'S *et al.* 2003) for a fish muscle tissue which means a mean of 2.9

‰ $\pm$ 0.32 se for $\delta^{15}N$ and 1.3 ‰ $\pm$ 0.3 se of for $\delta^{13}C$. Tropic position (TP) was estimated using a λ

(lambda) = 2 for using also secondary consumers as baseline as no primary producer was available. The

conventional range of trophic levels ranges from 1 (primary producers) to 5 (2 to 4.5 for fishes in

freshwater ecosystems) (TRITES, 2001; YOĞURTÇUOĞLU et al., 2020). The plant material collected were

several species but had to be pooled together to increase weight and thus has only informative value. The

model was run with 4 chains and for 20000 iterations. To access the significance of the differences

observed in muscle isotopic signals mean values between different seasons one-way ANOVAs as well as

Tukey tests to judge the results of the ANOVA were conducted. Alpha was set to 0.01 and the critical

value q was found accordingly in available tables online (see Table 3).

3. Results

3.1. Muscle samples analysis

In total, we analysed 288 samples of 11 species of the mormyrid community in the Sanaga River, and we

found that the stable isotope signals partly differ among the species (Fig. 2). The *Marcusenius* species and

Paramormyrops batesii share lower δ^{13}C and δ^{15}N values in comparison to all other species. Interestingly,

some species from different genera possess similar isotopic signatures like *Hippopotamyrus castor* and

Mormyrus tapirus, or *Mormyrops caballus* and *Campylomormyrus phantasticus* whereas the species from

the same genus are more separated from each other. This is especially obvious in the *Mormyrus* genus

(Figs 2-3).

3.2. Trophic samples

As a baseline, we analyzed trophic samples of possible prey items, i.e. selected items from the fish

stomachs as well as benthic samples. We analysed the stable isotope values of 16 potential prey groups.

Some individual samples had to be pooled together in order to reach the threshold weight needed for the

analysis, like the group Plecoptera, Trichoptera and Ephemeroptera. The δ^{15}N values range from 4.9 to

5.0 parts per thousand (‰) and the δ^{13}C values from -17,5 to -28 parts per thousand. Gastropoda has the

highest δ^{13}C values and Coleoptera the lowest. The plant material, Zygoptera and Hemiptera display the

highest δ^{15}N values whereas Plecoptera and Elmidae (Coleoptera) the lowest. Only three groups

(Anisoptera, Gastropoda and Elmidae (Coleoptera)) were simultaneously present in the stomach samples

and the benthic items. The isotopic signatures did only marginally differ, mainly in Elmidae (Figure S1

for the benthic items and Figure S2 for the stomach samples). We noticed expected pattern of higher

trophic position (δ^{15}N) in mormyrid fish muscles compared to the prey items (Fig. 3).

3.3. Calculations of trophic position

The analysis performed with the tRophicPosition R package (QUEZADA-ROMEGIALLI C. et al., 2018)

with a benthic prey baseline revealed the trophic positions of all mormyrid species between 3 and 4.

Pairwise comparisons within genera revealed only marginal differences in the trophic position of two

species of *Mormyrops* (mean TP of 3.8 vs. 3.7) (Fig. 4A) and *Marcusenius* (3.612 vs. 3.545) (Fig. 4B).

On the other hand, we found differences among the three *Mormyrus* species. Namely, *Mormyrus* sp.

"short snout" shows a notable difference (3.912) from *Mormyrus* sp. "long snout" (3.605) and from

Mormyrus tapirus (3.772; Fig. 4C,D,E). Accordingly, statistical analysis (t-tests) revealed significant

differences of the stable isotope values only in the *Mormyrus* species (see Table 2 below). This is in

accordance with the nearly congruent trophic positions of the *Mormyrops* and the *Marcusenius* species

but also reflects the separation of *Mormyrus* sp. "short snout" from *Mormyrus tapirus* and from

Mormyrus sp. "long snout" nicely.

3.4. The impact of the season

We analyzed samples from different seasons, namely during the dry season (February and March), the

wet season (August, September, October and November) and the transition phase between these two

seasons (in the months of May, June and July). We investigated the influence of the season on the stable

isotope values and found no difference when comparing dry and wet season values. We did find

differences though between the samples collected during the transition phase and the two main seasons.

Albeit the low number of samples from the transition phase (only for five species; Fig. 5), four species

show significant differences between the wet vs. transition and/or dry vs. transition phase (Table 2). The

one-way ANOVAs and the subsequent Tukey-Kramer tests revealed that in two species, *Mormyrus*

tapirus and *Marcusenius sanagaensis*, both stable isotopes δ^{15}N and δ^{13}C were significantly different

between the seasons (pairwise comparisons of wet-vs.-transition and dry-vs.-transition). In

Campylomormyrus phantasticus a difference was found only in the δ^{13}C signal for the dry-vs.-transition

samples, while in *Petrocephalus christyi* the δ^{15}N values are different, although the statistical analysis

could not be performed for all pairs (only one wet-season sample was available). Only in *Hippopotamyrus*

castor no differences in the stable isotope signal was noticed. (Table 3).

4. Discussion and Conclusions

Trophic niche partitioning in mormyrid fishes:

We provide the first comprehensive overview of the mormyrid stable isotope signals at the community level. Our trophic ecology results are in accord with previous evidence of mormyrids feeding mostly on insect larvae (VAN DER WAAL, 1985; WINEMILLER & ADITE, 1997). There are, however, noticeable differences among species, such as the two *Marcusenius* species and *P. batesii* having a lower trophic position than the other mormyrids. We also observe that the mean isotopic signal of different species within one genus often differs more than when compared across other genera. This is the case for *Mormyrops caballus* and *Mormyrops anguilloides* with distinct signals in both δ^{13}C and δ^{15}N and more strikingly among the three species of the *Mormyrus* genus. *Mormyrus tapirus*, and two formally undescribed species, *M.* sp. "long snout", and *M.* sp. "short snout" differ in δ^{13}C and δ^{15}N signals, and, accordingly, *Mormyrus* sp. "short snout" also differs from the other two other *Mormyrus* species in the trophic position analysis (Fig. 4). Our findings therefore suggest different trophic specializations for these closely related species. Trophic niche partitioning among closely related species from the same habitat has been commonly observed in African fishes (e.g. NAGELKERKE, 1997; POLÁČIK et al., 2014), while resource partitioning has been a general question in ecology since a long time (HARDIN, 1960; SHOENER, 1974). Stable isotope data have already proven to be helpful to visualize trophic niche partitioning within closely related fish species (WANG et. al, 2022) and our study is the first case of stable isotope analysis showing trophic niche partitioning in mormyrids. Differences in trophic preferences can be further observed on the jaw/mouth shape, which is often adapted to its specific purpose. For example, in croakers (Sciaenidae; WANG et al., 2022) the species with the most unique isotopic signatures also has a unique feeding strategy associated with different jaw shape (ZHANG et al., 2020). Based on our results, mormyrid species from Sanaga with snout protrusion tend to show higher δ^{13}C and/or δ^{15}N signals when compared to species without elongated snouts, or, from another angle, species with higher δ^{15}N values (~ trophic level) show higher variability in the snout elongation and shapes (Fig. 2D). Notably, the three *Mormyrus*

species have different isotopic signals and at the same time express anatomical differences in their trunk

shape and jaw and lip morphology. *Mormyrus* sp. "short snout" has a more truncated upper lip protrusion,

while *M. tapirus* has a longer lower lip, and *M.* sp. "long snout" has much longer snout than the two other

Mormyrus species (Fig. 2). The overall mouth shape and orientation probably reflect subtle differences in

microhabitat and prey preferences in these species. Our data therefore illustrates the differential trophic

specialization associated with the snout shape differences even on the shorter evolutionary scale in the

three closely related species. This seems to be in accord with the hypothesis of Feulner et al. (2007) who

suggested possible variations in trophic ecology based on morphological differences in the snout

protrusions in the *Campylomormyrus* genus from the lower Congo River (FEULNER et al., 2007).

The trophic position analysis is mostly based on the δ^{15}N values since it has been shown that trophic

position is corresponding more to the trophic fractionation of δ^{15}N than to δ^{13}C (POST, 2002). The

environmental benthic samples were set as a baseline for the analysis and these samples represent

possible sources of carbon and nitrogen from the environment. We also tested the trophic position with

the stomach samples with very similar results (see Supplementary material S3). Ideally, a primary

producer is used as a baseline for such analysis, unfortunately it was not available in our study as the

collected plant material was insufficient. Some authors have also claimed that macrophytes, the main

primary producers in many aquatic ecosystems, were revealed to contribute less to aquatic food webs than

assumed (HAMILTON et al., 1992; BUNN & BOON, 1993; FORSBERG et al., 1993; THORP & DELONG,

1994; THORP et al., 1998; LEWIS et al., 2001) or may have only local effects (GROSBOIS et al., 2017a)

(PEEL et al., 2019). Working on an understudied group of tropical fishes, such as mormyrids, may be

challenging as other factors contribute to the final patterns. In our case, seasonal changes (possible

migration), differing turnover rates, or unknown discrimination factors for Bayesian analysis. Turnover

rates are generally tissue-dependent and are slower in muscle tissue or bone collagen (two to three

months) than in plasma or blood (few days) (e.g., MADIGAN et al., 2021, POLISCHUK et al., 2001) but

there are hints that the turnover rates can also differ among species, fish size and life stage (BOSLEY et al.,

270 2002; GASTON et al., 2004; MADIGAN et al., 2012 and MADIGAN et al., 2021). For discrimination factors

271 (i.e., the exact isotopic shift between prey and consumer), we used McCutchan's (MCCUTCHAN et al.

272 2003) values for fish muscles with a mean of 2.9 ‰ ± 0.32 se for $\delta^{15}N$ and 1.3 ‰ ± 0.3 se of for $\delta^{13}C$,

273 although values of the trophic shifts differ especially for N depending on diet. Interestingly, McCutchan

274 et al. (2003) found lower ^{15}N trophic shift mean values (around 1.4‰) for consumers feeding exclusively

275 on invertebrates in contrast to consumers of high-protein diets. In our study, we found the trophic

276 discrepancy for $\delta^{15}N$ at a minimum of 2.5‰ between mormyrid consumers and trophic samples (Fig. 3)

277 and this could speak for a more high-protein diet in some mormyrid species. In order to reveal the trophic

278 resources and more details on mormyrid prey, we suggest that our results could be complemented by a

279 long-term stomach content analyses in the future

280 as the two methods can complement each other (DAVIS et al., 2012; LAYMAN et al., 2012; POLÁČIK et al.,

281 2014; DILLON et al., 2021).

283 Effect of seasonality

284 Dry and wet seasons in tropical regions can be reflected in fish behaviour, migration, reproduction or

285 food/prey availability. Impact of the season on the stable isotope values has been reported from multiple

286 fish groups (NASHIMA et al., 2020; MCMEANS et al., 2019; TAYLOR, 2016 [doctoral thesis]; WANTZEN et

287 al., 2002) and seasonal variability seems to be especially relevant for partly flooded aquatic systems

288 (TAYLOR et al., 2017; MERRON & MANN, 1995; DE ALMEIDA et al. 1997; WANTZEN et al. 2002;

289 EGGLETON & SCHRAMM, 2004). The isotopic signal can be influenced by various factors, not only by the

290 actual changes in the prey items. Some studies reported on fish to become more generalized feeders

291 during the wet season (COSTA-PEREIRA et al., 2017; POOL et al. 2017) and there are hints for increasing

292 trophic position during the dry season (WANTZEN et al. 2002), whereas other studies showed diverse

293 foraging strategies with no universal seasonal influence pattern (NOVAKOWSKI et al. 2008), (MCMEANS

294 et al., 2019). In this study, we have specifically focused on the stable isotope signatures in different

seasons throughout the year (Table 1 and S1). We found differences in isotopic signal between the transition phase (samples from May-July) compared to the wet season (August-November) or the dry season (February-March), whereas no difference was found between dry and wet season. The different isotope signal is present in four out of five tested species, i.e., in *Mormyrus tapirus*, *Marcusenius sanagaensis, Petrocephalus christyi* and *Campylomormyrus phantasticus,* while *Hippopotamyrus castor* did not show any differences throughout seasons. Therefore, the transition period seems to influence the diet of some, but not all mormyrid species. The habitats of the Sanaga River and tributaries are characterized by seasonal flooding during the wet season and are exposed to a rising water level already in the transition phase. Multiple factors could contribute to the seasonality differences. First, species seasonal migration could cause that individuals sampled in the transition phase may originate from different population than the dry- and wet-season samples. In general, mormyrids are known to migrate (VAN DER WAAL, 1996; OLOPADE, 2013), mostly in relation with the wet seasons spawning (OKEDI, 1965; BLAKE; 1977) but little is known for the tested species from the Sanaga River. At least one of the species, *Marcusenius sanagaensis,* is known to be more residing in the tributaries than in the main stream (personal observations on the study by Njom and others during several years of fieldwork). An alternative explanation would be in the turnover rates. If turnover rates are a few months for white muscle tissue like in most other more intensively studied fish species (TAYLOR et al., 2017), it could mean that a change of diet at the very beginning of the dry season and during the dry season would affect the transition phase values. Interestingly, this could be an anthropogenic induced cause due to the phytosanitary treatment of the plantations along the tributaries carried out towards the end of the major dry season (NJOM et al., 2022). Such unusually large terrestrial input to the water may possibly impact the signal of some (but not all) species. Lastly, we can also explain the pattern partially by different sampling locations in the transition period for some species (Table S1). For example, the samples of *Marcusenius sanagaensis* from the Mekono River (a tributary with a pollution impact largely increased during the dry season) are different from the main Sanaga River and this seems to influence the results more than the season in this species (revealed by ANOVAs shown in Table S3). In *Mormyrus tapirus* both season and locality seem to

both play a role. The difference in δ^{13}C values (but not in δ^{15}N) remains after exclusion of the most

different samples are from Mpem River (as seen in the Supplementary table S3). Interestingly, the

transition samples from the same locality (Mpem River) are not significantly different in *Marcusenius*

sanagaensis, which illustrates again differences among species. Some differences of the transition

samples, e.g. δ^{13}C values of *Campylomormyrus phantasticus* (all samples are from one locality) or the

δ^{15}N values of *Petrocephalus christyi* remain irrespective of locality issue. Unfortunately, our sampling is

not robust enough to test properly the effects of all aforementioned factors possibly involved in the

seasonally different signal of the stable isotopes in mormyrids.

To conclude, here we present the first comprehensive study applying stable isotope analysis on mormyrid

fishes at a species community level. We found differences in the isotopic signal among eleven species

corresponding to putative feeding specialization and we also report on putative role of the snout shape in

the trophic ecology of different species. We have specifically focused on the difference between three

species of the *Mormyrus* genus with the clear trophic niche differentiation after the recent evolution of

these species. We further report on seasonal differences in the isotopic signal in four species. Our study

attempts to shed light on the biology of the evolutionarily successful and diverse - and yet understudied –

group of tropical freshwater fishes.

Supporting Information:

Figure S1: Stable isotope analysis of the trophic samples from the benthic samples

Figure S2: Stable isotope analysis of the trophic samples from the stomach contents

Figure S3: Trophic positions comparisons based on stomach content baseline

Table S1: sample list overview of muscle samples isotope values and details about sampling

Table S2: sample list overview of trophic samples

Table S3: results of single-factor ANOVAs and Tukey-Kramer test (if applicable) for species with transition season samples testing the factor of locality

Funding:

The project has been funded by the Swiss National Science Foundation (PROMYS – 166550) and the PRIMUS Research Programme (Charles University). ZM was further supported by the Czech Science Foundation (21-31712S). GMS was supported by GAUK (251299). PH was further supported by a grant of the Czech Health Research Council (AZV CR, number NU21-04-00405).

Acknowledgements

We deeply thank Monika Kłodawska, Dima Omelchenko and Hassan Bassirou for the field assistance as well as their valuable support as colleagues. We are very thankful to the local fisherman Raphael Abanda without whom no successful fishing would have been possible and whose knowledge of the area was indispensable. We further thank our field guides Jean-Claude Remix and Gregoire Kayoum from the "Institut de Recherche pour le Développement (IRD)", as well as our hosts in the Batchenga village. We sincerely express gratitude to the local people of the Batchenga community to allow us to fish in their territory, the Nachtigal HydroPower Company (NHPC) for the support to sample in the Sanaga and the Mpem and Djim National Park rivers. We thank all the students and people from the Institute for Environmental Studies and the Department of Botany of Charles University for sharing space and giving us the possibility to work there and offering kind advice and friendliness. A special thanks goes to Marc Schmid for providing the map in Figure 1.

Conflict of Interest Statement:

The authors declare no conflict of interests.

Author contribution statement:

Conceptualisation: ZM, ARBN. Developing methods: ZM, ARBN, PH, JK Conducting the research: ZM,

AI, SDN, PH, GMS. Data analysis: PH, JK, GMS. Data interpretation: ZM, SDN, GMS. Preparation

figures and tables: GMS. Writing – initial draft - GMS and ZM, Writing – final revision and approval –

all authors.

Data Availability statement:

COI sequences used for the species phylogeny are available on GenBank for *Marcusenius mento* with

accession numbers: HM882757 and HM882727, and (will be available) for the other species with the

accession numbers (insert). Raw data of stable isotope analysis of the mass spectrometer for muscle

samples are imbedded in sample overview table in Table S1 in Supporting information and for trophic

samples in Table S2.

5. References:

Adjibade K. N., Adite A., Arame H., Imorou R. S. & Sonon S. P. (2019) Feeding ecology of *Marcusenius senegalensis* (Pisces: Osteoglossiformes: Mormyridae; Steindachner, 1870) from Niger River in Northern Benin. International Journal of Fisheries and Aquatic Studies, 7(4), 86-92.

Amougou, J. A., Nndam Ngoupayou, J. R., Djocgoue, P. F. & Bessoh Bell, S. (2015) «Variabilité climatique et régime hydrologique dans un milieu bioclimatique de transition: cas du bassin fluvial de la Sanaga». Afrique Science, 11(5). http://www.afriquescience.info/document.php?id=5175. ISSN 1813-548X.

Arnegard, M. E., & Carlson, B. A. (2005) Electric organ discharge patterns during group hunting by a mormyrid fish. Proceedings. Biological sciences, 272(1570), 1305–1314. https://doi.org/10.1098/rspb.2005.3101

Arnegard, M. E.,McIntyre, P. B., Harmon L. J., Zelditch M. L., Crampton, W. G. R., Davis, J. K., Sullivan, J. P., Lavoue, S. & Hopkins, C. D. (2010) Sexual Signal Evolution Outpaces Ecological Divergence during Electric Fish Species Radiation. The American Naturalist, 176(3), 335–356.

Bailey, R.G., (1994) Guide to the fishes of the River Nile in the Republic of the Sudan. J. Nat. Hist. 28, 937-970.

Bitja Nyom, A.R & Pariselle A. Etat Initial des Peuplements Piscicoles. Rapport Final (2015) Etat Initial

des   Peuplements   Piscicoles.   Rapport   Final   http://documents.worldbank.

org/curated/pt/930081498550117079/pdf/116.

Blake B. F. (1977) Aspects of the reproductive biology of *Hippopotamyrus pictus* from Lake Kainji West

Africa with notes on four other mormyrid species, Fish Biology, 11, 437-445.

Boden, G., G. G. Teugels & Hopkins, C. D. (1997) A systematic revision of the large-scaled Marcusenius

with description of a new species from Cameroon (Teleostei; Osteoglossomorpha; Mormyridae). Journal

of Natural History, 31 (11), 1645-1682.

Bosley, K. L., Witting, D. A., Chambers, R. C. & Wainright, S. C. Estimating turnover rates of carbon

and nitrogen in recently metamorphosed winter flounder *Pseudopleuronectes americanus* with stable

isotopes (2002) Mar. Ecol. Prog. Ser. 236, 233–240.

Boulenger, G. A. (1890) Description of a new species of Mormyrus. Annals and Magazine of Natural

History, 6 (32), (art. 25), 193.

Boulenger, G. A. (1906) Description of a new mormyrid fish from South Cameroon. Ann. Mag. nat. Hist.,

7, 18, 103, 36-37.

Boulenger, G. A.   (1920) Poissons recueillis au Congo Belge par l'expédition du Dr C. Christy.

(Matériaux pour la Faune du Congo.). Annales du Musée du Congo Belge Sér. C (Zoologie) (Sér. 1) v. 2

(fasc. 4): 1-38, Pls. 23-25.

Bunn, S. E. & BoonP. I. (1993) What sources of organic carbon drive food webs in billabongs? A study

based on stable isotope analysis. Oecologia, 96, 85–94.

Carlson, B. A., Hasan, S. M., Hollmann, M., Miller, D. B., Harmon, L- J. & Arnegard, M. E. (2011) Brain

evolution triggers increased diversification of electric fishes. Science, 332(6029), 583-586. DOI:

10.1126/science.1201524

Coplen, T.B. (1996) New guidelines for reporting stable hydrogen, carbon, and oxygen isotope-ratio data.

Geochim. Cosmochim. Acta, 60, 3359-3360.

Costa-Pereira, R., L. E. R. Tavares, P. B. de Camargo, and M. Araújo (2017) Seasonal population and

individual niche dynamics in a tetra fish in the Pantanal Wetlands. Biotropica, 49, 531–538.

Crawford, J. D., Philippe Jacob, & Vincent Bénech. (1997) Sound Production and Reproductive Ecology

of Strongly Acoustic Fish in Africa: Pollimyrus isidori, Mormyridae. Behaviour, 134(9/10), 677–725.

http://www.jstor.org/stable/4535465

Davis A. M., Blanchette M. L., Pusey, B. J., Jardines, T. D. & Pearson R. G. (2012) Gut content and

stable isotope analyses provide complementary understanding of ontogenetic dietary shifts and trophic

relationships among fishes in a tropical river. Freshwater Biology, 57, 2156–2172. doi:10.1111/j.1365-

2427.2012.02858.x

de Almeida, V. L. L., Hahn, N. S., & de Vazzoler, A. E. A. M. (1997). Feeding patterns in five predatory

fishes of the high Parana River floodplain (PR, Brazil). Ecology of Freshwater Fish, 6, 123–133.

Dillon, K.S., Fleming, C.R., Slife, C. & Leaf, R.T. (2021), Stable Isotopic Niche Variability and Overlap

across Four Fish Guilds in the North-Central Gulf of Mexico. Mar Coast Fish, 13, 213-227.

https://doi.org/10.1002/mcf2.10148

Dubreuil, P., Guiscafre, J., Nouvelot, J.-F., Olivry, J.-C. (1975) Le bassin de la rivière Sanaga, ORSTOM,

Paris, 431.

Eggleton M. A. & Schramm H. L. (2004) Feeding ecology and energetic relationships with habitat of blue

catfish, Ictalurus furcatus, and flathead catfish, *Pylodictis olivaris*, in the Lower Mississippi River, U.S.A.

Environ Biol Fishes 70, 107-121.

Feulner P. G. D., Kirschbaum F., Tiedemann R. (2008) Adaptive radiation in the Congo River: An

ecological speciation scenario for African weakly electric fish (Teleostei; Mormyridae;

Campylomormyrus). Journal of Physiology, 102 (4–6), 340-346.

https://doi.org/10.1016/j.jphysparis.2008.10.002.

Forsberg, B. R., Araujo-Lima, C. A. R. M., Martinelli, L. A., Victoria, R. L. & Bonassi, J. A. (1993).

Autotrophic carbon sources for fish of the central Amazon. Ecology 74, 643–652. doi: 10.2307/1940793

Fricke, R., Eschmeyer, W. N. & Van der Laan, R. (eds) 2023. ESCHMEYER'S CATALOG OF FISHES:

GENERA, SPECIES, REFERENCES.

(http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp). Electronic version

accessed 31.01.2023.

Gaston, T. F. & Suthers, I. M. Spatial variation in $\delta 13C$ and $\delta 15N$ of liver, muscle and bone in a rocky

reef planktivorous fish: the relative contribution of sewage (2004) J. Exp. Mar. Biol. Ecol. 304, 17–33.

Grosbois, G., Del Giorgio, P. A., and Rautio, M. (2017a). Zooplankton allochthony is spatially

heterogeneous in a boreal lake. Freshwater Biology. 62, 474–490. doi: 10.1111/fwb.12879

Hamilton, S. K., Lewis, W. M., and Sippel, S. J. (1992). Energy sources for aquatic animals in the

Orinoco River floodplain: evidence from stable isotopes. Oecologia, 89, 324–330. doi:

10.1007/BF00317409

Hardin, G. (1960) The Competitive Exclusion Principle. Science, 131 (3409), 1292–1297.

Hyslop, E. J. (1986). The food habits of four small‐sized species of Mormyridae from the floodplain

pools of the Sokoto‐Rima river basin, Nigeria. Journal of fish biology, 28(2), 147-151.

Jardine, T. D., McGeachy, S. A., Paton, C. M., Savoie, M. & Cunjak, R. A. (2003) Stable isotopes in

aquatic systems: sample preparation, analysis, and interpretation. Can. Manuscr. Rep. Fish. Aquat. Sci.,

2656, 39.

Kouamélan E. P., Koné T., N'Douba V. & Ollevier F. (2006) Food habits and trophic resource

partitioning among three mormyrid fishes from man-made Lake Ayame, Ivory Coast, African Zoology,

41(2), 266-274, DOI: 10.1080/15627020.2006.11407362

Layman, C. A., Araujo, M. S., Boucek, R. , Hammerschlag-Peyer, C. M., Harrison, E. , Jud, Z. R. ,

Matich, P. , Rosenblatt, A. E. , Vaudo, J. J., Yeager, L. A. , Post, D. M. & Bearhop, S. (2012) Applying

stable isotopes to examine food-web structure: an overview of analytical tools. Biol Rev Camb Philos

Soc. 87(3), 545-62. doi: 10.1111/j.1469-185X.2011.00208.x. Epub 2011 Nov 2. PMID: 22051097.

Lewis, W. M. J., Hamilton, S. K., Rodríguez, M. A., Saunders, J. F. & Lasi, M. A. (2001). Foodweb

analysis of the Orinoco Floodplain based on production estimates and stable isotope data. J. North Am.

Benthol. Soc. 20, 241. doi: 10.2307/1468319

Linnaeus, C. (1758) Systema Naturae, Ed. X. (Systema naturae per regna tria naturae, secundum classes,

ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Tomus I. Editio decima,

reformata.) Holmiae. v. 1: i-ii + 1-824 [Nantes and Pisces in Tom. 1, 230-338.

Madigan, D. J. et al. Tissue turnover rates and isotopic trophic discrimination factors in the endothermic

teleost,   Pacific   bluefin   tuna   (*Thunnus   orientalis*)   (2012)   PLoS   ONE,   7,   e49220.

https://doi.org/10.1371/journ al.pone.00492 20

Madigan D. J., Snodgrass O. E., Hyde J. R. & Dewar H. (2021) Stable isotope turnover rates and

fractionation in captive California yellowtail (*Seriola dorsalis*): insights for application to field studies.

Scientific Reports, 11(4466). https://doi.org/10.1038/s41598-021-83880-z

Mary, N. (2017) Les macro-invertébrés des cours d'eau de Nouvelle-Calédonie. ETHYCO (version

révisée),

Moisan, J. et al. (2010) Guide d'identification des principaux macroinvertébrés benthiques d'eau douce

du Québec, 2010 — Surveillance volontaire des cours d'eau peu profonds, Direction du suivi de l'état de

l'environnement, ministère du Développement durable, de l'Environnement et des Parcs, ISBN: 978-2-

550-58416-2 (version imprimée) 82 p. (incluant 1 ann.)

McCutchan Jr J. H., Lewis Jr W. M., Kendall C. & McGrath C. C. (2003) Variation in trophic shift for

stable isotope ratios of carbon, nitrogen, and sulfur. OIKOS, 102(2), 378–390.

McMeans B. C., T. Kadoya, T. K. Pool, G.W. Holtgrieve, S. Lek, H. Kong, K.Winemiller, V. Elliot, N.

Rooney, P. Laffaille & K. S. McCann (2019) Consumer trophic positions respond variably to seasonally

fluctuating environments. Ecology 100(2):e02570. 10.1002/ecy.2570

Merron, G. S., & Mann, B. Q. (1995). The reproductive and feeding biology of Schilbe intermedius

Ruppell in the Okavango Delta, Botswana. Hydrobiologia, 308, 121–129.

Muschick, M., Indermaur, A., & Salzburger, W. (2012). Convergent evolution within an adaptive

radiation of cichlid fishes. Current biology, 22(24), 2362-2368.

Nagelkerke, L. (1997). The barbs of Lake Tana, Ethiopia: morphological diversity and its implications for

taxonomy, trophic resource partitioning, and fisheries. PhD thesis, Wageningen University and Research.

Nashima, F. P., Strydomb, N. A., Connan, M. & Lamberth, S. J. (2020) Stable isotopes reveal trophic

linkages among fish species utilizing the Orange River Estuary Continuum. FoodWebs, 24, e00145.

Njom, S.D., Bitja Nyom, A.R., Hassan, B., Bissek, J.P., Bêche, L., Pariselle, A., Bilong Bilong, C.F.

(2022) Inventory of the Ichthyofauna of the Mpem and Djim National Park (Center, Cameroon) Provides

Baseline Data for a Conservation Project. Diversity, 14, 1029. https://doi.org/10.3390/d14121029

Novakowski, G. C., Hahn N. S. & Fugi R. (2008) Diet seasonality and food overlap of the fish

assemblage in a Pantanal pond. Neotropical Ichthyology, 6, 567–576.

Okedi, J. (1965) The biology and habitats of mormyrid fishes. EAFFRO Annual Report for 1964, 58-66.

Olivry J.-C. (1986) Fleuves et rivières du Cameroun. Monographies Hydrologiques, ORSTOM, Paris,

733.

Olopade, O. A. (2013) Preliminary Observations on the Family Mormyridae in Oyan Dam Lake

(Nigeria). Transylvanian Review of Systematical and Ecological Research, 15(1), 33-48.

https://doi.org/10.2478/trser-2013-0004

Pappenheim, P. (1905) Ein neuer "pseudo"-elektrischer Fisch aus Kamerum, Mormyrus tapirus, sp. n.

Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin 1905 (8-9): 217-219.

Pappenheim, P. (1906) Neue und ungenügend bekannte elektrische Fische (Fam. Mormyridae) aus den

deutsch-afrikanischen Schutzgebieten. Sitzungsberichte der Gesellschaft Naturforschender Freunde zu

Berlin 1906, 10, 260-264.

Peel, R. A., Hill, J. M., Taylor, G. C. & Weyl, O. L. F. (2019) Food Web Structure and Trophic Dynamics

of a Fish Community in an Ephemeral Floodplain Lake. Frontiers in Environmental Sciences, 7(192). doi:

10.3389/fenvs.2019.00192

Pellegrin, J. (1927) Mormyridés du Cameroun recueillis par M. Th. Monod. Description d'un genre, de

quatre espèces et d'une variété. Bull. Soc. Zool. Fr., 52, 294-300.

Polačik, M., Harrod, C., Blažek, R., & Reichard, M. (2014). Trophic niche partitioning in communities of

African annual fish: evidence from stable isotopes. Hydrobiologia, 721(1), 99-106.

Polischuk S. C., Hobson K. A. & Ramsay M. A. (2001) Use of stable-carbon and -nitrogen isotopes to

assess weaning and fasting in female polar bears and their cubs. Canadian Journal of Zoology, 79(3), 499-

511. https://doi.org/10.1139/z01-007

Pool, T., Holtgrieve G., Elliott V., McCann K.,McMeans B., Rooney N., Smits A., Phanara T.,

Cooperman M. & Clark S. (2017) Seasonal increases in fish trophic niche plasticity within a flood-pulse

river ecosystem (Tonle Sap Lake, Cambodia). Ecosphere, 8:e01881.

Post, D. M. (2002) Using stable isotopes to estimate trophic position: models, methods, and assumptions. Ecology, 83(3), 703–71.

Sullivan, J. P., Lavoué, S. & Hopkins, C. D. (2002) Discovery and phylogenetic analysis of a riverine species flock of African electric fishes (Mormyridae: Teleostei). Evolution, 56, 597-616.

Quezada-Romegialli C., Jackson a. L., Hayden B., Kahilainen K. K., Lopes C. & Harrod C. (2018) tRophicposition, an R package for the Bayesian estimation of trophic position from consumer stable isotope ratios. Methods in Ecology and Evolution, 9, 1592–1599. https://doi.org/10.1111/2041-210X.13009

Rau, G. H., Mearns, A. J., Young, D. R., Olson, R .J., Schafer, H. A. & Kaplan, I. R. (1983) Animal 13 C/12 C correlates with trophic level in pelagic food webs. Ecology, 64, 1314-1318.

Schoener, T. W. (1974). Resource Partitioning in Ecological Communities: Research on how similar species divide resources helps reveal the natural regulation of species diversity. Science, 185(4145), 27-39.

Skelton, P.H., (2001) A complete guide to the freshwater fishes of southern Africa. Cape Town (South Africa): Struik Publishers, 395 p.

Soto, D. X., Decru, E., Snoeks, J., Verheyen, E., Van de Walle, L., Bamps, J., Mambo, T. & Bouillon, S. (2019) Terrestrial contributions to Afrotropical aquatic food webs: The Congo River case. Ecol Evol., 9, 10746–10757. https://doi.org/10.1002/ece3.5594

Taylor, G. C. (2016) Comparative fish ecology in three periodically connected rivers in the Upper Zambezi and Okavango ecoregions. PhD thesis, Rhodes University.

Thorp, J. H., and Delong, M. D. (1994). The riverine productivity model: an heuristic view of carbon sources and organic processing in large river ecosystems. Oikos, 70, 305–308. doi: 10.2307/3545642

Thorp, J. H., Delong, M. D., Greenwood, K. S., and Casper, A. F. (1998). Isotopic analysis of three food

web theories in constricted and floodplain regions of a large river. Oecologia, 117, 551–563. doi:

10.1007/s004420050692

Trites, A. W. (2001) Marine Mammal Trophic Levels and Interactions: Trophic Levels (Diet

Composition). Chapter in: Encyclopedia of Ocean Sciences. (Eds: J. Steele, S. Thorpe & K. Turekian).

Academic Press, San Diego, USA.

van der Waal, B. C. W. (1985). Aspects of the biology of larger fish species of Lake Liambezi, Caprivi,

South West Africa. Madoqua, 14, 101–144.

van der Waal, B.V. (1996). SOME OBSERVATIONS ON FISH MIGRATIONS IN CAPRIVI,

NAMIBIA. Southern African Journal of Aquatic Sciences, 22, 62-80.

Wang, J., Jiang, R-J., Xiao, Y., Yin, R., Chen, F., Zhou, Y-d. & Xu, H-X. (2022) Trophic Niche

Partitioning of Five Sciaenidae Species Sampled in Zhoushan Archipelago Waters via Stable Isotope

Analysis. Front. Mar. Sci., 9: 880123. doi: 10.3389/fmars.2022.880123

Wantzen, K. M., de Arruda Machado, F., Voss, M., Boriss, H., & Junk, W. J. (2002). Seasonal isotopic

shifts in fish of the Pantanal wetland, Brazil. Aquatic Sciences, 64, 239–251.

Winemiller, K. O. & Adite, A. (1997). Convergent evolution of weakly electric fishes from floodplain

habitats in Africa and South America. Environ. Biol. Fishes, 49, 175–186. doi:

10.1023/A:1007376826609

Yoğurtçuoğlu, B.; Ekmekçi, F.G.; Karachle, P.K. (2020) A review and assessment of fish trophic levels in

a large reservoir of central Anatolia, Turkey. Mar. Freshw. Res., 72, 311–320.

Zhang, Y. L., Xu, B. D., Zhang, C. L., Ji, Y. P., Ren, Y. P., Cheng, Y., et al. (2020). Spatial

Heterogeneity in the Feeding Habits and Feeding Ground Distribution of *Johnius Belangerii* in Haizhou

Bay During Spring. J. Fish. Sci. China 27, 315–326. doi: CNKI:SUN:ZSCK.0.2020-03-006

Table 1: overview of collected species and their representation. Please note that *Mormyrus* sp. "short snout" and *Mormyrus* sp. "long snout" refer to so far undescribed species and thus have these preliminary names in this study.

species	number of individuals	Season
Campylomormyrus phantasticus	32	dry 2017 and dry, transition, wet 2018
Hippopotamyrus castor	18	dry 2017 and dry, transition, wet 2018
Marcusenius mento	4	wet 2017 and transition 2018
Marcusenius sanagaensis	43	dry, wet 2017 and dry, transition, wet 2018
Mormyrops anguilloides	31	dry, wet 2017 and 2018, one transition sample from 2018
Mormyrops caballus	21	wet, dry 2017 and 2018
Mormyrus tapirus	52	dry, wet 2017 and dry, transition, wet 2018
Mormyrus sp. "long snout"	3	dry 2017
Mormyrus sp. "short snout"	36	dry, wet 2017 and dry, wet 2018
Paramormyrops batesii	4	wet 2017 and dry 2018
Petrocephalus christyi	44	dry 2017 and dry, transition 2018, one wet sample from 2018

Table 2: t-tests performed for delta nitrogen values as well as delta carbon values for species comparisons of the same genus

t-test with different variances	p-value (two-tailed)
δ^{13}C *Mormyrops caballus* vs. *Mormyrops anguilloides*	0.116
δ^{15}N *Mormyrops caballus* vs. *Mormyrops anguilloides*	0.650
δ^{13}C *Marcusenius mento* vs. *Marcusenius sanagaensis*	0.126
δ^{15}N *Marcusenius mento* vs. *Marcusenius sanagaensis*	0.515
δ^{13}C *Mormyrus tapirus* vs. *Mormyrus* sp."short snout"	4.42E-12*
δ^{15}N *Mormyrus tapirus* vs. *Mormyrus* sp."short snout"	0.002*
δ^{13}C *Mormyrus tapirus* vs. *Mormyrus* sp."long snout"	0.04
δ^{15}N *Mormyrus tapirus* vs. *Mormyrus* sp."long snout"	0.002*
δ^{13}C *Mormyrus long "snout"* vs. *Mormyrus* sp."short snout"	1.471E-05*
δ^{15}N *Mormyrus long "snout"* vs. *Mormyrus* sp."short snout"	1.003E-05*

* indicates a significant p-value (p-value < 0.025)

Table 3: results of single-factor ANOVAs and Tukey-Kramer test (if applicable) for all species with transition season samples

single-factor ANOVA	p-value	df	Tukey-Kramer q-stat values	dry vs wet	dry vs transition	wet vs transition	critical value with α=0.01
Mormyrus tapirus δ^{13}C	1.288E-07*	50		2.476	8.353**	9.242**	2.678
Mormyrus tapirus δ^{15}N	0.001*	50		0.677	5.253**	5.233**	2.678
Marcusenius sanagaensis δ^{13}C	0.027*	40		0.669	3.157**	3.861**	2.704
Marcusenius sanagaensis δ^{15}N	0.011*	40		0.382	3.796**	4.268**	2.704
Petrocephalus christyi δ^{13}C dry vs transition	0.520	19					
Petrocephalus christyi δ^{15}N dry vs transition	0.011*	19					
Campylomormyrus phantasticus δ^{13}C	0.016*	31		0.378	4.354**	3.53**	2.750
Campylomormyrus phantasticus δ^{15}N	0.654	31					
Hippopotamyrus castor δ^{13}C	0.232	14					
Hippopotamyrus castor δ^{15}N	0.169	14					

610   *indicates a p-value below 0.05 ; ** indicate q-stat value > critical value

Figure captions:

Figure 1: Overview of the study site: Map of the section of the middle Sanaga river close to the

Nachtigal Falls (Chutes de Nachtigal) in Cameroon, Africa. The most frequented sampled sites are up-

and downstream of Nachtigal falls and are in the vicinity of the town Batchenga (black dot). (Basemap:

https://www.esri.com/en-us/maps-we-love/overview, borders: https://www.diva-gis.org/gdata, rivers,

streams and places: openstreetmap.org).

Figure 2: Stable isotope signature of eleven Sanaga river mormyrid species. A) Results of the stable

isotope analysis of the muscle samples - mean and standard errors shown. δ^{15} nitrogen values on the y-

axis and δ^{13} carbon values on the x-axis; unit in parts per thousand (‰). B) Stable isotopic signatures

shown for all individuals. Note that the plots do not include the transition season samples specifically

addressed in the text and in Figure 5. Note the different scale of plot A and B. C) Phylogenetic

relationships of the studied mormyrid species based on the COI sequences and reconstructed by Bayesian

inference. The colour code corresponds to the species in the plots above. D) Head shape of different

species projected on the results of the stable isotope analysis (plot A) to visualize that species with snout

protrusion tend to exhibit higher values in δ^{13} C and/or δ^{15} N.

Figure 3: combined isotopes plot showing both the muscle isotope means and standard errors

together with the trophic samples values. δ^{15} nitrogen values on the y-axis and δ^{13} carbon values on the

x-axis; unit in parts per thousand (‰).

Figure 4: **TP (Trophic Position) plots comparing TPs between two species within one genus.** A)

Mormyrops anguilloides vs. *Mormyrops caballus*, B) *Marcusenius sanagaensis* vs. *Marcusenius mento*.

C)-E) Three species of *Mormyrus* show larger differences among themselves, and in this case species

with shorter protrusion tend to have a higher trophic position. C) *Mormyrus tapirus* vs. *Mormyrus* sp.

"short snout", D) *Mormyrus* sp. "long snout" vs. *Mormyrus* sp. "short snout", E) *Mormyrus tapirus* vs.

Mormyrus sp. "long snout". The baseline was generated from all benthic-sourced samples from the

trophic samples.

Figure 5: season impact plot of stable isotopes: plotted means and standard errors of five mormyrid

species separated by wet (blue-coloured), dry (red-coloured) and transition (green-coloured) season. δ^{15}

nitrogen values on the y-axis and δ^{13} carbon values on the x-axis; unit in parts per thousand (‰). Note

that the *Petrocephalus christyi* wet sample is only one sample and thus no standard errors are available.

The dry season is encompassing the months of February and March, the wet season the months August,

September, October and November and the transition season the months of May, June and July.