Computational analysis of long-range allosteric communications in CFTR

Ersoy Ayca[1,2], Altintel Bengi [1,2*], Livnat Levanon Nurit[3*], Ben-Tal Nir[4], Haliloglu Turkan,[1,2], and*

Lewinson Oded[3].

[1] Department of Chemical Engineering, Bogazici University, 34342 Istanbul, Turkey

[2] Polymer Research Center, Bogazici University, 34342 Istanbul, Turkey

[3] Department of Molecular Microbiology, Bruce and Ruth Rappaport Faculty of Medicine, Technion-Israel Institute

of Technology

[4] Department of Biochemistry and Molecular Biology, Faculty of Life Sciences, Tel-Aviv University, Tel-Aviv, Israel

* Equal contribution

Classification Physical science

Keywords Cystic fibrosis, CFTR, ABC transporter, allosteric regulation, conformational

dynamics.

Abstract. Malfunction of the CFTR protein results in cystic fibrosis, one of the most common

hereditary diseases. CFTR functions as an anion channel, the gating of which is controlled by long-

range allosteric communications. Allostery also has direct bearings on CF treatment: the most

effective CFTR drugs modulate its activity allosterically. Herein, we integrated Gaussian Network

Model, Transfer Entropy, and Anisotropic Normal Mode-Langevin dynamics and investigated the

allosteric communications network of CFTR. The results are in remarkable agreement with

experimental observations and mutational analysis and provide extensive novel insight. We

identified residues that serve as pivotal allosteric sources and transducers, many of which

correspond to disease causing mutations. We find that in the ATP-free form, dynamic fluctuations

of the residues that comprise the ATP binding sites facilitate the initial binding of the nucleotide.

Subsequent binding of ATP then brings to the fore and focuses dynamic fluctuations that were

present in a latent and diffuse form in the absence of ATP. We demonstrate that drugs that

potentiate CFTR's conductance do so not by directly acting on the gating residues, but rather by

mimicking the allosteric signal sent by the ATP binding sites. We have also uncovered a previously

undiscovered allosteric "hotspot" located proximal to the docking site of the phosphorylated

Regulatory (R) domain, thereby establishing a molecular foundation for its phosphorylation-

dependent excitatory role. This study unveils the molecular underpinnings of allosteric

connectivity within CFTR and highlights a novel allosteric "hotspot" that could serve as a

promising target for the development of novel therapeutic interventions.

Introduction. Cystic fibrosis (CF) is a multifaceted disorder impacting the epithelial tissues of
various organs, including the lungs, intestinal tract, and pancreas. It stands as one of the prevailing
hereditary afflictions, affecting approximately one in every 3,000 newborns(1). Notwithstanding
the significant advancements in therapeutic interventions, individuals with cystic fibrosis (CF)
continue to endure pronounced morbidity and a reduced life expectancy. (2). The disease is caused
by mutations in the gene encoding the cystic fibrosis transmembrane conductance regulator
(CFTR) protein. Most CF patients carry a deletion of Phe 508 in NBD1 which leads to protein
instability and premature degradation. More than 2000 other disease-associated mutations have
been documented (3), many of which remain uncharacterized. CFTR functions as a chloride
channel, the excretion of which is important for mucus hydration. Malfunction of CFTR leads to
mucus dehydration and a subsequent decrease in mucus clearance. The over-accumulation of
mucus leads to persistent bacterial infections, chronic inflammations, and to organ failure (4).
CFTR is a member of the ABC transporter protein family, considered to be one of the largest
protein families of any proteome (5). CFTR is a unique family member, as all other ABC
transporters function as active pumps, moving biomolecules against their concentration gradients
(6,7). The domain organization of CFTR is similar to that of canonical ABC transporters,
comprising two transmembrane domains (TMDs) and two cytosolic nucleotide binding domains
(NBDs). The TMDs form the translocation channel and the NBDs control channel opening and
closing through processive cycles of ATP binding, hydrolysis, and release. A unique feature of
CFTR is the regulatory domain (R) which contains multiple phosphorylation sites and connects
the two NBDs giving rise to a TMD1–NBD1–R–TMD2–NBD2 topology. Decades of structure-
function studies enabled the formalization of a mechanistic model for the function of CFTR: In
the resting (non-conducting) state, the funnel-shaped pore is inward-facing and is solvent
accessible only to the cytoplasm. The NBDs are separated, and the R domain is wedged between
them, preventing their dimerization (8,9). Upon phosphorylation of the R domain by PKA, it
disengages from its inhibitory position and shifts aside (10,11). The NBDs are now free to bind
ATP and form a ''head-to-tail'' dimer (6,12,13), with the two ATP molecules sandwiched at
composite binding sites formed at the dimer interface. In CFTR, like in some other ABC
transporters, only one of the ATP sites is hydrolysis-competent while the other (termed the
degenerate site) can only bind ATP but not hydrolyze it (14–16). Formation of the NBDs dimer
leads to rearrangements of transmembrane gating residues and subsequent channel opening.

Hydrolysis of ATP at the catalytic site and release of hydrolysis products leads to opening of the

NBDs, channel closure, and resetting of the system. This mechanism bears resemblance to that of

bona fide ABC transporters, where ATP binding and hydrolysis at the NBDs drive the conversion

of the TMDs between inward- and outward-facing conformations (17–19).

It is now broadly recognized that most proteins and enzymes are allosteric, where ligand

binding at one site affects the functional properties of another (often distant) site (20–23). Two

conceptual models dominated the field of allostery for the past 50 years. According to Koshland,

Neḿethy, and Filmer (KNF model, (24)), the protein exists in a single conformation (an inactive

one) to which the allosteric ligand binds, inducing a conformational change that leads to the active

conformation. In contrast, Monod, Wyman, and Changeux (MWC model, (25)) introduced the

concept of "conformational selection", according to which proteins pre-exist in multiple

conformation, and the allosteric ligand preferentially binds to the active conformer, leading to a

redistribution of the conformational distributions towards the active one (26). The simplicity of

these models is appealing, yet we now begin to understand that they alone cannot explain the

complexity of allosteric signal generation and transduction. More recently, an alternative allosteric

mechanism was demonstrated, termed "dynamic mechanism" (27,28), where allostery is mediated

by changes in dynamic fluctuations around a mean conformation in the absence of a noticeable

conformational change (29,30). In some proteins, as we believe is the case of CFTR, multiple

allosteric mechanisms combine to generate the observed allosteric phenomenon.

There are abundant examples that long-range allosteric communications are essential to the

function of CFTR. One such example is the control of channel opening and closing, where the

binding site of the effector molecule (ATP) and the affected gating residues are separated by 60-

70Å (8,12). Very recently, Levring *et al.,* demonstrated that ATP-dependent dimerization of the

NBDs is insufficient for channel opening which additionally depends on an allosteric pathway that

connects the NBDs and the channel pore (31). Based on rate-equilibrium free-energy relationship

analysis Sorum *et al.,* reached similar conclusions, and suggested that this allosteric signal

originates at the NBDs and gradually propagates towards the gating residues along a clear

longitudinal protein axis (32). Another example of long-range allostery in CFTR is the finding that

point mutations at the cytosolic loops increase channel open probability (even in the absence of

ATP) by shifting a pre-existing conformational equilibrium towards the ligand bound conducting

conformation (33). Notably, allostery in CFTR has direct therapeutic relevance: The potentiator

drug ivacaftor and the dual-function modulator elexacaftor do not bind in the proximity of the
gating residues (34,35) yet increase the open probability (Po) of the channel (36,37). These and
other studies (38–40) significantly advanced our understanding of the function of CFTR and its
allosteric control.

However, some important features of the underlying allosteric network remain unknown.
For example, the complete allosteric map of CFTR remains to be determined, as are the causal
relations (driving vs. driven) between residues and domains. Similarly, the allosteric mechanism
for modulation of CFTR by commercial drugs is only partially understood. Herein, we sought to
further our understanding of the allosteric network of CFTR. For this, we combined two
complementary computational approaches: Gaussian network model-Transfer Entropy (GNM-TE
(41,42)), and ANM-LD simulations which combine Anisotropic Network Model (ANM) (43–46)
and Langevin Dynamics (LD) (47).

GNM reduces the complexity of a protein polymer to a network of nodes (representing the main
chain α carbons) connected by springs with a spring constant of unity. Further simplifications are
the assumptions of force linearity and isotropy of all fluctuations. These simplifications are the
source of both its advantages and limitations. On the one hand, GNM is computationally
economical, allowing for rapid analytical derivation of conformational dynamics even of large
proteins. This computational economy, while offering valuable insights, does entail a limitation.
GNM proves highly effective in predicting fluctuations around a predefined native state and offers
valuable information regarding allosteric communication and causality. Nevertheless, due to its
reliance on topology, GNM is unsuitable to monitor the conformational transitions from one native
state to another. However, it does have the capacity to suggest potential modes of motion
accessible within a given protein topology that may contribute to such conformational changes. In
addition, GNM cannot provide any information on side chains dynamics as these are omitted from
the model. The robustness of the model makes GNM a highly effective tool in studying global
dynamics. Despite, or because of its simplicity GNM based approaches have been very successful
in describing global conformational dynamics of proteins (48–50). More details on the
fundamentals of GNM, its advantages and limitations can be found here (51). To supplement
GNM we integrated it with Transfer Entropy calculations. The concept of Transfer Entropy (TE)
is derived from information theory and is used to measure the amount of information transferred

between two processes (41). In the context of proteins' conformational changes, TE can be used

to determine whether two residues are allosterically connected (42). TE for each pair of i and j

residues can be viewed as the degree to which the present movement of residue i decreases the

uncertainty regarding the future movement of residue j within a specified time delay τ. If TEij (τ)

$>$ TEji (τ), then the dynamics of residue i affects the dynamics of residue j, representing a causal

relationship between residues i and j. If a residue transfers information to many other residues in

the protein, it is considered as an information source; if it accepts information from much of the

protein, it is considered as an information receiver or a 'sink' (for more details about the correlation

between TE and allostery see (52)). In the present work, we performed TE calculations based on

Gaussian Network Model (GNM), which considers harmonic interactions between residue pairs

(41,50). Unlike simulation trajectory approaches that infer cause-and-effect relations from the

observed sequence of events (which may be circumstantial), GNM-based TE provides a direct

measure, at the linear approximation, of causality relations between residues and domains.

Quantitative inference of cause-and-effect relations from simulation trajectory approaches is in

principle possible using enhanced kinetic sampling techniques but is highly computationally

demanding for a large transmembrane transporter such as CFTR. Here we opted to use a

complimentary GNM-based TE approach that readily provides a quantitative measure of sources

and receivers of allosteric signals.

To complement GNM, we used a hybrid molecular simulations approach that integrates global

elastic modes with atomic-scale information derived from all-atom simulations. Several such

approaches have emerged in recent years (*e.g.,* collective MD (CoMD), Perturb-Scan-Pull,

eBDIMS, H-REMD) (53–56) with the aim of finding a "sweet spot" between the precision of

whole atom simulations and the computational efficacy of coarse-grained elastic networks. The

novelty of our approach, ANM-LD, is that it integrates a low-resolution α-carbon-based

anisotropic network model with stochastic all-atom implicit solvent Langevin dynamics (LD)

simulations (57). ANM-LD reflects only internal dynamic modes, free of any external bias. More

importantly, ANM-LD provides experimentally testable predictions, and has been successfully

used to compute conformational trajectories of proteins (50,58).

By combining ANM-LD and GNM-TE we present here the allosteric network of CFTR.

The results align very well with experimental and clinical data, disclose the allosteric mechanism

of CFTR, and identify a novel allosteric hotspot that may be used for therapeutic intervention.

Results

Transfer Entropy calculations identify allosteric hotspots in CFTR.

First, we performed GNM-TE calculations on non-phosphorylated ATP-free human CFTR

(PDB ID 5UAK, (8)) using the ten slowest GNM "modes". As explained in the methods section

and elsewhere (50,59), GNM decomposes a complex motion (such as a conformational change, or

a person walking) to discrete dynamic building blocks, historically termed GNM "modes", the

ensemble of which comprise the complex motion. These include modes of motion that involve

large parts of the protein and lead to major conformational changes (termed "slow modes"). The

"slow modes" often display the greatest degree of collective movements. Modes that involve local

conformational adjustments and a smaller number of residues and display fewer collective

movements are considered "fast modes". We expected that the ten slowest modes, those

comprising the largest of motions, will be the main allosteric effectors, and their ensemble will

suffice to capture the overall dynamics of CFTR. Indeed, Eigenvalue analysis demonstrated that

the ten slowest modes of CFTR encompass its complete dynamic spectrum (Figure S1).

Figure 1 shows the residues identified as the main allosteric sources in ATP-free human CFTR,

obtained by using an interaction cutoff distance (R_{cut}) of 10Å. Very similar results were obtained

when R_{cut} of 7Å was used instead (Figure S2), suggesting that the results are not very sensitive to

this parameter. The residues that GNM-TE identified as main allosteric determinants showed

partial overlap with the hinge residues identified by GNM: Of the 60 residues that serve as main

allosteric sources, 25 (~41%) were located within 4Å of a hinge residue, and this co-localization

was highly statistically significant (Figure S3). This is perhaps not unexpected, as hinge residues

likely play an important role in transmitting the strain energy induced by binding of allosteric

ligands.

As GNM-based TE has not been extensively used to study membrane proteins as large and

complex as CFTR we began by evaluating its utility in extracting meaningful functional

information. Allosteric residues often coincide with functional epitopes, *i.e.,* residues that   are

involved in ligand binding, catalysis, or serve as conformational hinges (21,60). Therefore, we

first evaluated the performance of GNM-based TE by examining its ability to identify such sites

of functional importance. For this, we first compared the locations of the GNM-TE identified

allosteric sources to the positions of 14 residues that have been experimentally demonstrated to be

directly responsible for the function of CFTR. This subset of 14 residues includes the most

common CFTR gating mutations (G178, S549, G551, G970, G1244, S1251, S1255, G1349),

residues of the intracellular loops that are essential for conformational changes (M265, N268,

Y275, W1063, L1065), and the NBD2 Walker B aspartate D1370 which is essential for ATP

hydrolysis (8,15,32,38). As shown in Figure 1 we observed that these 14 residues tightly colocalize

with the main allosteric peaks. To assess the statistical significance of this co-segregation we

compared it to a random identification of residues of functional importance, as follows: GNM-TE

identified 60 residues (out of 1480) that are the main allosteric sources (*i.e.,* allosteric peaks) in

dephosphorylated ATP-free CFTR (Figure 1). Accordingly, we generated 100,000 sets of 60

randomized positions and counted how many times the positions of the functionally essential

residues were correctly predicted using this random allocation. As shown in Figure S4A, the

random allocation has a mean value of 0.7 correct guesses while the GNM-TE based identification

has a mean value of 8 ($p=9\times e^{-13}$). Next, we expanded this analysis to include not only exact

matches (between the random or GNM-TE assignment of functional sites and their real location)

but also first- or second- coordination sphere interactions (*i.e.,* peaks that are located <4Å or <7Å

from the functionally essential residues). Using these cutoffs, we also observed a highly significant

correlation between the location of the GNM-TE peaks and that of the functionally essential

residues (Figure S4B-C). We then compiled a second set of reference residues, one which includes

all the residues that provide the first ATP coordination sphere in ATP-bound CFTR (12). We

reasoned that since CFTR gating is controlled by ATP binding and hydrolysis these residues (that

are allosteric by definition) should be readily identified by the GNM-TE analysis. Indeed, as shown

in Figure 1 the ATP residues also closely segregated to the allosteric peaks identified by GNM-

TE, and this co-segregation outperformed the one obtained via a random allocation of allosteric

peaks (Figure S5A). This co-segregation was observed whether we considered only exact matches

(Figure S5A), or whether we included also first- or second- coordination sphere interactions

(Figures S5B and S5C, respectively). Next, we considered a more challenging reference set of

residues, one which includes the 93 CFTR missense point mutations that lead to cystic fibrosis (3).

This reference set of residues holds some inherent limitations: many of the mutations that lead to

CF have no direct functional role *per se*, rather, they cause CF due to compromised mRNA

synthesis/stability, protein processing, or protein targeting (61). The inclusion of such residues in

the benchmark dataset will lead to apparent false-negative identifications of the GNM-TE analysis. In addition, it is likely that not all the allosteric residues have been naturally mutated, leading to false-positive identifications. Nevertheless, despite these limitations, we expected to see some degree of colocalization between the identified allosteric hotspots and the disease-causing mutations. From the 93 mutations listed in the cfrt2 database (3) we removed the cluster of seven N-terminal residues (S13, L15, W60, E56, E60, P67, G85) that are known to be involved in interactions with cellular filamins or ER retention (62). As shown in Figure S6A we observed considerable association between the positions of 86 disease-causing mutations and the allosteric peaks identified by GNM-TE. To test the statistical significance of this association we generated 100,000 sets of randomly positioned "allosteric peaks". As shown in Figure S6B, the random distribution is roughly normal with a mean of 4.5 incidents of exact matches. As shown in Figure S6B the GNM-TE analysis is significantly different from the random distribution, with a mean of 9 exact matches. As observed with the previous two reference set of residues (Figures S4 and S5) the co-segregation between the allosteric peaks identified by GNM-TE and the positions of the disease-causing mutations was also observed when we included also first- and second-coordination sphere interactions (Figures S6C and S6D, respectively). In further support of the biological relevance of the GNM-TE calculations, and despite the structural similarity between the two NBDs, a clear asymmetry can be observed between them, with NBD2 which harbors the catalytic Walker B glutamate providing the higher allosteric signaling (Figures 1 and 2A). This observed asymmetry is in line with a recent single molecule conformational study where the degenerate and catalytic ATP sites were shown to asymmetrically affect NBD dimerization and gating (31). That GNM-TE identifies this asymmetry is quite remarkable, as it is based solely on the positions of the α-carbons and does not consider the side chains. Taken together, these results (Figures 1, 2A, S2, S4, S5, and S6) suggest that GNM-based TE can provide meaningful functional information on CFTR.

As shown in Figures 1 and 2A, the entropy sources residues (*i.e.,* residues that drive allostery) are found predominantly on the NBDs and are largely absent from the TMDs. The only TMDs residues that serve as entropic sources are those found at or near intracellular loops 2 and 3 (Figure 2A). Notably, these observations were made for de-phosphorylated ATP-free CFTR. This demonstrates that even when the NBDs are completely separated and the composite ATP binding sites are yet to form, the dynamic infrastructure for nucleotide binding is already present, poised

for binding of the ligand. This pre-existing dynamic infrastructure likely facilitates ATP binding

and will come into full play at a later stage of the conformational change, following

phosphorylation and binding of ATP. Interestingly, some of the main entropy peaks (*i.e.,* entropy

sources) located in the NBDs are slightly shifted from the ATP binding residues (Figure 1) and

correspond to adjacent residues. This shift disappears once ATP is bound, and the entropy source

role is transferred from the neighboring residues to the ATP binding residues themselves (see next

results section dealing with the effects of ATP binding).

The results shown in Figure 2A were obtained using the 10 slowest GNM modes. However,

during the calculations the entropic contribution of the slowest modes may overshadow that of the

faster (less collective) ones, which may nevertheless have important functional roles (for more

details see (42)). To circumvent this problem, one may repeat the calculations while omitting the

slowest GNM modes, enabling the detection of latent allosteric interactions that may otherwise be

hidden in the global fluctuations. For ATP-free human CFTR, the greatest effect was observed

upon removal of the first and second slowest modes: Upon removal of the slowest GNM mode, a

new cluster of residues that serve as allosteric sources was revealed (Figure 2B). These residues,

which are distributed roughly symmetrically in the TMDs, include the putative gating residues

(*e.g.,* P99, T338, S341, (8)) and additional residues that line the ion permeation pathway. The

residues that form the ivacaftor binding site (34) were also revealed as entropy sources by removal

of the slowest GNM mode (Figure 2B), demonstrating the utility of this approach. Removal of the

next slowest GNM mode reveals another cluster of TMD residues that serve as entropy/allostery

sources. Unlike the above-described symmetric cluster, this cluster is highly asymmetric, and is

comprised solely of residues in TMD1 (Figure 2C). Interestingly, this cluster of residues is located

at and just above the future site for docking of the regulatory domain ((63,64) and see also later in

Figure 3C-D).

The above analysis revealed two latent entropy hotspots: one surrounding the ion

permeation pathway (Figure 2B) and the other adjacent to the docking site of the regulatory domain

(Figure 2C). Both sites were hidden by the more dominant global fluctuations of the protein (Figure

2A). Their identification in this conformation, which is non-conducting and where the R domain

is still wedged between the NBDs (8), highlights once again the pre-existence of a dynamic

infrastructure which will assume a central role following phosphorylation and repositioning of the

R domain.

ATP binding and phosphorylation rewires and focuses the allosteric connectivity of CFTR

To study the effects of ATP binding and phosphorylation we conducted Transfer Entropy analysis of phosphorylated ATP-bound CFTR (PDB ID 6MSM, (12)). We observed that phosphorylation and binding of ATP greatly rewires the overall allosteric connectivity in CFTR (Figure S7 and compare Figures 1 and 3A): residues that serve as entropy sources are now found in all the domains (TMD1, NBD1, TMD2, NBD2) and the entropy peaks are sharp, sharper than those observed for the unphosphorylated ATP-free form (compare Figures 1 and 3A), indicating that information is originating from distinct residues/cluster of residues. This suggests that phosphorylation and binding of ATP focuses the dynamics of CFTR, which pre-existed in a more diffuse manner in the unphosphorylated ATP-free state. This observation suggests that this conformation is the preferred template/target for structure-guided development of allosteric modulators.

As discussed above and shown in Figure 1, in the ATP-free state the entropy peaks at the NBDs are slightly shifted from the ATP binding residues. This offset largely disappears upon binding of ATP, and the entropy peaks in the NBDs now closely match the location of the ATP binding residues (compare Figures 1 and 3A). In the TMDs, the residues that serve as main entropy sources are those that surround the ion permeation channel (Figure 3B). This means that upon phosphorylation and binding of ATP the gating and pore lining residues assume a more dominant role in allosteric signaling. Notably, these transmitting TMD residues were identified as latent entropy sources in the unphosphorylated ATP-free state (Figure 2B). This result again highlights that the underlying dynamics in CFTR pre-exist, prior to phosphorylation and ATP binding. To test if the effect of ATP binging is specific to human CFTR and/or is a result of the specific experimental conditions in which its ATP-free and ATP-bound structures were determined (8,12) we conducted similar calculations with the corresponding conformations of zebrafish CFTR (9,65). As shown in Figure S8, binding of ATP had a very similar effect in both CFTR homologues, suggesting that the allosteric effects of ATP binding are robust and conserved.

As explained above (see analysis of the ATP-free conformation), to identify latent entropy sources we repeated the TE calculations for the phosphorylated ATP-bound state after removing the two slowest modes. Several striking features were revealed by these calculations (Figure 3C). First, the entropy sources in the NBDs have now migrated from the ATP binding sites at the dimer interface towards the intracellular loops, perfectly positioned to transmit the signal to the TMDs.

Second, the TMDs allosteric peaks surrounding the gating residues and the ion permeation
pathway have migrated towards the extracellular exit, as if to prepare for ion exit. Lastly, a cluster
of residues that serve as strong entropy sources appeared in the TMDs in an asymmetric manner
(Figures 3C-D). These residues are found exclusively in TMD1 and comprise or are adjacent to
the site of interaction of the regulatory (R) domain (63,64). This observation suggests that the role
of the R domain may not be limited to steric obstruction of NBD dimer formation in the
unphosphorylated ATP-free state. The presence of this cluster of information-transmitting residues
in the vicinity of the docking site of the R domain may explain previous observations that deletion
of the R domain leads to fewer channel opening events, and that exogenous addition of the
phosphorylated R domain peptide increases channel open probability of CFTR construct deleted
of the R domain (66,67).

Concerted movements in CFTR. The Transfer Entropy (TE) analysis presented above provides a
measure of information transfer between residues and domains. However, it gives no information
of the physical nature of the conformational changes that are involved. To complement the GNM-
TE calculations, we used ANM-LD, a molecular simulations approach which computes the
trajectory of a conformational change between two known states of a protein (43,47). We simulated
the transition between non-phosphorylated ATP-free and phosphorylated ATP-bound human
CFTR (8,12). To understand the allosteric connectivity that underlies this conformational change,
we analyzed the degree of correlated movements between all CFTR residues. At the basis of this
analysis is the assumption that residues that are allosterically connected will move in synchrony
in both space and time. An extreme example of such synchronization is the perfect correlation of
each residue with itself (diagonal in Figure 4A). Similarly, two residues that are sequential in the
amino acid sequence will also display high correlation. However, when highly synchronized
movements are observed for residues or domains that are distant in sequence and in 3-D space,
this likely represents allosteric connectivity.

The 2-D cross correlation map obtained for the transition between non-phosphorylated
ATP-free and phosphorylated ATP-bound human CFTR is shown in Figure 4A. In this map red
colors indicate high positive correlations and blue colors high negative correlations. Red colors
thus indicate residues that move in parallel vectors in space and time, and blue colors indicate
residues that move in anti-parallel vectors in space and time. The map is dominated by large and

continuous patches of red (positive correlation) and blue (negative correlation) areas and is very

similar to the 2-D cross correlation map calculated for PglK (50), a lipid-linked oligosaccharide

flippase that adopts the canonical ABC exporter 3-D fold (68). These observations suggest that, as

inferred from the cryo-EM structures of CFTR (8,12), global rigid body motions with a relatively

small number of hinges underlie this conformational change, and that the conformational changes

of CFTR are similar to those of canonical ABC exporters. The two NBDs display asymmetric

allosteric connectivity: NBD1 shows a tighter correlation with the residues that surround the

permeation pathway (Figure 4B), while NBD2 (harboring the catalytic Walker B glutamate) is

more strongly correlated with ECL1 (extra cellular loop 1) (Figure 4C) which is known to stabilize

the conducting state of the channel (69).

Sequence of allosteric transduction in CFTR. The ANM-LD simulations offer an opportunity

to examine mechanistic features that are difficult to determine experimentally. One such feature is

the pre-equilibrium kinetics of allosteric interactions: if residues A and B are allosterically

connected, does the conformational change of A follow that of B, or *vice versa*? To investigate the

sequence of allosteric transduction in CFTR we re-analyzed the simulations while imposing a time-

delay between the conformational changes of allosterically coupled residues (or domains) of

interest. For any two allosterically coupled residues, if we define that the conformational change

in A occurs before that of B, and still observe a positive correlation in the 2-D correlations map,

this means that the conformational change of A precedes that of B. If the correlation is lost, this

means that B likely precedes A, which can be confirmed by imposing the opposite time-delay (*i.e.,*

imposing that the change in B precedes) and verifying that the positive correlation is again

observed (see *Methods* for a mathematical expression of these relations). This approach can be

used to track the entire route of allosteric signal transduction as detailed below.

Rate-equilibrium free-energy relationship analysis and single molecule FRET studies both

suggest that the allosteric signal originates at the NBDs and propagates to the TMDs (32,70). Our

results agree with these observations: in parallel simulations we repeatedly observed that the

conformational change from the dephosphorylated ATP-free state to the phosphorylated ATP-

bound one begins with the movement of the NBDs. In addition, the TE analysis of the inward-

facing ATP-free state demonstrated that in this conformation the allosteric peaks are located almost

exclusively at the NBDs (Figure 1). Of the two NBDs, the catalytic NBD2 seems to provide the

greater allosteric input (Figures 2A and S7A) and therefore as detailed below we began tracing the

allosteric signal at NBD2. As shown above (Figure 4C), in the absence of a time delay the

movements of NBD2, ICL2/3, TMD2 (including TM8 up to its hinge) and ECL1 are highly

synchronized. However, this correlation persists only if NBD2 leads this motion and ICL2/3,

TMD2/TM8 and ECL1 follow (Figure S9 green rectangles, Figure 5 step I). This observation

strongly suggests that the movement of NBD2 is the leading event and those of ICL2/3,

TMD2/TM8 and ECL1 follow, and that the allosteric signal propagates from NBD2 to ICL2/3,

TMD2/TM8, and ECL1 (Figure 5 center, green arrows). Notably, based on their recent single

molecule studies Levring *et al.,* proposed similar cause and effect relations between the catalytic

ATP site and TM8 (31).

From ECL1 the allosteric signal propagates to NBD1: The movements of these two

domains are highly correlated, but only if ECL1 leads and NBD1 follows (Figure S9 magenta

rectangles, Figure 5 step II, and Figure 5 center, magenta arrow). Similar considerations suggest

that from NBD1 the allosteric signal propagates to the gating residues and to the TM helices that

surround the permeation pathway: The movements of these domains are only synchronized if the

former leads and the latter follows (Figure S9 blue rectangles, Figure 5 step III, and Figure 5 center,

blue arrow). In turn, the movements of the gating residues and the permeation pathway TM helices

are correlated with those of NBD2, but only if the TM helices lead and NBD2 follows (Figure S9

orange rectangles, Figure 5 Step IV). We therefore conclude that the allosteric signal propagates

from the permeation pathway to NBD2, thus completing the cycle of allosteric transduction

(Figure 5 center, orange arrow).

Modulation of CFTR by drugs. In the past decade, the development of small-molecule CFTR

modulators revolutionized cystic fibrosis treatment and led to a dramatic reduction in patient

morbidity and increased life expectancy (71). There are currently four FDA-approved drugs that

are used to treat CF: The type I correctors lumacaftor (VX-809) and tezacaftor (VX-661) stabilize

mutant CFTR variants (predominantly the most common ΔF508 mutation) reducing premature

protein degradation and improving surface presentation of the mature channel (72,73). The

potentiator ivacaftor (VX-770) enhances the activity of conductance-defective mutants by

increasing the open probability (Po) of the channel (31,34,36), and the dual-function type III

corrector elexacaftor (VX-445) stabilizes unstable variants and increases channel open probability

(37,74). The recently published structures of WT and ΔF508CFTR bound to various drugs (35)

enabled to us to investigate the allosteric effects of the drugs. First, we conducted GNM-TE

analysis on ΔF508CFTR bound to Trikafta (PDB ID 8EIQ, (35)), a trivalent drug which combines

ivacaftor, tezacaftor, and elexacaftor and is currently the most advanced cystic fibrosis treatment.

As shown in Figure 6A, the type I correctors lumacaftor and tezacaftor bind at 'valleys', *i.e.,* to

residues that are not allosteric sources. These results are consistent with the mechanism of these

drugs which improve the function of CFTR by lowering its $\Delta G_{(folding)}$ rather than by modulating its

activity (73). In contrast, the residues that comprise the binding sites for ivacaftor and elexacaftor

also include ones that serve as main allosteric sources (Figure 6A), alluding to their role as

allosteric modulators.

In CFTR, channel opening depends on phosphorylation of the regulatory domain and subsequent

binding of ATP by the NBDs. Accordingly, mutations that interfere with binding of ATP reduce

channel opening probability and result in cystic fibrosis (CF). The potentiator drug ivacaftor is

used to treat CF patients who carry mutations in the NBDs (most commonly the G551D mutation

at the highly conserved LSGGQ ATP binding/signature motif) (34,36,75,76). Ivacaftor increases

channel opening probability despite impaired ATP binding and hydrolysis. The molecular basis

for the beneficial effects of ivacaftor is not fully understood, especially since it does not bind in

the vicinity of either the ATP- or ion-coordinating residues but rather peripherally at the protein-

lipid interface (34). In addition, unlike the folding correcting drugs, ivacaftor does not induce a

significant conformational change in ΔF508 CFTR (35). To better understand the mechanism

underlying the CFTR's modulation by ivacaftor we mapped the residues that receive information

from the ivacaftor binding site. Surprisingly, the gating residues and those that line the permeation

pathway were not among the residues that receive information from the drug binding site (Figure

6B), leaving the molecular basis of the drug's action unclear. Thus, in attempt to resolve this issue,

we compared the profile of the residues that receive information from the ivacaftor binding site to

those that receive information from the ATP binding sites. As shown in Figure 6C, the two profiles

showed remarkable overlap: The ivacaftor docking site and the ATP binding site send information

to a very similar set of residues (Pearson's correlation coefficient of 0.83). These observations

suggest that ivacaftor increases the opening probability not by directly affecting the gating residues

but rather indirectly by mimicking the allosteric signaling evoked by ATP binding. Further insight

for the mechanism of action of ivacaftor can be obtained from the structural representation of the

allosteric signaling shown in Figure 6D-F. The gating region (*e.g.,* gating residue F337) (12)) sends

information mostly to the extracellular side of TM helices 2 and 11 and to the cytoplasmic side of

helices 3, 4, 7 including intracellular loops 2 and 3 (Figure 6D). In comparison, G551 of the ATP

binding signature motif (mutations in which are treated with ivacaftor) sends information

predominantly to the intracellular side of TM helices 3, 4, 7, 8 and to the C-terminal intracellular

helix of NBD2 (Figure 6E). Superimposition of these effects, which represents the combined

allosteric signaling of the ion-coordinating and ATP-binding residues (shown in Figure 6F) is very

similar to the allosteric signaling sent by the ivacaftor binding residue F312 (Figure 6B). This

result suggests that ivacaftor exerts its potentiating effect by mimicking the combined allosteric

output of the ATP-binding and gating residues, compensating for any potential allosteric

transduction short circuit cause by mutations.

Discussion – the allosteric mechanism of CFTR

Allostery in proteins is most often studied through the lenses of the classical KNF or MWC

models, and more recently the dynamic mechanism (see introduction and (25,27,28,77)). The

results presented herein suggest that none of these models can singly explain the allosteric

mechanism of CFTR.

In the absence of phosphorylation and ATP, CFTR adopts solely the NBDs-open

conformation and spontaneous conversions to the NBDs-closed conformation are not seen (70). In

addition, multiple phosphorylation events are required to dislodge the R domain from its

obstructing position, presenting a very large energetic barrier (8,78). This conformational

homogeneity of dephosphorylated ATP-free CFTR suggests that the first step of the

conformational change, that of movement of the R domain, cannot efficiently proceed via an

ensemble selection mechanism. In the absence of definitive experimental/computational evidence,

we currently propose that this step occurs by an induced fit mechanism, where the reversible

covalent phosphorylation is the allosteric ligand (Figure 7, step I). The next step of the

conformational change, binding of ATP and closure of the NBDs, is more complex. On the one

hand, single molecule studies showed that even after phosphorylation the NBDs solely adopt the

open conformation and do not spontaneously close unless ATP is added. Once ATP was added,

all the molecules converted to the closed-NBD conformation and were never seen to spontaneously

open their NBDs (70). We observed very similar conformational homogeneity and stability with

another member of the ABC transporter family (58). Along the same lines, MD simulations showed that the closed state with dimerized NBDs is stable over hundreds of nanoseconds (79). Taken together, these results strongly argue in favor of an induced-fit mechanism, and against MWC/ensemble-selection/population shift models which assume pre-existence of multiple conformations and a spontaneous dynamic equilibrium between them). Direct experimental evidence supporting an induced fit mechanism was also provided by macroscopic current recordings combined with double mutant cycles (80). On the other hand, when the NBDs are separated the ATP binding sites are yet to form it is hard to envision how the nucleotide binds to its non-existing binding site. Such large conformational changes are considered to occur only via an ensemble selection mechanism (81). We suggest that our results provide the solution: In the ATP free form, despite the separation of the NBDs, the main allosteric sites are located near the ATP binding sites (Figures 1 and 2A). Moreover, many of the dynamic features that manifest following ATP binding exist in a latent form in the ATP-free conformation (Figure 2B and C). This means that the ATP binding residues fluctuate considerably in the absence of nucleotide, sampling conformations in the vicinity of the NBDs-separated native state. Some of these conformations likely provide an initial state for the interaction with the nucleotide (Figure 7, step II). ATP will selectively bind to these conformers, as in ensemble selection allosteric mechanism. Once the docking of the nucleotide is initiated by this combination of dynamic fluctuations and ensemble selection closure of the NBDs can proceed by an induced fit mechanism (Figure 7, step III) as suggested by experimental evidence (80). The resulting conformational changes do not yet lead to channel opening and conductance: The phosphorylated ATP bound structure observed in the cryo-EM studies is non conducting: the pore is fully dehydrated, and an ion translocation pathway is missing. Therefore, phosphorylation, ATP binding, and closure of the NBDs are essential yet insufficient for channel opening. Several lines of evidence suggest that the final missing component that enables conductance are the dynamic fluctuations of the NBDs-closed state: MD simulations of chloride conductance showed that a major conformational change is not required for channel opening. Instead, slight side chain re-orientations of pore residues enable sporadic pulses of chloride conductance (79). Single molecule measurements also failed to identify any conformational changes between the conducting and non-conducting states (70). Along the same lines, allosteric potentiators that increase CFTR's channel-open probabilities do so without noticeable effects on its conformation (34). These studies suggest that the conformations of the

conducting and non-conducting states are very similar and that a major conformational change is not required to open the channel. Rather, transition to the conducting state occurs by a dynamic allosteric mechanism with rapid conformational sampling around the phosphorylated ATP bound NBDs-closed state (Figure 7, step IV).

With this combination of allosteric models in mind, we adopted a structural/dynamic view to study the molecular basis of allosteric connectivity in CFTR. The premise of the structural/dynamic perspective is that strain energy generated by binding of the allosteric ligand (phosphorylation and ATP in CFTR) is mechanically transmitted through a network of interacting residues to be relieved by a conformational switch. To identify this network of interacting residues we used two elastic network approaches: ANM-LD is a trajectory molecular simulation approach which identifies residues that interact during a conformational change (50). It is therefore particularly useful for studying induced fit mechanisms that assume the existence of such networks of interacting residues. GNM-TE on the other hand models the intrinsic dynamics of a protein around its mean native conformations and provides a useful tool to study the dynamic/ensemble selection components of CFTR's allostery. Both approaches are computationally economical, and for a protein as large and complex as CFTR provide an attractive alternative to atomistic MD simulations.

Despite their simplifications, and without any input of functional information, the ANM-LD/ GNM-TE combination clearly distinguishes the three main functional domains of the protein (*i.e.,* NBDs, gating residues, docking site of the R domain) from the rest of the protein (Figures 2 and 3). In further support of the utility of the calculations we observe a remarkable correlation between the residues identified as main allosteric sources and the positions of functionally essential residues (Figure S4), ATP-binding residues (Figure S5), and disease-causing mutations (Figure S6). In addition, despite the high structural similarity between the NBDs the calculations identify the functional singularity of the catalytic NBD2 and identifies it as the dominant allosteric source (Figure 2A). Taken together, these results show that the computations can capture the structural dynamics of CFTR.

The allosteric effects of ATP binding and hydrolysis are not always easily distinguishable. For example, Liu *et al.,* investigated the effects of the force originating from ATP hydrolysis on the chaperonin GroEL. They applied elastic network models and principal component analysis on a series of GroEL structures and found that in this processive enzyme ATP binding is insufficient

to induce the conformational changes, which require nucleotide hydrolysis (82). In CFTR, a large

body of evidence suggests that the allosteric driver is binding *per-se* and not hydrolysis: binding

of ATP under non hydrolyzing conditions is sufficient to dimerize the NBDs and to open the

channel (12,70). In addition, mutations and experimental conditions that stabilize the ATP-bound

state by reducing hydrolysis rates stabilize the NBDs-closed conformation and conducting state of

the channel. At first glance, the allosteric connectivity networks of the ATP-free and -bound

conformations seem very different (Figure S7). However, a closer examination reveals that binding

of ATP fine-tunes and brings to the fore dynamics that pre-existed in the ATP-free conformation.

For example, in the dephosphorylated ATP-free state conformation, the allosteric signaling

originates from many residues in the NBDs, mostly from those that surround the future ATP

binding sites (Figures 1, 2). Binding of ATP focuses these relatively diffuse dynamics to the exact

location of the ATP binding residues (Figure 3A). Similarly, in the ATP-free conformation the

residues that line the permeation pathway and the gating residues have a minor allosteric role and

are detected only once the more dominant fluctuations of the NBDs were omitted from the

calculations (Figure 2B). Following ATP binding the allosteric role of these residues becomes

much more dominant (Figure 3B). We propose that in CFTR the role of ATP binding is 1) To

induce the conversion to the NBDs-closed conformation via a combination of ensemble selection

and induced fit and mechanisms and 2) To modulate conformational dynamics that are present to

different degrees in other conformations via a dynamic allosteric mechanism. In CFTR, ATP

binding, rather than hydrolysis is sufficient to exert these effects: binding of ATP is sufficient to

induce the closure of the NBDs and to open the channel (70). ATP hydrolysis and product release

are not required for channel opening, but rather lead to separation of the NBDs and termination of

conductance. This differs from the role of ATP in processive enzymes such as the chaperonin

GroEL, where the exothermic energy released by ATP hydrolysis is required to drive the

conformational changes (82).

A new allosteric hotspot was identified in TMD1, in a cleft formed by TM helices 2, 3 and

6 (Figures 3C-D). Its proximity to the interaction site of the phosphorylated R domain may provide

the molecular basis for the dual role of the R domain: an inhibitory role while dephosphorylated,

and a conductance-stimulatory role once phosphorylated (66,83,84). Perhaps more importantly,

this allosteric cleft is highly druggable and provides a new target for structure-guided screening of

small molecules.

Materials and methods

GNM-based Transfer Entropy

GNM is a minimalist 1D elastic network model that assumes a Gaussian distribution of

fluctuations that is isotropic in nature, operating within the confines of a given protein's

topology. The GNM model relies on a pair of parameters: force constant governing harmonic

interactions and the cutoff distance within which these interactions occur. In this model, the force

constant is uniform for all interactions and is conventionally set to unity. The range of interaction

is defined by a cutoff distance, typically ranging between 7 to 10 Å, with a value of 10 Å being

employed in this study. Transfer Entropy (TE) formulation (52) which is based on GNM (59,85)

reveals the direction of information flow between two residues i and j. Given a certain movement

in residue i, TE is a measure of the reduction of the uncertainty in the movements of residue j

with a time delay τ between the movements of both residues.

$TE_{i,j}(\tau)$ between residues i and j at time τ is formulated as (41)

570
$$T_{i \to j}(\tau) = S\left(\Delta R_j(t+\tau) \middle| \Delta R_i(t)\right) - S\left(\Delta R_j(t+\tau) \middle| \Delta R_i(t), \Delta R_j(t)\right) \tag{1}$$

571   where the Ss are conditional entropies, given by,

572
$$S\left(\Delta R_j(t+\tau) \middle| \Delta R_i(t)\right) = - \left\langle \ln p\left(\Delta R_i(0), \Delta R_j(\tau)\right)\right\rangle + \left\langle \ln p(\Delta R_i(0))\right\rangle \tag{2}$$

573     $S\left(\Delta R_j(t+\tau) \middle| \Delta R_i(t), \Delta R_j(t)\right)$

574
$$= - \left\langle \ln p\left(\Delta R_j(0), \Delta R_j(\tau)\right)\right\rangle + \left\langle \ln p\left(\Delta R_i(0), \Delta R_j(0), \Delta R_j(\tau)\right)\right\rangle \tag{3}$$

575   For more details of the definitions of GNM time correlation of the residue fluctuations in Eqs. 4

576   and 5 see (41,59).

577   The net TE from residue i to j at a given τ is described as

578
$$\Delta T_{i \to j}(\tau) = T_{i \to j}(\tau) - T_{j \to i}(\tau) \tag{4}$$

where $\Delta T_{i \to j}(\tau)$ estimates the direction of Transfer Entropy between residues i and j in a certain

time delay τ. Thus, the TE calculations reveal entropy/information sources and receivers: Entropy

sources have positive net TE values and send information to many other residues, and entropy

receivers have negative net TE values and receive information from the rest of the structure.

The choice of τ between the movement of residues is important to identify causal interrelations:

when τ is too small, only local cause-and-effect relations (between adjacent amino acids) will be

revealed. if τ is big enough, few (if any) long range cause-and-effect relations will manifest. In a

previous work (50), we studied in detail the effects of choosing different τ values and found that

an optimal value of τ which maximizes the degree of collectivities of net TE values is in most

cases $3 \times \tau_{opt}$ (τ_{opt} is the time window in which the total TE of residues is maximized). (See also

Figure S1 in (50)).

Degree of Collectivity and TECol Score

The degree of collectivity is a measure of the number of residues that are influenced by the

movement of one residue, *i.e.,* how collective is the information transfer? Utilizing Bruschweiler's

study(86)(87), collectivity (K) values of residues in information transfer are calculated based on

the positive net TE values (42) as

$$K_{i,s} = \frac{1}{N} \exp\left(- \sum_{j=1}^{N} \alpha \left(\Delta T_{ij,s}(\tau) \right)^2 \log\left(\alpha \left(\Delta T_{ij,s}(\tau) \right)^2 \right) \right) \qquad (5)$$

where s is a selected subset of slow GNM modes, N is the total number of residues, and α is the

normalization factor which is determined as

$$\sum_{i=1}^{N} a \left(\Delta T_{ij,s}(\tau) \right)^2 = 1 \qquad (6)$$

Here, s covers three subsets of the ten slowest modes: Modes 1-10 (*i.e.,* all ten), modes 2-

10, and modes 3-10. τ is taken as three times the optimum tau value that maximizes the

collective information transfer(42)(42). The 10 slowest modes are considered since the

eigenvalue distribution of the dynamic modes shows they fully represent the slow end of the

dynamic spectrum (Figure S1).

604
605   TECol Score of each residue i is calculated by the multiplication of its cumulative positive net TE

606   value (the sum of positive net TE values that residue i sends to the other residues) with its

607   collectivity $K_{i,s}$ value in each subset s as

$$608 \qquad TECol\ Score_{i,s} = K_{i,s} \cdot \sum_{j=1}^{N} \Delta T_{ij,s} \qquad\qquad (7)$$

The TECol Score, combining TE and collectivity values, is used in the determination of the most

functionally plausible global information source residues that are powerful effectors.

ANM-LD Simulations

The conformational change between dephosphorylated ATP-free (PDB ID: 5UAK)(8) and ATP-

bound (PDB ID: 6MSM)(12)(12) human CFTR was simulated using the ANM-LD approach

(50,58)(50,58). ANM-LD combines a low-resolution alpha-carbon based Anisotropic Network

Model (ANM) with stochastic all-atom implicit solvent Langevin Dynamics (LD) simulations.

Intrinsic dynamics characterized by ANM dynamic modes and followed short-time LD simulations

guide the conformational transition from the initial to the target state. That is, the conformational

transition pathway is successfully evolved from the dephosphorylated ATP-free to the

phosphorylated ATP-bound CFTR, starting with 7.5 Å and converging to 2.3 Å of RMSD. Due to

the lack of a characterized structure, the regulatory domain is not included in the ANM-LD

simulations. The time window in ANM–LD simulations does not reflect real time of simulated

transitions yet provides information on transition trajectory and sequence of events, using

accessible internal dynamics as the only bias towards the target conformation.

ANM-LD Simulation Flow

The hessian matrix (**H**) with a distance threshold radius (R_{cut}=13 Å) is constructed at each cycle

of the ANM-LD simulation to calculate 3N-6 ANM dynamic modes, i.e., eigenvectors $\mathbf{U}_k$ and

eigenvalue λ_k, k = 1, 3N-6, where N is the number of residues(43). The ANM mode that best

overlaps with the difference vector of the aligned initial/intermediate and target conformations is

selected ($\mathbf{U}_{best}$) and the initial conformation is deformed along this vector using a deformation

factor (DF: 0.4 Å). Energy minimization is performed for 500 steps and 100 steps of LD

simulations are performed with a time step of 0.2 fs at a temperature of 310 K using the Amber

11-Sander 4 biomolecular simulation program(88)(89).

The simplified position conformational sampling of ANM-LD cycles is given below as

$$R_{new} = R_{old} + \Delta R_{ANM} + \Delta R_{Eng.Min} + \Delta R_{LD} \qquad (8)$$

where $\Delta R_{ANM} = U_{best} \cdot DF$.

$\Delta R_{Eng.Min}$ and ΔR_{LD} are energy minimization and LD simulations contributions to the sampling,

respectively.

The ANM steps and LD steps are iteratively performed for a predetermined number of ANM-LD

cycles or until the RMSD between intermediate and target states converges. In the present ANM-

LD simulations of CFTR as the RMSD converged to a value of 2.3 Å.

Equal Time and Time Delay Cross Correlations

Each residue has a fluctuation vector *ΔR* that corresponds to the difference between its current

position to the mean position of its alpha carbons. The mean positions are calculated from the

selected time window of the ANM-LD simulation cycles. The pseudo correlation between the

fluctuations of residues *i* and *j* is calculated as

$$C_{ij}(\tau) = \frac{<\Delta R_i(t) \cdot \Delta R_j(t+\tau)>}{\sqrt{<\Delta R_i(t)^2>}\sqrt{<\Delta R_j(t+\tau)^2>}} \qquad (9)$$

where *t* represents an instantaneous pseudo time and τ is an additive time delay in ANM-LD

cycles. When τ equals to 0, *Cij* reduces to standard correlation between the movements of

residues *i* and *j*, with values between -1 and 1, where 0 corresponding to no correlation, and

positive and negative values corresponding to movement in the same or opposite directions,

respectively. When τ is greater than 0, the correlation between the fluctuations of residues *i* at

time *t* and *j* at $t+\tau$ indicates that the movement of residue *i* is leading and of residue *j* is following

in the case of $C_{ij}(\tau) > C_{ji}(\tau)$. The time window in the conformational transition (part of the

ANM-LD simulation trajectory from which the conformations are extracted) and the time delay τ

are the two main parameters in defining the correlation behavior.

The time delay, τ, an order parameter in a running time window, is a measure of changes in the

dynamics that allows a temporal coarse graining in the transition pathway, i.e. quantifying

directional correlations. Leading/following events, depending on the length of the time delay and

the time window, reveal cues about the complex hierarchical reorganization of residues in their

motion. Here, a time delay of 16 cycles is taken in a transition window of 48 cycles from the

ATP-free to the ATP-bound states of CFTR. This time delay corresponds to a time window at

which the autocorrelation average over all residues shifts from a positive value to a negative

value with respect to the mean conformation within 48 cycles.

Statistical Analysis

In dephosphorylated ATP-free human CFTR (PDB ID 5UAK) we identified a total of 60

allosteric peaks with (a) above average TECol score and (b) above average peak prominence.

We then counted how many times the positions of the functionally essential residues or ATP

binding sites spatially correlate with these peaks using cut-off distances of 0 Å (i.e., exact

match), 4 Å, or 7 Å.

To evaluate the statistical significance of these predictions, we generated 100,000 sets of 60

randomly localized allosteric peaks and counted how many times the positions of the

functionally essential residues or ATP sites were in the vicinity of these random allocations.

Thus, Z scores of the predictions were determined using the definition of

678
$$Z = \frac{X - \mu}{\sigma} \tag{10}$$

where X is the count of correct guesses of our prediction while μ and σ are the mean and the

standard deviation of the correct matches of random samples, respectively. From the obtained Z

scores, p-values of the predictions were calculated by one-tailed hypothesis test using a

significance level of 0.05.

The same statistical analysis is performed for the spatial correlation of the hinge positions with

the entropy source peaks using cut-off distances of 4 Å or 7 Å.

Visualization

The 3D structural figures presented in this work are created using PyMol 2.5.4 (The PyMOL

Molecular Graphics System).

References

1.   Farrell PM. The prevalence of cystic fibrosis in the European Union. Journal of Cystic
Fibrosis. 2008 Sep;7(5):450–3.
2.   McBennett KA, Davis PB, Konstan MW. Increasing life expectancy in cystic fibrosis:
Advances and challenges. Pediatr Pulmonol. 2022 Feb 1;57(Suppl 1):S5.
3.   Cftr2.org. The Clinical and Functional TRanslation of CFTR (CFTR2) [Internet]. 2011 [cited
2021 Dec 6]. Available from: https://cftr2.org/
4.   Fahy J V., Dickey BF. Airway Mucus Function and Dysfunction. New England Journal of
Medicine. 2010 Dec 2;363(23):2233–47.
5.   Dean M, Rzhetsky A, Allikmets R. The Human ATP-Binding Cassette (ABC) Transporter
Superfamily. Genome Res. 2001 Jul 1;11(7):1156–66.
6.   Rees DC, Johnson E, Lewinson O. ABC transporters: the power to change. Nat Rev Mol
Cell Biol. 2009 Mar;10(3):218–27.
7.   Gadsby DC, Vergani P, Csanády L. The ABC protein turned chloride channel whose failure
causes cystic fibrosis. Vol. 440, Nature. Nature Publishing Group; 2006. p. 477–83.
8.   Liu F, Zhang Z, Csanády L, Gadsby DC, Chen J. Molecular Structure of the Human CFTR Ion
Channel. Cell [Internet]. 2017 Mar 23 [cited 2020 Jul 19];169(1):85-95.e8. Available from:
https://pubmed.ncbi.nlm.nih.gov/28340353/
9.   Zhang Z, Chen J. Atomic Structure of the Cystic Fibrosis Transmembrane Conductance
Regulator. Cell. 2016 Dec 1;167(6):1586-1597.e9.
10.   Picciotto MR, Cohn JA, Bertuzzi G, Greengard P, Nairn AC. Phosphorylation of the cystic
fibrosis transmembrane conductance regulator. J Biol Chem. 1992;267(18):12742–52.
11.   Tabcharani JA, Chang XB, Riordan JR, Hanrahan JW. Phosphorylation-regulated Cl-
channel in CHO cells stably expressing the cystic fibrosis gene. Nature.
1991;352(6336):628–31.
12.   Zhang Z, Liu F, Chen J. Molecular structure of the ATP-bound, phosphorylated human
CFTR. Proc Natl Acad Sci U S A [Internet]. 2018 Dec 11 [cited 2021 Nov
19];115(50):12757–62. Available from: /pmc/articles/PMC6294961/
13.   Mense M, Vergani P, White DM, Altberg G, Nairn AC, Gadsby DC. In vivo phosphorylation
of CFTR promotes formation of a nucleotide-binding domain heterodimer. EMBO J. 2006
Oct 18;25(20):4728–39.
14.   Hohl M, Briand C, Grütter MG, Seeger MA. Crystal structure of a heterodimeric ABC
transporter in its inward-facing conformation. Nat Struct Mol Biol. 2012 Mar
25;19(4):395–402.
15.   Csanády L, Vergani P, Gadsby DC. Structure, gating, and regulation of the CFTR anion
channel. Physiol Rev. 2019 Jan 1;99(1):707–38.
16.   Stockner T, Gradisch R, Schmitt L. The role of the degenerate nucleotide binding site in
type I ABC exporters. Vol. 594, FEBS Letters. Wiley Blackwell; 2020. p. 3815–38.
17.   Johnson ZL, Chen J. ATP Binding Enables Substrate Release from Multidrug Resistance
Protein 1. Cell. 2018 Jan 11;172(1–2):81-89.e10.
18.   Kim Y, Chen J. Molecular structure of human P-glycoprotein in the ATP-bound, outward-
facing conformation. Science. 2018 Feb 23;359(6378):915–9.

19. Nguyen PT, Lai JY, Lee AT, Kaiser JT, Rees DC. Noncanonical role for the binding protein in
substrate uptake by the MetNI methionine ATP Binding Cassette (ABC) transporter.
Proceedings of the National Academy of Sciences. 2018 Nov 6;115(45):E10596–604.

20. Kessel A, Ben-Tal N. Introduction to Proteins: Structure, Function, and Motion. 2018th
ed. CRC Press; 2018.

21. Liu J, Nussinov R. Allostery: An Overview of Its History, Concepts, Methods, and
Applications. PLoS Comput Biol. 2016 Jun 1;12(6).

22. Gerek ZN, Ozkan SB. Change in allosteric network affects binding affinities of PDZ
domains: Analysis through perturbation response scanning. PLoS Comput Biol.
2011;7(10).

23. Campitelli P, Modi T, Kumar S, Banu Ozkan S. The Role of Conformational Dynamics and
Allostery in Modulating Protein Evolution. Vol. 49, Annual Review of Biophysics. Annual
Reviews Inc.; 2020. p. 267–88.

24. Koshland DE, Nemethy JG, Filmer D. Comparison of experimental binding data and
theoretical models in proteins containing subunits. Biochemistry [Internet]. 1966 Jan 1
[cited 2023 Sep 14];5(1):365–85. Available from:
https://pubmed.ncbi.nlm.nih.gov/5938952/

25. Monod J, Wyman J, Changeux JP. ON THE NATURE OF ALLOSTERIC TRANSITIONS: A
PLAUSIBLE MODEL. J Mol Biol [Internet]. 1965 [cited 2023 Sep 14];12(1):88–118.
Available from: https://pubmed.ncbi.nlm.nih.gov/14343300/

26. Ma B, Kumar S, Tsai CJ, Nussinov R. Folding funnels and binding mechanisms. Protein Eng
[Internet]. 1999 [cited 2023 Sep 14];12(9):713–20. Available from:
https://pubmed.ncbi.nlm.nih.gov/10506280/

27. Motlagh HN, Wrabl JO, Li J, Hilser VJ. The ensemble nature of allostery [Internet]. Vol.
508, Nature. Nature Publishing Group; 2014 [cited 2021 Nov 19]. p. 331–9. Available
from: /pmc/articles/PMC4224315/

28. Nussinov R, Tsai CJ. Allostery without a conformational change? Revisiting the paradigm.
Curr Opin Struct Biol [Internet]. 2015 Feb 1 [cited 2023 Sep 14];30:17–24. Available from:
https://pubmed.ncbi.nlm.nih.gov/25500675/

29. Popovych N, Sun S, Ebright RH, Kalodimos CG. Dynamically driven protein allostery. Nat
Struct Mol Biol [Internet]. 2006 Sep [cited 2023 Sep 14];13(9):831–8. Available from:
https://pubmed.ncbi.nlm.nih.gov/16906160/

30. Tsai CJ, del Sol A, Nussinov R. Allostery: Absence of a Change in Shape Does Not Imply
that Allostery Is Not at Play. J Mol Biol. 2008 Apr 18;378(1):1–11.

31. Levring J, Terry DS, Kilic Z, Fitzgerald G, Blanchard S, Chen J. CFTR function, pathology and
pharmacology at single-molecule resolution. Nature. 2023;

32. Sorum B, Czégé D, Csanády L. Timing of CFTR Pore Opening and Structure of Its Transition
State. Cell. 2015 Oct 22;163(3):724–33.

33. Wang W, Wu J, Bernard K, Li G, Wang G, Bevensee MO, et al. ATP-independent CFTR
channel gating and allosteric modulation by phosphorylation. Proc Natl Acad Sci U S A.
2010 Feb 23;107(8):3888–93.

34. Liu F, Zhang Z, Levit A, Levring J, Touhara KK, Shoichet BK, et al. Structural identification
of a hotspot on CFTR for potentiation. Science (1979) [Internet]. 2019 [cited 2021 Nov
19];364(6446):1184–8. Available from: https://pubmed.ncbi.nlm.nih.gov/31221859/

35. Fiedorczuk K, Chen J. Molecular structures reveal synergistic rescue of D508 CFTR by
Trikafta modulators. Science (1979). 2022 Oct 21;378(6617):284–90.
36. Eckford PDW, Li C, Ramjeesingh M, Bear CE. Cystic fibrosis transmembrane conductance
regulator (CFTR) potentiator VX-770 (ivacaftor) opens the defective channel gate of
mutant CFTR in a phosphorylationdependent but ATP-independent manner. Journal of
Biological Chemistry. 2012 Oct 26;287(44):36639–49.
37. Shaughnessy CA, Zeitlin PL, Bratcher PE. Elexacaftor is a CFTR potentiator and acts
synergistically with ivacaftor during acute and chronic treatment. Sci Rep. 2021 Dec
1;11(1).
38. Sorum B, Töröcsik B, Csanády L. Asymmetry of movements in CFTR's two ATP sites during
pore opening serves their distinct functions. Elife. 2017 Sep 25;6.
39. Scholl D, Sigoillot M, Overtus M, Martinez RC, Martens C, Wang Y, et al. A topological
switch in CFTR modulates channel activity and sensitivity to unfolding. Nat Chem Biol.
2021 Sep 1;17(9):989–97.
40. Wei S, Roessler BC, Chauvet S, Guo J, Hartman IV JL, Kirk KL. Conserved allosteric hot
spots in the transmembrane domains of Cystic Fibrosis Transmembrane Conductance
Regulator (CFTR) channels and Multidrug Resistance Protein (MRP) pumps. Journal of
Biological Chemistry. 2014;289(29):19942–57.
41. Hacisuleyman A, Erman B. Causality, transfer entropy, and allosteric communication
landscapes in proteins with harmonic interactions. Proteins: Structure, Function, and
Bioinformatics [Internet]. 2017 Jun [cited 2019 Aug 27];85(6):1056–64. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/28241380
42. Altintel B, Acar B, Erman B, Haliloglu T. Subsets of Slow Dynamic Modes Reveal Global
Information Sources as Allosteric Sites. J Mol Biol [Internet]. 2022 Sep 15 [cited 2023 Mar
22];434(17). Available from: https://pubmed.ncbi.nlm.nih.gov/35644497/
43. Atilgan AR, Durell SR, Jernigan RL, Demirel MC, Keskin O, Bahar I. Anisotropy of
Fluctuation Dynamics of Proteins with an Elastic Network Model. Biophys J [Internet].
2001 Jan [cited 2017 Apr 4];80(1):505–15. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/11159421
44. Wang Y, Rader AJ, Bahar I, Jernigan RL. Global ribosome motions revealed with elastic
network model. J Struct Biol. 2004 Sep;147(3):302–14.
45. Keskin O, Durell SR, Bahar I, Jernigan RL, Covell DG. Relating molecular flexibility to
function: A case study of tubulin. Biophys J. 2002;83(2):663–80.
46. Keskin O. Binding induced conformational changes of proteins correlate with their
intrinsic fluctuations: A case study of antibodies. BMC Struct Biol. 2007;7.
47. Brünger A, Brooks CL, Karplus M. Stochastic boundary conditions for molecular dynamics
simulations of ST2 water. Chem Phys Lett. 1984 Mar 30;105(5):495–500.
48. Yang LW, Bahar I. Coupling between catalytic site and collective dynamics: a requirement
for mechanochemical activity of enzymes. Structure [Internet]. 2005 Jun [cited 2023 Sep
14];13(6):893–904. Available from: https://pubmed.ncbi.nlm.nih.gov/15939021/
49. Li H, Sakuraba S, Chandrasekaran A, Yang LW. Molecular binding sites are located near
the interface of intrinsic dynamics domains (IDDs). J Chem Inf Model [Internet]. 2014 Aug
25 [cited 2023 Sep 14];54(8):2275–85. Available from:
https://pubmed.ncbi.nlm.nih.gov/25089914/

50.   Acar B, Rose J, Aykac Fas B, Ben-Tal N, Lewinson O, Haliloglu T. Distinct Allosteric
820   Networks Underlie Mechanistic Speciation of ABC Transporters. Structure [Internet].
821   2020 Apr 2 [cited 2020 Jul 19];28(6):651-663.e5. Available from:
822   https://pubmed.ncbi.nlm.nih.gov/32320672/
823   51.   Cui Q, Bahar Ivet. Normal mode analysis : theory and applications to biological and
824   chemical systems [Internet]. Chapman & Hall/CRC; 2006 [cited 2019 Sep 2]. 406 p.
825   Available from: https://www.crcpress.com/Normal-Mode-Analysis-Theory-and-
826   Applications-to-Biological-and-Chemical/Cui-Bahar/p/book/9781584884729
827   52.   Hacisuleyman A, Erman B. Entropy Transfer between Residue Pairs and Allostery in
828   Proteins: Quantifying Allosteric Communication in Ubiquitin. PLoS Comput Biol [Internet].
829   2017 Jan 1 [cited 2022 Dec 27];13(1). Available from:
830   https://pubmed.ncbi.nlm.nih.gov/28095404/
831   53.   Krieger JM, Doruker P, Scott AL, Perahia D, Bahar I. Towards gaining sight of multiscale
832   events: utilizing network models and normal modes in hybrid methods. Curr Opin Struct
833   Biol [Internet]. 2020 Oct 1 [cited 2023 Sep 14];64:34–41. Available from:
834   https://pubmed.ncbi.nlm.nih.gov/32622329/
835   54.   Gur M, Zomot E, Bahar I. Global motions exhibited by proteins in micro- to milliseconds
836   simulations concur with anisotropic network model predictions. J Chem Phys [Internet].
837   2013 Sep 28 [cited 2023 Sep 14];139(12). Available from:
838   https://pubmed.ncbi.nlm.nih.gov/24089724/
839   55.   Jalalypour F, Sensoy O, Atilgan C, Atilgan C. Perturb-Scan-Pull: A Novel Method
840   Facilitating Conformational Transitions in Proteins. J Chem Theory Comput [Internet].
841   2020 Jun 9 [cited 2023 Sep 14];16(6):3842–55. Available from:
842   https://pubmed.ncbi.nlm.nih.gov/32324386/
843   56.   Kandzia F, Ostermeir K, Zacharias M. Global Dynamics of Yeast Hsp90 Middle and C-
844   Terminal Dimer Studied by Advanced Sampling Simulations. Front Mol Biosci [Internet].
845   2019 Sep 1 [cited 2023 Sep 14];6(SEP). Available from:
846   https://pubmed.ncbi.nlm.nih.gov/31681792/
847   57.   Brünger A, Brooks CL, Karplus M. Stochastic boundary conditions for molecular dynamics
848   simulations of ST2 water. Chem Phys Lett. 1984 Mar 30;105(5):495–500.
849   58.   Yang M, Livnat Levanon N, Acar B, Aykac Fas B, Masrati G, Rose J, et al. Single-molecule
850   probing of the conformational homogeneity of the ABC transporter BtuCD. Nat Chem
851   Biol [Internet]. 2018 Jul 18 [cited 2018 Dec 10];14(7):715–22. Available from:
852   http://www.ncbi.nlm.nih.gov/pubmed/29915236
853   59.   Haliloglu T, Bahar I, Erman B. Gaussian Dynamics of Folded Proteins. Phys Rev Lett
854   [Internet]. 1997 Oct 20 [cited 2018 Dec 11];79(16):3090–3. Available from:
855   https://link.aps.org/doi/10.1103/PhysRevLett.79.3090
856   60.   Wodak SJ, Paci E, Dokholyan N V., Berezovsky IN, Horovitz A, Li J, et al. Allostery in Its
857   Many Disguises: From Theory to Applications. Vol. 27, Structure. Cell Press; 2019. p. 566–
858   78.
859   61.   Chen Q, Shen Y, Zheng J. A review of cystic fibrosis: Basic and clinical aspects. Animal
860   Model Exp Med. 2021 Sep 1;4(3):220–32.

861   62.   Thelin WR, Chen Y, Gentzsch M, Kreda SM, Sallee JL, Scarlett CO, et al. Direct interaction
862   with filamins modulates the stability and plasma membrane expression of CFTR. J Clin
863   Invest. 2007 Feb 1;117(2):364–74.
864   63.   Bozoky Z, Krzeminski M, Muhandiram R, Birtley JR, Al-Zahrani A, Thomas PJ, et al.
865   Regulatory R region of the CFTR chloride channel is a dynamic integrator of phospho-
866   dependent intra- and intermolecular interactions. Proc Natl Acad Sci U S A. 2013 Nov
867   19;110(47).
868   64.   Chappe V, Irvine T, Liao J, Evagelidis A, Hanrahan JW. Phosphorylation of CFTR by PKA
869   promotes binding of the regulatory domain. EMBO Journal. 2005 Aug 3;24(15):2730–40.
870   65.   Zhang Z, Liu F, Chen J. Conformational Changes of CFTR upon Phosphorylation and ATP
871   Binding. Cell [Internet]. 2017 Jul 27 [cited 2023 Sep 14];170(3):483-491.e8. Available
872   from: https://pubmed.ncbi.nlm.nih.gov/28735752/
873   66.   Winter MC, Welsh MJ. Stimulation of CFTR activity by its phosphorylated R domain.
874   Nature. 1997;389(6648):294–6.
875   67.   Ma J, Zhao J, Drumm ML, Xie J, Davis PB. Function of the R domain in the cystic fibrosis
876   transmembrane conductance regulator chloride channel. Journal of Biological Chemistry.
877   1997 Oct 31;272(44):28133–41.
878   68.   Perez C, Gerber S, Boilevin J, Bucher M, Darbre T, Aebi M, et al. Structure and mechanism
879   of an active lipid-linked oligosaccharide flippase. Nature. 2015 Aug 12;524(7566):433–8.
880   69.   Infield DT, Cui G, Kuang C, McCarty NA. Positioning of extracellular loop 1 affects pore
881   gating of the cystic fibrosis transmembrane conductance regulator. Am J Physiol Lung
882   Cell Mol Physiol. 2016 Mar 1;310(5):L403–14.
883   70.   Levring J, Terry DS, Kilic Z, Fitzgerald G, Blanchard S, Chen J. CFTR function, pathology and
884   pharmacology at single-molecule resolution. Nature [Internet]. 2023 Apr 20 [cited 2023
885   Sep 14];616(7957):606–14. Available from: https://pubmed.ncbi.nlm.nih.gov/36949202/
886   71.   Clancy JP. Rapid therapeutic advances in CFTR modulator science. Pediatr Pulmonol.
887   2018 Nov 5;53(S3):S4–11.
888   72.   Van Goor F, Hadida S, Grootenhuis PDJ, Burton B, Stack JH, Straley KS, et al. Correction of
889   the F508del-CFTR protein processing defect in vitro by the investigational drug VX-809.
890   Proc Natl Acad Sci U S A. 2011 Nov 15;108(46):18843–8.
891   73.   Fiedorczuk K, Chen J. Mechanism of CFTR correction by type I folding correctors. Cell.
892   2022 Jan 6;185(1):158-168.e11.
893   74.   Veit G, Vaccarin C, Lukacs GL. Elexacaftor co-potentiates the activity of F508del and
894   gating mutants of CFTR. Journal of Cystic Fibrosis. 2021 Sep 1;20(5):895–8.
895   75.   Hadida S, Van Goor F, Zhou J, Arumugam V, McCartney J, Hazlewood A, et al. Discovery
896   of N -(2,4-Di- tert -butyl-5-hydroxyphenyl)-4-oxo-1,4-dihydroquinoline-3-carboxamide
897   (VX-770, Ivacaftor), a potent and orally bioavailable CFTR potentiator. J Med Chem. 2014
898   Dec 11;57(23):9776–95.
899   76.   Yu H, Burton B, Huang CJ, Worley J, Cao D, Johnson JP, et al. Ivacaftor potentiation of
900   multiple CFTR channels with gating mutations. J Cyst Fibros. 2012 May;11(3):237–45.
901   77.   Ribeiro AAST, Ortiz V. A Chemical Perspective on Allostery. Chem Rev [Internet]. 2016 Jun
902   8 [cited 2023 Sep 14];116(11):6488–502. Available from:
903   https://pubmed.ncbi.nlm.nih.gov/26741913/

78.   Gadsby DC, Nairn AC. Control of CFTR channel gating by phosphorylation and nucleotide
hydrolysis. Physiol Rev [Internet]. 1999 [cited 2023 Sep 14];79(1 Suppl). Available from:
https://pubmed.ncbi.nlm.nih.gov/9922377/

79.   Zeng ZW, Linsdell P, Pomès R. Molecular dynamics study of Cl- permeation through cystic
fibrosis transmembrane conductance regulator (CFTR). Cell Mol Life Sci [Internet]. 2023
Feb 1 [cited 2023 Sep 14];80(2). Available from:
https://pubmed.ncbi.nlm.nih.gov/36694009/

80.   Szollosi A, Vergani P, Csanády L. Involvement of F1296 and N1303 of CFTR in induced-fit
conformational change in response to ATP binding at NBD2. J Gen Physiol [Internet].
2010 Oct [cited 2023 Sep 14];136(4):407–23. Available from:
https://pubmed.ncbi.nlm.nih.gov/20876359/

81.   Nussinov R, Ma B, Tsai CJ. Multiple conformational selection and induced fit events take
place in allosteric propagation. Biophys Chem [Internet]. 2014 [cited 2023 Sep
18];186:22. Available from: /pmc/articles/PMC6361548/

82.   Liu J, Sankar K, Wang Y, Jia K, Jernigan RL. Directional Force Originating from ATP
Hydrolysis Drives the GroEL Conformational Change. Biophys J [Internet]. 2017 Apr 25
[cited 2023 Sep 14];112(8):1561–70. Available from:
https://pubmed.ncbi.nlm.nih.gov/28445748/

83.   Ostedgaard LS, Baldursson O, Welsh MJ. Regulation of the Cystic Fibrosis
Transmembrane Conductance Regulator Cl- Channel by Its R Domain. Vol. 276, Journal of
Biological Chemistry. J Biol Chem; 2001. p. 7689–92.

84.   Csanády L, Chan KW, Seto-Young D, Kopsco DC, Nairn AC, Gadsby DC. Severed channels
probe regulation of gating of cystic fibrosis transmembrane conductance regulator by its
cytoplasmic domains. Journal of General Physiology. 2000;116(3):477–500.

85.   Bahar I, Atilgan AR, Erman B. Direct evaluation of thermal fluctuations in proteins using a
single-parameter harmonic potential. Fold Des [Internet]. 1997 Jun [cited 2018 Dec
11];2(3):173–81. Available from:
http://linkinghub.elsevier.com/retrieve/pii/S1359027897000242

86.   Brüschweiler R. Collective protein dynamics and nuclear spin relaxation. J Chem Phys
[Internet]. 1995 Feb 22 [cited 2023 Mar 29];102(8):3396–403. Available from:
http://aip.scitation.org/doi/10.1063/1.469213

87.   Brüschweiler R. Collective protein dynamics and nuclear spin relaxation. J Chem Phys.
1995 Feb 22;102(8):3396–403.

88.   D.A. Case, H.M. Aktulga, K. Belfon, I.Y. Ben-Shalom, J.T. Berryman, S.R. Brozell, D.S.
Cerutti, T.E. Cheatham, III, G.A. Cisneros, V.W.D. Cruzeiro, T.A. Darden, R.E. Duke, G.
Giambasu, M.K. Gilson, H. Gohlke, A.W. Goetz, R. Harris, S. Izadi, S.A. Izmailov and PAK.
AMBER 11 [Internet]. San Francisco.: University of California; 2010. Available from:
http://ambermd.org/

89.   D.A. Case, H.M. Aktulga, K. Belfon, I.Y. Ben-Shalom, J.T. Berryman, S.R. Brozell, D.S.
Cerutti, T.E. Cheatham, III, G.A. Cisneros, V.W.D. Cruzeiro, T.A. Darden, R.E. Duke, G.
Giambasu, M.K. Gilson, H. Gohlke, A.W. Goetz, R. Harris, S. Izadi, S.A. Izmailov and PAK.
AMBER 11. San Francisco.: University of California; 2010.

Figure 1

Figure 2

Figure 3

Figure 4

Step IV: gating residues and the TM helices surrounding permeation pathway lead NBD2

Step II: ECL1 leads NBD1

Step III: NBD1 leads the gating residues and the TM helices surrounding permeation pathway

Step I: NBD2 leads ICL2/3 and ECL1

T338

D110

ECL1

ICL2/3

NBD2

NBD1

S1251

T460

Figure 5

Figure 6

D

Received Entropy

x10⁻²

-2   2

E

Received Entropy

x10⁻²

-1.4   1.8

F

Received Entropy

x10⁻²

-1.2   1.8

Figure 6 (cont.)

I   II   III   IV

Induced fit   Dynamic   Ensemble selection   Dynamic
                                        + Induced fit

ATP

Figure 7

Figure 1. Functional sites of CFTR segregate to allosteric hotspots. Shown is the amount of information (TE score) transmitted by each residue of dephosphorylated ATP-free human CFTR (PDB ID 5UAK) calculated using the ten most collective GNM modes (solid black trace). The positions of the 14 functionally important residues (see text for details) and of 30 ATP binding residues are shown as cyan and magenta spheres, respectively.

Figure 2. Identification of allosteric hotspots in dephosphorylated ATP-free CFTR.
A cartoon representation of dephosphorylated ATP-free human CFTR (PDB ID 5UAK), where each residue is colored according to the amount of information it transmits (TE score), expressed as Transfer Entropy x Collectivity (TECol) calculated using the ten most collective GNM modes (A), or upon removal of the first (B) or first and second (C) most collective GNM modes. Red and blue colors represent high and low levels of information output, respectively.

Figure 3. ATP binding and phosphorylation rewires and focuses the allosteric connectivity of CFTR. (A) Shown is the amount of information transmitted (TE score) by each residue of phosphorylated ATP-bound human CFTR (PDB ID 6MSM) calculated using the ten most collective GNM modes (solid black trace). The positions of the ATP binding residues are shown as magenta spheres. (B,C) A cartoon representation of phosphorylated ATP-bound human CFTR (PDB ID 6MSM), where each residue is colored according to the amount of information it transmits (TE score), calculated using the ten most collective GNM modes (B), or upon removal of the first most collective GNM mode (C). Red and blue colors represent high and low levels of information output, respectively, and the ATP molecules are shown as black/grey spheres. (D) a magnified view of the allosteric cluster that is adjacent to the docking site of the R domain.

Figure 4. Synchronized movements of residues and domains of CFTR.
(A) Shown is a 2-D map of the dynamic cross-correlations between all residues of CFTR during the conformational transiting from the dephosphorylated ATP-free state to the phosphorylated ATP-bound state (PDB ID 5UAK and 6MSM, respectively). Red colors indicate strong positive correlation, meaning that the residues move in parallel vectors in space and time, blue colors indicate negative correlations. Domain boundaries are indicated by white lines. (B, C) Cartoon representation of CFTR where each residue is colored according to its dynamic cross-correlations with the ATP binding residues (black spheres) of NBD1 (B) or NBD2 (C).

Figure 5. Trajectory and sequence of allosteric transduction in CFTR.
Dynamic cross-correlations were calculated using a time delay (τ) of 16 cycles out of 50 required to complete the conformational transition between the dephosphorylated ATP-free and phosphorylated ATP-bound states. Red and blue colors indicate high and low correlations, respectively.
Steps I-IV: Each residue is colored according to the degree of the correlation of its movement at time (t + τ) relative to the movement at time (t) of S1251 of NBD2 (Step 1), D110 of ECL1 (Step 2), T460 of NBD1 (Step 3), and the gating residue T338 (Step 4).
Center (large) panel: shown is a summary of the allosteric trajectory, originating from NBD2 (green) to ICL2/3 and ECL1 (magenta), from ECL1 to NBD1 (blue), from NBD1 to the permeation pathway helices (orange), and finally back to NBD2.

Figure 6. Allosteric modulation of CFTR by drugs. (A) Shown is the amount of information TRANSMITTED (TE score) by each residue of CFTR ΔF508 bound to Trikafta (PDB ID 8EIQ) calculated using the ten most collective GNM modes (solid black trace). The binding sites of ivacaftor, elexacaftor, and lumacaftor/tezacaftor are shown as magenta, green, or cyan spheres, respectively. (B) Cartoon representations of ivacaftor bound CFTR (PDB ID 6O2P), with ivacaftor and ATP shown as magenta and black/grey spheres, respectively. Residues are colored according to the amount of entropy RECEIVED from the ivacaftor binding residue F312 (C) Shown is the amount of entropy RECEIVED by each residue of CFTR from the ATP binding residue G551 (blue trace) or the ivacaftor binding residue F312 (orange trace).
(D-F) Same as B, only the residues are colored according to the amount of entropy RECEIVED from the gating residue F337 (D), the ATP binding residue G551 (E), and the sum of the entropy received from both G551 and F337 (F).

Figure 7. Model of CFTR's conformational changes and their allosteric control. CFTR is shown in a grey space-filling model, the regulatory (R) domain as a red ribbon, and ATP in shades of green. Following its phosphorylation, the R domain dislodges from the NBDs dimer interface (Step I of the conformational change). This allows the ATP binding residues to fluctuate and sample conformers near the NBDs-open state (Step II), some of which provide an initial ATP binding site. In step III, this subset of conformers is selected by ATP to initiate binding (*i.e.,* ensemble selection), which then proceeds to complete closure of the NBDs by an induced fit mechanism. In step IV, the channel bursts to the conducting state by a dynamic mechanism, without a major conformational change.