A diversity of diversities: do complex environmental effects underpin associations between below- and aboveground taxa?

Authors: Fiona M. Seaton[1,2], Paul B. L. George[2,3], Jamie Alison[4,5], Davey L. Jones[2], Simon Creer[2], Simon M. Smart[1], Bridget Emmett[4] and David A. Robinson[4]

1. UK Centre for Ecology & Hydrology, Lancaster, UK

2. School of Natural Sciences, Bangor University, Bangor, Gwynedd, UK

3. Département de biochimie, de microbiologie et de bio-informatique, Université Laval, Quebec City, Canada

4. UK Centre for Ecology & Hydrology, Bangor, UK

5. Department of Ecoscience, Aarhus University, Aarhus, Denmark

Corresponding author: Fiona M. Seaton, UK Centre for Ecology & Hydrology, Lancaster Environment Centre, Library Avenue, Bailrigg, Lancaster, LA1 4AP, UK. fseaton@ceh.ac.uk

Abstract

1. To predict how biodiversity will respond to global change, it is crucial to understand the relative roles of abiotic drivers and biotic interactions in driving associations between the biodiversity of disparate taxa. It is particularly challenging to understand diversity-diversity links across domains and habitats, because data are rarely available for multiple above- and below-ground taxa across multiple sites.

2. Here we analyse data from a unique biodiversity dataset gathered across a variety of oceanic temperate terrestrial habitats in Wales, comprising 300 sites with co-located soil microbial, plant, bird, and pollinator surveys along with climate and soil physicochemical information. Soil groups are analysed using metabarcoding of the 16S, ITS1 and 18S DNA regions, allowing in depth characterisation of microbial and soil animal biodiversity.

3. We explore biodiversity relationships along three aspects of community composition: First, we assess correlation between the alpha diversity of different groups. Second, we assess whether biotic turnover between sites is correlated across different groups. Finally, we investigate the co-occurrence of individual taxa across sites. In each analysis, we assess the contribution of linear or non-linear environmental effects.

4. We find that a positive correlation between alpha diversity of plants, soil bacteria, soil fungi, soil heterotrophic protists, bees and butterflies is in fact driven by complex non-linear responses to abiotic drivers. In contrast, environmental variation did not account for positive associations between the diversity of plants and both birds and AM fungi, suggesting a role for biotic interactions.

5. Both the diversity and taxon-level associations between the differing soil groups remained even after accounting for non-linear environmental gradients. Aboveground, spatial factors played larger roles in driving biotic communities, while linear environmental gradients were sufficient to explain many group- and taxon-level relationships.

6. **Synthesis.** Our results show how non-linear responses to environmental gradients drive many of the relationships between plant biodiversity and the biodiversity of above- and belowground biological communities. Our work shows how different aspects of biodiversity might respond non-linearly to changing environments and identifies cases where management-induced changes in one community could either influence other taxa or lead to loss of apparent biological associations.

Keywords: above-belowground, bird diversity, joint species distribution modelling, plant diversity, pollinator diversity, soil microbial communities

Introduction

Ecosystem functionality and stability are underpinned by biodiversity (IPBES, 2019) and interactions between different components of the biosphere are essential to ecosystem maintenance. Correlation between the different biological communities of an ecosystem could be due to the direct influence of one taxonomic group on another or due to shared response to environmental conditions (Schulze et al., 2004; Wolters et al., 2006). Consequently, taxonomic or environmental mechanisms may lead to associated patterns of biodiversity but have differing implications for land use management. Indeed, opposing trajectories of above- and below-ground biodiversity as well as weakened biotic correlations have been observed in response to land use intensification (Gossner et al., 2016; Manning et al., 2015). Therefore, disentangling the relative influence of abiotic and biotic impacts upon the different components of the biosphere is key to understanding and predicting future responses of biological communities to environmental change.

There is a prevailing notion in ecology that diversity begets diversity. For example, a more diverse plant community can support a more diverse pollinator community through a proliferation of ecological niches (Schulze et al., 2004). There is still uncertainty over the strength of this synergistic effect, as relationships between the diversity of different taxa aboveground have been shown to be variable and weak (Wolters et al., 2006). Results investigating the relationship between above- and belowground biodiversity have also been variable, with either positive or no correlations commonly observed between plant diversity and soil microbial diversity (Hiiesalu et al., 2014; Leff et al., 2018; Prober et al., 2015). Certain taxonomic groups appear to show more positive relationships with plant diversity. For example, fungal diversity, particularly mycorrhizal diversity, is often found to be

positively related to plant diversity (Hiiesalu et al., 2014; Milcu et al., 2013; Nguyen et al., 2016; Peay et al., 2013; Ren et al., 2017; T. Yang et al., 2017). The type of plants that are considered in plant diversity inventories can also be important, for example the number of flowering plant species and their abundance has been shown to influence bee and butterfly abundance and richness (Kearns & Oliveras, 2009; Potts et al., 2009).

While relationships between the diversity of different taxonomic groups may be variable, there have been many results demonstrating that the species composition of plant assemblages may be related to the composition of other taxonomic groups even when diversity is not. Relationships between the community turnover of plants and the turnover of soil bacteria, fungi and protist communities have been found in both experimental and observational studies (Barberán et al., 2015; Cline et al., 2017; Delgado-Baquerizo et al., 2018; Leff et al., 2018; Prober et al., 2015). These correlations in composition and turnover in observational results can persist even after controlling for environmental drivers such as climate or soil physicochemical properties (Delgado-Baquerizo et al., 2018; Prober et al., 2015). Such changes in composition could be due to species-level associations between taxa, such as trophic interactions, symbioses or parasitic links.

Shared responses to environmental gradients could explain many of the measured associations between the biodiversity of different taxonomic groups. Some of the positive correlations that have been found between plant diversity and soil bacterial, fungal and protistan diversity are explained by shared response to environmental variables, e.g. soil pH and fertility (Goberna et al., 2016; Yashiro et al., 2018; Yuan et al., 2017). However, there are still many ecosystems, especially in extreme environments such as sub-polar regions and intensively managed agricultural systems where environmental factors may in fact cause

divergent patterns and disconnectance of biodiversity (Cameron et al., 2019; Manning et al., 2015). Identifying the role of environmental factors in determining cross-taxa biodiversity associations can be challenging, with a vast array of potential environmental drivers that can each have direct, indirect, linear, non-linear, synergistic, and/or antagonistic effects on the species within an ecosystem. The scale of the environmental gradient also impacts the relative role of the environment in driving biodiversity change, as larger environmental gradients can have greater, or even non-linear, effects (Chase et al., 2018). Experimental manipulations, such as those established to evaluate the impact of plant diversity upon the biodiversity of the other groups, can yield useful information on the potential role of biodiversity interactions under relatively constant environmental conditions. Manipulative studies have found positive relations between plant diversity and the diversity of some other organisms, particularly for aboveground herbivores but less so for soil microbial groups (Cline et al., 2017; Dassen et al., 2017; Lind et al., 2015; Scherber et al., 2010; Weisser et al., 2017). However, by necessity empirical studies are limited in scope, covering a limited set of environmental conditions and plant species combinations. Furthermore, too few studies consider complex, non-linear environmental effects when assessing relationships between diversity or composition of disparate taxonomic groups.

Using a large multi-trophic field dataset spanning all terrestrial habitats in Wales, UK, here we attempt to unpick the relationships between a breadth of above and below-ground taxa. Specifically, we explore associations between plants, birds, pollinators and belowground soil bacteria, fungi (total fungi and arbuscular mycorrhizal fungi), heterotrophic protists and animals, both before and after accounting for linear or complex non-linear environmental effects. We assess the relationships between taxa across three axes of biodiversity, addressing aspects of alpha diversity, community turnover and taxon co-occurrence. We

investigate (1) correlation between the alpha diversity of different groups using Bayesian hierarchical regression models; (2) whether turnover between sites is correlated across different groups using ecological distance analysis; and (3) the co-occurrence of individual taxa across sites, irrespective of higher groupings. In each analysis, we assess the contribution of linear or complex non-linear environmental effects. We hypothesise that (a) many inter- and intra-group relationships are readily explained by environmental variation, especially when complex, non-linear effects are considered; (b) those relationships remaining after accounting for environmental variation are more likely to be between groups and taxa that are known to have biotic interactions (e.g. nectar producing plants and pollinators, or forbs and soil fungi); and (c) that accounting for environmental effects would be more likely to cause associations between the alpha diversity of different groups to disappear compared to associations between community turnover or co-occurrence of individual taxa.

Methods

Field measurement programme

The data was collected as part of the Glastir Monitoring and Evaluation Programme (GMEP) field measurement program in Wales (Emmett et al., 2017; Wood et al., 2021). In total, 300 individual 1 km squares were randomly selected from within land classification strata in proportion to their extent across Wales in order to be representative of the range of habitat types across Wales or targeted to areas with high potential for agri-environment scheme uptake (Wood et al., 2021). Sampling occurred over a five-month period across each of the summers of 2013 to 2016; each square was only surveyed once over the four years with different squares being surveyed each year. Every square was subjected to a habitat survey,

bird survey, two pollinator survey transects and multiple plant survey plots. A representative schematic of the survey layout within a 1km square is shown in Supplementary Fig. 1. For each square there were up to five 200 m^2 square plant survey plots that also had soil samples taken. Soil samples were analysed for a variety of soil physicochemical properties, including pH in 1:2.5 $CaCl_2$ suspension, total carbon, nitrogen and phosphorus, bulk density, water content, water repellency and electrical conductivity in 1:2.5 distilled water suspension. The data and full methods are available on the Environmental Information Data Centre (EIDC) (Robinson et al., 2019). Within the first two years of the survey soil samples were taken for microbial and eukaryotic community composition analysis from three of the 200 m^2 plots, randomly selected per each 1 km square.

The habitats of each square were mapped and each plot was assigned to a habitat according to the UK Joint Nature Conservation Committee criteria (Jackson, 2000). The plot level measurements of microbial diversity were derived from 31% improved grassland; 23% neutral grassland; 12% acid grassland; 7% broadleaved woodland; 7% coniferous woodland; 7% bog; 4% arable; 3% dwarf shrub heath; 3% fen, marsh and swamp; and 2% bracken. Precipitation was the annual average rainfall and temperature was the annual average daily temperature for 1981-2010 calculated on a 1 km grid. All climate data came from the Met Office © Crown copyright 2019 (Met Office et al., 2019).

Biological data

Plant survey

Vegetation surveys were conducted for multiple plots per square, with the data being publicly available on the EIDC (Smart et al., 2020). Overall, there were ten types of sampling plots in total. Some plots were targeted to specific landscape features (e.g. hedges), meaning the total number and proportion of plot types varies across squares. The total number of vascular plant species recorded across the entire 1 km square was used as plant species richness for the square level analyses. Due to the differences in sampling effort across the squares, rarefaction curves were constructed to confirm that the number of plant species found had reached saturation. To assess whether different subsets of plants had greater associations with the diversity of other groups, the number of forb species within each plot was calculated (for associations with the soil communities), and for the aboveground communities the richness of plant species that were important to the diet of lowland birds, butterfly larvae and nectar provision was also calculated (Baude et al., 2016; Smart et al., 2000).

Bird and pollinator surveys

Birds were surveyed based on four morning visits to each square, equally spaced through mid-March to mid-July. Surveyors walked a route that passed within 50 m of all parts of each survey square, varying the start point for each survey in order to visit all parts of the square at least once before 08:00 h. All bird species seen and heard were recorded within each visit, and the total number of bird species recorded across all visits per square was used to calculate total bird species richness. Pollinator surveys were split into two independent parts: two 1 km transect routes separated by at least 500 m and where

possible 250 m from the edge of the 1 km square; and a 20-minute timed search in a 150 m^2

flower rich area within the 1 km square. The total number of butterfly species and bee

groups recorded per square were used to calculate richness. Pollinator surveys were

managed by Butterfly Conservation and bird surveys managed by the British Trust for

Ornithology, and both datasets are publicly available on the EIDC (Botham et al., 2020;

Siriwardena et al., 2020). Further details on survey methods are available within the data

documentation upon the EIDC, and also within Wood et al. (2021).

Soil microbial diversity and community composition

Soil samples for microbial biodiversity analyses were taken using a gouge auger at 5 points

around the physicochemical soil core location down to 15 cm, and then bulking together the

samples. DNA was extracted from the combined, homogenised samples using mechanical

lysis following homogenisation in triplicate from 0.25 g of soil per sample. The 16S (V4), ITS1

and 18S regions of the rRNA marker gene were targeted for amplicon sequencing to analyse

the bacterial, fungal, and general eukaryotic diversity respectively. Diversity of arbuscular

mycorrhizal fungi (AM fungi) was assessed separately using 18S data due to poor resolution

with the ITS region (Berruti et al., 2017). For a full description of the methods used see

George et al. (2019). Sequences with associated sample metadata have been uploaded to

The European Nucleotide Archive with the following primary accession codes: PRJEB27883

(16S), PRJEB28028 (ITS1) and PRJEB28067 (18S).

Amplicon Sequence Variants (ASVs) were identified from the Illumina output using the

DADA2 algorithm in R (Callahan et al., 2016). First, Illumina adapters were trimmed from

sequences using Cutadapt (Martin, 2011). Then, the DADA2 package was used to filter all

16S, 18S and ITS1 reads to be truncated after the first instance of a quality score of 2, to

remove all Ns and to have no more than 2 errors. Based on the examination of the change in quality scores with read length for a random subset of samples 16S reads were truncated to 200 bases, 18S reads truncated to 250 bases and ITS reads fewer than 100 bases in length removed. The DADA2 algorithm was then used to de-noise and merge the paired reads at standard settings. Sequence tables were constructed from the resultant ASVs and chimeric sequences were removed using default settings. Taxonomy was assigned using the DECIPHER R package (Wright, 2016), with both 16S and 18S reads being matched to SILVA r138 (Glöckner et al., 2017) and ITS being matched to UNITE v2019 (Nilsson et al., 2019). Non-bacterial ASVs were removed from the 16S data, and non-fungal ASVs from the ITS data. All ASVs appearing in the negative controls were removed from the analysis.

To account for the differences in read depth between samples, rarefaction was used as it has been shown to preserve the microbial relationships with biological origin (Weiss et al., 2017). The fungal data was rarefied to 10000 reads, bacteria to 20000 reads, and 18S eukaryotes to 20000 reads. Samples below this threshold were discarded, resulting in 437 fungal (ITS) measurements, 437 bacterial measurements and 438 eukaryote (18S) measurements. Eukaryote data was separated in to three sections based on higher order identification in the 18S dataset: AM fungi (phylum Glomeromycota), heterotrophic protists (Cercozoa, Ciliophora and Amoebozoa) and soil animals. Richness was calculated after rarefaction, which was repeated 100 times and the average result used in further analysis. For analysis of community composition the average of 20 rarefaction repeats was used.

Statistics

Alpha diversity

All statistical analyses were performed within R version 4.3.0 (R Core Team, 2023). For analysis (1), assessing correlations between alpha diversity of different groups, we tested for effects of environmental variables using Bayesian multivariate hierarchical modelling within the brms package in R (Bürkner, 2017). Group richness (for the plot-level model with soil groups and plant community) or group Shannon diversity (for the 1km square-level plant, bird and pollinator model) were modelled in a multivariate framework as either a function of a) solely the hierarchical spatial grouping, b) a linear function of abiotic variables, or c) as a non-linear function of abiotic variables. For the plot-level model with linear abiotic drivers we included soil pH, total soil carbon, water content, total phosphorus and soil water repellency as predictors, with a unit normal prior on all effects. When including non-linear environmental effects, we assumed a sigmoidal relationship between microbial richness and pH due to the pre-existing literature on the non-linear effects of soil pH on soil biochemical properties. This sigmoidal model took the form shown in equation 1, where parameters α (upper asymptote), β (growth rate), γ (x value at maximum growth), and δ (intercept) were fit using weakly informative priors.

$$y = \frac{\alpha}{1 + \exp\left(-\beta(x - \gamma)\right)} + \delta \qquad (1)$$

For the square-level model with linear abiotic drivers we included annual temperature and precipitation as predictors, again with a unit normal prior on all effects. To investigate a role for complex environmental effects, the plant, bird and pollinator model with linear environmental effects was compared to a model where the exponentiated Shannon diversity values were modelled as a function of interacting temperature and precipitation.

The square identity was included as a group-level effect on the intercept (δ) in order to account for the spatial element of the data.

To evaluate whether associations between the alpha diversity of the different groups remained after accounting for environment the residual correlations between groups were extracted from the models. All response variables were modelled as normal, to allow for both the rarefaction process for microbial groups detailed above resulting in non-integer responses and the inclusion of residual correlations within the model structure. Satisfactory recovery of the data was confirmed using graphical posterior predictive checks. No correlations between the group-level effects were included within the model.

Community turnover

For analysis (2), assessing the correlations in turnover between sites across different groups, we carried out an ordination analysis using non-metric dimensional scaling (NMDS). Distance calculation and NMDS were performed using the vegan package (Oksanen et al., 2020). Binary Jaccard distance was used for plant, bird and butterfly species composition and Bray-Curtis distance used for bee and hoverfly groups due to their lower richness. All soil microbial groups were compared based upon Bray-Curtis distance. Direct correlation coefficients between the turnover of the different groups were calculated using Procrustes analysis upon the ordination scores for each group. To unpick the relative contribution of shared environmental drivers, interaction with the plant community and spatial autocorrelation that could be driving changes in the main axes of variation in the communities the variance in the ordination scores of each group was partitioned using the varpart function within the vegan package. The response variables were the NMDS scores of each group based on Jaccard distance for aboveground groups and Bray-Curtis distance for

belowground groups. The NMDS for the plot-level analysis were mostly performed using 3 dimensions, and had stress values of 0.067 for bacteria, 0.117 for fungi, 0.095 for heterotrophic protists, 0.192 for soil animals, and 0.076 for plants (over 4 dimensions). No stable NMDS solution was found for AM fungi even after varying the number of dimensions, ecological distance used, and convergence criteria so they were omitted from this analysis. The NMDS for the square level analysis were mostly performed using 4 dimensions and had stress values of 0.098 for plants, 0.089 for birds, 0.139 for bees and hoverflies (3 dimensions), and 0.139 for butterflies. The predictors within the variance partitioning were climate (represented by temperature and precipitation), spatial distance (represented by principal coordinates of neighbour matrices), plant community composition (represented by the NMDS scores). For the plot level data only the first four dimensions of a PCA upon soil physicochemical properties (pH, carbon, nitrogen, total phosphorus, bulk density, electrical conductivity, water content, water repellency) were also used as predictors. These dimensions explained a total of 88% of the variance in the measured soil physicochemical properties.

Taxon-level relationships

For analysis (3), assessing co-occurrence of taxa irrespective of higher groupings, relationships were evaluated using joint species distribution models to identify whether co-occurrences could be explained by shared responses to environmental drivers and spatial autocorrelation. Joint species distribution models were fitted using the sjSDM package in R (Pichler & Hartig, 2021). Only common taxa were included within the analysis to reduce the number of false positive associations (Weiss et al., 2016), which we define as being present in 25% of sites for bacteria, heterotrophic protists, fungi and soil animals and 10% for plants, birds and pollinators. In total there were 2309 bacterial taxa, 236 heterotrophic protists, 190

fungal taxa, 22 animals and 28 plants included in the plot-level belowground analysis while the square-level aboveground analysis included 348 plant species, 101 bird species, 22 butterfly species, 4 bee groups and 4 hoverfly groups. All models had the spatial relationships between sites modelled as a linear function of the spatial eigenvectors, all taxa converted to presence/absence and fit with a binomial model with probit link, and all other parameters set to their defaults. The responses to environmental gradients were either excluded (intercept-only model) or modelled as either linear responses or deep neural networks. Within the plot-level model (i.e. including microbial groups and plant measurements from individual plots) the environmental predictors were soil pH, organic matter, carbon, nitrogen, available phosphorus (Olsen-P), conductivity, water repellency, moisture content and bulk density. Within the aboveground model (i.e. including plants at the square level, birds and pollinating insects) the environmental predictors were average annual precipitation (1981-2010), daily temperature (1981-2010) and elevation.

Results

Alpha diversity

We found that bacterial, fungal, heterotrophic protistan, bird, bee and butterfly richness were all positively correlated with plant diversity. Within the soil groups, AM fungi showed the strongest correlation with plants, whereas bacteria, general fungi and heterotrophic protists all had lower correlations, and no correlation with animal richness (Table 1, upper triangle). All soil groups were positively correlated with each other; however animal richness was only weakly associated with the other groups. Soil microbial richness was more strongly positively correlated to the richness of forb plant species as opposed to overall plant richness. Microbial richness was also negatively correlated to woody plant richness,

particularly in AM fungi (ρ = -0.65) and heterotrophic protists (ρ = -0.51). Plant diversity in the 1 km squares positively correlated most strongly with bird species richness (ρ = 0.50), followed by butterfly species richness (ρ = 0.43), then bee group and hoverfly group (ρ ~ 0.35) richness. Limiting the plant community to those species known to be important to the different aboveground animal groups through food provision resulted in an increase in correlation coefficients of 0.08 for birds but not for the other groups. The different pollinator groups were positively associated with each other, with butterflies and bees associated more strongly with each other than hoverflies with the other two groups (ρ = 0.56 compared to ρ 0.42-0.44). Bird species richness was positively correlated with butterfly and bee richness (ρ ~ 0.5). Average soil microbial diversity per 1 km square was also positively correlated with bird, butterfly and hoverfly richness, with correlations ranging from ~0.35 to ~0.55.

Table 1: The correlations between the different microbial groups and the plant community, the upper triangle shows the Spearman rank correlation coefficient for alpha diversity (richness) and the lower triangle shows the Procrustes correlation coefficient for compositional turnover (Bray-Curtis distance).

Plant	0.22	0.21	0.19	-0.02	0.40	0.81
0.54	**Fungi**	0.51	0.78	0.20	0.53	0.42
0.50	0.71	**Bacteria**	0.56	0.07	0.32	0.39
0.56	0.72	0.78	**Hetero. protists**	0.13	0.54	0.45
0.41	0.45	0.50	0.52	**Soil animals**	-0.05	-0.16
0.43	0.54	0.52	0.58	0.35	**AM Fungi**	0.57
0.55	0.47	0.46	0.51	0.32	0.40	**Forbs**

The effects of environmental factors upon alpha diversity relationships

Accounting for environmental gradients reduced the magnitude of the correlations between the diversity of the different taxonomic groups both above and belowground (Fig. 1). Aboveground, only the correlation between plant diversity and bird diversity and the correlation between bee diversity and butterfly diversity remained after accounting for the temperature and rainfall gradients (Fig. 1e-f). It might be expected that the diversity of plant groups that are important to the diets of the different aboveground animal groups would show stronger relationships with these groups, and this was tested by comparing direct models of plant richness as a predictor of bird, butterfly and bee richness versus models with the richness of plants important in the diet of lowland birds, butterfly larval food plants and nectar producing plants respectively. However, we found no improvement in model predictive performance compared to models with overall plant richness.

Figure 1: The residual correlations between the diversity of the different groups if no abiotic drivers are taken into account (a, d), if abiotic drivers are included as linear effects only (b, e), and if non-linear effects are included (c, f) are included. Blue lines indicate positive residual correlations, red lines negative. The width of each line is proportional to the estimated residual correlation, lines are solid if the 95% interval did not include zero or dashed if the 95% interval included zero but the 60% interval did not. Residual correlations where the 60% interval included zero are not shown in the figure but are included within the model. For the full model summary see Supplementary tables 1-6.

Belowground, the richness of the soil groups did maintain correlations with each other even after accounting for changes in soil properties, and in the case of soil animal richness the residual correlation with the other groups actually increased once a variety of soil properties including soil carbon were accounted for (Fig. 1a-b, Supplementary tables 1-2). For all other soil groups, the model with a sigmoidal pH gradient accounted for more residual correlation (Fig. 1c) than the linear models of multiple soil properties (Fig. 1b). All soil microbial groups, in addition to plant richness, showed a good fit of the sigmoidal pH model to the data while

soil animal richness showed a linear decrease with increasing pH (Fig. 2). Accounting for soil

properties removed the residual correlation between plant richness and soil richness for all

soil groups other than AM fungi, which retained a positive relationship with plant richness.

Figure 2: Predicted effect of soil pH on soil and plant diversity in the multivariate model with a sigmoidal pH

effect. Plots show the actual values as black dots and the conditional pH effect as a blue line with the

95% interval of the effect as a shaded grey ribbon.

Community turnover

Plant community turnover was positively associated with fungal, bacterial, heterotrophic

protistan, bird and butterfly community turnover. Bacteria, heterotrophic protists and

general fungi showed a slightly stronger correlation with plant community turnover than

AM fungi did (Table 1, lower triangle); all were significant at p = 0.001. Turnover in

heterotrophic protistan communities strongly tracked turnover in bacterial communities,

with a Procrustes correlation statistic of 0.78. Fungal community turnover also correlated with bacterial turnover, with a Procrustes correlation statistic of 0.71. Bird and butterfly community turnover were more strongly related to plant community turnover than the link between bee, hoverfly and plant turnover. This was reflected by both the spearman rank correlations and the Procrustes correlation statistic (0.73, 0.59, 0.23 and 0.32 for bird, butterflies, bees and hoverflies respectively). However, all the Procrustes analyses were found to be significant at p = 0.001.

The effects of environmental factors upon community turnover relationships

Plant community turnover explained unique variance in the bacterial, fungal and heterotrophic protist communities even after accounting for spatial and environmental factors (Fig. 3a). Plant community turnover at the plot scale was partially explained by soil, climate and spatial factors but had considerable unexplained variance (71%). Climate was initially included but did not explain any unique variance in turnover of the soil microbial communities and so was omitted from this analysis. The pattern of variance explained was similar across the soil groups, with the soil physicochemical properties and plant community jointly explaining more variation than the distance between samples. However, there was a large proportion of variation in fungi, bacteria, animals and heterotrophic protists that was explained by soil, plant and spatial factors together.

Figure 3: The properties that explain the variation in community turnover at the plot level (a) and the square level (b). The height of each coloured bar section represents the proportion of variation explained by the category, with the "Shared" category representing the variation explained by all factors together in addition to the variation explained by each single or two-part combination of factors.

Plant community turnover explained some of the variance in bird, butterfly and bee and hoverfly community turnover (Fig. 3b). Plant community turnover at the 1 km square level was largely explained by variation in precipitation and temperature (38%), with some effect of the distance between squares interacting with climate (19%). Turnover in the bird community was explained comparatively well by the plant community and the climate (28% without any spatial element), whilst also incorporating a spatial interaction between the two. The pollinator communities were explained relatively poorly; plant community turnover, climate and spatial factors did appear to be important together but overall ~60% of variation was unexplained. Precipitation and temperature were less important to bees and hoverflies than they are to the other groups, with precipitation and temperature in general being the least important factor for the aboveground animal groups but much more

important for plants. Spatial factors appeared to be relatively more important for the animal groups than they were for the plants.

Taxon-level relationships

Figure 4 The density distributions of taxon correlations from the belowground/plot-level (a) and aboveground/square level (b) joint species distribution models that took no account of environment (No env), linear functions of environment only (Linear), or that used deep neural nets to allow for non-linear effects of environment (DNN). Butterflies, bees and hoverflies are combined into a pollinator category for graphical simplicity, as are general fungi and AM fungi.

Residual co-occurrence relationships between common below- and aboveground taxa were identified by joint species distribution modelling both with and without accounting for shared responses to environmental variables. In all cases the average magnitude of taxon correlations was decreased once environmental gradients were accounted for, with non-linear environmental gradients removing more apparent species correlations than linear

environmental gradients (Fig. 4). The effect of incorporating environmental conditions was greater upon apparent negative associations between taxa, with proportionally more negative associations lost after changes in environmental conditions were accounted for. Within the soil groups more positive associations were retained after accounting for environmental conditions, with the mode of the correlation distribution remaining above zero compared to it being centred on zero in all plant-related and aboveground correlation distributions. Soil groups showed on average differing levels of association with differing plant growth forms, with forbs showing more positive associations with soil microbial groups than grasses. However, accounting for non-linear environmental conditions removed the strength of the forb relationship in average correlation magnitude between plant growth forms (Supplementary Fig. 2a). Only one woody species (*Vaccinium myrtillus*) appeared in enough plots to be included in this analysis, and so those results are unlikely to be representative of all woody species. There was very little difference between the correlation strength of plants that are, or are not, food sources for birds and pollinators across the different models, with only a slight tendency to have a higher average magnitude of correlation for food source plants in the no environment and non-linear models but not in the non-linear model (Supplementary Fig. 2b).

Despite the overall trend towards lower average magnitude of species correlations after accounting for environmental conditions there was a small proportion of species correlations that actually increased in magnitude after accounting for environmental gradients. Around 7% of correlations increased above- and belowground by over 0.05 units in the non-linear environmental model compared to the model with only spatial and biotic components. The interactions between soil animal and other soil taxa showed proportionally more correlations that increased, with 17% of correlations increasing by

>0.05, while the interactions between the plant and soil taxa showed proportionally more decreases in magnitude. The relative importance of biotic factors, environmental factors, and spatial factors varied considerably by taxa, although on average biotic factors appeared more important belowground than aboveground while spatial factors were more important aboveground than belowground (Supplementary Figs 3 and 4).

Discussion

Consistent with our hypothesis we find that many alpha diversity relationships dissolve after accounting for abiotic effects, with non-linear environmental gradients clearly affecting belowground alpha diversity relationships. Our results indicate that the positive correlation of plant and soil microbial diversity is driven by changes in the soil environment and are in agreement with previous work showing the importance of soil pH and fertility in jointly influencing plant and microbial diversity (Goberna et al., 2016; Tedersoo et al., 2016; Yashiro et al., 2018; Yuan et al., 2017). Importantly, accounting for the influence of soil pH upon microbial diversity in a non-linear way led to the loss of a plant ~ microbe diversity relationship, with the exception of AM fungi. AM fungi are more dependent upon plant inputs and are capable of growing through large volumes of soil, meaning that they would be expected to show stronger relations with the plant community (Hiiesalu et al., 2014; Nguyen et al., 2016). Our results also clearly indicate a strong linkage between bacterial, fungal and heterotrophic protistan communities that could represent the dominant role of trophic dynamics over edaphic properties and plant inputs in structuring those communities.

The diversity of the pollinator groups showed no residual correlation with plant communities after accounting for environmental gradients, and in contrast to our hypothesis we found no evidence of stronger links between foodplants and the aboveground animals that feed on them compared to overall plant diversity. The lack of relationship between pollinator and plant diversity was unexpected due to the role of plants in providing food resources and niche space, and we found no evidence of stronger associations in what should be trophically interacting species at either the alpha diversity or the taxon-level. However, previous research has shown variable relationships between plant and pollinator diversity, potentially related to vegetation structure and the greater importance of floral abundance and seasonality compared to plant or food-plant diversity (Alison et al., 2022; Berg et al., 2019; Lowe et al., 2022; Potts et al., 2009). In contrast, increasing bird diversity in conjunction with plant diversity was only partially confounded by the climate gradient, indicating that plant diversity could potentially have a direct impact upon bird diversity. This could be due to the provision of a greater variety of food sources, through increased landscape heterogeneity or potentially through the influence of an unidentified confounding variable such as anthropogenic activity (Le Provost et al., 2021; Stein & Kreft, 2014).

We found that the composition of above- and belowground communities was influenced by plant community composition even after accounting for environmental gradients, in contrast to the alpha diversity results. The proportion of variance in community turnover explained by plants was much higher in the bacterial, fungal and heterotrophic protist communities than in the bird, butterfly, bee and hoverfly communities. The finding that plant community turnover explains bacterial, fungal and heterotrophic protistan community turnover is consistent with previous findings at the field (Leff et al., 2018), regional

(Barberán et al., 2015; Chen et al., 2017; T. Yang et al., 2017), and global scales (Delgado-Baquerizo et al., 2018; Prober et al., 2015). The higher proportion of explained variance in bird communities compared to butterfly, bee and hoverfly communities was unexpected as pollinators have in general been found to be closely related to plant composition (Hofmann et al., 2017; Kearns & Oliveras, 2009; Weisser et al., 2017). Overall, the majority of the variation in butterfly, bee and hoverfly community turnover was unexplained by climate, plant communities or spatial factors. This may be due to anthropogenic influences or other factors not incorporated into our model being important in driving pollinator community assembly.

The closer linkages of the soil groups to each other compared to the plant linkages above- and belowground were apparent in both the overall diversity and species-level results. In particular, the soil animal community showed stronger linkages with other soil groups after accounting for environmental gradients. Soil animal groups are strongly influenced by soil carbon and other environmental factors (Caruso et al., 2019; George et al., 2017; Lavelle et al., 2022), and it appears that once these are accounted for the secondary role of biotic associations in driving community assembly becomes clearer. Some of these identified linkages between the taxa within and between groups may represent shared response to environmental or geographic drivers not included within our analysis, while some may instead represent true interactions across domains (Blanchet et al., 2020; Carr et al., 2019; Seaton et al., 2020).

The different taxonomic groups within our analysis operate upon different spatial and temporal scales, and this appears to have affected the relative influence of spatial factors upon biological communities that we have found (Chase et al., 2018). In our results we

found that spatial distance appeared to have a greater influence upon the species composition of the aboveground communities measured at the 1km square level. Previous work has shown that the scale at which biodiversity is measured, impacts the strength of the correlation found, with 10 km^2 areas showing the strongest relationship for aboveground diversity (Wolters et al., 2006). However, the life history traits of the community in question should be expected to impact the scale of interest, and within our dataset we have organisms that are whose mobility is limited to very short distances such as soil bacteria (P. Yang & van Elsas, 2018) ranging up to bird species that can travel vast distances. Therefore, the spatial factors we measured may not have been at the appropriate scale to influence the soil microbial communities at the plot level. It is also worth considering the role of the spatio-temporal scale of survey in determining the range of the environmental drivers of biodiversity (Chase et al., 2018). Larger areas contain a wider range of environmental conditions where different parts of the range could have contrasting effects on the biosphere. Our survey covers a range of environmental conditions that is representative of the Welsh landscape, however it only contains a limited sampling of feasible potential environmental conditions such as intense agricultural landscapes more common in other areas of Europe where we might expect different influences upon soil and aboveground communities (Billeter et al., 2008; Gossner et al., 2016).

We found that accounting for non-linear environmental effects had a greater influence on inferred biotic correlations at both the overall diversity and species level compared to only incorporating linear environmental effects. This emphasises the importance of considering nonlinear effects to prevent underestimation of the impacts of environmental gradients as well as increase the applicability of the results outside the environmental range of each study, particularly within soil communities (Austin, 2007). In contrast, including only linear

effects of the environment proved sufficient to remove most of the aboveground diversity and positive species associations. However, it is possible that other environmental drivers or landscapes may show a greater role of non-linear environmental effects in driving plant, pollinator, and bird community dynamics. It is also possible that changes in land use intensity or other factors could change the very nature of the relationships between the diversity and composition of differing components of the ecosystem (Felipe-Lucia et al., 2020; Le Provost et al., 2021). Within this work we take a step towards disentangling the relative cross domain biotic and environmental associations governing community assembly across the terrestrial-soil habitat boundary. Importantly, our results provide insight into the importance of non-linear environmental effects upon ecological communities across a wide variety of entwined habitats and environmental conditions and provide potential future avenues of research in evaluating whole-ecosystem state and change, in relation to complex environmental drivers.

Acknowledgements

We thank all the GMEP team who contributed to collecting all the data. The research was funded by the Welsh Government through the Glastir Monitoring and Evaluation Program (GMEP) with underpinning funding from the Natural Environment Research Council award number NE/R016429/1 as part of the UKScaPE programme delivering National Capability. Contract reference: C147/2010/11. NERC/Centre for Ecology & Hydrology (CEH Projects: NEC04780/NEC05371/NEC05782). F.M.S. and P.B.L.G were supported by the award of studentship grants by the NERC and BBSRC funded STARS (Soils Training And Research Studentships) Centre for Doctoral Training and Research Programme; a consortium

consisting of Bangor University, British Geological Survey, Centre for Ecology and Hydrology, Cranfield University, James Hutton Institute, Lancaster University, Rothamsted Research and the University of Nottingham. Grant Ref: NE/M009106/1.

Author contributions

Seaton, George, Creer, Jones, and Robinson conceived the ideas and Seaton designed the statistical methodology with input from Alison, George and Smart; Emmett and Robinson led the collection of the data; Seaton analysed the data and led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

Data availability

The data that support the findings of this study are openly available, with the soil physicochemical data at DOI: 10.5285/0fa51dc6-1537-4ad6-9d06-e476c137ed09, the vegetation data at DOI: 10.5285/71d3619c-4439-4c9e-84dc-3ca873d7f5cc, the pollinator data at DOI: 10.5285/3c8f4e46-bf6c-4ea1-9340-571fede26ee8, the bird data at DOI: 10.5285/31da0a94-62be-47b3-b76e-4bdef3037360 and the soil microbial sequences data in the European Nucleotide Archive with primary accession codes: PRJEB27883 (16S), PRJEB28028 (ITS1) and PRJEB28067 (18S).

References

Alison, J., Botham, M., Maskell, L. C., Garbutt, A., Seaton, F. M., Skates, J., Smart, S. M., Thomas, A. R. C., Tordoff, G., Williams, B. L., Wood, C. M., & Emmett, B. A. (2022). Woodland, cropland and hedgerows promote pollinator abundance in intensive grassland landscapes, with saturating benefits of flower cover. *Journal of Applied Ecology*, *59*(1), 342–354. https://doi.org/10.1111/1365-2664.14058

Austin, M. (2007). Species distribution models and ecological theory: A critical assessment and some possible new approaches. *Ecological Modelling*, *200*(1), 1–19. https://doi.org/10.1016/j.ecolmodel.2006.07.005

Barberán, A., McGuire, K. L., Wolf, J. A., Jones, F. A., Wright, S. J., Turner, B. L., Essene, A., Hubbell, S. P., Faircloth, B. C., & Fierer, N. (2015). Relating belowground microbial composition to the taxonomic, phylogenetic, and functional trait distributions of trees in a tropical forest. *Ecology Letters*, *18*(12), 1397–1405. https://doi.org/10.1111/ele.12536

Baude, M., Kunin, W. E., Boatman, N. D., Conyers, S., Davies, N., Gillespie, M. A. K., Morton, R. D., Smart, S. M., & Memmott, J. (2016). Historical nectar assessment reveals the fall and rise of floral resources in Britain. *Nature*, *530*(7588), 85–88. https://doi.org/10.1038/nature16532

Berg, Å., Cronvall, E., Eriksson, Å., Glimskär, A., Hiron, M., Knape, J., Pärt, T., Wissman, J., Żmihorski, M., & Öckinger, E. (2019). Assessing agri-environmental schemes for semi-natural grasslands during a 5-year period: Can we see positive effects for vascular plants and pollinators? *Biodiversity and Conservation*. https://doi.org/10.1007/s10531-019-01861-1

Berruti, A., Desirò, A., Visentin, S., Zecca, O., & Bonfante, P. (2017). ITS fungal barcoding

    primers versus 18S AMF-specific primers reveal similar AMF-based diversity patterns

    in roots and soils of three mountain vineyards. *Environmental Microbiology Reports*,

    9(5), 658–667. https://doi.org/10.1111/1758-2229.12574

Billeter, R., Liira, J., Bailey, D., Bugter, R., Arens, P., Augenstein, I., Aviron, S., Baudry, J.,

    Bukacek, R., Burel, F., Cerny, M., De Blust, G., De Cock, R., Diekötter, T., Dietz, H.,

    Dirksen, J., Dormann, C., Durka, W., Frenzel, M., … Edwards, P. J. (2008). Indicators

    for biodiversity in agricultural landscapes: A pan-European study. *Journal of Applied*

    Ecology, *45*(1), 141–150. https://doi.org/10.1111/j.1365-2664.2007.01393.x

Blanchet, F. G., Cazelles, K., & Gravel, D. (2020). Co-occurrence is not evidence of ecological

    interactions. *Ecology Letters*, ele.13525. https://doi.org/10.1111/ele.13525

Botham, M., Peyton, J., Tordoff, G., Edwards, M., Alison, J., Astbury, S., Burden, A., Emmett,

    B. A., Garbutt, A., Henrys, P., Owen, A., Smart, S. M., Sharps, K., Williams, B., &

    Wood, C. M. (2020). *Insect pollinator and flower data from the Glastir Monitoring*

    and Evaluation Programme, Wales, 2013-2016. NERC Environmental Information

    Data Centre. https://doi.org/10.5285/3c8f4e46-bf6c-4ea1-9340-571fede26ee8

Bürkner, P.-C. (2017). brms: An R package for Bayesian multilevel models using Stan. *Journal*

    of Statistical Software, *80*(1), 1–28. https://doi.org/10.18637/jss.v080.i01

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P.

    (2016). DADA2: High-resolution sample inference from Illumina amplicon data.

    Nature Methods, *13*(7), 581–583. https://doi.org/10.1038/nmeth.3869

Cameron, E. K., Martins, I. S., Lavelle, P., Mathieu, J., Tedersoo, L., Bahram, M., Gottschall,

    F., Guerra, C. A., Hines, J., Patoine, G., Siebert, J., Winter, M., Cesarz, S., Ferlian, O.,

    Kreft, H., Lovejoy, T. E., Montanarella, L., Orgiazzi, A., Pereira, H. M., … Eisenhauer, N.

(2019). Global mismatches in aboveground and belowground biodiversity.

Conservation Biology, *33*(5), 1187–1192. https://doi.org/10.1111/cobi.13311

Carr, A., Diener, C., Baliga, N. S., & Gibbons, S. M. (2019). Use and abuse of correlation

analyses in microbial ecology. *The ISME Journal*, *13*(11), 2647–2655.

https://doi.org/10.1038/s41396-019-0459-z

Caruso, T., Schaefer, I., Monson, F., & Keith, A. M. (2019). Oribatid mites show how climate

and latitudinal gradients in organic matter can drive large-scale biodiversity patterns

of soil communities. *Journal of Biogeography*, *46*(3), 611–620.

https://doi.org/10.1111/jbi.13501

Chase, J. M., McGill, B. J., McGlinn, D. J., May, F., Blowes, S. A., Xiao, X., Knight, T. M.,

Purschke, O., & Gotelli, N. J. (2018). Embracing scale-dependence to achieve a

deeper understanding of biodiversity and its change across communities. *Ecology*

Letters, *21*(11), 1737–1751. https://doi.org/10.1111/ele.13151

Chen, Y.-L., Xu, T.-L., Veresoglou, S. D., Hu, H.-W., Hao, Z.-P., Hu, Y.-J., Liu, L., Deng, Y., Rillig,

M. C., & Chen, B.-D. (2017). Plant diversity represents the prevalent determinant of

soil fungal community structure across temperate grasslands in northern China. *Soil*

Biology and Biochemistry, *110*, 12–21. https://doi.org/10.1016/j.soilbio.2017.02.015

Cline, L. C., Hobbie, S. E., Madritch, M., Buyarski, C. R., Tilman, D., & Cavender-Bares, J. M.

(2017). Resource availability underlies the plant-fungal diversity relationship in a

grassland ecosystem. *Ecology*, *99*(1), 204–216. https://doi.org/10.1002/ecy.2075

Dassen, S., Cortois, R., Martens, H., de Hollander, M., Kowalchuk, G. A., van der Putten, W.

H., & De Deyn, G. B. (2017). Differential responses of soil bacteria, fungi, archaea and

protists to plant species richness and plant functional group identity. *Molecular*

Ecology, *March*, 4085–4098. https://doi.org/10.1111/mec.14175

Delgado-Baquerizo, M., Fry, E. L., Eldridge, D. J., de Vries, F. T., Manning, P., Hamonts, K., Kattge, J., Boenisch, G., Singh, B. K., & Bardgett, R. D. (2018). Plant attributes explain the distribution of soil microbial communities in two contrasting regions of the globe. *New Phytologist*. https://doi.org/10.1111/nph.15161

Emmett, B. A., Abdalla, M., Anthony, S., Astbury, S., August, T., Barrett, G., Beckman, B., Biggs, J., Botham, M., Bradley, D., Brown, M., Carter, H., Chadwick, D., Cigna, F., Collier, R., Cooper, D., Cooper, J., Cosby, B. J., Creer, S., … Wright, S. (2017). *Glastir Monitoring & Evaluation Programme. Final Report to Welsh Government. Contract reference: C147/2010/11.* NERC/Centre for Ecology & Hydrology (CEH Projects: NEC04780/NEC05371/NEC05782) Short.

Felipe-Lucia, M. R., Soliveres, S., Penone, C., Fischer, M., Ammer, C., Boch, S., Boeddinghaus, R. S., Bonkowski, M., Buscot, F., Fiore-Donno, A. M., Frank, K., Goldmann, K., Gossner, M. M., Hölzel, N., Jochum, M., Kandeler, E., Klaus, V. H., Kleinebecker, T., Leimer, S., … Allan, E. (2020). Land-use intensity alters networks between biodiversity, ecosystem functions, and services. *Proceedings of the National Academy of Sciences of the United States of America*, *117*(45), 28140–28149. https://doi.org/10.1073/pnas.2016210117

George, P. B. L., Keith, A. M., Creer, S., Barrett, G. L., Lebron, I., Emmett, B. A., Robinson, D. A., & Jones, D. L. (2017). Evaluation of mesofauna communities as soil quality indicators in a national-level monitoring programme. *Soil Biology and Biochemistry*, *115*, 537–546. https://doi.org/10.1016/j.soilbio.2017.09.022

George, P. B. L., Lallias, D., Creer, S., Seaton, F. M., Kenny, J. G., Eccles, R. M., Griffiths, R. I., Lebron, I., Emmett, B. A., Robinson, D. A., & Jones, D. L. (2019). Divergent national-scale trends of microbial and animal biodiversity revealed across diverse temperate

soil ecosystems. *Nature Communications*, *10*(1), 1107.

https://doi.org/10.1038/s41467-019-09031-1

Glöckner, F. O., Yilmaz, P., Quast, C., Gerken, J., Beccati, A., Ciuprina, A., Bruns, G., Yarza, P.,

Peplies, J., Westram, R., & Ludwig, W. (2017). 25 years of serving the community

with ribosomal RNA gene reference databases and tools. *Journal of Biotechnology*,

261(February), 169–176. https://doi.org/10.1016/j.jbiotec.2017.06.1198

Goberna, M., Navarro-Cano, J. A., Verdú, M., Wardle, DA., Bardgett, RD., Klironomos, JN.,

Setälä, H., Putten, WH. van der, Wall, DH., Balvanera, P., Pfisterer, AB., Buchmann,

N., He, J.-S., Nakashizuka, T., Raffaelli, D., Schmid, B., Prober, SM., Tedersoo, L.,

Mouquet, N., … Zheng, Y.-M. (2016). Opposing phylogenetic diversity gradients of

plant and soil bacterial communities. *Proceedings Royal Society B*, *283*(1825),

20153003. https://doi.org/10.1098/rspb.2015.3003

Gossner, M. M., Lewinsohn, T. M., Kahl, T., Grassein, F., Boch, S., Prati, D., Birkhofer, K.,

Renner, S. C., Sikorski, J., Wubet, T., Arndt, H., Baumgartner, V., Blaser, S., Blüthgen,

N., Börschig, C., Buscot, F., Diekötter, T., Jorge, L. R., Jung, K., … Allan, E. (2016).

Land-use intensification causes multitrophic homogenization of grassland

communities. *Nature*, *540*(7632), 266–269. https://doi.org/10.1038/nature20575

Hiiesalu, I., Pärtel, M., Davison, J., Gerhold, P., Metsis, M., Moora, M., Öpik, M., Vasar, M.,

Zobel, M., & Wilson, S. D. (2014). Species richness of arbuscular mycorrhizal fungi:

Associations with grassland plant richness and biomass. *The New Phytologist*, *203*(1),

233–244. https://doi.org/10.1111/nph.12765

Hofmann, S., Everaars, J., Schweiger, O., Frenzel, M., Bannehr, L., & Cord, A. F. (2017).

Modelling patterns of pollinator species richness and diversity using satellite image

texture. *PLOS ONE*, *12*(10), e0185591.

https://doi.org/10.1371/journal.pone.0185591

IPBES. (2019). *Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.* (S. Díaz, J. Settele, E. S. Brondízio E.E., H. T. Ngo, M. Guèze, J. Agard, A. Arneth, P. Balvanera, K. A. Brauman, S. H. M. Butchart, K. M. A. Chan, L. A. Garibaldi, K. Ichii, J. Liu, S. M. Subramanian, G. F. Midgley, P. Miloslavich, Z. Molnár, D. Obura, … C. N. Zayas, Eds.). IPBES secretariat, Bonn, Germany.

Jackson, D. L. (2000). *Guidance on the interpretation of the Biodiversity Broad Habitat Classification (terrestrial and freshwater types): Definitions and the relationships with other classifications, JNCC Report 307* (p. 73). JNCC, Peterborough, UK. http://jncc.defra.gov.uk/page-2433

Kearns, C. A., & Oliveras, D. M. (2009). Environmental factors affecting bee diversity in urban and remote grassland plots in Boulder, Colorado. *Journal of Insect Conservation*, *13*(6), 655–665. https://doi.org/10.1007/s10841-009-9215-4

Lavelle, P., Mathieu, J., Spain, A., Brown, G., Fragoso, C., Lapied, E., De Aquino, A., Barois, I., Barrios, E., Barros, M. E., Bedano, J. C., Blanchart, E., Caulfield, M., Chagueza, Y., Dai, J., Decaëns, T., Dominguez, A., Dominguez, Y., Feijoo, A., … Zhang, C. (2022). Soil macroinvertebrate communities: A world-wide assessment. *Global Ecology and Biogeography*, *31*(7), 1261–1276. https://doi.org/10.1111/geb.13492

Le Provost, G., Thiele, J., Westphal, C., Penone, C., Allan, E., Neyret, M., van der Plas, F., Ayasse, M., Bardgett, R. D., Birkhofer, K., Boch, S., Bonkowski, M., Buscot, F., Feldhaar, H., Gaulton, R., Goldmann, K., Gossner, M. M., Klaus, V. H., Kleinebecker, T., … Manning, P. (2021). Contrasting responses of above- and belowground diversity

to multiple components of land-use intensity. *Nature Communications*, *12*(1), 3918.

https://doi.org/10.1038/s41467-021-23931-1

Leff, J. W., Bardgett, R. D., Wilkinson, A., Jackson, B. G., Pritchard, W. J., De Long, J. R.,

Oakley, S., Mason, K. E., Ostle, N. J., Johnson, D., Baggs, E. M., & Fierer, N. (2018).

Predicting the structure of soil communities from plant community taxonomy,

phylogeny, and traits. *The ISME Journal*, *12*(7), 1794–1805.

https://doi.org/10.1038/s41396-018-0089-x

Lind, E. M., Vincent, J. B., Weiblen, G. D., Cavender-Bares, J., & Borer, E. T. (2015). Trophic

phylogenetics: Evolutionary influences on body size, feeding, and species

associations in grassland arthropods. *Ecology*, *96*(4), 998–1009.

https://doi.org/10.1890/14-0784.1

Lowe, A., Jones, L., Brennan, G., Creer, S., & de Vere, N. (2022). Seasonal progression and

differences in major floral resource use by bees and hoverflies in a diverse

horticultural and agricultural landscape revealed by DNA metabarcoding. *Journal of

Applied Ecology*, *59*(6), 1484–1495. https://doi.org/10.1111/1365-2664.14144

Manning, P., Gossner, M. M., Bossdorf, O., Allan, E., Zhang, Y.-Y., Prati, D., Blüthgen, N.,

Boch, S., Böhm, S., Börschig, C., Hölzel, N., Jung, K., Klaus, V. H., Klein, A. M.,

Kleinebecker, T., Krauss, J., Lange, M., Müller, J., Pašalić, E., … Fischer, M. (2015).

Grassland management intensification weakens the associations among the

diversities of multiple plant and animal taxa. *Ecology*, *96*(6), 1492–1501.

https://doi.org/10.1890/14-1307.1

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing

reads. *EMBnet.Journal*, *17*(1), 10. https://doi.org/10.14806/ej.17.1.200

Met Office, Hollis, D., McCarthy, M., Kendon, M., Legg, T., & Simpson, I. (2019). *HadUK-Grid gridded and regional average climate observations for the UK*. Centre for Environmental Data Analysis, 2019. http://catalogue.ceda.ac.uk/uuid/4dc8450d889a491ebb20e724debe2dfb

Milcu, A., Allan, E., Roscher, C., Jenkins, T., Meyer, S. T., Flynn, D., Bessler, H., Buscot, F., Engels, C., Gubsch, M., König, S., Lipowsky, A., Loranger, J., Renker, C., Scherber, C., Schmid, B., Thébault, E., Wubet, T., Weisser, W. W., … Eisenhauer, N. (2013). Functionally and phylogenetically diverse plant communities key to soil biota. *Ecology*, *94*(8), 1878–1885. https://doi.org/10.1890/12-1936.1

Nguyen, N. H., Williams, L. J., Vincent, J. B., Stefanski, A., Cavender-Bares, J., Messier, C., Paquette, A., Gravel, D., Reich, P. B., & Kennedy, P. G. (2016). Ectomycorrhizal fungal diversity and saprotrophic fungal diversity are linked to different tree community attributes in a field-based tree experiment. *Molecular Ecology*, *25*(16), 4032–4046. https://doi.org/10.1111/mec.13719

Nilsson, R. H., Larsson, K.-H., Taylor, A. F. S., Bengtsson-Palme, J., Jeppesen, T. S., Schigel, D., Kennedy, P., Picard, K., Glöckner, F. O., Tedersoo, L., Saar, I., Kõljalg, U., & Abarenkov, K. (2019). The UNITE database for molecular identification of fungi: Handling dark taxa and parallel taxonomic classifications. *Nucleic Acids Research*, *47*(D1), D259–D264. https://doi.org/10.1093/nar/gky1022

Oksanen, J., Blanchet, G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., & Wagner, H. (2020). *vegan: Community Ecology Package* [Computer software]. http://cran.r-project.org/package=vegan

Peay, K. G., Baraloto, C., & Fine, P. V. A. (2013). Strong coupling of plant and fungal

community structure across western Amazonian rainforests. *The ISME Journal*, *7*(9),

1852–1861. https://doi.org/10.1038/ismej.2013.66

Pichler, M., & Hartig, F. (2021). A new joint species distribution model for faster and more

accurate inference of species associations from big community data. *Methods in

Ecology and Evolution*, 2041-210X.13687. https://doi.org/10.1111/2041-210X.13687

Potts, S. G., Woodcock, B. A., Roberts, S. P. M., Tscheulin, T., Pilgrim, E. S., Brown, V. K., &

Tallowin, J. R. (2009). Enhancing pollinator biodiversity in intensive grasslands.

Journal of Applied Ecology, *46*(2), 369–379. https://doi.org/10.1111/j.1365-

2664.2009.01609.x

Prober, S. M., Leff, J. W., Bates, S. T., Borer, E. T., Firn, J., Harpole, W. S., Lind, E. M.,

Seabloom, E. W., Adler, P. B., Bakker, J. D., Cleland, E. E., DeCrappeo, N. M.,

DeLorenze, E., Hagenah, N., Hautier, Y., Hofmockel, K. S., Kirkman, K. P., Knops, J. M.

H., La Pierre, K. J., … Fierer, N. (2015). Plant diversity predicts beta but not alpha

diversity of soil microbes across grasslands worldwide. *Ecology Letters*, *18*(1), 85–95.

https://doi.org/10.1111/ele.12381

R Core Team. (2023). *R: A language and environment for statistical computing* [Computer

software]. R Foundation for Statistical Computing. http://www.r-project.org/

Ren, C., Zhao, F., Shi, Z., Chen, J., Han, X., Yang, G., Feng, Y., & Ren, G. (2017). Differential

responses of soil microbial biomass and carbon-degrading enzyme activities to

altered precipitation. *Soil Biology and Biochemistry*, *115*, 1–10.

https://doi.org/10.1016/j.soilbio.2017.08.002

Robinson, D. A., Astbury, S., Barrett, G., Burden, A., Carter, H., Emmett, B. A., Garbutt, A.,

Giampieri, C., Hall, J., Henrys, P., Hughes, S., Hunt, A., Jarvis, S., Jones, D. L., Keenan,

P., Lebron, I., Nunez, D., Owen, A., Patel, M., … Wood, C. (2019). *Topsoil physico-chemical properties from the Glastir Monitoring and Evaluation Programme, Wales 2013-2016*. NERC Environmental Information Data Centre. https://doi.org/10.5285/0fa51dc6-1537-4ad6-9d06-e476c137ed09

Scherber, C., Eisenhauer, N., Weisser, W. W., Schmid, B., Voigt, W., Fischer, M., Schulze, E.-D., Roscher, C., Weigelt, A., Allan, E., Beßler, H., Bonkowski, M., Buchmann, N., Buscot, F., Clement, L. W., Ebeling, A., Engels, C., Halle, S., Kertscher, I., … Tscharntke, T. (2010). Bottom-up effects of plant diversity on multitrophic interactions in a biodiversity experiment. *Nature*, *468*(7323), 553–556. https://doi.org/10.1038/nature09492

Schulze, C. H., Waltert, M., Kessler, P. J. A., Pitopang, R., Shahabuddin, Veddeler, D., Mühlenberg, M., Gradstein, S. R., Leuschner, C., Steffan-Dewenter, I., & Tscharntke, T. (2004). Biodiversity indicator groups of tropical land-use systems: Comparing plants, birds, and insects. *Ecological Applications*, *14*(5), 1321–1333. https://doi.org/10.1890/02-5409

Seaton, F. M., George, P. B. L., Lebron, I., Jones, D. L., Creer, S., & Robinson, D. A. (2020). Soil textural heterogeneity impacts bacterial but not fungal diversity. *Soil Biology and Biochemistry*, *144*, 107766. https://doi.org/10.1016/j.soilbio.2020.107766

Siriwardena, G. M., Taylor, R. M., Norfolk, D., Jones, K., Alison, J., Astbury, S., Burden, A., Emmett, B. A., Garbutt, A., Henrys, P., Owen, A., Smart, S. M., Sharps, K., Williams, B., & Wood, C. M. (2020). *Bird counts from the Glastir Monitoring and Evaluation Programme, Wales 2013-2016*. NERC Environmental Information Data Centre. https://doi.org/10.5285/31da0a94-62be-47b3-b76e-4bdef3037360

Smart, S. M., Astbury, S., Burden, A., Emmett, B. A., Garbutt, A., Henrys, P., Maskell, L. C., Owen, A., Sharps, K., Wallace, H., Williams, B., & Wood, C. M. (2020). *Vegetation plot data from the Glastir Monitoring and Evaluation Programme, Wales 2013-2016.* NERC Environmental Information Data Centre. https://doi.org/10.5285/71d3619c-4439-4c9e-84dc-3ca873d7f5cc

Smart, S. M., Firbank, L. G., Bunce, R. G. H., & Watkins, J. W. (2000). Quantifying changes in abundance of food plants for butterfly larvae and farmland birds. *Journal of Applied Ecology*, *37*(3), 398–414.

Stein, A., & Kreft, H. (2014). Terminology and quantification of environmental heterogeneity in species-richness research. *Biological Reviews of the Cambridge Philosophical Society*. https://doi.org/10.1111/brv.12135

Tedersoo, L., Bahram, M., Cajthaml, T., Põlme, S., Hiiesalu, I., Anslan, S., Harend, H., Buegger, F., Pritsch, K., Koricheva, J., & Abarenkov, K. (2016). Tree diversity and species identity effects on soil fungi, protists and animals are context dependent. *The ISME Journal*, *10*(2), 346–362. https://doi.org/10.1038/ismej.2015.116

Weiss, S., Van Treuren, W., Lozupone, C., Faust, K., Friedman, J., Deng, Y., Xia, L. C., Xu, Z. Z., Ursell, L., Alm, E. J., Birmingham, A., Cram, J. A., Fuhrman, J. A., Raes, J., Sun, F., Zhou, J., & Knight, R. (2016). Correlation detection strategies in microbial data sets vary widely in sensitivity and precision. *The ISME Journal*, *10*(7), 1669–1681. https://doi.org/10.1038/ismej.2015.235

Weiss, S., Xu, Z. Z., Peddada, S., Amir, A., Bittinger, K., Gonzalez, A., Lozupone, C., Zaneveld, J. R., Vázquez-Baeza, Y., Birmingham, A., Hyde, E. R., & Knight, R. (2017). Normalization and microbial differential abundance strategies depend upon data characteristics. *Microbiome*, *5*(1), 27. https://doi.org/10.1186/s40168-017-0237-y

Weisser, W. W., Roscher, C., Meyer, S. T., Ebeling, A., Luo, G., Allan, E., Beßler, H., Barnard,

R. L., Buchmann, N., Buscot, F., Engels, C., Fischer, C., Fischer, M., Gessler, A.,

Gleixner, G., Halle, S., Hildebrandt, A., Hillebrand, H., de Kroon, H., … Eisenhauer, N.

(2017). Biodiversity effects on ecosystem functioning in a 15-year grassland

experiment: Patterns, mechanisms, and open questions. *Basic and Applied Ecology*,

23, 1–73. https://doi.org/10.1016/j.baae.2017.06.002

Wolters, V., Bengtsson, J., & Zaitsev, A. S. (2006). Relationship among the species richness of

different taxa. *Ecology*, *87*(8), 1886–1895. https://doi.org/10.1890/0012-

9658(2006)87[1886:RATSRO]2.0.CO;2

Wood, C. M., Alison, J., Botham, M. S., Burden, A., Edwards, F., Garbutt, R. A., George, P. B.

L., Henrys, P. A., Hobson, R., Jarvis, S., Keenan, P., Keith, A. M., Lebron, I., Maskell, L.

C., Norton, L. R., Robinson, D. A., Seaton, F. M., Scarlett, P., Siriwardena, G. M., …

Emmett, B. A. (2021). Integrated ecological monitoring in Wales: The Glastir

Monitoring and Evaluation Programme field survey. *Earth System Science Data*,

13(8), 4155–4173. https://doi.org/10.5194/essd-13-4155-2021

Wright, E. (2016). Using DECIPHER v2.0 to analyze big biological sequence data in R. *The R

Journal*, *8*(1), 352–359.

Yang, P., & van Elsas, J. D. (2018). Mechanisms and ecological implications of the movement

of bacteria in soil. *Applied Soil Ecology*, *129*, 112–120.

https://doi.org/10.1016/J.APSOIL.2018.04.014

Yang, T., Adams, J. M., Shi, Y., He, J., Jing, X., Chen, L., Tedersoo, L., & Chu, H. (2017). Soil

fungal diversity in natural grasslands of the Tibetan Plateau: Associations with plant

diversity and productivity. *New Phytologist*, *215*, 756–765.

https://doi.org/10.1111/nph.14606

Yashiro, E., Pinto-Figueroa, E., Buri, A., Spangenberg, J. E., Adatte, T., Niculita-Hirzel, H.,

Guisan, A., & van der Meer, J. R. (2018). Meta-scale mountain grassland

observatories uncover commonalities as well as specific interactions among plant

and non-rhizosphere soil bacterial communities. *Scientific Reports*, *8*(1), 5758.

https://doi.org/10.1038/s41598-018-24253-x

Yuan, X., Knelman, J. E., Wang, D., Goebl, A., Gasarch, E., Seastedt, T. R., Yuan, X., Knelman,

J. E., Wang, D., Goebl, A., Gasarch, E., & Seastedt, T. R. (2017). Patterns of soil

bacterial richness and composition tied to plant richness, soil nitrogen, and soil

acidity in alpine tundra. *Arctic, Antarctic, and Alpine Research*, *49*(3), 441–453.

https://doi.org/10.1657/AAAR0016-050 URL:

Supplementary information

Supplementary figure 1

A representative schematic showing the layout of the vegetation plots and pollinator and bird transects within a 1km square. There are 5 randomly located plots with both plant and soil surveyed in each square, while the number and location of plant only plots varies by square as some plot types are habitat specific (e.g. hedgerow plots). Size of plots and transect sections is not to scale.

Supplementary figure 2

The density distributions of species correlations from the belowground/plot-level (a) and aboveground/square level (b) joint species distribution models that took no account of environment (No env), linear functions of environment only (Linear), or that used deep neural nets to allow for non-linear effects of environment (DNN). Residual correlations are split by plant growth form for the belowground analysis (a) and split by whether the plants are food plants for birds or pollinators for the aboveground analysis (b). Note that despite butterflies, bees and hoverflies being combined into a pollinator category for graphical simplicity the food plants for butterflies are those plants that are important for butterfly larvae while the food plants for bees and hoverflies are those that produce nectar. Also note that there is only woody plant species that appeared in enough plots to be included in the belowground analysis (Vaccinium myrtillus), so those results are unlikely to be representative of all woody species.

Supplementary figure 3

The relative importance (y axis) of the different joint species distribution model components for every bacterial, plant, fungal, animal, and heterotrophic protist taxa (x axis). The importance of the biotic interactions is shown in blue, soil variables (carbon, nitrogen, pH, conductivity, organic matter, water repellency, Olsen-P phosphorus, water content, bulk density) is shown in orange and spatial variation is shown in grey. Taxa are shown in order of decreasing contribution of the biotic component. Results shown are for the model with linear effects of the environment only.

Supplementary figure 4

The relative importance (y axis) of the different joint species distribution model components for every bee/hoverfly, bird, butterfly and plant species (x axis). The importance of biotic interactions is shown in blue, the environment (rainfall, temperature and elevation) is shown in orange and spatial variation is shown in grey. Taxa are shown in order of decreasing contribution of the biotic component. Results shown are for the model with linear effects of the environment only.

Supplementary table 1: Model parameter estimates from belowground no-

environment model

Parameter	Mean	Est.Error	l-95% CI	u-95% CI	Rhat	Bulk_ESS	Tail_SS
Group-level effects							
sd(Bacteria)	477.66	34.57	413.59	549.56	1.00	5254	9557
sd(Fungi)	50.15	5.48	39.98	61.53	1.00	3875	7184
sd(AM Fungi)	2.28	0.58	0.91	3.28	1.00	2847	2653
sd(Heterotrophs)	36.65	7.52	21.08	50.81	1.00	1976	2852
sd(Animals)	17.83	1.79	14.41	21.50	1.00	5863	8827
sd(Plants)	2.17	0.38	1.37	2.88	1.00	3704	5497
Population-level effects							
Bacteria Intercept	2307.14	44.22	2220.48	2395.41	1.00	7659	9978
Fungi Intercept	342.62	6.20	330.51	354.88	1.00	16093	12548
AM Fungi Intercept	10.06	0.39	9.29	10.81	1.00	22616	13035
Heterotroph Intercept	277.93	5.97	266.31	289.59	1.00	22786	10836
Animal Intercept	64.27	1.88	60.55	67.93	1.00	15502	12845
Plant Intercept	10.54	0.29	9.98	11.10	1.00	20980	12641
Family specific parameters							
Sigma Bacteria	435.93	23.67	392.12	484.28	1.00	6285	9714
Sigma Fungi	92.01	4.82	82.68	101.66	1.00	5046	8322
Sigma AM Fungi	6.94	0.31	6.35	7.56	1.00	6818	8536
Sigma Heterotrophs	101.76	5.34	91.32	112.19	1.00	3529	6672
Sigma Animal	24.50	1.06	22.50	26.64	1.00	11884	12125
Sigma Plants	4.59	0.21	4.20	5.03	1.00	6767	10143
Residual Correlations							
Bacteria, Fungi	0.60	0.04	0.51	0.68	1.00	7008	10831
Bacteria, AM Fungi	0.37	0.05	0.26	0.46	1.00	13007	12551
Fungi, AM Fungi	0.56	0.04	0.48	0.63	1.00	19891	14261
Bacteria, Heterotrophs	0.67	0.04	0.59	0.74	1.00	7798	10733
Fungi, Heterotrophs	0.80	0.02	0.75	0.84	1.00	8596	10582
AM Fungi, Heterotrophs	0.52	0.04	0.43	0.59	1.00	21013	13808
Bacteria, Animals	0.24	0.06	0.12	0.35	1.00	16516	13739

Fungi, Animal	0.24	0.06	0.12	0.35	1.00	17571	13457
AM Fungi, Animal	-0.01	0.06	-0.13	0.11	1.00	17916	13082
Heterotroph, Animal	0.19	0.06	0.07	0.31	1.00	17929	13777
Bacteria, Plant	0.27	0.06	0.16	0.38	1.00	18851	13946
Fungi, Plant	0.14	0.06	0.02	0.25	1.00	19447	13581
AM Fungi, Plant	0.31	0.05	0.20	0.40	1.00	18497	12706
Heterotroph, Plant	0.17	0.06	0.06	0.28	1.00	21841	13432
Animal, Plant	-0.01	0.06	-0.12	0.11	1.00	18000	13017

Supplementary table 2: Model parameter estimates from belowground linear

environment model

Parameter	Mean	Est.Error	l-95% CI	u-95% CI	Rhat	Bulk_ESS	Tail_SS
Group-level effects							
sd(Bacteria)	485.09	35.34	420.42	558.35	1.00	4365	8353
sd(Fungi)	51.58	5.69	41.09	63.34	1.00	2639	5804
sd(AM Fungi)	2.39	0.52	1.23	3.29	1.00	2502	2074
sd(Heterotrophs)	35.91	8.7	16.53	51.76	1.00	1323	1465
sd(Animals)	13.93	1.75	10.53	17.44	1.00	4820	8324
sd(Plants)	2.15	0.36	1.41	2.83	1.00	4075	5802
Population-level effects							
Bacteria Intercept	2308.10	45.41	2218.16	2396.38	1.00	5004	7864
Fungi Intercept	347.58	8.91	330.37	364.81	1.00	15531	11916
AM Fungi Intercept	10.07	2.35	5.40	14.68	1.00	18448	12795
Heterotroph Intercept	285.12	8.69	268.24	302.26	1.00	21240	11439
Animal Intercept	51.25	5.52	40.56	62.15	1.00	23026	11874
Plant Intercept	5.86	1.97	1.98	9.68	1.00	15542	11882
Bacteria pH	0.24	1.00	-1.72	2.20	1.00	31809	11227
Bacteria Carbon	-0.12	1.00	-2.08	1.83	1.00	30469	11496
Bacteria Water	-0.01	0.98	-1.93	1.88	1.00	28298	11831
Bacteria Total P	-0.05	0.99	-1.98	1.88	1.00	32645	11249
Bacteria SWR	-0.25	0.99	-2.18	1.68	1.00	29401	11595
Fungi pH	0.49	0.98	-1.43	2.41	1.00	28923	11225
Fungi Carbon	-1.28	0.98	-3.23	0.65	1.00	27564	11324
Fungi Water	-0.25	1.01	-2.22	1.73	1.00	31189	11011
Fungi Total P	-0.27	0.99	-2.23	1.65	1.00	28801	11403
Fungi SWR	-0.86	0.84	-2.53	0.81	1.00	22227	11631
AM Fungi pH	0.54	0.40	-0.24	1.31	1.00	16135	12384
AM Fungi Carbon	-2.02	0.44	-2.86	-1.16	1.00	13846	12595
AM Fungi Water	-0.05	0.91	-1.83	1.73	1.00	26581	11602
AM Fungi Total P	1.40	0.55	0.32	2.47	1.00	17351	12966
AM Fungi SWR	0.33	0.14	0.05	0.60	1.00	17349	12711
Heterotrophs pH	1.42	0.99	-0.52	3.37	1.00	27496	11069
Heterotrophs Carbon	-1.02	0.98	-2.93	0.90	1.00	29679	11271

Heterotrophs Water	-0.07	0.99	-2.01	1.87	1.00	27640	10821
Heterotrophs Total P	0.54	1.00	-1.43	2.50	1.00	29504	11148
Heterotrophs SWR	-2.21	0.87	-3.90	-0.52	1.00	23872	11840
Animal pH	-2.09	0.89	-3.85	-0.31	1.00	21494	12070
Animal Carbon	3.87	0.90	2.10	5.61	1.00	23362	12676
Animal Water	0.78	1.01	-1.21	2.75	1.00	29958	11553
Animal Total P	0.08	0.94	-1.78	1.91	1.00	26089	11870
Animal SWR	2.72	0.5	1.74	3.69	1.00	19709	11363
Plant pH	1.12	0.34	0.48	1.79	1.00	15392	11842
Plant Carbon	-0.71	0.37	-1.42	0.01	1.00	14657	11767
Plant Water	1.02	0.86	-0.69	2.71	1.00	25028	12902
Plant Total P	-0.56	0.46	-1.48	0.36	1.00	18584	12316
Plant SWR	0.15	0.12	-0.08	0.38	1.00	17736	12660
Family specific parameters							
Sigma Bacteria	429.28	23.30	386.31	476.99	1.00	4138	7342
Sigma Fungi	89.27	4.95	79.82	99.15	1.00	3043	6223
Sigma AM Fungi	6.14	0.30	5.58	6.76	1.00	5698	7606
Sigma Heterotrophs	98.32	5.67	87.5	109.68	1.00	1933	3557
Sigma Animal	24.71	1.10	22.68	26.93	1.00	8999	11458
Sigma Plants	4.36	0.20	3.99	4.77	1.00	6737	9422
Residual Correlations							
Bacteria, Fungi	0.58	0.05	0.48	0.67	1.00	4469	8046
Bacteria, AM Fungi	0.27	0.06	0.15	0.39	1.00	9362	11687
Fungi, AM Fungi	0.46	0.05	0.36	0.56	1.00	12405	11997
Bacteria, Heterotrophs	0.65	0.04	0.56	0.72	1.00	4542	8129
Fungi, Heterotrophs	0.78	0.03	0.73	0.83	1.00	4276	7634
AM Fungi, Heterotrophs	0.41	0.06	0.29	0.51	1.00	11636	12142
Bacteria, Animals	0.41	0.05	0.30	0.51	1.00	10812	12481
Fungi, Animal	0.43	0.05	0.32	0.53	1.00	12233	11743
AM Fungi, Animal	0.12	0.06	0.00	0.23	1.00	16028	13084
Heterotroph, Animal	0.40	0.06	0.28	0.5	1.00	12937	12049
Bacteria, Plant	0.12	0.07	-0.02	0.25	1.00	12923	13151
Fungi, Plant	-0.02	0.07	-0.15	0.12	1.00	13761	12593

AM Fungi, Plant	0.21	0.06	0.09	0.32	1.00	13550	12713
Heterotroph, Plant	0.01	0.07	-0.13	0.15	1.00	12678	12114
Animal, Plant	0.02	0.06	-0.09	0.14	1.00	17748	12603

Supplementary table 3: Model parameter estimates from belowground sigmoidal pH

model, where richness ~ delta + alpha/(1 + exp(-beta*(pH – gamma)))

Parameter	Mean	Est.Error	l-95% CI	u-95% CI	Rhat	Bulk_ESS	Tail_SS
Group-level effects							
sd(Bacteria)	483.85	32.61	424.03	551.00	1.00	4819	8526
sd(Fungi)	48.89	5.00	39.42	59.01	1.00	4652	8230
sd(AM Fungi)	2.35	0.47	1.34	3.20	1.00	3846	4596
sd(Heterotrophs)	36.15	5.80	24.51	47.32	1.00	3450	5180
sd(Animals)	13.91	1.90	10.12	17.66	1.00	4880	7739
sd(Plants)	1.90	0.38	1.09	2.59	1.00	3367	4043
Population-level effects							
Bacteria delta	-815.07	504.36	-1810.22	180.76	1.00	12744	11036
Bacteria alpha	3729.08	500.27	2740.88	4711.39	1.00	13426	11601
Bacteria beta	0.85	0.14	0.60	1.13	1.00	13886	12847
Bacteria gamma	2.21	0.31	1.62	2.86	1.00	11008	10699
Fungi delta	187.06	25.97	131.31	232.50	1.00	9484	9775
Fungi alpha	220.03	29.32	167.45	282.80	1.00	9839	11044
Fungi beta	1.84	0.35	1.25	2.60	1.00	13547	11455
Fungi gamma	3.65	0.18	3.28	3.98	1.00	11472	10999
AM Fungi delta	0.77	1.69	-3.20	3.12	1.00	7903	5083
AM Fungi alpha	12.73	1.76	10.17	16.88	1.00	8319	5456
AM Fungi beta	3.37	0.59	2.24	4.56	1.00	9362	6480
AM Fungi gamma	3.70	0.14	3.37	3.92	1.00	8174	5392
Heterotrophs delta	40.24	45.66	-55.61	119.52	1.00	7675	10751
Heterotrophs alpha	342.78	50.95	249.80	447.29	1.00	8057	11253
Heterotrophs beta	1.38	0.28	0.95	2.06	1.00	9392	11202
Heterotrophs gamma	3.59	0.23	3.10	4.00	1.00	7968	10348
Animal delta	44.81	7.24	29.88	57.68	1.00	13041	12309
Animal alpha	46.50	10.16	26.82	66.37	1.00	15245	9742
Animal beta	-0.62	0.33	-1.44	-0.18	1.00	11085	8079
Animal gamma	3.80	0.92	2.04	5.67	1.00	19688	12257
Plant delta	-3.68	6.46	-19.27	4.90	1.00	8989	9031
Plant alpha	15.8	6.49	7.10	31.42	1.00	8944	9041
Plant beta	2.08	0.53	1.22	3.29	1.00	13516	12529
Plant gamma	2.77	0.39	1.98	3.44	1.00	9098	10654

Parameter	Mean	Est.Error	l-95% CI	u-95% CI	Rhat	Bulk_ESS	Tail_SS
Family specific parameters							
Sigma Bacteria	324.95	15.71	295.73	356.94	1.00	9070	12594
Sigma Fungi	74.92	3.65	68.09	82.43	1.00	6450	10399
Sigma AM Fungi	5.63	0.25	5.16	6.14	1.00	7722	8874
Sigma Heterotrophs	76.35	3.86	69.2	84.32	1.00	5292	7938
Sigma Animal	25.66	1.16	23.48	28.05	1.00	8109	11683
Sigma Plants	4.29	0.19	3.94	4.68	1.00	6650	7518
Residual Correlations							
Bacteria, Fungi	0.38	0.06	0.26	0.48	1.00	8392	11520
Bacteria, AM Fungi	0.06	0.06	-0.06	0.17	1.00	15094	13628
Fungi, AM Fungi	0.36	0.05	0.25	0.45	1.00	16897	13100
Bacteria, Heterotrophs	0.42	0.05	0.31	0.52	1.00	8724	12094
Fungi, Heterotrophs	0.68	0.03	0.61	0.74	1.00	9808	12968
AM Fungi, Heterotrophs	0.26	0.05	0.16	0.37	1.00	18659	12906
Bacteria, Animals	0.35	0.05	0.24	0.45	1.00	12776	13129
Fungi, Animal	0.36	0.05	0.25	0.46	1.00	13437	13853
AM Fungi, Animal	0.03	0.06	-0.08	0.14	1.00	20372	13474
Heterotroph, Animal	0.34	0.05	0.23	0.44	1.00	14176	12433
Bacteria, Plant	0.06	0.06	-0.05	0.17	1.00	18625	13447
Fungi, Plant	-0.10	0.06	-0.21	0.01	1.00	18670	13472
AM Fungi, Plant	0.12	0.05	0.02	0.23	1.00	21454	13134
Heterotroph, Plant	-0.10	0.06	-0.21	0.02	1.00	20854	13135
Animal, Plant	0.00	0.06	-0.11	0.11	1.00	21656	13670

Supplementary table 4: Model parameter estimates for the square-level model with

no environmental variables

Parameter	Mean	Est.Error	l-95% CI	u-95% CI	Rhat	Bulk_ESS	Tail_SS
Group-level effects							
sd(Bird)	0.19	0.06	0.10	0.33	1.00	4778	7544
sd(Butterfly)	0.18	0.06	0.07	0.32	1.00	3436	3102
sd(Bee)	0.13	0.03	0.08	0.19	1.00	6108	9760
sd(Plant)	0.04	0.03	0.00	0.12	1.00	5522	6382
Population-level effects							
Bird Intercept	2.87	0.06	2.75	2.99	1.00	5834	7708
Butterfly Intercept	1.31	0.06	1.20	1.43	1.00	6805	9244
Bee Intercept	0.37	0.04	0.29	0.44	1.00	4657	7232
Plant Intercept	2.95	0.03	2.90	3.01	1.00	13772	10780
Family specific parameters							
Sigma Bird	0.60	0.03	0.55	0.65	1.00	11991	11968
Sigma Butterfly	0.55	0.02	0.51	0.60	1.00	11464	11140
Sigma Bee	0.31	0.01	0.28	0.33	1.00	15334	12013
Sigma Plant	0.44	0.02	0.40	0.48	1.00	17887	12348
Residual Correlations							
Bird, Butterfly	0.44	0.05	0.35	0.54	1.00	12775	11460
Bird, Bee	0.40	0.05	0.30	0.50	1.00	12517	12002
Butterfly, Bee	0.40	0.05	0.30	0.50	1.00	14866	12384
Bird, Plant	0.41	0.05	0.31	0.50	1.00	16577	12828
Butterfly, Plant	0.27	0.06	0.16	0.38	1.00	16249	12317
Bee, Plant	0.24	0.06	0.13	0.35	1.00	16463	12440

Supplementary table 5: Model parameter estimates for the square-level model with

linear environmental effects

Parameter	Mean	Est.Error	l-95% CI	u-95% CI	Rhat	Bulk_ESS	Tail_SS
Group-level effects							
sd(Bird)	0.07	0.04	0.00	0.16	1.00	4906	7476
sd(Butterfly)	0.08	0.04	0.00	0.17	1.00	4597	7665
sd(Bee)	0.13	0.03	0.08	0.19	1.00	7743	10212
sd(Plant)	0.07	0.04	0.00	0.16	1.00	4496	6900
Population-level effects							
Bird Intercept	2.87	0.03	2.80	2.94	1.00	18563	11184
Butterfly Intercept	1.31	0.04	1.24	1.38	1.00	17306	11421
Bee Intercept	0.36	0.04	0.28	0.42	1.00	9322	10928
Plant Intercept	2.95	0.03	2.89	3.02	1.00	18104	11705
Bird Rainfall	-0.16	0.04	-0.23	-0.08	1.00	17187	13273
Bird Temperature	0.28	0.04	0.20	0.34	1.00	18637	12865
Butterfly Rainfall	-0.21	0.04	-0.28	-0.13	1.00	17696	11189
Butterfly Temperature	0.17	0.04	0.10	0.24	1.00	17741	12927
Bee Rainfall	-0.04	0.03	-0.09	0.01	1.00	16468	12460
Bee Temperature	0.14	0.02	0.09	0.19	1.00	16750	12887
Plant Rainfall	-0.02	0.03	-0.09	0.04	1.00	19208	13308
Plant Temperature	0.14	0.03	0.08	0.21	1.00	13754	11802
Family specific parameters							
Sigma Bird	0.46	0.02	0.42	0.50	1.00	28206	12267
Sigma Butterfly	0.45	0.02	0.41	0.49	1.00	24489	12021
Sigma Bee	0.26	0.01	0.24	0.28	1.00	24091	12640
Sigma Plant	0.41	0.02	0.37	0.44	1.00	22690	12417
Residual Correlations							
Bird, Butterfly	0.11	0.06	0.00	0.23	1.00	28912	11151
Bird, Bee	0.10	0.06	-0.02	0.22	1.00	29749	11640
Butterfly, Bee	0.15	0.06	0.03	0.26	1.00	27279	11895
Bird, Plant	0.24	0.06	0.13	0.34	1.00	27174	11776
Butterfly, Plant	0.07	0.06	-0.04	0.19	1.00	30723	11240
Bee, Plant	0.04	0.06	-0.08	0.16	1.00	31735	11764

Supplementary table 6: Model parameter estimates from aboveground

temperature*rainfall model to exponentiated diversity values

Parameter	Mean	Est.Error	l-95% CI	u-95% CI	Rhat	Bulk_ESS	Tail_SS
Group-level effects							
sd(Bird)	0.61	0.45	0.03	1.67	1.00	5710	6653
sd(Butterfly)	0.42	0.18	0.08	0.80	1.00	4361	4016
sd(Bee)	0.22	0.05	0.13	0.33	1.00	6282	9516
sd(Plant)	0.64	0.49	0.03	1.83	1.00	6781	6545
Population-level effects							
Bird Intercept	21.68	0.44	20.80	22.53	1.00	15293	11855
Butterfly Intercept	4.19	0.16	3.87	4.52	1.00	9384	10145
Bee Intercept	1.51	0.06	1.38	1.63	1.00	5189	8224
Plant Intercept	21.94	0.53	20.90	22.98	1.00	15717	11762
Bird Rainfall	-1.03	0.42	-1.87	-0.22	1.00	11612	10824
Bird Temperature	3.15	0.39	2.37	3.90	1.00	10825	10758
Bird Rain:Temp	1.82	0.28	1.26	2.37	1.00	16958	12921
Butterfly Rainfall	-0.69	0.15	-0.99	-0.40	1.00	10031	11251
Butterfly Temperature	0.66	0.14	0.39	0.93	1.00	10132	11386
Butterfly Rain:Temp	-0.12	0.09	-0.29	0.06	1.00	15630	12336
Bee Rainfall	-0.05	0.05	-0.14	0.04	1.00	8262	9444
Bee Temperature	0.22	0.04	0.14	0.30	1.00	8853	10129
Bee Rain:Temp	0.01	0.02	-0.04	0.06	1.00	14405	12199
Plant Rainfall	-0.13	0.50	-1.12	0.82	1.00	12470	11396
Plant Temperature	1.44	0.45	0.55	2.34	1.00	13752	11963
Plant Rain:Temp	1.42	0.35	0.74	2.11	1.00	16997	11535
Family specific parameters							
Sigma Bird	5.70	0.24	5.25	6.18	1.00	18974	12314
Sigma Butterfly	1.64	0.07	1.51	1.79	1.00	15431	11038
Sigma Bee	0.44	0.02	0.41	0.48	1.00	16565	11169
Sigma Plants	7.19	0.30	6.63	7.81	1.00	19782	12479
Residual Correlations							
Bird, Butterfly	0.04	0.06	-0.08	0.15	1.00	18679	12017
Bird, Bee	0.10	0.06	-0.02	0.21	1.00	18293	12300
Butterfly, Bee	0.15	0.06	0.03	0.26	1.00	17104	11691
Bird, Plant	0.21	0.06	0.10	0.32	1.00	17690	12319
Butterfly, Plant	0.03	0.06	-0.08	0.15	1.00	18300	12066

Parameter	Mean	Est.Error	l-95% CI	u-95% CI	Rhat	Bulk_ESS	Tail_SS
Bee, Plant	-0.01	0.06	-0.13	0.11	1.00	18750	11857