Exposure to false cardiac feedback alters pain perception and anticipatory cardiac frequency.

Eleonora Parrotta[1,2,3], Patric Bach[1], Giovanni Pezzulo[4], Andrea Zaccaro[3], Mauro Gianni Perrucci[3,5], Marcello Costantini [5,6], Francesca Ferri[3,5]

[1] School of Psychology, University of Aberdeen

[2] Department of Psychology, Sapienza University of Rome, Italy

[3] Department of Neuroscience, Imaging and Clinical Sciences, "G. d'Annunzio" University of Chieti-Pescara, Chieti, Italy

[4] Institute of Cognitive Sciences and Technologies, National Research Council, Rome, Italy

[5] Institute for Advanced Biomedical Technologies ☐ ITAB, "G. d'Annunzio" University of Chieti-Pescara, Chieti, Italy

[6] Department of Psychology, "G. d'Annunzio" University of Chieti-Pescara, Chieti, Italy

[7] UdA-TechLab, Research Center, University "G. d'Annunzio" of Chieti-Pescara, Chieti, Italy

Correspondence:

Eleonora Parrotta
eleonora.parrotta@uniroma1.it
Department of Psychology
Sapienza University of Rome
Rome, Italy

Keywords: embodied predictive coding, perceptual inference, active inference, pain perception, false cardiac feedback

Abstract

The experience of pain, like other interoceptive processes, has recently been conceptualized in terms of predictive coding and free energy frameworks. In these views, the brain integrates sensory, proprioceptive, and interoceptive signals to generate probabilistic inferences about upcoming events, which shape both the state and the perception of our inner body. Here we ask whether it is possible to induce pain expectations by providing false faster (vs. slower) acoustic cardiac feedback before administering electrical cutaneous shocks. We test whether these expectations will shape both the perception of pain and the body's physiological state toward prior predictions. Results confirmed that faster cardiac feedback elicited pain expectations that affected both perceptual pain judgments and the body's physiological response. Perceptual pain judgments were biased towards the expected level of pain, such that participants illusorily perceived identical noxious stimuli as more intense and unpleasant. Physiological changes mirrored the predicted level of pain, such that participants' actual cardiac response in anticipation of pain stimuli showed a deceleration in heart rate, in line with the well-known orienting cardiac response in anticipation of threatening stimuli (Experiment 1). In a control experiment, such perceptual and cardiac modulations were dramatically reduced when the feedback reproduced an exteroceptive, instead of interoceptive, cardiac feedback (Experiment 2). These findings show that cardiac perception can be

45 understood as interoceptive inference that modulates both our perception and the physiological state
46 of the body, thereby actively generating the interoceptive and autonomic consequences that have
47 been predicted.
48
49 **Introduction**
50
Far more complex than the mere transmission of nociceptive inputs, pain is a component of
the interoceptive system (Craig, 2003b). It is defined as an unpleasant sensory and emotional
experience associated with, or resembling, actual or potential tissue damage (International
Association for the Study of Pain, IASP, 2020). It can occur in the absence of physical harm
(Loeser & Treede, 2008) and is permeated by all the available sources of information, including
sensory, emotional, cognitive, and social components as well as prior information and expectations
(Atlas & Wager, 2012; Tracey, 2010; Williams & Craig, 2016). While the last decades provided
important advances in the understanding of pain, developing an overarching theory that accounts for
all the dimensions of pain still remains challenging (for a full discussion, see Moayedi and Davis,
2013).
Current views propose predictive processing models as a suitable candidate to account for
the multifaceted experience of pain (Büchel et al., 2014; Kiverstein et al., 2022; Song et al., 2019,
2021). Interoceptive inference and the Embodied Predictive Coding frameworks (e.g., Barrett,
2017; Barrett and Simmons, 2015; Pezzulo, 2014; Seth, 2013; Seth and Friston, 2016) conceptualise
the brain as an active generator of inferences, which maintains an internal model of both the internal
and the external world and attempts to fit it with incoming inputs through a process of Bayesian
hypothesis testing and revision. Interoceptive sensations are thought to be derived from the
integration of different sources of information into an expectation of the body's upcoming changes,
which are kept in check by the actual state of the body (Barrett & Simmons, 2015). However,
people's predictions do not always mirror reality, but can be inaccurate, leading to a gap between
the expected and the actual present state. The goal of inferential processes is to reduce this
difference between one's internal models and the actual interoceptive input, thus converging
towards the *brain's best guess* about the body's state (Ainley et al., 2016; Seth et al., 2012; Seth &
Friston, 2016).
An important feature of these accounts is that the difference between one's internal models
and the sensory data can be minimized not only by updating internal models to better fit the data
(i.e., perceptual inference), but also by performing actions, changing bodily states themselves (i.e.,
active inference, see Friston, 2010, 2005; Parr et al., 2022). For example, an expectation of a
threatening, painful stimulus could be realised both by making the painful stimulus *appear* more
painful than it really is, thereby fitting one's internal model to bodily reality, or by *actually*
modulating the state of the body and shape the physiologic response to pain towards previous
predictions, for example by engaging autonomic reflexes (e.g., typical decrease in the heart rate in
relation to pain anticipation, e.g., Bradley et al., 2008, 2005; Colloca et al., 2006; Lykken et al.,
1972; Taggart et al., 1976; Tracy et al., 2017).
There is suggestive evidence for both components in the extant literature on pain. The
assumption that prior information modulates pain perception underlines decades of work on placebo
and nocebo effects (i.e., the expectation-effect; Crombez et al., 1998; Keltner et al., 2006; Petrovic
& Ingvar, 2002; Price et al., 1999; Wager, 2005; Wager et al., 2004). On a neuronal level, expecting
pain alters the neural mechanisms in pain-processing regions, thought to reflect a combination of

nociceptive inputs and top-down information (Atlas et al., 2010; Keltner et al., 2006; Koyama et al.,
2005; Wiech et al., 2008), even at a very early stage of processing (Eippert et al., 2009; Geuter &
Büchel, 2013). Moreover, such changes can persist - or even grow over time - in the absence of
disconfirming evidence (Atlas et al., 2010; Colloca et al., 2010; Craggs et al., 2008; Jepma &
Wager, 2015; Koban & Wager, 2016; Montgomery & Kirsch, 1997; Vase et al., 2005, 2011),
turning pain experience into potentially a self-fulfilling prophecy (Jepma et al., 2018).

As for the assumption that unfulfilled predictions can be resolved by engaging autonomic
reactions, there is evidence that prior pain expectations, and pain itself, modulate physiological
responses. Expecting pain induces augmented activity in the sympathetic system (e.g. increases in
blood pressure and skin conductance) to prepare potential avoidance responses (Barlow et al., 1996;
Oka et al., 2007; Tousignant-Laflamme & Marchand, 2006; Yang et al., 2003), as well as a
characteristic decrease in heart rate to promote orienting responses, attention and sensory
processing (Bradley et al., 2008, 2005; Colloca et al., 2006; Lykken et al., 1972; Taggart et al.,
1976; Tracy et al., 2017; for a full discussion, see Skora et al., 2022). Once pain is actually
experienced, however, heart rate rises, reflecting robust sympathetic activation and a well-
documented autonomic response to nociceptive input (Tousignant-Laflamme et al., 2005; Terkelsen
et al., 2005; for a review, see Forte et al., 2022).

Together, these findings support the idea that the experience of pain may rely on top-down
predictive mechanisms, which shape not only the subjective perception of pain, but also its neural
and bodily correlates, both when pain is experienced and merely expected. However, to our
knowledge, no study exists that tracks both the perceptual and physiological responses to
mismatches between pain expectation and bodily reality. Moreover, in previous work, pain
expectations were typically induced by exteroceptive cues (e.g., visual, Wiech et al., 2014; Jepma et
al., 2019; auditory, Colloca et al., 2006; Atlas et al., 2010). Although interoceptive processes may
have contributed to the observed effects, these studies did not specifically target interoceptive
sources of information within the inferential process.

If one assumes that pain and bodily states are integrated within a unified interoceptive
system, then information about the internal condition of the body should be tightly linked to the
experience of pain. Among such interoceptive streams, cardiac activity appears to be closely
intertwined with the experience of pain, likely due to a close integration between the pain system
and the neural circuits involved in cardiovascular regulation (Craig, 2003b, 2003a, 2008). For
example, pain-related evoked potentials and nociception are modulated across the cardiac cycle
(Edwards et al., 2001, 2002, 2008; Martins et al., 2009; McIntyre et al., 2006), and both the
anticipation of and the response to pain are associated with changes in heart rates (Colloca et al.,
2006; Mischkowski et al., 2018; for a review, see Kyle and McNeil, 2014). Moreover, when
estimating their own cardiac frequency, people's estimates integrate the level of threat (i.e., pain)
associated with upcoming events (Parrotta et al., 2024), reflecting the (fictitious) belief that
anticipating threat increases heart rates. Even though anticipating pain typically decreases real heart
rates (the well-known orienting response, Sokolov et al., 2002), people therefore illusorily perceive
a higher cardiac frequency when expecting painful stimulation (Parrotta et al., 2024).

The tight coupling between heart rate and pain experience suggests that it should be possible
to modulate the experience of pain - and the anticipatory bodily response - by manipulating
interoceptive cardiac feedback, that is, by misleading people into believing that their heart rate is
increasing in anticipation of a noxious stimulus. Under predictive architectures, persuading one's
internal model towards the fictitious evidence of increased heart rates should then induce changes

both in *pain perception* (i.e., perceptual inference) and in the *actual state of the body* (i.e., active inference). Specifically, the former process would correspond to an interoceptive illusion of increased pain, whereas the latter would correspond to physiological and autonomic adaptations to the predicted level of pain. We hypothesized that a cardiac feedback manipulation that renders interoceptive streams more similar to those expected during pain would produce an interoceptive illusion of pain as well as ensuing autonomic adjustments - as hypothesized by interoceptive inference and Embodied Predictive Coding theories. To test this idea, Experiment 1 provided participants with auditory cardiac feedback, via headphones, that could diverge from their actual heart rate by being either faster (as when experiencing pain or threat) or slower (as in relaxed states), before the administration of a noxious electrical stimulation, while their ECG was recorded. Once the pain stimulus was administered, subjects were asked to rate its intensity and unpleasantness.

We predicted, first, that the exposure to faster (vs. slower) cardiac feedback would induce expectations of increased levels of pain, such that noxious stimuli are felt and reported as more unpleasant and intense. Accelerated heart rate is a salient, evolutionarily conserved signal of arousal, threat, and nociception, a mapping continually reinforced physiologically when responding to *actual* pain (e.g. for a review, see Forte et al., 2022). If these expectations are integrated with interoceptive inference, they should therefore heighten assessment of pain.

Second, expectations of threat induced by the faster feedback should be fulfilled through anticipatory parasympathetic reflexes, such that participants' real heart rates decrease when exposed to the faster, compared to the slower cardiac feedback, in line with the well-known orienting cardiac response when expecting threat or pain (Bradley et al., 2008, 2005; Colloca et al., 2006; Lykken et al., 1972; Taggart et al., 1976; Tracy et al., 2017; for a full discussion, see Skora et al., 2022). This anticipatory cardiac deceleration reflects an allostatic adjustment implemented by the brain in response to predicted (instead of experienced) threat.

Our hypotheses were theoretically derived from Embodied Predictive Coding accounts of interoception. Our previous findings (Parrotta et al., 2024) suggest that when the brain estimates cardiac activity, it may assign special relevance to pain and anticipated threat, implying that cardiac signals are weighted within the interoceptive schema of pain (Schoeller et al., 2022). In particular, we previously demonstrated that the mere expectation of pain can induce a cardiac interoceptive illusion: participants reported increased heart rate when anticipating pain, even in the absence of any actual physiological change (Parrotta et al., 2024). Here, we tested the counterpart of this illusion: if pain expectations can bias cardiac perception, then manipulating cardiac feedback should, in turn, influence pain perception. Presenting participants with artificially accelerated heartbeat feedback may act as a misleading prior, enhancing perceived pain intensity despite unchanged nociceptive input. Evidence for such a mechanism is further suggested by the evidence that providing mismatched cardiac feedback in contexts where a faster heart rate is normally expected (i.e. sexual arousal, Rupp & Wallen, 2008; Valin, 1996; physical exercise, Iodice et al 2019) can enhance the perception of physiological states associated with those experiences (Crucian et al., 2000; Iodice et al., 2019).

To rule out that potential modulations in participants' pain perception and cardiac state were merely associated with the frequency of the feedback, rather than a heartbeat, Experiment 2 replicated the design in a second group of participants, with the only difference being that the congruent or incongruent feedback would not be a heartbeat tone, but an unrelated exteroceptive stimulus. We predicted that, as participants would have no expectation that such an external

stimulus signals a preparation for a threatening, painful stimulus, variations in the rate of this
exteroceptive stimulus should lead to no (or reduced) perceptual and cardiac changes.

If confirmed, the results would provide the first evidence that predictive processes are
generated by the simulation of interoceptive streams, influencing not only our perception of pain
but also its heart-related physiological response: two changes that jointly minimize the discrepancy
between the expected and the actual homeostatic (interoceptive) state. This evidence would
corroborate the assumption that predictions originate along the embodied multisensory sphere by
integrating multiple sources of information (i.e., interoceptive) within the inference. Ultimately, the
results may provide important insights into how the experience of pain is actively built through
predictive processes.

Experiment 1
Experiment 1 tests whether false cardiac feedback of an accelerated (i.e., faster), relative to a
decelerated (i.e., slower) heart rate, alters both pain perception and the cardiac anticipatory response
to noxious stimuli.
Methods
Participants
Thirty-five participants (mean age 25.11, SD= 2.94, 21 women) took part in the experiment,
recruited from Gabriele D'Annunzio University and the wider community. All were right-handed
with normal or corrected-to-normal vision. Exclusion criteria for taking part were self-reported
chronic and acute pain, neurological disease, serious cardiovascular disease (i.e., any type of disease
involving the heart or blood vessels that might result in life-threatening medical emergencies, e.g.,
arrhythmias, infarction, stroke), or conditions that could potentially interfere with pain sensitivity
(e.g. drug intake or skin diseases). Participants were instructed not to drink coffee or smoke
cigarettes. After taking part in the experiment, participants were excluded if the correlation between
the desired five levels of the nociceptive stimulus intensity and their ratings in the no-feedback
condition was below r=0.75, suggesting insensitivity to the varying nociceptive degrees of the
electrical stimulus. No participants were excluded based on this criterion. All gave written informed
consent, were unaware of the purposes of the study, and were fully debriefed about it at the end of
the experiment. Ethical approval from the local ethics board was obtained. One participant was
excluded as, at the end of the experiment, they declared to suffer from a cardiac defect that made
them realize the manipulation of the cardiac feedback.
A sensitivity analysis with G*Power 3.1 (Faul et al., 2007) showed that the final sample size
of 34 provides .90 power to detect effects with Cohen's $d = 0.57$ (SESOI of $\delta = 0.34$). Following
recommendations, we do not estimate power based on previously reported effect sizes as this
neglects uncertainty around reported effect size measurements, especially for new effects for which
no reliable effect sizes can be estimated across studies (Albers & Lakens, 2018; Anderson et al.,
2017). Instead, we report sensitivity analyses that reveal the effect sizes that the crucial comparison
between faster and slower feedback is well-powered to detect, given our experimental parameters
(i.e., target power, sample size and type of test), as well as the smallest effect size of interest it can,
in principle, reveal (SESOI, Lakens, 2022). Note that these values are well below the observed top-
down effects on pain ratings of $d = .7$ reported before (e.g., Iodice et al., 2019).
Apparatus
Painful stimuli were electrical pulses delivered using a constant-current electrical stimulator

(Digitimer DS7A) controlling a pair of neurological electrodes attached to the phalanx of the middle
finger of the participant's left hand, which provide a precise constant current, isolated stimulus,
controllable in pulse duration and amplitude. The duration of the electrical stimuli (2 ms) was kept
fixed over the experiment and it was established during a calibration phase before the experiment.

Cardiac recording was performed by a Biopac MP 160 system (Biopac Systems Inc., USA).
ECG was recorded continuously from two electrodes attached to the lower ribs and one over the
right mid-clavicle bone (reference electrode). The ECG signal was sampled at 2 kHz with the
Biopac Acqknowledge 3.7.1 software (Biopac Systems Inc., USA) according to the manufacturer's
guidelines. The ECG signal was then fully analysed in MATLAB (R2020a). The tone used for the
creation of the feedback was the sound of a single heartbeat, gathered from https://freesound.org
and manipulated in Audacity. The feedback audio was then created in MATLAB (R2020a),
repeating the single heartbeat sound according to the desired frequency (see *Procedure*).

Stimulus presentation was controlled through E-Prime (Psychology Software Tools Inc.,
Pittsburgh, USA), which interfaced with the pain stimulators and the Biopac system via a parallel
port.

Procedure

Upon arrival at the lab, participants were briefed by the experimenter. After providing
consent, they were placed in a comfortable chair with their left hand placed on a table, and the ECG
electrodes were applied after cleaning the skin. To increase the ambiguity and thus predictive
influences acting on the pain stimulus (Yon & Frith, 2021), the intensity of the noxious input varied
between five intensities, which were identified for each participant in an initial calibration session.
To do so, participants were informed that they would undergo a psychophysical calibration
procedure to determine their subjective response to increasing stimulus intensities. The first
stimulus was delivered at a low intensity, which is below the threshold for pain perception in most
people. The intensity increased in a ramping procedure up to a maximum of five volts. Participants
verbally rated the pain intensity for each stimulus on a 0-100 Numerical Pain Scale (NPS).

Following previous research (Atlas et al., 2014; Colloca et al., 2006; Hird et al., 2019) a pain
intensity rating of NPS 20 denoted "just painful", NPS 50 denoted "medium pain", and NPS 80
marked the point at which the stimulus was "just tolerable". We identified a 'low' pain level of 10,
a 'low-medium' pain level of 30, a 'medium' pain level of 50, a 'medium-high' pain level of 70,
and a 'high' pain level of 90. Level 100 was considered the point where the participant did not wish
to experience a higher stimulation level in the experimental session and was not used. We repeated
this procedure three times and computed the average stimulus intensities over these three repetitions
corresponding to NPSs 10, 30, 50, 70, and 90.

Participants then underwent a pre-experiment test procedure: stimulus intensities
corresponding to their pain intensity ratings NPS 10 to 90 were delivered in a pseudo-randomised
order four times and participants were instructed to identify the intensity of each pulse. Participants
had to correctly identify 75% of stimulus intensities to proceed to the main experiment. If they did
not achieve this in the test procedure, the intensities were adjusted, and the test was repeated until
participants correctly identified 75% of stimulus intensities (Hird et al., 2019). Participants were
excluded if the correlation between their ratings of the five levels of the nociceptive stimulus
intensities and the actual stimulation intensities within the baseline phase was below $r=0.75$,
suggesting insensitivity to the varying nociceptive degrees of the electrical stimulus, and an
essentially flat response profile. No participants were excluded with this criterion. This procedure
was needed because, although the paradigm introduced ambiguity by using different levels of actual

stimulus intensity, it was important to ensure that participants developed a stable internal
representation of the pain scale and to ensure that illusion effects were not due to an inability to
reliably distinguish between intensities. Note that this correlation criterion only requires that all
participants' ratings generally increase with actual stimulus intensities; participants are still free to
vary in absolute values and the range they use on the pain scale. Supplementary Figure 11 shows
the stimulation intensities corresponding to each of the five pain levels obtained through this
calibration.

Having established the individual levels of intensity for each stimulation, participants were
informed that their task was to rate the intensity and unpleasantness of nociceptive stimulations. We
carefully described the distinction between intensity and unpleasantness ratings using the standard
language developed by Price et al. (1989), emphasizing that pain intensity and unpleasantness
should be rated independently.

The experiment started with a no-feedback session in which participants' heart rate in
anticipation of the shock (measured with ECG) and perceived level of pain (measured with Numeric
Pain Scale and Likert scale) were assessed. These measures were used to normalize our dependent
variables for our analysis (see *Data analysis*).

Participants were seated in front of the computer, with the ECG electrodes attached, their
left hand placed on the table with the electrodes of the cutaneous electrical stimulation fixed on the
phalanx of the middle finger, and their right hand placed on the mouse, ready to start the task. The
ECG was recorded for the entire session.

Each trial of the no-feedback phase started with the presentation of a fixation cross
appearing on the screen. After 60 seconds, a single electrical shock was administered, randomly
varying the level of intensity in each trial - as established in the previous calibration phase (i.e.,
NPS 10, 30, 50, 70, and 90). After each stimulation, participants first rated the intensity of the
painful stimulus (instruction: "How intense was the painful stimulation?") on a continuous
Numerical Pain Scale (NPS) appearing on the screen, from 0 ("not at all painful") to 100
("extremely painful"). They then rated the unpleasantness of the painful stimulus (instruction:
"How unpleasant was the painful stimulation?") on a 5-point Likert scale (1=no pain, 2=weak,
3=moderate, 4=severe, 5=extremely severe). The two scales were presented on the screen in
succession, and participants made their judgments through a mouse click on the scales. Between
trials, a pause screen was shown for 45 seconds, in order to avoid the subsequent trial to be
contaminated by the heart rate response to the shock. Then, a new screen appeared with the
instructions to press the spacebar to start the new trial.

The no-feedback phase comprised 10 trials, after which participants were asked to wait and
rest for 15 minutes before participating in the second experimental feedback phase. This time was
used by the experimenter to prepare the acoustic stimuli for use in the second session for each
participant. For this purpose, we assessed the individual mean heart rate during the anticipation of
the painful stimulus by measuring the mean R-R intervals of the ECG for the whole 1-minute
interval preceding the shock, and by transforming them into frequency (1/R-R, Colloca et al., 2006).
The congruent feedback consisted of tones of heartbeats reproduced at the same individual HR
frequency recorded in the no feedback phase. The incongruent faster and slower feedback consisted
of tones of heartbeats reproduced at a frequency corresponding to an R-R interval obtained by either
reducing or increasing the length of the original mean R-R interval over the no-feedback phase of
25%, respectively. The choice of a ±25% variation in heart rate feedback was guided by prior
research using false cardiac feedback paradigms. Studies in this field have used a range of

modulation strategies, from fixed-frequency symbolic feedback (e.g., Valins, 1966; Tajadura-Jiménez et al., 2008) to real-time individualised adjustments without specific percentage values (e.g., Iodice et al., 2019). More recent work has adopted proportional manipulations of instantaneous heart rate, using variations such as -20% (Azevedo et al., 2017), ±30% (Dey et al., 2018), and ±50% (Gray et al., 2007). Our choice of ±25% falls within this established range, aiming to strike a balance between producing a salient modulation and preserving the plausibility of the feedback as a reflection of one's actual physiology. Specifically, the generation of feedback was tailored to each participant. The feedback was set to be 25% above or below their baseline heart rate, with the feedback gradually increasing or decreasing. This individualized approach ensured that each participant experienced feedback relative to their own baseline heart rate. Moreover, to make the incongruent feedback manipulation credible, the incongruent acoustic feedback always started at a frequency rate that reproduced the individual's heart rate recorded over the no feedback phase, to then gradually increase or decrease the R-R length according to whether the feedback was slower or faster, respectively. In the feedback phase, participants were informed that their task was similar to the previous no-feedback procedure, namely, to rate the intensity and unpleasantness of pain stimuli. They were also informed that in the 60 seconds preceding the administration of the shock, they would hear acoustic feedback, which was equivalent to their ongoing heart rate. This phase consisted of 18 trials (6 trials per experimental condition) of 60 seconds (i.e., fixation) each, after which the electrical shock was administered, randomly varying its intensity in each trial (i.e., NPS 30, 50, and 70), either congruent, slower or faster than the participant's heart rate recorded in the no-feedback phase.

This relatively low number of repetitions was a deliberate design choice based on both theoretical and practical considerations. As observed in other body illusion paradigms (e.g., Botvinick & Cohen, 1998; Ehrsson, 2007; Pratviel et al., 2022), repeated exposures can reduce illusion strength due to habituation, decreased plausibility, or increased awareness of the manipulation. Moreover, sessions lasted approximately 1.5 to 2 hours and involved cognitively demanding tasks, limiting the feasibility of increasing trial numbers without introducing participant fatigue or disengagement. Finally, as we used explicit pain ratings - measures typically less noisy than implicit physiological indices - fewer trials were sufficient to detect reliable effects (e.g., Corneille et al., 2024).

As for the use of only these three pain intensities in the test phase, the rationale was to focus on a manageable subset that still covered a meaningful portion of the stimulus spectrum. Following the approach of Iodice et al. (2019, PNAS), we deliberately excluded the extreme calibrated levels (NPS 10 and NPS 90) to avoid floor- and ceiling-effects. Moreover, by restricting the range we ensured that every test intensity could be paired with both a "slower" and a "faster" cardiac-feedback condition derived from an adjacent level - pairings that would have been impossible at the extremes where no neighbouring level exists.

Specifically, each of the three levels of intensity of the nociceptive stimulus was associated with one of the three different types of acoustic feedback from the previous no-feedback phase. As in the previous no-feedback phase, participants' individual mean heart rate before the painful stimulus (i.e., ECG) and the perceived intensity and unpleasantness of pain (i.e., NPS and Likert scale) were assessed. The order of the experimental conditions (i.e., congruent, slower and faster) was randomly generated for each participant by a web-based computer program (www.randomization.com).

369
370
Figure 1. **Trial timeline of the experiment**. Participants are seated with the electrodes of
both the electrical stimulation and the ECG attached. Each trial starts with a fixation cross (60
seconds), after which the noxious stimulation is administered and the pain intensity (i.e., Numerical
Rating Scale) and unpleasantness (i.e., Likert scale) ratings are collected. After each trial, a pause
screen of 45 seconds was shown (i.e., ISI, Inter Stimulus Interval), after which participants
proceeded to the next trial by pressing the spacebar. The trial timeline was identical for both the no-
feedback and feedback phases, with the only exception that in the feedback phase participants were
provided with the acoustic feedback (i.e., slower, faster, congruent) about their HR, reproduced for
the whole fixation period (60 seconds) preceding the shock.
380
Body Perception Questionnaire
After the experiment, all participants completed the Body Perception Questionnaire (Short form
BPQ- SF) (Poli et al., 2021), to investigate whether either perceptual or autonomic modulation as
induced by our task would be predictive of self-reported measures of bodily awareness and
reactivity. The Body Perception Questionnaire is a 22-item self-administered questionnaire that

assesses awareness and reactivity of the autonomic nervous system, that is, the subjective ability to
perceive bodily states and bodily reactions to stress. High scores on the BPQ reflect high awareness
of internal bodily signals (i.e., high interoceptive sensibility) and high perceived reactivity of the
visceral nervous system. Items ask participants to rate, on a 5-point scale (from 1 = never to 5 =
always), the frequency with which they feel aware of bodily sensations (e.g., body awareness
subscale "My mouth being dry"), experience supradiaphragmatic reactivity (e.g.,
supradiaphragmatic reactivity subscale "I feel shortness of breath"), and subdiaphragmatic
reactivity (e.g., subdiaphragmatic reactivity subscale reactivity subscale "I have indigestion"). In
this work, we focus on the body awareness and supradiaphragmatic reactivity subscales, following
prior research (Petzschner et al., 2019). Body awareness plays a critical role in how individuals
perceive and interpret bodily signals, which in turn affects emotional regulation and self-awareness.
Supradiaphragmatic reactivity refers specifically to organs located or occurring above the
diaphragm - such as the heart - compared to subdiaphragmatic reactivity subscales further down.
Our decision to include these subscales was further supported by recent work from Petzschner et al.
(2021), showing that supradiaphragmatic reactivity predicts attentional modulation of the heartbeat-
evoked potential (HEP), an electroencephalographic signature of the heartbeating. Thus, this
subscale, and the more general body awareness scale, most closely reflect the interplay between
cardiac attention, bodily awareness and the development of the interoceptive illusion.
404
Data Analysis
Measures of the heart rate recorded with the ECG (beats per minute) in the feedback phase
were normalized relative to the baseline (i.e., no-feedback phase) by subtracting and then dividing
by the baseline (Lambert et al., 2024; Mirmoosavi et al., 2024; Sundararaj et al., 2025) using the
formula: normalized value = (X- bX)/bX, where X represents the mean value of each measure
assessed in the experimental feedback phase and bX the mean value of the measure calculated over
the no-feedback phase. As in prior research (e.g., Bartolo et al., 2013; Cecchini et al., 2020; Riello
et al., 2019), the same normalization was also applied to pain intensity and unpleasantness
measures. We did not have specific hypotheses about how the different levels of noxious stimulus
intensity affect the perceptual illusion.
For all three dependent variables (heart rate, pain intensity, pain unpleasantness), our main
hypothesis is tested by comparing the effects of faster against the effects of slower feedback.
Statistically, this comparison collapses our prediction onto a single and most powerful test of our
hypothesis for each of the dependent measures, as faster and slower feedback are assumed to
emerge from changes in opposite direction, in contrast to separate comparisons against the
congruent feedback for which effect sizes will by necessity be on average half of the combined
influence of the faster vs slower comparison. Moreover, as the slower and faster feedback
manipulation reflect analogous deviations from the real heart rate, this comparison keeps any effect
of a perceived absolute deviation (in either direction) from the real heart rate constant. Assume, for
example, that in one participant, heart rates generally decrease as soon as a deviation from the real
heart rate is noticed (i.e., in either direction). The comparison between faster and slower feedback
will still measure whether this decrease is larger for faster than slower feedback, just as in a
participant who does not show such a general decrease or one who shows an increase.
The data from each of our dependent measures (heart rate, pain intensity ratings, pain
unpleasantness ratings) were analysed similarly. First, to confirm that the relevant measure of
interest was affected by the different types of heart rate feedback, the data were entered into a 1-

factor repeated measures ANOVA (Greenhouse-Geisser corrected) with the levels faster feedback,
congruent feedback, and slower feedback.
Second, once such an influence was confirmed, a planned paired t-test tested our main
prediction: that there would be a difference in objective heart rate as well as subjective measures of
pain intensity and pain unpleasantness between trials with faster and slower feedback.
Finally, we tested whether any such effect was mostly driven by the faster instead of the
slower feedback, using separate t-tests comparing the faster feedback against the congruent
feedback and the slower feedback against the congruent feedback.
Please note that, as noted above, this latter comparison is not necessary to test our
hypothesis, which only requires a difference between faster and slower feedback. Indeed, many
influential studies omitted such a control entirely (Hill et al., 2024; Makkar & Grisham, 2013;
Phillips et al., 1999; Valins, 1966), or if included, typically treat it as a baseline control (e.g.,
Patchitt et al., 2025). Nevertheless, we included the congruent condition here to provide an
informative baseline reference, and situating our findings clearly within the broader experimental
context, which suggests a specific role of heartrate deviations upwards (i.e. faster) than downwards
(slower) (Iodice, 2019). For all analyses, the assumption of normality was systematically assessed
using the Kolmogorov-Smirnov (K-S) test.
448
449 **Results**
450
ECG Heart Rate measurement
We first tested whether the heart rate feedback was treated as an interoceptive stimulus and
induced a compensatory adjustment of the real heart rate. If so, faster feedback should elicit a
comparative slowdown of real heart rates compared to slower feedback (see Figure 2c). An initial
ANOVA confirmed that the different types of heart rate feedback (faster, congruent, slower)
affected heart rates, $F(2, 66) = 4.071$, $p = .022$, $\eta p^2 = .110$. A planned paired sample t-test showed
that heart rates decreased during faster cardiac feedback compared to slower heart rate feedback,
$t(33) = 2.074$, $p = .046$, $d = .36$ (Figure 2c). When compared with congruent feedback, significant
heart rate differences emerged for faster feedback, $t(33) = 2.228$, $p = .033$, $d = .38$, but not for
slower feedback, $t(33) = .92$, $p = .364$, $d = .16$.
461
Subjective pain intensity ratings
Participants' pain intensity ratings were analysed analogously. If faster heart rates are
treated as an interoceptive signal of threat, then faster cardiac feedback should lead to increased
pain intensity ratings compared to slower feedback. A repeated measures ANOVA with three levels
(faster feedback, congruent feedback, slower feedback) revealed that the types of heart rate
feedback affected pain intensity ratings, $F(2, 66) = 7.017$, $p = .002$, $\eta p^2 = .175$. A planned t-test then
revealed that participants' perception of pain intensity increased after faster relative to slower
cardiac feedback, $t(33) = 3.339$, $p = .002$, $d = .573$ (Figure 2a). Additional t-tests showed that, like
in the heart rate recordings, this difference was largely driven by the faster feedback. When
compared to congruent feedback, significant differences emerged for faster feedback, $t(33) = 2.66$,
$p = .012$, $d = .457$, but not for the slower feedback, which showed a slight numerical decrease, $t(33)$
$= .477$, $p = .636$, $d = .082$.
474

Subjective pain unpleasantness ratings

Analogous tests run on pain unpleasantness ratings replicated all findings. The initial
ANOVA confirmed that the different types of cardiac feedback (faster, congruent, slower) affected
pain unpleasantness ratings, $F(2, 66) = 6.092$, $p = .004$, $\eta p^2 = .156$. A paired sample t-test showed
that pain unpleasantness ratings were higher after faster feedback than slower cardiac feedback,
$t(33) = 2.771$, $p = .009$, $d = .48$ (Figure 2b). When compared to congruent feedback, significant
differences emerged for faster feedback, $t(33) = 3.045$, $p = .005$, $d = .522$, but not for the slower
feedback, which showed a slight numerical decrease, $t(33) = .206$, $p = .838$, $d = .035$.

485
486

487   *Figure 2.* **a** Intensity pain ratings (Numeric Pain Scale), **b** Unpleasantness ratings (5-points Likert
488   Scale), **c** Real cardiac frequency. Values consist of the central tendency of change of pain intensity (a),
489   unpleasantness (b) and heart rate (c) associated with the exposure to the faster, slower and congruent cardiac
490   feedback, relative to the no-feedback. Values of zero on the vertical axis would represent no change relative
491   to the exposure to the no-feedback. Positive and negative values would represent an increase and a decrease,

respectively. Brackets denote significance: = significant one-way RM-ANOVA; short bracket = planned Faster–Slower contrast. Asterisks indicate two-tailed p values (*< .05, ** < .01, *** < .001, "n.s." = not significant).

Body Perception Questionnaire

The Short form Body Perception Questionnaire (Poli et al., 2021), administered at the end of the experiment, allowed us to explore whether individual differences in body awareness and supradiaphragmatic reactivity were related to the extent to which participants developed (1) the perceptual illusion of pain (2) the physiological changes in their cardiac frequency. Overall, the average score obtained at the Body awareness subscale and Supradiaphragmatic subscale was 5.22 (SD = 1.62) and 2.05 (SD = 1.96), respectively.

We examined Pearson's correlations between BPQ scores and changes in heart rate, pain intensity, and pain unpleasantness in response to the *faster vs. slower* feedback conditions - our main contrast of interest which was a key driver of effects. No significant associations were found between either BPQ subscale and the change in heart rate (all p > 0.05). Likewise, no correlations were observed between the BPQ subscales and changes in pain unpleasantness or pain intensity across these two conditions (all p > 0.05).

Overall, these findings do not suggest reliable association between individual differences in self-reported body perception and the perceptual or physiological effects of the experimental manipulation. Please note that the administration of this questionnaire was fully exploratory, as the current sample size is likely underpowered for robust individual difference analyses, which typically require substantially larger samples to detect meaningful associations (Hedge et al., 2018; Schönbrodt & Perugini, 2013).

Experiment 2

Experiment 1 established that false accelerated cardiac feedback induced both an illusory increase in the perception of pain and a modulation in the pain-anticipatory related cardiac state (i.e., heart rate) towards the expected level of the noxious stimulus, based on the frequency rate of the cardiac feedback. Experiment 2 tests whether the same perceptual and physiological changes are observed if participants are exposed to non-interoceptive feedback (i.e., unrelated to biological human sounds), which should be less likely to induce expectations of threat. To address this question, we substituted the interoceptive cardiac feedback with an exteroceptive tone but kept the experiment otherwise identical. Thus, as in Experiment 1, participants' pain intensity (i.e., Numeric Pain Scale), unpleasantness (i.e., Likert scale) ratings and cardiac frequency (i.e., ECG) were collected both in a no-feedback as well as in a feedback phase, which could be either congruent, faster or slower than participants' individual heart rate recorded over the no-feedback phase. We reasoned that participants would have no expectation that an external acoustic sound signals a preparation of their body to a threatening, high painful stimulus. Consequently, perceptual and cardiac modulations associated with the feedback manipulation should be reduced or eliminated with exteroceptive feedback.

Methods

Participants

Thirty-five participants (mean age 25.17, SD = 4.02, 23 women) were recruited from Gabriele D'Annunzio University and the wider community. All were right-handed with normal or corrected-to-normal vision. Exclusion criteria were identical to Experiment 1. All gave written

informed consent, were unaware of the purposes of the study, and were fully debriefed at the end of
the experiment. Ethical approval from the local ethics board was obtained.
As in Experiment 1, participants were excluded if the correlation between the desired five
levels of the nociceptive stimulus intensity and their ratings in the no-feedback condition was below
r=0.75, suggesting insensitivity to the varying nociceptive degrees of the electrical stimulus. Five
participants were excluded with this criterion. Moreover, participants were excluded if, for any
reason, they linked the exteroceptive sound to a simulation of an interoceptive, cardiac acoustic
feedback, in their post-experiment debriefing. One additional participant was excluded with this
criterion. Hence, the final sample size was 29. A sensitivity analysis with G*Power 3.1 (Faul et al.,
2007) showed that a sample size of 29 provides .90 power to detect effects with Cohen's d >.62
(SESOI of $\delta = 0.37$).
Apparatus and Stimuli
The apparatus and stimuli were the same as Experiment 1. The only difference was the
selection of the tone used to present the heart rate feedback. Instead of using a heartbeat, the sound
was a single percussion obtained by knocking two woods, gathered from https://freesound.org and
manipulated in Audacity. The feedback audio was then created in MATLAB (R2020a), repeating
the single tone according to the desired frequency (see *Procedure*).
Procedure
The procedure used in Experiment 2 was the same as Experiment 1, with the only exception
that the sound used for the feedback was an exteroceptive, instead of interoceptive (i.e., cardiac)
tone. Unlike in Experiment 1, where participants were told they would hear their own heartbeats in
real time, in Experiment 2 participants were not informed that the sound was meant to represent
their heart rate.
Data Analysis
Data analysis was identical to Experiment 1. Note that we predicted an absence/reduction of
effects here, compared to the findings of Experiment 1. In standard analyses, a non-significant
result does not allow one to conclude that there is no effect; it simply indicates a lack of evidence
for a statistically detectable difference. The predicted reduction/absence of effects with
exteroceptive cues here relative to the interoceptive cues in Experiment 1 were tested in two ways.
First, to test for the predicted absence of effects with exteroceptive cues, equivalence tests
(i.e., TOST 'two one-sided t-tests' procedure; Lakens, 2017; Lakens et al., 2018) were performed
for the differences between each dependent (i.e., pain unpleasantness, pain intensity, heart rate).
The TOST procedure addresses the above limitation by assessing whether a given (typically non-
significant) effect is smaller than the effective smallest effect size of interest (SESOI) that the
current design has power to detect, using two one-sided t-tests against positive and negative
equivalence bounds. If both tests are statistically significant, it confirms that the differences in heart
rate, pain intensity and pain unpleasantness induced by the target conditions (faster and slower
feedback) are smaller than the smallest effect size of interest detectable in the current study.
Second, to confirm that the exteroceptively induced changes in heart rate and pain
perception differ from those with interoceptive stimuli in Experiment 1, we compared our main
contrast of interest - each participant's difference between faster and slower feedback in each of the
three dependent measures - between the two experiments, with independent-samples t-tests. A
significant result from this t-test shows that the changes induced by faster vs slower exteroceptive
feedback here differ from the analogous changes induced by interoceptive feedback in Experiment
1.

Finally, we checked whether the different sounds (interoceptive heartbeat sounds in
Experiment 1, exteroceptive control sounds in Experiment 2) induced any overall changes in heart
rates, painfulness or unpleasantness, which would be expected if these sounds differ systematically
in the amount of arousal or alertness they induce, using between-group t-tests on overall heart rate,
painfulness or unpleasantness averaged across all feedback conditions (equivalent to a main effect
of Experiment in an ANOVA).
588
Results
ECG Heartrate measurement
As in Experiment 1, we first tested whether participants' heartrate was affected by the
feedback manipulation, now using non-interoceptive stimuli, using a 1-factor repeated measures
ANOVA with three feedback levels (faster, congruent, slower). In contrast to Experiment 1, and as
hypothesized, no robust differences between the three conditions emerged, $F(2, 56) = 2.54$, $p =$
$.088$, $\eta p^2 = .083$. Our main hypothesis test, which directly compares cardiac frequency in trials with
faster and slower feedback revealed no robust differences either, $t(28) = 1.48$, $p = .317$, $d = .19$. A
TOST equivalence test was conducted to determine whether the difference in heart rate between the
Incongruent Faster and *Incongruent Slower* conditions was statistically equivalent within a range of
$d = -0.57$ to $d = 0.57$. The observed effect size was $d = -0.19$ (90% CI [-0.36, -0.03]). Both one-
sided tests were significant, $t(28) = 2.05$, $p = .025$; $t(28) = -3.71$, $p < .001$, indicating statistical
equivalence between conditions, relative to the smallest detectable differences in our design.
Note that, if anything, the induced difference in heart rate went in the numerically opposite
direction than in Experiment 1, showing a heart rate *increase* after faster compared to slower
exteroceptive feedback, in contrast to the decrease induced by interoceptive feedback cues. Indeed,
comparing this increase to the equivalent change in Experiment 1 with a between-groups t-test
revealed robust differences, $t(61) = 2.187$, $p = .033$, $d = .553$. Thus, while faster vs. slower
interoceptive heart rate feedback led to compensatory changes in participants' own heart rate, no
such effects were obtained for exteroceptive feedback.
As in Experiment 1, we also investigated whether any differences were apparent when faster
and slower feedback were compared to congruent feedback. No significant differences emerged for
faster feedback $t(28) = 1.202$, $p = .239$, $d = .22$. However, the slower feedback induced a decrease
$t(28) = 2.291$, $p = .030$, $d = .43$, again showing the opposite pattern than in Experiment 1.
Finally, we checked whether the interoceptive and exteroceptive stimuli in Experiments 1
and 2 induced overall changes in heart rate response, but no such differences were apparent, $t(61)$
$=.140$, $p = .889$, $d = .035$.
616
Subjective pain intensity ratings
Data were analysed as in Experiment 1. We first tested whether pain intensity ratings were
generally affected by the feedback manipulation, using a 1-factor repeated measures ANOVA with
the levels (faster feedback, congruent feedback, slower feedback). Consistent with the results of the
heart rate data, there was no robust difference, $F(2, 56) = 2.54$, $p = 0.088$, $\eta p^2 = .083$. In particular,
there was no difference in pain ratings between trials with faster and slower feedback, $t(28) = .64$, p
$= .53$, $d = .12$, and a A TOST equivalence test was conducted to examine whether the difference in
external NPS ratings between the *faster* and *slower* conditions was statistically equivalent within
bounds of Cohen's d = ±0.57. The observed effect size was $d = -0.12$ (89% CI [-0.34, 0.10]). Both
one-sided tests were significant, $t(28) = 2.43$, $p = .011$; $t(28) = -3.33$, p = .001, indicating that the

effect was statistically equivalent within the bounds of our experimental design.
When the difference in pain intensity ratings induced by faster and slower exteroceptive
feedback was compared to that induced by interoceptive feedback in Experiment 1, a significant
difference emerged, $t(61) = 2.052$, $p = .044$, $d = .519$. Thus, while faster (relative to slower) heart
rate feedback increased pain ratings, delivering the same feedback through an exteroceptive
stimulus induced no such changes.
Relative to congruent feedback, intensity ratings were increased for both faster and slower
feedback, even though the difference was robust only for the faster feedback, $t(28) = 2.12$, $p =$
0.043, $d = 0.39$, not the slower cardiac feedback, $t(28) = 1.58$, $p = 0.13$, $d = 0.29$.
Finally, we checked whether the interoceptive and exteroceptive stimuli in Experiments 1
and 2 induced overall changes in pain intensity ratings, but no such differences were apparent, $t(61)$
$=.538$, $p = .593$, $d = .136$.
639
640
641
Subjective pain unpleasantness ratings
Pain unpleasantness ratings showed the same pattern as pain intensity ratings. First, a 1-
factor repeated measures ANOVA with the levels faster feedback, congruent feedback, and slower
feedback revealed no robust differences, $F(2, 56) = 1.01$, $p = .371$, $\eta p^2 = .035$. Moreover, there was
no difference between trials with faster and slower feedback, $t(28) = .864$, $p = .395$, $d = .16$ (Figure
3b) and the A TOST procedure (Lakens, 2017) was conducted to test whether the difference in
external liking ratings between the *faster* and *slower* conditions was statistically equivalent within
bounds of Cohen's d = ±0.57. The observed effect size was d = -0.02 (89% CI [-0.05, 0.02]). Both
one-sided tests were significant, t(28) = 2.21, p = .018; t(28) = -3.56, p < .001, indicating statistical
equivalence between the two conditions within the bounds of our experimental design.
Comparing these effects of exteroceptive feedback to the same stimulation delivered
interoceptively in Experiment 1 revealed suggestive evidence for a difference in line with the
previous measures, but it was not significant, $t(61) = 1.679$, $p = .098$, $d = .424$.
Further simple t-tests showed that, relative to the congruent feedback, differences did neither
emerge for faster feedback, $t(28) = 1.428$, $p = .164$, $d = .27$, nor for the slower feedback, $t(28) =$
$.571$, $p = .573$, $d = .06$.
Finally, we checked whether the interoceptive and exteroceptive stimuli in Experiments 1
and 2 induced overall changes in pain unpleasantness ratings, but no such differences were
apparent, $t(61) =.435$, $p = .665$, $d = .110$.

662
663

Figure 3. **a** Intensity pain ratings (Numeric Pain Scale), **b** Unpleasantness ratings (5-points Likert
Scale), **c** Real cardiac frequency. Values consist of the central tendency of change of pain intensity (a),
unpleasantness (b) and heart rate (c) associated with the exposure to the faster, slower and congruent
exteroceptive feedback, relative to the no-feedback. Values of zero on the vertical axis would represent no
change relative to the exposure to the no-feedback. Positive and negative values would represent an increase
and a decrease, respectively. Brackets denote significance: = significant one-way RM-ANOVA; short
bracket = planned Faster–Slower contrast. Asterisks indicate two-tailed *p* values (*< .05, ** < .01, *** <
.001, "n.s." = not significant).

Supplementary analyses

To rule out potential confounds and confirm the robustness of our findings, we conducted additional
analyses on the raw data (see *Supplementary materials*), using the mixed linear models approach,

which offer increased power and reliability with limited trials and conditions (Kliegl et al., 2011;
Baayen et al., 2010; Ambrosini et al., 2019). These analyses included all levels of feedback for
heart rate and pain ratings (unpleasantness and intensity), stimulus intensity levels, as well as trial
number (i.e., the rank order of each trial) as a variable to assess potential time-related effects such
as learning, adaptation, or fatigue (e.g., Möckel et al., 2015). In addition, we incorporated the initial
no-feedback phases as condition to test for potential baseline differences, and we analyzed all
stimulus intensities separately to test whether feedback effects were specific to particular
nociceptive levels or generalized across them. These analyses fully replicate the findings of the
main analysis reported above, but offer more detailed insights for interested readers into other, not
theoretically relevant, factors that may affect the results. Due to the large number of comparisons,
and the associated alpha inflation, these additional, not a priori predicted, results should be
interpreted with caution, however, before being replicated in further studies.

687
General Discussion

Interoceptive inference and the Embodied Predictive Coding frameworks (Barrett, 2017;
Barrett & Simmons, 2015; Pezzulo, 2014; Seth, 2013; Seth & Friston, 2016) propose that
interoception is not merely a passive mirror of the internal milieu but rather the outcome of active,
probabilistic inferences. Through this mechanism, the brain constantly estimates and regulates
homeostatic states by integrating prior expectations with incoming interoceptive, proprioceptive and
exteroceptive signals. To test this framework in the context of pain, participants were given auditory
cardiac feedback, which they believed reflected their actual physiological state, before the
administration of pain stimuli.. Unbeknownst to them, we varied the frequency of this heart rate
feedback, making it either faster or slower. This manipulation aimed to create a false interoceptive
prior, suggesting an increase or decrease in heart rate just before they received a painful stimulus.
In predictive models of interoception, such differences between expected and actual
interoceptive information can be reduced by both changing one's perceptual state to encompass the
new information, and by actively changing the body in response to the unexpected input, by
engaging autonomous reflexes (i.e., perceptual and active inference, Parr et al., 2022, Friston, 2005;
2010; Ainley et al., 2016, Paulus et al., 2019). If so, then sensing an accelerated heart rate
(relatively to a decelerated one) should have two consequences: first, it may falsely lead participants
to expect, and then perceive, a more intense pain stimulus then is really delivered. Second, it may
induce changes in autonomic states themselves, compensating for the perceived changes in heart
rate by actually down-regulating the real heart rate, towards the expected decreasing orienting
cardiac response when expecting threat (Lykken et al., 1972; Taggart et al., 1976; Bradley et al.,
2005; Bradley et al., 2008; Colloca et al., 2012; Tracy et al., 2017).
Experiment 1 provides evidence for both components of the hypothesized response. It
showed, first, that faster cardiac feedback induced an illusory perception of increased
unpleasantness and intensity of subsequent pain stimuli compared to slower cardiac feedback. Thus,
as predicted, the mismatch between people's expectations and the actual input led to changes in the
perception of pain, towards the *expected* input associated with increased heart rates, in line with
Bayesian/predictive processing accounts of pain perception. Second, false cardiac feedback also
changed participants' real heart rate. When hearing faster (relative to slower) feedback, the real
heart rate decreased, assuming the pattern of anticipatory compensatory response typically enacted

when humans prepare to face a threatening stimulus (Lykken et al., 1972; Taggart et al., 1976;
Bradley et al., 2005; Bradley et al., 2008; Colloca et al., 2006; Tracy et al., 2017).
Importantly, both perceptual and cardiac frequency changes after faster (vs. slower)
feedback were not observed when the heartbeat tones were replaced by exteroceptive sounds in
Experiment 2. This finding supports the hypothesis that the observed perceptual and physiological
modulations are specifically tied to interoceptive feedback, which provides evidence of the current
state of the body in preparation for a potential upcoming (threatening) stimulus. In contrast, as the
frequency of an external sound is not informative about the current state of the body, it should not
induce any nociceptive stimulus expectations. Indeed, no discrepancy in the pain unpleasantness
and intensity ratings, as well as in participants' actual heart rates, were observed in Experiment 2.
Importantly, the change from interoceptive to exteroceptive feedback did not induce any overall
changes in pain perception or heart rate between experiments, only affecting the sensitivity to its
rate (i.e., faster or slower). This suggests that the two types of cues were otherwise similarly
processed, and did not vary substantially in overall arousal or sensitivity to pain they induced by
themselves.
To our knowledge, this is the first study to successfully demonstrate that both *perceptual*
and *active inference* components of pain are modulated through false interoceptive feedback. This
provides evidence for the proposed mechanism through which, based on prior knowledge and
expectations, the organism infers and actively implements the interoceptive consequences that have
been predicted (Barrett & Simmons, 2015; Pezzulo, 2014; Seth & Friston, 2016). Our findings
therefore support predictive coding accounts of interoception, which propose such effects on
perception and bodily state, and extend them beyond exteroceptive cues (visual, Wiech et al., 2014;
Jepma et al., 2019; auditory, Colloca et al., 2006; Atlas et al., 2010). They indicate that cardiac
signals are weighted in the brain's generative model of pain and serve as key variables in allostatic
regulation (Barrett et al., 2016), driving both the perception of further interoceptive signals and the
regulation of adaptive bodily states.
A conceptual parallel can be drawn with the Rubber Hand Illusion (RHI), where mismatches
between seen and felt touch are resolved both perceptually and through autonomic adjustments
(Botvinick & Cohen, 1998; Moseley et al., 2008). Crucially, these responses can emerge from
expectations alone (Ferri et al., 2013). Similarly, here, faster cardiac feedback created a prediction
of threat that was resolved perceptually as higher pain ratings and actively as a parasympathetic
deceleration of heart rate. Thus, just as the RHI illustrates predictive integration of visual and
proprioceptive signals, the present findings reveal similar mechanisms in the visceral domain,
showing that interoceptive predictions about cardiac state can reshape both the subjective and
autonomic dimensions of pain.
An interesting observation was that, when compared to congruent cardiac feedback, the
current effects on pain perception and autonomic responses were mainly driven by accelerated, not
decelerated, feedback. Similar asymmetries of faster cardiac feedback have been reported in other
domains. For instance, Iodice et al. (2019) showed that participants overestimate the effort they
exerted when provided with faster cardiac feedback, whereas slower feedback had no effect. We
speculate that, here, the asymmetry arises because increased heart rates are a biologically conserved
(and culturally reinforced) cue for arousal, threat, and pain, which occurs for a wide range of
stressors. Faster feedback therefore, conveys the expectation of pain by mimicking the bodily
response that follows its actual experience (i.e., heart-rate acceleration). In contrast, a slowed heart
rate is the body's intended autonomic response when *expecting* (but not yet experiencing) pain in

both human and non-human species, to prepare for the forthcoming stressor (for reviews, see
Livermore et al., 2021; Roelofs, 2017; Vila et al., 2007). It is considered an evolutionarily
conserved, adaptive response that allows the organism to orient attention, enhance information
processing and action preparation (Hashemi et al., 2019; Klaassen et al., 2021; Lojowska et al.,
2015). In our case, therefore, decelerated cardiac feedback signals that this intended autonomic
response for dealing with the anticipated pain stimulus is already achieved. An accelerated heart
rate, in contrast, signals the opposite, providing interoceptive evidence of a misaligned bodily state
that mandates immediate allostatic responses, in terms of both pain perception and cardiac
regulation (Barrett & Simmons, 2015; Pezzulo, 2014; Seth & Friston, 2016).Our findings therefore
blend well with recent proposals that frame cardiac deceleration under a Bayesian inference
approach. Under these views, the cardiac deceleration mechanism allows to adjust precision of
sensory evidence accumulation relative to precision of bodily information, optimising perception
and action (Skora et al., 2022). Thus, decelerated cardiac activity when exposed to the faster
feedback may reflect an autonomic adjustment to prioritize incoming noxious input information in
light of the expectation of the threatening painful event.
778   Together, our evidence therefore supports the hypothesis that not only interoceptive states
modulate interoceptive inference and perception, but the converse is also true: inference dynamics
can modify or produce new internal states, to the extent that the actual cardiac frequency rate
decreased as it would have done over the anticipation of a real threat. More broadly, our findings
are in line with recent proposals of *interoceptive schemas* (Barca & Pezzulo, 2020; Iodice et al.,
2019; Schoeller et al., 2022; Tschantz et al., 2022), internal models akin to the body schema (Head
& Holmes, 1911), which consist of a central representation of interoceptive variables (e.g., body
temperature, cardiac activity etc.) along with prior assumptions or "set points" for these variables.
This internal model would support homeostatic and allostatic regulation by optimally weighting
multiple (i.e., proprioceptive, interoceptive, exteroceptive) streams of information to predict
incoming interoceptive signals (Barca & Pezzulo, 2020; Iodice et al., 2019; Schoeller et al., 2022;
Tschantz et al., 2022). Interoceptive illusions, i.e., contextual manipulations that shift the precision
(i.e., estimated reliability) of interoceptive expectations (Schoeller et al., 2024), offer a direct probe
of this schema. Our paradigm is one of such manipulation, showing how a transient change in
cardiac evidence reveals how the precision weighting of priors and their prediction errors reshapes
both the perceptual and autonomic dimensions of pain. Such illusions open a window onto the
internal generative models that guide perception and action. In this regard, the contribution of the
study of illusions offers important insights not only for a converging model of brain functioning
under the principle of optimal predictive architectures, but it also provides implications for
understanding psychopathology in terms of aberrant predictions and compounding allostatic
consequences (for a full discussion, see Barrett & Simmons, 2015). Because manipulating
contextual cues selectively alters the precision-weighting of interoceptive expectations, such
illusions constitute the kind of diagnostic "stress tests" that reveal how much confidence the brain
assigns to priors versus bodily evidence, a parameter now believed to underlie dysfunctional
conditions, such as anxiety, PTSD and chronic pain (Schoeller et al., 2022). Our experimental
paradigm provides a probe of this mechanism: the coupled shift in pain ratings and heart-rate
deceleration indexes the degree to which threat-related priors dominate over incoming cardiac
signals. Emerging perspectives (Schoeller et al., 2022) frame precisely this imbalance (i.e.,
excessively precise priors or under-weighted sensory channels) as a form of *false inference* that
underlies psychopathology, and identify interoceptive illusions as tools both for phenotyping such

aberrant precision control and for retraining it through repeated, controlled exposure.
809

Accounting for general unspecific contributions

One possibility is that the different effects in Experiment 1 and 2 may not reflect
interoceptive vs exteroceptive processing, but more general features of the sounds used in both
experiments. For example, the heartbeat sounds' cultural meaning and salience (e.g., as signals of
anxiety or distress), or their acoustic properties (e.g., loomingness, roughness), could have increased
arousal, alertness or attentional engagement more generally. Several aspects of our data argue
against this proposal, however. First, if heartbeat sounds had such general alerting/arousing effects,
they should have induced robust between-experiment differences in pain ratings or heart rates when
such heartbeat (Experiment 1) or non-heart related sounds were presented (Experiment 2).
However, no such differences were obtained for any of our measures (see between-experiment t-
tests in the main analyses; cross-experiment analysis in supplementary analyses). In contrast, all
effects we obtained reflected not the overall presence or absence of cardiac feedback, but responses
to a *change* in cardiac feedback (faster or slower), which was absent for identical changes in
exteroceptive feedback.

Second, one may suspect that the acceleration of cardiac feedback, rather than cardiac
feedback more generally, had simply increased arousal, instead of functioning as a misaligned
interoceptive signal. However, if so, one would expect to see an overall *acceleration* in participants'
actual heart rate, as this is a typical physiological marker of heightened arousal (Azarbarzin et al.,
2014; Wascher, 2021; Yang et al., 2017), not the slowing of the heart rate we had predicted and
observed. Thus, our findings are *opposite* to what would be expected under a generic arousal-based
explanation but consistent with the well-known anticipatory physiological regulatory response
predicted by interoceptive inference frameworks. Note also that for such changes in heart rate to be
obtained, very strong manipulations of threat or arousal are required. For example, in Parrotta et al.
(2024), even highly salient and reliable threat cues (i.e., images that predicted upcoming pain with
100% certainty) did not produce any measurable change in heart rate. Our pattern of results is
therefore not readily accounted for by general arousal or salience mechanisms, but is fully in line
with the intrinsic capacity of cardiac feedback to engage autonomic regulation.

Finally, the idea that the interoceptive relevance of the cue, rather than its low-level acoustic
properties or general human-related salience, is the key driver of the observed effects is also
reflected in neurophysiological studies. For example, fMRI studies have shown that heartbeat
sounds selectively activate interoceptive regions (e.g., anterior insula, frontal operculum; Kleint et
al., 2015), while EEG work (Vicentin et al., 2024) has demonstrated enhanced cortical activation to
faster heartbeats, particularly over frontocentral regions, suggesting enhanced processing in
networks associated with interoceptive attention. Moreover, heartbeat sounds have been shown to
attenuate the auditory N1 component (van Elk et al., 2014), a neural signature typically linked to
self-generated or predicted bodily signals.
846

Limitations and future directions

The current results provide a platform for further investigation of the link between
perceptual and active inference in interoception. As this was the first study using our paradigm, it
was optimised to detect robust within-subject effects of pain expectation. By definition, such robust
within-subject effects minimize between-participant differences and are therefore difficult to link to
cross-participant variability (i.e., the "reliability paradox", Hedge, Powell & Sumner, 2018),

requiring substantially larger samples than used here and in the literature (i.e., N>200). Potentially
informative associations, such as those between anticipatory heart-rate dynamics and pain ratings,
could not be tested here. Similarly, the absence of a robust relationship between pain modulation
and BPQ subscales of body awareness and supradiaphragmatic reactivity should be interpreted with
caution, given that these measures were included for exploratory purposes and not sufficiently
powered to draw strong conclusions, especially about the absence of relationships. Future work
designed and powered for individual differences should determine how interoceptive awareness and
cardiac attention shape pain experience. Such studies could also investigate how the current
findings depend on sex/gender. In our sample, like in most psychology research, women were
somewhat over-represented. Prior research suggests that men and women differ in pain processing
and interoceptive sensitivity (Sorge et al., 2018; Mogil, 2012). Future studies with larger samples
could investigate whether such differences modulate the integration of interoceptive signals and
pain perception, especially as predictive processing accounts assume that the precision of sensory
information determines how much it is weighted against prior expectations (Yon & Frith, 2021).

A promising avenue is integrating perceptual, neuroimaging and physiological measures
(e.g., the Heartbeat Evoked Potential, Petzschner et al., 2019) into computational models of
interoceptive inference (e.g., Allen & Friston, 2018; Owens et al., 2018; Allen et al., 2022; Eckert et
al., 2022). Similar inferential frameworks as proposed here have been implemented in
computational models of cardiac (Smith et al., 2020) and gastrointestinal (Smith et al., 2021)
interoception, thermoregulation (Tschantz et al., 2022), and psychopathological conditions, such as
anorexia nervosa (Barca & Pezzulo, 2020), panic disorder (Maisto et al., 2021), self-injury (Barca et
al., 2023), depression (Barrett et al., 2016; Stephan et al., 2016), and others (Paulus et al., 2019).
Such frameworks would allow latent parameters like prior precision and sensory gain to be
estimated for each participant (Unal et al., 2021), allowing us to test whether individuals who rely
more heavily on cardiac priors also exhibit larger pain illusions and autonomic adjustments, further
enabling individual differences research and extending it to clinical applications.

Model comparison could also establish whether faster cardiac feedback shifts the mean of
the pain prior or instead alters the precision of nociceptive evidence. Systematically manipulating
the precision of priors or prediction errors, or explicitly measuring participants' interoceptive
beliefs and their precision, offers a powerful means to uncover the mechanisms that facilitate the
development of the interoceptive illusion. They may reveal whether perceptual biases in the
experience of pain emerge from overlearned priors that do not update to incoming sensory data, or
whether it is possible to directly act on interoceptive illusions and their autonomic consequences by
changing prior knowledge.

The latter would have far-reaching implications, making it possible to both modify the
perception of our internal milieu and impact autonomic states by simply reconfiguring individuals'
beliefs and knowledge. Such findings would have profound consequences for understanding and
potentially intervening in psychopathological conditions. On this basis, manipulating contextual
cues can selectively up- or down-weight interoceptive expectations. In our paradigm, the coupled
shift in pain ratings and heart-rate deceleration quantifies the extent to which threat-related priors
override incoming bodily signals (i.e., cardiac input), a precision parameter that recent accounts
frame as the very locus of dysfunctional predictive processes in anxiety, PTSD and chronic pain.

From this perspective, the pain illusion documented here, together with other interoceptive
illusions, provide a direct window into the brain's predictive architecture (Barrett & Simmons,
2015), offering a mechanistic framework for understanding psychopathology as a failure to revise

inaccurate interoceptive priors, leading to maladaptive regulation and compounding allostatic
consequences. At the same time, they serve as diagnostic "stress tests," revealing how much
confidence the brain assigns to priors versus bodily evidence and thereby offering a mechanistic
foothold for targeted intervention. Finally, we note that in our unpleasantness scale, the lowest
anchor was labeled "no pain." Although this label was not ideal, as unpleasantness and pain can in
principle be dissociated, the effect of this choice is likely minimal given that unpleasantness ratings
were always provided after intensity ratings, and our main findings rely on the interplay between
intensity measures and physiological cardiac modulations.

Conclusions

This study is the first to show that false cardiac signals reshape both pain perception and
anticipatory autonomic responses. False cardiac feedback amplified pain intensity and
unpleasantness while inducing parasympathetic deceleration of heart rate, revealing how
expectations actively construct interoceptive experiences and drive allostatic adjustments. These
findings support predictive models by emphasizing the pivotal role of interoceptive schemas in
predictive architectures and highlight cardiac signals as key variables in the generative models
governing pain and autonomic regulation. Importantly, the lack of effects with non-interoceptive
feedback highlights the specificity of this mechanism to cardiac interoceptive signals, providing a
platform to explore whether other visceral channels, such as respiratory cues, similarly recalibrate
pain perception and physiological states. Beyond advancing theoretical models of interoception, our
paradigm offers a novel tool for probing the precision-weighting of priors and sensory evidence in
healthy and clinical populations, which is consistent with proposals that interoception provides
actionable targets for mental-health interventions (Nord & Garfinkel, 2022). Future work may
explore its translational potential in understanding and modulating maladaptive pain experiences,
where aberrant interoceptive predictions contribute to chronic pain and related disorders.
922
923

Acknowledgements
The work was funded by Leverhulme Trust grant RPG-2019-248 to Patric Bach, and PhD
studentship awarded to Eleonora Parrotta from the Universities of Plymouth and Aberdeen. This
research received funding from the European Union's Horizon 2020 Framework Programme for
Research and Innovation under the Specific Grant Agreement 945539 (Human Brain Project SGA3)
and 952215 (TAILOR) to Giovanni Pezzulo, the European Research Council under the Grant
Agreement No. 820213 (ThinkAhead) to Giovanni Pezzulo and the PNRR MUR project
PE0000013-FAIR to Giovanni Pezzulo. This work was supported by the "Departments of
Excellence 2018-2022" initiative of the Italian Ministry of Education, University and Research for
the Department of Neuroscience, Imaging and Clinical Sciences (DNISC) of the University of
Chieti-Pescara, and by the "Search for Excellence" initiative of the University of Chieti-Pescara.
Below we list the main authors' contribution, in accordance with the CRediT taxonomy:
Eleonora Parrotta: Conceptualization, Methodology, Software, Formal Analysis, Investigation,
Writing - Original Draft, Visualization.
Patric Bach: Conceptualization, Methodology, Writing – Original Draft, Resources, Supervision,
Project Administration, Funding Acquisition.
Giovanni Pezzulo: Conceptualization, Methodology, Writing - Review & Editing, Resources,

Supervision, Funding Acquisition.

Andrea Zaccaro: Methodology, Software, Formal Analysis, Writing - Review & Editing

Mauro Gianni Perrucci: Methodology, Software, Formal Analysis, Investigation, Writing - Review
& Editing

Marcello Costantini: Conceptualization, Methodology, Writing - Review & Editing, Supervision,
Resources, Project Administration, Funding Acquisition.

Francesca Ferri: Conceptualization, Methodology, Writing - Review & Editing, Supervision,
Resources, Project Administration, Funding Acquisition.

Competing interests

The authors declare no competing financial or academic interest.

Data availability

The data used for the analyses is available upon publication in an open repository at the following
link: https://github.com/eleonorap17/painperception. The behavioural and physiological raw data
can be shared by the corresponding author upon request if data privacy can be guaranteed according
to the rules of the European General Data Protection Regulation (EU GDPR).

References

Ainley, V., Apps, M. A., Fotopoulou, A., & Tsakiris, M. (2016). 'Bodily precision': A predictive
coding account of individual differences in interoceptive accuracy. Philosophical Transactions of
the Royal Society B: Biological Sciences, 371(1708), 20160003.

Albers, C., & Lakens, D. (2018). When power analyses based on pilot data are biased: Inaccurate
effect size estimators and follow-up bias. Journal of experimental social psychology, 74, 187-195.

Allen, M., & Friston, K. J. (2018). From cognitivism to autopoiesis: towards a computational
framework for the embodied mind. Synthese, 195(6), 2459-2482.

Allen, M., Levy, A., Parr, T., & Friston, K. J. (2022). In the body's eye: the computational anatomy
of interoceptive inference. PLoS Computational Biology, 18(9), e1010490.

Ambrosini, E., Peressotti, F., Gennari, M., Benavides-Varela, S., & Montefinese, M. (2023). Aging-
related effects on the controlled retrieval of semantic information. Psychology and Aging, 38(3),
219.

Anderson, S. F., Kelley, K., & Maxwell, S. E. (2017). Sample-Size Planning for More Accurate
Statistical Power: A Method Adjusting Sample Effect Sizes for Publication Bias and Uncertainty.
Psychological Science, 28(11), 1547-1562. https://doi.org/10.1177/0956797617723724

Asai, T. (2015). Illusory body-ownership entails automatic compensative movement: For the
unified representation between body and action. Experimental Brain Research, 233(3), 777-785.
https://doi.org/10.1007/s00221-014-4153-0

Atlas, L. Y., & Wager, T. D. (2012). How expectations shape pain. Neuroscience letters, 520(2),
140-148.

Atlas, L. Y., Bolger, N., Lindquist, M. A., & Wager, T. D. (2010). Brain mediators of predictive cue
effects on perceived pain. Journal of Neuroscience, 30(39), 12964-12977.

Atlas, L. Y., Wielgosz, J., Whittington, R. A., & Wager, T. D. (2014). Specifying the non-specific
factors underlying opioid analgesia: Expectancy, attention, and affect. Psychopharmacology,
231(5), 813-823.

Azarbarzin, A., Ostrowski, M., Hanly, P., & Younes, M. (2014). Relationship between arousal
intensity and heart rate response to arousal. Sleep, 37(4), 645-653.

Azevedo, R., Bennett, N., Bilicki, A., Hooper, J., Markopoulou, F., & Tsakiris, M. (2017). The
calming effect of a new wearable device during the anticipation of public speech. *Scientific reports*,
7(1), 2285.

Baayen, R. H., & Milin, P. (2010). Analyzing reaction times. *International Journal of*
Psychological Research, 3(2), 12-28. https://doi.org/10.21500/20112084.807.

Barca, L., & Pezzulo, G. (2020). Keep your interoceptive streams under control: An active
inference perspective on anorexia nervosa. Cognitive, Affective, & Behavioral Neuroscience, 20(2),
427-440. https://doi.org/10.3758/s13415-020-00777-6

Barca L., Maisto D., Pezzulo G. (2023) Modeling and controlling the body in maladaptive ways: an
active inference perspective on non-suicidal self-injury behaviors. *Neuroscience of Consciousness.*

Barlow, D. H., Chorpita, B. F., & Turovsky, J. (1996). Fear, panic, anxiety, and disorders of
emotion.

Barrett, L. F. (2017). The theory of constructed emotion: An active inference account of
interoception and categorization. Social cognitive and affective neuroscience, 12(1), 1-23.

Barrett, L. F., & Simmons, W. K. (2015). Interoceptive predictions in the brain. Nature reviews
neuroscience, 16(7), 419-429.

Barrett, L. F., Quigley, K. S., & Hamilton, P. (2016). An active inference theory of allostasis and
interoception in depression. Philosophical Transactions of the Royal Society B: Biological
Sciences, 371(1708), 20160011. https://doi.org/10.1098/rstb.2016.0011

Bartolo, M., Serrao, M., Gamgebeli, Z., Alpaidze, M., Perrotta, A., Padua, L., Pierelli, F., Nappi,
G., & Sandrini, G. (2013). Modulation of the human nociceptive flexion reflex by pleasant and
unpleasant odors. PAIN®, 154(10), 2054-2059.

Benson, H., Beary, J. F., & Carol, M. P. (1974). The relaxation response. Psychiatry, 37(1), 37-46.

Botvinick, M., & Cohen, J. (1998). Rubber hands 'feel' touch that eyes see. Nature, 391(6669),
756-756.

Bradley, M. M., Moulder, B., & Lang, P. J. (2005). When good things go bad: The reflex physiology of defense. Psychological science, 16(6), 468-473.

Bradley, M. M., Silakowski, T., & Lang, P. J. (2008). Fear of pain and defensive activation. PAIN®, 137(1), 156-163.

Büchel, C., Geuter, S., Sprenger, C., & Eippert, F. (2014). Placebo analgesia: A predictive coding perspective. Neuron, 81(6), 1223-1239.

Cecchini, M. P., Riello, M., Sandri, A., Zanini, A., Fiorio, M., & Tinazzi, M. (2020). Smell and taste dissociations in the modulation of tonic pain perception induced by a capsaicin cream application. European Journal of Pain, 24(10), 1946-1955.

Colloca, L., Benedetti, F., & Pollo, A. (2006). Repeatability of autonomic responses to pain anticipation and pain stimulation. European Journal of Pain, 10(7), 659-665.

Colloca, L., Petrovic, P., Wager, T. D., Ingvar, M., & Benedetti, F. (2010). How the number of learning trials affects placebo and nocebo responses. Pain®, 151(2), 430-439.

Corlett, P. R., Horga, G., Fletcher, P. C., Alderson-Day, B., Schmack, K., & Powers III, A. R. (2019). Hallucinations and strong priors. Trends in cognitive sciences, 23(2), 114-127.

Craggs, J. G., Price, D. D., Perlstein, W. M., Verne, G. N., & Robinson, M. E. (2008). The dynamic mechanisms of placebo induced analgesia: Evidence of sustained and transient regional involvement. Pain, 139(3), 660-669.

Craig, A. D. (2003a). A new view of pain as a homeostatic emotion. Trends in neurosciences, 26(6), 303-307.

Craig, A. D. (2003b). Interoception: The sense of the physiological condition of the body. Current opinion in neurobiology, 13(4), 500-505.

Craig, A. D. (2008). Interoception and emotion: A neuroanatomical perspective. Handbook of emotions, 3(602), 272-288.

Crombez, G., Eccleston, C., Baeyens, F., & Eelen, P. (1998). Attentional disruption is enhanced by the threat of pain. Behaviour research and therapy, 36(2), 195-204.

Crucian, G. P., Hughes, J. D., Barrett, A. M., Williamson, D. J. G., Bauer, R. M., Bowers, D., & Heilman, K. M. (2000). Emotional and physiological responses to false feedback. *Cortex*, *36*(5), 623-647.

Dey, A., Chen, H., Billinghurst, M., & Lindeman, R. W. (2018, October). Effects of manipulating physiological feedback in immersive virtual environments. In *Proceedings of the 2018 Annual Symposium on Computer-Human Interaction in Play* (pp. 101-111).

Eckert, A. L., Pabst, K., & Endres, D. M. (2022). A Bayesian model for chronic pain. Frontiers in Pain Research, 3, 966034.

Edwards, L., Inui, K., Ring, C., Wang, X., & Kakigi, R. (2008). Pain-related evoked potentials are
modulated across the cardiac cycle. PAIN®, 137(3), 488-494.

Edwards, L., McINTYRE, D., Carroll, D., Ring, C., & Martin, U. (2002). The human nociceptive
flexion reflex threshold is higher during systole than diastole. Psychophysiology, 39(5), 678-681.

Edwards, L., Ring, C., McIntyre, D., & Carroll, D. (2001). Modulation of the human nociceptive
flexion reflex across the cardiac cycle. Psychophysiology, 38(4), 712-718.

Ehrsson, H. H. (2007). The experimental induction of out-of-body experiences. Science, 317(5841),
1048-1048.

Eippert, F., Finsterbusch, J., Bingel, U., & Büchel, C. (2009). Direct evidence for spinal cord
involvement in placebo analgesia. Science, 326(5951), 404-404.

Faul, F., Erdfelder, E., Lang, A., & Buchner, A. (2007). G* Power 3: A flexible statistical power
analysis for the social, behavioral, and biomedical sciences. Behavioral Research Methods, 39 (2),
175-191.

Ferri, F., Chiarelli, A. M., Merla, A., Gallese, V., & Costantini, M. (2013). The body beyond the
body: Expectation of a sensory event is enough to induce ownership over a fake hand. Proceedings
of the Royal Society B: Biological Sciences, 280(1765), 20131140.

Forte, G., Troisi, G., Pazzaglia, M., Pascalis, V. D., & Casagrande, M. (2022). Heart rate variability
and pain: a systematic review. Brain sciences, 12(2), 153.

Friston, K. (2005). A theory of cortical responses. Philosophical transactions of the Royal Society
B: Biological sciences, 360(1456), 815-836.

Friston, K. (2010). The free-energy principle: A unified brain theory? Nature reviews neuroscience,
11(2), 127-138.

Gagne, C., Dayan, P., & Bishop, S. J. (2018). When planning to survive goes wrong: Predicting the
future and replaying the past in anxiety and PTSD. Current Opinion in Behavioral Sciences, 24, 89-
95.

Geuter, S., & Büchel, C. (2013). Facilitation of pain in the human spinal cord by nocebo treatment.
Journal of Neuroscience, 33(34), 13784-13790.

Gray, M. A., Harrison, N. A., Wiens, S., & Critchley, H. D. (2007). Modulation of emotional
appraisal by false physiological feedback during fMRI. *PLoS one*, *2*(6), e546.

Hashemi, M. M., Gladwin, T. E., de Valk, N. M., Zhang, W., Kaldewaij, R., van Ast, V., Koch, S.
B., Klumpers, F., & Roelofs, K. (2019). Neural dynamics of shooting decisions and the switch from
freeze to fight. Scientific reports, 9(1), 4240.

Head, H., & Holmes, G. (1911). Sensory disturbances from cerebral lesions. Brain, 34(2-3), 102-
254.

Hedge, C., Powell, G., & Sumner, P. (2018). The reliability paradox: Why robust cognitive tasks do not produce reliable individual differences. Behavior Research Methods, 50(3), 1166-1186. https://doi.org/10.3758/s13428-017-0935-1

Hill, M. W., Johnson, E., & Ellmers, T. J. (2024). The influence of false interoceptive feedback on emotional state and balance responses to height-induced postural threat. Biological Psychology, 189, 108803. https://doi.org/10.1016/j.biopsycho.2024.108803

Hird, E. J., Charalambous, C., El-Deredy, W., Jones, A., & Talmi, D. (2019). Boundary effects of expectation in human pain perception. Scientific reports, 9(1), 1-13.

Iodice, P., Porciello, G., Bufalari, I., Barca, L., & Pezzulo, G. (2019). An interoceptive illusion of effort induced by false heart-rate feedback. Proceedings of the National Academy of Sciences, 116(28), 13897-13902.

Jennings, J. R., & van der Molen, M. W. (2005). Preparation for speeded action as a psychophysiological concept. Psychological bulletin, 131(3), 434.

Jennings, J. R., Van Der Molen, M. W., & Tanase, C. (2009). Preparing hearts and minds: Cardiac slowing and a cortical inhibitory network. Psychophysiology, 46(6), 1170-1178.

Jepma, M., & Wager, T. D. (2015). Conceptual conditioning: Mechanisms mediating conditioning effects on pain. Psychological science, 26(11), 1728-1739.

Jepma, M., Koban, L., van Doorn, J., Jones, M., & Wager, T. D. (2018). Behavioural and neural evidence for self-reinforcing expectancy effects on pain. Nature Human Behaviour, 2(11), 838-855.

Keltner, J. R., Furst, A., Fan, C., Redfern, R., Inglis, B., & Fields, H. L. (2006). Isolating the modulatory effect of expectation on pain transmission: A functional magnetic resonance imaging study. Journal of Neuroscience, 26(16), 4437-4443.

Kiverstein, J., Kirchhoff, M. D., & Thacker, M. (2022). An embodied predictive processing theory of pain experience. Review of Philosophy and Psychology, 1-26.

Klaassen, F. H., Held, L., Figner, B., O'Reilly, J. X., Klumpers, F., de Voogd, L. D., & Roelofs, K. (2021). Defensive freezing and its relation to approach-avoidance decision-making under threat. Scientific Reports, 11(1), 1-12.

Kleint, N. I., Wittchen, H. U., & Lueken, U. (2015). Probing the interoceptive network by listening to heartbeats: an fMRI study. PloS one, 10(7), e0133164.

Kliegl, R., Wei, P., Dambacher, M., Yan, M., & Zhou, X. (2011). Experimental effects and individual differences in linear mixed models: Estimating the relation of spatial, object, and attraction effects in visual attention. *Frontiers in Psychology, 1*, 238. https://doi.org/10.3389/fpsyg.2010.00238

Koban, L., & Wager, T. D. (2016). Beyond conformity: Social influences on pain reports and physiology. Emotion, 16(1), 24.

Koyama, T., McHaffie, J. G., Laurienti, P. J., & Coghill, R. C. (2005). The subjective experience of pain: Where expectations become reality. Proceedings of the National Academy of Sciences, 102(36), 12950-12955.

Kyle, B. N., & McNeil, D. W. (2014). Autonomic arousal and experimentally induced pain: A critical review of the literature. Pain Research and Management, 19(3), 159-167.

Lacey, J. L. (1970). Some autonomic-central nervous system interrelationships. Physiological correlates of emotion, 205-227.

Lakens, D. (2017). Equivalence Tests: A Practical Primer for t Tests, Correlations, and Meta-Analyses. Social Psychological and Personality Science, 8(4), 355-362. https://doi.org/10.1177/1948550617697177

Lakens, D. (2022). Sample size justification. Collabra: psychology, 8(1), 33267.

Lakens, D., Scheel, A. M., & Isager, P. M. (2018). Equivalence Testing for Psychological Research: A Tutorial. Advances in Methods and Practices in Psychological Science, 1(2), 259-269. https://doi.org/10.1177/2515245918770963

Lambert, T. P., Chan, M., Sanchez-Perez, J. A., Nikbakht, M., Lin, D. J., Nawar, A., Bashar, S. K., Kimball, J. P., Zia, J. S., & Gazi, A. H. (2024). A Comparison of Normalization Techniques for Individual Baseline-Free Estimation of Absolute Hypovolemic Status Using a Porcine Model. Biosensors, 14(2), 61.

Lissek, S., & van Meurs, B. (2015). Learning models of PTSD: Theoretical accounts and psychobiological evidence. International journal of psychophysiology, 98(3), 594-605.

Livermore, J. J., Klaassen, F. H., Bramson, B., Hulsman, A. M., Meijer, S. W., Held, L., Klumpers, F., de Voogd, L. D., & Roelofs, K. (2021). Approach-avoidance decisions under threat: The role of autonomic psychophysiological states. Frontiers in Neuroscience, 15, 621517.

Loeser, J. D., & Treede, R.-D. (2008). The Kyoto protocol of IASP basic pain terminology. PAIN®, 137(3), 473-477.

Lojowska, M., Gladwin, T. E., Hermans, E. J., & Roelofs, K. (2015). Freezing promotes perception of coarse visual features. Journal of Experimental Psychology: General, 144(6), 1080.

Lykken, D., Macindoe, I., & Tellegen, A. (1972). Preception: Autonomic response to shock as a function of predictability in time and locus. Psychophysiology, 9(3), 318-333.

Maisto, D., Barca, L., Van den Bergh, O., & Pezzulo, G. (2021). Perception and misperception of bodily symptoms from an active inference perspective: Modelling the case of panic disorder. Psychological Review, 128(4), 690-710. https://doi.org/10.1037/rev0000290

Makkar, S. R., & Grisham, J. R. (2013). Effects of false feedback on affect, cognition, behavior, and postevent processing: The mediating role of self-focused attention. Behavior therapy, 44(1), 111-124.

Martins, A. Q., Ring, C., McIntyre, D., Edwards, L., & Martin, U. (2009). Effects of unpredictable
stimulation on pain and nociception across the cardiac cycle. PAIN®, 147(1-3), 84-90.

Maselli, A., Lanillos, P., & Pezzulo, G. (2022). Active inference unifies intentional and conflict-
resolution imperatives of motor control. PLOS Computational Biology, 18(6), e1010095.
https://doi.org/10.1371/journal.pcbi.1010095

McIntyre, D., Edwards, L., Ring, C., Parvin, B., & Carroll, D. (2006). Systolic inhibition of
nociceptive responding is moderated by arousal. Psychophysiology, 43(3), 314-319.

Mirmoosavi, M., Aminitabar, A., Mirfathollahi, A., & Shalchyan, V. (2024). Exploring altered
oscillatory activity in the anterior cingulate cortex after nerve injury: Insights into mechanisms of
neuropathic allodynia. Neurobiology of Disease, 190, 106381.
https://doi.org/10.1016/j.nbd.2023.106381

Mischkowski, D., Palacios-Barrios, E. E., Banker, L., Dildine, T. C., & Atlas, L. Y. (2018). Pain or
nociception? Subjective experience mediates the effects of acute noxious heat on autonomic
responses. Pain, 159(4), 699.

Moayedi, M., & Davis, K. D. (2013). Theories of pain: From specificity to gate control. Journal of
neurophysiology, 109(1), 5-12.

Möckel, T., Beste, C., & Wascher, E. (2015). The effects of time on task in response selection-an
ERP study of mental fatigue. Scientific reports, 5(1), 10113.

Montgomery, G. H., & Kirsch, I. (1997). Classical conditioning and the placebo effect. Pain, 72(1-
2), 107-113.

Nord, C. L., & Garfinkel, S. N. (2022). Interoceptive pathways to understand and treat mental
health conditions. *Trends in Cognitive Sciences*, 26(6), 499–513.

Oka, S., Chapman, C. R., Kim, B., Nakajima, I., Shimizu, O., & Oi, Y. (2007). Pupil dilation
response to noxious stimulation: Effect of varying nitrous oxide concentration. Clinical
neurophysiology, 118(9), 2016-2024.

Owens, A. P., Allen, M., Ondobaka, S., & Friston, K. J. (2018). Interoceptive inference: From
computational neuroscience to clinic. Neuroscience & Biobehavioral Reviews, 90, 174-183.

Parr, T., Pezzulo, G., & Friston, K. J. (2022). Active inference: The free energy principle in mind,
brain, and behavior. MIT Press.

Parrotta, E., Bach, P., Perrucci, M. G., Costantini, M., & Ferri, F. (2024). Heart is deceitful above
all things: Threat expectancy induces the illusory perception of increased heartrate. Cognition, 245,
105719.

Patchitt, J., Garfinkel, S., Strawson, W. H., Miller, M., Tsakiris, M., Clark, A., & Critchley, H. D.
(2025). Somatosensory false feedback biases emotional ratings through interoceptive embodiment.
Scientific Reports, 15(1), 11472.

Paulus, M. P., Feinstein, J. S., & Khalsa, S. S. (2019). An active inference approach to interoceptive psychopathology. Annual review of clinical psychology, 15, 97-122.

Pennebaker, J. W., & Lightner, J. M. (1980). Competition of internal and external information in an exercise setting. *Journal of personality and social psychology*, *39*(1), 165.

Petrovic, P., & Ingvar, M. (2002). Imaging cognitive modulation of pain processing. Pain, 95(1), 1-5. https://doi.org/10.1016/S0304-3959(01)00467-5

Petzschner, F. H., Weber, L. A., Wellstein, K. V., Paolini, G., Do, C. T., & Stephan, K. E. (2019). Focus of attention modulates the heartbeat evoked potential. NeuroImage, 186, 595-606.

Pezzulo, G. (2014). Why do you fear the bogeyman? An embodied predictive coding model of perceptual inference. Cognitive, Affective & Behavioral Neuroscience, 14(3), 902-911. https://doi.org/10.3758/s13415-013-0227-x

Phillips, G. C., Jones, G. E., Rieger, E. J., & Snell, J. B. (1999). Effects of the presentation of false heart-rate feedback on the performance of two common heartbeat-detection tasks. Psychophysiology, 36(4), 504-510.

Poli, A., Maremmani, A. G. I., Chiorri, C., Mazzoni, G.-P., Orrù, G., Kolacz, J., Porges, S. W., Conversano, C., Gemignani, A., & Miccoli, M. (2021). Item Reduction, Psychometric and Biometric Properties of the Italian Version of the Body Perception Questionnaire-Short Form (BPQ-SF): The BPQ-22. International Journal of Environmental Research and Public Health, 18(7), Articolo 7. https://doi.org/10.3390/ijerph18073835

Powers, A. R., Mathys, C., & Corlett, P. R. (2017). Pavlovian conditioning-induced hallucinations result from overweighting of perceptual priors. Science, 357(6351), 596-600.

Pratviel, Y., Bouni, A., Deschodt-Arsac, V., Larrue, F., & Arsac, L. M. (2022). Avatar embodiment in VR: Are there individual susceptibilities to visuo-tactile or cardio-visual stimulations?. Frontiers in Virtual Reality, 3, 954808.

Price, D. D., McHaffie, J. G., & Larson, M. A. (1989). Spatial summation of heat-induced pain: Influence of stimulus area and spatial separation of stimuli on perceived pain sensation intensity and unpleasantness. Journal of Neurophysiology, 62(6), 1270-1279. https://doi.org/10.1152/jn.1989.62.6.1270

Price, D. D., Milling, L. S., Kirsch, I., Duff, A., Montgomery, G. H., & Nicholls, S. S. (1999). An analysis of factors that contribute to the magnitude of placebo analgesia in an experimental paradigm. PAIN, 83(2), 147-156. https://doi.org/10.1016/S0304-3959(99)00081-0

Riello, M., Cecchini, M. P., Zanini, A., Di Chiappari, M., Tinazzi, M., & Fiorio, M. (2019). Perception of phasic pain is modulated by smell and taste. European Journal of Pain, 23(10), 1790-1800.

Roelofs, K. (2017). Freeze for action: Neurobiological mechanisms in animal and human freezing. Philosophical Transactions of the Royal Society B: Biological Sciences, 372(1718), 20160206.

Rupp, H. A., & Wallen, K. (2008). Sex differences in response to visual sexual stimuli: A review. *Archives of sexual behavior*, *37*(2), 206-218.

Schoeller, F., Horowitz, A., Maes, P., Jain, A., Reggente, N., Moore, L. C., Trousselard, M., Klein, A., Barca, L., & Pezzulo, G. (2022). Interoceptive technologies for clinical neuroscience.

Schönbrodt, F. D., & Perugini, M. (2013). At what sample size do correlations stabilize? Journal of Research in Personality, 47(5), 609-612.

Series, P. (2019). Post-traumatic stress disorder as a disorder of prediction. Nature Neuroscience, 22(3), 334-336.

Seth, A. K. (2013). Interoceptive inference, emotion, and the embodied self. Trends in Cognitive Sciences, 17(11), 565-573. https://doi.org/10.1016/j.tics.2013.09.007

Seth, A. K., & Friston, K. J. (2016). Active interoceptive inference and the emotional brain. Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences, 371(1708), 20160007. https://doi.org/10.1098/rstb.2016.0007

Seth, A., Suzuki, K., & Critchley, H. (2012). An Interoceptive Predictive Coding Model of Conscious Presence. Frontiers in Psychology, 2. https://www.frontiersin.org/articles/10.3389/fpsyg.2011.00395

Shahidi, S., & Baluch, B. (1991). False heart-rate feedback, social anxiety and self-attribution of embarrassment. *Psychological reports*, *69*(3), 1024-1026

Skora, L. I., Livermore, J. J. A., & Roelofs, K. (2022). The functional role of cardiac activity in perception and action. Neuroscience & Biobehavioral Reviews, 104655.

Smith, R., Kuplicki, R., Feinstein, J., Forthman, K. L., Stewart, J. L., Paulus, M. P., Investigators, T. 1000, & Khalsa, S. S. (2020). A Bayesian computational model reveals a failure to adapt interoceptive precision estimates across depression, anxiety, eating, and substance use disorders. PLOS Computational Biology, 16(12), e1008484.

Smith, R., Mayeli, A., Taylor, S., Al Zoubi, O., Naegele, J., & Khalsa, S. S. (2021). Gut inference: A computational modelling approach. Biological Psychology, 164, 108152.

Sokolov, E. N., Spinks, J. A., Näätänen, R., & Lyytinen, H. (2002). *The orienting response in information processing*. Lawrence Erlbaum Associates Publishers.

Song, Y., Kemprecos, H., Wang, J., & Chen, Z. (2019). A predictive coding model for evoked and spontaneous pain perception. 2019 41st Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC), 2964-2967.

Song, Y., Yao, M., Kemprecos, H., Byrne, A., Xiao, Z., Zhang, Q., Singh, A., Wang, J., & Chen, Z. S. (2021). Predictive coding models for pain perception. Journal of computational neuroscience, 49, 107-127.

Stephan, K. E., Manjaly, Z. M., Mathys, C. D., Weber, L. A. E., Paliwal, S., Gard, T., Tittgemeyer, M., Fleming, S. M., Haker, H., Seth, A. K., & Petzschner, F. H. (2016). Allostatic Self-efficacy: A Metacognitive Theory of Dyshomeostasis-Induced Fatigue and Depression. Frontiers in Human Neuroscience, 10, 550. https://doi.org/10.3389/fnhum.2016.00550

Sundararaj, S., Gazi, A. H., Vaccarino, V., Shah, A. J., Inan, O. T., & Bremner, J. D. (2025). Accrued reductions in heart rate following transcutaneous vagal nerve stimulation in adults with posttraumatic stress disorder. Frontiers in Neuroscience, 19, 1456662. https://doi.org/10.3389/fnins.2025.1456662

Taggart, P., Hedworth-Whitty, R., Carruthers, M., & Gordon, P. D. (1976). Observations on electrocardiogram and plasma catecholamines during dental procedures: The forgotten vagus. British Medical Journal, 2(6039), 787-789. https://doi.org/10.1136/bmj.2.6039.787

Tajadura-Jiménez, A., Väljamäe, A., & Västfjäll, D. (2008). Self-representation in mediated environments: the experience of emotions modulated by auditory-vibrotactile heartbeat. *CyberPsychology & Behavior*, *11*(1), 33-38.

Terkelsen, A. J., Mølgaard, H., Hansen, J., Andersen, O. K., & Jensen, T. S. (2005). Acute pain increases heart rate: differential mechanisms during rest and mental stress. Autonomic Neuroscience, 121(1-2), 101-109.

Tousignant-Laflamme, Y., Rainville, P., & Marchand, S. (2005). Establishing a link between heart rate and pain in healthy subjects: a gender effect. *The journal of pain*, *6*(6), 341-347.

Tousignant-Laflamme, Y., & Marchand, S. (2006). Sex differences in cardiac and autonomic response to clinical and experimental pain in LBP patients. European Journal of Pain (London, England), 10(7), 603-614. https://doi.org/10.1016/j.ejpain.2005.09.003

Tracey, I. (2010). Getting the pain you expect: Mechanisms of placebo, nocebo and reappraisal effects in humans. Nature Medicine, 16(11), 1277-1283. https://doi.org/10.1038/nm.2229

Tracy, L. M., Gibson, S. J., Georgiou-Karistianis, N., & Giummarra, M. J. (2017). Effects of explicit cueing and ambiguity on the anticipation and experience of a painful thermal stimulus. PloS One, 12(8), e0183650. https://doi.org/10.1371/journal.pone.0183650

Tsakiris, M., Tajadura-Jiménez, A., & Costantini, M. (2011). Just a heartbeat away from one's body: Interoceptive sensitivity predicts malleability of body-representations. Proceedings. Biological Sciences, 278(1717), 2470-2476. https://doi.org/10.1098/rspb.2010.2547

Tschantz, A., Barca, L., Maisto, D., Buckley, C. L., Seth, A. K., & Pezzulo, G. (2022). Simulating homeostatic, allostatic and goal-directed forms of interoceptive control using active inference. Biological Psychology, 169, 108266. https://doi.org/10.1016/j.biopsycho.2022.108266

Unal, O., Eren, O. C., Alkan, G., Petzschner, F. H., Yao, Y., & Stephan, K. E. (2021). Inference on homeostatic belief precision. Biological Psychology, 165, 108190.

Valins, S. (1966). Cognitive effects of false heart-rate feedback. Journal of personality and social psychology, 4(4), 400.

Van der Veen, F. M., Van Der Molen, M. W., & Jennings, J. R. (2000). Selective inhibition is indexed by heart rate slowing. Psychophysiology, 37(5), 607-613.

Vase, L., Nørskov, K. N., Petersen, G. L., & Price, D. D. (2011). Patients' direct experiences as central elements of placebo analgesia. Philosophical Transactions of the Royal Society B: Biological Sciences, 366(1572), 1913-1921.

Vase, L., Robinson, M. E., Verne, G. N., & Price, D. D. (2005). Increased placebo analgesia over time in irritable bowel syndrome (IBS) patients is associated with desire and expectation but not endogenous opioid mechanisms. Pain, 115(3), 338-347.

Vila, J., Guerra, P., Muñoz, M. Á., Vico, C., Viedma-del Jesús, M. I., Delgado, L. C., Perakakis, P., Kley, E., Mata, J. L., & Rodríguez, S. (2007). Cardiac defense: From attention to action. International Journal of Psychophysiology, 66(3), 169-182.

Wager, T. D. (2005). Expectations and anxiety as mediators of placebo effects in pain. In Pain (Vol. 115, Fascicolo 3, pp. 225-226). LWW.

Wager, T. D., Rilling, J. K., Smith, E. E., Sokolik, A., Casey, K. L., Davidson, R. J., Kosslyn, S. M., Rose, R. M., & Cohen, J. D. (2004). Placebo-induced changes in FMRI in the anticipation and experience of pain. Science, 303(5661), 1162-1167.

Wascher, C. A. F. (2021). Heart rate as a measure of emotional arousal in evolutionary biology. Philosophical Transactions of the Royal Society B: Biological Sciences, 376(1831), 20200479. https://doi.org/10.1098/rstb.2020.0479

Wiech, K., Ploner, M., & Tracey, I. (2008). Neurocognitive aspects of pain perception. Trends in cognitive sciences, 12(8), 306-313.

Wiech, K., Vandekerckhove, J., Zaman, J., Tuerlinckx, F., Vlaeyen, J. W., & Tracey, I. (2014). Influence of prior information on pain involves biased perceptual decision-making. Current Biology, 24(15), R679-R681.

Williams, A. C. de C., & Craig, K. D. (2016). Updating the definition of pain. Pain, 157(11), 2420-2423.

Yang, L. L., Niemann, C. U., & Larson, M. D. (2003). Mechanism of pupillary reflex dilation in awake volunteers and in organ donors. Anesthesiology, 99(6), 1281-1286. https://doi.org/10.1097/00000542-200312000-00008

Yang, Z., Jia, W., Liu, G., & Sun, M. (2017). Quantifying mental arousal levels in daily living using additional heart rate. Biomedical Signal Processing and Control, 33, 368-378.

Yon, D., & Frith, C. D. (2021). Precision and the Bayesian brain. Current Biology, 31(17), R1026-R1032.

Valins, S. (1966). Cognitive effects of false heart-rate feedback. Journal of personality and social
psychology, 4(4), 400.

van Elk, M., Lenggenhager, B., Heydrich, L., & Blanke, O. (2014). Suppression of the auditory N1-
component for heartbeat-related sounds reflects interoceptive predictive coding. Biological
psychology, 99, 172-182.

van Langen, J. (2020). Open-visualizations in R and Python. Jordy Van Lagen.

Vicentin, S., Guglielmi, S., Stramucci, G., Bisiacchi, P., & Cainelli, E. (2024). Listen to the beat:
behavioral and neurophysiological correlates of slow and fast heartbeat sounds. International
Journal of Psychophysiology, 206, 112447.
1335
1336
1337