Title: A CD24+CD271+ melanoma cancer stem cell generates a diffuse hierarchy of attributes that promote metastasis and therapeutic resistance

Authors: Olivia Knowles[1]*, Patricio Doldan[1]*, Isabella Hillier-Richardson[1]*, Stephanie Lunt[1], Gehad Youssef[1], Luke Gammon[1], Ian C. Mackenzie[1], Michael P. Philpott[1], Hasan Rizvi[2], Daniele Bergamaschi[1], Catherine A. Harwood[1], Adrian Biddle[1]

Affiliation and address of authors: [1]Blizard Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, UK. [2]Department of Cellular Pathology, Bart's Health NHS Trust, London, UK.

*These authors contributed equally.

Corresponding author: Adrian Biddle, Centre for Cell Biology and Cutaneous Research, Blizard Institute, 4 Newark Street, London, E1 2AT, UK. E-mail: a.biddle@qmul.ac.uk

Abstract

An important role for phenotype switching has been demonstrated in metastasis and therapeutic resistance of both melanoma and epithelial tumours. Phenotype switching in epithelial tumours is driven by a minority cancer stem cell sub-population with lineage plasticity, but such a sub-population has not been identified in melanoma. We investigated whether cell surface markers used to identify cancer stem cells in epithelial tumours could help to identify a cancer stem cell sub-population with lineage plasticity in melanoma. We identified a CD24+CD271+ minority sub-population in melanoma that possesses enhanced stem cell characteristics and lineage plasticity. We further found that that, unlike in epithelial

tumours, more differentiated sub-populations in melanoma also possessed these attributes to a lesser extent. The CD24+CD271+ stem cell sub-population was observed in only 10% of human melanomas, mainly at the invasive front. The lack of this stem cell sub-population in the majority of human melanoma specimens led us to conclude that it may not be required for melanoma progression. This may be due to the observed diffuse nature of stem cell characteristics in melanoma. However, the enhanced self-renewal, lineage plasticity, invasive ability and drug resistance of the CD24+CD271+ sub-population may signal a contextual requirement for these stem cells when melanomas face challenging environments both in clinical melanoma and in experimental systems.

Introduction

Melanoma is a highly aggressive form of skin cancer. Invasive melanoma accounts for about 1% of all skin cancer cases, but is responsible for around 60% of skin cancer deaths (American Cancer Society.). Melanoma is unusual in that it is often very invasive at an early stage of tumour development, setting it apart from other solid tumours that typically become invasive at a later stage. This single attribute is a major contributor to melanoma's high mortality rate, despite recent therapeutic advances. Even when a patient presents with a small primary lesion, the tumour has often already invaded beyond the primary site and is thus incurable with surgery. Melanoma develops from melanocytes, the pigment forming cells of the skin, and these are distinct from the other cells of the skin in that they are not derived from an epithelial developmental lineage. Instead, they derive from the neural crest, a lineage that undergoes an epithelial-mesenchymal-transition (EMT) and migrates throughout the embryo to produce a number of different cell types in development (Morrison et al., 1999).

EMT is a major driver of tumour invasion in epithelial cancers (Biddle et al., 2011, Lu and Kang, 2019), with activation of an EMT program during tumour development greatly enhancing

metastatic spread (Tsai et al., 2012, Pastushenko et al., 2021). Activation of an EMT program is restricted to cancer stem cells (CSCs), a minority sub-population of cells within the tumour that possess tumour-initiating potential by virtue of their self-renewal and lineage plasticity. Previous work in epithelial cancers has shown that tumour invasion and metastasis is driven by phenotypic switching between two CSC states: 1) a stationary, highly proliferative, drug-sensitive epithelial state that is $CD44^{high}$, $EpCAM^{high}$ and $Vimentin^{low}$, and 2) an invasive, slow-growing, drug-tolerant mesenchymal-like state that is $CD44^{high}$, $EpCAM^{low}$ and $Vimentin^{high}$ (Biddle et al., 2016, Biddle et al., 2011, Liu et al., 2014, Hermann et al., 2007). There is further heterogeneity within the sub-population of cells that have transitioned through EMT into a mesenchymal-like state, with only some of these cells retaining the CSC attribute of lineage plasticity that enables them to regenerate a proliferative epithelial tumour at a metastatic site and thus propagate metastatic dissemination (Youssef et al., 2020, Pastushenko et al., 2018). In our own work on oral squamous cell carcinoma (OSCC), we have shown that these metastatic EMT-CSCs can be identified by their expression of the plasticity marker CD24 alongside the epithelial marker EpCAM, the mesenchymal marker Vimentin, and the CSC marker CD44 (Youssef et al., 2020).

As melanomas derive from a lineage that has already undergone a developmental EMT, they do not need to undergo the phenotypic switching described above. It might be expected that this is a key underlying reason for their propensity for early tumour invasion. Indeed, melanoma cells do not transition between epithelial and mesenchymal states, and ubiquitously express high levels of the mesenchymal marker Vimentin (Rambow et al., 2019). However, numerous studies have demonstrated that melanomas nevertheless exhibit lineage plasticity and transition between two states that are analogous to those seen in epithelial tumours: 1) a stationary, highly proliferative, drug-sensitive melanocytic state that is $MITF^{high}$, Axl^{low}, $Zeb1^{low}$ and $CD271^{low}$, and 2) an invasive, slow-growing, drug-tolerant neural crest-like state that is $MITF^{low}$, Axl^{high}, $Zeb1^{high}$ and $CD271^{high}$ (Carreira et al., 2006, Richard et al., 2016, Müller et

al., 2014, Restivo et al., 2017, Sensi et al., 2011). This demonstrates the same important role for phenotypic switching as has been seen in epithelial tumours.

The existence of CSCs in melanoma is controversial. In a study of tumour initiation, 25% of unfractionated melanoma cells possessed tumour-initiating potential (Quintana et al., 2008). Further, the same team failed to find any evidence for a CSC hierarchy when assessing tumour-initiating potential using previously-described melanoma CSC markers (Quintana et al., 2010). This raises the question of whether there is any other basis for a phenotypic hierarchy in melanoma: is lineage plasticity restricted to a minority sub-population, or is it common to all melanoma cells?

We hypothesised that the minority CSC sub-populations we have identified in epithelial tumours may have parallel states in melanoma, whose retained lineage plasticity is responsible for driving phenotypic switching and thus propagating metastatic dissemination. In order to investigate this, we incorporated a previously-reported melanoma CSC marker, CD271 (Boiko et al., 2010), alongside markers we previously identified in epithelial carcinomas. As well as being a reported melanoma CSC marker, CD271 is a key cell surface marker for the invasive slow-growing state in melanoma (Restivo et al., 2017). It is also a marker for resistance to BRAF inhibitors (Richard et al., 2016) and resistance to immunotherapy (Landsberg et al., 2012). We therefore selected CD271 for investigation alongside the cell surface markers for metastatic CSCs in epithelial tumours - CD24, EpCAM and CD44 (Youssef et al., 2020). We sought to determine whether any of these markers in combination could discriminate a metastatic CSC sub-population in melanoma.

Results

We initially cultured 4 established melanoma cell lines – LM4, LM32, CHL-1 and A375M. We assessed each cell line for cell surface expression of CD44, EpCAM, CD24 and CD271 by flow cytometry (Figure 1A). All cell lines were homogeneously CD44$^+$EpCAM$^-$, except for CHL-1 which contained a small number of CD44$^-$EpCAM$^-$ cells. This demonstrates the mesenchymal character of all cell lines, in agreement with the aforementioned ubiquitously high expression of the mesenchymal marker Vimentin in melanoma (Rambow et al., 2019). It also suggests that CD44, a CSC marker in epithelial cancers, does not discriminate sub-populations in melanoma. The surface expression of CD24 and CD271 was more variable between cell lines; the 4 lines expressed varying levels of CD271, whereas only CHL-1 expressed CD24. This gave rise to both CD24$^+$CD271$^+$ and CD24$^+$CD271$^-$ sub-populations in the CHL-1 cell line only. Brightfield imaging (Figure 1B) demonstrated a mesenchymal morphology for all cell lines, further underlining their mesenchymal character. However, there was heterogeneity between cell lines, with LM4 and LM32 having a homogenously mesenchymal morphology whereas CHL-1 and A375M were more heterogeneous in their morphology, with clustering cobblestone regions. These were more pronounced in CHL-1, which overall showed a greater level of morphological heterogeneity than the other three lines. In order to assess the CSC attributes of the four lines, we performed a tumoursphere assay. The tumoursphere assay tests ability to survive and grow in suspension, and is a surrogate assay for CSC identity (Mukherjee et al., 2021); it enriches for cells that exhibit resistance to the oxidative stress and anoikis triggered by suspension culture (Zhu et al., 2020). We performed the tumoursphere assay under two conditions; in fully supplemented FAD medium with 10% serum, as used in adherent culture (Figure 1C, top), or in a defined serum-free medium (Figure 1C, bottom) (Sato et al., 2011). Interestingly, behaviour was starkly different under the two conditions; all four cell lines produced both primary and secondary spheres in

the FAD medium, whereas only CHL-1 produced spheres in the more stringent serum-free conditions.

Thus, of the four melanoma cell lines, only CHL-1 exhibits heterogeneous sub-populations based on any of the metastatic CSC markers derived from epithelial tumours. Both CD44 and CD24 were heterogeneously expressed in CHL-1, but only CD24 identified a minority positive sub-population that was absent from the other cell lines. CHL-1 was also the only cell line capable of producing tumourspheres under the more stringent serum-free conditions. We therefore focussed on CD24 as a potential marker of a sphere-forming CSC sub-population in CHL-1.

Figure 1 – Cellular heterogeneity and stem cell attributes differ between melanoma cell lines. **A**, Flow cytometric analysis of the four cell lines for CD44 and EpCAM (left) and CD24 and CD271 (right). The graphs show mean +/- SEM of sub-populations in each of 4 quadrants based on +/- gating set using isotype controls, as a percentage of the total cells, presented as stacked bars. Representative flow cytometry plots for each cell line are to the left of each graph. **B**, Brightfield images of four melanoma cell lines in culture. **C**, Sphere counts for the four cell lines, for primary spheres and secondary spheres after dissociation and re-plating, in serum-containing FAD medium (top) and serum-free medium (bottom). **D**, Gene expression in FACS sorted CD24+ vs CD24- sub-populations. For each gene, expression was measured by QPCR and is represented as the expression level in CD24+ cells relative to that in CD24- cells. For each experiment, the number of biological repeats is indicated next to the graph. P-values were calculated using two-way ANOVA.

We performed QPCR analysis on FACS sorted CD24+ and CD24- sub-populations from CHL-1, for markers of the two recognised melanoma phenotypic states – melanocytic (proliferative) and neural crest-like (invasive) (Figure 1D). Compared to CD24- cells, CD24+ cells had heightened expression of markers of the neural crest-like state (Zeb1 and Axl) and reduced expression of the melanocytic state marker MITF. The mesenchymal marker Vimentin was stably expressed across the two sub-populations. Interestingly, and contrary to the documented growth-arrested status of the neural crest-like state, the CD24+ sub-population had greatly increased Cyclin D expression. This suggested that CD24 may mark an alternative phenotypic state that has a dedifferentiated neural crest-like status whilst maintaining proliferative competence, such as has been recently proposed (Rambow et al., 2019). Therefore, we focussed on CD24 alongside CD271 for further investigations in the CHL-1 cell line in order to determine whether this cell line contains a metastatic CSC sub-population that is absent from the other melanoma cell lines.

We FACS sorted four sub-populations (CD24$^+$CD271$^+$, CD24$^+$CD271$^-$, CD271$^+$CD24$^-$ and CD24$^-$CD271$^-$) from the CHL-1 sub-line, and performed a range of functional CSC assays. Using the double-negative CD24$^-$CD271$^-$ sub-population as a baseline, we found that only the CD24$^+$CD271$^+$ sub-population produced a significantly increased number of tumourspheres (utilising the more stringent serum-free condition) (Figure 2A) and adherent colonies (Figure 2B). Therefore, the CD24$^+$CD271$^+$ sub-population exhibits enhanced anoikis resistance and self-renewal. To assess lineage plasticity, we cultured each sorted sub-population for 7 days and then re-assessed the sub-population distribution (Figure 2C). Plasticity was much more pronounced in the CD24$^+$CD271$^+$ sub-population, which gave rise to large numbers of cells in each of the other three sub-populations. In contrast, the CD24$^+$CD271$^-$ and CD24$^-$CD271$^+$ sub-populations gave rise mostly to CD24$^-$CD271$^-$ cells, and the CD24$^-$CD271$^-$ sub-population produced few cells in any of the other sub-populations. Therefore, the CSC attributes of self-renewal and lineage plasticity reside mainly in the CD24$^+$CD271$^+$ sub-population. However,

unlike in epithelial tumours (Biddle et al., 2011, Youssef et al., 2020), there is still a significant baseline retention of these CSC attributes in the other sub-populations in melanoma, with even the double-negative CD24$^-$CD271$^-$ sub-population possessing some degree of self-renewal and lineage plasticity.

Figure 2 – CD24 and CD271 mark a stem cell sub-population in the CHL-1 melanoma line. **A**, Primary sphere counts for four FACS sorted sub-populations from the 4 quadrants based on +/- gating set using isotype controls. Data is presented as paired individual data points (due to considerable baseline variation between repeats), with each of the three marker-positive sub-populations compared individually to the CD24-CD271-sub-population. P-values were calculated using paired ANOVA (due to the variation between repeats). **B**, Colony counts in 2D culture for the four FACS sorted sub-populations, with representative images of crystal violet stained colonies for each sub-population. P-values were calculated using unpaired ANOVA. **C**, Flow cytometric analysis for CD24 and CD271, 7 days after sorting and re-plating the four sub-populations, with accompanying brightfield images of the four sorted sub-populations in culture. The graph shows mean +/- SEM of sub-populations in each of 4 quadrants based on +/- gating set using isotype controls, as a percentage of the total cells, presented as stacked bars. P-values were calculated using two-way ANOVA. Representative flow cytometry plots for each re-plated sub-population are to the left of the graph. For each experiment, the number of biological repeats is indicated next to the graph.

We next assessed the ability of the four CHL-1 sub-populations to drive metastatic dissemination. The CD24$^+$CD271$^+$ sub-population exhibited enhanced migration in a transwell migration assay (Figure 3A). For a more stringent assessment of invasive ability, we developed a 3D matrigel-collagen invasion assay through adaptation of a previously published method (Katz et al., 2011). In this assay, the melanoma cells are embedded in a matrigel 'plug'

that is then itself embedded within a larger collagen outer gel. In this way, the degree of invasion of melanoma cells from the matrigel plug into the surrounding collagen can be assessed. The matrigel is rich in basement membrane components, and thus provides an environment somewhat similar to the *in situ* tumour environment for a stringent assessment of invasive ability. Flattening the resulting 3D invasion into a 2D maximum intensity projection, and plotting the average distance of each invading cell from the centre of the matrigel plug, enabled assessment of the invasive ability of each sorted sub-population (Figure 3B). Both the CD24$^+$CD271$^+$ and CD24$^+$CD271$^-$ sub-populations had enhanced invasive ability compared to the CD24$^-$CD271$^-$ sub-population. However, again unlike epithelial tumours (Biddle et al., 2011, Hermann et al., 2007), all four sub-populations possessed some invasive potential.

Figure 3 – The CD24+CD271+ sub-population in the CHL-1 melanoma line has enhanced ability to migrate and invade. **A**, The number of cells that have migrated to the other side of the membrane in transwell assays, for the four FACS sorted sub-populations. **B**, Invasion of the four FACS sorted sub-populations into a collagen gel, when seeded in an inner matrigel gel. Left, Representative images, with each blue diamond representing a cell, projected onto x-y co-ordinates (z co-ordinates are flattened to zero). Units are pixel numbers in the x-y plane, and the orange star indicates the centre of the inner matrigel gel from which distance was calculated for each invading cell. Cells remaining in the inner gel are excluded from the visualisation. Top right, Average invasion distance for the four FACS sorted sub-populations. Bottom right, A representative brightfield image of cells breaking away from the inner gel and invading into the surrounding collagen. For each experiment, the number of biological repeats is indicated next to the graph. P-values were calculated using unpaired ANOVA.

Melanoma exhibits broad chemotherapeutic resistance, and also quickly develops resistance to targeted therapies (Kozar et al., 2019). CSCs have been reported to exhibit enhanced therapeutic resistance in many tumour types, including melanoma, and drive the development of tumour drug resistance (Biddle et al., 2016, Li et al., 2008, Phi et al., 2018). In particular, CD24+ CSCs in epithelial tumours exhibit the highest resistance to chemotherapy drugs (Biddle et al., 2016, Goldman et al., 2015), although the Src tyrosine kinase inhibitor Dasatinib has been reported to target CD24+ CSCs in epithelial tumours (Goldman et al., 2015). Therefore, to complete the functional characterisation of the four CHL-1 sub-populations, we assessed their relative responses to therapeutic challenge with the cytotoxic chemotherapeutic drugs Paclitaxel and Cisplatin and the Src tyrosine kinase inhibitor Dasatinib (Figure 4A). The $CD24^+CD271^+$ and $CD24^+CD271^-$ sub-populations exhibited heightened resistance to Dasatinib, becoming enriched after treatment (contrary to previous findings in epithelial tumours). Conversely, heightened resistance to Paclitaxel and Cisplatin was exhibited by the $CD24^+CD271^+$ and $CD24^-CD271^+$ sub-populations. Therefore, the $CD24^+CD271^-$ and $CD24^-CD271^+$ sub-populations exhibit differing resistance profiles, but these converge on the $CD24^+CD271^+$ sub-population which exhibits resistance to all three drugs.

Acquired resistance to targeted therapies is a serious problem in melanoma (Rambow et al., 2019), and we therefore used long-term Dasatinib treatment to create Dasatinib resistant CHL-1 (Figure 4B, top). This caused a marked shift in the sub-population distribution, with a big increase in the $CD24^+CD271^-$ sub-population and corresponding loss of the $CD24^-CD271^+$ and $CD24^-CD271^-$ sub-populations. This underlines the role of CD24 in marking Dasatinib resistant cells in this cell line. However, we were also able to create Dasatinib resistant A375M using the same method, despite the absence of CD24+ sub-populations in this line. In A375M, the $CD24^-CD271^+$ sub-population was lost, resulting in an entirely $CD24^-CD271^-$ cell line (Figure 4B, bottom). Therefore, $CD271^+CD24^-$ cells are particularly sensitive to Dasatinib, just

as CD24+ cells are particularly resistant. The fact that we were able to create a resistant cell line from A375M, which lacks CD24+ sub-populations, indicates that resistance to targeted therapy can also occur within the CD24⁻CD271⁻ sub-population.

Figure 4 – The CD24+CD271+ sub-population is drug resistant. **A**, Surviving cells in the CHL-1 melanoma line, after 3 days treatment with the indicated drugs and a 2 day recovery. **B**, Changes in distribution of sub-populations after generation of Dasatinib resistant CHL-1 (top) and A375M (bottom) melanoma lines. **C**, Changes in distribution of sub-populations after TGFβ treatment for 6 days on CHL-1 (left) and A375M (right) melanoma lines. For all experiments, flow cytometric analysis for CD24 and CD271 is shown. The graphs show mean +/- SEM of sub-populations in each of 4 quadrants based on +/- gating set using isotype controls, as a percentage of the total cells, presented as stacked bars. Note that Dasatinib resistant A375M exhibited increased isotype control staining, necessitating altered +/- gating compared to untreated cells. P-values were calculated using two-way ANOVA. Representative flow cytometry plots for each condition are to the left of each graph. For each experiment, the number of biological repeats is indicated next to the graph.

We next considered the potential role for an amoeboid phenotype in the enhanced invasive ability of the CD24+ sub-populations. Amoeboid cells are known to cluster at the tumour invasive front in melanoma and drive metastatic dissemination (Georgouli et al., 2019), and this phenotype is induced by TGFβ treatment (Cantelli et al., 2015). We therefore treated the CHL-1 and A375M cell lines with TGFβ and observed the effect on the CD24 and CD271

marked sub-populations (Figure 4C). In both lines, TGFβ caused an expansion of the CD24$^-$CD271$^+$ sub-population. In CHL-1, the CD24$^+$CD271$^-$ sub-population was reduced and there was no change in the CD24$^+$CD271$^+$ sub-population. Therefore, TGFβ induces the CD24$^-$CD271$^+$ sub-population, indicating that the invasive amoeboid phenotype is not a specific attribute of the CD24$^+$CD271$^+$ CSC sub-population.

Finally, we investigated whether a CD24$^+$CD271$^+$ CSC sub-population exists in human primary melanoma specimens. FFPE sections of 31 melanomas were examined by a pathologist, and included 19 superficial spreading, 11 nodular, and 1 acral lentiginous melanomas (Table 1). We developed a method for immunofluorescent co-staining for CD24 and CD271, combined with automated imaging and image stitching to create high resolution images of entire cut sections. This enabled accurate assessment of CD24 and CD271 co-staining in each of the specimens, which was performed by both a researcher and a pathologist alongside the H&Es to ensure that analysis of co-staining was restricted to melanoma cells (Figure 5). Of the 31 tumours, 15 (48%) had evidence of CD24$^-$CD271$^+$ melanoma cells. These were specific to the invasive front in 5 tumours. Only 3 tumours (10%) had evidence of CD24$^+$CD271$^+$ melanoma cells and, interestingly, CD24$^+$CD271$^-$ melanoma cells were also restricted to these same 3 tumours. In 2 of these tumours, the staining was specific to the invasive front. These findings confirm the existence of a CD24$^+$CD271$^+$ sub-population that localises to the invasive front in human melanoma but, in agreement with our panel of melanoma cell lines, is only present in a minority of tumours.

Tumour	CD24+/CD271+ cells	CD24+ cells	CD271+ cells	Notes	Breslow Thickness (mm)
1				Superficial spreading melanoma	1
2				Superficial spreading melanoma	5
3				Superficial spreading with vertical growth/ possibly nodular	2.7
4	Present	Present	Present	Nodular melanoma. CD271 and CD24 are present in melanoma capsules	7.5
5			Present	Nodular melanoma, more positive in deeper areas	5.5
6				Superficial spreading melanoma in vertical growth phase. Spindle cell.	4
7				Superficial spreading melanoma, with microinvasion. In situ. Intradermal cells.	0.7
8				Superficial spreading melanoma, in situ	0.5
9				Nodular melanoma	13.5
10			Present	Nodular melanoma, CD271 present in deep parts of nodule (invading)	7.2
11				Superficial spreading melanoma, in situ	0.5
12				Superficial spreading melanoma, with invasion	0.5
13				Nodular melanoma	2.76
14				Superficial spreading melanoma	3.5
15			Present	Superficial spreading melanoma in vertical growth phase, some nests with CD271	2.6
16			Present	Superficial spreading melanoma with clear CD271 staining	1.2
17			Present	Superficial Spreading, CD271 very positive in 2 large nests of melanoma	7
18	Present	Present	Present	Superficial spreading in situ. Co-staining superficially in melanoma - at invasive edge	0.5
19			Present	Acral lentiginous melanoma from foot, CD271 staining is in nests.	1.1
20				Superficial spreading melanoma	1
21			Present	Superficial spreading Melanoma in situ	0.22
22				Nodular melanoma	8
23			Present	Superficial Spreading Melanoma in situ	4.5
24			Present	Nodular melanoma	6
25			Present	Nodular melanoma. CD271 present at deep invading edge of melanoma	3.9
26				Nodular melanoma	5
27				Superficial spreading melanoma	0.8
28			Present	Nodular melanoma, CD271 positive	4.1
29	Present	Present	Present	Superficial spreading melanoma, in situ. Co-staining is at invasive edge.	0.9
30				Nodular melanoma.	1.7
31			Present	Superficial spreading melanoma	2.5

Table 1 – Pathological assessment of CD24 and CD271 staining in 31 melanoma specimens.

Figure 5 – A CD24+CD271- sub-population in human melanoma tumour specimens. **A**, H&E of a human melanoma specimen, showing regions of melanoma cells (black arrows). **B**, Isotype control staining. **C**, Melanoma specimen with a region of CD24+CD271+ staining. Blue – DAPI nuclear stain, Green – CD24, Red – CD271. **D**, Magnification of the region of CD24+CD271+ staining. Displayed images are image-stitches of multiple fields of view at 40x magnification.

Discussion

Previous investigations of cellular heterogeneity in melanoma have led to a developing understanding of the distribution of functional sub-populations and how switching between phenotypes underlies metastasis and therapeutic resistance (Rambow et al., 2019, Müller et al., 2014). However, there are still many unanswered questions about how these different sub-populations contribute to metastasis and therapeutic resistance. In particular, it is not known which (if any) of the identified sub-populations correspond to a melanoma CSC sub-population, or even if CSC sub-populations exist in melanoma (Rambow et al., 2019, Quintana et al., 2010). We, and others, have previously identified a metastatic CSC sub-population that drives phenotype switching in epithelial tumours and through this propagates metastatic dissemination and therapeutic resistance (Biddle et al., 2016, Youssef et al., 2020, Pastushenko et al., 2021). In this report, we sought to test the hypothesis that a similar minority sub-population exists in melanoma, whose retained lineage plasticity is responsible for driving phenotypic switching and thus propagating metastatic dissemination. Using the epithelial tumour CSC marker CD24 alongside the melanoma CSC marker CD271, we have identified a minority CSC sub-population that exhibits enhanced self-renewal and lineage plasticity. It is also more invasive and exhibits heightened drug resistance. However, this CSC sub-population only occurred in 1 out of 4 established melanoma cell lines and 3 out of 31 melanoma tumour specimens. It is also notable that, whilst exhibiting heightened CSC attributes, this sub-population was not absolutely required for any of the functional behaviours tested; tumoursphere formation, adherent colony formation, migration and invasion, and drug resistance.

Interestingly, in epithelial tumours, tumoursphere formation in serum-containing FAD medium is restricted to sub-populations that have undergone EMT (Biddle et al., 2013, Biddle et al., 2011). The mesenchymal character of the melanoma cell lines maybe explains their ubiquitous sphere-forming abilities under these conditions. Conversely, under more stringent serum-free

conditions sphere-forming ability was restricted to the CHL-1 cell line that contains the CD24+CD271+ sub-population, and was heightened in this sub-population. Given that sphere-forming cells must resist oxidative stress, this draws parallels with work demonstrating that metastasising melanoma cells must resist oxidative stress in circulation and undergo reversible metabolic changes to achieve this (Piskounova et al., 2015). Therefore, the CD24+CD271+ cells may represent a minority CSC sub-population that is primed for efficient metastatic dissemination.

Previous findings have shown that CD24 expression in primary melanoma is one of the most predictive markers for metastatic disease (Gschaider et al., 2012). Similarly, Tang et al. (2014) demonstrated that CD24 was increased in melanoma patient samples compared to normal tissue, and that this marker was prognostic, with elevated CD24 correlating with decreased survival (Tang et al., 2014). A pro-metastatic CD24+ sub-population has also been characterised in a mouse model of melanoma (Liu et al., 2012).

In conclusion, in this study we have identified a CSC hierarchy in melanoma but, unlike in epithelial tumours, this may be dispensable for tumour progression, invasion and therapy resistance. This is highlighted by the fact that the CD24+CD271+ CSC sub-population is absent from a majority of melanoma tumour specimens and melanoma cell lines. Thus we propose that melanomas possess a phenotypic hierarchy, with full lineage plasticity restricted to a sub-population of CD24+CD271+ CSCs. However, unlike in epithelial tumours, functional CSC attributes are not exclusive to this CSC sub-population but rather exhibit a diffuse pattern; they are most concentrated on the CSC sub-population but also displayed to a lesser degree by more differentiated sub-populations. This diffuse nature of CSC attributes in melanoma may mean that CSCs are dispensable for melanomas. Functional outcomes of a CSC hierarchy may only manifest under particular conditions, for example the increased oxidative

stress associated with metastatic dissemination (Piskounova et al., 2015). However, even here there may be other ways for melanoma to subvert the requirement for a CSC sub-population, for example by metastasising via the lymphatics where they experience less oxidative stress (Ubellacker et al., 2020). The diffuse nature of CSC attributes in melanoma may also explain conflicting findings in the literature on whether melanomas are hierarchically organised on a CD271+ CSC (Boiko et al., 2010, Quintana et al., 2010). Our findings suggest that the manifestation of a CSC hierarchy may be dependent on local conditions and, thus, differences in experimental conditions between studies may have an important impact on whether a functional CSC hierarchy is identified.

Methods

Cell culture

Four human melanoma lines were used in this study: CHL-1 (provided by Penny Lovat, Newcastle), A375M (provided by John Marshall, Barts & the London), LM4 and LM32 (provided by Monica Rodolfo, Inst. Tumori Milano). LM4 was from a human primary tumour of the trunk region (Daniotti et al., 2004). CHL-1 (RRID: CVCL_1122) and LM32 (Daniotti et al., 2004) were from human metastatic sites (pleural effusion and cutaneous metastasis, respectively). A375M was formed from pooled lung metastatic deposits of a human melanoma cell line in a mouse metastatic model (Kozlowski et al., 1984). Cell culture was performed as previously described (Biddle et al., 2011). Cell removal from adherent culture was performed using 1x Trypsin-EDTA (Sigma, T3924) at 37ºC.

Flow cytometry and fluorescence-activated cell sorting (FACS)

Flow cytometry was performed as previously described (Biddle et al., 2011). Antibodies for cell line staining were CD44-PE (clone G44-26, BD Bioscience), CD24-FITC (clone ML5, BD Bioscience), CD271-PerCP/Cy5.5 and CD271-APC (both clone C40-1457, BD Bioscience), and EpCAM-APC (clone HEA-125, Miltenyi Biotec). Single stained controls were performed

for compensation, and isotype controls were performed to set negative gating. Flow cytometry data was analysed using FlowJo software, and statistical testing performed using 2-way ANOVA with multiple comparisons in GraphPad Prism software.

Sphere assay

For suspension cultures, 48-well plate wells were coated with polyhema (Sigma) (12mg/ml in 95% ethanol). Cells were then plated at a density of 1000 cells/ml, incubated (37C, 5% CO_2), and spheres were counted after 7 days. 1x Trypsin-EDTA was used to break down spheres for secondary sphere plating. The medium for suspension culture was either fully supplemented FAD medium (Biddle et al., 2011) or defined serum free medium (Sato et al., 2011), as indicated, with addition of 1% methylcellulose (Sigma). Statistical testing was performed using paired ANOVA with multiple comparisons.

QPCR

RNA extraction, cDNA synthesis and QPCR were performed as previously described (Biddle et al., 2011). Primer sequences were; Zeb1 F: GTCCAAGAACCACCCTTGAA R: TTTTTGGGCGGTGTAGAATC; Vimentin F: CCCTCACCTGTGAAGTGGAT R: GACGAGCCATTTCCTCCTTC; Axl F: CAAGTGGATTGCCATTGAGA R: GATATGGGGTTTGGCCTCTT; Cyclin D1 F: CTTCCTCTCCAAAATGCCAG R: TGAGGCGGTAGTAGGACAGG; MITF F: GGGCTTGATGGATCCTGCTT R: TATGTTGGGAAGGTTGGCTG; GAPDH F: GTGAACCATGAGAAGTATGACAAC R: CATGAGTCCTTCCACGATACC.

Colony assay

Cells were plated at a concentration of 1,000 cells per ml and cultured for seven days. They were then fixed in 4% paraformaldehyde and stained with crystal violet for colony counting. Statistical testing was performed using unpaired ANOVA with multiple comparisons.

Migration assay

1000 cells were plated in Transwell tissue culture inserts (8 mm membrane, Corning) in 24-well plates. After 24 hours, the membranes were fixed in 4% paraformaldehyde, stained with crystal violet, the non-migrated cells on the top of the membrane removed with a cotton-wool bud, and the migrated cells on the underside of the membrane counted. Statistical testing was performed using unpaired ANOVA with multiple comparisons.

Collagen invasion assay

10,000 cells were suspended in 50μl 4mg/ml collagen (BD Bioscience) in 96-well plate wells, which was allowed to set before being removed and inserted into the middle of 500μl 4mg/ml collagen in 48-well plate wells coated with polyhema. Once this had set, medium was added and the cells were cultured for 7 days. For the final day of culture, 20 µg/ml Hoescht 33342 (Sigma) was added to the medium. The cultures were then fixed in 4% paraformaldehyde, and imaged through z-stacks using an InCell 2200 automated epifluorescent microscope (GE) to detect all stained nuclei. For each stained nucleus, distance from the centre of the small inner gel in the x-y plane was calculated and reconstructed in a graphical plot. Average distance was calculated for each sorted sub-population, and statistical testing was performed using unpaired ANOVA with multiple comparisons.

Drug treatment, generation of Dasatinib resistant lines, and TGFβ treatment

Cells were plated at a density of 4000 cells per ml in T25 flasks. For CHL-1 drug treatment, drugs were added to the following final concentrations: Cisplatin 1.5 µM (Sigma), Paclitaxel 10 µM (Sigma), Dasatinib 0.5 µM (BioVision). Medium was replaced after 3 days to remove the drugs and cells were analysed by flow cytometry after a further 2 day recovery period. For the generation of Dasatinib resistant CHL-1 and A375M, cells were grown in 1 µM Dasatinib continuously with the medium and drug changed every 3 days until robust growth was re-established. Cells were then analysed by flow cytometry. For TGF-β treatment, 2 ng/ml TGF-β (R&D Systems) was added. TGF-β was replaced once during a six days incubation period, followed by flow cytometric analysis.

Immunofluorescent staining of tumour tissue sections

FFPE blocks from Barts pathology archive were obtained under UK HRA consent 07/Q0604/23. H&Es were examined by a pathologist to confirm presence of melanoma. 5 μm sections on poly-lysine slides were then de-waxed in xylene and ethanol and hot antigen retrieval was performed in Tris-EDTA pH9 for 10 minutes. Permeabilisation was performed for 10 minutes in 0.25% Triton X-100 in PBS. Washes were in PBS, and blocking was overnight at 4 degrees C with 3% BSA in PBS. Primary antibody incubation was overnight at 4 degrees C with the following antibodies at 1:100 dilution in PBS + 3% BSA: CD24-FITC (clone ML5, BD Bioscience), CD271-APC (clone C40-1457, BD Bioscience). Isotype controls were included. After washing in PBS + 3% BSA, sections were incubated for 1 hr at room temp in the following secondary antibodies at 1:500 dilution in PBS + 3% BSA: Alexa Fluor 488 Goat Anti-mouse IgG2a (Invitrogen) (binds the CD24 primary antibody), Alexa Fluor 647 Goat Anti-mouse IgG1 (Invitrogen) (binds the CD271 primary antibody). Antibody combinations were tested for cross reactivity, and none was seen. After washing again in PBS + 3% BSA, sections were incubated in 1 μg/ml DAPI (Sigma) for 10 minutes, washed in PBS, and mounted using Immu-mount (Shandon). Immunofluorescent imaging was performed using an INCA2200 Analyser (GE), capturing between 80 and 108 overlapping 40x fields in a region with confirmed melanoma as judged by pathological assessment of the H&E. This was performed in three blocks per sample. Images were stitched into blocks of 36 fields of view for further analysis.

Identifying sub-populations in tumour tissue sections

Stitched images were examined by a researcher and a pathologist to identify areas of positive, true (non-background) CD24+ and CD271+ staining, and these were compared to the corresponding area of the H&E. Areas of single staining (CD24+ or CD271+) and double staining (CD24+/CD271+) on the stained images were recorded if they coincided with melanoma cells. The nature and location of this staining was also recorded.

Conflict of interest

The authors declare no conflicts of interest.

Acknowledgements

We thank Gary Warnes for technical assistance and discussion. Cell lines were kindly provided by Penny Lovat, Newcastle; John Marshall, Barts & the London; Monica Rodolfo, Inst. Tumori Milano. Adrian Biddle has received funding support from Animal Free Research UK, as part of the Animal Replacement Centre of Excellence at Queen Mary University of London, and from the UK Medical Research Council (MR/V009494/1).

References

AMERICAN CANCER SOCIETY. Cancer facts & figures. Atlanta, GA: The Society.

BIDDLE, A., GAMMON, L., FAZIL, B. & MACKENZIE, I. C. 2013. CD44 staining of cancer stem-like cells is influenced by down-regulation of CD44 variant isoforms and up-regulation of the standard CD44 isoform in the population of cells that have undergone epithelial-to-mesenchymal transition. *PLoS One,* 8**,** e57314.

BIDDLE, A., GAMMON, L., LIANG, X., COSTEA, D. E. & MACKENZIE, I. C. 2016. Phenotypic Plasticity Determines Cancer Stem Cell Therapeutic Resistance in Oral Squamous Cell Carcinoma. *EBioMedicine,* 4**,** 138-45.

BIDDLE, A., LIANG, X., GAMMON, L., FAZIL, B., HARPER, L. J., EMICH, H., COSTEA, D. E. & MACKENZIE, I. C. 2011. Cancer stem cells in squamous cell carcinoma switch between two distinct phenotypes that are preferentially migratory or proliferative. *Cancer Res,* 71**,** 5317-26.

BOIKO, A. D., RAZORENOVA, O. V., VAN DE RIJN, M., SWETTER, S. M., JOHNSON, D. L., LY, D. P., BUTLER, P. D., YANG, G. P., JOSHUA, B., KAPLAN, M. J., LONGAKER, M. T. & WEISSMAN, I. L. 2010. Human melanoma-initiating cells express neural crest nerve growth factor receptor CD271. *Nature,* 466**,** 133-7.

CANTELLI, G., ORGAZ, J. L., RODRIGUEZ-HERNANDEZ, I., KARAGIANNIS, P., MAIQUES, O., MATIAS-GUIU, X., NESTLE, F. O., MARTI, R. M., KARAGIANNIS, S. N. & SANZ-MORENO, V. 2015. TGF-beta-Induced Transcription Sustains Amoeboid Melanoma Migration and Dissemination. *Curr Biol,* 25**,** 2899-914.

CARREIRA, S., GOODALL, J., DENAT, L., RODRIGUEZ, M., NUCIFORO, P., HOEK, K. S., TESTORI, A., LARUE, L. & GODING, C. R. 2006. Mitf regulation of Dia1 controls melanoma proliferation and invasiveness. *Genes Dev,* 20**,** 3426-39.

DANIOTTI, M., OGGIONNI, M., RANZANI, T., VALLACCHI, V., CAMPI, V., DI STASI, D., TORRE, G. D., PERRONE, F., LUONI, C., SUARDI, S., FRATTINI, M., PILOTTI, S., ANICHINI, A., TRAGNI, G., PARMIANI, G., PIEROTTI, M. A. & RODOLFO, M. 2004. BRAF alterations are associated with complex mutational profiles in malignant melanoma. *Oncogene,* 23**,** 5968-5977.

GEORGOULI, M., HERRAIZ, C., CROSAS-MOLIST, E., FANSHAWE, B., MAIQUES, O., PERDRIX, A., PANDYA, P., RODRIGUEZ-HERNANDEZ, I., ILIEVA, K. M., CANTELLI, G., KARAGIANNIS, P., MELE, S., LAM, H., JOSEPHS, D. H., MATIAS-GUIU, X., MARTI, R. M., NESTLE, F. O., ORGAZ, J. L., MALANCHI, I., FRUHWIRTH, G. O., KARAGIANNIS, S. N. & SANZ-MORENO, V. 2019. Regional Activation of Myosin II in Cancer Cells Drives Tumor Progression via a Secretory Cross-Talk with the Immune Microenvironment. *Cell,* 176**,** 757-774 e23.

GOLDMAN, A., MAJUMDER, B., DHAWAN, A., RAVI, S., GOLDMAN, D., KOHANDEL, M., MAJUMDER, P. K. & SENGUPTA, S. 2015. Temporally sequenced anticancer drugs overcome adaptive resistance by targeting a vulnerable chemotherapy-induced phenotypic transition. *Nat Commun,* 6**,** 6139.

GSCHAIDER, M., NEUMANN, F., PETERS, B., LENZ, F., CIBENA, M., GOISER, M., WOLF, I., WENZEL, J., MAUCH, C., SCHREINER, W. & WAGNER, S. N. 2012. An attempt at a molecular prediction of metastasis in patients with primary cutaneous melanoma. *PLoS One,* 7**,** e49865.

HERMANN, P. C., HUBER, S. L., HERRLER, T., AICHER, A., ELLWART, J. W., GUBA, M., BRUNS, C. J. & HEESCHEN, C. 2007. Distinct populations of cancer stem cells determine tumor growth and metastatic activity in human pancreatic cancer. *Cell Stem Cell,* 1**,** 313-23.

KATZ, E., DUBOIS-MARSHALL, S., SIMS, A. H., GAUTIER, P., CALDWELL, H., MEEHAN, R. R. & HARRISON, D. J. 2011. An in vitro model that recapitulates the epithelial to mesenchymal transition (EMT) in human breast cancer. *PLoS One,* 6**,** e17083.

KOZAR, I., MARGUE, C., ROTHENGATTER, S., HAAN, C. & KREIS, S. 2019. Many ways to resistance: How melanoma cells evade targeted therapies. *Biochim Biophys Acta Rev Cancer,* 1871**,** 313-322.

KOZLOWSKI, J. M., HART, I. R., FIDLER, I. J. & HANNA, N. 1984. A human melanoma line heterogeneous with respect to metastatic capacity in athymic nude mice. *J Natl Cancer Inst,* 72**,** 913-7.

LANDSBERG, J., KOHLMEYER, J., RENN, M., BALD, T., ROGAVA, M., CRON, M., FATHO, M., LENNERZ, V., WÖLFEL, T., HÖLZEL, M. & TÜTING, T. 2012. Melanomas resist T-cell therapy through inflammation-induced reversible dedifferentiation. *Nature,* 490**,** 412-6.

LI, X., LEWIS, M. T., HUANG, J., GUTIERREZ, C., OSBORNE, C. K., WU, M. F., HILSENBECK, S. G., PAVLICK, A., ZHANG, X., CHAMNESS, G. C., WONG, H., ROSEN, J. & CHANG, J. C. 2008. Intrinsic resistance of tumorigenic breast cancer cells to chemotherapy. *J Natl Cancer Inst,* 100**,** 672-9.

LIU, S., CONG, Y., WANG, D., SUN, Y., DENG, L., LIU, Y., MARTIN-TREVINO, R., SHANG, L., MCDERMOTT, S. P., LANDIS, M. D., HONG, S., ADAMS, A., D'ANGELO, R., GINESTIER, C., CHARAFE-JAUFFRET, E., CLOUTHIER, S. G., BIRNBAUM, D., WONG, S. T., ZHAN, M., CHANG, J. C. & WICHA, M. S. 2014. Breast Cancer Stem Cells Transition between Epithelial and Mesenchymal States Reflective of their Normal Counterparts. *Stem Cell Reports,* 2**,** 78-91.

LIU, W., MONAHAN, K. B., PFEFFERLE, A. D., SHIMAMURA, T., SORRENTINO, J., CHAN, K. T., ROADCAP, D. W., OLLILA, D. W., THOMAS, N. E., CASTRILLON, D. H., MILLER, C. R., PEROU, C. M., WONG, K. K., BEAR, J. E. & SHARPLESS, N. E. 2012. LKB1/STK11 inactivation leads to expansion of a prometastatic tumor subpopulation in melanoma. *Cancer Cell,* 21**,** 751-64.

LU, W. & KANG, Y. 2019. Epithelial-Mesenchymal Plasticity in Cancer Progression and Metastasis. *Dev Cell,* 49**,** 361-374.

MORRISON, S. J., WHITE, P. M., ZOCK, C. & ANDERSON, D. J. 1999. Prospective identification, isolation by flow cytometry, and in vivo self-renewal of multipotent mammalian neural crest stem cells. *Cell,* 96**,** 737-49.

MUKHERJEE, N., LAMBERT, K. A., NORRIS, D. A. & SHELLMAN, Y. G. 2021. Enrichment of Melanoma Stem-Like Cells via Sphere Assays. *Methods Mol Biol,* 2265**,** 185-199.

MÜLLER, J., KRIJGSMAN, O., TSOI, J., ROBERT, L., HUGO, W., SONG, C., KONG, X., POSSIK, P. A., CORNELISSEN-STEIJGER, P. D., GEUKES FOPPEN, M. H., KEMPER, K., GODING, C. R., MCDERMOTT, U., BLANK, C., HAANEN, J., GRAEBER, T. G., RIBAS, A., LO, R. S. & PEEPER, D. S. 2014. Low MITF/AXL ratio predicts early resistance to multiple targeted drugs in melanoma. *Nat Commun,* 5**,** 5712.

PASTUSHENKO, I., BRISEBARRE, A., SIFRIM, A., FIORAMONTI, M., REVENCO, T., BOUMAHDI, S., VAN KEYMEULEN, A., BROWN, D., MOERS, V., LEMAIRE, S., DE CLERCQ, S., MINGUIJON, E., BALSAT, C., SOKOLOW, Y., DUBOIS, C., DE COCK, F., SCOZZARO, S., SOPENA, F., LANAS, A., D'HAENE, N., SALMON, I., MARINE, J. C., VOET, T., SOTIROPOULOU, P. A. & BLANPAIN, C. 2018. Identification of the tumour transition states occurring during EMT. *Nature,* 556**,** 463-468.

PASTUSHENKO, I., MAURI, F., SONG, Y., DE COCK, F., MEEUSEN, B., SWEDLUND, B., IMPENS, F., VAN HAVER, D., OPITZ, M., THERY, M., BARECHE, Y., LAPOUGE, G., VERMEERSCH, M., VAN EYCKE, Y. R., BALSAT, C., DECAESTECKER, C., SOKOLOW, Y., HASSID, S., PEREZ-BUSTILLO, A., AGREDA-MORENO, B., RIOS-BUCETA, L., JAEN, P., REDONDO, P., SIEIRA-GIL, R., MILLAN-CAYETANO, J. F., SANMATRTIN, O., D'HAENE, N., MOERS, V., ROZZI, M., BLONDEAU, J., LEMAIRE, S., SCOZZARO, S., JANSSENS, V., DE TROYA, M., DUBOIS, C., PÉREZ-MORGA, D., SALMON, I., SOTIRIOU, C., HELMBACHER, F. & BLANPAIN, C. 2021. Fat1 deletion promotes hybrid EMT state, tumour stemness and metastasis. *Nature,* 589**,** 448-455.

PHI, L. T. H., SARI, I. N., YANG, Y. G., LEE, S. H., JUN, N., KIM, K. S., LEE, Y. K. & KWON, H. Y. 2018. Cancer Stem Cells (CSCs) in Drug Resistance and their Therapeutic Implications in Cancer Treatment. *Stem Cells Int,* 2018**,** 5416923.

PISKOUNOVA, E., AGATHOCLEOUS, M., MURPHY, M. M., HU, Z., HUDDLESTUN, S. E., ZHAO, Z., LEITCH, A. M., JOHNSON, T. M., DEBERARDINIS, R. J. & MORRISON, S. J. 2015. Oxidative stress inhibits distant metastasis by human melanoma cells. *Nature,* 527**,** 186-91.

QUINTANA, E., SHACKLETON, M., FOSTER, H. R., FULLEN, D. R., SABEL, M. S., JOHNSON, T. M. & MORRISON, S. J. 2010. Phenotypic heterogeneity among tumorigenic melanoma cells from patients that is reversible and not hierarchically organized. *Cancer Cell,* 18**,** 510-23.

QUINTANA, E., SHACKLETON, M., SABEL, M. S., FULLEN, D. R., JOHNSON, T. M. & MORRISON, S. J. 2008. Efficient tumour formation by single human melanoma cells. *Nature,* 456**,** 593-8.

RAMBOW, F., MARINE, J. C. & GODING, C. R. 2019. Melanoma plasticity and phenotypic diversity: therapeutic barriers and opportunities. *Genes Dev,* 33**,** 1295-1318.

RESTIVO, G., DIENER, J., CHENG, P. F., KIOWSKI, G., BONALLI, M., BIEDERMANN, T., REICHMANN, E., LEVESQUE, M. P., DUMMER, R. & SOMMER, L. 2017. low neurotrophin receptor CD271 regulates phenotype switching in melanoma. *Nat Commun,* 8**,** 1988.

RICHARD, G., DALLE, S., MONET, M. A., LIGIER, M., BOESPFLUG, A., POMMIER, R. M., DE LA FOUCHARDIÈRE, A., PERIER-MUZET, M., DEPAEPE, L., BARNAULT, R., TONDEUR, G., ANSIEAU, S., THOMAS, E., BERTOLOTTO, C., BALLOTTI, R., MOURAH, S., BATTISTELLA, M., LEBBÉ, C., THOMAS, L., PUISIEUX, A. & CARAMEL, J. 2016. ZEB1-mediated melanoma cell plasticity enhances resistance to MAPK inhibitors. *EMBO Mol Med,* 8**,** 1143-1161.

SATO, T., STANGE, D. E., FERRANTE, M., VRIES, R. G., VAN ES, J. H., VAN DEN BRINK, S., VAN HOUDT, W. J., PRONK, A., VAN GORP, J., SIERSEMA, P. D. & CLEVERS, H. 2011. Long-term expansion of epithelial organoids from human colon, adenoma, adenocarcinoma, and Barrett's epithelium. *Gastroenterology,* 141**,** 1762-72.

SENSI, M., CATANI, M., CASTELLANO, G., NICOLINI, G., ALCIATO, F., TRAGNI, G., DE SANTIS, G., BERSANI, I., AVANZI, G., TOMASSETTI, A., CANEVARI, S. & ANICHINI, A. 2011. Human cutaneous melanomas lacking MITF and melanocyte differentiation antigens express a functional Axl receptor kinase. *J Invest Dermatol,* 131**,** 2448-57.

TANG, M. R., WANG, Y. X., GUO, S., HAN, S. Y., LI, H. H. & JIN, S. F. 2014. CD24 expression predicts poor prognosis for patients with cutaneous malignant melanoma. *Int J Clin Exp Med,* 7**,** 4337-41.

TSAI, J. H., DONAHER, J. L., MURPHY, D. A., CHAU, S. & YANG, J. 2012. Spatiotemporal regulation of epithelial-mesenchymal transition is essential for squamous cell carcinoma metastasis. *Cancer Cell,* 22**,** 725-36.

UBELLACKER, J. M., TASDOGAN, A., RAMESH, V., SHEN, B., MITCHELL, E. C., MARTIN-SANDOVAL, M. S., GU, Z., MCCORMICK, M. L., DURHAM, A. B., SPITZ, D. R., ZHAO, Z., MATHEWS, T. P. & MORRISON, S. J. 2020. Lymph protects metastasizing melanoma cells from ferroptosis. *Nature,* 585**,** 113-118.

YOUSSEF, G., GAMMON, L., AMBLER, L., WICKER, B., PATEL, S., COTTOM, H., PIPER, K., MACKENZIE, I. C., PHILPOTT, M. P. & BIDDLE, A. 2020. Disseminating cells in human tumours acquire an EMT stem cell state that is predictive of metastasis. *bioRxiv***,** 2020.04.07.029009.

ZHU, G., XU, P., GUO, S., YI, X., WANG, H., YANG, Y., LIU, L., SHI, Q., GAO, T. & LI, C. 2020. Metastatic Melanoma Cells Rely on Sestrin2 to Acquire Anoikis Resistance via Detoxifying Intracellular ROS. *J Invest Dermatol,* 140**,** 666-675.e2.