1

Combinatorial fedratinib and venetoclax treatment is effective on human B cell acute lymphoblastic leukemia with high Flt3 expression

Sean P Rinella[1], Haley C Bell[1], David P Turicek[1], Lei Shi[1], Nguyet-Minh Hoang[2], Lixin Rui[2], Nicholas J Hess[3] and Christian M Capitini[3]

1. Department of Pediatrics, University of Wisconsin School of Medicine and Public Health, Madison, WI

2. Department of Medicine and Carbone Cancer Center, University of Wisconsin School of Medicine and Public Health, Madison, WI

3. Department of Pediatrics and Carbone Cancer Center, University of Wisconsin School of Medicine and Public Health, Madison, WI

Abstract

Treatment of relapsed/refractory B cell acute lymphoblastic leukemia (B-ALL) remains a challenge, particularly in patients who do not respond to traditional chemotherapy or immunotherapy. The objective of this study was to assess the efficacy of fedratinib, a semi selective JAK2 inhibitor and venetoclax, a selective BCL-2 inhibitor, on human B-ALL using both single-agent and combinatorial treatments. The combination treatment of fedratinib and venetoclax improved killing of the human B-ALL cell lines RS4;11 and SUPB-15 in vitro over single-agent treatments. This combinatorial effect was not detected in the human B-ALL cell line NALM-6, which was less responsive to fedratinib due to the absence of Flt3 expression. The combination treatment induces a unique gene expression profile relative to single-agent treatment and with an enrichment in apoptotic pathways. Finally, the combination treatment was superior to single agent treatment in an in vivo xenograft model of human B-ALL with a two-week treatment regimen significantly improving overall survival. Overall, our data demonstrates the

efficacy of a combinatorial treatment strategy of fedratinib and venetoclax against

human B-ALL expressing high levels of Flt3.

Key points

1. Combination fedratinib and venetoclax reduces the survival and proliferation of

$FLT3^+$ B-ALL in vitro.

2. Gene set enrichment analysis of RNA from B-ALL treated with fedratinib and

venetoclax identified dysregulation of pathways associated with apoptosis, DNA

repair and proliferation.

3. Combination fedratinib and venetoclax reduces the number of peripheral blood

B-ALL blasts in vivo, improving overall survival while also increasing CD19

expression.

Introduction

Pediatric B cell acute lymphoblastic leukemia (B-ALL) patients who relapse after initial

treatment suffer from a poor prognosis (1,2). While many relapsed/refractory pediatric

patients are salvaged with CD19-directed immunotherapies like blinatumomab or

tisagenlecleucel, the 12 month event free survival is only 50%, with CD19 loss

representing a major barrier to the long-term success of these immunotherapies (3).

Many of these immunotherapies are also associated with a high cost, making them

unattainable to some patients. It is thus essential that we investigate novel

combinatorial treatments with targeted pharmaceuticals to improve patient outcomes

and quality of life.

The dysregulation and/or activating mutations in the protein Janus kinase 2 (JAK2) lead

to downstream phosphorylation and activation of signal transducer and activator of

transcription (STAT) proteins that are predominant in many cancers (4,5). In a recent

Children's Oncology Group (COG) study (6) genomic alterations involving activation of

the JAK/STAT pathway were found in 22% of high-risk B cell ALL patient samples,

including Philadelphia (Ph)-like B-ALL.

The BCL-2 family of proteins are central regulators of apoptosis. BCL-2 inhibits

apoptosis by the preservation of mitochondrial membrane integrity, specifically

preventing the pro-apoptotic proteins BAX and BAK from release into the mitochondria

membrane and the subsequent release of cytochrome C (7). Furthermore, some high-

risk subsets of B-ALL, such as KMT2A-rearranged B-ALL or B-ALL with hypodiploidy,

express high levels of BCL-2 to prevent apoptosis (8). We and others have previously

shown that dual targeting of the JAK/STAT pathway and BCL-2 can treat acute

myelogenous leukemia (9) and T cell ALL (10), however the impact of targeting both

these pathways in B-ALL is unknown.

The overall goal of this study was to investigate the direct anti-leukemic activity of the

JAK2 inhibitor fedratinib and the BCL-2 inhibitor venetoclax either as a single-agent

treatment or combinatorial treatment. By inhibiting both of these pathways at the same

time, we aim to enhance their anti-leukemic effects on B-ALL.

Materials and Methods:

B-ALL cell lines

NALM-6 (B-ALL), RS4;11 (KMT2A-rearranged B-ALL) and SUP-B15 (Ph-like B-ALL)

were purchased from ATCC. NALM-6 and RS4;11 were cultured in RPMI 1640

(Corning) supplemented with 10% fetal bovine serum (FBS) (GeminiBio), 2 mmol/L L-

glutamine (Gibco), and 100 mg/mL streptomycin/100 U/mL penicillin (Lonza). SUP-B15

cells were cultured in McCoy 5A (modified) media (Gibco) supplemented with 20% FBS,

100 mg/mL streptomycin and 100 U/mL penicillin. RS4;11 cells were transduced to

express the luminescent reporter luciferase using pGL4.51[luc2/CMV/Neo] vector

(Promega). Cell authentication was performed using short tandem repeat analysis

(Idexx BioAnalytics, Westbrook, ME) and per ATCC guidelines using morphology,

growth curves, and *Mycoplasma* testing within 6 months of use (InVivoGen). Cell lines

were maintained in culture at 37°C in 5% CO_2.

Fedratinib and Venetoclax

Venetoclax (1 mL*10 mM in DMSO) and powdered compound were purchased from

TargetMol. Fedratinib (1 mL*10 mM in DMSO) was purchased from Selleckchem and

powdered was purchased from MedChemExpress. All concentrations for in vitro

studies were made at 2x or 4x for combination studies and diluted in RPMI-10 for

plating assays. Cells were treated with a range of doses ranging from 1nM to 10uM.

Live/dead discrimination assays

Cells were collected after a 48 hour incubation with either vehicle control, fedratinib,

venetoclax, or combination and live/dead discrimination was assessed through flow

cytometry. Samples were labeled with Ghost Dye Red 780 (Tonbo Biosciences) and

analyzed via an Attune NxT Flow Cytometer (ThermoFisher) and analysis on FlowJo

V10.

Proliferation assays

Cells were collected after a 72 hour incubation with either vehicle control, fedratinib,

venetoclax, or combination, and then proliferation was assessed through a colorimetric

MTS assay for determining cell viability. The CellTiter 96 AQueous One Solution Cell

Proliferation Assay (Promega) was used and 96 well plates were read using a

CLARIOStar microplate reader (BMG Labtech).

RNA seq

RS4;11 cells were treated for 24 hours in triplicate (untreated, fedratinib alone,

venetoclax alone, or combination). All RNA isolation and sequencing was performed by

the Gene Expression Center at the University of Wisconsin-Madison. RNA was isolated

with a RNeasy kit (Qiagen) prior to a quality control check using an Agilent Bioanalyzer

and NanoDrop One spectrophotometer. mRNA libraries were prepared using the

TruSeq stranded mRNA kit and sequenced on an Illumina NovaSeq 6000 platform to 30

million reads per sample. Sample reads were processed by the University of Wisconsin

Biotechnology Center.

In vivo studies

NOD/SCID/*Il2rg^tmwjl*/Szj (NSG) breeder mice were purchased from Jackson Labs (Bar

Harbor, ME) and bred internally at the Biomedical Research Model Services (BRMS)

Breeding Core at the University of Wisconsin-Madison in accordance with the Guide for

the Care and Use of Laboratory Mice and experiments were performed under an IACUC

approved animal protocol (M005915). Male and female NSG mice were used at 8-16

weeks of age and injected via tail vein with 2 x10^6 RS4;11 cells on Day 0. Peripheral

blood was obtained via retro-orbital bleeds conducted on Days 10 (engraftment), Day

21 (pre-treatment), and Day 35 (post-treatment). Mice were treated with fourteen days

of once daily treatment with vehicle, fedratinib, venetoclax, or combination starting at

Day 21 (5 days on, 2 days off). Venetoclax was administered via intraperitoneal (IP)

injections at a dose of 6.25 mg/kg. Fedratinib was administered via oral gavage at a

dose of 100 mg/kg. Mice were monitored for survival and percent weight change and

euthanized when a humane endpoint was reached. Peripheral blood was treated with

ACK lysis buffer (Lonza) and then stained in flow buffer (PBS, 10% FBS) with CD19

(HIB19), CD20 (2H7), CD22 (HIB22), PD-L1 (MIH2), FAS (DX2), HLA-DR (L243), BCL2

(100) and BAX (2D2), all purchased from Biolegend, prior to quantification on an Attune

NxT Flow Cytometer with analysis on FlowJo V10.

<u>RNA-Seq analysis</u>

Bioinformatic analysis of transcriptomic data adhere to recommended ENCODE

guidelines and best practices for RNA-Seq. Alignment of adapter-trimmed (Skewer

v0.1.123; (11)) 2x150 (paired-end; PE) bp strand-specific Illumina reads to the *Homo*

*sapiens*GRCh38 genome (assembly accession NCBI:GCA_000001405.25) was

achieved with the Spliced Transcripts Alignment to a Reference (STAR v2.5.3a)

software (12), a splice-junction aware aligner, using annotation provided by Ensembl

release 104. Expression estimation was performed with RSEM v1.3.0 (RNASeq by

Expectation Maximization; (13)). To test for differential gene expression among

individual group contrasts, expected read counts obtained from RSEM were used as

input into edgeR (v3.16.5; (14)). Inter-sample normalization was achieved with the

trimmed mean of M-values (TMM; (15)) method. Statistical significance of the negative-

binomial regression test was adjusted with a Benjamini-Hochberg FDR correction at the

5% level (16). Prior to statistical analysis with edgeR, independent filtering was applied

and required genes to have a count-per-million (CPM) above k in n samples, where k is

determined by minimum read count (10 reads) and by the sample library sizes

where n is determined by the number of biological replicates in each group. The validity

of the Benjamini-Hochberg FDR multiple testing procedure was evaluated by inspection

of the uncorrected p-value distribution.

<u>Statistical analysis</u>

Statistics were performed using GraphPad Prism version 9.0 for the Macintosh OS

(GraphPad Software). Data were expressed as mean ± SEM. For analysis of three or

more groups, a non-parametric one-way ANOVA test was performed with the Bonferroni

or Sidak's multiple comparisons post-test. Analysis of differences between two normally

distributed test groups was performed using a two-sided Mann Whitney test. A p-value

less than 0.05 was considered statistically significant.

Results:

Combinatorial treatment of fedratinib and venetoclax effectively kills human B-ALL in

vitro

To assess the efficacy of JAK inhibitors to kill human B-ALL cells, we tested ruxolitinib

(JAK1/2 inhibitor), lestaurtinib (semi-selective JAK2 inhibitor) and fedratinib (semi-

selective JAK2 inhibitor) on the human B-ALL cell line RS4;11. Cells were treated with a

range of doses ranging from 1nM to 10uM for 48-72 hours prior to analysis for cell

survival (Figure 1A-B, Suppl Figure 1). Survival and proliferation of RS4;11 was

assessed by flow cytometry at 48hrs or through a colorimetric MTS assay (Figure 1A-

B). Based on the dose-curves, the calculated IC50 for lestaurtinib was 60nM, fedratinib

was 600nM but the IC50 for ruxolitinib was not reached.

We next wanted to explore whether or not fedratinib would be more efficacious when
combined with venetoclax. To test this, we treated RS4;11 with single agent fedratinib at
700nm, single agent venetoclax at 7.5nM and the combinatorial treatment (Figure 1D-
E). Combinatorial treatment was significantly more effective than either single agent
treatment and similar to the positive control staurosporine. We next tested if ruxolitinib
had a similar beneficial effect as fedratinib (Fig 1F-G). Ruxolitinib, used at a higher dose
than fedratinib, did not improve the killing of RS4;11 over single-agent venetoclax (Fig
1F-G). To ensure that the combinatorial effect of fedratinib and venetoclax was not
specific to RS4;11, we tested both the single agent and combinatorial treatments on the
human B-ALL cell lines SUPB-15 and NALM-6. The combinatorial treatment was
equally effective on SUPB-15 (Fig 2A-B) but surprisingly, the combinatorial treatment
did not improve the killing of NALM-6 cells compared to venetoclax alone (Fig 2C-D).
These data suggest that while the combinatorial treatment of fedratinib and venetoclax
can be highly effective in human B-ALL, there are a subset of B-ALLs that remain
resistant to this combinatorial treatment.
198
199   *Resistance to fedratinib treatment is correlated with absence of Flt3 expression*
200
To determine the precise activity of and contribution to cell killing by fedratinib and
venetoclax, we performed several additional dose-escalation studies on all three human
B-ALL cell lines. Keeping a constant venetoclax dose of 5 nM, both RS4;11 and SUPB-
15 experienced significant cell death that was further enhanced with increasing doses of
fedratinib. Conversely, NALM-6 was mostly refractive to venetoclax at 5 nM with
increasing concentrations of fedratinib only modestly improving the killing efficiency (Fig
3A-C, left panels). Using a constant dose of fedratinib of 525 nM, increasing doses of
venetoclax was able to significantly increase cell death in both RS4;11 and SUP-B15
while NALM-6 remains refractory as before (Fig 3 A-C, right panels).
210
To assess a potential mechanism for NALM-6 resistance to both venetoclax and
fedratinib treatment, we measured the expression of BCL-2, BRD4 and FLT3 in each
human B-ALL cell line. Notably, NALM-6 had no expression of FLT3, which is also a

target of fedratinib while RS4;11, which has the greatest sensitivity to fedratinib, had the

highest expression of Flt3 (Fig 3D-E). We also observed that NALM-6 was more

resistant to venetoclax treatment than either RS4;11 or SUPB-15 (Fig 1, 2).

Interestingly, NALM-6 had a significantly lower ratio of the pro-apoptotic protein BAX to

the anti-apoptotic protein BCL2 relative to RS4;11, which correlates with their

venetoclax sensitivity (Fig 3F).

Combination therapy differentially enriches pathways related to cell death

Next, we wanted to identify the dominant pathways altered in RS4;11 post single-agent

and combination treatment via bulk RNA sequencing. RS4;11 cells were plated similarly

as described above and exposed for 24 hours to fedratinib (525 nM), venetoclax (5 nM),

or combination and compared to untreated cells. Cells were harvested after 24 hours for

RNAseq. Whole transcriptome analysis revealed that each treatment regimen induced a

unique expression profile clustered image map (Fig 4A-B). Comparing each treatment

regimen to untreated cells, fedratinib had the fewest number of significantly different

genes while the combinatorial treatment had the most (Fig 4C-E). Interestingly in all

cases, most of the differentially expressed genes were down-regulated and not up-

regulated as a result of treatment (Fig 4C-E). We also completed a gene set enrichment

analysis (GSEA) that identified pathways associated with apoptosis, DNA repair and

proliferation predominantly altered in the combinatorial treatment versus all three other

conditions (Fig 4F). A heatmap of the differentially enriched genes primarily impacting

the enrichment into the apoptotic pathway is also shown (Fig 4G).

Combination therapy reduces leukemic burden and improves overall survival in RS4;11

derived xenograft model of B-ALL

Based on the efficacy of fedratinib and venetoclax observed *in vitro*, a xenograft model

of B-ALL was established utilizing RS4;11 cells and NSG mice to test these drugs as a

combination therapy *in vivo*. NSG mice were injected with luciferase[+] RS4;11 cells on

Day 0 and then randomized to one of the following groups: untreated, single agent

venetoclax, single agent fedratinib, or combination. Peripheral blood was obtained via

retroorbital bleeds on days 10 (engraftment), 21 (pretreatment), and 35 (post treatment)

to assess B-ALL disease burden (Fig 5A). Treatment started on day 21 and lasted for

14 days (5 days on, 2 days off) with 6.25 mg/kg venetoclax IP and/or 100 mg/kg

fedratinib via gavage. Combination treatment prolonged the survival of the mice and

decreased the RS4;11 burden the most compared to the other treatment groups (Fig

5B). Peripheral blood counts and IVIS imaging throughout the experiment revealed that

combination treatment was highly effective at decreasing the RS4;11 burden in vivo (Fig

5C-F). Venetoclax alone also significantly improved overall survival when compared to

fedratinib alone and untreated mice (Figure 5B-F). Flow cytometric analysis of RS4;11

taken from the peripheral blood also showed a significant increase in MFI of FAS and

CD19 in the combination group. Overall, these data support the role of fedratinib and

venetoclax as a combination therapy against FLT3$^+$ B-ALL.

Discussion:

Treatment of relapsed/refractory B cell acute lymphoblastic leukemia (B-ALL) remains a

challenge, particularly in patients who do not respond to traditional chemotherapy or

immunotherapy. Targeting BCL-2 in B-ALL has been shown previously to be efficacious

using venetoclax. One group showed a single dose of 12.5 mg/kg venetoclax in a

RS4;11-derived xenograft model led to a maximal tumor growth inhibition of 47% and

tumor growth delay of 26%. However, maximal effects were seen when combining

venetoclax with chemotherapy (17). Other studies demonstrating efficacy of venetoclax

in B-ALL include combining with MCL-1 inhibitors (18, 19), monoclonal antibodies such

as daratumumab in relapsed/refractory B-ALL (20), AKT inhibitors (21, 22), MDM2

inhibitors (23), and also with BCR-ABL positive B-ALL (24). No other studies have

previously looked at the combination of JAK2 inhibitors with venetoclax in B-ALL. The

dysregulation of JAK/STAT related genes has been previously shown to be found in

high risk patient samples of B-ALL (6). One group identified rearrangements of JAK2 in

Ph like B-ALL involving the activation of mutations in IL7R and FLT3, and deletion of
SH2B3, which encodes the JAK2 negative regulator LNK (25).
276
The goal of this study was to suppress leukemia proliferation related to JAK/STAT
dysregulation while at the same time inhibiting BCL-2 to prevent escape from apoptosis
in models of human B-ALL. To do this, we tested fedratinib, a semi selective JAK2
inhibitor, and venetoclax, a potent BCL2 inhibitor, to assess the efficacy of this
combination in 3 models of human B-ALL in vitro and with the RS4;11 cell line in vivo.
Fedratinib has been FDA approved for patients with myeloproliferative disorders,
primarily myelofybrosis (26, 27) and has been used to salvage patients that have failed
post ruxolitinib therapy (28). Furthermore, fedratinib has been used to decrease tumor
burden in Hodgkin lymphoma and mediastinal large B-cell lymphoma in vitro and in vivo
(29). However, fedratinib has not previously been used for B-ALL. Analysis of BRD4
and FLT3 was included for all 3 cell lines in our study as fedratinib has been shown to
target both BRD4 (30, 31) and FLT3 (30) in addition to its activity against JAK2.
289
Our data both in vitro and in vivo demonstrated that the combination of fedratinib and
venetoclax more effectively treated B-ALL than single agent treatment. We also
examined the use of other JAK inhibitors, including ruxolitinib, which we had previously
shown synergized with venetoclax to treat T cell ALL (10). Similar to observations by
others (18), we found that ruxolitinib alone was ineffective in doses up to 10 uM. That
group did demonstrate that there may be some efficacy when ruxolitinib was combined
with venetoclax in Ph+ B-ALL models of disease in vitro. Combination fedratinib and
venetoclax did demonstrate efficacy in another high-risk Ph+ B-ALL cell line SUP-B15,
perhaps suggesting that the combination of targeting the JAK/STAT and BCL-2
pathways simultaneously is particularly effective in Ph-like and FLT3+ high risk subsets
compared to more standard B-ALL models of disease such as NALM-6. Furthermore,
lestaurtinib, another semi selective JAK inhibitor was tested in preliminary studies.
Lestaurtinib, similar to fedratinib, also inhibits FLT3 which is commonly overexpressed
in KMT2A-rearranged B-ALL (32) and was found to be highly expressed in RS4;11. The
association of FLT3 and BCL2 in B-ALL is unclear but our RNA GSEA suggest the

potential roles of proliferation, apoptosis and DNA repair pathways with combination

fedratinib and venetoclax treatment.

Finally, we demonstrated efficacy of combination fedratinib and venetoclax therapy in a

B-ALL xenograft model. In fact, with only a 2-week treatment we were able to identify

reduced leukemic burden and an increase in overall survival benefit with combination

therapy. Our group attempted to reduce potential toxicity in our in vivo model by

reducing DMSO concentration in our venetoclax drug preparation for in vivo use and

gavaging with sucrose-soaked gavage tips to reduce mice stress as much as possible

using techniques described by others (33). Our group also identified an increase in

CD19 expression in peripheral blood samples obtained from mice that received either

venetoclax or combination therapy. One group recently demonstrated pre-sensitizing B

cell malignancies, including B-ALL, through venetoclax exposure increases cytotoxic

efficacy of CD19 directed CAR-T therapy (34). These findings along with our data with

combination venetoclax plus fedratinib suggest these targeted therapies may serve as a

potential bridge therapy to sensitize B-ALL to CD19 targeted treatments such as

blinatumomab or chimeric antigen receptor (CAR) T cells.

The development of novel treatment strategies for relapsed/refractory B-ALL is essential

for the subset of patients with poor outcomes due to failure to respond to initial

conventional chemotherapy. The combination of fedratinib and venetoclax is an

effective treatment strategy to reduce the burden of disease, particularly in FLT3$^+$ B-

ALL, through reduction in proliferation, apoptosis and DNA repair pathways.

Combination fedratinib and venetoclax may also potentially serve as a bridge to CD19-

directed therapies such as CAR-T cell and future studies should explore if these

therapies can help B-ALL avoid potential CD19 antigen escape.

Acknowledgements:

We would like to thank the University of Wisconsin Carbone Cancer Center (UWCCC)

Flow Cytometry and Small Animal Imaging and Radiotherapy core facilities, who are

supported in part through NCI/NIH P30 CA014520 as well as the UW Biotechnology
Center.

Author contributions:

341 SPR, NJH and CMC designed the experiments, analyzed and interpreted results and
342 wrote the manuscript; SPR, NJH, HCB, LS, NMH, LR, DPT, and NJH conducted
343 experiments and analyzed data; SPR, NJH, DPT and HCB analyzed data and
344 generated figures; and SPR, NJH, and HCB discussed and interpreted results. All
345 authors read and approved the manuscript.

Disclosure statement:

348 C.M.C receives honorarium for advisory board membership for Bayer, Elephas, Nektar
349 Therapeutics, Novartis and WiCell Research Institute, who had no input in the study
350 design, analysis, manuscript preparation or decision to submit. All other authors declare
351 they have no competing interests.

Funding:

This work was supported by grants from the NIH TL1 TR002375 (S.P.R), NIH/NCATS
UL1-TR002373, the Cormac Pediatric Leukemia Fellowship and the Stem Cell and
Regenerative Medicine Center Fellowship (N.J.H.), NCI/NIH R01 CA187299 (L.R.), St.
Baldrick's Empowering Pediatric Immunotherapy for Childhood Cancers Grant, NCI/NIH
R01 CA215461 and the MACC Fund (C.M.C). The contents of this article do not
necessarily reflect the views or policies of the Department of Health and Human
Services, nor does mention of trade names, commercial products, or organizations
imply endorsement by the US Government. None of these funding sources had any
input in the study design, analysis, manuscript preparation or decision to submit for
publication.

References

References

1. Hunger SP, Lu X, Devidas M, et al. Improved survival for children and adolescents

with acute lymphoblastic leukemia between 1990 and 2005: a report from the Children's

Oncology Group. *JCO*. 2012;30(14):1663-1669. doi:10.1200/JCO.2011.37.8018

2. Bhojwani D, Pui CH. Relapsed childhood acute lymphoblastic leukaemia. *The Lancet*

Oncology. 2013;14(6):e205-e217. doi:10.1016/S1470-2045(12)70580-6

3. Leahy AB, Elgarten CW, Grupp SA, Maude SL, Teachey DT. Tisagenlecleucel for the

treatment of B-cell acute lymphoblastic leukemia. *Expert Review of Anticancer Therapy*.

2018;18(10):959-971. doi:10.1080/14737140.2018.1512411

4. Loh CY, Arya A, Naema AF, Wong WF, Sethi G, Looi CY. Signal transducer and

activator of transcription (STAT) proteins in cancer and inflammation: functions and

therapeutic implication. *Front Oncol*. 2019;9:48. doi:10.3389/fonc.2019.00048

5. Bousoik E, Montazeri Aliabadi H. "Do we know jack" about JAK? A closer look at

JAK/STAT signaling pathway. *Front Oncol*. 2018;8:287. doi:10.3389/fonc.2018.00287

6. Reshmi SC, Harvey RC, Roberts KG, et al. Targetable kinase gene fusions in high-

risk B-ALL: a study from the Children's Oncology Group. *Blood*. 2017;129(25):3352-

3361. doi:10.1182/blood-2016-12-758979

7. Shamas-Din A, Kale J, Leber B, Andrews DW. Mechanisms of action of BCL-2 family

proteins. *Cold Spring Harbor Perspectives in Biology*. 2013;5(4):a008714-a008714.

doi:10.1101/cshperspect.a008714

8. Diaz-Flores E, Comeaux EQ, Kim KL, et al. Bcl-2 is a therapeutic target for

hypodiploid B-lineage acute lymphoblastic leukemia. *Cancer Research*.

2019;79(9):2339-2351. doi:10.1158/0008-5472.CAN-18-0236

9. Karjalainen R, Pemovska T, Popa M, et al. JAK1/2 and BCL2 inhibitors synergize to

counteract bone marrow stromal cell–induced protection of AML. *Blood*.

2017;130(6):789-802. doi:10.1182/blood-2016-02-699363

10. Walker KL, Rinella SP, Hess NJ, et al. CXCR4 allows T cell acute lymphoblastic

leukemia to escape from JAK1/2 and BCL2 inhibition through CNS infiltration. *Leukemia*

& Lymphoma. 2021;62(5):1167-1177. doi:10.1080/10428194.2021.1910684

11. Jiang H, Lei R, Ding SW, Zhu S. Skewer: a fast and accurate adapter trimmer for

next-generation sequencing paired-end reads. *BMC Bioinformatics*. 2014;15(1):182.

doi:10.1186/1471-2105-15-182

12. Dobin A, Davis CA, Schlesinger F, et al. STAR: ultrafast universal RNA-seq

aligner. *Bioinformatics*. 2013;29(1):15-21. doi:10.1093/bioinformatics/bts635

13. Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with

or without a reference genome. *BMC Bioinformatics*. 2011;12(1):323. doi:10.1186/1471-

2105-12-323

14. Robinson MD, McCarthy DJ, Smyth GK. Edger : a bioconductor package for

differential expression analysis of digital gene expression data. *Bioinformatics*.

2010;26(1):139-140. doi:10.1093/bioinformatics/btp616

15. Robinson MD, Oshlack A. A scaling normalization method for differential expression

analysis of RNA-seq data. *Genome Biol*. 2010;11(3):R25. doi:10.1186/gb-2010-11-3-r25

16.Reiner A, Yekutieli D, Benjamini Y. Identifying differentially expressed genes using

false discovery rate controlling procedures. *Bioinformatics*. 2003;19(3):368-375.

doi:10.1093/bioinformatics/btf877

17. Souers AJ, Leverson JD, Boghaert ER, et al. ABT-199, a potent and selective BCL-
2 inhibitor, achieves antitumor activity while sparing platelets. *Nat Med*. 2013;19(2):202-
208. doi:10.1038/nm.3048
430
431   18. Moujalled DM, Hanna DT, Hediyeh-zadeh S, et al. Cotargeting BCL-2 and MCL-1 in
432   high-risk B-ALL. *Blood Advances*. 2020;4(12):2762-2767.
433   doi:10.1182/bloodadvances.2019001416
434
435   19. Mayumi A, Tomii T, Kanayama T, et al. The combination of ruxolitinib and Bcl-2/Mcl-
436   1 inhibitors has a synergistic effect on leukemic cells carrying a SPAG9::JAK2
437   fusion. *Cancer Gene Ther*. 2022;29(12):1930-1938. doi:10.1038/s41417-022-00511-z
438
439   20. Voruz S, Blum S, De Leval L, Schoumans J, Solly F, Spertini O. Daratumumab and
440   venetoclax in combination with chemotherapy provide sustained molecular remission in
441   relapsed/refractory CD19, CD20, and CD22 negative acute B lymphoblastic leukemia
442   with KMT2A-AFF1 transcript. *Biomark Res*. 2021;9(1):92. doi:10.1186/s40364-021-
443   00343-3
444
445   21. Richter A, Fischer E, Holz C, et al. Combined application of pan-AKT inhibitor MK-
446   2206 and BCL-2 antagonist venetoclax in B-cell precursor acute lymphoblastic
447   leukemia. *IJMS*. 2021;22(5):2771. doi:10.3390/ijms22052771
448
449   22. Holz C, Lange S, Sekora A, et al. Combined BCL-2 and PI3K/AKT pathway
450   inhibition in KMT2A-rearranged acute B-lymphoblastic leukemia cells. *IJMS*.
451   2023;24(2):1359. doi:10.3390/ijms24021359
452
453   23. Hohtari H, Kankainen M, Adnan-Awad S, et al. Targeting apoptosis pathways with
454   BCL2 and MDM2 inhibitors in adult B-cell acute lymphoblastic leukemia. *HemaSphere*.
455   2022;6(3):e701. doi:10.1097/HS9.0000000000000701
456

24. Massimino M, Tirrò E, Stella S, et al. Targeting BCL-2 as a therapeutic strategy for

primary p210 BCR-ABL1-positive B-ALL cells. *In Vivo*. 2020;34(2):511-516.

doi:10.21873/invivo.11802

25. Roberts KG, Morin RD, Zhang J, et al. Genetic alterations activating kinase and

cytokine receptor signaling in high-risk acute lymphoblastic leukemia. *Cancer Cell*.

2012;22(2):153-166. doi:10.1016/j.ccr.2012.06.005

26. Pardanani A, Harrison C, Cortes JE, et al. Safety and efficacy of fedratinib in

patients with primary or secondary myelofibrosis: a randomized clinical trial. *JAMA*

Oncol. 2015;1(5):643. doi:10.1001/jamaoncol.2015.1590

27. Harrison CN, Schaap N, Vannucchi AM, et al. Safety and efficacy of fedratinib, a

selective oral inhibitor of Janus kinase-2 (JAK2), in patients with myelofibrosis and low

pretreatment platelet counts. *Br J Haematol*. 2022;198(2):317-327.

doi:10.1111/bjh.18207

28. Harrison CN, Schaap N, Mesa RA. Management of myelofibrosis after ruxolitinib

failure. *Ann Hematol*. 2020;99(6):1177-1191. doi:10.1007/s00277-020-04002-9

29. Hao Y, Chapuy B, Monti S, Sun HH, Rodig SJ, Shipp MA. Selective JAK2 inhibition

specifically decreases Hodgkin lymphoma and mediastinal large B-cell lymphoma

growth in vitro and in vivo. *Clinical Cancer Research*. 2014;20(10):2674-2683.

doi:10.1158/1078-0432.CCR-13-3007

30. Talpaz M, Kiladjian JJ. Fedratinib, a newly approved treatment for patients with

myeloproliferative neoplasm-associated myelofibrosis. *Leukemia*. 2021;35(1):1-17.

doi:10.1038/s41375-020-0954-2

31. Ciceri P, Müller S, O'Mahony A, et al. Dual kinase-bromodomain inhibitors for
rationally designed polypharmacology. *Nat Chem Biol*. 2014;10(4):305-312.
doi:10.1038/nchembio.1471
489

490 32. Armstrong SA, Kung AL, Mabon ME, et al. Inhibition of FLT3 in MLL. *Cancer Cell*.
491 2003;3(2):173-183. doi:10.1016/S1535-6108(03)00003-5
492

493 33. Hoggatt AF, Hoggatt J, Honerlaw M, Pelus LM. A spoonful of sugar helps the
494 medicine go down: a novel technique to improve oral gavage in mice. *J Am Assoc Lab*
495 *Anim Sci*. 2010;49(3):329-334
496

497 34. Yang M, Wang L, Ni M, et al. Pre-sensitization of malignant B cells through
498 venetoclax significantly improves the cytotoxic efficacy of CD19 CAR-T cells. *Front*
499 *Immunol*. 2020;11:608167. doi:10.3389/fimmu.2020.608167
500

Figure 1. Fedratinib, but not ruxolitinib, combines with venetoclax to improve
killing of human B-ALL
(A-C) The human B-ALL cell line RS4;11 was treated with a dose curve of single agent
ruxolitinib (A), fedratinib (B) or venetoclax (C) and number of live cells quantified by flow
cytometry after 48hrs. (D-G) RS4;11 cells were treated with the indicated concentration
of fedratinib and venetoclax (D-E) or ruxolitinib and venetoclax (F-G) for 48hrs. Live
cells were quantified by flow cytometry and proliferation rate measured by colorimetric
MTS assay. Fed = fedratinib. Rux = ruxolitinib. Ven = venetoclax.
509

Figure 2. Combinatorial treatment of venetoclax and fedratinib is efficacious in
the human B-ALL cell line SUP-B15, but not NALM-6
Human B-ALL cell lines NALM-6 and SUP-B15 were treated with venetoclax in
combination with fedratinib. Live cells were quantified by flow cytometry as previously
described (A,C) and proliferation was assessed by an MTS assay as previously
described (B,D).
516

Figure 3. Fedratinib activity is associated with Flt3 expression

(A-C) Combination dose curves were performed on RS4;11 (A), SUP-B15 (B) and

NALM-6 (C), keeping either the venetoclax dose (left panel) or fedratinib dose (right

panel) constant to determine the potency of the combinatorial treatment on different

human B-ALL cell lines. (D-F) Expression of both Flt3 and Brd4 (D-E) and the ratio of

BAX to BCL2 (F) were measured in each cell line by flow cytometry. Fold MFI

represents the median fluorescent intensity of the marker divided by the isotype control.

*** $p < 0.001$.

Figure 4. Combination treatment induces a unique gene expression profile

RS4;11 cells were left untreated or treated with 5 nM ventoclax, 525 nM fedratinib or the

combination treatment for 24 hrs prior to RNA isolation and bulk sequencing. (A-B)

Transcriptome analysis of each treatment group by two-dimensional multidimensional

scaling (MDS) (A) and a clustered image map reveals each treatment group induces a

unique transcriptomic profile. (C-E) Volcano plots showing the number of positively and

negatively regulated genes in each treatment group relative to the untreated control. (F)

Gene set enrichment analysis of the combination treatment against all other treatments

was performed with three gene sets shown. (G) Heatmap showing the genes

contributing to the enrichment score of the apoptosis gene set.

Figure 5. Fedratinib and venetoclax combination treatment is superior to single-

agent treatment in vivo

(A) RS4;11 was transplanted into non-irradiated NSG mice via tail vein injection at 2E6

cells per mouse and allowed to engraft for 21 days prior to starting treatment.

Venetoclax was given IP at a dose of 6.25 mg/kg while fedratinib was given gavage at a

dose of 100 mg/kg. Retro-orbital blood draws were taken at day 10, 21 and 35 to

assess cancer burden. (B) Kaplan-meier survival curve of each treatment. (C-D)

Representative dot plot of RS4;11 taken from the blood at day 35 (C) and quantification

across all times (D). (E-F) Representative IVIS images (E) and total flux quantification of

all mice at the indicated time points (F). (G) A select number of B-ALL cancer markers

expression profiles were assessed at day 35 across all treatment conditions. Fold MFI

represents the median fluorescent intensity of the marker divided by the isotype control.

Data is representative of two independent experiments. * $p < 0.05$, ** $p < 0.01$, *** $p <$

0.001.

Supplementary Figure 1. Single agent lestaurtinib and in combination with

venetoclax show efficacy against RS4;11

The human B-ALL cell line RS4;11 was treated with single agent lestaurtinib for 48hrs

(A) or in combination with venetoclax (B) and live cells quantified by flow cytometry.

Supplementary Figure 2. FISH of bone marrow and spinal cord samples from

untreated Day 51 RS4;11 transplanted NSG mice show very limited CNS infiltrate

during late stage leukemic disease

Bone marrow and spinal cord tissue were harvested on untreated Day 51 transplanted

RS4;11 NSG mice. Human and mouse CEP probes via FISH were used to quantify

human RS4;11 cells versus mouse cells to determine disease burden in bone marrow

and spinal cord tissues and to confirm level of CNS infiltrate for subsequent treatment

with small molecule inhibitors.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5