Combination fedratinib and venetoclax has activity against human B-ALL with high FLT3 expression

Sean P Rinella[1], Haley C Bell[1], Nicholas J Hess[1], Nguyet-Minh Hoang[2], Thao Trang Nguyen[3], David P Turicek[1], Lei Shi[1], Lixin Rui[2,4], James L LaBelle[3], and Christian M Capitini[1,3]

1. Department of Pediatrics, University of Wisconsin School of Medicine and Public Health, Madison, WI

2. Department of Medicine, University of Wisconsin School of Medicine and Public Health, Madison, WI

3. Department of Pediatrics, Section of Hematology/Oncology, University of Chicago, Chicago, IL

4. Carbone Cancer Center, University of Wisconsin School of Medicine and Public Health, Madison, WI

Running title

Fedratinib and venetoclax have activity against B-ALL

Corresponding author

Sean Rinella, 1111 Highland Ave, WIMR 4118, Madison WI 53705 srinella@wisc.edu

Phone 608-265-2445

Disclosures:

C.M.C receives honorarium for advisory board membership for Bayer, Elephas, Nektar Therapeutics, Novartis and WiCell Research Institute; J.L.L receives funding from

27 AbbVie as part of the University of Chicago Collaborative Grant. These entities had no

28 input in the study design, analysis, manuscript preparation or decision to submit. All

29 other authors declare they have no competing interests.

Abstract

Treatment of relapsed/refractory B cell acute lymphoblastic leukemia (B-ALL) remains a challenge, particularly in patients who do not respond to traditional chemotherapy or immunotherapy. The objective of this study was to assess the efficacy of fedratinib, a semi selective JAK2 inhibitor and venetoclax, a selective BCL-2 inhibitor, on human B-ALL using both single-agent and combinatorial treatments. The combination treatment of fedratinib and venetoclax improved killing of the human B-ALL cell lines RS4;11 and SUPB-15 in vitro over single-agent treatments. This combinatorial effect was not detected in the human B-ALL cell line NALM-6, which was less responsive to fedratinib due to the absence of Flt3 expression. The combination treatment induces a unique gene expression profile relative to single-agent treatment and with an enrichment in apoptotic pathways. Finally, the combination treatment was superior to single agent treatment in an in vivo xenograft model of human B-ALL, with a two-week treatment regimen significantly improving overall survival while inducing CD19 expression. Overall, our data demonstrates the efficacy of a combinatorial treatment strategy of fedratinib and venetoclax against human B-ALL expressing high levels of Flt3.

Statement of Implication

The combination treatment of fedratinib and venetoclax has activity against human B-ALL with high FLT3 expression and has the potential to be a salvage therapy for patients with relapsed/refractory B-ALL.

Introduction

While significant strides have been made to treating B cell acute lymphoblastic leukemia (B-ALL) with now over 80% children cured, patients who relapse after initial treatment suffer a poor prognosis (1,2). While many relapsed/refractory pediatric patients are salvaged with CD19-directed immunotherapies like blinatumomab or tisagenlecleucel, the 12-month event free survival is only 50%, with CD19 loss representing a major barrier to the long-term success of these immunotherapies (3). It is thus essential to investigate novel combinatorial treatments with targeted pharmaceuticals to improve patient outcomes and quality of life.

The dysregulation and/or activating mutations in the protein Janus kinase 2 (JAK2) lead to downstream phosphorylation and activation of signal transducer and activator of transcription (STAT) proteins that are predominant in many cancers (4,5). In a recent Children's Oncology Group (COG) study (6) genomic alterations involving activation of the JAK/STAT pathway were found in 22% of high-risk B cell ALL patient samples, including Philadelphia (Ph)-like B-ALL.

The BCL-2 family of proteins are central regulators of apoptosis. BCL-2 inhibits apoptosis by the preservation of mitochondrial membrane integrity, specifically preventing the pro-apoptotic proteins BAX and BAK from release into the mitochondria membrane and the subsequent release of cytochrome C (7). Furthermore, some high-risk subsets of B-ALL, such as KMT2A-rearranged B-ALL or B-ALL with hypodiploidy, express high levels of BCL-2 to prevent apoptosis (8). We and others have previously shown that dual targeting of the JAK/STAT pathway and BCL-2 can treat acute myelogenous leukemia (9) and T cell ALL (10), however the impact of targeting both these pathways in B-ALL is unknown.

The overall goal of this study was to investigate the direct anti-leukemic activity of the JAK2 inhibitor fedratinib and the BCL-2 inhibitor venetoclax either as a single-agent treatment or combinatorial treatment. By inhibiting both of these pathways at the same time, we aim to enhance their anti-leukemic effects on B-ALL.

Materials and Methods:

B-ALL cell lines

NALM-6 (B-ALL), RS4;11 (KMT2A-rearranged B-ALL) and SUP-B15 (Ph-like B-ALL)
were purchased from ATCC. NALM-6 and RS4;11 were cultured in RPMI 1640
(Corning) supplemented with 10% fetal bovine serum (FBS) (GeminiBio), 2 mmol/L L-
glutamine (Gibco), and 100 mg/mL streptomycin/100 U/mL penicillin (Lonza). SUP-B15
cells were cultured in McCoy 5A (modified) media (Gibco) supplemented with 20% FBS,
100 mg/mL streptomycin and 100 U/mL penicillin. RS4;11 cells were transduced to
express the luminescent reporter luciferase using pGL4.51[luc2/CMV/Neo] vector
(Promega). Cell authentication was performed using short tandem repeat analysis
(Idexx BioAnalytics, Westbrook, ME) and per ATCC guidelines using morphology,
growth curves, and *Mycoplasma* testing within 6 months of use (InVivoGen). Cell lines
were maintained in culture at 37°C in 5% CO_2.

Fedratinib and Venetoclax

Venetoclax (1 mL*10 mM in DMSO) and powdered compound were purchased from
TargetMol. Fedratinib (1 mL*10 mM in DMSO) was purchased from Selleckchem and
powdered was purchased from MedChemExpress. All concentrations for in vitro
studies were made at 2x or 4x for combination studies and diluted in RPMI with 10%
FBS for plating assays. Cells were treated with a range of doses ranging from 1nM to
10uM.

Live/dead discrimination assays

Cells were collected after a 48 hour incubation with either vehicle control, fedratinib,
venetoclax, or combination and live/dead discrimination was assessed through flow
cytometry. Samples were labeled with Ghost Dye Red 780 (Tonbo Biosciences) and
analyzed via an Attune NxT Flow Cytometer (ThermoFisher) with analysis on FlowJo
V10.

<u>Proliferation assays</u>

Cells were collected after a 72 hour incubation with either vehicle control, fedratinib,

venetoclax, or combination using the CellTiter 96 AQueous One Solution Cell

Proliferation Assay (Promega). Plates were read using a CLARIOStar microplate reader

(BMG Labtech).

<u>RNA seq</u>

RS4;11 cells were treated for 24 hours in triplicate (DMSO, fedratinib alone, venetoclax

alone, or combination). All RNA isolation and sequencing was performed by the Gene

Expression Center at the University of Wisconsin-Madison. RNA was isolated with a

RNeasy kit (Qiagen) prior to a quality control check using an Agilent Bioanalyzer and

NanoDrop One spectrophotometer. mRNA libraries were prepared using the TruSeq

stranded mRNA kit and sequenced on an Illumina NovaSeq 6000 platform to 30 million

reads per sample. Sample reads were processed by the University of Wisconsin

Biotechnology Center.

<u>RNA-Seq analysis</u>

Bioinformatic analysis of transcriptomic data adhere to recommended ENCODE

guidelines and best practices for RNA-Seq. Alignment of adapter-trimmed (Skewer

v0.1.123; (11)) 2x150 (paired-end; PE) bp strand-specific Illumina reads to the *Homo*

*sapiens*GRCh38 genome (assembly accession NCBI:GCA_000001405.25) was

achieved with the Spliced Transcripts Alignment to a Reference (STAR v2.5.3a)

software (12), a splice-junction aware aligner, using annotation provided by Ensembl

release 104. Expression estimation was performed with RSEM v1.3.0 (RNASeq by

Expectation Maximization; (13)). To test for differential gene expression among

individual group contrasts, expected read counts obtained from RSEM were used as

input into edgeR (v3.16.5; (14)). Inter-sample normalization was achieved with the

trimmed mean of M-values (TMM; (15)) method. Statistical significance of the negative-

binomial regression test was adjusted with a Benjamini-Hochberg FDR correction at the

5% level (16). Prior to statistical analysis with edgeR, independent filtering was applied

and required genes to have a count-per-million (CPM) above k in n samples, where k is

determined by minimum read count (10 reads) and by the sample library sizes

where n is determined by the number of biological replicates in each group. The validity

of the Benjamini-Hochberg FDR multiple testing procedure was evaluated by inspection

of the uncorrected p-value distribution.

Gene Set Enrichment Analysis (GSEA) software MSigDB_2023.1 was used to test for

enriched pathways in the combination group compared to the other three experimental

groups. A total of 1000 permutations were chosen using the Ensembl_Human_Gene_ID

platform and tested against the Hallmark, KEGG and Reactome data sets, with relevant

significantly enriched pathways shown (17,18). The GSEA and MSigDB are available

under a Creative Commons style license and is a joint project of UC San Diego and the

Broad Institute.

Apoptosis and cell cycle assays

RS4;11 cells were treated for 24 hours (DMSO, fedratinib alone, venetoclax alone, or

combination) prior to both analyses. BrdU-7AAD and Annexin V-PI analyses were

performed according to manufacturer's protocols (BD Biosciences) and analyzed via an

Attune NxT Flow Cytometer (ThermoFisher, Waltham, MA) with analysis on FlowJo v10

(Ashland, OR).

BH3 Profiling Assay

BH3 profiling is a functional assay that measures apoptotic priming of cells of interest

after pre-treatment with a drug. The BH3 profiling assay measures the extent of

mitochondria outer membrane permeability that occurs in response to pro-apoptotic

BH3 peptides, which mimic the activity of pro-apoptotic BH3 only proteins from the BCL-

2 family methods. Methods have been previously described by others(19).

In vivo studies

NOD/SCID/$Il2rg^{tm1wjl}$/Szj (NSG) breeder mice were purchased from Jackson Labs (Bar

Harbor, ME) and bred internally at the Biomedical Research Model Services (BRMS)

Breeding Core at the University of Wisconsin-Madison in accordance with the Guide for

the Care and Use of Laboratory Mice and experiments were performed under an IACUC

approved animal protocol (M005915). Male and female NSG mice were used at 8-16

weeks of age and injected via tail vein with 2 x10^6 RS4;11 cells on Day 0. Peripheral

blood was obtained via retro-orbital bleeds conducted on Days 10 (engraftment), Day

21 (pre-treatment), and Day 35 (post-treatment). Mice were treated with fourteen days

of once daily treatment with vehicle, fedratinib, venetoclax, or combination starting at

Day 21 (5 days on, 2 days off). Venetoclax was prepared for in vivo use by dissolving

drug into the following: 5% DMSO, 30% PEG400, 7.5% Tween20, 7.5% Tween80 and

50% sterile water. Fedratinib was prepared for in vitro use by dissolving drug into 100%

sterile water. Venetoclax was administered via intraperitoneal (IP) injections at a dose of

6.25 mg/kg. Fedratinib was administered via oral gavage at a dose of 100 mg/kg. Mice

were monitored for survival and percent weight change and euthanized when a humane

endpoint was reached. Peripheral blood was treated with ACK lysis buffer (Lonza) and

then stained in flow buffer (PBS, 10% FBS) with CD19 (HIB19), CD20 (2H7), CD22

(HIB22), PD-L1 (MIH2), FAS (DX2), HLA-DR (L243), BCL2 (100) and BAX (2D2), all

purchased from Biolegend, prior to quantification on an Attune NxT Flow Cytometer with

analysis on FlowJo V10.

Statistical analysis

Statistics were performed using GraphPad Prism version 9.0 for the Macintosh OS

(GraphPad Software). Data were expressed as mean ± SEM. For analysis of three or

more groups, a non-parametric one-way ANOVA test was performed with the Bonferroni

or Sidak's multiple comparisons post-test. Analysis of differences between two normally

distributed test groups was performed using a two-sided Mann Whitney test. A p-value

less than 0.05 was considered statistically significant.

Data availability statement

By the time of publication all RNA seq data will be reposited in the Gene Expression
Omnibus (GEO). All other data will be made available upon reasonable request.

Results:

Combinatorial treatment of fedratinib and venetoclax effectively kills human B-ALL in
vitro

To assess the efficacy of JAK inhibitors to kill human B-ALL cells, we tested ruxolitinib
(JAK1/2 inhibitor) and fedratinib (semi-selective JAK2 inhibitor) on the human B-ALL cell
line RS4;11. Cells were treated with a range of doses from 1nM to 10uM for 48 hours
prior to analysis for cell survival (Figure 1A-B). Cells were also treated with single agent
venetoclax (Figure 1C). Based on the dose-curves, the calculated IC50 for fedratinib
was 600nM but the IC50 for ruxolitinib was not reached.

We next wanted to explore whether or not fedratinib would be more efficacious when
combined with venetoclax. To test this, we treated RS4;11 with single agent fedratinib at
700nm, single agent venetoclax at 7.5nM and the combinatorial treatment (Figure 1D-
E). Combinatorial treatment was significantly more effective than either single agent
treatment and similar to the positive control staurosporine. We next tested if ruxolitinib
had a similar beneficial effect as fedratinib (Figure 1F-G). Ruxolitinib did not improve the
killing of RS4;11 over single-agent venetoclax (Figure 1F-G), even at higher doses. To
ensure that the combinatorial effect of fedratinib and venetoclax was not specific to
RS4;11, we tested both the single agent and combinatorial treatments on the human B-
ALL cell lines SUPB-15 and NALM-6. The combinatorial treatment was equally effective
on SUPB-15 (Figure 2A-B) but surprisingly, the combinatorial treatment did not improve
the killing of NALM-6 cells compared to venetoclax alone (Figure 2C-D). These data
suggest that while the combinatorial treatment of fedratinib and venetoclax can be
highly effective in human B-ALL, there are a subset of B-ALLs that remain resistant to
this combinatorial treatment.

Resistance to fedratinib treatment is correlated with absence of Flt3 expression

To determine the precise activity of and contribution to cell killing by fedratinib and

venetoclax, we performed several additional dose-escalation studies on all three human

B-ALL cell lines. Keeping a constant venetoclax dose of 5 nM, both RS4;11 and SUPB-

15 experienced significant cell death that was further enhanced with increasing doses of

fedratinib. Conversely, NALM-6 was mostly refractive to venetoclax at 5 nM with

increasing concentrations of fedratinib modestly improving the killing efficiency (Figure

3A-C, left panels). Using a constant dose of fedratinib of 525 nM, increasing doses of

venetoclax was able to significantly increase cell death in both RS4;11 and SUP-B15

while NALM-6 remains refractory as before (Figure 3 A-C, right panels).

To assess a potential mechanism for NALM-6 resistance to both venetoclax and

fedratinib treatment, we measured the expression of downstream targets BCL-2, BRD4

and FLT3 in each human B-ALL cell line. Notably, NALM-6 had no expression of FLT3,

which is also a target of fedratinib while RS4;11, which has the greatest sensitivity to

fedratinib, had the highest expression of Flt3 (Figure 3D-E). We also observed that

NALM-6 was more resistant to venetoclax treatment than either RS4;11 or SUPB-15

(Figure 1, 2). Interestingly, NALM-6 had a significantly lower ratio of the pro-apoptotic

protein BAX to the anti-apoptotic protein BCL2 relative to RS4;11, which correlated with

venetoclax sensitivity (Figure 3F).

Combination therapy differentially enriches pathways related to cell death

Because there are many potential downstream targets impacted by both fedratinib and

venetoclax, we wanted to identify the dominant pathways altered in RS4;11 post single-

agent and combination treatment via bulk RNA sequencing. RS4;11 cells were plated

similarly as described above and exposed for 24 hours to fedratinib (525 nM),

venetoclax (5 nM), or combination and compared to untreated cells. Cells were

harvested after 24 hours for RNAseq. Whole transcriptome analysis revealed that each

treatment regimen induced a unique gene expression profile (Figure 4A-B). Comparing

each treatment regimen to untreated cells, fedratinib had the fewest number of

significantly different genes while the combinatorial treatment had the most (Figure 4C-

E). Interestingly in all cases, most of the differentially expressed genes were down-

regulated and not up-regulated as a result of treatment (Figure 4C-E). We also

completed a gene set enrichment analysis (GSEA) that identified pathways associated

with apoptosis, DNA repair and proliferation predominantly altered in the combinatorial

treatment versus all three other conditions (Figure 4F). A heatmap of the differentially

enriched genes primarily impacting the enrichment into the apoptotic pathway is also

shown (Figure 4G). Functional assays of both apoptosis and cell cycle were consistent

with RNA-seq analyses. A higher fraction of cell death was observed in combination

treatment in RS4;11 cells compared to single agent fedratinib or venetoclax (Figure 5A).

There also was evidence of G2-M arrest in combination treatment compared to single

agent treatment (Figure 5B). Finally, BH3 profiling suggests venetoclax differentially

contributes to BIM-mediated apoptosis (Figure 5C-G).

Combination therapy reduces leukemic burden and improves overall survival in RS4;11

derived xenograft model of B-ALL

Based on the efficacy of fedratinib and venetoclax observed *in vitro*, a xenograft model

of B-ALL was established utilizing RS4;11 cells and NSG mice to test these drugs as a

combination therapy *in vivo*. Our group has previously published data in T-ALL using

similar combination therapy, however, unlike in T-ALL we did not see evidence of

leukemia related CNS disease with the RS4:11 B-ALL model (Suppl Figure 1). NSG

mice were injected with luciferase[+] RS4;11 cells on Day 0 and then randomized to one

of the following groups: DMSO, single agent venetoclax, single agent fedratinib, or

combination. Peripheral blood was obtained via retroorbital bleeds on days 10

(engraftment), 21 (pretreatment), and 35 (post treatment) to assess B-ALL disease

burden (Figure 6A). Treatment started on day 21 and lasted for 14 days (5 days on, 2

days off) with 6.25 mg/kg venetoclax IP and/or 100 mg/kg fedratinib via gavage.

Combination treatment prolonged the survival of the mice and decreased the RS4;11

burden the most compared to the other treatment groups (Figure 6B). Venetoclax alone

also significantly improved overall survival when compared to fedratinib alone and

untreated mice (Figure 6B). Peripheral blood counts and IVIS imaging throughout the

experiment revealed that combination treatment was highly effective at decreasing the

RS4;11 burden in vivo (Figure 6C-D). Flow cytometric analysis of RS4;11 taken from

the peripheral blood also showed a significant increase in MFI of FAS and CD19 in the

combination group (Figure 6E). Overall, these data support the role of fedratinib and

venetoclax as a combination therapy against FLT3[+] B-ALL, with potential modulation of

cell surface marker expression of CD19.

Discussion:

Treatment of relapsed/refractory B-ALL remains a challenge, particularly in patients who

do not respond to traditional chemotherapy or immunotherapy. Targeting BCL-2 using

venetoclax may be a viable option in subsets of B-ALL like KMT2A-rearranged and

hypodiploid B-ALL. In previous work, a single dose of 12.5 mg/kg venetoclax in a

RS4;11-derived xenograft model led to a maximal tumor growth inhibition of 47% and

tumor growth delay of 26%. However, maximal effects were seen when combining

venetoclax with chemotherapy (20). Other studies demonstrating efficacy of venetoclax

in B-ALL include combining with MCL-1 inhibitors (21, 22), monoclonal antibodies such

as daratumumab in relapsed/refractory B-ALL (23), AKT inhibitors (24, 25), MDM2

inhibitors (26), and also with BCR-ABL inhibitors (27). No prior studies have looked at

the combination of JAK2 inhibitors with venetoclax in B-ALL. The dysregulation

of JAK/STAT related genes has been previously shown to be found in high-risk patient

samples of B-ALL (6). One group identified rearrangements of JAK2 in Ph like B-ALL

involving the activation of mutations in IL7R and FLT3, and deletion of SH2B3, which

encodes the JAK2 negative regulator LNK (28).

The goal of this study was to suppress leukemia proliferation related to JAK/STAT

dysregulation while at the same time inhibiting BCL-2 to prevent escape from apoptosis

in models of human B-ALL. To do this, we tested fedratinib, a semi selective JAK2

inhibitor, and venetoclax, a potent BCL2 inhibitor, to assess the efficacy of this

combination in 3 models of human B-ALL in vitro and with the RS4;11 cell line in vivo.

Fedratinib is FDA approved for patients with myeloproliferative disorders, primarily

myelofibrosis (29, 30) and has been used to salvage patients that have failed ruxolitinib

(31). Furthermore, fedratinib has been used to decrease tumor burden in Hodgkin

lymphoma and mediastinal large B-cell lymphoma in vitro and in vivo (32). However,

fedratinib has not previously been used for B-ALL. Analysis of BRD4 and FLT3 was

included for all 3 cell lines in our study as fedratinib has been shown to target both

BRD4 (33, 34) and FLT3 (33) in addition to its activity against JAK2.

Our data both in vitro and in vivo demonstrated that the combination of fedratinib and

venetoclax more effectively treated B-ALL than single agent treatment. We also

examined the use of another JAK2 inhibitor, ruxolitinib, which we had previously shown

synergized with venetoclax to treat T cell ALL (10). Similar to observations by others

(21), we found that ruxolitinib alone was ineffective in doses up to 10 uM. In contrast,

combination fedratinib and venetoclax did demonstrate efficacy in the Ph+ B-ALL cell

line SUP-B15, suggesting that inhibiting the JAK/STAT and BCL-2 pathways

simultaneously is particularly effective in Ph-like and FLT3+ high risk subsets compared

to more standard B-ALL models of disease such as NALM-6. Fedratinib also inhibits

FLT3, which is commonly overexpressed in KMT2A-rearranged B-ALL (35) and was

found to be highly expressed in RS4;11. The association of FLT3 and BCL2 in B-ALL is

unclear but GSEA suggests changes primarily in proliferation, apoptosis and DNA repair

pathways with combination fedratinib and venetoclax treatment. Functional assays of

both apoptosis and cell cycle were consistent with these RNA-Seq analyses. A higher

fraction of cell death was observed in combination treatment compared to single agent

fedratinib or venetoclax. Furthermore, there also was evidence of G2-M arrest in

combination treatment compared to single agent treatment. BH3 profiling suggests

venetoclax differentially contributed to BIM-mediated apoptosis. The mitochondrial outer

membrane priming of the combination therapy appears to reflect that of venetoclax

alone. The addition of fedratinib, while not increasing mitochondrial sensitivity to the

BIM BH3 peptide, does synergize with venetoclax to increase cell death (Figures 1, 5A,

6B) but appears to do so via a different mechanism. Further testing utilizing a FLT3

and/or BIM knock in or knock out lines could help determine the role of these molecules

in response to the combination therapy.

Finally, we demonstrated efficacy of combination fedratinib and venetoclax therapy in a

B-ALL xenograft model. In fact, with only a 2-week treatment we were able to identify

reduced leukemic burden and an increase in overall survival benefit with combination

therapy. Our group attempted to reduce potential toxicity in our in vivo model by

reducing the DMSO concentration in our venetoclax drug preparation for in vivo use and

gavaging with sucrose-soaked gavage tips to reduce stress on mice as much as

possible, using techniques described by others (36). Our group also identified an

increase in CD19 expression in peripheral blood samples obtained from mice that

received either venetoclax or combination therapy. One group recently demonstrated

pre-sensitizing B cell malignancies, including B-ALL, through venetoclax exposure

increases cytotoxic efficacy of CD19 directed CAR-T therapy (37). These findings along

with our data with combination venetoclax plus fedratinib suggest these targeted

therapies may serve as a potential bridge therapy to sensitize B-ALL to CD19 targeted

treatments such as blinatumomab or CAR T cells.

The development of novel treatment strategies for relapsed/refractory B-ALL is essential

for the subset of patients with poor outcomes due to failure to respond to initial

conventional chemotherapy. The combination of fedratinib and venetoclax is an

effective treatment strategy to reduce the burden of disease, particularly in FLT3$^+$ B-

ALL, through reduction in proliferation and DNA repair pathways and induction of

apoptosis. Combination fedratinib and venetoclax may also potentially serve as a bridge

to CD19-directed therapies such as CAR T cell and future studies should explore if

these therapies can help B-ALL avoid potential CD19 antigen escape.

Acknowledgements:

We would like to thank the University of Wisconsin Carbone Cancer Center (UWCCC)

Flow Cytometry and Small Animal Imaging and Radiotherapy core facilities, who are

supported in part through NCI/NIH P30 CA014520 as well as the UW Biotechnology

Center.

Author contributions:

SPR, NJH and CMC designed the experiments, analyzed and interpreted results and

wrote the manuscript; SPR, NJH, HCB, LS, NMH, LR, DPT, TN and NJH conducted

experiments and analyzed data; SPR, NJH, DPT, TN and HCB generated figures; LR,

JLL and CMC supervised experiments. All authors read and approved the manuscript.

Funding:

This work was supported by grants from the NIH TL1 TR002375 (S.P.R.), NIH/NCATS

UL1-TR002373, the Cormac Pediatric Leukemia Fellowship and the Stem Cell and

Regenerative Medicine Center Fellowship (N.J.H.), NCI/NIH R01 CA187299 (L.R.), St.

Baldrick's Empowering Pediatric Immunotherapy for Childhood Cancers Grant, the

MACC Fund (C.M.C) and the University of Chicago Collaborative Grant (J.L.L). The

contents of this article do not necessarily reflect the views or policies of the Department

of Health and Human Services, nor does mention of trade names, commercial products,

or organizations imply endorsement by the US Government. None of these funding

sources had any input in the study design, analysis, manuscript preparation or decision

to submit for publication.

References

References

1. Hunger SP, Lu X, Devidas M, et al. Improved survival for children and adolescents

with acute lymphoblastic leukemia between 1990 and 2005: a report from the Children's

Oncology Group. *JCO.* 2012;30(14):1663-1669. doi:10.1200/JCO.2011.37.8018

2. Bhojwani D, Pui CH. Relapsed childhood acute lymphoblastic leukaemia. *The Lancet*

Oncology. 2013;14(6):e205-e217. doi:10.1016/S1470-2045(12)70580-6

3. Leahy AB, Elgarten CW, Grupp SA, Maude SL, Teachey DT. Tisagenlecleucel for the

treatment of B-cell acute lymphoblastic leukemia. *Expert Review of Anticancer Therapy*.

2018;18(10):959-971. doi:10.1080/14737140.2018.1512411

4. Loh CY, Arya A, Naema AF, Wong WF, Sethi G, Looi CY. Signal transducer and

activator of transcription (STAT) proteins in cancer and inflammation: functions and

therapeutic implication. *Front Oncol*. 2019;9:48. doi:10.3389/fonc.2019.00048

5. Bousoik E, Montazeri Aliabadi H. "Do we know jack" about JAK? A closer look at

JAK/STAT signaling pathway. *Front Oncol*. 2018;8:287. doi:10.3389/fonc.2018.00287

6. Reshmi SC, Harvey RC, Roberts KG, et al. Targetable kinase gene fusions in high-

risk B-ALL: a study from the Children's Oncology Group. *Blood*. 2017;129(25):3352-

3361. doi:10.1182/blood-2016-12-758979

7. Shamas-Din A, Kale J, Leber B, Andrews DW. Mechanisms of action of BCL-2 family

proteins. *Cold Spring Harbor Perspectives in Biology*. 2013;5(4):a008714-a008714.

doi:10.1101/cshperspect.a008714

8. Diaz-Flores E, Comeaux EQ, Kim KL, et al. Bcl-2 is a therapeutic target for

hypodiploid B-lineage acute lymphoblastic leukemia. *Cancer Research*.

2019;79(9):2339-2351. doi:10.1158/0008-5472.CAN-18-0236

9. Karjalainen R, Pemovska T, Popa M, et al. JAK1/2 and BCL2 inhibitors synergize to

counteract bone marrow stromal cell–induced protection of AML. *Blood*.

2017;130(6):789-802. doi:10.1182/blood-2016-02-699363

10. Walker KL, Rinella SP, Hess NJ, et al. CXCR4 allows T cell acute lymphoblastic

leukemia to escape from JAK1/2 and BCL2 inhibition through CNS infiltration. *Leukemia*

& Lymphoma. 2021;62(5):1167-1177. doi:10.1080/10428194.2021.1910684

11. Jiang H, Lei R, Ding SW, Zhu S. Skewer: a fast and accurate adapter trimmer for

next-generation sequencing paired-end reads. *BMC Bioinformatics*. 2014;15(1):182.

doi:10.1186/1471-2105-15-182

12. Dobin A, Davis CA, Schlesinger F, et al. STAR: ultrafast universal RNA-seq

aligner. *Bioinformatics*. 2013;29(1):15-21. doi:10.1093/bioinformatics/bts635

13. Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with

or without a reference genome. *BMC Bioinformatics*. 2011;12(1):323. doi:10.1186/1471-

2105-12-323

14. Robinson MD, McCarthy DJ, Smyth GK. Edger□: a bioconductor package for

differential expression analysis of digital gene expression data. *Bioinformatics*.

2010;26(1):139-140. doi:10.1093/bioinformatics/btp616

15. Robinson MD, Oshlack A. A scaling normalization method for differential expression

analysis of RNA-seq data. *Genome Biol*. 2010;11(3):R25. doi:10.1186/gb-2010-11-3-r25

16.Reiner A, Yekutieli D, Benjamini Y. Identifying differentially expressed genes using

false discovery rate controlling procedures. *Bioinformatics*. 2003;19(3):368-375.

doi:10.1093/bioinformatics/btf877

17. Mootha, Vamsi K., Cecilia M. Lindgren, Karl-Fredrik Eriksson, Aravind

Subramanian, Smita Sihag, Joseph Lehar, Pere Puigserver, et al. "PGC-1α-Responsive

Genes Involved in Oxidative Phosphorylation Are Coordinately Downregulated in

Human Diabetes." *Nature Genetics* 34, no. 3 (July 2003): 267–73.

https://doi.org/10.1038/ng1180.

18. Subramanian, Aravind, Pablo Tamayo, Vamsi K. Mootha, Sayan Mukherjee,

Benjamin L. Ebert, Michael A. Gillette, Amanda Paulovich, et al. "Gene Set Enrichment

Analysis: A Knowledge-Based Approach for Interpreting Genome-Wide Expression

Profiles." *Proceedings of the National Academy of Sciences* 102, no. 43 (October 25,

2005): 15545–50. https://doi.org/10.1073/pnas.0506580102.

19. Fraser, Cameron, Jeremy Ryan, and Kristopher Sarosiek. "BH3 Profiling: A

Functional Assay to Measure Apoptotic Priming and Dependencies." In *BCL-2 Family*

Proteins, edited by Evripidis Gavathiotis, 1877:61–76. New York, NY: Springer New

York, 2019. https://doi.org/10.1007/978-1-4939-8861-7_4.

20. Souers AJ, Leverson JD, Boghaert ER, et al. ABT-199, a potent and selective BCL-

2 inhibitor, achieves antitumor activity while sparing platelets. *Nat Med*. 2013;19(2):202-

208. doi:10.1038/nm.3048

21. Moujalled DM, Hanna DT, Hediyeh-zadeh S, et al. Cotargeting BCL-2 and MCL-1 in

high-risk B-ALL. *Blood Advances*. 2020;4(12):2762-2767.

doi:10.1182/bloodadvances.2019001416

22. Mayumi A, Tomii T, Kanayama T, et al. The combination of ruxolitinib and Bcl-2/Mcl-

1 inhibitors has a synergistic effect on leukemic cells carrying a SPAG9::JAK2

fusion. *Cancer Gene Ther*. 2022;29(12):1930-1938. doi:10.1038/s41417-022-00511-z

23. Voruz S, Blum S, De Leval L, Schoumans J, Solly F, Spertini O. Daratumumab and

venetoclax in combination with chemotherapy provide sustained molecular remission in

relapsed/refractory CD19, CD20, and CD22 negative acute B lymphoblastic leukemia

with KMT2A-AFF1 transcript. *Biomark Res*. 2021;9(1):92. doi:10.1186/s40364-021-

00343-3

24. Richter A, Fischer E, Holz C, et al. Combined application of pan-AKT inhibitor MK-

2206 and BCL-2 antagonist venetoclax in B-cell precursor acute lymphoblastic

leukemia. *IJMS*. 2021;22(5):2771. doi:10.3390/ijms22052771

25. Holz C, Lange S, Sekora A, et al. Combined BCL-2 and PI3K/AKT pathway

inhibition in KMT2A-rearranged acute B-lymphoblastic leukemia cells. *IJMS*.

2023;24(2):1359. doi:10.3390/ijms24021359

26. Hohtari H, Kankainen M, Adnan-Awad S, et al. Targeting apoptosis pathways with

BCL2 and MDM2 inhibitors in adult B-cell acute lymphoblastic leukemia. *HemaSphere*.

2022;6(3):e701. doi:10.1097/HS9.0000000000000701

27. Massimino M, Tirrò E, Stella S, et al. Targeting BCL-2 as a therapeutic strategy for

primary p210 BCR-ABL1-positive B-ALL cells. *In Vivo*. 2020;34(2):511-516.

doi:10.21873/invivo.11802

28. Roberts KG, Morin RD, Zhang J, et al. Genetic alterations activating kinase and

cytokine receptor signaling in high-risk acute lymphoblastic leukemia. *Cancer Cell*.

2012;22(2):153-166. doi:10.1016/j.ccr.2012.06.005

29. Pardanani A, Harrison C, Cortes JE, et al. Safety and efficacy of fedratinib in

patients with primary or secondary myelofibrosis: a randomized clinical trial. *JAMA*

Oncol. 2015;1(5):643. doi:10.1001/jamaoncol.2015.1590

30. Harrison CN, Schaap N, Vannucchi AM, et al. Safety and efficacy of fedratinib, a

selective oral inhibitor of Janus kinase-2 (JAK2), in patients with myelofibrosis and low

pretreatment platelet counts. *Br J Haematol*. 2022;198(2):317-327.

doi:10.1111/bjh.18207

31. Harrison CN, Schaap N, Mesa RA. Management of myelofibrosis after ruxolitinib

failure. *Ann Hematol*. 2020;99(6):1177-1191. doi:10.1007/s00277-020-04002-9

32. Hao Y, Chapuy B, Monti S, Sun HH, Rodig SJ, Shipp MA. Selective JAK2 inhibition

specifically decreases Hodgkin lymphoma and mediastinal large B-cell lymphoma

growth in vitro and in vivo. *Clinical Cancer Research*. 2014;20(10):2674-2683.

doi:10.1158/1078-0432.CCR-13-3007

33. Talpaz M, Kiladjian JJ. Fedratinib, a newly approved treatment for patients with

myeloproliferative neoplasm-associated myelofibrosis. *Leukemia*. 2021;35(1):1-17.

doi:10.1038/s41375-020-0954-2

34. Ciceri P, Müller S, O'Mahony A, et al. Dual kinase-bromodomain inhibitors for

rationally designed polypharmacology. *Nat Chem Biol*. 2014;10(4):305-312.

doi:10.1038/nchembio.1471

35. Armstrong SA, Kung AL, Mabon ME, et al. Inhibition of FLT3 in MLL. *Cancer Cell*.

2003;3(2):173-183. doi:10.1016/S1535-6108(03)00003-5

36. Hoggatt AF, Hoggatt J, Honerlaw M, Pelus LM. A spoonful of sugar helps the

medicine go down: a novel technique to improve oral gavage in mice. *J Am Assoc Lab*

Anim Sci. 2010;49(3):329-334

37. Yang M, Wang L, Ni M, et al. Pre-sensitization of malignant B cells through

venetoclax significantly improves the cytotoxic efficacy of CD19 CAR-T cells. *Front*

Immunol. 2020;11:608167. doi:10.3389/fimmu.2020.608167

Figure Legends

Figure 1. Fedratinib, but not ruxolitinib, combines with venetoclax to improve

killing of human B-ALL

The human B-ALL cell line RS4;11 was treated with a dose curve of single agent

ruxolitinib (A), fedratinib (B) or venetoclax (C) and number of live cells quantified by flow

cytometry after 48hrs. (D-G) RS4;11 cells were treated with the indicated concentration

of fedratinib and venetoclax (D-E) or ruxolitinib and venetoclax (F-G) for 48 or 72hrs.

Live cells were quantified by flow cytometry and proliferation rate measured by

colorimetric MTS assay. Fed = fedratinib. Rux = ruxolitinib. Ven = venetoclax.

* = p < 0.05, ** = p < 0.01, **** = p < 0.0001

Figure 2. Combinatorial treatment of venetoclax and fedratinib is efficacious in

the human B-ALL cell line SUP-B15, but not NALM-6

Human B-ALL cell lines NALM-6 and SUP-B15 were treated with venetoclax in

combination with fedratinib. Live cells were quantified by flow cytometry as previously

described (A,C) and proliferation was assessed by an MTS assay as previously

described (B,D). ns = not significant, * = p < 0.05, ** = p < 0.01, *** = p < 0.001, **** = p

< 0.0001

Figure 3. Fedratinib activity is associated with Flt3 expression

Combination dose curves were performed on RS4;11 (A), SUP-B15 (B) and NALM-6

(C), keeping either the venetoclax dose (left panel) or fedratinib dose (right panel)

constant to determine the potency of the combinatorial treatment on different human B-

ALL cell lines. (D-F) Expression of both Flt3 and Brd4 (D-E) and the ratio of BAX to

BCL2 (F) were measured in each cell line by flow cytometry. Fold MFI represents the

median fluorescent intensity of the marker divided by the isotype control. *** p < 0.001.

Figure 4. Combination venetoclax and fedratinib treatment induces a unique gene

expression profile. RS4;11 cells were left untreated or treated with 5 nM ventoclax,

525 nM fedratinib or the combination treatment for 24 hrs prior to RNA isolation and

bulk sequencing. (A-B) Transcriptome analysis of each treatment group by two-

dimensional multidimensional scaling (MDS) (A) and a clustered image map reveals

each treatment group induces a unique transcriptomic profile. (C-E) Volcano plots

showing the number of positively and negatively regulated genes in each treatment

group relative to the untreated control. (F) Gene set enrichment analysis of the

combination treatment against all other treatments was performed with three gene sets

shown. (G) Heatmap showing the genes contributing to the enrichment score of the apoptosis gene set.

Figure 5. Apoptosis, cell cycle and BH3 profiling is dysregulated after combinatorial treatment with venetoclax and fedratinib. Flow cytometric assay of apoptosis (A) and cell cycle (B) after treatment with (need doses and target cell line). BH3 profiling assay examining BIM depolarization after exposure to (C) DMSO control, (D) 7.5 nM venetoclax, (E) 700 nM fedratinib and (F) combination. (G) AUC analysis of depolarization BH3 profiling curves in RS4;11 B-ALL.

Figure 6. Fedratinib and venetoclax combination treatment is superior to single-agent treatment in vivo. (A) RS4;11 was transplanted into non-irradiated NSG mice via tail vein injection at 2E6 cells per mouse and allowed to engraft for 21 days prior to starting treatment. On days 21-35, Venetoclax was given IP at a dose of 6.25 mg/kg while fedratinib was given gavage at a dose of 100 mg/kg. Retro-orbital blood draws were taken at day 10, 21 and 35 to assess cancer burden. (B) Kaplan-Meier survival curve of each treatment. (C-D) Representative dot plot of RS4;11 taken from the blood at day 35 (C) and quantification across all times (D). Representative IVIS images (E) and total flux quantification of all mice at the indicated time points (F). (G) A select number of B-ALL cancer markers expression profiles were assessed at day 35 across all treatment conditions. Fold MFI represents the median fluorescent intensity of the marker divided by the isotype control. Data is representative of two independent experiments. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Supplementary Figure 1. FISH of bone marrow and spinal cord samples from untreated Day 51 RS4;11 transplanted NSG mice show very limited CNS infiltration during late stage leukemic disease. Bone marrow and spinal cord tissue were harvested on untreated Day 51 transplanted RS4;11 NSG mice. Human and mouse CEP probes were used to quantify human RS4;11 cells versus mouse cells via FISH to determine disease burden in bone marrow and spinal cord tissues.

Figure 4

Figure 6

Figure 5

Figure 3

Figure 2

Figure 1