zol & fai: large-scale targeted detection and evolutionary investigation of gene clusters

Rauf Salamzade[1,2], Patricia Q. Tran[3,4], Cody Martin[2,3], Abigail L. Manson[5], Michael S.
Gilmore[5,6,7], Ashlee M. Earl[5], Karthik Anantharaman[3], Lindsay R. Kalan[1,8,9]

[1]Department of Medical Microbiology and Immunology, School of Medicine and Public Health, University of
Wisconsin-Madison, Madison, WI, USA
[2]Microbiology Doctoral Training Program, University of Wisconsin-Madison, Madison, WI, USA
[3]Department of Bacteriology, University of Wisconsin-Madison, Madison, WI, USA
[4]Freshwater and Marine Science Doctoral Program, University of Wisconsin-Madison, WI, USA
[5]Infectious Disease and Microbiome Program, Broad Institute of MIT and Harvard, Cambridge, Massachusetts, USA
[6]Department of Ophthalmology, Harvard Medical School and Mass Eye and Ear, Boston, Massachusetts, USA
[7]Department of Microbiology, Harvard Medical School and Mass Eye and Ear, Boston, Massachusetts, USA
[8]Department of Medicine, Division of Infectious Disease, School of Medicine and Public Health, University of
Wisconsin-Madison, Madison, WI, USA
[9]M.G. DeGroote Institute for Infectious Disease Research, David Braley Centre for Antibiotic Discovery, Department
of Biochemistry and Biomedical Sciences, McMaster University, Hamilton, Ontario, Canada

Address for Correspondence: Lindsay R. Kalan; kalanlr@mcmaster.ca

Abstract

Many universally and conditionally important genes are genomically aggregated within
clusters. Here, we introduce fai and zol, which together enable large-scale comparative analysis
of different types of gene clusters and mobile-genetic elements (MGEs), such as biosynthetic
gene clusters (BGCs) or viruses. Fundamentally, they overcome a current bottleneck to reliably
perform comprehensive orthology inference at large scale across broad taxonomic contexts and
thousands of genomes. First, fai allows the identification of orthologous instances of a query
gene cluster of interest amongst a database of target genomes. Subsequently, zol enables
reliable, context-specific inference of ortholog groups for individual protein-encoding genes
across gene cluster instances. In addition, zol performs functional annotation and computes a
variety of evolutionary statistics for each inferred ortholog group. Importantly, in comparison to
tools for visual exploration of homologous relationships between gene clusters, zol can scale to
thousands of gene cluster instances and produce detailed reports that are easy to digest. To
showcase fai and zol, we apply them for: (i) longitudinal tracking of a virus in metagenomes, (ii)
discovering novel population-level genetic insights of two common BGCs in the fungal species
Aspergillus flavus, and (iii) uncovering large-scale evolutionary trends of a virulence-associated
gene cluster across thousands of genomes from a diverse bacterial genus.

Background

De novo ortholog grouping typically involves searching for reciprocal best hits of proteins
between pairs of genomes, indicative of orthology, and subsequently clustering pairs of inferred
orthologs and in-paralogs across multiple genomes[1–4]. Initial methods for orthology inference
were designed to be able to identify orthologs between distinct species but limited in the number
of genomes they could process[1–3]. This limitation is largely due to the all-vs-all alignment of

proteomes, core to most methods for *de novo* ortholog grouping, which is an $O(n^2)$ operation
and a major computational bottleneck. Approaches to overcome this procedure include limiting
proteome comparisons by using a guiding-phylogeny[5,6], adapting alignment searching
parameters and heuristics to further boost speeds[7,8], or preliminary aggressive clustering of
proteins into coarse homolog groups[9]. Recently, graph-based and iterative-clustering
approaches have also allowed vast scalability to thousands of bacterial genomes, but are
primarily designed for application to a single species[10–13].
Available orthology inference methods struggle to infer ortholog groups across large
datasets of taxonomically diverse genomes, potentially representing thousands of species, such
as a set of metagenome-assembled genomes (MAGs) related to a common microbiome. While
multiple methods exist to identify instances of previously established ortholog groups within the
predicted proteome of a metagenome[14–17], these are unable to account for proteins not
represented in their database. Recently, independent advancements in methods to collapse
large protein sets based on sequence similarity have enabled rapid clustering of millions of
sequences[18–20]. These approaches have even been used on massive protein datasets gathered
from across multiple metagenomic datasets[21]; however, more resolute delineation of functionally
analogous ortholog groups across thousands of genomes from multiple species remains difficult
to perform *de novo*.
Of relevance, within bacterial genomes, genes are often co-located within smaller,
discrete, multi-gene units, which we will broadly refer to as gene clusters. Examples of gene
clusters include operons[22,23], phages[24], metabolic gene clusters[25], biosynthetic gene clusters
(BGCs)[26–29], and pathogenicity islands[30,31]. Although less common, eukaryotic genomes can
also contain genes aggregated within discrete clusters[32–34]. Sometimes gene clusters are highly
conserved, encoding for products essential to the survival of the organism[35]. In other cases, a
single gene cluster can exhibit variability in gene carriage and order across different strains or
species[36–38]. This is often the case for BGCs encoding specialized metabolites or virulence-
associated gene clusters, where evolution of gene content and sequence divergence can
influence fitness and contribute to adaptation within a changing ecosystem[39–41].
Syntenic conservation has been used to assist *de novo* identification of homologous
instances of a gene cluster of interest in diverse target genomes[42–45]. Homologous gene cluster
instances can then be comprehensively investigated to delineate homolog or ortholog groups of
the proteins found across them[44,46]. While such targeted approaches can alleviate time and
computational resources by avoiding more comprehensive identification of orthologs at genome-
wide scales, currently available methods are mostly designed for specific types of gene clusters,
such as BGCs[42,44,45]. Many of the software implementing such approaches also do not provide
support for uniform annotation of coding sequences in target genomes, which can decrease
sensitivity for gene cluster detection. In addition, most methods do not account for gene cluster
paralogy, which has been observed for BGCs in bacterial[38] and fungal genomes[33], or provide
specialized capabilities for finding gene clusters across fragmented genomes or metagenomic
assemblies[38].
Following identification of homologous gene clusters in target genomes, software to
understand the evolutionary relationships between gene cluster instances and infer protein
ortholog groups have largely applied coarse protein clustering and aimed to provide
visualization based exploration to users[44,46–48]. Visual assessment of related gene clusters and

manual refinement of ortholog groups work well at smaller scales but become impractical when
dealing with hundreds to thousands of gene cluster instances. Scalability challenges are due to
both computational costs needed to render visuals as well as the figures becoming convoluted
and difficult to interpret. An effective solution to ease the identification of evolutionary trends
amongst homologous gene clusters is to first identify ortholog groups[44] and present information
pertaining to their conservation and sequence divergence within tabular reports[10,38]. Such
tabular reports scale by the number of unique ortholog groups and can be organized by their
consensus order along gene cluster instances. We recently introduced construction of such
reports in a software suite for exploring microdiversity amongst homologous BGCs from a single
taxon[38]; however, the functionality was difficult to use outside of the suite and reliant on
orthologous relationships between proteins of gene clusters being known in advance.
Here, we introduce the zol suite, providing functionalities for gene cluster detection and
subsequent inference and investigation of protein ortholog groups across homologous gene
clusters. The versatility and scalability of these programs is demonstrated through application to
three types of gene clusters within different genomic contexts including a virus within
environmental metagenomes, fungal secondary metabolite encoding biosynthetic gene clusters,
and a conserved polysaccharide antigen locus from the diverse bacterial genus of
Enterococcus.
109
110   **Results**
111
112   **fai and zol allow for the rapid inference of gene cluster orthologs across diverse**
113   **genomes**
114
The zol suite consists of three major programs: prepTG (*prepare target genomes*), fai
(*find additional instances*), and zol (*zoom on locus*) (**Figure 1A**). First, prepTG and fai can be
run to process a set of target genomes and rapidly search for a query gene cluster within them,
respectively. Afterwards, zol can perform reliable and efficient context-limited inference of
ortholog groups across homologous gene cluster instances identified using a flexible
InParanoid-type algorithm[3]. For each ortholog group, zol will further compute evolutionary
statistics, such as Tajima's D[49], and functional annotations, using several, diverse databases
suitable for a variety of gene clusters, including those specific to phages[50], virulence elements[51],
and BGCs[52]. Ultimately, zol will summarize data in a table report where each row corresponds
to a distinct ortholog group. This report is automatically color formatted and provided as an
XLSX spreadsheet to allow for easy interpretation of the data, which can span thousands of
gene cluster instances.
To promote consistency in gene calling across target genomes, we have incorporated
computationally light-weight dependencies for *de novo* gene prediction in bacterial genomes[53,54]
and protein-mapping in eukaryotic genomes[55] within prepTG, to prepare and format target
genomes for optimized gene cluster searching in fai (**Figure 1B**). prepTG also aims to provide a
convenient interface to transform genomic or metagenomic datasets into a format ready for
searching using fai. Options are available to download pre-built databases of distinct
representative genomes for 18 commonly studied bacterial taxa[56] or to build comprehensive

databases for any genus or species in the latest release of the Genome Taxonomy Database
(GTDB)[57].
fai features two key features which are absent in most existing methods for gene cluster
detection (**Figure 1C; Table S1; Supplementary Text**). First, it has an option to automatically
filter secondary instances of query gene clusters identified in target genomes, removing
potentially paralogous gene clusters from downstream investigations. Second, fai implements a
mode for searching for gene clusters in draft quality genomes, MAGs, or unbinned
metagenomic assemblies, where gene clusters might be fragmented across multiple scaffolds.
When this mode is activated, fai relaxes requirements for reporting a gene cluster as present in
a genome or metagenome if multiple homologous gene cluster regions are identified near
scaffold edges in a target genome and instead assesses whether reporting criteria are met in
unison across such instances (**Figure S1**). Similar to prepTG, fai also aims to provide
convenience for users and can accept query gene clusters in different formats to ease
searching for gene clusters and genomic islands cataloged in databases such as ICEberg[58],
MIBiG[52], or IslandViewer[59]. Query gene clusters can be provided as a coordinate along a
reference genome, in GenBank format, or as a set of proteins in FASTA format. In addition, to
simplify conservation and novelty assessment of a single isolate's BGCs, phages, and plasmids
relative to other genomes from the same genus or species, specialized wrapper programs of fai
are also provided within the zol suite (**Figure S2**).
zol will infer ortholog groups for proteins across homologous gene clusters and then
construct a tabular report with information on conservation, evolutionary trends, and annotation
for each individual ortholog group (**Figure 1D**). To make annotated reports generated by zol
more comprehensive for different types of gene clusters, several databases have been
included, such as VOGs[50], VFDB[51], ISFinder[60], and CARD[61]. In addition, zol incorporates
HyPhy[62] as a dependency and calculates various evolutionary statistics. Ultimately, beyond
high-throughput inference of ortholog groups across diverse genomic datasets, the rich tabular
report produced by zol provides complementary information to figures generated by
comparative visualization software such as clinker[46], CORASON[44], gggenomes[63], and Easyfig[64].
A key feature in zol is the ability to dereplicate gene clusters directly using skani[65], which
was recently shown to be more reliable at estimating average nucleotide identity (ANI) between
genomes of variable contiguity relative to comparative methods. Dereplication can allow for
more appropriate inference of evolutionary statistics to overcome availability or sampling biases
in genomic databases[66]. It can also be used to subset distinct representative gene cluster
instances to make investigation using visualization software more tractable. Another important
ability of zol is a mode where users can provide a handful of known instances for a gene cluster
to estimate optimal parameters to search for additional instances of the gene cluster using fai.
We applied this functionality of zol on sets of homologous BGCs and phages to determine
distributions for search parameters in fai which users could consult as priors (**Figure S3;**
Supplementary Text).
Finally, zol allows for comparative investigations of gene clusters based on taxonomic or
ecological groupings[67–69]. For instance, users can designate a subset of gene clusters as
belonging to a specific population to allow zol to calculate ortholog group conservation across
just the focal set of gene clusters. In addition, zol will compute the fixation index[70], F_{ST}, for each
ortholog group to assess gene flow between the focal and complementary sets of gene clusters.

178
Longitudinal tracking of a virus within lake metagenomic assemblies
180

Metagenomic datasets represent a large reservoir of underexplored sequence
space[71,72]. To demonstrate the ability of the zol suite to identify and investigate gene clusters in
metagenomes, we applied it to track a virus in a longitudinal metagenomic dataset profiling a
lake's microbiome over space and time[73].
We first identified large ($\geq$20kb) viruses, that were also predicted to represent circular
molecules, across a subset of the metagenomic assemblies corresponding to the earliest
sampling date[74]. Afterwards, clustering based on the sequence and syntenic similarity of protein
domains led to the identification of a ~36kb highly conserved virus in two of the metagenomes
sampled from lower lake depths.
All 16 metagenomic assemblies, spanning five distinct sampling timepoints and four
distinct sampling depths, were processed through prepTG to identify coding sequences and
construct a database ready to search for gene clusters using fai. GenBank files with coding
sequence annotations for metagenomic assemblies generated by prepTG, amassing 27 Gb
total in size, were further provided as input for cblaster makedb, which serves a similar role to
prepTG in the cblaster suite to format genomic data for downstream gene cluster searches.
However, cblaster makedb does not feature the ability to perform *de novo* gene-calling for either
genomes or metagenomes and is not designed to accommodate the size of metagenomic
assemblies. During database construction, cblaster makedb required around 30 Gb of memory,
while prepTG needed less than 3 Gb of memory (**Figure S4A**).
Next, fai was used to perform a rapid, targeted search for this ~36 kb *Caudovirales* virus
across the full set of 16 metagenomes to identify additional instances of the virus. fai completed
its search of the metagenomes, featuring >20 million proteins and 10.7 million contigs, in less
than four minutes using 20 threads, performing similarly to cblaster, run using similar settings as
fai (**Figure S4B**). Of the 16 total metagenomes, the virus was found in ten metagenomes,
including all nine metagenomes surveying anoxic conditions (p<0.001; one-sided Fisher's exact
test; **Figure 2A**). This is concordant with inferences for the host for the virus being *Rhodoferax*,
which are purple bacterium featuring species classified as anaerobic photoheterotrophs[73,75,76]. In
addition, *Rhodoferax* classified MAGs from the metagenomic dataset were exclusively obtained
from anoxic conditions[73]. To investigate how the gene repertoire of the virus evolved over time,
we next applied zol. zol-based analysis revealed that 45 (72.6%) of the 62 total distinct ortholog
groups were core to all instances of the virus across ten metagenomes with most completely
conserved in sequence over the course of 2.5 months (**Figure 2B; Table S2**). Furthermore, 15
of the 62 ortholog groups were not observed in the query viruses from the earliest sampling
date, suggesting the potential acquisition or duplication of genes in the virus during the span of
sampling at the lake.
216

Investigating population-level and species-wide evolutionary trends of BGCs in the eukaryotic species *Aspergillus flavus*
219

Low sensitivity for gene cluster detection in eukaryotic genome assemblies can arise
from their incompleteness, leading to gene clusters being fragmented across multiple

scaffolds[77,78], as well as challenges in *ab initio* gene prediction due to alternative splicing[79,80].
Therefore, many gene cluster detection software are either specific for bacterial genomes or
require coding sequence annotations for eukaryotic genomes to be provided by the user. To
overcome such challenges to user application, we integrated miniprot[55] into prepTG which
allows for mapping high-quality protein annotations from a reference genome to the remainder
of the genomes available for a species or genus. We showcase the ability of prepTG and fai to
simplify the reliable identification of gene clusters in eukaryotic genomes by using them to find
instances of two BGCs across genomes belonging to the fungal species *Aspergillus flavus*.
The genus of *Aspergillus* is a source of several natural products, including aflatoxins, a
common and economically impactful contaminant of food[81]. The genus also contains species
that are model organisms for studying fungal secondary metabolism[34,82,83]. Examination of the
secondary metabolome of *A. flavus* has revealed that different clades or populations can exhibit
variability in their metabolite production despite high conservation of core BGC genes encoding
enzymes for synthesis of these metabolites[37,84]. For instance, population B *A. flavus* were
identified as producing a greater abundance of the insecticide leporin B relative to populations A
and C[37,85]. We showcase zol's ability to aid comparative analysis of gene clusters from different
populations through application to the leporin BGC. We further show how zol can detect
variation in sequence conservation for different genes from the aflatoxin BGC and be inclusive
of genes present in target genome annotations but missing in the query gene cluster, allowing
for comprehensive profiling of BGC auxiliary content.
Based on read alignment to a reference genome, the leporin cluster was recently
identified to be a core component of the *A. flavus* genome[37]. However, a restricting factor in the
direct prediction of gene clusters in *A. flavus* assemblies is the lack of gene annotations, with
only 11 (5.1%) of 216 genomes from the species in NCBI's GenBank database having coding
sequence predictions (**Figure 3A**). Therefore, we mapped high-quality protein predictions for a
reference *A. flavus* genome[86] to the remainder of the 216 genomes available for the species.
Running fai in "draft mode" led to the identification of the leporin BGC within 212 (98.1%)
assemblies, consistent with the prior read mapping-based investigation suggesting that the BGC
was core to the species[37]. In comparison, the CAGECAT server[87], which runs cblaster[45], was
limited to genomes with protein coding annotations available on NCBI and thus unable to
assess the remaining 205 genomes for the presence of the leporin BGC (**Figure 3B**). We also
investigated the ability of non-targeted approaches for BGC detection to identify the leporin
BGC by applying antiSMASH followed by BiG-SCAPE for clustering related BGCs and matching
them to characterized BGCs in the MIBiG database. When this approach was applied using
GenBank files prepared by prepTG, the gene cluster clan corresponding containing the leporin
BGC was found in all *A. flavus* genomes provided as input. However, when antiSMASH was run
using *de novo* gene prediction in antiSMASH based on GlimmerHMM[88] with *Cryptococcus* gene
annotation models, recovery of the leporin BGC was limited (**Figure 3B**).
Of the 212 genomes with the leporin BGC identified by fai, 202 contained instances that
were high-quality and not near scaffold edges. This set of 202 instances of the gene cluster was
further investigated using zol with options to perform comparative investigation of BGC
instances from *A. flavus* population B genomes to instances from other populations. High
sequence conservation was observed for all genes in the leporin gene cluster as previously
reported[37] (**Table S3**). Further, alleles for genes in the BGC from population B genomes were

generally more similar to each other than to alleles from outside the population, as indicated by
high F_{ST} values (>0.85 for 9 of 10 genes) (**Figure 3C; Table S3**). While regulation of secondary
metabolites in *Aspergillus* is complex[89], zol analysis showed that the three essential genes for
leporin production[85] also had the lowest variation in the 100 bps upstream their exonic
coordinates (**Figure S5**). This suggests higher variability is occurring in the transcription of the
accessory *lep* genes within the species. This supports experimental evidence that has shown
gene knockouts depleting certain leporin species will still permit the production of others[85].
fai and zol were also applied to the BGC encoding aflatoxin across *A. flavus*[90] (Table
S4). Similar to the leporin BGC, the aflatoxin BGC was highly prevalent in the species and found
in 71.8% of genomes. However, in contrast to the leporin BGC, the aflatoxin BGC contained
several genes with positive Tajima's D values, indicating greater sequence variability for these
coding regions across the species (**Figure 3D**). One of the genes with a positive Tajima's D
value was *aflX*, which has been shown to influence conversion of the precursor veriscolorin A to
downstream intermediates in the aflatoxin biosynthesis pathway[91] (**Figure 3E**). An abundance
of sites with mid-frequency alleles in the oxidoreductase encoding gene could represent
granular control for the amount of aflatoxin relative to intermediates produced. The polyketide
synthase gene *pksA* had the lowest Tajima's D value of -2.4, which suggests it is either highly
conserved or under purifying selection (**Figure 3F**). In addition, because the reference proteome
used to infer genomic coding regions was constructed recently[86], fai and zol detected several
highly conserved genes within the aflatoxin BGC that are not represented in the original
reference gene cluster input for fai[52]. This includes a gene annotated as a noranthrone
monooxygenase and recently characterized as contributing to aflatoxin biosynthesis[92,93] (**Figure
3D**).
289
290   **Identification of the Enterococcal polysaccharide antigen and assessment of context**
291   **restricted orthology inference**
292
To demonstrate the ability of zol and fai to reliably identify ortholog groups across
multiple species and thousands of genomes, we used the tools to assess the distribution of the
enterococcal polysaccharide antigen (Epa) and its individual genes across the diverse genus of
Enterococcus. Because previous comparative genomic investigations have been performed
between *epa* loci from different species[94,95], we also showcase how such prior insight can be
used to tailor parameters in fai for searching for the locus across the full genus and how results
from fai can be assessed for appropriate selection of parameter values in zol.
The Epa is a signature component of the cellular envelope of multiple species within
Enterococcus[94–97] and has mostly been characterized in the species *Enterococcus faecalis*[96,98–
101]. While molecular studies have provided evidence that the locus contributes to enterococcal
host colonization[100], evasion of immune systems[102], and sensitivity to antibiotics[103] and
phages[103,104], it was only recently that the structure of Epa was resolved and a model for its
biosynthesis and localization formally proposed[101]. A homologous instance of the *epa* locus was
identified in the other prominent pathogenic species from the genus, Enterococcus
faecium[94,95,105]; however, the prevalence and conservation of *epa* across the diverse genus of
Enterococcus[106–108] remains poorly studied.

We first assessed the performance of fai and zol to identify *epa* loci across
representative genomes for each of the 92 species of *Enterococcus* in GTDB R214[57] and
subsequently delineate protein ortholog groups relative to other methods. Specifically, we
compared the runtime and ortholog group predictions of fai and zol to the combination of
cblaster and clinker as well as OrthoFinder, an established software for multi-species ortholog
group delineation, run on full genomes. For this comparison, the parameter settings for fai and
cblaster as well as zol and clinker were adapted to match each other more closely, with an
exception being to run fai in draft-mode, which lacks an analogous feature in cblaster. The
combination of fai and zol was the fastest of the three methods tested and able to identify
ortholog groups for the *epa* locus in approximately one minute (**Figure 4A, S6**). Orthology
inferences from fai and zol exhibited high overlap with orthology predictions by the alternate two
methods, finding 96.3% of ortholog protein pairs identified by at least two of the three methods
(**Figure 4B**). We also applied all three methods to determine *epa* locus orthologs across low
quality representative genomes for each species to demonstrate the convenience of fai's ability
to be run in "draft mode" and improve sensitivity for detecting fragmented gene clusters in
comparison to cblaster. fai identified 2.1-fold more exclusive ortholog pairs in common with
OrthoFinder, expected to be relatively robust to the effects of assembly fragmentation, than the
number of ortholog pairs shared exclusively by cblaster and clinker with OrthoFinder (**Figure
4C**). In addition, we performed evolutionary-simulation of the *epa* locus, allowing for sequence
gains and losses, and assessed context-limited orthology inference by zol, clinker and
OrthoFinder (**Figure S7; Supplementary Text**). zol was able to recover a high fraction of true
positive ortholog relations and was the best method at avoiding prediction of false positive
orthologs.
Next, to properly and comprehensively assess the distribution of *epa* across the entire
set of 5,291 genomes in GTDB classified as one of the 92 *Enterococcus* species[57], we applied
fai with more careful consideration of parameter values and requested more advanced features
for gene cluster detection. A sensitive searching criterium was selected based on prior
comparative genomics for the locus[94,95] and its coordinates along the *E. faecalis* V583 genome
as a reference[99,101]. For detection of *epa* orthologous regions, co-location of at least seven of
the 14 *epa* genes previously identified as conserved in both *E. faecalis* and *E. faecium* was
required. The default threshold for syntenic conservation of homologous instances to the query
gene cluster was disregarded to increase sensitivity for the detection of *epa* in enterococcal
species more distantly related to *E. faecalis*. In addition, key proteins were specified and the
length of the flanking context to include as part of the loci was expanded. Using these criteria,
5,085 of the genomes assessed were found to possess an *epa* locus, with phylogenomic
investigations further revealing that the locus is highly conserved in three of the four major
clades of *Enterococcus* (**Figure 4D; Table S5**).
Based on fai's reports, we realized that to achieve optimal clustering for ortholog groups
across the diverse set of *epa* loci identified, we needed to lower the default thresholds for
percent identity and coverage that protein pairs needed to exhibit for being considered as
orthologs (**Figure 4D; Table S5**). We ran zol on both the full set of 5,052 high-quality *epa* loci
and only loci from species representative genomes. For the comprehensive analysis, zol was
able to identify 14 ortholog groups as core or near-core, found in >90% of loci instances (**Table
S6**). When provided 30 threads, zol completed in 30.7 hours and had a maximum memory

usage of 101.3 GB. The more restricted analysis of zol to investigate *epa* instances from 65 species representative genomes was to allow for assessing the quality of ortholog group predictions using phylogenetics (**Table S7**). After applying zol on *epa* from species representative genomes, orthology predictions were assessed through construction of a maximum-likelihood phylogeny of *epa* associated glycosyltransferases. Ortholog groups which corresponded to glycosyltransferases from *E. faecalis* V583 were labelled on the phylogeny and confirmed to match distinct phylogenetic clades, which suggests their appropriate delineation (**Figure 4EF**). zol further identified several *epa* associated glycosyltransferase ortholog groups that were absent in the *E. faecalis* representative genome and other representative genomes from the *E. faecalis* clade (**Figure 4G**). These distinct glycosyltransferases might impact the final structure or decoration of Epa in other *Enterococcus* species.

zol identifies genetic diversity of *epa*X-like glycosyltransferases in *E. faecalis*

zol features several options related to the dereplication of input gene clusters to retain only distinct representative instances for orthology inference and other downstream analytics (**Figure S8**). Importantly, the application of these methods can substantially reduce zol's runtime and impact some of the evolutionary statistics computed (**Figure S8, S9, S10, Supplementary Text**). Whether dereplication is appropriate for a particular analysis should thus be carefully considered by users depending on their research aims. In particular, dereplication can impact investigations for highly sequenced bacterial taxa, including the opportunistic pathogen *E. faecalis*. For such pathogens, certain lineages, such as those commonly isolated at clinics, might be overrepresented in genomic databases, and the researcher may find it beneficial for the analysis to apply dereplication.

To showcase the scalability of zol and its ability to expand knowledge for even well-studied gene clusters, we applied it to high-quality, complete *epa* loci from 1,232 *E. faecalis* genomes without dereplication. In accordance with prior studies[94,101], zol was able to distinguish core and strain-variable patterns. The report from zol showed that one end of the locus corresponds to genes which are highly conserved and core to *E. faecalis* (*epa*A-*epa*R), whereas the other end contained strain-specific genes (**Figure 5A; Table S8**). Using zol, we further found that variably conserved genes exhibit high sequence dissimilarity, as measured using both Tajima's D and average sequence entropy, in comparison to the core genes of the locus (**Figure 5BC**). These statistics were robust to the application of dereplication and thus unlikely to be heavily impacted by well-sequenced lineages (**Figure S9, S10**).

One ortholog group, corresponding to the glycosyltransferase *epaX*, exhibited substantially higher sequence variation than other *epa* associated glycosyltransferases (**Figure 5BD**). This finding was further validated through phylogenetic analysis of glycosyltransferases from the species, which highlighted the breadth of diversity observed for the *epaX* ortholog group relative to other *epa* associated glycosyltransferases (**Figure 5E**).

Discussion

Here fai and zol are introduced to enable large-scale evolutionary investigations of gene clusters in diverse taxa. Together these tools overcome current bottlenecks in computational

biology to infer orthologous sets of genes at scale across thousands of diverse genomes and
large metagenomic assemblies.
The set of input gene clusters for zol does not need to be produced by fai. cblaster[45] is
another tool that can identify instances of a query gene cluster within a set of target genomes
and extract them in GenBank format for downstream investigations using zol. For those lacking
computational resources needed for fai analysis, cblaster offers remote searching of BGCs
using NCBI's BLAST infrastructure and non-redundant databases. More recently, CAGECAT[87],
a highly accessible web-application for running cblaster, was also developed and can similarly
be used to identify and extract gene cluster instances from genomes represented in NCBI
databases. In contrast to these tools, prepTG and fai feature algorithms and options for users
interested in: (i) identification of gene clusters in metagenomes, (ii) performing standardized
gene annotation across target genomes, (iii) improved sensitivity for gene cluster detection in
draft-quality assemblies, and (iv) automated filtering of secondary, or paralogous, matches to
query gene clusters. In addition, users can apply zol to further investigate homologous sets of
gene clusters identified from IslandCompare[109], BiG-SCAPE[44], or vConTACT2[110] analyses,
which perform comprehensive clustering of predicted genomic islands, BGCs, or viruses.
The application of fai to identify gene clusters in metagenomes is demonstrated here
through rapid, targeted detection of a virus across lake metagenomic assemblies. We expect
that both fai and zol will gain greater relevance for metagenomic applications in the future as
long-read sequencing becomes cheaper. Importantly, the tools can be applied directly on
assemblies without the need for binning scaffolds into MAGs, avoiding complications associated
with binning[111]. In addition to their application to viral tracking, fai and zol's application to
metagenomes could be useful for assessing the presence of concerning transposons carrying
antimicrobial resistance traits[112–114] and identifying novel auxiliary genes within known BGCs
which may tailor the resulting specialized metabolites and expand chemical diversity[115,116].
Reidentifying gene clusters in eukaryotic genomes remains difficult due to technical
challenges in gene prediction owing to the presence of alternative splicing. The ability of fai and
zol to perform population-level genetics on BGCs from the eukaryotic species *A. flavus* was
demonstrated. While there are over 200 genomes of *A. flavus* in NCBI, only 5.1% have coding-
sequence information readily available. We used miniprot[55] to map high quality gene coordinate
predictions from a representative genome in the species[86] to the remainder of genomic
assemblies with prepTG which enabled high sensitivity detection of BGCs with fai. Our analysis
provides additional support that the leporin BGC is conserved across the species[37] using an
assembly-based approach.
The ability of zol to identify ortholog groups across 5,052 gene cluster instances from 71
distinct species using limited computational resources was demonstrated through investigation
of the *epa* locus across *Enterococcus*. While such large-scale investigations will be largely
limited to those with access to a server, we expect datasets to often feature some degree of
species level redundancy. For instance, 80.2% of the 5,052 *epa* instances were from only two
species, *E. faecalis* and *E. faecium*. Thus, to alleviate computational costs, we have included
functions for dereplication of gene clusters and reinflation of ortholog groups in zol. Applying
these features to the comprehensive set of *epa* loci using 30 threads, reduced runtime from
30.7 to 3.5 hours and maximum memory usage from 101.3 GB to 83.2 GB (**Table S9**).

We further assessed the quality of ortholog group predictions by fai and zol using
phylogenetic investigations and comparisons with other software for homology inference.
Specifically, we compared orthology inference results from fai and zol to predictions obtained
from the combination of cblaster and clinker as well as OrthoFinder[117], which was used to
detect ortholog groups at the genome-wide scale. Notably, clinker[46], which is developed by the
authors of cblaster, is primarily designed to produce interactive visualizations showing
relationships between related gene cluster instances. clinker's application of single-linkage
clustering to determine related sets of genes and to color matching genes in figures is expected
to produce relatively coarse ortholog groups. OrthoFinder was chosen as a representative
method for standard multi-species orthology inference because it has been shown to perform
well for several criteria in prior benchmarking studies[117,118]. Through application to identification
of ortholog groups for diverse *epa* loci from multiple distinct species and evolutionary simulation
of the locus from *E. faecalis*, we found zol produces reliable orthology predictions that are
mostly in accordance with alternate orthology inference methods while exhibiting restraint for
over clustering. In the future, we are considering further improving the algorithm for ortholog
group classifications within zol. Specifically, we might take a similar approach to OrthoFinder in
which coarse ortholog groups are first identified and later refined using phylogenetics.
Our investigation of *epa* loci from multiple species revealed the presence of a multitude
of glycosyltransferases associated with production or decoration of the polysaccharide,
including some that are absent in the representative *E. faecalis* genome, the species in which
the polysaccharide has been most extensively characterized. Through population-genetic
investigations of the locus in *E. faecalis* using zol, we further determined that an ortholog group
containing *epa*X-like glycosyltransferases possessed high sequence divergence relative to other
glycosyltransferases associated with the locus. In addition to influencing the ability of *E. faecalis*
to colonize hosts[100], mutations in *epa*X and other genes from the ortholog group have also been
shown to impact susceptibility to phage predation[119–122]. Therefore, we hypothesize that
extensive evolution of the *epa*X ortholog group is a result of contrasting selective forces,
pressuring *E. faecalis* to retain or (re-)acquire the glycosyltransferase to gain a fitness
advantage within hosts but also lose the gene to escape phage predation.
469
470 **Conclusions**
471
Practically, zol presents a comprehensive analysis tool for comparative genetics of
related gene clusters to facilitate detection of evolutionary patterns that might be less apparent
from visual analysis. Fundamentally, the algorithms presented within fai and zol enable the
reliable detection of orthologous gene clusters, and subsequently orthologous proteins, across
multi-species datasets spanning thousands of genomes and help overcome a key barrier in
scalability for comparative genomics.
478
479 **Methods**
480
481 **Software availability**
482

zol is provided as an open-source software suite, developed primarily in Python3 on GitHub at:
https://github.com/Kalan-Lab/zol. Docker and Bioconda[123] based installations of the suite are
supported. For the analyses presented in this manuscript, we used v1.4.1 of the zol software
package[124]. Version information for major dependencies of the zol suite[53,55,62,65,125–132] and other
software used[44,74,133] for analyses in this study is provided in Table S10. Code and input files for
generation of figures in this manuscript are provided separately on GitHub at:
https://github.com/Kalan-Lab/Salamzade_etal_zol.
490
491 **Availability of data and materials**
492
Genomes and metagenomes used to showcase the application of fai and zol are listed with
GenBank accession identifiers in Table S11. Total metagenomes and their associated
information from Lake Mendota microbiome samplings were originally described in Tran *et al.*
2023[73] and deposited in NCBI under BioProject PRJNA758276. Genomic assemblies available
for *A. flavus* in NCBI's GenBank database on Jan 31st, 2023 were downloaded in FASTA
format using ncbi-genome-download (https://github.com/kblin/ncbi-genome-download).
Genomic assemblies for *Enterococcus* that met quality and taxonomic criteria for belonging to
the genus or related genera (e.g. Enterococcus_A, Enterococcus_B, etc.) in GTDB[57] release
R207 were similarly downloaded from NCBI's GenBank database using ncbi-genome-download
in FASTA format.
503
Assessment of compute time, memory usage, and disk space: The UNIX *time* command was
applied to measure the runtime and memory usage of programs. Specifically, the "Elapsed (wall
clock) time" was regarded as the runtime and the "Maximum resident set size (kbytes)" as the
maximum memory usage. The UNIX *du* command was used to measure the final disk space
used by various programs. All analyses were computed on the same server running Ubuntu
18.04.06 LTS with AMD EPYC 7451 24-Core processors, 472 GB of 288-Pin DDR4 random-
access memory, and a Samsung 970 Pro solid disk drive.
511
512 **Overview of tools and algorithms**
513
prepTG - processing and preparing target genomes for searching with fai: prepTG allows users
to create a database of target genomes that can be searched for homologous instances of
query gene clusters with fai. In addition to formatting and producing files for optimizing fai
searches, prepTG integrates pyrodigal[53], prodigal[54], and miniprot[55] for gene-calling or protein-
mapping in prokaryotic and eukaryotic genomes as well as metagenomes to aid consistency in
fai's performance and limit bias due to potential differences in gene-calling methods. For
miniprot-based protein-mapping, coding sequence predictions are required to exhibit an identity
of at least 80% to the reference protein and instances of overlapping mRNA and exon features
are resolved by retaining only the highest scoring mappings.
prepTG also features options to download pre-built databases for select bacterial taxa
that are commonly studied[56], such as ESKAPE pathogens, or to download all genomes
belonging to any genus or species in GTDB R214[57] and subsequently construct a database *ab*
initio.

527
fai - automated identification of homologous instances of gene clusters: fai allows for rapid
detection of gene clusters in target genomes. It accepts a target genomes database prepared
by prepTG and query gene cluster(s). Query gene cluster(s) can be provided in one of three
formats: (i) GenBank file(s) with CDS features, (ii) a coordinate along a reference genome, or
(iii) a set of proteins. When using coordinates along a reference genome to define a gene
cluster, fai reperforms gene-calling along the reference using pyrodigal[53] and extracts a local
GenBank file corresponding to the specified region.
zol implements HMM-based and CDS separation-based approaches for determining
homologous gene cluster instances in target genomes, which can further be combined in a
hybrid approach. For both approaches, homologs of proteins from query gene clusters are first
searched for in predicted proteomes of target genomes using DIAMOND alignment[130]. Then, in
"Gene-Clumper" mode, which is the default, scaffolds with homologs of query proteins are
dynamically assessed for whether homologs are within a maximum number of CDS predictions
to be regarded as belonging to the same gene cluster. In "HMM" mode, scaffolds of target
genomes are instead scanned gene-by-gene using an HMM and neighborhoods or sets of
genes are regarded as being in a state of homology to the query gene cluster if several
individual genes depict homology to the proteins from the query gene cluster(s). The algorithm
is similar to *lsa*BGC-Expansion[38], however, it is not dependent on a preliminary genome-wide
orthology grouping analysis and thus features a different set of filters to still enable high-
throughput automated detection of homologous gene cluster segments as a result. *lsa*BGC-
Expansion is reliant on a preliminary orthology analysis to identify BGC-specific genes that
could be used to differentiate true homologous instances of BGCs and customize weighting of
HMM emission probabilities for distinct genes. It further requires the length of genes within
putative homologous regions to be within a certain deviation from the median length of known
gene instances. In contrast, fai has preconfigured emission probabilities which can be
customized by users and has no length requirement for potential homologous instances of
genes. fai further allows the "HMM-based" approach to be run with the parameter for
aggregating CDS predictions for the "Gene-Clumper" mode, whereby, gene cluster segments
detected by the HMM can be joined with other such segments if they are withing a certain
number of CDS features from each other. Similar to *lsa*BGC-Expansion, syntenic similarity
between candidate and query gene cluster segments can also be used to filter candidate
segments using a gene cluster-wide correlation metric[38].
By default, fai requires filters pertaining to the number of genes from query gene clusters
to be met for each homologous gene cluster candidate segment. However, in "draft mode",
thresholds for detection of gene clusters within target genomes are assessed in aggregate for
putative gene cluster segments found near scaffold edges (< 2,000 bp). Visual reports produced
by fai showcasing the sequence similarity of target genome proteins to the query protein(s) can
then be manually investigated by users to assess the validity of fragmented gene cluster
instances. In addition, fai features an option to filter for paralogous, overlapping candidate
segments of a gene cluster in target genomes and offers an intuitive visualization of gene
cluster segments, if requested, to allow users to assess their quality, including proximity of
candidate segments to scaffold edges. Together, these options enable the large-scale

identification of orthologous gene clusters across genomes which can then be leveraged by zol
to perform context-specific inference of protein ortholog groups.
In addition to a directory of homologous gene clusters in GenBank format, to serve as
input for zol analysis, and a small set of visual PDF files, fai generates an in-depth report on
which target genomes have the query gene cluster as an XLSX spreadsheet. This spreadsheet
includes information such as the average amino acid identity (AAI), syntenic similarity, and
number of conserved genes for gene clusters from target genomes relative to the query gene
cluster. The spreadsheet allows for easy sorting of various columns to assist identification of
which target genomes feature a gene cluster to the desired degree of similarity for the user.
579
zol - computes a variety of evolutionary statistics and can perform gene cluster specific
dereplication: The zol workflow begins by processing the input directory of gene cluster
GenBank files to assess validity and perform filtering of gene clusters or individual proteins.
Filtering can be performed at the gene cluster level by requesting filtering of draft-quality gene
clusters, those marked as being near scaffold edges, or low-quality gene clusters, those with
≥10% missing base-pairs (e.g. Ns) in their sequence. Filtering of individual proteins which are
near scaffold edges can also be performed if fai was used to identify the input gene cluster set,
because fai marks these proteins with a special feature tag in the resulting gene cluster
GenBank files.
Next, zol will perform dereplication of gene clusters, if requested by users, with skani[65]
by clustering gene clusters which depict some user-defined coverage and identity thresholds
using single linkage clustering or more resolved MCL-based clustering, for which the inflation
parameter can be adjusted. Representative gene clusters are selected from each cluster as part
of the dereplication based on maximum length and, if comparative analysis is requested,
whether the representative gene cluster is part of the focal or focal-complement set of gene
cluster instances specified by the user.
The input set of gene clusters or set of dereplicated representative gene clusters is then
used to identify protein ortholog groups with an InParanoid-type approach[3]. Briefly,
DIAMOND[130] is used to perform all vs. all pairwise alignment between proteins from the set of
gene clusters after which the alignments are processed to identify reciprocal best hits (RBH)
between pairs of gene clusters. In-paralogs are identified within each gene cluster based on
whether two coding sequences depict more similarity to each other than one does to an RBH
with a different gene cluster. Bitscores, standardized through division by reflexive bitscore
values for query proteins, are used to assess homology. Specifically, the average normalized
bitscore between each pair of orthologs and in-paralogs is recorded. Afterwards, bitscores
between such protein pairs are further standardized through dividing them with the average
values between pairs of gene clusters to aid proper clustering of proteins downstream. This is
akin to the genome-wide normalization procedure recommended in OrthoMCL, owing to the
realization that orthologs between distantly related species are also more likely to exhibit lower
sequence similarity, which should be corrected for prior to MCL clustering[2]. This information is
input into MCL with the inflation parameter set to 1.5, similar to other orthology inference
methods[7,117]. The inflation parameter and minimum identity and coverage cutoffs to consider
valid pairs of in-paralogs and orthologs are adjustable by users.

Reinflation can also be requested by users to expand ortholog groups to include proteins
from the full input set of gene clusters if gene cluster dereplication was requested[10]. Reinflation
of ortholog groups is performed by first performing comprehensive and granular clustering of
proteins from all input gene clusters using CD-HIT[128], requiring proteins to depict >98%
sequence similarity and > 95% bi-directional coverage to the representative sequences of
clusters. Proteins in CD-HIT clusters are then mapped to ortholog groups if they co-cluster with
proteins from dereplicated gene clusters which are already assigned to ortholog groups.
Dereplication and reinflation are not recommended if sequence redundancy amongst the set of
input gene clusters is low. Stringent cutoffs used for CD-HIT clustering during reinflation assume
that dereplication was also run with stringent parameters to only collapse highly similar gene
clusters. Otherwise, reinflation could miss more distant instances of ortholog groups, resulting in
an underestimation of ortholog group conservation amongst gene clusters.
Next, zol will partition protein and nucleotide sequences from gene clusters according to
ortholog groups, perform protein alignment using MUSCLE[132], and create codon alignments
using PAL2NAL[134]. We also offer an option to use reference proteins to refine and filter
sequences based on multiple sequence alignment using MUSCLE[132], which might be useful to
further filter intronic sequences in eukaryotic ORFs. Codon alignments are filtered for regions
with high ambiguity ($\geq$10% gaps) using trimAL[126] which are then used downstream for
calculation of evolutionary statistics and to construct approximate maximum-likelihood
phylogenies using FastTree 2[127] for each ortholog group. Consensus protein sequences for
each ortholog group are finally constructed using HMMER3[129].
Using protein consensus sequences of each ortholog group, zol is next able to linearize
annotation of ortholog groups with various annotation databases including KOfam[14], the PGAP
database[135], VFDB[51], CARD[61], MIBiG[52], ISfinder[60], the PaperBLAST database[136], and Pfam[137].
A custom FASTA file can also be provided by users to annotate ortholog groups. The best hit
per ortholog group for each annotation database is selected by score, if annotation is HMM
based[138], or bitscore, if it is DIAMOND alignment based[130], and a default E-value cutoff of 1e-5.
The E-value of the alignment is provided in the zol report for each putative annotation except
Pfam domains. However, for Pfam annotations, only domains meeting trusted thresholds are
reported.
Next, zol will compute basic statistics per ortholog group including the consensus order,
consensus directionality, whether proteins are single-copy across gene clusters, the median
length of ortholog group sequences, their median GC% percentage, and GC skew values. The
consensus order and directionality are performed similarly to *lsa*BGC-PopGene[38]. Afterwards, in
the sixth step, zol will calculate evolutionary statistics for each ortholog group including Tajima's
D[49], the proportion of filtered codon alignments which correspond to segregating sites, the
average sequence entropy of the filtered codon alignment and the 100 upstream region, and the
median and maximum Beta-RDgc. Beta-RDgc is a statistic that is derived from the Beta-RD
statistic which we described in *lsa*BGC[38] and measures the divergence of a pair of protein
sequences based on the expected divergence between the gene clusters. Values below one
suggest that protein divergence is larger for the pair than expected based on other shared
proteins between the two gene clusters; conversely, the opposite trend might suggest high
conservation of the particular protein between the gene clusters and potentially gene-specific
horizontal gene transfer. Finally, we perform site-specific selection analyses using the FUBAR[139]

and GARD[140] methods offered in the HyPhy suite. While highly scalable relative to comparable
methods[139], these analyses can still take considerable time and are turned off by default.
Importantly, GARD recombination detection[140] and partitioning of input alignments for ortholog
groups can also be used for alternate HyPhy analyses with HyPhy Vision[62], to extend beyond
the site-specific selection analyses using FUBAR[139] supported directly in zol.
Prior to the generation of a final report, zol allows users to perform an optional
comparative analysis between user-defined set(s) of focal and complementary or alternate gene
cluster instances. In these comparative analyses, the conservation and fixation index[70] is
calculated for each ortholog group.
Finally, we generate a consensus report and a spreadsheet in XLSX format where each
row corresponds to an ortholog group and columns correspond to basic statistics, evolutionary
statistics, and annotation information. Quantitative fields are automatically colored to make
visual detection of patterns easier for users. A basic heatmap showing the presence of ortholog
groups across gene clusters is also produced.
zol additionally features two alternate modes that can be triggered via specific
arguments. First, the "only-orthologs" argument will invoke zol to only compute ortholog groups
and exist after determining them. Second, the "select_fai_params_mode" argument allows
users to provide a handful of known instances for a gene cluster and determine appropriate
thresholds for searching for additional instances of the gene cluster using fai. This mode
assumes that the known instances provided are representative of the breadth of diversity
expected for the gene cluster amongst the target genomes being searched.
678
abon, atpoc, and apos – tools for assessing novelty and conservation of BGCs, phages, and
plasmids from a single strain: The zol suite features three small wrapper programs called abon,
atpoc, and apos which assess the conservation and novelty of a single genome's BGC-ome,
phage-ome, and plasmid-ome, respectively, relative to a target genome database constructed
by prepTG. The target genomes database could be all other genomes belonging to the focal
genome's species or genus. The three programs are wrappers of fai but also offer a simple
BLAST search alternative, to more thoroughly check for whether individual genes from BGCs,
phages, and plasmids are present in the target genomes being searched. These tools accept
results from standard software for annotation of BGCs[133,141], phages[74,142,143], and plasmids[143,144]
but do not integrate them within the suite. Similar to fai and zol they produce auto-formatted
XLSX spreadsheets as primary results.
690
691 **Application of fai and zol to track a virus within lake metagenomes**
692
693 VIBRANT was used to identify viral contigs or sub-contigs in the three total metagenomes from
694 Tran *et al.* 2023[73] sampled on the earliest date of 07/24. Afterwards, predicted circular contigs
695 were clustered using BiG-SCAPE[44] which revealed a ~36 kb virus was found in two of the three
696 metagenomes.
697
prepTG was run on all 16 total metagenomic assemblies from the Tran *et al.* 2023 study,
performing gene calling with pyrodigal in metagenomics mode[53] to prepare for comprehensive
targeted searching of the virus with fai. fai was run with largely default settings, with filtering of

secondary instances of the virus requested to retain only the best matching scaffold or scaffold
segment resembling the queries. In addition, the syntenic correlation requirement of hits to the
query gene clusters was turned off to account for the circular nature of the virus, which the
assessment is not designed for. To assess the performance of cblaster for preparing the target
metagenomes database and subsequently searching for the virus, we provided GenBank files
with CDS features produced by prepTG as input for cblaster makedb and adjusted searching
parameters for cblaster search to more closely match what we used for fai.
708
709 **Microevolutionary investigations of leporin and aflatoxin BGCs in *Aspergillus flavus***
710
Genomic assemblies downloaded from NCBI GenBank were processed using prepTG. Of the
217 genomic assemblies downloaded, one, GCA_000006275.3, was dropped from the analysis
because the original GenBank file had multiple CDS features with the same name, leading to
difficulties in performing BGC prediction with antiSMASH[133], and because alternate assemblies
were available for the isolate. prepTG was run on all assemblies with miniprot[55] based gene-
mapping of the high-quality gene coordinate predictions available for *A. flavus* NRRL 3357
(GCA_009017415.1)[86] requested. Target genomes were then searched for the leporin
(BGC0001445) and aflatoxin (BGC0000008) BGCs using GenBank files downloaded from
MIBiGv3[52] as queries. For leporin, AFLA_066840, as represented in the MIBiG database, was
treated as a key protein required for detection of the BGC. Similarly, for aflatoxin, PksA
(AAS90022.1), as represented in the MIBiG database, was treated as a key protein required for
detection of the BGC. Draft-mode and filtering of paralogous segments was requested. For both
analyses, ortholog groups found in fewer than 5% of gene cluster instances were disregarded.
724
725 We reidentified population B as previously delineated[37] using k-mer based ANI estimation[145] and
726 neighbor-joining tree construction[146]. A discrete clade (n=81) in the tree was validated to feature
727 all isolates previously determined as part of population B[37] and thus regarded as such.
728
For comprehensive and *de novo* BGC prediction, antiSMASH was run on the 216 genomic
assemblies with 'glimmerhmm' requested for the option '--genefinding-tool'. Similarly,
antiSMASH was also run on full GenBank files for genomes generated by prepTG from
reference proteome-mapping via miniprot. For one genome, antiSMASH was unable to process
the full GenBank created by prepTG due to an error related to "inconsistent exon ordering".
BGCs from each set of genome annotations were independently clustered using BiG-SCAPE
with "mix" clustering analysis and MIBiG reference BGC integration requested. The gene cluster
family and clan matching the reference leporin BGC in MIBiG (BGC0001445) were regarded as
the leporin BGC. For remote cblaster[45] analysis, CAGECAT[87] was used to search NCBI's nr
database with proteins from the leporin BGC representative (BGC0001445) provided as a
query. Only 13 scaffolds, belonging to 12 assemblies (including GCA_000006275.3), were
identified.
741
742 **Evolutionary investigations of the *epa* locus across *Enterococcus***
743

744   All *Enterococcus* genomes represented in GTDB R207[57] (n=5,291) were downloaded using
745   ncbi-genome-download[53]. The same query for *epa* was used for all analyses. Specifically,
746   coordinates extending from 2,071,671 to 2,115,174 along the *E. faecalis* V583 chromosome,
747   corresponding to genes EF2164 to EF2200, were used as a query for the *epa* locus in fai to
748   identify homologous instances in target genomes[99,101].
749
Comparing orthology/homology inferences between fai & zol, cblaster & clinker, and
OrthoFinder: Representative genome assemblies were selected for each of the 92 species of
Enterococcus in GTDB R214[57] based on the N50 metric. One set of species representative
genomes corresponded to those with the largest N50 values and the other set was comprised of
genomes with the lowest N50 values. The two sets of species representative genomes were
processed and investigated identically but independently. Gene calling was first performed for
genomes using prepTG with pyrodigal[53]. To generate the input for OrthoFinder, proteins from
prepTG's genome-wide GenBank files were extracted in FASTA format. After, OrthoFinder was
run with default settings. Phylogenetic hierarchical orthogroups inferred by OrthoFinder were
used for comparisons. To perform gene cluster specific homology prediction with cblaster and
clinker, we first used cblaster makedb to convert the genome-wide GenBank files from prepTG
into a database that could be searched with cblaster search. cblaster search was run using the
criteria: (i) DIAMOND alignment sensitivity mode set to very-sensitive, (ii) the percentage of
query genes required to be present in a cluster set to 25%, (iii) 1e-10 as the maximum E-value
for protein hits to be considered, (iv) 0% as the minimum coverage for protein hits to be
considered, (v) 0% as the minimum identity for protein hits to be considered, (vi) the maximum
flanking context for the gene cluster to gather set to 0 bp, (vii) request for intergenic proteins to
be included, and (viii) a maximum of 4620 bp allowed to separate protein hits for them to be
considered as part of the same gene cluster, which should approximately correspond to the
aggregate length of 5 bacterial genes on average[147]. Next, cblaster extract_clusters was used to
extract gene clusters found in target genomes by cblaster in GenBank format and provide them
as input for clinker. clinker was run using default settings but with only an output and matrix
output file requested to cut time needed to render an interactive figure, its primary intended
result file. To aid appropriate comparisons in orthology prediction, fai was largely run using
similar criteria as cblaster search: (i) DIAMOND alignment sensitivity mode set to very-sensitive,
(ii) the percentage of query genes required to be present in a cluster set to 25%, (iii) 1e-10 as
the maximum E-value for protein hits to be considered, (iv) the maximum flanking context for the
gene cluster to gather set to 0 bp, (v) a maximum of 5 proteins allowed to separate hits for
them to be considered as part of the same gene cluster, and (vi) syntenic similarity assessment
between target gene clusters and the query gene cluster turned off. However, draft-mode was
enabled in fai, which is not available in cblaster, to showcase the program's ability to improve
sensitivity for draft-quality assemblies. zol was applied with mostly default settings but with the
flags "only-orthologs", to stop after it determined ortholog groups, and "allow_edge_cds", to
allow usage of CDS features marked by fai to be near scaffold edges. All three methods were
provided 20 threads wherever possible.
785
Comprehensive and tailored usages of fai and zol for finding epa *in* Enterococcus*:* Based on
prior comparative analyses that had shown that gene conservation and gene order can be

slightly variable between *epa* loci from *E. faecalis* and *E. faecium*[94,95], we relaxed the syntenic
similarity requirement of candidate gene cluster matches in target genomes to the query in fai
from 0.6 to 0.0. In addition, we relaxed the minimum percentage of query proteins needed to
report a homologous instance of the *epa* locus to 10%. Instead, we required the presence of
50% of key *epa* proteins found in both *E. faecalis* and *E. faecium*, defined as
epaABCDEFGHLMOPQR, for the identification of valid homologous instances of the *epa* locus.
The E-value cutoff to determine presence for the key *epa* proteins was lowered from 1e-20 to
1e-10 to be inclusive of shorter genes and allow for higher levels of sequence divergence
across the *Enterococcus* genus. To gather auxiliary genes flanking the core *epa* region in target
genomes, we further requested the inclusion of CDS features found within 20 kb of the
boundary genes in detected instances of the *epa* locus within the resulting GenBank files
produced by fai. A phylogenetic heatmap was constructed for the presence of the *epa* locus
across a species tree using species representative genomes, selected based on largest
assembly N50, where the values of the heatmap corresponded to the maximum percent identity
of a query protein to their best match in target genomes. Because EF2173 and EF2185 are
identical transposases, they were shown as one column in the heatmap. The species tree was
constructed using GToTree[148] using HMMs for proteins regarded as largely single-copy core to
the phylum Bacillota. The phylogenetic heatmap visual was created using iTol[149].
From inspection of fai's resulting XLSX spreadsheet, zol's parameters were adjusted to
relax identity and coverage thresholds for assessing protein pairs for orthology prior to MCL
clustering to 20% and 25%, respectively. Identical processing was performed for the full set of
epa loci and *epa* loci from only species representative genomes. During the comprehensive
processing of all high-quality *epa* loci identified, one instance was dropped during zol analysis
despite meeting requirements because all CDS features in it were found near scaffold edges
and, by default, such features are not used in zol to aid more accurate inference of ortholog
groups and assessment of their sequence variation. A third run of zol was performed using
identical settings and all the gene cluster instances but leveraging the dereplication and
reinflation options to showcase how the combination of the options can reduce the runtime
needed for comprehensive processing. For dereplication of gene clusters, alignment fraction
was increased from the default of 95% to 99% and MCL was used for clustering to gather more
resolute representative gene clusters. Major ortholog groups determined between the
comprehensive and the dereplication + reinflation runs were found to be similarly conserved
based on matching to known *epa* genes.
821
Phylogenetic assessment of glycosyltransferase orthology predictions: Proteins from ortholog
groups determined by zol analysis of species representative genomes were extracted based on
whether the ortholog group was annotated as featuring the keywords: "glycosyl" and
"transferase" in Pfam protein domain annotations[150]. Two additional ortholog groups were
included and featured the Pfam domain "Bacterial sugar transferase", including *epaR*, which is
also regarded as a glycosyltransferase[101]. The comprehensive set of glycosyltransferases were
next aligned using MUSCLE with the default align mode[132]. Filtering of the alignment was next
performed using trimal with options "-keepseqs -gt 0.9" to filter sites composed largely of gaps
and further filtered for sequences which were composed of >10% gaps or ambiguous
characters ("X"). IQ-TREE[151] was used to construct a maximum-likelihood phylogeny with

ModelFinder limited to the WAG and LG substitution models. The phylogeny was visualized
using iTol[149] with classifications for ortholog groups most closely matching *E. faecalis* V583 *epa*
glycosyltransferases marked on leaves. Ortholog groups were assigned to specific *epa* gene
designations based on sequence alignment of their consensus sequences to *E. faecalis* V583
epa-associated proteins. Best matching ortholog groups for each *E. faecalis* V583 *epa*
glycosyltransferase were identified based on E-value.

Large-scale evolutionary investigations of *epa* loci from *E. faecalis*

The full set of *epa* loci identified by fai in *E. faecalis* genomes were processed through zol
requesting for retention of only complete instances that were also distant from scaffold edges.
For projection of conservation, Tajima's D, and sequence entropy statistics onto genes for the
epa locus in *E. faecalis* V583, sequence alignment was used to identify the best matching
ortholog groups based on E-value. For the identical transposases, EF2173 and EF2185, data
from a common ortholog group was used for both.

Investigation of glycosyltransferase phylogenetic diversity: A similar phylogeny of
glycosyltransferases was constructed for the *E. faecalis* analysis as was done for the
investigation of *epa* glycosyltransferases across species representatives of *Enterococcus*.
Glycosyltransferase ortholog groups were identified based on Pfam domains featuring the
keywords "glycosyl transferase" or because they matched *epa* genes regarded as
glycosyltransferases in prior studies[101]. To accommodate for the larger number of sequences: (i)
only ortholog groups found in >1% of *epa* loci instances were regarded, (ii) MUSCLE[132] super5
mode was used for alignment, and (iii) FastTree 2[127] was used for approximate maximum-
likelihood phylogeny construction. After trimal based filtering of sites, only sequences which
featured greater than 20% gaps or ambiguous characters ("X") were filtered to retain *epa*A in
the final alignment prior to phylogeny construction.

Abbreviations

Biosynthetic gene cluster (BGC), mobile-genetic element (MGE), Enterococcal polysaccharide
antigen (Epa), coding sequence (CDS), average nucleotide identity (ANI), metagenome-
assembled genome (MAG).

Declarations

Ethics approval and consent to participate: Not applicable

Consent for publication: Not applicable

Availability of data and materials: All genomic and metagenomic datasets used for showcasing
the application of fai and zol are publicly available on NCBI with accessions provided in
Supplementary Table S11.

Competing interests: The authors declare that they have no competing interests.

Funding: This work was supported by grants from the National Institutes of Health awarded to
L.R.K (NIAID U19AI142720 and NIGMS R35GM137828) and the Broad Institute
(U19AI110818). The content is solely the responsibility of the authors and does not necessarily
represent the official views of the National Institutes of Health.

Authors' contributions: RS & LRK designed the bioinformatics toolkit. RS, KA, & LRK conceived
of showcase applications. RS developed the software and performed all analyses. PQT
performed assembly of metagenomes. RS & LRK performed the initial drafting of the
manuscript. RS, PQT, CM, ALM, MSG, AME, KA, and LRK performed interpretation of results
and revision of the manuscript. All authors read and approved the final manuscript.

Acknowledgments: The authors are grateful to James Kosmopoulos, Dr. Caitlin Pepperell, Dr.
Caitlin Sande, and Dr. Mary Hannah Swaney for feedback and assistance with data acquisition
as well as Dr. Devon Ryan and Dr. Robert A. Petit III for assistance with incorporation of the
suite into Bioconda.

893
References

1.   Enright, A. J., Kunin, V. & Ouzounis, C. A. Protein families and TRIBES in genome

sequence space. *Nucleic Acids Res.* **31**, 4632–4638 (2003).

2.   Li, L., Stoeckert, C. J., Jr & Roos, D. S. OrthoMCL: identification of ortholog groups for

eukaryotic genomes. *Genome Res.* **13**, 2178–2189 (2003).

3.   Remm, M., Storm, C. E. & Sonnhammer, E. L. Automatic clustering of orthologs and in-

paralogs from pairwise species comparisons. *J. Mol. Biol.* **314**, 1041–1052 (2001).

4.   van Dongen, S. M. Graph clustering by flow simulation. (2000).

5.   Schreiber, F. & Sonnhammer, E. L. L. Hieranoid: hierarchical orthology inference. *J. Mol.*

Biol. **425**, 2072–2081 (2013).

6.   Georgescu, C. H. *et al.* SynerClust: a highly scalable, synteny-aware orthologue clustering

tool. *Microb Genom* **4**, (2018).

7.   Hu, X. & Friedberg, I. SwiftOrtho: A fast, memory-efficient, multiple genome orthology

classifier. *Gigascience* **8**, (2019).

8.   Cosentino, S. & Iwasaki, W. SonicParanoid: fast, accurate and easy orthology inference.

Bioinformatics **35**, 149–151 (2019).

9.   Ding, W., Baumdicker, F. & Neher, R. A. panX: pan-genome analysis and exploration.

Nucleic Acids Res. **46**, e5 (2018).

10. Page, A. J. *et al.* Roary: rapid large-scale prokaryote pan genome analysis. *Bioinformatics*

31, 3691–3693 (2015).

11. Bayliss, S. C., Thorpe, H. A., Coyle, N. M., Sheppard, S. K. & Feil, E. J. PIRATE: A fast and

scalable pangenomics toolbox for clustering diverged orthologues in bacteria. *bioRxiv*

(2019) doi:10.1101/598391.

12. Tonkin-Hill, G. *et al.* Producing polished prokaryotic pangenomes with the Panaroo

pipeline. *Genome Biol.* **21**, 180 (2020).

13. Gautreau, G. *et al.* PPanGGOLiN: Depicting microbial diversity via a partitioned

pangenome graph. *PLoS Comput. Biol.* **16**, e1007732 (2020).

14. Aramaki, T. *et al.* KofamKOALA: KEGG Ortholog assignment based on profile HMM and

adaptive score threshold. *Bioinformatics* **36**, 2251–2252 (2019).

15. Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P. & Huerta-Cepas, J.

eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain

Prediction at the Metagenomic Scale. *Mol. Biol. Evol.* **38**, 5825–5829 (2021).

16. Tatusov, R. L., Galperin, M. Y., Natale, D. A. & Koonin, E. V. The COG database: a tool for

genome-scale analysis of protein functions and evolution. *Nucleic Acids Res.* **28**, 33–36

(2000).

17. Melnyk, R. A., Hossain, S. S. & Haney, C. H. Convergent gain and loss of genomic islands

drive lifestyle changes in plant-associated Pseudomonas. *ISME J.* **13**, 1575–1588 (2019).

18. Steinegger, M. & Söding, J. MMseqs2 enables sensitive protein sequence searching for the

analysis of massive data sets. *Nat. Biotechnol.* **35**, 1026–1028 (2017).

19. Buchfink, B., Ashkenazy, H., Reuter, K., Kennedy, J. A. & Drost, H.-G. Sensitive clustering

of protein sequences at tree-of-life scale using DIAMOND DeepClust. *bioRxiv* (2023)

doi:10.1101/2023.01.24.525373.

20. Coelho, L. P. *et al.* Towards the biogeography of prokaryotic genes. *Nature* **601**, 252–256

(2022).

21. Steinegger, M. & Söding, J. Clustering huge protein sequence sets in linear time. *Nat.*

Commun. **9**, 2542 (2018).

22. Snyder, L., Henkin, T. M., Peters, J. E. & Champness, W. Molecular Genetics of Bacteria,

4th Edition. Preprint at https://doi.org/10.1128/9781555817169 (2013).

23. Price, M. N., Arkin, A. P. & Alm, E. J. The life-cycle of operons. *PLoS Genet.* **2**, e96 (2006).

24. Ptashne, M. *A Genetic Switch: Gene Control and Phage. Lambda.* (Palo Alto, CA (US);

Blackwell Scientific Publications, 1986).

25. Andreu, V. P. *et al.* gutSMASH predicts specialized primary metabolic pathways from the

human gut microbiota. *Nature Biotechnology* Preprint at https://doi.org/10.1038/s41587-

023-01675-1 (2023).

26. Cortes, J., Haydock, S. F., Roberts, G. A., Bevitt, D. J. & Leadlay, P. F. An unusually large

multifunctional polypeptide in the erythromycin-producing polyketide synthase of

Saccharopolyspora erythraea. *Nature* **348**, 176–178 (1990).

27. Donadio, S., Staver, M. J., McAlpine, J. B., Swanson, S. J. & Katz, L. Modular organization

of genes required for complex polyketide biosynthesis. *Science* **252**, 675–679 (1991).

28. Walsh, C. T. & Fischbach, M. A. Natural products version 2.0: connecting genes to

molecules. *J. Am. Chem. Soc.* **132**, 2469–2493 (2010).

29. Medema, M. H. *et al.* antiSMASH: rapid identification, annotation and analysis of secondary

metabolite biosynthesis gene clusters in bacterial and fungal genome sequences. *Nucleic*

Acids Res. **39**, W339-46 (2011).

30. Gal-Mor, O. & Finlay, B. B. Pathogenicity islands: a molecular toolbox for bacterial

virulence. *Cell. Microbiol.* **8**, 1707–1719 (2006).

31. Kaper, J. B., Nataro, J. P. & Mobley, H. L. Pathogenic Escherichia coli. *Nat. Rev. Microbiol.*

2, 123–140 (2004).

32. Bolwell, G. P. & Paul Bolwell, G. Biochemistry & Molecular Biology of Plants.

Phytochemistry vol. 58 185 Preprint at https://doi.org/10.1016/s0031-9422(01)00095-4

(2001).

33. Rokas, A., Mead, M. E., Steenwyk, J. L., Raja, H. A. & Oberlies, N. H. Biosynthetic gene

clusters and the evolution of fungal chemodiversity. *Nat. Prod. Rep.* **37**, 868–878 (2020).

34. Robey, M. T., Caesar, L. K., Drott, M. T., Keller, N. P. & Kelleher, N. L. An interpreted atlas

of biosynthetic gene clusters from 1,000 fungal genomes. *Proc. Natl. Acad. Sci. U. S. A.*

118, (2021).

35. Lindahl, L. & Zengel, J. M. Operon-specific regulation of ribosomal protein synthesis in

Escherichia coli. *Proc. Natl. Acad. Sci. U. S. A.* **76**, 6542–6546 (1979).

36. Cordero, O. X. & Polz, M. F. Explaining microbial genomic diversity in light of evolutionary

ecology. *Nat. Rev. Microbiol.* **12**, 263–273 (2014).

37. Drott, M. T. *et al.* Microevolution in the pansecondary metabolome of Aspergillus flavus and

its potential macroevolutionary implications for filamentous fungi. *Proc. Natl. Acad. Sci. U.*

S. A. **118**, (2021).

38. Salamzade, R. *et al.* Evolutionary investigations of the biosynthetic diversity in the skin

microbiome using lsaBGC. *Microb Genom* **9**, (2023).

39. Ziemert, N. *et al.* Diversity and evolution of secondary metabolism in the marine

actinomycete genus Salinispora. *Proc. Natl. Acad. Sci. U. S. A.* **111**, E1130-9 (2014).

40. van Bergeijk, D. A., Terlouw, B. R., Medema, M. H. & van Wezel, G. P. Ecology and

genomics of Actinobacteria: new concepts for natural product discovery. *Nat. Rev.*

Microbiol. **18**, 546–558 (2020).

41. Chevrette, M. G. *et al.* Evolutionary dynamics of natural product biosynthesis in bacteria.

Nat. Prod. Rep. **37**, 566–599 (2020).

42. Medema, M. H., Takano, E. & Breitling, R. Detecting sequence homology at the gene

cluster level with MultiGeneBlast. *Mol. Biol. Evol.* **30**, 1218–1223 (2013).

43. Abby, S. S., Néron, B., Ménager, H., Touchon, M. & Rocha, E. P. C. MacSyFinder: a

program to mine genomes for molecular systems with an application to CRISPR-Cas

systems. *PLoS One* **9**, e110726 (2014).

44. Navarro-Muñoz, J. C. *et al.* A computational framework to explore large-scale biosynthetic

diversity. *Nat. Chem. Biol.* **16**, 60–68 (2020).

45. Gilchrist, C. L. M. *et al.* Cblaster: A remote search tool for rapid identification and

visualization of homologous gene clusters. *Bioinformatics Advances* **1**, (2021).

46. Gilchrist, C. L. M. & Chooi, Y.-H. clinker & clustermap.js: automatic generation of gene

cluster comparison figures. *Bioinformatics* **37**, 2473–2475 (2021).

47. Hackl, T. & Ankenbrand, M. J. gggenomes: a grammar of graphics for comparative

genomics. *R package version 0.9*.

48. moshi. *PyGenomeViz: A Genome Visualization Python Package for Comparative*

Genomics. (Github).

49. Tajima, F. Statistical method for testing the neutral mutation hypothesis by DNA

polymorphism. *Genetics* **123**, 585–595 (1989).

50. Grazziotin, A. L., Koonin, E. V. & Kristensen, D. M. Prokaryotic Virus Orthologous Groups

(pVOGs): a resource for comparative genomics and protein family annotation. *Nucleic*

Acids Res. **45**, D491–D498 (2017).

51. Liu, B., Zheng, D., Jin, Q., Chen, L. & Yang, J. VFDB 2019: a comparative pathogenomic

platform with an interactive web interface. *Nucleic Acids Res.* **47**, D687–D692 (2019).

52. Terlouw, B. R. *et al.* MIBiG 3.0: a community-driven effort to annotate experimentally

validated biosynthetic gene clusters. *Nucleic Acids Res.* **51**, D603–D610 (2023).

53. Larralde, M. Pyrodigal: Python bindings and interface to Prodigal, an efficient method for

gene prediction in prokaryotes. *J. Open Source Softw.* **7**, 4296 (2022).

54. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site

identification. *BMC Bioinformatics* **11**, 119 (2010).

55. Li, H. Protein-to-genome alignment with miniprot. *Bioinformatics* **39**, (2023).

56. Salamzade, R. & Kalan, L. R. skDER: microbial genome dereplication approaches for

comparative and metagenomic applications. *bioRxivorg* (2023)

doi:10.1101/2023.09.27.559801.

57. Parks, D. H. *et al.* GTDB: an ongoing census of bacterial and archaeal diversity through a

phylogenetically consistent, rank normalized and complete genome-based taxonomy.

Nucleic Acids Res. (2021) doi:10.1093/nar/gkab776.

58. Liu, M. *et al.* ICEberg 2.0: an updated database of bacterial integrative and conjugative

elements. *Nucleic Acids Res.* **47**, D660–D665 (2019).

59. Bertelli, C. *et al.* IslandViewer 4: expanded prediction of genomic islands for larger-scale

datasets. *Nucleic Acids Res.* **45**, W30–W35 (2017).

60. Siguier, P., Perochon, J., Lestrade, L., Mahillon, J. & Chandler, M. ISfinder: the reference

centre for bacterial insertion sequences. *Nucleic Acids Res.* **34**, D32-6 (2006).

61. Alcock, B. P. *et al.* CARD 2023: expanded curation, support for machine learning, and

resistome prediction at the Comprehensive Antibiotic Resistance Database. *Nucleic Acids*

Res. **51**, D690–D699 (2023).

62. Kosakovsky Pond, S. L. *et al.* HyPhy 2.5—A Customizable Platform for Evolutionary

Hypothesis Testing Using Phylogenies. *Mol. Biol. Evol.* **37**, 295–299 (2019).

63. Hackl, T., Duponchel, S., Barenhoff, K., Weinmann, A. & Fischer, M. G. Virophages and

retrotransposons colonize the genomes of a heterotrophic flagellate. *Elife* **10**, (2021).

64. Sullivan, M. J., Petty, N. K. & Beatson, S. A. Easyfig: a genome comparison visualizer.

Bioinformatics **27**, 1009–1010 (2011).

65. Shaw, J. & Yu, Y. W. Fast and robust metagenomic sequence comparison through sparse

chaining with skani. *Nat. Methods* **20**, 1661–1665 (2023).

66. Blackwell, G. *et al.* Exploring bacterial diversity via a curated and searchable snapshot of

1040   archived DNA. *Access Microbiol.* **4**, (2022).

1041 67. Lebreton, F. *et al.* Emergence of epidemic multidrug-resistant Enterococcus faecium from

1042   animal and commensal strains. *MBio* **4**, (2013).

1043 68. Lieberman, T. D. *et al.* Genetic variation of a bacterial pathogen within individuals with

1044   cystic fibrosis provides a record of selective pressures. *Nat. Genet.* **46**, 82–87 (2014).

1045 69. Crits-Christoph, A., Olm, M. R., Diamond, S., Bouma-Gregson, K. & Banfield, J. F. Soil

1046   bacterial populations are shaped by recombination and gene-specific selection across a

1047   grassland meadow. *ISME J.* **14**, 1834–1846 (2020).

1048 70. Hudson, R. R., Slatkin, M. & Maddison, W. P. Estimation of levels of gene flow from DNA

1049   sequence data. *Genetics* **132**, 583–589 (1992).

1050 71. Pavlopoulos, G. A. *et al.* Unraveling the functional dark matter through global

1051   metagenomics. *Nature* **622**, 594–602 (2023).

1052 72. Vanni, C. *et al.* Unifying the known and unknown microbial coding sequence space. *Elife*

1053   **11**, (2022).

1054 73. Tran, P. Q. *et al.* Viral impacts on microbial activity and biogeochemical cycling in a

1055   seasonally anoxic freshwater lake. *bioRxiv* 2023.04.19.537559 (2023)

1056   doi:10.1101/2023.04.19.537559.

1057 74. Kieft, K., Zhou, Z. & Anantharaman, K. VIBRANT: automated recovery, annotation and

1058   curation of microbial viruses, and evaluation of viral community function from genomic

1059   sequences. *Microbiome* **8**, 90 (2020).

1060 75. Willems, A. The Family Comamonadaceae. in *The Prokaryotes* 777–851 (Springer Berlin

1061   Heidelberg, Berlin, Heidelberg, 2014).

1062 76. Roux, S. *et al.* iPHoP: An integrated machine learning framework to maximize host

1063   prediction for metagenome-derived viruses of archaea and bacteria. *PLoS Biol.* **21**,

1064   e3002083 (2023).

1065 77. Klassen, J. L. & Currie, C. R. Gene fragmentation in bacterial draft genomes: extent,

consequences and mitigation. *BMC Genomics* **13**, 14 (2012).

78. Thomma, B. P. H. J. *et al.* Mind the gap; seven reasons to close fragmented genome

assemblies. *Fungal Genet. Biol.* **90**, 24–30 (2016).

79. Drăgan, M.-A., Moghul, I., Priyam, A., Bustos, C. & Wurm, Y. GeneValidator: identify

problems with protein-coding gene predictions. *Bioinformatics* **32**, 1559–1561 (2016).

80. Scalzitti, N., Jeannin-Girardon, A., Collet, P., Poch, O. & Thompson, J. D. A benchmark

study of ab initio gene prediction methods in diverse eukaryotic organisms. *BMC Genomics*

21, 293 (2020).

81. Jallow, A., Xie, H., Tang, X., Qi, Z. & Li, P. Worldwide aflatoxin contamination of agricultural

products and foods: From occurrence to control. *Compr. Rev. Food Sci. Food Saf.* **20**,

2332–2381 (2021).

82. Bok, J. W. *et al.* Genomic mining for Aspergillus natural products. *Chem. Biol.* **13**, 31–37

(2006).

83. Vadlapudi, V. *et al.* Aspergillus Secondary Metabolite Database, a resource to understand

the Secondary metabolome of Aspergillus genus. *Sci. Rep.* **7**, 7325 (2017).

84. Hatmaker, E. A. *et al.* Genomic and Phenotypic Trait Variation of the Opportunistic Human

Pathogen Aspergillus flavus and Its Close Relatives. *Microbiol Spectr* **10**, e0306922 (2022).

85. Cary, J. W. *et al.* An Aspergillus flavus secondary metabolic gene cluster containing a

hybrid PKS-NRPS is necessary for synthesis of the 2-pyridones, leporins. *Fungal Genet.*

Biol. **81**, 88–97 (2015).

86. Skerker, J. M. *et al.* Chromosome assembled and annotated genome sequence of

Aspergillus flavus NRRL 3357. *G3* **11**, jkab213 (2021).

87. van den Belt, M. *et al.* CAGECAT: The CompArative GEne Cluster Analysis Toolbox for

rapid search and visualisation of homologous gene clusters. *BMC Bioinformatics* **24**, 181

(2023).

88. Majoros, W. H., Pertea, M. & Salzberg, S. L. TigrScan and GlimmerHMM: two open source

ab initio eukaryotic gene-finders. *Bioinformatics* **20**, 2878–2879 (2004).

89. Yang, K., Tian, J. & Keller, N. P. Post-translational modifications drive secondary

metabolite biosynthesis in Aspergillus: a review. *Environ. Microbiol.* **24**, 2857–2881 (2022).

90. Klich, M. A. Aspergillus flavus: the major producer of aflatoxin. *Mol. Plant Pathol.* **8**, 713–

722 (2007).

91. Cary, J. W., Ehrlich, K. C., Bland, J. M. & Montalbano, B. G. The Aflatoxin Biosynthesis

Cluster Gene, *aflX* , Encodes an Oxidoreductase Involved in Conversion of Versicolorin A

to Demethylsterigmatocystin. *Applied and Environmental Microbiology* vol. 72 1096–1101

Preprint at https://doi.org/10.1128/aem.72.2.1096-1101.2006 (2006).

92. Cleveland, T. E. *et al.* Potential of Aspergillus flavus genomics for applications in

biotechnology. *Trends Biotechnol.* **27**, 151–157 (2009).

93. Ehrlich, K. C., Li, P., Scharfenstein, L. & Chang, P.-K. HypC, the anthrone oxidase involved

in aflatoxin biosynthesis. *Appl. Environ. Microbiol.* **76**, 3374–3377 (2010).

94. Palmer, K. L. *et al.* Comparative Genomics of Enterococci: Variation in Enterococcus

faecalis, Clade Structure in E. faecium, and Defining Characteristics of *E* . *gallinarum* and *E*

. *casseliflavus. mBio* vol. 3 Preprint at https://doi.org/10.1128/mbio.00318-11 (2012).

95. Qin, X. *et al.* Complete genome sequence of Enterococcus faecium strain TX16 and

comparative genomic analysis of Enterococcus faecium genomes. *BMC Microbiol.* **12**, 135

(2012).

96. Xu, Y., Murray, B. E. & Weinstock, G. M. A cluster of genes involved in polysaccharide

biosynthesis from Enterococcus faecalis OG1RF. *Infect. Immun.* **66**, 4313–4323 (1998).

97. Hancock, L. E., Murray, B. E. & Sillanpää, J. Enterococcal Cell Wall Components and

Structures. in *Enterococci: From Commensals to Leading Causes of Drug Resistant*

Infection (eds. Gilmore, M. S., Clewell, D. B., Ike, Y. & Shankar, N.) (Massachusetts Eye

and Ear Infirmary, Boston, 2014).

98. Teng, F., Jacques-Palaz, K. D., Weinstock, G. M. & Murray, B. E. Evidence that the

Enterococcal Polysaccharide Antigen Gene (*epa*) Cluster Is Widespread in *Enterococcus*

faecalis and Influences Resistance to Phagocytic Killing of *E. faecalis*. *Infection and*

Immunity vol. 70 2010–2015 Preprint at https://doi.org/10.1128/iai.70.4.2010-2015.2002

(2002).

99. Teng, F., Singh, K. V., Bourgogne, A., Zeng, J. & Murray, B. E. Further Characterization of

the *epa* Gene Cluster and Epa Polysaccharides of *Enterococcus faecalis*. *Infection and*

Immunity vol. 77 3759–3767 Preprint at https://doi.org/10.1128/iai.00149-09 (2009).

100. Rigottier-Gois, L. *et al.* The surface rhamnopolysaccharide epa of Enterococcus faecalis is

a key determinant of intestinal colonization. *J. Infect. Dis.* **211**, 62–71 (2015).

101. Guerardel, Y. *et al.* Complete structure of the enterococcal polysaccharide antigen (EPA) of

vancomycin-resistant Enterococcus faecalis V583 reveals that EPA decorations are

teichoic acids covalently linked to a rhamnopolysaccharide backbone. *MBio* **11**, (2020).

102. Smith, R. E. *et al.* Decoration of the enterococcal polysaccharide antigen EPA is essential

for virulence, cell surface charge and interaction with effectors of the innate immune

system. *PLoS Pathog.* **15**, e1007730 (2019).

103. Singh, K. V. & Murray, B. E. Loss of a Major Enterococcal Polysaccharide Antigen (Epa) by

Enterococcus faecalis Is Associated with Increased Resistance to Ceftriaxone and

Carbapenems. *Antimicrob. Agents Chemother.* **63**, (2019).

104. Ho, K., Huo, W., Pas, S., Dao, R. & Palmer, K. L. Loss-of-Function Mutations in *epaR*

Confer Resistance to □NPV1 Infection in Enterococcus faecalis OG1RF. *Antimicrobial*

Agents and Chemotherapy vol. 62 Preprint at https://doi.org/10.1128/aac.00758-18 (2018).

105. Fiore, E., Van Tyne, D. & Gilmore, M. S. Pathogenicity of Enterococci. *Microbiol Spectr* **7**,

(2019).

106. Lebreton, F., Willems, R. J. L. & Gilmore, M. S. Enterococcus Diversity, Origins in Nature,

and Gut Colonization. in *Enterococci: From Commensals to Leading Causes of Drug*

Resistant Infection (eds. Gilmore, M. S., Clewell, D. B., Ike, Y. & Shankar, N.)

(Massachusetts Eye and Ear Infirmary, Boston, 2014).

107. Lebreton, F. *et al.* Tracing the Enterococci from Paleozoic Origins to the Hospital. *Cell* **169**,

849-861.e13 (2017).

108. Schwartzman, J. A. *et al.* Global diversity of enterococci and description of 18 novel

species. *bioRxiv* 2023.05.18.540996 (2023) doi:10.1101/2023.05.18.540996.

109. Bertelli, C. *et al.* Enabling genomic island prediction and comparison in multiple genomes to

investigate bacterial evolution and outbreaks. *Microb. Genom.* **8**, (2022).

110. Bin Jang, H. *et al.* Taxonomic assignment of uncultivated prokaryotic virus genomes is

enabled by gene-sharing networks. *Nat. Biotechnol.* **37**, 632–639 (2019).

111. Meyer, F. *et al.* Critical Assessment of Metagenome Interpretation: the second round of

challenges. *Nat. Methods* **19**, 429–440 (2022).

112. Salamzade, R. *et al.* Inter-species geographic signatures for tracing horizontal gene

transfer and long-term persistence of carbapenem resistance. *Genome Med.* **14**, 37 (2022).

113. Sheppard, A. E. *et al.* Nested Russian Doll-Like Genetic Mobility Drives Rapid

Dissemination of the Carbapenem Resistance Gene blaKPC. *Antimicrob. Agents*

Chemother. **60**, 3767–3778 (2016).

114. Groussin, M. *et al.* Elevated rates of horizontal gene transfer in the industrialized human

microbiome. *Cell* **184**, 2053-2067.e18 (2021).

115. Crits-Christoph, A., Diamond, S., Butterfield, C. N., Thomas, B. C. & Banfield, J. F. Novel

soil bacteria possess diverse genes for secondary metabolite biosynthesis. *Nature* **558**,

440–444 (2018).

116. Bickhart, D. M. *et al.* Generation of lineage-resolved complete metagenome-assembled

genomes by precision phasing. *bioRxiv* 2021.05.04.442591 (2021)

doi:10.1101/2021.05.04.442591.

117. Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative

genomics. *Genome Biol.* **20**, 238 (2019).

118. Nevers, Y. *et al.* The Quest for Orthologs orthology benchmark service in 2022. *Nucleic*

Acids Res. (2022).

119. Chatterjee, A. *et al.* Bacteriophage Resistance Alters Antibiotic-Mediated Intestinal

Expansion of Enterococci. *Infect. Immun.* **87**, (2019).

120. Chatterjee, A. *et al.* Parallel genomics uncover novel enterococcal-bacteriophage

interactions. Preprint at https://doi.org/10.1101/858506.

121. Canfield, G. S. *et al.* Lytic bacteriophages facilitate antibiotic sensitization of *Enterococcus*

faecium. Preprint at https://doi.org/10.1101/2020.09.22.309401.

122. Kirsch, J. M. *et al.* Targeted IS-element sequencing uncovers transposition dynamics

during selective pressure in enterococci. *PLoS Pathog.* **19**, e1011424 (2023).

123. Grüning, B. *et al.* Bioconda: sustainable and comprehensive software distribution for the life

sciences. *Nat. Methods* **15**, 475–476 (2018).

124. Salamzade, R. & Kalan, L. *Zol.* (Zenodo, 2024). doi:10.5281/ZENODO.10828137.

125. Cock, P. J. A. *et al.* Biopython: freely available Python tools for computational molecular

biology and bioinformatics. *Bioinformatics* **25**, 1422–1423 (2009).

126. Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973

(2009).

127. Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2--approximately maximum-likelihood

trees for large alignments. *PLoS One* **5**, e9490 (2010).

128. Huang, Y., Niu, B., Gao, Y., Fu, L. & Li, W. CD-HIT Suite: a web server for clustering and

comparing biological sequences. *Bioinformatics* **26**, 680–682 (2010).

129. Eddy, S. R. Accelerated Profile HMM Searches. *PLoS Comput. Biol.* **7**, e1002195 (2011).

130. Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using DIAMOND.

Nat. Methods **12**, 59–60 (2014).

131. Schreiber, J. Pomegranate: fast and flexible probabilistic modeling in python. *J. Mach.*

Learn. Res. (2017).

132. Edgar, R. C. Muscle5: High-accuracy alignment ensembles enable unbiased assessments

of sequence homology and phylogeny. *Nat. Commun.* **13**, 6968 (2022).

133. Blin, K. *et al.* antiSMASH 6.0: improving cluster detection and comparison capabilities.

Nucleic Acids Res. **49**, W29–W35 (2021).

134. Suyama, M., Torrents, D. & Bork, P. PAL2NAL: robust conversion of protein sequence

alignments into the corresponding codon alignments. *Nucleic Acids Res.* **34**, W609-12

(2006).

135. Li, W. *et al.* RefSeq: expanding the Prokaryotic Genome Annotation Pipeline reach with

protein family model curation. *Nucleic Acids Res.* **49**, D1020–D1028 (2021).

136. Price, M. N. & Arkin, A. P. PaperBLAST: Text Mining Papers for Information about

Homologs. *mSystems* **2**, (2017).

137. Finn, R. D. *et al.* Pfam: the protein families database. *Nucleic Acids Res.* **42**, D222-30

(2014).

138. Larralde, M. & Zeller, G. PyHMMER: a Python library binding to HMMER for efficient

sequence analysis. *Bioinformatics* **39**, (2023).

139. Murrell, B. *et al.* FUBAR: a fast, unconstrained bayesian approximation for inferring

selection. *Mol. Biol. Evol.* **30**, 1196–1205 (2013).

140. Kosakovsky Pond, S. L., Posada, D., Gravenor, M. B., Woelk, C. H. & Frost, S. D. W.

GARD: a genetic algorithm for recombination detection. *Bioinformatics* **22**, 3096–3098

(2006).

141. Carroll, L. M. *et al.* Accurate de novo identification of biosynthetic gene clusters with

GECCO. *bioRxiv* 2021.05.03.442509 (2021) doi:10.1101/2021.05.03.442509.

142. Akhter, S., Aziz, R. K. & Edwards, R. A. PhiSpy: a novel algorithm for finding prophages in

bacterial genomes that combines similarity- and composition-based strategies. *Nucleic*

Acids Res. **40**, e126 (2012).

1222 143. Camargo, A. P. *et al.* Identification of mobile genetic elements with geNomad. *Nat.*

1223 *Biotechnol.* (2023) doi:10.1038/s41587-023-01953-y.

1224 144. Robertson, J. & Nash, J. H. E. MOB-suite: software tools for clustering, reconstruction and

1225 typing of plasmids from draft assemblies. *Microb Genom* **4**, (2018).

1226 145. Ondov, B. D. *et al.* Mash: fast genome and metagenome distance estimation using

1227 MinHash. *Genome Biol.* **17**, 132 (2016).

1228 146. Paradis, E., Claude, J. & Strimmer, K. APE: Analyses of Phylogenetics and Evolution in R

1229 language. *Bioinformatics* **20**, 289–290 (2004).

1230 147. Xu, L. *et al.* Average gene length is highly conserved in prokaryotes and eukaryotes and

1231 diverges only between the two kingdoms. *Mol. Biol. Evol.* **23**, 1107–1108 (2006).

1232 148. Lee, M. D. GToTree: a user-friendly workflow for phylogenomics. *Bioinformatics* **35**, 4162–

1233 4164 (2019).

1234 149. Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v4: recent updates and new

1235 developments. *Nucleic Acids Res.* **47**, W256–W259 (2019).

1236 150. Mistry, J. *et al.* Pfam: The protein families database in 2021. *Nucleic Acids Res.* **49**, D412–

1237 D419 (2021).

1238 151. Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference

1239 in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

1240

Figure 1: Overviews of fai and zol. A) A cartoon schematic of how prepTG, fai, and zol are integrated to perform evolutionary investigations by searching for gene-clusters. Certain statistics in the zol report will not be calculated if not enough instances of an ortholog group are identified, resulting in non-available (NA) values being reported. Squiggles correspond to arbitrary text pertaining to functional annotation information, etc. **B**) An overview of the prepTG, **C**) fai, and **D**) zol algorithms and workflows. Inputs and outputs for the programs are indicated with bolder coloring.

Figure 2: Targeted viral detection in metagenomes using fai. A) Total metagenomes from a single site in Lake Mendota across multiple depths and timepoints from Tran et al. 2023 were investigated using fai for the presence of a virus found in two of the three earliest microbiome samplings (red box). The presence of the virus is indicated by a virus icon. Metagenome samples are colored according to whether they corresponded to oxic, oxycline, or anoxic. The most shallow sampling depths varied for different dates and consolidated as a single row corresponding to a sampling depth of either 5 or 10 meters. **D)** The pangenome of the virus is shown based on the consensus order and directionality of coding sequences inferred by zol. Bar heights correspond to the conservation of the ortholog groups across the ten metagenomes the virus was detected in. BioRender was used in generation of this figure.

Figure 3: Evolutionary trends of common BGCs in *A. flavus*. A) The proportion of 216 A. flavus genomes from NCBI's GenBank database with coding-sequence predictions available. **B)** Comparison of the sensitivity of fai and alternate approaches based on assemblies for detecting the leporin BGC. The dashed violet line indicates the total number of genomes (n=216) assessed and the dashed pink line indicates the number of genomes with CDS features available on NCBI (n=11). Dark grey indicates instances identified by CAGECAT/cblaster or fai or as belonging to the same GCF as the reference leporin BGC from MIBiG by antiSMASH and BiG-SCAPE analysis. Lighter grey indicates the number of similar BGCs identified by BiG-SCAPE, belonging to the same clan but not to the same GCF as the reference leporin BGC. A schematic of the (**C**) leporin and (**D**) aflatoxin BGCs is shown with genes present in ≥ 10% of samples shown in consensus order and relative directionality. Coloring of genes in (**C**) corresponds to FST values and in (**D**) to Tajima's D values, as calculated by zol. Grey bars in the legends, at (C) 0.92 and (D) -0.98, indicate the mean values for the statistics across genes in the BGC. *For the leporin BGC, *lepB* corresponds to an updated open-reading frame (ORF) prediction by Skerker *et al.* 2021 which was the combination of AFLA_066860 and AFLA_066870 ORFs in the MIBiG entry BGC0001445 used as the query for fai. For the aflatoxin BGC, ORFs which were not represented in the MIBiG entry BGC0000008 but predicted to be within the aflatoxin BGC by mapping of gene-calls from *A. flavus* NRRL 3357 by Skerker et al. 2021 are shown in gold. The major allele frequency distributions are shown for (**E**) *aflX* and (**F**) *pksA*, which depict opposite trends in sequence conservation according to their respective Tajima's D calculations.

Figure 4: Searching for the *epa* locus across the diverse genus of *Enterococcus*. A) Overview of the time needed to run orthology/homology inference methods on the 92 genomes with the highest N50 for each distinct *Enterococcus* species. OrthoFinder was run at the genome-wide scale, while fai and cblaster were used to first identify genomic regions corresponding to the *epa* locus from *E. faecalis* V583 and subsequently zol and clinker were applied to determine ortholog groups, respectively. The red asterisks denote that manual assessment or filtering of homologous gene clusters identified by fai and cblaster is encouraged and thus additional time might be required for them. Counts showing the overlap in orthologous protein pair predictions by the three different methods are shown following their application to representative genomes from GTDB R214 with the **B**) highest N50 and **C**) lowest N50 for the 92 different species. **D**) The distribution of the *epa* locus, based on criteria used for running fai, is shown across a species phylogeny for 92 genomes representative of distinct *Enterococcus* species in GTDB R214. The coloring of the heatmap corresponds to the percent identity of the best matching protein from each genome to the query *epa* proteins from *E. faecalis* V583. **E**) A schematic of the *epa* gene cluster from *E. faecalis* V583 (from EF2164 to EF2200) with glycosyltransferase encoding genes shown in color. **F**) A maximum-likelihood phylogeny of zol-identified ortholog groups corresponding to glycosyltransferases in *epa* loci across *Enterococcus*. **G**) Distribution of different glycosyltransferase ortholog groups across the four major clades of *Enterococcus* are shown. For **D** and **F** the tree scales correspond to the number of amino acid substitutions along the alignments used for phylogeny construction.

Figure 5: High sequence diversity of *epaX*-like glycosyltransferases amongst *E. faecalis* . A schematic of the *epa* locus from *E. faecalis* V583 with evolutionary statistics, **A**) conservation, **B**) Tajima's D and **C**) sequence entropy, gathered from the best corresponding ortholog group for each protein. Ortholog groups were inferred from zol investigation of 1,232 *epa* loci from the species. Genes upstream of and including *epaR* were recently proposed to be involved in Epa decoration by Guerardel *et al.* 2020. "//" indicates that the ortholog group was not single-copy in the context of the gene-cluster and calculation of evolutionary statistics for these genes was avoided (grey in panels B and C). Note, the same ortholog group was regarded for EF2173 and EF2185 which correspond to an identical *ISEf1* transposase. The length of proteins in the locus schematic are the median lengths of the corresponding ortholog groups. **D**) The major allele frequency is depicted across the alignment for the ortholog group featuring *epaX*. Sites predicted to be under negative selection by FUBAR, Prob($\alpha > \beta$) $\geq$ 0.9, are marked in red. **E**) An approximate maximum-likelihood phylogeny of glycosyltransferase ortholog groups identified by zol which were found in >1% of *epa* instances. Ortholog groups identified by zol are indicated by colored circular nodes with names of *epa* genes from *E. faecalis* V583 noted where possible. The number of leaves/proteins for each clade is provided for labeled ortholog groups. The tree scale corresponds to the number of amino acid substitutions along the input protein alignment used for phylogeny construction.