zol & fai: large-scale targeted detection and evolutionary investigation of gene clusters

Rauf Salamzade[1,2], Patricia Tran[3,4], Cody Martin[2,3], Abigail L. Manson[5], Michael S. Gilmore[5,6,7],
Ashlee M. Earl[5], Karthik Anantharaman[3], Lindsay R. Kalan[1,8,9]

[1]Department of Medical Microbiology and Immunology, School of Medicine and Public Health, University of
Wisconsin-Madison, Madison, WI, USA
[2]Microbiology Doctoral Training Program, University of Wisconsin-Madison, Madison, WI, USA
[3]Department of Bacteriology, University of Wisconsin-Madison, Madison, WI, USA
[4]Freshwater and Marine Science Doctoral Program, University of Wisconsin-Madison, WI, USA
[5]Infectious Disease and Microbiome Program, Broad Institute of MIT and Harvard, Cambridge, Massachusetts, USA
[6]Department of Ophthalmology, Harvard Medical School and Mass Eye and Ear, Boston, Massachusetts, USA
[7]Department of Microbiology, Harvard Medical School and Mass Eye and Ear, Boston, Massachusetts, USA
[8]Department of Medicine, Division of Infectious Disease, School of Medicine and Public Health, University of
Wisconsin-Madison, Madison, WI, USA
[9]M.G. DeGroote Institute for Infectious Disease Research, David Braley Centre for Antibiotic Discovery, Department
of Biochemistry and Biomedical Sciences, McMaster University, Hamilton, Ontario, Canada

Address for Correspondence: Lindsay R. Kalan; kalanlr@mcmaster.ca

Abstract

Many universally and conditionally important genes are genomically aggregated within
clusters. Here, we introduce fai and zol, which together enable large-scale comparative analysis
of different types of gene clusters and mobile-genetic elements (MGEs), such as biosynthetic
gene clusters (BGCs) or viruses. Fundamentally, they overcome a current bottleneck to reliably
perform comprehensive orthology inference at large scale across broad taxonomic contexts and
thousands of genomes. First, fai allows the identification of orthologous or homologous
instances of a query gene cluster of interest amongst a database of target genomes.
Subsequently, zol enables reliable, context-specific inference of protein-encoding ortholog
groups for individual genes across gene cluster instances. In addition, zol performs functional
annotation and computes a variety of statistics for each inferred ortholog group. These
programs are showcased through application to: (i) longitudinal tracking of a virus in
metagenomes, (ii) discovering novel population-genetic insights of two common BGCs in a
fungal species, and (iii) uncovering large-scale evolutionary trends of a virulence-associated
gene cluster across thousands of genomes from a diverse bacterial genus.

Introduction

Within bacterial genomes, genes are often co-located within smaller genetic structures
such as operons[1,2], phages[3], metabolic gene clusters[4], biosynthetic gene clusters (BGCs)[5], and
pathogenicity islands [6,7]. Although less prevalent, eukaryotic genomes also contain genes
aggregated within discrete clusters[5,8].
Sometimes gene clusters are highly conserved, encoding for products essential to the
survival of the organism[9]. In other cases, a single gene cluster can exhibit variability in gene
carriage and order across different strains or species[10–12]. This is often the case for BGCs

encoding specialized metabolites or virulence-associated gene clusters, where evolution of
gene content and sequence divergence can influence fitness and contribute to adaptation within
a changing ecosystem.
Bioinformatic toolkits to perform accurate pangenomic and comparative genomic
analyses have been heavily developed over the past two decades[13–18]; however, tool
development to aid the identification and comparative analysis of smaller homologous gene
clusters has been more limited and largely designed for specific types of gene clusters[19–22]. In
addition, while methods for comprehensive comparative genomics within species exist and are
scalable[17,23,24], methods for reliable, large-scale comparative genomics of thousands of
genomes representing a greater breadth of taxonomic diversity are lacking and still bear heavy
computational costs[25–27]. Context-specific inference of orthologous genes within homologous
gene clusters offers a targeted and reliable solution to overcome challenges with scalability[28,29].
Such an approach was recently taken to infer orthologous genes between instances of
homologous BGCs[22].
Here, we introduce fai (*find-additional-instances*) and zol (*zoom-on-locus*), which are
designed for the identification (fai) and in-depth evolutionary genomics investigations (zol) of a
wide array of gene cluster types. We demonstrate the utility of these programs through
application to three types of gene clusters within different genomic contexts including a novel
bacteriophage within environmental metagenomes, a fungal secondary metabolite encoding
biosynthetic gene clusters, and a conserved polysaccharide antigen locus within the diverse
bacterial genus of *Enterococcus.*
68
69   **Results**
70
71   **fai and zol allow for the rapid inference of gene cluster orthologs across diverse**
72   **genomes**
73
The two programs, fai and zol, build upon approaches we recently reported in *lsa*BGC[12]
that were developed to investigate evolutionary trends of BGCs in a single taxon. Within fai and
zol, algorithmic adjustments have been implemented to broaden the application for searching
any type of gene cluster across a diverse set of target genomes (Figure 1A). First, fai allows
users to rapidly search for gene cluster instances in a target set of genomes. Then, zol can be
used to compute evolutionary statistics and functional annotations of gene cluster content in
table-based reports. Importantly, because fai has an option to filter secondary, potentially
paralogous, instances of gene clusters found in target genomes, downstream *ab initio* clustering
of encoded proteins using a flexible, InParanoid-type algorithm[14] by zol can be used to reliably
infer ortholog groups.
In addition to filtering secondary instances of query gene clusters identified in target
genomes, detection criteria in fai can be adjusted by assessing whether gene cluster homologs
lie near scaffold edges in target genomic assemblies. This feature overcomes challenges
inherent to the identification of full gene clusters in metagenomic assemblies or metagenome-
assembled genomes, which can be highly fragmented (Figure S1). fai can further accept query
gene clusters in different formats to ease searching for gene clusters and genomic islands
cataloged in databases such as ICEberg[30], MIBiG[31], or IslandViewer[32]. In addition, to promote

consistency in gene calling across target genomes, we have incorporated computationally light-
weight dependencies for *de novo* gene prediction in prokaryotic genomes[33,34] and gene-
mapping in eukaryotic genomes[35] within prepTG, to prepare and format target genomes for
optimized gene cluster searching in fai (Figure 1B). Together these unique features and options
differentiate fai from other software with similar functionalities, such as cblaster[21] (Figure 1C,
S1; Table S1; Supplementary Text).
zol is differentiated from *lsa*BGC[12], where ortholog groups are inferred across full
genomes using OrthoFinder[18], by delineating ortholog groups within the context of a
homologous or orthologous set of gene clusters, similar to the approach taken within
CORASON[22] to visualize similarities between BGCs. While CORASON uses bidirectional best-
hits to identify direct orthologs, zol accounts for the presence of in-paralogs and
comprehensively partitions proteins into ortholog groups. Similar to *lsa*BGC-PopGene[12], zol will
then construct a tabular report with information on conservation, evolutionary trends, and
annotation for individual ortholog groups (Figure 1D). To make annotated reports generated by
zol more comprehensive for different types of gene clusters, several databases have been
included, such as VOGs[36], VFDB[37], ISFinder[38], and CARD[39]. In addition, zol incorporates
HyPhy[40] as a dependency and calculates evolutionary statistics not previously reported in
*lsa*BGC-PopGene, such as sequence entropy in the 100 bp upstream of an ortholog group,
where important regulatory differences could exist[41]. Ultimately, beyond high-throughput
inference of ortholog groups across diverse genomic datasets, the rich tabular report produced
by zol provides complementary information to figures generated by comparative visualization
software such as clinker[42], CORASON[22], gggenomes[43], and Easyfig[44].
Another key feature in zol is the ability to dereplicate gene clusters directly using skani[45],
which was recently shown to be more reliable at estimating ANI between genomes of variable
contiguity relative to comparative methods. Dereplication allows for more appropriate inference
of evolutionary statistics to overcome availability or sampling biases in genomic databases[46].
Finally, zol allows for comparative investigations of gene clusters based on taxonomic or
ecological groupings[47–49]. For instance, users can designate a subset of gene clusters as
belonging to a specific population to allow zol to calculate ortholog group conservation across
just the focal set of gene clusters. In addition, if comparative investigations are requested, zol
will also compute the fixation index[50], F_{ST}, for each ortholog group to assess gene flow between
the focal and complementary sets of gene clusters.
123
124　**Longitudinal tracking of a virus within lake metagenomic assemblies**
125
Viruses are important members of host and environmental microbiomes[51–53], influencing
the microbial composition and participating in several metabolic pathways. Targeted
identification of a specific virus or bacteriophage within metagenomes can thus offer greater
insight into their elusive functional roles in microbiomes.
Recently, changes in the composition and function of the metagenome at three different
depths of a lake was reported using longitudinal shotgun metagenomics[54]. Using metagenome
assemblies generated from this dataset, large ($\geq$20kb) and predicted-circular phages were
identified independently across a subset of metagenomes from the three different depths at the
the earliest sampling date using VIBRANT[55]. Subsequent clustering based on the sequence and

syntenic similarity of protein domains identified a ~36kb highly conserved virus in two
metagenomes sampled from lower lake depths.
fai was then used to perform a rapid, targeted search for this ~36kb *Caudovirales* virus
across the full set of 16 metagenomes to identify additional instances of the virus. fai completed
its search of the metagenomes, featuring >20 million proteins and 10.7 million contigs, in less
than seven minutes using 20 threads. Of the 16 total metagenomes, spanning five distinct
sampling timepoints and four distinct sampling depths, nine metagenomes containing the virus
were identified (Figure 2A) exclusively from anoxic conditions (p=8.7e-5; two-sided Fisher's
exact test). This suggests the viral host likely performs anaerobic respiration. Application of zol
further revealed that 34 (64%) of the 53 total distinct ortholog groups were core to all instances
of the virus across nine metagenomes and completely conserved in sequence over the course
of 2.5 months (Figure 2B; Table S2). Furthermore, seven of the 53 ortholog groups were not
observed in the query viruses from the earliest sampling date, demonstrating the ability of fai to
identify new genes within additional instances of known gene clusters.
149
150   **Investigating population-level and species-wide evolutionary trends of BGCs in the**
151   **eukaryotic species *Aspergillus flavus***
152
The fungal genus of *Aspergillus* is a source of several natural products, including
aflatoxins, a common and economically impactful contaminant of food. The genus also contains
species that are model organisms for studying fungal secondary metabolism[56–58]. Examination
of the secondary metabolome of *A. flavus* has revealed that different clades or populations
comprising certain species can exhibit variability in their metabolite production despite high
conservation of core BGC genes encoding enzymes for synthesis of these metabolites[11,59,60].
For instance, population B *A. flavus* were identified as producing a greater abundance of the
insecticide leporin B relative to populations A and C[11,61].
To further understand the genomic basis for differences in metabolite content between
populations, we investigated the leporin BGC using fai and zol. While the leporin cluster was
previously identified as a core component of the *A. flavus* genome[11], a recent study suggested
that the full BGC was specific to a single clade from the species[60]. Low sensitivity in direct
assessment of gene cluster presence in eukaryotic genome assemblies can arise from their
incompleteness, leading to gene clusters being fragmented across multiple scaffolds, and
challenges in *ab initio* gene prediction[62,63]. Further deterring the direct prediction of gene
clusters in eukaryotic assemblies is the lack of gene annotations, with only 11 (5.1%) of 216 *A.*
flavus genomes in NCBI's GenBank database having coding sequence predictions (Figure 3A).
Therefore, we used miniprot[35], which is integrated within prepTG, to directly map high-quality
coding genes predictions based on transcriptomics data from the genome of strain *A. flavus*
NRRL 3357[64] to the 216 genomes available for the species. Running fai in "draft mode" led to
the identification of the leporin BGC within 212 (98.1%) assemblies, consistent with prior read
mapping-based investigations[11]. This increase in sensitivity when fai is run with miniprot-based
gene-mapping is substantial when compared to common alternate approaches for identifying
homologous instances of BGCs across genomes (Figure 3B; Supplementary Text).
Of the 212 genomes with the leporin BGC, 202 contained instances that were high-
quality and not near scaffold edges. This set of 202 instances of the gene cluster were further

investigated using zol, with comparative investigation of BGC instances from *A. flavus*
population B genomes to instances from other populations requested. High sequence
conservation was observed for all genes in the leporin gene cluster as previously reported[11]
(Table S3). Further, alleles for genes in the BGC from population B genomes were generally
more similar to each other than to alleles from outside the population as indicated by high F_{ST}
values (>0.85 for 9 of 10 genes) (Figure 3C; Table S3). While regulation of secondary
metabolites in *Aspergillus* is complex[65], zol analysis showed that the three essential genes for
leporin production[61] also had the lowest variation in the 100 bps upstream their exonic
coordinates (Figure S2). This suggests higher variability is occurring in the transcription of the
accessory *lep* genes within the species. This supports experimental evidence that has shown
gene knockouts depleting certain leporin species will still permit the production of others[61].
fai and zol were also applied to the BGC encoding aflatoxin across *A. flavus*[66] (Table
S4). Similar to the leporin BGC, the aflatoxin BGC was highly prevalent in the species and found
in 71.8% of genomes. However, in contrast to the leporin BGC, the aflatoxin BGC contained
several genes with positive Tajima's D values, indicating greater sequence variability for these
coding regions across the species (Figure 3D). One of the genes with a positive Tajima's D
value was *aflX*, which has been shown to influence conversion of the precursor veriscolorin A to
downstream intermediates in the aflatoxin biosynthesis pathway[67] (Figure 3E). An abundance of
sites with mid-frequency alleles in the oxidoreductase encoding gene could represent granular
control for the amount of aflatoxin relative to intermediates produced. The polyketide synthase
gene *pksA* had the lowest Tajima's D value of -2.4, which suggests it is either highly conserved
or under purifying selection (Figure 3F). In addition, because a recent predicted reference
proteome was used to infer genomic coding regions, fai and zol detected several highly
conserved genes within the aflatoxin BGC that are not represented in the original reference
gene cluster input for fai[31]. This includes a gene annotated as a noranthrone monooxygenase
and recently characterized as contributing to aflatoxin biosynthesis[68,69] (Figure 3D).
205
206   **Large-scale identification of the Enterococcal polysaccharide antigen and assessment of**
207   **context restricted orthology inference**
208
The Enterococcal polysaccharide antigen (Epa) is a signature component of the cellular
envelope of multiple species within *Enterococcus*[70–73], which has mostly been characterized in
the species *Enterococcus faecalis*[70,74–77]. While molecular studies have provided evidence that
the locus contributes to enterococcal host colonization[76], evasion of immune systems[78], and
sensitivity to antibiotics[79] and phages[79,80], it was only recently that the structure of Epa was
resolved and a model for its biosynthesis and localization formally proposed[77]. A homologous
instance of the *epa* locus was identified in the other prominent pathogenic species from the
genus, *Enterococcus faecium*[71,73,81]; however, the prevalence and conservation of *epa* across
the diverse genus of *Enterococcus*[82–84] remains poorly studied.
fai was used to search for homologous instances of *epa* across 5,291 enterococcal
genomes*,* estimated by GTDB to represent 92 species[85], using a sensitive searching criterium
and coordinates of the locus along the *E. faecalis* V583 genome as a reference[75,77]
(Supplementary Text). For detection of *epa* orthologous regions, co-location of at least seven of
the 14 *epa* genes previously identified as conserved in both *E. faecalis* and *E. faecium* was

required. The default threshold for syntenic conservation of homologous instances to the query
gene cluster was also disregarded to increase sensitivity for the detection of *epa* in enterococcal
species more distantly related to *E. faecalis*. To allow for capture and downstream analysis of
auxiliary genes, which might be species or strain-specific, but related to Epa production or
decoration, 20 kb flanking contexts of the core *epa* genes identified in each target genome were
extracted.
Using these criteria, 5,085 (96.1%) genomes from across the genus were found to
possess an *epa* locus, confirming the locus as nearly core to the genus. Visual inspection of the
epa genes among 463 representative *Enterococcus* genomes revealed that the core genes
*epa*A-*epa*R were highly conserved in three of four major clades (Figure 4; Supplementary Text).
Based on the detection criteria in fai, the *epa* locus in the fourth clade, previously referred to as
the *Enterococcus columbae* group[82], was either missing or encoded for highly divergent
homologs of these genes. This clade includes *Enterococcus gallinarum*, one of the only other
species in the genus besides *E. faecalis* and *E. faecium,* reported to cause nosocomial
outbreaks[86,87].
Evolutionary trends and sequence diversity for individual genes with the *epa* locus were
next computed using zol, after assessing zol's reliability for gene cluster context-limited
inference of orthology and the impact of dereplication on the calculation of evolutionary statistics
by zol.
242
243   **Gene-context specific orthology inference using fai and zol are concordant with genome-**
244   **wide ortholog group predictions**
245
Genome-wide orthology inference is currently difficult to scale to hundreds or thousands
of genomes belonging to multiple species. However, orthology inference can be made more
accessible if larger loci are first identified as orthologous between genomes, through leveraging
syntenic support[23,28]. To assess whether ortholog group inference was reliable when zol is
applied on orthologous gene clusters identified across multiple species, we ran zol on high-
quality instances of the *epa* locus from 42 different species (Figure 5A). Ortholog group
predictions by zol were then compared to genome-wide orthology predictions by OrthoFinder[18],
which has been shown to yield highly accurate predictions in benchmarking experiments
involving genomes from multiple species[88]. Orthology predictions were highly concordant
between zol and OrthoFinder for proteins from diverse instances of the *epa* locus. zol identified
23,623 pairs of proteins within ortholog groups, of which 22,843 (96.70%) were also grouped
together by OrthoFinder. Only 1,520 (6.24%) pairs of *epa*-associated proteins which were
identified by OrthoFinder to belong to the same ortholog group were missed by zol.
Because the *epa* locus encodes multiple characterized and putative
glycosyltransferases[89], we used phylogenetics to examine the relationship between proteins
belonging to ortholog groups with glycosyltransferase domains to confirm that major clades
correspond to distinct ortholog group designations (Figure 5BC). zol also has an option to "re-
inflate" ortholog groups, expanding them to include proteins from gene clusters which were
deemed redundant during dereplication. To demonstrate the scalability of zol to process
thousands of gene-clusters, this "re-inflation"-based approach was next applied on the full set of
high-quality and contiguous *epa* instances from across the genus and a comprehensive

phylogeny of ortholog groups corresponding to glycosyltransferases was constructed. In
concordance with our analysis using the 42 representative genomes, distinct phylogenetic
clades for the 21,586 glycosyltransferases from 2,442 genomes mostly corresponded to
individual ortholog groups identified by zol (Figure 5D).
271
272 **Dereplication can impact taxa-wide inferences of selection-informative statistics**
273
Dereplication, or removal of redundant gene cluster instances, is important to consider
when working with highly sequenced bacterial taxa, including *E. faecalis*, where certain
lineages, such as those commonly isolated at clinics, can be overrepresented in genomic
databases. Over-representation of select lineages due to sampling bias will skew estimates for
some evolutionary statistics, such as those informative of selective pressures, complicating
evaluation of evolutionary trends across the entire taxonomic group. We thus assessed the
impact of dereplication on the calculation of evolutionary statistics for instances of *epa* in *E.*
faecalis using two different approaches: (i) genome-wide dereplication with dRep[90,91] and (ii)
gene cluster specific dereplication with skani[45]. Dereplication at the gene cluster level with skani
was performed directly in zol. The "re-inflation" option was also used to simulate comprehensive
processing and calculation of evolutionary statistics. This option thus enables users to retain
selection-informative redundancy amongst gene clusters if appropriate for the analysis of
interest while avoiding excessive computation when inferring ortholog groups.
Regardless of the approach for dereplication, genome-wide or gene cluster-specific, the
estimates of evolutionary and genomic statistics for analogous ortholog groups were highly
concordant (Figure 6, S3). However, gene cluster based dereplication can lead to slight
differences for selection informative statistics, such as FUBAR-based[92] inference of the
proportion of sites under selection under positive selection, relative to the usage of genome-
wide dereplication. This is likely because the core *epa* locus is highly conserved across *E.*
faecalis which led to fewer representative gene clusters following dereplication and a lower
weight being placed on conserved alleles when estimating such statistics. In contrast, more
simplistic statistics, such as average sequence entropy and the proportion of total alignment
sites regarded as segregating sites, were closely estimated for genes regardless of the
dereplication method used.
298
299 **zol identifies genetic diversity of *epa*X-like glycosyltransferases**
300
Because Epa biosynthesis and its conditional importance has mostly been investigated
in *E. faecalis*[70,74,75,77], we first examined evolutionary trends for proteins across instances of the
epa locus from 75 *E. faecalis* representative genomes following genome-wide dereplication. In
accordance with prior studies[71,77], zol reported that one end of the locus corresponds to genes
which are highly conserved and core to *E. faecalis* (*epa*A-*epa*R), whereas the other end
contains strain-specific genes (Figure 7A; Table S5). Using zol, we further found that variably
conserved genes exhibit high sequence dissimilarity, as measured using both Tajima's D and
average sequence entropy, in comparison to the core genes of the locus (Figure 7BC).
Comparative and multi-species analysis of the *epa* locus between and across *E. faecalis* and *E.*
faecium was next performed using gene cluster based dereplication with re-inflation using zol

(Table S6). zol-reported conservation statistics were consistently in agreement with previous
studies[71,73]. Twenty genes were determined to be present in the majority (>95%) of *epa* clusters
across both species, including *epa*ABCDEFGH, *epa*LM, and *epa*OPQR. In addition, default
parameters for orthologous clustering of proteins in zol detected a known truncated variant of
the glycosyltransferase *epa*N in *E. faecium*.
The gene *epaX,* encoding a glycosyltransferase, was identified as one ortholog group
with the greatest sequence variation in *E. faecalis* (Figure 7BD, S4). *epa*X was previously
shown to be critical for *E. faecalis* host-gut colonization and proposed to be involved in the
decoration of the rhamnan backbone structure of Epa with galactose and N-acetyl
glucosamine[76]. Comparative analysis using *E. faecium* as the focal taxa further showed that the
*epa*X-containing ortholog group had a low F_{ST} value of 0.19, indicating alleles from *E. faecalis*
and *E. faecium* species are phylogenetically interspersed. This was confirmed through
phylogenetic assessment of the ortholog group (Figure 7E). In addition, although some allelic
clades encoded sequences from both species, genes remained sub-partitioned by species. This
phylogenetic structure for the ortholog group, together with our prior observation that the *epa*X-
containing ortholog group in *E. faecalis* has greater sequence variability relative to other
glycosyltransferases from the locus, suggests extensive and ancestral sequence evolution of
*epa*X-like glycosyltransferases. Further, while only 70% of *E. faecium* with contiguous instances
of the *epa* locus possessed an *epa*X-like ortholog group, approximately 7% of them encoded
the ortholog in multi-copy (Figure 7F), suggesting the occurrence of intra-locus gene duplication.
331
332   **Discussion**
333
Here fai and zol are introduced to enable large scale evolutionary investigations of gene
clusters in diverse taxa. Together these tools overcome current bottlenecks in computational
biology to infer orthologous sets of genes at scale across thousands of diverse genomes.
Both fai and cblaster[21] can be used to identify additional gene clusters within target
genomes and extract them in GenBank format for downstream investigations using zol. For
those lacking computational resources needed for fai analysis, cblaster offers remote searching
of BGCs using NCBI's BLAST infrastructure and non-redundant databases. More recently,
CAGECAT[93], a highly accessible web-application for running cblaster, was also developed and
can similarly be used to identify and extract gene cluster instances from genomes represented
in NCBI databases. In contrast to these tools, fai contains algorithms and options for users
interested in: (i) identifying gene clusters across a comprehensive or redundant set of genomic
assemblies, (ii) improved sensitivity for gene cluster detection in draft-quality assemblies, and
(iii) automated filtering of secondary, or paralogous, matches to query gene clusters. In addition,
users can apply zol to further investigate homologous sets of gene clusters identified from
IslandCompare[94], BiG-SCAPE[22], or vConTACT2[95] analyses, which perform comprehensive
clustering of predicted genomic islands, BGCs, or viruses.
The utility of fai is demonstrated here through rapid, targeted detection of a virus directly
from lake metagenomic assemblies. Targeted detection of specific viruses longitudinally
presents an efficient and tractable approach to understand how viral pangenomes evolve over
time. In addition, by permitting fragmented detection of gene clusters and detection of proximity
to scaffold edges, users can assess whether phages or other gene clusters corresponding to

MGEs are present in their metagenomes. fai and zol will continue to compliment metagenomic
applications as long-read sequencing becomes more economical and commonly used to profile
microbial communities. For example, their application could be useful for assessing the
presence of concerning MGEs conferring antimicrobial resistance traits[96–98] and identifying novel
auxiliary genes within known BGCs which may tailor the resulting specialized metabolites and
expand chemical diversity[99,100].
Reidentifying gene clusters in eukaryotic genomes remains difficult due to technical
challenges in gene prediction owing to the presence of alternative splicing. The ability of fai and
zol to perform population-level genetics on common BGCs from the eukaryotic species *A. flavus*
was demonstrated. While there are over 200 genomes of *A. flavus* in NCBI, only 5.1% have
coding-sequence information readily available. We used miniprot[35] to map high quality gene
coordinate predictions from a representative genome in the species[64] to the remainder of
genomic assemblies within prepTG which enabled high sensitivity detection of BGCs with fai.
Our analysis provides additional support that the leporin BGC is conserved in full across the
species[11] using an assembly-based approach.
Application of fai and zol to exopolysaccharide encoding gene clusters from pathogens
of interest allows a better understanding of their conservation and evolutionary trends. This
information can then aid the identification of potential genes to target for antivirulence
efforts[101,102] or genes underlying host-pathogen interactions[76,103]. fai was used to identify
orthologous instances of the *epa* locus, encoding for an extracellular polysaccharide antigen,
across thousands of diverse genomes from the genus of *Enterococcus*. Subsequently,
application of zol reliably produced orthology predictions comparable to OrthoFinder, a highly
dependable genome-wide orthology inference software[18,88]. While zol missed a small
percentage of orthologous instances identified by OrthoFinder in our testing, this could be due
to threshold settings for percent identity and coverage between pairs of proteins set in zol. Such
thresholds are not enforced in OrthoFinder. However, parameters controlling these thresholds
are adjustable in zol and allow users to increase or decrease orthology sensitivity at the
expense of incurring false positives as they deem appropriate for their research objective.
Using zol, it was determined that an ortholog group containing *epa*X-like
glycosyltransferases possessed high sequence divergence relative to other
glycosyltransferases within the *epa* locus in *E. faecalis*. In addition to influencing the ability of *E.*
faecalis to colonize hosts[76], mutations in *epa*X and other genes from the ortholog group have
also been shown to impact susceptibility to phage predation[104–107]. Thus, because similar *epa*X-
like glycosyltransferases are found in both *E. faecalis* and *E. faecium*, we hypothesize that
extensive ancestral evolution of the *epa*X-containing ortholog group may have occurred to
support evasion from phages and confer colonization of new hosts. In this study, we further
found that the *E. columbae* group might lack or possess highly divergent versions of core *epa*
genes found in *E. faecalis* and *E. faecium*, suggesting that development of anti-virulence
approaches to broadly target Epa in all pathogenic enterococci might be difficult to achieve.
Similar investigations with fai and zol can readily be performed for other exopolysaccharide
encoding gene clusters of pathogens to better understand their conservation, evolutionary
trends, identify appropriate genes to target for antivirulence efforts[101,102], and infer whether
certain genes underlie host-pathogen interactions[76,103].

Options for dereplication and re-inflation provided within zol enable scalability to
thousands of gene cluster instances. The usage of these options can further aid in performing
more accurate evolutionary investigations for genes broadly across focal taxa or between
clades, by overcoming biases due to overrepresentation of certain lineages in genomic
databases[11,47]. Depending on the underlying origin of input gene clusters, zol can also be used
to assess temporal[48,108] or spatial[49] evolutionary trends.
Practically, zol presents a comprehensive analysis tool for comparative genetics of
related gene clusters to facilitate detection of evolutionary patterns that might be less apparent
from visual analysis. Fundamentally, the algorithms presented within fai and zol enable the
reliable detection of orthologous gene clusters, and subsequently orthologous proteins, across
multi-species datasets spanning thousands of genomes and help overcome a key barrier in
scalability for comparative genomics.
410

Acknowledgments

This work was supported by grants from the National Institutes of Health awarded to L.R.K
(NIAID U19AI142720 and NIGMS R35GM137828) and the Broad Institute (U19AI110818). The
content is solely the responsibility of the authors and does not necessarily represent the official
views of the National Institutes of Health. The authors are grateful to James Kosmopoulos, Dr.
Caitlin Sande, and Mary Hannah Swaney for feedback and assistance with data acquisition as
well as Dr. Devon Ryan and Dr. Robert A. Petit III for assistance with incorporation of the suite
into Bioconda.

Methods

Software availability

425   zol is provided as an open-source software suite, developed primarily in Python3 on GitHub at:
426   https://github.com/Kalan-Lab/zol. Docker and Bioconda[109] based installations of the suite are
427   supported. For the analyses presented in this manuscript, we used v1.3.6 of the zol software
428   package[110]. Version information for major dependencies of the zol suite[33,35,40,45,111–118] and other
429   software used[22,55,119] for analyses in this study is provided in Supplementary Table S7.

Availability of data and materials

Genomes and metagenomes used to showcase the application of fai and zol are listed with
GenBank accession identifiers in Supplementary Table S8. Total metagenomes and their
associated information from Lake Mendota microbiome samplings were originally described in
Tran *et al.* 2023[54] and deposited in NCBI under BioProject PRJNA758276. Genomic assemblies
available for *A. flavus* in NCBI's GenBank database on Jan 31st, 2023 were downloaded in
GenBank format using ncbi-genome-download (https://github.com/kblin/ncbi-genome-
download). Genomic assemblies for *Enterococcus* that met quality and taxonomic criteria for
belonging to the genus or related genera (e.g. Enterococcus_A, Enterococcus_B, etc.) in
GTDB[85] release R207 were similarly downloaded from NCBI's GenBank database using ncbi-
genome-download in FASTA format.

Application of fai and zol to identify phages within metagenomes

VIBRANT was used to identify viral contigs or sub-contigs in the three total metagenomes from Tran *et al.* 2023[54] sampled on the earliest date of 07/24. Afterwards, predicted circular contigs were clustered using BiG-SCAPE[22] which revealed a ~36 kb virus was found in two of the three metagenomes.

prepTG was run on all 16 total metagenomic assemblies from the Tran *et al.* 2023 study, performing gene calling with pyrodigal in metagenomics mode[33] to prepare for comprehensive targeted searching of the virus with fai. fai was run with default settings, with filtering of secondary instances of the phage requested to retain only the best matching scaffold or scaffold segment resembling the queries.

Microevolutionary investigations of leporin and aflatoxin BGCs in *Aspergillus flavus*

Genomic assemblies downloaded from NCBI GenBank were processed using prepTG. Of the 217 genomic assemblies downloaded, one, GCA_000006275.3, was dropped from the analysis because the original GenBank had multiple CDS features with the same name, leading to difficulties in performing BGC prediction with antiSMASH[119], and because alternate assemblies were available for the isolate. prepTG was run on all assemblies with miniprot[35] based gene-mapping of the high-quality gene coordinate predictions available for *A. flavus* NRRL 3357 (GCA_009017415.1)[64] requested. Target genomes were then searched for the leporin (BGC0001445) and aflatoxin (BGC0000008) BGCs using GenBank files downloaded from MIBiGv3[31] as queries. For leporin, AFLA_066840, as represented in the MIBiG database, was treated as a key protein required for detection of the BGC. Similarly, for aflatoxin, PksA (AAS90022.1), as represented in the MIBiG database, was treated as a key protein required for detection of the BGC. Draft-mode and filtering of paralogous segments was requested. For both analyses, ortholog groups found in fewer than 5% of gene cluster instances were disregarded.

We reidentified population B as previously delineated[11] using k-mer based ANI estimation[120] and neighbor-joining tree construction[121]. A discrete clade (n=81) in the tree was validated to feature all isolates previously determined as part of population B[11] and thus regarded as such.

For comprehensive and *de novo* BGC prediction, antiSMASH was run on the 216 genomic assemblies with 'glimmerhmm' requested for the option '--genefinding-tool'. BGCs were clustered using default settings in BiG-SCAPE with MIBiG reference BGC integration requested and a PKS-NRPS hybrid GCF was found to feature the leporin B BGC representative (BGC0001445). Only 65 (30.1%) of the 216 genomic assemblies featured this GCF, likely resulting from the use of distant gene models based on *Cryptococcus* genomes with glimmerhmm[122]. For remote cblaster[21] analysis, CAGECAT[93] was used to search NCBI's nr database with proteins from the leporin BGC representative (BGC0001445) provided as a query. Only 13 scaffolds, belonging to 12 assemblies (including GCA_000006275.3), were identified.

Evolutionary investigations of the *epa* locus across *Enterococcus*

All *Enterococcus* genomes represented in GTDB R207[85] (n=5,291) were downloaded using
ncbi-genome-download and processed in prepTG with gene-calling performed using pyrodigal[33].
Coordinates extending from 2,071,671 to 2,115,174 along the *E. faecalis* V583 chromosome,
corresponding to genes EF2164 to EF2200, were used as a query for the *epa* locus in fai to
identify homologous instances in target genomes. When using direct coordinates along a
reference, fai reperforms gene-calling along the reference using pyrodigal and extracts a local
GenBank corresponding to the region between the coordinates. Because prior comparative
analyses had shown that gene-conservation and gene-order can be slightly variable between
epa loci from *E. faecalis* and *E. faecium*[71], we relaxed the syntenic similarity requirement of
candidate regions in target genomes to the query in fai from 0.6 to 0.0 and minimum percentage
of query proteins needed to report a homologous instance of the *epa* locus to 10%. Instead, we
required the presence of 50% of key *epa* proteins found in both *E. faecalis* and *E. faecium*,
epaABCDEFGHLMOPQR, for the identification of valid homologous instances of the *epa* locus.
The E-value cutoff to determine presence for the key *epa* proteins was lowered from 1e-20 to
1e-10 to be inclusive of shorter genes and account for high levels of sequence divergence
observed across the *Enterococcus* genus. To gather auxiliary genes flanking the core *epa*
region in target genomes, we further requested the inclusion of CDS features found within 20 kb
of the boundary genes in detected instances of the *epa* locus within the resulting GenBank files
produced by fai.

Representative genome selection for visualizing epa *gene detection using fai across the*
Enterococcus *genus*: Genome-wide dereplication of all *Enterococcus* genomes using dRep[90]
with fastANI[91] and a secondary ANI clustering threshold of 99.0% led to the identification of 463
distinct genomes, including 101 *E. faecalis* genomes (75 of which were high-quality *epa*
instances that were not located near scaffold edges). A phylogeny of these 463 representative
genomes was constructed using GToTree[123] based on 119 inferred single-copy genes for
Firmicutes and FastTree2[113].

Genome selection for comparing ortholog grouping of proteins by zol with OrthoFinder:
Another round of dereplication of the 463 representative *Enterococcus* genomes was performed
using dRep[90] with fastANI[91] and a coarser secondary ANI clustering threshold of 95.0% to
approximate selection of representative genomes from different species[124]. This led to the
identification of 89 representative genomes, 42 of which had high-quality *epa* instances that
were not located near scaffold edges. Each of the 42 genomes belonged to a different species
based on taxonomic classifications in GTDB R207[85]. zol was run on the 42 high-quality *epa*
instances using default ortholog grouping parameters. Similarly, OrthoFinder v2.5.4 was run
using default settings on the full, genome-wide set of 42 proteomes. Phylogenetic hierarchical
orthogroups inferred by OrthoFinder were used for comparison with zol determined ortholog
groups.

Phylogenetic analysis of glycosyltransferases found in or near the epa locus: Ortholog groups
from the zol analysis on the 42 representative and 2,442 comprehensive multi-species *epa*
instances (Figure 5CD), as well as the 75 representative *E. faecalis epa* instances (Figure S4),
were identified as glycosyltransferases if they featured the key words: "glycosyl" and
"transferase" in Pfam protein domain annotations[125]. For Figure S4, the ortholog group
corresponding to *epaC* was not determined as a glycosyl transferase based on Pfam protein
domain annotations at the default required E-value threshold and thus manually included in the
analysis. For each gene cluster set, protein sequences belonging to the ortholog groups were
extracted, retaining association information with particular ortholog groups, and subsequently
aligned using MUSCLE[118]. Alignment filtering was next performed using trimal with options "-
keepseqs -gt 0.9", sequences with $\geq$25% of sites being gaps were filtered, and an approximate
maximum-likelihood phylogeny was finally constructed using FastTree2[113]. The phylogeny was
midpoint rooted and visualized using iTol[126]. Ortholog groups were assigned to specific *epa*
gene designations based on BLASTp sequence alignment of their consensus sequences to *E.*
faecalis V583 *epa*-associated proteins. Only the best matching ortholog group for each *epa*
gene was assigned the gene designation and only alignments with E-values of at most 1e-50
were considered.
547
Assessing the impact of dereplication on the calculation of evolutionary statistics computed by
zol: To assess the impact of dereplication on the estimation of evolutionary statistics using zol,
we focused on high-quality instances (<10% of bases ambiguous) of the *epa* locus that were not
near scaffold edges from *E. faecalis* genomes. We ran dereplication at the genome scale using
dRep[90] with fastANI[91] and a secondary ANI clustering threshold of 99.0% and dereplication at
the gene cluster scale using skani[45] at 99.0% identity and 99.0% coverage with single-linkage
clustering. We additionally simulated comprehensive processing of all high-quality gene clusters
distant from scaffold edges using the re-inflation option in zol, which allows expansion of
ortholog groups determined in the dereplicated gene cluster set to the full listing of gene
clusters. Comparisons of estimates for various evolutionary statistics by zol between the
different dereplication approaches were performed by first identifying the best matching ortholog
groups from the three distinct analyses to each *epa*-associated gene from EF2164 to EF2200 in
the *E. faeacalis* V583 reference genome based on E-value. Only ortholog groups which were
found in single-copy within the *epa* context were considered.
562
563 **Figure Legends**
564
Figure 1: Overviews of fai and zol. A) A schematic of how prepTG, fai, and zol are integrated
to perform evolutionary investigations by searching for gene clusters. An overview of the
prepTG (**B**), fai (**C**) and zol (**D**) algorithms and workflows.
Figure 2: Targeted viral detection in metagenomes using fai. A) Total metagenomes from a
single site in Lake Mendota across multiple depths and timepoints from Tran et al. 2023 were
investigated using fai for the presence of a virus found in two of the three earliest microbiome
samplings (red box). The presence of the virus is indicated by a phage icon. Metagenome
samples are colored according to whether they corresponded to oxic, oxycline, or anoxic. The
most shallow sampling depths varied for different dates and consolidated as a single row
corresponding to a sampling depth of either 5 or 10 meters. **B**) The pangenome of the virus is
shown based on the consensus order and directionality of coding sequences inferred by zol. Bar
heights correspond to the median length of coding sequences and are colored based on the

percentages of the nine metagenomes the virus was detected in. Figure 2a was created with
BioRender.com.

Figure 3: Evolutionary trends of common BGCs in *A. flavus*. A) The proportion of 216 *A.*
flavus genomes from NCBI's GenBank database with coding-sequence predictions available. **B)**
Comparison of the sensitivity of fai and alternate approaches based on assemblies for detecting
the leporin BGC. The red line indicates the total number of genomes (n=216) assessed. A
schematic of the (**C**) leporin and (**D**) aflatoxin BGCs is shown with genes present in $\geq$ 10% of
samples shown in consensus order and relative directionality. Coloring of genes in (**C**)
corresponds to FST values and in (**D**) to Tajima's D values, as calculated by zol. Grey bars in
the legends, at (**C**) 0.92 and (**D**) -0.98, indicate the mean values for the statistics across genes
in the BGC. *For the leporin BGC, *lepB* corresponds to an updated open-reading frame (ORF)
prediction by Skerker *et al.* 2021 which was the combination of AFLA_066860 and
AFLA_066870 ORFs in the MIBiG entry BGC0001445 used as the query for fai. For the
aflatoxin BGC, ORFs which were not represented in the MIBiG entry BGC0000008 but
predicted to be within the aflatoxin BGC by mapping of gene-calls from *A. flavus* NRRL 3357 by
Skerker et al. 2021 are shown in gold. The major allele frequency distributions are shown for (**E**)
aflX and (**F**) *pksA*, which depict opposite trends in sequence conservation according to their
respective Tajima's D calculations.

Figure 4: The *epa* locus is conserved across most enterococcal species. The distribution
of the *epa* locus and associated genes, based on criteria used for running fai, is shown across
463 representative *Enterococcus* genomes. Coloring of the heatmap corresponds to the
normalized bitscore of the best alignment to coding sequences from *E. faecalis* V583.
599
Figure 5: Assessment of gene cluster restricted ortholog grouping by fai and zol. A) zol
gene cluster constricted ortholog group predictions for *epa* locus proteins from 42 distinct
representative enterococcal species were compared to genome-wide predictions of ortholog
groups by OrthoFinder. **B)** A schematic of the *epa* gene cluster from *E. faecalis* V583 (from
EF2164 to EF2200) with glycosyltransferase encoding genes shown in color. A phylogeny
based on gap-filtered protein alignments of conserved ortholog with domains featuring "glycosyl"
and "transferase" as key words is shown from (**C**) *epa* loci in the 42 representative genomes
and (**D**) a more comprehensive set of 2,442 *epa* loci. Each leaf represents a specific protein and
coloring of the track corresponds to their ortholog group designations by zol. Coloring is only
shown for ortholog groups which are the best matches for each of the *epa*-associated
glycosyltransferases from *E. faecalis* V583. Note, for (**C**) 7 (2.0%) and (**D**) 70 (0.3%) proteins
were removed prior to phylogeny construction due to an abundance of gaps in the trimmed
alignment.
613
614   **Figure 6: Effects of dereplication on the calculation of evolutionary statistics by zol.** The
615   heatmap shows the correlation of values for analogous ortholog groups for various evolutionary
616   statistics computed by zol when different approaches to dereplication are used. See Methods
617   for further details. *To simulate no dereplication, gene cluster dereplication with re-inflation
618   parameters were used in zol.
619
Figure 7: Distribution of the *epa* locus and associated genes across the genus of
***Enterococcus*.** **A**) A schematic is shown for the *epa* locus in *E. faecalis* for genes which were
found in $\geq$ 25% of 75 representative genomes for the species, presented in consensus order
with consensus directionality as inferred by zol. Only ortholog groups which are predicted to
occur between and including EF2198 (*epaA*) and EF2164 based on their consensus order as
calculated by zol are shown. The coloring corresponds to the conservation of individual genes.

Genes upstream and/or including *epaR* were recently proposed to be involved in decoration of
Epa by Guerardel *et al.* 2020. "//" indicates that the ortholog group was not single-copy in the
context of the gene cluster. The tracks below the gene showcase their sequence similarity
across the *E. faecalis* genomes measured using (**B**) Tajima's D and (**C**) the average sequence
alignment entropy. **D**) The major allele frequency is depicted across the alignment for the
ortholog group featuring *epaX*. Sites predicted to be under negative selection by FUBAR,
Prob($\alpha > \beta$) $\geq$ 0.9, are marked in red. **E**) An approximate maximum-likelihood phylogeny based
on gap-filtered codon alignments for the ortholog group corresponding to *epaX* and *epaX*-like
proteins in the joint *E. faecalis* and *E. faecium* investigation of the epa locus using zol. **F**)
Barplot showcasing the percentage of high-quality *epa* gene clusters identified in *E. faecalis* and
E. faecium with the *epaX*-like ortholog group detected. Coloring of the bars corresponds to the
copy-count of the *epaX*-like ortholog group.
638
639 **References**
640

1. Snyder, L., Henkin, T. M., Peters, J. E. & Champness, W. Molecular Genetics of Bacteria,

4th Edition. Preprint at https://doi.org/10.1128/9781555817169 (2013).

2. Price, M. N., Arkin, A. P. & Alm, E. J. The life-cycle of operons. *PLoS Genet.* **2**, e96 (2006).

3. Ptashne, M. *A genetic switch: Gene control and phage. lambda.* (Palo Alto, CA (US);

Blackwell Scientific Publications, 1986).

4. Andreu, V. P. *et al.* gutSMASH predicts specialized primary metabolic pathways from the

human gut microbiota. *Nature Biotechnology* Preprint at https://doi.org/10.1038/s41587-

023-01675-1 (2023).

5. Fischbach, M. A., Walsh, C. T. & Clardy, J. The evolution of gene collectives: How natural

selection drives chemical innovation. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 4601–4608

(2008).

6. Gal-Mor, O. & Finlay, B. B. Pathogenicity islands: a molecular toolbox for bacterial

virulence. *Cell. Microbiol.* **8**, 1707–1719 (2006).

7. Kaper, J. B., Nataro, J. P. & Mobley, H. L. Pathogenic Escherichia coli. *Nat. Rev. Microbiol.*

2, 123–140 (2004).

8. Bolwell, G. P. & Paul Bolwell, G. Biochemistry & Molecular Biology of Plants.

Phytochemistry vol. 58 185 Preprint at https://doi.org/10.1016/s0031-9422(01)00095-4

(2001).

659 9. Lindahl, L. & Zengel, J. M. Operon-specific regulation of ribosomal protein synthesis in

660   Escherichia coli. *Proc. Natl. Acad. Sci. U. S. A.* **76**, 6542–6546 (1979).

661 10. Cordero, O. X. & Polz, M. F. Explaining microbial genomic diversity in light of evolutionary

662   ecology. *Nat. Rev. Microbiol.* **12**, 263–273 (2014).

663 11. Drott, M. T. *et al.* Microevolution in the pansecondary metabolome of Aspergillus flavus and

664   its potential macroevolutionary implications for filamentous fungi. *Proc. Natl. Acad. Sci. U.*

665   *S. A.* **118**, (2021).

666 12. Salamzade, R. *et al.* Evolutionary investigations of the biosynthetic diversity in the skin

667   microbiome using lsaBGC. *Microb Genom* **9**, (2023).

668 13. Tatusov, R. L., Galperin, M. Y., Natale, D. A. & Koonin, E. V. The COG database: a tool for

669   genome-scale analysis of protein functions and evolution. *Nucleic Acids Res.* **28**, 33–36

670   (2000).

671 14. Remm, M., Storm, C. E. & Sonnhammer, E. L. Automatic clustering of orthologs and in-

672   paralogs from pairwise species comparisons. *J. Mol. Biol.* **314**, 1041–1052 (2001).

673 15. Li, L., Stoeckert, C. J., Jr & Roos, D. S. OrthoMCL: identification of ortholog groups for

674   eukaryotic genomes. *Genome Res.* **13**, 2178–2189 (2003).

675 16. Edwards, D. J. & Holt, K. E. Beginner's guide to comparative bacterial genome analysis

676   using next-generation sequence data. *Microb. Inform. Exp.* **3**, 2 (2013).

677 17. Page, A. J. *et al.* Roary: rapid large-scale prokaryote pan genome analysis. *Bioinformatics*

678   **31**, 3691–3693 (2015).

679 18. Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative

680   genomics. *Genome Biol.* **20**, 238 (2019).

681 19. Medema, M. H., Takano, E. & Breitling, R. Detecting sequence homology at the gene

682   cluster level with MultiGeneBlast. *Mol. Biol. Evol.* **30**, 1218–1223 (2013).

683 20. Abby, S. S., Néron, B., Ménager, H., Touchon, M. & Rocha, E. P. C. MacSyFinder: a

684   program to mine genomes for molecular systems with an application to CRISPR-Cas

systems. *PLoS One* **9**, e110726 (2014).

21. Gilchrist, C. L. M. *et al.* Cblaster: A remote search tool for rapid identification and

visualization of homologous gene clusters. *Bioinformatics Advances* **1**, (2021).

22. Navarro-Muñoz, J. C. *et al.* A computational framework to explore large-scale biosynthetic

diversity. *Nat. Chem. Biol.* **16**, 60–68 (2020).

23. Georgescu, C. H. *et al.* SynerClust: a highly scalable, synteny-aware orthologue clustering

tool. *Microb Genom* **4**, (2018).

24. Tonkin-Hill, G. *et al.* Producing polished prokaryotic pangenomes with the Panaroo

pipeline. *Genome Biol.* **21**, 180 (2020).

25. Cosentino, S. & Iwasaki, W. SonicParanoid: fast, accurate and easy orthology inference.

Bioinformatics **35**, 149–151 (2019).

26. Hu, X. & Friedberg, I. SwiftOrtho: A fast, memory-efficient, multiple genome orthology

classifier. *Gigascience* **8**, (2019).

27. Melnyk, R. A., Hossain, S. S. & Haney, C. H. Convergent gain and loss of genomic islands

drive lifestyle changes in plant-associated Pseudomonas. *ISME J.* **13**, 1575–1588 (2019).

28. Vallenet, D. *et al.* MaGe: a microbial genome annotation system supported by synteny

results. *Nucleic Acids Res.* **34**, 53–65 (2006).

29. Stam, M. *et al.* NetSyn: genomic context exploration of protein families. *bioRxiv* (2023)

doi:10.1101/2023.02.15.528638.

30. Liu, M. *et al.* ICEberg 2.0: an updated database of bacterial integrative and conjugative

elements. *Nucleic Acids Res.* **47**, D660–D665 (2019).

31. Terlouw, B. R. *et al.* MIBiG 3.0: a community-driven effort to annotate experimentally

validated biosynthetic gene clusters. *Nucleic Acids Res.* **51**, D603–D610 (2023).

32. Bertelli, C. *et al.* IslandViewer 4: expanded prediction of genomic islands for larger-scale

datasets. *Nucleic Acids Res.* **45**, W30–W35 (2017).

33. Larralde, M. Pyrodigal: Python bindings and interface to Prodigal, an efficient method for

711  gene prediction in prokaryotes. *J. Open Source Softw.* **7**, 4296 (2022).

712 34. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site

713  identification. *BMC Bioinformatics* **11**, 119 (2010).

714 35. Li, H. Protein-to-genome alignment with miniprot. *Bioinformatics* **39**, (2023).

715 36. Grazziotin, A. L., Koonin, E. V. & Kristensen, D. M. Prokaryotic Virus Orthologous Groups

716  (pVOGs): a resource for comparative genomics and protein family annotation. *Nucleic*

717  *Acids Res.* **45**, D491–D498 (2017).

718 37. Liu, B., Zheng, D., Jin, Q., Chen, L. & Yang, J. VFDB 2019: a comparative pathogenomic

719  platform with an interactive web interface. *Nucleic Acids Res.* **47**, D687–D692 (2019).

720 38. Siguier, P., Perochon, J., Lestrade, L., Mahillon, J. & Chandler, M. ISfinder: the reference

721  centre for bacterial insertion sequences. *Nucleic Acids Res.* **34**, D32-6 (2006).

722 39. Alcock, B. P. *et al.* CARD 2023: expanded curation, support for machine learning, and

723  resistome prediction at the Comprehensive Antibiotic Resistance Database. *Nucleic Acids*

724  *Res.* **51**, D690–D699 (2023).

725 40. Kosakovsky Pond, S. L. *et al.* HyPhy 2.5—A Customizable Platform for Evolutionary

726  Hypothesis Testing Using Phylogenies. *Mol. Biol. Evol.* **37**, 295–299 (2019).

727 41. Thorpe, H. A., Bayliss, S. C., Sheppard, S. K. & Feil, E. J. Piggy: a rapid, large-scale pan-

728  genome analysis tool for intergenic regions in bacteria. *Gigascience* **7**, 1–11 (2018).

729 42. Gilchrist, C. L. M. & Chooi, Y.-H. clinker & clustermap.js: automatic generation of gene

730  cluster comparison figures. *Bioinformatics* **37**, 2473–2475 (2021).

731 43. Hackl, T., Duponchel, S., Barenhoff, K., Weinmann, A. & Fischer, M. G. Virophages and

732  retrotransposons colonize the genomes of a heterotrophic flagellate. *Elife* **10**, (2021).

733 44. Sullivan, M. J., Petty, N. K. & Beatson, S. A. Easyfig: a genome comparison visualizer.

734  *Bioinformatics* **27**, 1009–1010 (2011).

735 45. Shaw, J. & Yu, Y. W. Fast and robust metagenomic sequence comparison through sparse

736  chaining with skani. *bioRxiv* 2023.01.18.524587 (2023) doi:10.1101/2023.01.18.524587.

46. Blackwell, G. *et al.* Exploring bacterial diversity via a curated and searchable snapshot of

archived DNA. *Access Microbiol.* **4**, (2022).

47. Lebreton, F. *et al.* Emergence of epidemic multidrug-resistant Enterococcus faecium from

animal and commensal strains. *MBio* **4**, (2013).

48. Lieberman, T. D. *et al.* Genetic variation of a bacterial pathogen within individuals with

cystic fibrosis provides a record of selective pressures. *Nat. Genet.* **46**, 82–87 (2014).

49. Crits-Christoph, A., Olm, M. R., Diamond, S., Bouma-Gregson, K. & Banfield, J. F. Soil

bacterial populations are shaped by recombination and gene-specific selection across a

grassland meadow. *ISME J.* **14**, 1834–1846 (2020).

50. Hudson, R. R., Slatkin, M. & Maddison, W. P. Estimation of levels of gene flow from DNA

sequence data. *Genetics* **132**, 583–589 (1992).

51. Tran, P. Q. & Anantharaman, K. Biogeochemistry Goes Viral: towards a Multifaceted

Approach To Study Viruses and Biogeochemical Cycling. *mSystems* **6**, e0113821 (2021).

52. Barr, J. J. *et al.* Bacteriophage adhering to mucus provide a non-host-derived immunity.

Proc. Natl. Acad. Sci. U. S. A. **110**, 10771–10776 (2013).

53. Lefeuvre, P. *et al.* Evolution and ecology of plant viruses. *Nat. Rev. Microbiol.* **17**, 632–644

(2019).

54. Tran, P. Q. *et al.* Viral impacts on microbial activity and biogeochemical cycling in a

seasonally anoxic freshwater lake. *bioRxiv* 2023.04.19.537559 (2023)

doi:10.1101/2023.04.19.537559.

55. Kieft, K., Zhou, Z. & Anantharaman, K. VIBRANT: automated recovery, annotation and

curation of microbial viruses, and evaluation of viral community function from genomic

sequences. *Microbiome* **8**, 90 (2020).

56. Bok, J. W. *et al.* Genomic mining for Aspergillus natural products. *Chem. Biol.* **13**, 31–37

(2006).

57. Vadlapudi, V. *et al.* Aspergillus Secondary Metabolite Database, a resource to understand

the Secondary metabolome of Aspergillus genus. *Sci. Rep.* **7**, 7325 (2017).

58.   Robey, M. T., Caesar, L. K., Drott, M. T., Keller, N. P. & Kelleher, N. L. An interpreted atlas

of biosynthetic gene clusters from 1,000 fungal genomes. *Proc. Natl. Acad. Sci. U. S. A.*

118, (2021).

59.   Hatmaker, E. A. *et al.* Genomic and Phenotypic Trait Variation of the Opportunistic Human

Pathogen Aspergillus flavus and Its Close Relatives. *Microbiol Spectr* **10**, e0306922 (2022).

60.   Xie, H. *et al.* Global multi-omics profiling reveals evolutionary drivers of phylogeographic

diversity of fungal specialized metabolism. (2023) doi:10.21203/rs.3.rs-2471999/v1.

61.   Cary, J. W. *et al.* An Aspergillus flavus secondary metabolic gene cluster containing a

hybrid PKS-NRPS is necessary for synthesis of the 2-pyridones, leporins. *Fungal Genet.*

Biol. **81**, 88–97 (2015).

62.   Drăgan, M.-A., Moghul, I., Priyam, A., Bustos, C. & Wurm, Y. GeneValidator: identify

problems with protein-coding gene predictions. *Bioinformatics* **32**, 1559–1561 (2016).

63.   Scalzitti, N., Jeannin-Girardon, A., Collet, P., Poch, O. & Thompson, J. D. A benchmark

study of ab initio gene prediction methods in diverse eukaryotic organisms. *BMC Genomics*

21, 293 (2020).

64.   Skerker, J. M. *et al.* Chromosome assembled and annotated genome sequence of

Aspergillus flavus NRRL 3357. *G3* **11**, jkab213 (2021).

65.   Yang, K., Tian, J. & Keller, N. P. Post-translational modifications drive secondary

metabolite biosynthesis in Aspergillus: a review. *Environ. Microbiol.* **24**, 2857–2881 (2022).

66.   Klich, M. A. Aspergillus flavus: the major producer of aflatoxin. *Mol. Plant Pathol.* **8**, 713–

722 (2007).

67.   Cary, J. W., Ehrlich, K. C., Bland, J. M. & Montalbano, B. G. The Aflatoxin Biosynthesis

Cluster Gene, *aflX* , Encodes an Oxidoreductase Involved in Conversion of Versicolorin A

to Demethylsterigmatocystin. *Applied and Environmental Microbiology* vol. 72 1096–1101

Preprint at https://doi.org/10.1128/aem.72.2.1096-1101.2006 (2006).

68. Cleveland, T. E. *et al.* Potential of Aspergillus flavus genomics for applications in

biotechnology. *Trends Biotechnol.* **27**, 151–157 (2009).

69. Ehrlich, K. C., Li, P., Scharfenstein, L. & Chang, P.-K. HypC, the anthrone oxidase involved

in aflatoxin biosynthesis. *Appl. Environ. Microbiol.* **76**, 3374–3377 (2010).

70. Xu, Y., Murray, B. E. & Weinstock, G. M. A cluster of genes involved in polysaccharide

biosynthesis from Enterococcus faecalis OG1RF. *Infect. Immun.* **66**, 4313–4323 (1998).

71. Palmer, K. L. *et al.* Comparative Genomics of Enterococci: Variation in Enterococcus

faecalis, Clade Structure in E. faecium, and Defining Characteristics of *E . gallinarum* and *E*

. casseliflavus. mBio vol. 3 Preprint at https://doi.org/10.1128/mbio.00318-11 (2012).

72. Hancock, L. E., Murray, B. E. & Sillanpää, J. Enterococcal Cell Wall Components and

Structures. in *Enterococci: From Commensals to Leading Causes of Drug Resistant*

Infection (eds. Gilmore, M. S., Clewell, D. B., Ike, Y. & Shankar, N.) (Massachusetts Eye

and Ear Infirmary, 2014).

73. Qin, X. *et al.* Complete genome sequence of Enterococcus faecium strain TX16 and

comparative genomic analysis of Enterococcus faecium genomes. *BMC Microbiol.* **12**, 135

(2012).

74. Teng, F., Jacques-Palaz, K. D., Weinstock, G. M. & Murray, B. E. Evidence that the

Enterococcal Polysaccharide Antigen Gene (*epa*) Cluster Is Widespread in *Enterococcus*

faecalis and Influences Resistance to Phagocytic Killing of *E. faecalis. Infection and*

Immunity vol. 70 2010–2015 Preprint at https://doi.org/10.1128/iai.70.4.2010-2015.2002

(2002).

75. Teng, F., Singh, K. V., Bourgogne, A., Zeng, J. & Murray, B. E. Further Characterization of

the *epa* Gene Cluster and Epa Polysaccharides of *Enterococcus faecalis. Infection and*

Immunity vol. 77 3759–3767 Preprint at https://doi.org/10.1128/iai.00149-09 (2009).

76. Rigottier-Gois, L. *et al.* The surface rhamnopolysaccharide epa of Enterococcus faecalis is

a key determinant of intestinal colonization. *J. Infect. Dis.* **211**, 62–71 (2015).

77. Guerardel, Y. *et al.* Complete structure of the enterococcal polysaccharide antigen (EPA) of

vancomycin-resistant Enterococcus faecalis V583 reveals that EPA decorations are

teichoic acids covalently linked to a rhamnopolysaccharide backbone. *MBio* **11**, (2020).

78. Smith, R. E. *et al.* Decoration of the enterococcal polysaccharide antigen EPA is essential

for virulence, cell surface charge and interaction with effectors of the innate immune

system. *PLoS Pathog.* **15**, e1007730 (2019).

79. Singh, K. V. & Murray, B. E. Loss of a Major Enterococcal Polysaccharide Antigen (Epa) by

Enterococcus faecalis Is Associated with Increased Resistance to Ceftriaxone and

Carbapenems. *Antimicrob. Agents Chemother.* **63**, (2019).

80. Ho, K., Huo, W., Pas, S., Dao, R. & Palmer, K. L. Loss-of-Function Mutations in *epaR*

Confer Resistance to ☐NPV1 Infection in Enterococcus faecalis OG1RF. *Antimicrobial*

Agents and Chemotherapy vol. 62 Preprint at https://doi.org/10.1128/aac.00758-18 (2018).

81. Fiore, E., Van Tyne, D. & Gilmore, M. S. Pathogenicity of Enterococci. *Microbiol Spectr* **7**,

(2019).

82. Lebreton, F., Willems, R. J. L. & Gilmore, M. S. Enterococcus Diversity, Origins in Nature,

and Gut Colonization. in *Enterococci: From Commensals to Leading Causes of Drug*

Resistant Infection (eds. Gilmore, M. S., Clewell, D. B., Ike, Y. & Shankar, N.)

(Massachusetts Eye and Ear Infirmary, 2014).

83. Lebreton, F. *et al.* Tracing the Enterococci from Paleozoic Origins to the Hospital. *Cell* **169**,

849-861.e13 (2017).

84. Schwartzman, J. A. *et al.* Global diversity of enterococci and description of 18 novel

species. *bioRxiv* 2023.05.18.540996 (2023) doi:10.1101/2023.05.18.540996.

85. Parks, D. H. *et al.* GTDB: an ongoing census of bacterial and archaeal diversity through a

phylogenetically consistent, rank normalized and complete genome-based taxonomy.

Nucleic Acids Res. (2021) doi:10.1093/nar/gkab776.

86. Reid, K. C., Cockerill, F. R., III & Patel, R. Clinical and epidemiological features of

Enterococcus casseliflavus/flavescens and Enterococcus gallinarum bacteremia: a report of

20 cases. *Clin. Infect. Dis.* **32**, 1540–1546 (2001).

87. Monticelli, J., Knezevich, A., Luzzati, R. & Di Bella, S. Clinical management of non-faecium

non-faecalis vancomycin-resistant enterococci infection. Focus on Enterococcus gallinarum

and Enterococcus casseliflavus/flavescens. *J. Infect. Chemother.* **24**, 237–246 (2018).

88. Nevers, Y. *et al.* The Quest for Orthologs orthology benchmark service in 2022. *Nucleic*

Acids Res. (2022).

89. Dale, J. L., Cagnazzo, J., Phan, C. Q., Barnes, A. M. T. & Dunny, G. M. Multiple Roles for

Enterococcus faecalis Glycosyltransferases in Biofilm-Associated Antibiotic Resistance,

Cell Envelope Integrity, and Conjugative Transfer. *Antimicrobial Agents and Chemotherapy*

vol. 59 4094–4105 Preprint at https://doi.org/10.1128/aac.00344-15 (2015).

90. Olm, M. R., Brown, C. T., Brooks, B. & Banfield, J. F. dRep: a tool for fast and accurate

genomic comparisons that enables improved genome recovery from metagenomes through

de-replication. *ISME J.* **11**, 2864–2868 (2017).

91. Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High

throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries.

Nat. Commun. **9**, 1–8 (2018).

92. Murrell, B. *et al.* FUBAR: a fast, unconstrained bayesian approximation for inferring

selection. *Mol. Biol. Evol.* **30**, 1196–1205 (2013).

93. van den Belt, M. *et al.* CAGECAT: The CompArative GEne Cluster Analysis Toolbox for

rapid search and visualisation of homologous gene clusters. *BMC Bioinformatics* **24**, 181

(2023).

94. Bertelli, C. *et al.* Enabling genomic island prediction and comparison in multiple genomes to

investigate bacterial evolution and outbreaks. *Microb. Genom.* **8**, (2022).

95. Bin Jang, H. *et al.* Taxonomic assignment of uncultivated prokaryotic virus genomes is

enabled by gene-sharing networks. *Nat. Biotechnol.* **37**, 632–639 (2019).

96. Salamzade, R. *et al.* Inter-species geographic signatures for tracing horizontal gene

transfer and long-term persistence of carbapenem resistance. *Genome Med.* **14**, 37 (2022).

97. Sheppard, A. E. *et al.* Nested Russian Doll-Like Genetic Mobility Drives Rapid

Dissemination of the Carbapenem Resistance Gene blaKPC. *Antimicrob. Agents*

Chemother. **60**, 3767–3778 (2016).

98. Groussin, M. *et al.* Elevated rates of horizontal gene transfer in the industrialized human

microbiome. *Cell* **184**, 2053-2067.e18 (2021).

99. Crits-Christoph, A., Diamond, S., Butterfield, C. N., Thomas, B. C. & Banfield, J. F. Novel

soil bacteria possess diverse genes for secondary metabolite biosynthesis. *Nature* **558**,

440–444 (2018).

100.Bickhart, D. M. *et al.* Generation of lineage-resolved complete metagenome-assembled

genomes by precision phasing. *bioRxiv* 2021.05.04.442591 (2021)

doi:10.1101/2021.05.04.442591.

101.van Tilburg Bernardes, E., Charron-Mazenod, L., Reading, D. J., Reckseidler-Zenteno, S.

L. & Lewenza, S. Exopolysaccharide-repressing small molecules with antibiofilm and

antivirulence activity against Pseudomonas aeruginosa. *Antimicrob. Agents Chemother.* **61**,

(2017).

102.Shen, Y. & Loessner, M. J. Beyond antibacterials - exploring bacteriophages as

antivirulence agents. *Curr. Opin. Biotechnol.* **68**, 166–173 (2021).

103.Shankar-Sinha, S. *et al.* The Klebsiella pneumoniae O antigen contributes to bacteremia

and lethality during murine pneumonia. *Infect. Immun.* **72**, 1423–1430 (2004).

104.Chatterjee, A. *et al.* Bacteriophage Resistance Alters Antibiotic-Mediated Intestinal

Expansion of Enterococci. *Infect. Immun.* **87**, (2019).

105.Chatterjee, A. *et al.* Parallel genomics uncover novel enterococcal-bacteriophage

interactions. Preprint at https://doi.org/10.1101/858506.

106.Canfield, G. S. *et al.* Lytic bacteriophages facilitate antibiotic sensitization of *Enterococcus*

893   *faecium*. Preprint at https://doi.org/10.1101/2020.09.22.309401.

894 107. Kirsch, J. M. *et al.* Targeted IS-element sequencing uncovers transposition dynamics

895   during selective pressure in enterococci. *PLoS Pathog.* **19**, e1011424 (2023).

896 108. Zhao, S. *et al.* Adaptive evolution within gut microbiomes of healthy people. *Cell Host*

897   *Microbe* **25**, 656-667.e8 (2019).

898 109. Grüning, B. *et al.* Bioconda: sustainable and comprehensive software distribution for the life

899   sciences. *Nat. Methods* **15**, 475–476 (2018).

900 110. Salamzade, R. & Kalan, L. *zol.* (Zenodo, 2023). doi:10.5281/ZENODO.8132329.

901 111. Cock, P. J. A. *et al.* Biopython: freely available Python tools for computational molecular

902   biology and bioinformatics. *Bioinformatics* **25**, 1422–1423 (2009).

903 112. Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: a tool for automated

904   alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973

905   (2009).

906 113. Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2--approximately maximum-likelihood

907   trees for large alignments. *PLoS One* **5**, e9490 (2010).

908 114. Huang, Y., Niu, B., Gao, Y., Fu, L. & Li, W. CD-HIT Suite: a web server for clustering and

909   comparing biological sequences. *Bioinformatics* **26**, 680–682 (2010).

910 115. Eddy, S. R. Accelerated Profile HMM Searches. *PLoS Comput. Biol.* **7**, e1002195 (2011).

911 116. Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using DIAMOND.

912   *Nat. Methods* **12**, 59–60 (2014).

913 117. Schreiber, J. Pomegranate: fast and flexible probabilistic modeling in python. *J. Mach.*

914   *Learn. Res.* (2017).

915 118. Edgar, R. C. Muscle5: High-accuracy alignment ensembles enable unbiased assessments

916   of sequence homology and phylogeny. *Nat. Commun.* **13**, 6968 (2022).

917 119. Blin, K. *et al.* antiSMASH 6.0: improving cluster detection and comparison capabilities.

918   *Nucleic Acids Res.* **49**, W29–W35 (2021).

919 120. Ondov, B. D. *et al.* Mash: fast genome and metagenome distance estimation using

920   MinHash. *Genome Biol.* **17**, 132 (2016).

921 121. Paradis, E., Claude, J. & Strimmer, K. APE: Analyses of Phylogenetics and Evolution in R

922   language. *Bioinformatics* **20**, 289–290 (2004).

923 122. Majoros, W. H., Pertea, M. & Salzberg, S. L. TigrScan and GlimmerHMM: two open source

924   ab initio eukaryotic gene-finders. *Bioinformatics* **20**, 2878–2879 (2004).

925 123. Lee, M. D. GToTree: a user-friendly workflow for phylogenomics. *Bioinformatics* **35**, 4162–

926   4164 (2019).

927 124. Olm, M. R. *et al.* Consistent Metagenome-Derived Metrics Verify and Delineate Bacterial

928   Species Boundaries. *mSystems* **5**, (2020).

929 125. Mistry, J. *et al.* Pfam: The protein families database in 2021. *Nucleic Acids Res.* **49**, D412–

930   D419 (2021).

931 126. Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v4: recent updates and new

932   developments. *Nucleic Acids Res.* **47**, W256–W259 (2019).

Figure 1: Overviews of fai and zol. A) A schematic of how prepTG, fai, and zol are integrated to perform evolutionary investigations by searching for gene-clusters. An overview of the prepTG (**B**), fai (**C**) and zol (**D**) algorithms and workflows.

Figure 2: Targeted viral detection in metagenomes using fai. A) Total metagenomes from a single site in Lake Mendota across multiple depths and timepoints from Tran et al. 2023 were investigated using fai for the presence of a virus found in two of the three earliest microbiome samplings (red box). The presence of the virus is indicated by a phage icon. Metagenome samples are colored according to whether they corresponded to oxic, oxycline, or anoxic. The most shallow sampling depths varied for different dates and consolidated as a single row corresponding to a sampling depth of either 5 or 10 meters. **B**) The pangenome of the virus is shown based on the consensus order and directionality of coding sequences inferred by zol. Bar heights correspond to the median length of coding sequences and are colored based on the percentages of the nine metagenomes the virus was detected in. BioRender was used in generation of this figure.

Figure 3: Evolutionary trends of common BGCs in *A. flavus*. A) The proportion of 216 A. flavus genomes from NCBI's GenBank database with coding-sequence predictions available. **B)** Comparison of the sensitivity of fai and alternate approaches based on assemblies for detecting the leporin BGC. The red-line indicates the total number of genomes (n=216) assessed. A schematic of the (**C**) leporin and (**D**) aflatoxin BGCs is shown with genes present in ≥ 10% of samples shown in consensus order and relative directionality. Coloring of genes in (**C**) corresponds to FST values and in (**D**) to Tajima's D values, as calculated by zol. Grey bars in the legends, at (C) 0.92 and (D) -0.98, indicate the mean values for the statistics across genes in the BGC. *For the leporin BGC, *lepB* corresponds to an updated open-reading frame (ORF) prediction by Skerker *et al.* 2021 which was the combination of AFLA_066860 and AFLA_066870 ORFs in the MIBiG entry BGC0001445 used as the query for fai. For the aflatoxin BGC, ORFs which were not represented in the MIBiG entry BGC0000008 but predicted to be within the aflatoxin BGC by mapping of gene-calls from *A. flavus* NRRL 3357 by Skerker et al. 2021 are shown in gold. The major allele frequency distributions are shown for (**E**) *aflX* and (**F**) *pksA*, which depict opposite trends in sequence conservation according to their respective Tajima's D calculations.

Figure 4: The *epa* locus is conserved across most enterococcal species. The distribution of the *epa* locus and associated genes, based on criteria used for running fai, is shown across 463 representative genomes across *Enterococcus*. Coloring of the heatmap corresponds to the normalized bitscore of the best alignment to coding sequences from *E. faecalis* V583.

Figure 5: Assessment of gene cluster restricted ortholog grouping by fai and zol. A) zol gene cluster constricted ortholog group predictions for *epa* locus proteins from 42 distinct representative enterococcal species were compared to genome-wide predictions of ortholog groups by OrthoFinder. **B**) A schematic of the *epa* gene cluster from *E. faecalis* V583 (from EF2164 to EF2200) with glycosyltransferase encoding genes shown in color. A phylogeny based on gap-filtered protein alignments of conserved ortholog with domains featuring "glycosyl" and "transferase" as key words is shown from (**C**) *epa* loci in the 42 representative genomes and (**D**) a more comprehensive set of 2,442 *epa* loci. Each leaf represents a specific protein and coloring of the track corresponds to their ortholog group designations by zol. Coloring is only shown for ortholog groups which are the best matches for each of the *epa*-associated glycosyltransferases from *E. faecalis* V583. Note, for (**C**) 7 (2.0%) and (**D**) 70 (0.3%) proteins were removed prior to phylogeny construction due to an abundance of gaps in the trimmed alignment.

Figure 6: Effects of dereplication on the calculation of evolutionary statistics by zol. The heatmap shows the correlation of values for analogous ortholog groups for various evolutionary statistics computed by zol when different approaches to dereplication are used. See Methods for further details. *To simulate no dereplication, gene-cluster dereplication with re-inflation parameters were used in zol.

Figure 7: Distribution of the *epa* locus and associated genes across the genus of *Enterococcus*. A) A schematic is shown for the *epa* locus in *E. faecalis* for genes which were found in ≥ 25% of 75 representative genomes for the species presented in consensus order with consensus directionality as inferred by zol. Only ortholog groups which are predicted to occur between and including EF2198 (*epaA*) and EF2164 based on their consensus order as calculated by zol are shown. The coloring corresponds to the conservation of individual genes. Genes upstream and/or including *epaR* were recently proposed to be involved in decoration of Epa by Guerardel *et al.* 2020. "//" indicates that the ortholog group was not single-copy in the context of the gene-cluster. The tracks below the gene showcase their sequence similarity across the *E. faecalis* genomes measured using (**B**) Tajima's D and (**C**) the average sequence alignment entropy. **D**) The major allele frequency is depicted across the alignment for the ortholog group featuring *epaX*. Sites predicted to be under negative selection by FUBAR, Prob($\alpha > \beta$) ≥ 0.9, are marked in red. **E**) An approximate maximum-likelihood phylogeny based on gap-filtered codon alignments for the ortholog group corresponding to *epaX* and *epaX*-like proteins in the joint *E. faecalis* and *E. faecium* investigation of the epa locus using zol. **F**) Barplot showcasing the percentage of high-quality *epa* gene clusters identified in *E. faecalis* and *E. faecium* with the *epa*X-like ortholog group detected. Coloring of the bars corresponds to the copy-count of the *epa*X-like ortholog group.