Circular viral copy DNA of Dengue virus (cvcDENV-2) isolated from infected mosquito cell cultures

Warachin Gangnonngiw [1,2], Wilawan Thongda[1,2], Malinee Bunnontae[1],

Nipaporn Kanthong[3], T.W. Flegel[1]

[1]Center of Excellence for Shrimp Molecular Biology and Biotechnology (CENTEX Shrimp),
Faculty of Science, Mahidol University, Bangkok, 10400, Thailand

[2]National Center for Genetic Engineering and Biotechnology (BIOTEC), National Science
and Technology Development Agency (NSTDA), Pathum Thani, 12120, Thailand

[3]Department of Biotechnology, Faculty of Science and Technology, Rajamangala University
of Technology Tawan-ok, Sriracha, Chonburi, 20110, Thailand

Correspondence

Nipaporn Kanthong; E-mail: Nipaporn_ka@rmutto.ac.th

T.W. Flegel; tim.flegel@gmail.com

ABSTRACT

Dengue virus serotype 2 (DENV-2) is a mosquito-borne disease in the family *Flaviviridae*. It has been previously shown that DENV-2 can infect C6/36 mosquito cells and cause initial cytopathic effects that dissipate upon serial split-passage to yield persistently infected cultures with normal growth and morphology. In other words, the cell line accommodated persistent DENV-2 infections. It has recently been found that insect viral infections induce the production of viral copy DNA (vcDNA) fragments via host reverse transcriptase (RT). The vcDNA occurs in both linear (lvcDNA) and circular (cvcDNA) forms and produces small interfering RNA (siRNA) transcripts that can result in an immediate protective RNA interference (RNAi) response. The vcDNA can lead to the host acquiring endogenous viral elements (EVE) in genomic DNA. Thus, we hypothesized that DENV-2 cvcDNA and DENV-2-EVE would arise in C6/36 insect cells challenged with DENV-2 virus *in vitro*. Here we describe the successful isolation and characterization of cvcDNA constructs homologous to DENV-2 from laboratory challenges with C6/36 cells. At least 1 of these appeared to arise from a DENV-2-EVE. We also show that a cvcDNA preparation derived from the DENV-2 infected in C6/36 insect cells

and subjected to rolling circle amplification (RCA) significantly reduced DENV-2 replication when applied to naive C6/36 cells prior to the DENV-2 challenge. This preliminary work lays the groundwork for further studies using the C6/36 cell model to screen and characterize protective EVE against insect and shrimp viruses.

Keywords: Anti-viral immunity, Dengue virus, C6/36 mosquito cells, circular viral copy DNA (cvcDNA), endogenous viral element(s)(EVE)

1. INTRODUCTION

It was formerly believed that invertebrates were incapable of specific adaptive immune responses and depended solely on generic responses via innate immune mechanisms (Cerenius et al. 2010). It is well known that insects and crustaceans lack a mechanism for a specific adaptive immune response via antibodies. However, they do have a specific adaptive immune response against viruses via nucleic acids. A major feature of this response is persistent, relatively innocuous viral infections. The outcome of this response is usually persistent, but relatively innocuous viral infections. The response has been called viral accommodation and has been recently reviewed (Flegel 2020, 2022), as has similar work related specifically to insects (Bonning & Saleh 2021). To summarize, endogenous host reverse transcriptase (RT) recognizes and binds with viral RNA to produce viral copy DNA (vcDNA) in both linear (lvcDNA) and circular (cvcDNA) forms. This produces small interfering RNA (siRNA) that results in individual cell and systemic RNA interference (RNAi) responses. At the same time, some of the vcDNA fragments may be integrated into the cell genomic DNA as endogenous viral elements (EVE). The EVE may also produce antisense RNA, leading to an RNAi response. In addition, the EVE may give rise to cvcDNA, a feature that allows for its detetion in uninfected individuals because of the relative ease of isolating and sequencing it (Taengchaiyaphum et al. 2021).

It has been previously shown that C6/36 insect cells can be sequentially infected with *Aedes albopictus* densovirus (AaDNV), Dengue virus serotype 2 (DENV-2), and *Japanese encephalitis* virus to result in individual cells with innocuous, persistent triple infections that can be maintained by serial split passage (6). The mechanisms underlying the ability of these cells to accommodate 3 viruses without signs of cytopathic effects (CPE) or other signs of disease was unknown at that time. More recent work described in the previous paragraph suggested that induction of host RT and production of vcDNA might be involved. Thus, in this

communication, we show that cvcDNA homologous to DENV-2 can be isolated and
characterized from C6/36 cells challenged with DENV-2. We also show that DENV-2-EVE
arose in the challenged, cultured cells and that these cvcDNA fragments obtained from the
DENV-2 infected in C6/36 insect cells and subjected to rolling circle amplification (RCA)
could significantly reduce the severity of DENV-2 infection in pre-exposed, naive C6/36 cells
subsequently challenged with DENV-2.
73
2. MATERIALS AND METHODS
2.1 Cell cultures for DENV-2 infection
DENV-2 persistent cultures were established, as previously reported (Kanthong et al. 2008,
2010). Initially, naive C6/36 cells at 10^5 cells/well were cultured in 6-well plates with 2 ml
fresh Leibovitz's (L15) medium containing supplements of 10% fetal bovine serum (FBS),
0.3% tryptose phosphate broth (TPB), and 1% antibiotics and incubated for 24 hours in 28°C
incubator. After 24 hours of incubation, the old medium was removed from the C6/36 cell layer
that was rinsed with 1 ml fresh L15 medium without other supplements. The cell layers were
then challenged with DENV-2 stock at 10^5 copies/well in 1 ml L15 medium without
supplements and gentle swirling incubation at room temperature (RT= ~25 °C) for 2 hours.
After 2 hours, the 1 ml of L15 medium containing DENV-2 was removed, and 2 ml fresh L15
followed by further incubation in a 28°C incubator. Control cells were treated in the same
manner, except for no exposure to DENV-2. All cell cultures were then subjected to serial sub-
culture every 3 days for 7 passages (total 21 days). Each passage consisted of creating a 2 ml
cell suspension divided into two 1 ml portions, one for analysis and the other for ongoing cell
culture. Thus, there were 6 replicates for the test and control samples for each passage. The
collected cells were stored in a -80 °C freezer until DNA extraction.
91
2.2. Circular DNA extraction and verification
Before DNA extraction, each 1 ml of suspended cells was centrifuged at 10,000 rpm for 5 min
at room temperature to spin down cells. Cell samples from passages 1-7 were separately
extracted for the total DNA using a DNA Extraction kit (GeneAll, Korea). The total DNA
extracts of each passage were measured for DNA quantity and quality using NanoDrop™
2000/2000c Spectrophotometers (Thermo Fisher Scientific, USA). The exact amount of 2,000
ng DNA extract from each passage (7 passages) was pooled and processed to prepare circular
DNA. Total DNA was initially exposed to exonuclease V, which digests linear single-stranded-
and double-stranded DNA while circular DNA remains intact. The residual putative circular

DNA was isolated, purified, and collected using a DNA clean-up kit (NucleoSpin® Gel and PCR Clean-up Kit, Macherey-Nagel, Germany) following the manufacturers' instructions.

To verify linear DNA digestion in the exonuclease V reaction step to isolate C6/36 circular DNA, we used linear DNA derived from the pGEM vector cut with TstI restriction enzyme in the same DNA amount of the C6/36 DNA and compared this to similar samples contatining ordinary pGEM vector (circular DNA). C6/36 DNA preparations containing the linear pGEM vector gave negative PCR test results for pGEM while samples containing the circular form gave positive test results. Exonuclease V enzyme was added to the reaction until linear DNA was eliminated as evidenced by no linear DNA in gel electrophoresis (1.5% agarose gel) while the circular pGEM vector was unaffected (data not shown).

Another step to confirm the successful circular DNA extraction after the exonuclease V enzyme digestion step was negative PCR test results for C6/ 36 cell genes using specific primers (**Table 1**) via polymerase chain reaction (PCR) followed by gel electrophoresis. These included use of primers for *Aedes* elongation factor gene (EF165_*Aedes*_F/ R, Table 1). In contrast, positive PCR test results were obtained using mitochondrial (mt) DNA genes of the *Aedes* ATP Synthase and cytochrome c oxidase primers (ATPSynthase_150F/ R and CYCS_*Aedes*_125F/R, Table 1).

Table 1. Primers were used to amplify genes of C6/36 cells in this study.

Primer name	Sequence (5'-3')	Expected amplicon size
linear DNA of C6/36 cell		
EF165_Aedes_F/R	CCCCAGTAAGGAGCAAGCAG CCAGCGGGATCGGATTGTAG	165
mtDNA of C6/36 cell		
ATPSynthase_150F/R	CCTCAAGGCACCCCTCCTAT AGGGCCGGTATTTCCTAAAAGA	150
CYCS_Aedes_125F/R	TGCTACAGGATTTCATGGACTT ATCAAGCTGCTGCTTCAAAACC	125

2.3. Sequencing of cvcDNA

To verify the result of cvcDNA of DENV-2 identification, the pooled circular DNA extracts after the exonuclease V enzyme digestion were subjected to metagenomic sequencing. The first pooled circular DNA isolation was submitted directly to GENEWIZ, Inc., Suzhou, China, for library construction and sequencing without any PCR amplification step. A second pooled

circular DNA preparation was subjected to circular DNA amplification using Phi29 DNA

polymerase rolling circle amplification (RCA) (TempliPhi™ 100 Amplification Kit, Cytiva,

USA) according to the manufacturer's instructions. This second pooled circular DNA sample

was submitted to NovogeneAIT Genomics, Singapore, for library preparation and

metagenomic sequencing. The specimens submitted were in the amount (Qubit®) ≥ 200 ng,

volume ≥ 20 µl, concentration ≥ 10 ng/ µl, and purity (Nanodrop^{TM}/Agarose Gel) OD260/280

= 1.8-2.0. Both companies confirmed that there was no degradation or contamination and that

the samples passed quality and quantity requirements for DNA sequencing.

2.2 Identification of cvcDNA of DENV-2 from circular DNA sequencing

Circular DNA library preparation and sequencing were separately performed by GENEWIZ,

Inc., Suzhou, China (1st pooled DNA extracts without any PCR amplification step), and

NovogeneAIT Genomics, Singapore (2nd pooled DNA extracts after rolling circle amplification

(RCA)). A TruSeq DNA Library Preparation Kit (Illumina) was used for cDNA library

preparation, following the TruSeq protocol. Briefly, DNA was fragmented by sonic disruption

using a Covaris S220 apparatus (Covaris). Subsequently, blunt-end fragments were created,

and the A-base was tailed to each 3'side. Dual indexes within P5 and P7 adapters were ligated

to the fragments. The final products were purified. Then, the size of the purified DNA was

selected before PCR amplification for library construction. Initial DNA concentration was

quantified using Qubit 3.0, and fragment size was determined using Agilent 2100. The final

library samples were additionally examined by Agilent 2100. Afterward, amplified libraries

were sequenced on an Illumina HiSeq 2000 instrument with a 150 bp pair-end (PE).

The raw reads were generated in FASTQ format. The company, moreover, cleaned up raw

reads by removing the adapter sequences. The reads with low-quality scores and those

containing ambiguous nucleotides (N) greater than 10 percent were removed. Clean reads were

obtained for further bioinformatics analysis. The clean raw reads were *de novo* assembled into

contigs using MEGAHIT (Version 1.2.9) by Li et al. (2015) under the Galaxy platform (The

Galaxy Community 2022). Subsequently, the assembled contigs were used as query

sequences to search BLASTN against the dengue virus serotype 2 (DENV-2) complete genome

database (accession no. M29095.1; 10723 bp ss-RNA). BLASTN (NCBI BLAST+ blastn). The

program was run under the Galaxy platform (The Galaxy Community 2022) with a cutoff E-

value of 1x10^{-30}.

The hits between contigs and DENV-2 (M29095.1) from BLASTN output were retrieved for

sequences that were further analyzed for the presence of cvcDNA. The retrieved sequences

were performed using the program Filter FASTA on the sequence headers with a list of IDs.

Afterward, the sequences that matched with DENV-2 (M29095.1) were further analyzed for

Aedes albopictus isolate C6/36 fragments. To detect *Aedes albopictus* isolate C6/36 fragments

in the retrieved contig sequences, the BLASTN program was applied using *Aedes albopictus*

isolate C6/36 whole genome (GCF_001876365.2) as the database (subject sequences).

Multiple sequence alignments were performed using CLUSTALW (https://

www.genome.jp/tools-bin/clustalw). The circular DNA diagrams were drawn by GenomeVx

(http://wolfe.ucd.ie/GenomeVx/). Moreover, the clean raw reads by GENEWIZ, Inc. were

analyzed for similarity to DENV-2 sequences using map reads and the reference program

QIAGEN CLC Genomics Workbench version 12 (QIAGEN, Denmark) to reconfirm the

similarity of raw reads and contig assembly compared to DENV-2 genome.

2.3 Validation for the presence of DENV-2 contigs in the circular DNA preparation

Multiple specific single primers were designed from selected contig sequences to confirm

DENV-2 cvcDNA in the persistently infected DENV-2 cultures (or circular DNA extract). PCR

amplification with each primer using a portion of the original, pooled circular DNA sample as

a template (30 ng) was carried out using KOD FX Neo polymerase (Toyobo, Japan), following

the manufacturer's instructions for PCR conditions. The thermal cycling profile consisted of

an initial denaturation at 98º C for 2 minutes, followed by 35 cycles of denaturation at 98º C

for 10 seconds, appropriate annealing temperature at 60 ℃ for 5 seconds, and extension at 68℃

for 1 minute. The PCR products were visualized by 1.5% agarose gel electrophoresis. There

was no confirmation by submitting PCR products for sequencing again.

2.4 Anti-DENV-2 activity tests using circular vcDNA (cvcDNA) produced by RCA

For the anti-DENV-2 activity test (**Figure 1**), naive C6/36 cells at 10^5 cells/well were seeded

into 6-well plates with 2 ml fresh Leibovitz's (L15) medium with supplements containing 10%

FBS, 0.3% TPB, and 1% antibiotics and incubated for 24 hours at 28°C before transfection

using liposome Escort IV transfection reagent (Sigma) according to the manufacturer's

instructions. After 24 hours of incubation, the medium was removed and the cells were rinsed

with 1 ml fresh L15 medium without supplements. Then, 800 µl fresh L15 cell culture medium

without supplements was added. At the same time, the transfection reagent was separately

prepared by (A) 4 µl Escort IV transfection reagent mixed in 100 µl L15 medium without

supplements and (B) 100 or 1000 ng of cvcDNA mixed in 100 μl L15 medium without supplements and then mixing (A) and (B) to form liposome-forming complexes for 45 minutes at room temperature. Then, the ~200 μl mixtures (A and B) were added dropwise to the cell monolayers in 800 μl fresh L15 medium without supplements with gentle swirling for 5 hours at room temperature. After 5 hours of incubation, they were challenged with DENV-2 stock at 16.85×10^6 copies/well in 1 ml of L15 medium without supplements followed by gentle swirling for 2 hours at room temperature. After incubation for 2 hours, the old medium was removed, and 2 ml fresh L15 medium with supplements (10% FBS, 0.3% TPB, and 1% antibiotics) was added for further incubation at 28°C until day 5.

Figure 1. Diagram of the protocol for anti-DENV-2 activity tests using circular vcDNA.

An experiment to test for anti-DENV-2 activity was carried out with 4 groups, including 1) negative control (naive C6/36 cells), 2) positive control (C6/36 cells+DENV-2), 3) treatment of 100 ng cvcDNA (C6/36 cells+100 ng cvcDNA+DENV-2), and 4) treatment of 1,000 ng cvcDNA (C6/36 cells+1,000 ng cvcDNA+DENV-2) with six replicates in each group. The presence of any cytopathic effect (CPE) was assessed on days 3 to 5 to examine the severity of viral pathology. Pictures of CPE were captured using an inverted microscope (Leica DMi1)

with 20X magnification. Three pictures of each replicate were captured. The seriousness of
CPE was assessed using a grid of 100 squares in one picture. The infected area was counted as
1 for squares containing syncytial cells and as 0 for squares without CPE. The total infected
areas were then summed. Statistical analysis was performed using the R program, with a one-
way ANOVA followed by the Tukey HSD post hoc test (***p-value ≤ 0.001, ns = non-
significant).

3. RESULTS

3.1 Confirmation of circular DNA isolation from DENV-2 infected C6/36 cell cultures

To verify the extraction of circular DNA from DENV-2 infected C6/36 cell cultures by our
isolation process, DNA extracts before and after exonuclease V digestion were examined for
the presence of linear DNA and circular DNA using PCR amplification with specific primers
followed by visualization with gel electrophoresis. Linear DNA was detected using the primers
EF165_Aedes_F/R (Table 1) that amplified host elongation factor gene. Circular DNAs were
detected using two primers of ATPSynthase_150F/R and CYCS_Aedes_125F/R (Table 1) that
amplified mtDNA genes of ATP synthase and cytochrome c oxidase, respectively. The result
in **Fig. 2.** showed the presence of linear DNA amplicons in lane 1 (positive amplification
confirming presence) derived from elongation factor (EF) gene primers before DNA extracts
were treated with exonuclease V. These amplicons were absent after DNA extracts digested
with exonuclease V (lane 4 negative amplification confirming absence). In contrast, there were
amplicons present for the circular DNAs from mtDNA genes (ATP synthase and cytochrome
c oxidase) both before (lanes 2 and 3) and after (lanes 5 and 6) exonuclease V digestion (Figure
2). These results confirmed that only host linear DNA was eliminated while circular DNA
forms remained. The presence of circular DNA initially supported the proposal for presence of
a cvcDNA construct corresponding to the dengue virus single-stranded positive-sense RNA
genome. Therefore, metagenomic sequencing was subsequently conducted to reveal the
sequence in the circular DNA in the cvcDNA

preparation.

Figure 2. Analysis of PCR assays by agarose
gel electrophoresis to confirm host circular
DNA presence but linear DNA absence
following exonuclease V (EXO V) digestion.

Before EXO V digestion, Lane 1 shows amplicons for the linear DNA of host elongation factor
gene (EF, Lane 1) and for two mitochondrial circular DNA genes, ATP synthase (ATP Lane
2) and cytochrome c oxidase (CYCS Lane 3). After EXO V digestion, the EF amplicon is
absent (Lane 4) while the two mitochondrial gene amplicons (ATP Lane 5 & CYCS Lane 6)
still remain. M is a GeneRuler DNA ladder mix.

3.2 Identification of DENV-2 cvcDNA from circular DNA metagenomic sequencing
3.2.1 Raw reads matching DENV-2 mapped irregularly across its whole genome
Based on the viral accommodation concept, we hypothesized that the RNAi and host reverse
transcriptase activities would generate vcDNA with high variation in size and sequence and
that a portion of this would be in circular form (i.e., cvcDNA), even though DENV-2 is an
RNA virus. Metagenomic sequencing was utilized to test this hypothesis. To ensure the next-
generation sequencing procedure, we submitted two separate circular DNA extracts to two
companies (see section Materials and Methods 2.2). The raw reads of two metagenomic
sequencings were deposited as NCBI short read archive (SRA) under BioProject of
PRJNA1003297 and PRJNA1160223 that are raw reads from GENEWIZ, Inc. with the ID of
D2C6P1-7, and raw reads from NovogeneAIT Genomics with the ID of CCDENV21. The
sample of D2C6P1-7 generated 29,182,977 pair-end (150 bp) raw reads, while the sample of
CCDENV21 generated 35,174,410 pair-end (150 bp) raw reads.

From each company, raw reads that matched the DEN-2 genome sequence were mapped to the
reference genome M29095 of DENV-2 (**Fig. 3**). Distribution of the DENV-2-matching raw
reads was irregular and spanned across most of the genome sequence. This is consistent with
the proposal that EVE acquisition is an individual cell process that leads to high variation in
the types and numbers obtained from a tissue sample or cell culture. In contrast, raw data from
circular DNA preparation (D2C6P1-7) revealed a high-frequency distribution of genes only
within the ENV-NS2A region of the DENV-2 genome. This roughly matched a similar map
region for the CCDENV21 data set.

Figure 3. Diagram showing the number, length and distribution of 122 raw reads obtained from the metagenomics of the D2C6P1-7 (122 raw reads) and CCDENV21 (191 Raw reads) cvcDNA samples when mapped against the reference DENV-2 genome (GenBank M29095).

Both raw read samples were separately assembled using MEGAHIT, yielding 110,350 and 450,310 contigs for D2C6P1-7 and CCDENV21, respectively. Given the emphasis on DENV-2 cvcDNA, we further analyzed the contigs from two samples using BLASTN. Each contig assembly was tested separately using BLASTN against the dengue virus serotype 2 (DENV-2) complete genome database (M29095.1). With a cut-off E-value of 1×10^{-10} when using the BLASTN program, the results showed that the D2C6P1-7 sample with 110,350 contigs yielded 3 contigs that matched the DENV-2 sequence. These were k141_3180 (368 bp), k141_92607 (433 bp), and k141_113249 (327 bp) (supplementary data **Table S1**). BLASTN output of the CCDENV21 sample with 450,310 contigs showed 5 DENV-2 matching contigs that included contigs k141_157363 (417 bp), k141_311629 (281 bp), k141_42362 (298 bp), k141_43783 (2061 bp), and k141_554965 (499 bp) (supplementary data **Table S1**). Sequences of all eight contigs were retrieved for further comprehensive analysis (supplementary data **Table S2**).

The 8 contig sequences were subsequently analyzed for host DNA fragments (*Aedes albopictus* isolate C6/36) using BLASTN. This process was run to confirm the presence of DENV-2 and *A. albopictus* isolate C6/36 sequences together in the same contig. The BLASTN output showed three contigs containing sequences of dengue virus serotype 2, joined with fragments of DNA from the *A. albopictus* isolate C6/36 (supplementary data **Table S3**). All three contig sequences, including contigs k141_157363, k141_42362, and

k141_43783, were derived sequences, including contigs k141_157363, k141_42362, and

k141_43783, were derived from the CCDENV21 sample. The feature summary of raw reads,

contigs from raw reads assembly, the retrieved contig sequences containing the DENV-2

sequences, and the *Aedes albopictus* isolate C6/36 fragments from each company are shown

in **Table 2**.

Table 2. The feature summary of raw reads and the number of contigs from the raw read
Assembly.

Company Sample ID	D2C6P1-7	CCDENV21
NCBI SRA under BioProject	PRJNA1003297	PRJNA1160223
Raw reads (PE150)	29,182,977	35,174,410
Number of contigs from MEGAHIT assembler	110,350	450,310
Number of contigs possessing DENV-2 sequences	3	5
Number of contigs possessing the sequences of DENV-2 and the *Aedes albopictus* isolate C6/36	0	3

3.2.2 Characterization of eight contigs of DENV-2 cvcDNA

As mentioned above, cvcDNA constructs would vary in size and sequence. BLASTN results

of **Table S3** demonstrated that the sequences of *A. albopictus* isolate C6/36 in the cvcDNA

contigs may have originated from several locations in the host genome. For example, the

sequences at 278-417 (140 bp) of contig k141_157363 and the sequence at positions 1-140

(140 bp) of contig k141_42362 gave hits at 49 and 47 different locations in the *A. albopictus*

isolate C6/36 genome (supplementary data **Table S3**). Contig k141_43783 showed positions

at 387-1026 (640 bp), 387-1031 (645 bp), 388-603 (216 bp), 1028-1122 (95 bp), 1029-1117

(89 bp), and 1110-1194 (85 bp) that mapped to 19 locations in host genome (supplementary

data **Table S3**). These results indicated that DENV-2 cvcDNA contained a wide variety of viral

and host gene fragments in variable numbers.

Subsequently, all eight contig sequences were examined for circular forms by searching for

overlaps within each sequence. The results showed that contigs k141_43783 (2061 bp) and

k141_554965 (499 bp) had complementary overlap sequences of 141 bp at the ends of their

contigs (supplementary data **Document S1, Figure S1**). In **Table 3**, in the far-right column,

the reading directions within the contigs are indicated by arrows with reference to the DENV-

2 5'v3' genome sequence, and the vertical bars indicate lack of continuity in the sequences

and reading directions. These features, especially when accompanied by linked host DNA

sequences, support the contention that the contigs arose from cvcDNA constructs derived

from the host genome and not directly from the DENV-2 genome. The circular diagram of

two contigs of DENV-2 cvcDNA is illustrated in **Fig. 5**. The 141 bp overlap regions were

merged. Thus, the contigs k141_43783 and k141_554965 represent cvcDNA constructs of

1921 and 358 bp, respectively.

Table 3. Summary features for eight contigs that hit the DENV-2 and/or *Aedes albopictus*
346 C6/36 cell genome sequences. The contigs derived from raw read set D2C6P1-7 are
highlighted 347 in a gray background, while those derived from set CCDENV21 are not
highlighted. In the 348 column for "Position and Reading Direction", arrows indicate reading
direction and vertical 349 bars indicate disjunction between the DENV-2 sequence
connections. Yellow highlights in the column entitled "Fragment Positions" indicate overlaps
(i.e., common bases at the junction of two fragments).

#	Contig Name	Length (bp)	C6/36 content	Circular (yes/no)	# Fragments	Fragment Positions	DENV-2 Genome Position and Reading Direction (← →)
1	k141_3180	368	No	No	3	26-253 (228) 248-323 (76) 321-368 (48)	1458-1685→ |1246-1321→ | 1082-1129→
2	k141_92607	433	No	No	2	13-128 (116) 116-413 (298)	←7302-7188 | 6374-6671→
3	K141_113249	327	No	No	1	21-327 (307)	1287-1567→
4	K141_157363	417	Yes	No	3	1-147 (147) 147-281 (135) 278-417(140)	←1422-1276 | ←1619-1485 C6/36 seq
5	K141_311629	281	No	No	1	1-281 (281)	1287-1567→
6	K141_42362	298	Yes	No	2	1-140 (140) 141-158 (18) 159-298 (140)	C6/36 seq No identity 1485-1346→
7	K141_43783	2061	Yes	Yes	6	1-141 (141) 141-387 (247) 387-1031 (645) 1028-1122 (95) 1110-1194 (85) 1179-2061 (883)	←1422-1282 | ←2758-2512 | C6/36 seq. C6/36 seq. C6/36 seq. ←2165-1282
8	K141_554965	499	No	Yes	5	4-96 (93) 95-272 (178) 277-361 (85) 362-454 (93) 453-499 (47)	←2757-2665 | ←3319-3142 | |3316-3400→ ←2757-2665 | ←3319-3273

Figure 4. Diagram of the reference DENV-2 polyprotein genome together with five

representative contigs from **Table 3**. The scales are relative, with the contigs magnified about

5 times compared to the whole genome to make the fragment features more visible. These

contigs illustrate the features of chimeric gene fragment compositions (including viral

genome fragments or viral plus host genome fragments), the linkage of disjunctive viral and

host genome fragments, and variable reading directions (arrows below the contig bars), which

can be compared to the reference genome diagram at the head of the illustration.

The feature summary of eight contigs in Table 3 and the BLASTN results in the

supplementary data (Table S1) revealed that the eight contig sequences originated from

positions scattered across the DENV-2 genome. Some contigs from two different company

pools of raw reads also shared common areas. The eight contigs generated 18 fragments that,

in a scrambled and scattered manner, matched their positions and reading directions in the

DENV-2 genome (Table 3 and Fig. 4).

Fig. 5. Diagram of 2 circular DNA contigs, k141_43783 and k141_554965. The grey fill
377 indicates C6/36 host DNA sequences, while the DENV-2 sequences are filled with colors
to match the DENV-2 genes illustrated in **Fig. 4** (envelope gene in yellow and NS1 gene in
green). **(A)** Contig k141_43783 (2,061 bp) contains three disrupted fragments of the DENV-2
envelope protein gene plus three disruptive fragments of C6/36 DNA. **(B)** Contig k141_554965
(499 bp) contains five DNA fragments derived from the DENV-2 genome (M29095), positions
varying from 2,665 to 3,400. The inner blue circle indicates the position of the nucleotides in
the contigs. Note the opposite reading directions in the two single contigs, each containing
disruptive sequences. The inner double-circle part represents overlapping sequences, each 141
bp, resulting in contig 7 (k141_43783) with 1,921 bp and contig 8 (k141_554965) with 358 bp
in circular form. The figure does not show the actual contig size scales in the comparisons, but
the figure still shows that contig 8 (k141_554965) is smaller than contig 7 (k141_43783),
although it is actually 5 times smaller.

3.3 Confirmation of the presence of circular DENV-2 DNA related to contig sequences

According to **Table 3** and **Figs. 4** and **5**, the data showed that the DENV-2 genome content in
the contigs originated from the region of the genome where most raw-read matches occurred
at positions 1000-2000. Multiple single primers were designed to detect circular DENV-2 DNA
using DENV-2 contig sequences at positions 1000-2000. The single primer of
cvcDENV2_1422 5'-GATTTCCTTGCCATGTTTTCCTGTGTCA-3' was directly designed
from the sequence of contig k141_43783, and another single primer of cvcDENV2_1422x 5'-
TGACACAGGAAAACATGGCAAGGAAATC-3' was designed from the antiparallel
complementary sequence of contig k141_43783. A portion of the final circular DNA
preparation obtained from using RCA (the same sample submitted for Novogene sequencing).

was used as a template to test these single primers. When each of these primers were used alone

for PCR amplification, they successfully produced PCR amplicons (**Fig. 6.**). For example, the

single primer for cvcDENV2_1422x gave several amplicon bands of different sizes, indicating

the presence of a variety of circular DENV-2 DNA targets, with the most robust band found at

an amplicon size of ~500 bp in both primers and the product size of ~2000 bp from a primer

of cvcDENV2_1422 (**Fig. 6**). These results also supported the idea that cvcDNA produced

from the same DENV2 genome regions were produced in variable sizes.

Fig. 6. Photograph of an agarose gel showing that PCR amplicons were obtained using 2
different single primers (cvcDENV2_1422 and cvcDENV2_1422X), indicating the presence
of palindromic repeats. Lane M is a GeneRuler DNA ladder mix.

3.4 DENV-2 tolerance in C6/36 cells accelerated by cvcDNA of DENV-2 treatment

In viral accommodation, vcDNA arises as linear (lvcDNA) and circular (cvcDNA) forms via

endogenous host RT activity, leading to a viral-tolerant, persistent infection state. Here, a

mixture of cvcDNA of DENV-2 obtained using the cvcDNA preparation amplified by RCA

(see section 2.1 of the Materials and Methods) was used in pre-exposure tests with naive C6/36

cells prior to the DENV-2 challenge (**Fig. 1)**. The consistent outcome for naive cells challenged

with DENV-2 is always a significant occurrence of CPE exemplified by syncytial cells on day

2, followed by a progressive decrease in CPE until day 5. This is shown in **Fig. 7**, where CPE

can be readily observed on day 2 in untreated cells. This contrasts with the cells treated with

DENV-2-cvcDNA (at 100 and 1,000 ng) on day 2, in which only a minor CPE was observed. On day 5, the untreated groups showed distinguishable morphological differences, with clearly visible, more-severe CPE in the untreated cells (**Fig. 7**) that showed. a significant reduction in CPE. This result was confirmed statistically by quantifying the infected areas, as shown in **Fig. 8**. It revealed that the presence of DENV-2-cvcDNA prior DENV-2 challenge resulted in the cells accommodating DENV-2 faster than untreated cells and confirmed it association with accommodation.

Fig. 7. The cell morphology of DENV-2-infected cells, 2,3,4, and 5 days after infection, treated or not treated with DENV-2 cvcDNA (100 and 1,000 ng) prior to the DENV-2 challenge. CPE was observed by syncytial cells (black arrows). The untreated cells exhibit significant CPE, as evidenced by the formation of syncytia (black arrows).

Fig. 8. The percentage of the infected areas indicated by CPE on days 3, 4, and 5, in cells

treated or not treated with DENV-2 cvcDNA (100 and 1,000 ng) prior to DENV-2 challenge.

CPE was assessed by the presence of syncytial cells. Statistical analysis was performed using

447 the R program and analyzed with one-way ANOVA, followed by the Tukey HSD post hoc

test (***p-value $\leq$ 0.001; ns = non-significant).

4. DISCUSSION

The tiny proportion of DNA raw reads homologous to DENV-2 (122 raw reads) in the total

of over 58 million raw reads from GENEWIZ, Inc., and the 284 raw reads in 70 million raw

reads from Novogene Co., Ltd, was not unexpected. However, these results were consistent

with the number of contigs from the MEGAHIT assembler that showed three contigs

homologous to DENV-2 obtained from a total of 110,350 contigs from GENEWIZ, Inc., and

the five contigs homologous to DENV-2 derived from a total of 450,310 contigs obtained by

Novogene Co., Ltd (Table 2). From **Figs. 5** and **6**, with the naked eye, there are clusters of

slightly higher frequency raw reads around the region of ENV (DENV-2 positions of 937-

2421), but no statistical analysis was carried out to determine whether these differences were

significant. However, in this brief communication, the results supported our hypothesis that

C6/36 cells infected with DENV-2 would produce cvcDNA constructs containing DENV-2

homologous sequence fragments.

As seen in **Figs. 3** and **4**, **Table 3**, and supplementary data **Table S2**, the sequences of these

eight contigs could not have arisen directly from a DENV-2 viral genome because of the

disjunctive and opposing reading directions of different portions of the viral genome that were
connected in the contigs. In addition, three contigs contained sequences matching disjunctive
fragments of the host genome sequence linked to the DENV-2 sequences. These features
support the contention that circular DNA entities may have arisen from the EVE of DENV-
2and been acquired after DENV-2 infection.

We have confirmed that DENV-2 infections in C6/36 cell cultures, as with DENconV-1
473 infections described earlier (Goic et al. 2016), result in a specific, adaptive immune
response that involves the production of vcDNA by the endogenous host reverse transcriptase.
We have confirmed that this vcDNA includes circular forms that can be differentially isolated
from whole DNA extracts. However, for contigs that contained only unidirectional DENV-2
sequences, it was impossible to determine whether they originated from the infecting virus or
from EVE. To do so, a procedure for selecting single-cell clones free of DENV-2 infection
would be needed, as previously recommended (Flegel, 2020, 2022). This would allow the
individual DEN2-free single-cell clones to be separately analyzed for comparison of their
cvcDNA constructs. The results from this study reveal that such a process would be successful.

Our previous work using C6/36 cells was carried out to test hypotheses regarding the shrimp
response to viral pathogens because no immortal crustacean cell lines existed then or even now
to do such work (Kanthong et al. 2008, 2010, Sriton et al. 2009, Gangnonngiw et al. 2010,
Arunrut et al. 2011). Those earlier studies revealed the ability of C6/36 cells to individually
support three different shrimp viruses (WSSV, TSV, and YHV) that remained infectious for
shrimp up to 5 passages but not thereafter. On the other hand, the insect cells remained
immunopositive for those 3 viruses for up to 30 split passages, and they could transfer
immunopositive status to the hemocytes of challenged shrimp. This suggested the possibility
of using insect cells as a source of attenuated viurses for injection into shrimp to protect (i.e.,
vaccinate) against disease caused by cognate viruses. A first attempt to do this with a yellow
head virus (YHV) failed (Gangnonngiw & Kanthong 2023). However, the hemocytes of the
injected shrimp in the failed vaccination attempt were converted to immunopositive status for
YHV, and there was a significant delay in their mortality. Since the "attenuated vaccine"
comprised a crude extract from whole insect cells, it is possible, based on the results described
herein, that vcDNA from YHV in those extracts contributed to this delay in mortality.

In addition, we have revealed here that cvcDNA prepared *in vitro* after RCA targeting an EVE of DENV-2 can reduce CPE in naive cells challenged with DENV-2. This experiment still exhibited the same level of DENV-2 viral replication as the untreated cells, with a previous report in shrimp showing that a crude cvcDNA extracted from shrimp infected with the parvovirus IHHNV could result in a significant decrease in IHHNV replication (5). Further investigation is warranted into the specific prevention of CPE by cvcDNA treatment, with the addition of quantifying DENV-2 replication. To us, the fact that pre-exposure of the naïve C6-36 cells to the crude DENV-2 cvcDNA preparation prior to DENV-2 challenge allowed them to reach accommodation quicker than the untreated cells is worthy of further investigation. It as if its may indicate the presence of some preliminary factor(s) necessary for accommodation to occur were already present. It is somehow similar to the earlier discovery that C6-36 cells already accommodating DENV-2 could more rapidly accommodate the densovirus *Aedes albopictus* densovirus (*Aal*DNV) and vice-versa (Burivong et al. 2004). It suggests the presence of a critical factor necessarily for accommodation to occur. One possibility may be host-derived reverse transcriptase (RT) or some mechanism related to it. It is a key enzyme in the viral accommodation process, and our recent work described at BioRxiv has revealed that RT inhibition with cells that have accommodated DENV-2 revert to the CPE state (Gangnonngiw et al. 2025).

These results have shown that C6/36 cells may be used as a convenient model to study the detailed mechanisms by which cvcDNA is produced directly from viral RNA and whether these mechanisms differ from those that give rise to cvcDNA from EVE. We hypothesize that C6/36 cells or other immortal insect cell lines may also produce cognate cvcDNA and EVE when challenged with shrimp viruses. If so, they might provide a convenient tool to develop EVE-clone libraries to facilitate screening for the most protective EVE or combinations thereof for use with shrimp. This is already possible, of course, for insect viruses, in general, using their matching immortal cell lines.

ACKNOWLEDGEMENTS

This research project was supported by Rajamangala University of Technology Tawan-ok (Basic Research Fund: fiscal year 2021, **Project no. 50063**) and Mahidol University (Fundamental Fund: fiscal year 2023 by National Science Research and Innovation Fund (NSRF) **Grant no. FF-056/2566**).

REFERENCES

Arunrut N, Phromjai J, Gangnonngiw W, Kanthong N, Sriurairatana S, Kiatpathomchai W. 2011. In vitro cultivation of shrimp Taura syndrome virus (TSV) in a C6/36 mosquito cell line. Journal of Fish Diseases 34:805-810.

Bonning BC, Saleh M-C. 2021. The interplay between viruses and RNAi pathways in insects. Annual Review of Entomology 66.

Burivong, P., Pattanakitsakul, S.-N., Thongrungkiat, S., Malasit, P., Flegel, T.W., 2004. Markedly reduced severity of Dengue virus infection in mosquito cell cultures persistently infected with Aedes albopictus densovirus (AalDNV). Virology. 329, 261-269.

Cerenius L, Jiravanichpaisal P, Liu H-p, Soderhall I. 2010. Crustacean immunity. Invertebrate immunity:239-259.

Flegel TW. 2020. Research progress on viral accommodation 2009 to 2019. Developmental & Comparative Immunology 112:103771.

Flegel TW. 2022. Viral accommodation: A New mechanism for disease control in shrimp aquaculture, p 427-445. *In* Alday-Sanz V (ed), The Shrimp Book II. 5m Books Ltd, United Kingdom.

Gangnonngiw W, Kanthong N. 2023. Failed shrimp vaccination attempt with yellow head virus (YHV) attenuated in an immortal insect cell line. Fish and Shellfish Immunology Reports:100084.

Gangnonngiw W, Kanthong N, Flegel TW. 2010. Successful propagation of shrimp yellow head virus in immortal mosquito cells. Diseases of aquatic orgalemnisms 90:77-83.

Gangnonngiw W, Flegel TW, Kanthong N. Host reverse transcriptase is required to maintain persistent dengue virus infections in C6/36 insect cells. bioRxiv 2022.12.20.521226; doi: https://doi.org/10.1101/2022.12.20.521226

Goic B, Stapleford KA, Frangeul L, Doucet AJ, Gausson V, Blanc H, Schemmel-Jofre N, Cristofari G, Lambrechts L, Vignuzzi M. 2016. Virus-derived DNA drives mosquito vector tolerance to arboviral infection. Nature communications 7:12410.

Kanthong N, Khemnu N, Pattanakitsakul S-N, Malasit P, Flegel TW. 2010. Persistent, triple-virus co-infections in mosquito cells. BMC microbiology 10:1-8.

Kanthong N, Khemnu N, Sriurairatana S, Pattanakitsakul S-N, Malasit P, Flegel TW. 2008. Mosquito cells accommodate balanced, persistent co-infections with a densovirus and Dengue virus. Developmental & Comparative Immunology 32:1063-1075.

Li, D., Liu, C.-M., Luo, R., Sadakane, K., Lam, T.-W. (2015). MEGAHIT: an ultra-fast single-
node solution for large and complex metagenomics assembly via succinct de Bruijn
graph. Bioinformatics, 31(10), 1674–1676.
R Core Team (2024). _R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
Sriton A, Kanthong N, Gangnonngiw W, Sriurairatana S, Ubol S, Flegel TW. 2009. Persistent
expression of shrimp-virus antigens in two insect cell lines challenged with two shrimp
viruses. Fish Pathology 44:86-93.
Taengchaiyaphum S, Buathongkam P, Sukthaworn S, Wongkhaluang P, Sritunyalucksana K,
Flegel TW. 2021. Shrimp parvovirus circular DNA fragments Arise from both
endogenous viral elements and the infecting virus. Frontiers in Immunology 12:729528.
The Galaxy Community. The Galaxy platform for accessible, reproducible and collaborative
biomedical analyses: 2022 update, Nucleic Acids Research, Volume 50, Issue W1, 5
July 2022, Pages W345–W351, doi:10.1093/nar/gkac247

Supplementary information

Yellow highlights indicate Dengue virus type 2 complete genome sequences (M29095.1).

Turquoise highlights indicate Aedes albopictus isolate C6/36 genome fragments.

Green highlights indicate repeat regions representing circular DNA.

>k141_157363 flag=0 multi=3.5761 len=417

BLASTN output showed that the positions of k141_157363, which matched the Dengue virus type 2
complete genome (M29095.1), were 1-147 (147 bp) and 147-281 (135 bp).

BLASTN output showed that the position of k141_157363, which matched the Aedes albopictus
isolate C6/36 (NW_017856292.1), was 278-417 (140 bp).

GATTTCCTTGCCATGTTTTCCTGTGTCATTTCCGACTGCATGCTCTTCCCCTGAGTGAGGT
GTTATCACAATGGTGTATTCCAAGTTTTCTGGTTGCACGACTTTTCCTTTCATGTTCTTTTT
GCATGTGAACATAGCACAGGTCACCTTGTGTGTCCGCTCCGGGCAGCCATGGCAACGGC
AGGTCTAGGAACCATTGCCTGTGCACCAGCCAAGCTTTATTTTCCATTTGCAGCAACACC
ATCTCATTGAAGTCGAGGCCCGTTCTCGGAGAGCACTCCCTTAAAGAAAATTCTGTTGCT
TAACTGTCTACACATATTTGGTTAGTGGATTAGTTCACAACACATTACAGGTCCAATCTA
TGCGGAGCCTTTCTTTCTCGAGTTGATCTTCTAACTGTCTCACCAGCATTATCTA

>k141_42362 flag=1 multi=1.0000 len=298

BLASTN output showed that the position of k141_42362, which matched the Dengue virus type 2
complete genome (M29095.1), was 159-298 (140 bp).

BLASTN output showed that the position of k141_42362, which matched Aedes albopictus isolate
C6/36 (NW_017858069.1), was 1-140 (140 bp).

GGATCCTGGCATCAACATTTCACGACGTCGCTGGATATCATCTGGAACCTGTTCACGCAG
TTCAGCACGCTGAGTAACGGGTCCTGGCACTGGTCCCGGCTTCAATTGGTCGACATGCCG
CTTCCACATTCTACCGTCATAGAGTTCCATCACAGAAGCATCGTGACAGTGCCATAGCCT
GTCAACTCTGCTTCTGTGATGGAACTCTGTGGTGTTATTTTGATTTCCTTGCCATGTTTTCC
TGTGTCATTTCCGACTGCATGCTCTTCCCCTGAGTGAGGTGTTATCACAATGGTGT

>k141_43783 flag=0 multi=7.4781 len=2061

BLASTN output showed that the positions of k141_43783, which matched the Dengue virus type 2
complete genome (M29095.1), were 1-141 (141 bp), 141-387 (247 bp), and 1179-2061 (884 bp).

BLASTN output showed that the positions of k141_43783, which matched the Aedes albopictus
isolate C6/36 (NW_017856292.1), were 387-1031 (645 bp), 1028-1122 (95 bp), and 1110-1194 (85
bp).

GATTTCCTTGCCATGTTTTCCTGTGTCATTTCCGACTGCATGCTCTTCCCCTGAGTGAGGT
GTTATCACAATGGTGTATTCCAAGTTTTCTGGTTGCACGACTTTTCCTTTCATGTTCTTTTT
GCATGTGAACATAGCACACTTCAGCTCAGTGGGCTGGGGCTGCAGAGATCGTTTTCCTGC
CTGCATGATTCCTTTGATGTCTCCTGTCATAATAGTCAACTTCACCTCATTTTCTGATAGA
ATGTGATTCAATTCTGGTGTTATTTGTTTCCACATCAGATTTTCCAGTCTTGTTACTGAGC
GGATTCCACAAATGCCCTCTTCATGAGCTTTCTGGATAGCTGAAGCTAGTTTTGAAGGGG
ATTCTGGTTGGAACTTGTATTGAGTTCGAAATGACGGATGAGGGCGAGATCAAACAGTTC
CTAGGTATGCGGATTGATCGTGACAGGAACGCAGGTGTTTTGAAAATCAGTCAACGTGA
CTACTTTGATGGGCTACTGCGGCGTTTCGGAATGATGGACTGCAAACCAATTTCGACGCC
GATGGAGAACCGGCTGAAGCTGCAGAAGGGTGACGAGAAGCAAAGGACATCGAAACCT
TACCGTGAGCTGGTCGGGTGTTTAACATACGCGTCTATCACGACGAGGCCGGATCTGGCT
GCGGCTGTAAACTTTTTCAGTCAGTTTCAGTCGTGTCCGAATGAAGAGCACTGGAATCAT
CTGAAGCGGATTCTCCGTTATATCAAGGGGACATTGGACGTCGGTCTGGTGTATAAGGTG
GAGGAGAATGAACCGTTGCTGCAGGTATTCTGTGACGCGGACTGGGCGAACGATCTTAC
CGATCGTCGTTCGGTTTCTGGCTACGTGGTGAAGCTGTATGGAAGTACTGTTGGTTGGAT
TACGAGAAAGCAACAGACCGTATCATTGTCGTCTACGGAAGCGGAACTAGCGGCATTGT
GCACAGCTGCATGCCACACGTTGTGGATGGTGCGCATACTGGAAGATCTTGGAATGCGG
TTGGACGCTTGTTAAGATGGCAACCACTGCACTCTAATTGCCCCCACTTAGTTCGACCCC
ACTGCGCAAGGTGCGCCTATGAACATTTTTTCACTTGGTAGTTAAACATCGACTGATTTA
TTCAACTGAGTATGGTATTACATTCGGTTCACACTCGGGTTCGCTCACTGATTGTTGTCTC
AATCATTTGGCCGATAGAACTTCCTTTCTTAAACCAGTTGAGCTTCAATTGTCCCGGCTCT
ACTCCTATGATGATGTAGCTGTCTCCGAATGGAGGTTCTGCTTCTATGTTGACTGGGCTAT
CTTTTTCTGTTACGATTGGGTTGACTGTAATCAGGCGACCTAAAACATGTCTTTTTTCCAA
ATCCATTATCTCAAAAGGGATCTTACATGGAGAACCGTCCCCTTCATATTGTACTCTGAT
AACTATTGTTCCATGTTGTGTTTCTGCTATTTCCTTCACAACTTTAAACTTTCCTGTGCACA
TAGAGTATGACATTCCTTTGAGCTGTAGTTTGTCCATCCTCAGCCTGCACTTGAGATGTCC
TGTGAACAGTAAGTTTCCTGATGACATCTGGATTTCTGTGGCCCCTGTGAGTGCTGTGTG
CATGGCCCCTTCTTGGGATCCCAAAACAACAACATCCTGTTTCTTCGCATGGGGATTTTTG
AAAGTGACCAATGTCTCTTTCTGTATCCAATTTGATCCTTGTGTGTCCGCTCCGGGCAGCC
ATGGCAACGGCAGGTCTAGGAACCATTGCCTGTGCACCAGCCAAGCTTTATTTTCCATTT
GCAGCAACACCATCTCATTGAAGTCGAGGCCCGTTCTCGGAGAGCACTCCATCGTGACA
GTGCCATAGCCTGTCAACTCTGCTTCTGTGATGGAACTCTGTGGTGTTATTTTGATTTCCT
TGCCATGTTTTCCTGTGTCATTTCCGACTGCATGCTCTTCCCCTGAGTGAGGTGTTATCAC
AATGGTGTATTCCAAGTTTTCTGGTTGCACGACTTTTCCTTTCATGTTCTTTTTGCATGTGA
ACATAGCACA

>k141_554965 flag=3 multi=22.0615 len=499

BLASTN output showed that the positions of k141_554965, which matched the Dengue virus type 2
complete genome (M29095.1), were 4-96 (93 bp), 95-272 (178 bp), 277-361 (85 bp), and 362-454 (93
bp).

AGGCTTCAGCTCAGTGGGCTGGGGCTGCAGAGATCGTTTTCCTGCCTGCATGATTCCTTT
GATGTCTCCTGTCATAATAGTCAACTTCACCTCATTCTTAAAGAGGGTCCTCTATTTCCAC
AGTCCTCAGTCACCACCACTGTGGTTCCTTCGCAGAAATCAAAGTCCATCTCAAGCTTAC
CTAGATGCCATGGTCCTGCTGTTTGTGTATGGTAGCCTGGTCTGTAGTTGTGTTGTGACAC
TGGTCCAGCGAAATTCTTTGGAATTATCATAGTCAGACCAACTACTGCCTCTGGAAAACT
CATAACAGAATGGTGCTGCCGATCTTGCACATTACCACCGCTAAGATACAGAGGTGAGG
CTTCAGCTCAGTGGGCTGGGGCTGCAGAGATCGTTTTCCTGCCTGCATGATTCCTTTGATG
TCTCCTGTCATAATAGTCAACTTCACCTCATTCTTAAAGAGGGTCCTCTATTTCCACAGTC
CTCAGTCACCACCACT

Supplementary Figure S1

141 bp overlap regions using sequence ailgnment

```
k141_43783_1-141          GATTTCCTTGCCATGTTTTCCTGTGTCATTTCCGACTGCATGCTCTTCCCCTGAGTGAGG
k141_43783_1921-2061      GATTTCCTTGCCATGTTTTCCTGTGTCATTTCCGACTGCATGCTCTTCCCCTGAGTGAGG
                          ************************************************************
k141_43783_1-141          TGTTATCACAATGGTGTATTCCAAGTTTTCTGGTTGCACGACTTTTCCTTTCATGTTCTT
k141_43783_1921-2061      TGTTATCACAATGGTGTATTCCAAGTTTTCTGGTTGCACGACTTTTCCTTTCATGTTCTT
                          ************************************************************
k141_43783_1-141          TTTGCATGTGAACATAGCACA
k141_43783_1921-2061      TTTGCATGTGAACATAGCACA
                          *********************

k141_554965_1-141         AGGCTTCAGCTCAGTGGGCTGGGGCTGCAGAGATCGTTTTCCTGCCTGCATGATTCCTTT
k141_554965_359-499       AGGCTTCAGCTCAGTGGGCTGGGGCTGCAGAGATCGTTTTCCTGCCTGCATGATTCCTTT
                          ************************************************************
k141_554965_1-141         GATGTCTCCTGTCATAATAGTCAACTTCACCTCATTCTTAAAGAGGGTCCTCTATTTCCA
k141_554965_359-499       GATGTCTCCTGTCATAATAGTCAACTTCACCTCATTCTTAAAGAGGGTCCTCTATTTCCA
                          ************************************************************
k141_554965_1-141         CAGTCCTCAGTCACCACCACT
k141_554965_359-499       CAGTCCTCAGTCACCACCACT
                          ********************
```

758   **Other supplementary information available upon request.**