1

Intra-lineage microevolution of *Wolbachia* leads to the emergence of new cytoplasmic incompatibility patterns

Short title: Molecular bases of CI patterns emergence

Alice Namias[1,*], Annais Ngaku[1], Patrick Makoundou[1], Sandra Unal[1], Mathieu Sicard[1], Mylène Weill[1,*]

[1] *ISEM, Université de Montpellier, CNRS, IRD, EPHE, Montpellier, France*

*Corresponding authors

Email: alice.namias@gmail.com (AN), mylene.weill@umontpellier.fr (MW)

Abstract

The mosquitoes of the *Culex pipiens* complex are worldwide vectors of arboviruses, filarial

nematodes and avian malaria agents. In these hosts, the endosymbiotic bacteria *Wolbachia* induce

cytoplasmic incompatibility (CI), a reduced embryo viability in so-called incompatible crosses. The

Wolbachia infecting *Culex* (*w*Pip) cause CI patterns of an unparalleled complexity, associated with the

amplification and diversification of their *cidA* and *cidB* genes, with up to six different gene copies

described in a single *w*Pip genome. By repeating crosses between *Culex* isofemale lines over 17 years,

we documented the emergence of a new compatibility type. Using a new sequencing method

adapted to multi-genes families to acquire *cid* genes, we show that some *w*Pip genomes lost specific

gene copies, giving rise to several sub-lineages segregating within a cage. Linking phenotypic changes

with their underlying genotypic bases, we show that gene copies which are key for CI phenotypes

originated from recombination, rather than point mutations. We show how new CI patterns can

emerge as a two-step process, with (i) local changes in CI genes, while maintaining compatibility with

surrounding lines (all sub-lineages being mutually compatible) and (ii) reveal of those incompatibility

by migration and exposure to hosts infected with incompatible lines.

Keywords

Wolbachia, *Culex pipiens*, cytoplasmic incompatibility, microevolution, recombination

Introduction

Wolbachia are maternally-transmitted endosymbiotic bacteria which infect up to 50% of arthropod species [1–3]. These bacteria are well known for their wide range of reproductive manipulations in arthropods. Their major manipulation is cytoplasmic incompatibility (CI), a reduction in embryos' hatching rates in crosses between infected males and uninfected females (in its simplest form). CI results from a *Wolbachia*-induced *modification*, perturbing embryos' first division, that infected females are able to *rescue* [4,5]. CI is thus formalized in a global *modification-rescue* framework.

CI can also occur between two infected individuals if they are infected with genetically different and incompatible *Wolbachia* strains. In *Culex pipiens* mosquitoes, where *Wolbachia* infection is fixed [6], the *w*Pip strains are responsible for a unique (to date) complexity of CI patterns based on multiple uni- and bidirectional incompatibilities [7–10]. A first analysis of *w*Pip genetic diversity in *Culex* identified 5 phylogenetic groups (named *w*Pip-I to *w*Pip-V, [11]). Confronting *w*Pip phylogeny with crossing results revealed that these groups were good predictors of compatibility outcomes [9]: while both compatible and incompatible crosses were reported in crosses among *w*Pip groups, intra-group crosses were, with very few exceptions, compatible.

In 2013, putative CI genes were identified by a combination of proteomics and genetics approaches [12]. It was four years later, in 2017, that two functional studies confirmed that these genes were key for CI, naming them *cif* for 'CI factors' [13,14]. The *cif* genes were described in *cifA-cifB* tandems, with a central role in CI, although precise molecular mechanisms of CI are still to be determined. Two main alternative models are still debated: a host-modification (HM) model, stating that toxicity occurs through modifications of the host, and rescue through a reversal of these modifications; and a toxin-antidote (TA) model, in which the sperm carries a *Wolbachia* factor which is toxic, unless there is the appropriate *Wolbachia* antidote in the egg that binds and neutralizes the toxin [15–18]. In *Culex,* the complexity of the crossing patterns makes the TA model far more likely: (in)compatibility would simply be based on different toxin-antidote binding abilities [16,19–21].

Recent publications all agree with *cifA* being the rescue (resc, or antidote) gene, and *cifB* a major

modification (mod, or toxin) gene [13,18,22]. Sequencing of *Wolbachia* genomes from different host

species revealed several different pairs of *cif* genes, with distinct functional domains, discriminating

them into 5 clades [23–25]. All *w*Pip genomes sequenced so far have *cif* genes from two of these

clades: clade I *cif*, in which the *cifB* gene bears a deubiquitinase domain (the tandem is thus also

called *cid*; *d* standing for deubiquitinase domain), and clade IV *cif,* in which *cifB* bears a nuclease

domain (thus called *cin*) [13,14]. Here, we use the functional-based nomenclature [19].

Before knowing the *cif* genes, models based on *Culex* crossing experiments predicted that several

factors or pairs of *mod/resc* genes were required to encode the complex crossing patterns induced

by *w*Pip [9,26]. Investigations on *cif* (*cid* and *cin*) present in *w*Pip genomes highlighted that *cin* genes

were always monomorphic while the *cid* genes were amplified and diversified in all sequenced *w*Pip

genomes [27], with up to 6 different copies of each gene within a single *Wolbachia w*Pip genome.

The different *cidA/cidB* copies in a given *w*Pip genome are called 'variants' and full set of variants

constitutes the '*cid* repertoire'. In *w*Pip genomes, polymorphism of both *cidA* and *cidB* genes are

located in two specific regions, which were named 'upstream' and 'downstream' regions that were

predicted to be involved in CidA-CidB specific interactions [27]. This has been confirmed by the

recently obtained structure of a CidA-CidB co-crystal: out of three interaction regions identified, two

perfectly match the previously identified 'upstream' and 'downstream' regions for both CidA and

CidB [20,21]. In addition to showing that CidA and CidB from *w*Pip bind together, this recent study

also showed that the different CidA-CidB variants have different binding abilities, further fueling the

toxin-antidote model for *w*Pip CI complexity [21].

Evolutionary changes of *Wolbachia* genomes, at large time scale, have been widely documented

among different host species. Comparison of *cif* genes in *Wolbachia* strains from different arthropod

hosts showed they are quite divergent, and highly subject to lateral gene transfers [28,29] so that the

congruence between *Wolbachia* and *cif* phylogenies is totally disrupted [23,24]. Such lateral transfers

have been suggested to be phage-linked, as *cif* genes are located in WO prophage regions [14,23]

and lateral transfers of phage WO and of genes in prophage WO regions have been documented [30–

33]. Yet, transposon-dependent (and thus phage-independent) transfers of *cif* genes have also been

described [28]. In contrast, although the ability of *Wolbachia* to rapidly adapt to new environmental

conditions has already been suggested [34], there are few studies of short-term *Wolbachia* evolution.

To our knowledge, the only studies linking rapid changes in phenotypes (in few host generations) to

underlying genomic variations were those reported on the 'Octomom region' in *w*Mel and *w*MelPop,

showing that variations in amplification level of this region were responsible for variations in both

Wolbachia virulence and viral protection conferred by *w*Mel [35–37].

CI patterns in *Culex* have also been shown to change over a few host generations [38]. Yet, these

changes were described before the discovery of the *cif* genes. Here, we observed a change in the CI

pattern between two isofemale lines maintained in our laboratory since 2005: Slab (*w*Pip III) and

Istanbul (*w*Pip IV) [10]. While crosses between Slab females and Ist males were compatible for 12

years (2005-2017), we observed fully incompatible crosses for the first time in 2021. Such CI shift was

an opportunity to study the underlying genetic basis of CI evolution in laboratory controlled

isofemale lines.

The emergence of a new CI phenotype could have resulted from either i) a contamination, ii) the

acquisition of a new toxin in some Istanbul's *Wolbachia*, or iii) the loss of antidotes in some Slab's

Wolbachia. We took advantage of a recent methodological development, enabling the rapid and

extensive acquisition of *cidA* and *cidB* repertoires by Nanopore Technologies sequencing [39] to

address these three hypotheses. We were able to rule out the contamination hypothesis and show

that no less than three distinct *Wolbachia* sublineages, displaying different *cidA* repertoires, now

coexist in the Slab isofemale line (but do not seem to co-infect the same individuals). We found that

the loss of *cidA* variants (i.e. antidotes) in Slab perfectly matches variations in rescuing ability. The

cidA variant whose presence/absence best matches CI variations displays original amino-acid

combinations at its interaction interface with CidB, likely resulting from recombination.

Results

From CI phenotype shift to underlying genomic bases

In 2021, for the first time, crosses between Slab females and Istanbul (Ist) males highlighted that a part of the Slab females were now fully incompatible with Istanbul males.

At least three *cidA* repertoires coexist within the Slab isofemale line

In order to decipher the genomic changes that led to this recent CI shift, we sequenced the *cidA* and *cidB* repertoires present in the *Wolbachia* infecting 4 individuals from the Ist isofemale line and 8 females from the Slab isofemale line (4 compatible and 4 incompatible), using Nanopore Technologies sequencing of *cidA* and *cidB* PCR products [39]. While *Wolbachia* present in all Ist individuals had the same *cid* repertoires (S1 Table), we found polymorphism in *cid* repertoires of Slab individuals: they all had the same *cidB* (i.e. toxins) diversity, composed of *cidB-III-ae3* and *cidB-III-ag1* variants, while *cidA* (i.e. antidotes) repertoires varied among individuals. *Wolbachia* present in compatible and incompatible Slab females all exhibited the variants *cidA-III-alpha(5)-25* and *cidA-III-gamma(3)-12,* but additional variants were found in compatible females: *cidA-III-beta(2)-25*, either alone or accompanied by *cidA-III-beta(2)-16.* The observed differences in *cidA* repertoires can be summed up by the presence/absence of two specific *cidA* regions: the upstream region *cidA-III-beta(2)* and the downstream region *cidA-III-16* (S1 Fig).

cidA repertoire variation fully explains compatibility polymorphism

In order to determine if compatibility variations were indeed linked to these specific *cidA* repertoire variations, we performed three replicates of the [Slab x Ist] cross. We isolated egg rafts from a total of 102 Slab females from the same isofemale line (28, 34 and 40 females for replicate 1, 2 and 3 respectively) and studied their hatching rates (HR, number of hatched larvae over the number of eggs laid). Eight females were determined as having produced unfertilized rafts and excluded from

further analyses. For the remaining 94 females, HRs ranged from 0% (fully incompatible raft) to

96.4%, highlighting strong differences in females' rescuing abilities. Out of these 94 females, 25 (i.e.

27%) were fully incompatible with Ist males.

Since repertoires differed in the presence/absence of two specific regions of *cidA* repertoires, we

designed primers and specifically quantified the 94 females (S2 Table). The PCR assay confirmed that

3 distinct *Wolbachia cidA* repertoires coexisted within the Slab isofemale line: (i) individuals hosting

Wolbachia bearing both the upstream '*beta(2)*' and the downstream '*16*' regions, named (β+,16+);

(ii) individuals hosting *Wolbachia* bearing only the upstream region *beta(2)*, but not the downstream

region *16*, named (β+,16-); and (iii) individuals hosting *Wolbachia* genomes bearing none of these

two regions, named (β-,16-) (S2 Table). For ease of reading, we will metonymically qualify

mosquitoes based on their *cidA* repertoires (β+,16+), (β+,16-) or (β-,16-). Out of the 94 females

analyzed, 61 were (β+,16+), 8 (β+,16-) and 25 (β-,16-) (Fig 1).

Fig 1. Intra-line polymorphism in *cidA* repertoires matches compatibility patterns. (β-,16-) females

have neither *cidA-III-beta(2)* nor *cidA-III-16* regions; (β+,16+) females have both and (β+,16-) females

have the *cidA-III-beta(2)* region but not the *cidA-III-16* one. Presence or absence of *cidA-III-beta(2)*

and *cidA-III-16* regions was determined by specific qPCRs. *N* = Number of individuals.

All but one (β-,16-) females produced egg-rafts with no hatching (the only one for which larvae

hatched corresponded to a HR of 1.8%). Such fully incompatible crosses were only observed with (β-

,16-) mothers, (β+,16+) and (β+,16-) mothers being compatible (to various extents) with Ist males (Fig

1).

cidA repertoire variations does not explain HR variability in compatible crosses

While it was long believed that CI in *Culex* was a "yes or no" phenomenon, a recent study explored HR variations in compatible crosses involving *w*Pip from different groups [40]. Here, looking at compatible crosses involving (β+,16+) or (β+,16-) Slab females and Ist males, we also found strong variations in HRs which ranged from 3.2% to 96.4% (S2 Table). While infection levels (quantified by qPCR) in full individuals were variable, with 0.11 to 10.1 *Wolbachia* genomes per host genome, with an average of 1.0 ± 0.2 (mean ± standard error), this variability did not correlate with HR variations (S2A Fig, Spearman correlation test, ρ = -0.09, p = 0.49, S2 Table). Similarly, HR variations were not correlated with the *cidA* copy numbers (total *cidA*, *cidA-III-beta* or *cidA-III-16* copy numbers) per *w*Pip-Slab genome investigated in Slab females (Spearman correlation test, ρ = -0.13, p = 0.28; ρ = 0.01, p = 0.94; and ρ = -0.17, p = 0.16 for the total *cidA*, *cidA-III-beta* and *cidA-III-16* respectively, S2B-D Figs, S2 Table).

Recombination in *cid* variants leads to changes in CidA-CidB interaction zones

Recent works disclosed that there were three interaction interfaces between CidA and CidB proteins, named interfaces I to III (Fig 2A, [20,21]). The upstream region *cidA-III-beta(2)*, present in both (β+,16+) and (β+,16-), seems key for Slab females to rescue Ist males. This region corresponds to the first interaction interface of the CidA protein (Fig 2A). Looking closer at the protein sequences at this interface, we found that CidA-III-beta(2) displays an original combination of aminoacids, possibly issued from a recombination between the CidA-alpha and CidA-gamma regions. Building a phylogenetic network of all described *cidA* upstream regions [22,27,40,41] confirmed that *beta*-like regions were rare (only found in *w*Pip group IV and in Slab to date) and issued from a recombination between the regions *cidA-gamma*-like and *cidA-alpha*-like, these last two regions being frequent and

found in all *w*Pip groups surveyed so far (Fig 2B). We also showed that the *cidA-delta* region was

recombinant.

Fig 2. Matching recombination in a CidA-CidB interaction interface I is key for rescue. (A) CidA and

CidB interact head to tail through three distinct interfaces, shown here in green, yellow and red

respectively. (B) The regions *cidA-III-beta(2)* and *cidB-IV-2*, key to in the (in)compatibility between

Slab and Istanbul isofemale lines, are both exposed in the interaction interface I and shown in green.

Alignments of the protein sequences of the interaction interface I are shown for *cidA* and *cidB* (the

sequence between the first and the last variable aminoacid belonging to interface I is represented;

Wang & Hochstrasser, pers. com.). (C) Network phylogenetic analysis of the *cidA* upstream and *cidB*

downstream region, using the Neighbor-net method on variants from [27,40,41] and this study. For

both regions, there are two major groups of variants (alpha/gamma for *cidA* and 1/3 for *cidB*), with

few intermediate recombinant variants (in green). Variants used in the present study are shown in

light green. Each edge (or set of parallel edges) corresponds to a split in the dataset and has a length

equal to the weight of the split.

The downstream region CidB-IV-2, present in *w*Pip-Ist, had previously been associated with

incompatibility when crossed with a wide range of females [22]. This region interacts with *cidA-beta*

at the interface I (Fig 2A) and also displays a pattern that could be explained by a recombination

(between CidB-1 and CidB-3 downstream regions) (Fig 2B). When looking at wider phylogenetic

scales, relying on all the previously published sequences, we found a pattern even clearer with only

two downstream CidB-1 and CidB-3 regions, identical among *w*Pip groups and CidB-IV-2 being the

only intermediate, recombinant region (Fig 2C).

cid repertoires evolution

The different *cidA* repertoires segregate into sublines

Since a new CI pattern emerged between 2017 and 2021, a closer look at Slab individuals' repertoires transmission and variation was required to better understand the precise nature of this evolution. For each of the three *cidA* repertoires found in the original isofemale Slab line, we established two isofemale sublines. We sequenced their *cidA* and *cidB* repertoires and confirmed that they were identical to those previously found for (β-,16-), (β+,16-) and (β+,16+) (S1 Fig). We confirmed that females of these new isofemale sublines had the same CI phenotype as their respective founding females when crossed with Ist males (S3Table). We also checked that they were compatible with each other in both directions showing that *cidA* repertoire polymorphism among sublines did not create incompatibilities (S4 Table).

To test whether changes in repertoires could occur at the scale of a few host generations, we genotyped the *Wolbachia* of 20 to 45 individuals of each Slab subline after about fifteen generations for both *cidA* markers and found that no change occurred since the establishment of the isofemale sublines.

We also genotyped individuals from the Slab isofemale line kept in liquid nitrogen since 2005, *i.e.* shortly after the isofemale line establishment in our insectaries. Out of 30 tested mosquitoes, there were 26 (β+,16-), 4 (β+,16+) but none of them was (β-,16-), in accordance with the hypothesis that this genotype, leading to new CI pattern, appeared afterwards.

No sign of stable coinfection

Using qPCR on Slab females, we found that total *cidA* copy numbers were variable, ranging from 2.2 to 6.2 copies per *w*Pip genome (quantifications normalized on the single copy *Wolbachia* gene *wsp*, S3A Fig). However, the distribution was bimodal with two peaks around three and five *cidA* copies (S3D Fig). The *cidA* copy number was homogeneous for females infected with a given *cidA* repertoire,

with around five *cidA* copies for all (β+,16+) and (β+,16-) individuals, and around three *cidA* copies for

all (β-,16-) (S3A Fig). We also quantified the number of *cidB* copies in 8-10 females of each repertoire

and found that *cidB* copy numbers equal *cidA* copy numbers: around five copies in both (β+,16+) and

(β+,16-) individuals and around three *cidB* copies in (β-,16-) individuals (S4 Fig). These results show

that, while Slab *cidB* repertoires do not differ between lines in terms of variants' identity, they differ

in terms of *cidB* copy numbers.

The variability in these qPCR results can be explained by qPCR intrinsic variability. The centered

distributions of data in S3 and S4 Figs support the conclusion that most individuals (if not all) from all

the different sublines [(β+,16+), (β+,16-) and (β-,16-)] are not co-infected by *Wolbachia* with different

cid repertoires.

Multiple copies of the same variant in the repertoires

qPCR results also revealed that the total number of *cidA* copies was always higher than the number

of distinct variants, showing that some variants must be present in multiple identical copies within a

given repertoire. By combining the qPCR data and relative reads coverage information from

Nanopore sequencing (e.g. a variant with two copies is covered twice as much as one with a single

copy), we estimated the copy numbers of each variant in the different *cidA* repertoires found in Slab.

We found that (β+,16+) individuals, which have a total of 5 *cidA* copies, had 2 copies of *cidA-III-*

alpha(5)-25, and a single copy of each of the three other variants (*cidA-III-beta(2)-16*, *cidA-III-beta(2)-*

25 and *cidA-III-gamma(3)-12*). (β+,16-) individuals also had 5 *cidA* copies, out of which there were 2

copies of *cidA-III-alpha(5)-25*, a single copy of *cidA-III-beta(2)-25*, and two copies of *cidA-III-*

gamma(3)-12. Lastly, (β-,16-) individuals, which had a total of 3 *cidA* copies, had 2 copies of *cidA-III-*

alpha(5)-25 and a single copy of *cidA-III-gamma(3)-12* (Fig 3).

Fig 3. *cidAB* repertoires of the *Wolbachia* infecting the 3 Slab sublines. *cidA* gene is represented in

light purple, and *cidB* in dark purple. Upstream and downstream variable regions are colored.

Regions colored in green are the key studied regions: *cidA-III-beta(2)* and *cidA-III-16.* Copy number of

each tandem was deduced by combining the relative coverages in nanopore sequencing data with

the qPCR data.

We confirmed the deduced copy numbers using qPCR specific of *cidA-III-beta(2)* and *cidA-III-16*:

quantifying those two markers in 89 single Slab females, we confirmed that (β+,16+) repertoires had

a single copy of *cidA-III-16* while other repertoires had none (S3B and S3E Figs). For *cidA-III-beta(2)*,

we found that (β+,16+) individuals had 2 copies (corresponding to 1 copy of *cidA-III-beta(2)-16* and 1

copy of *cidA-III-beta(2)-25*), while (β+,16-) individuals had a single copy (unique copy of *cidA-III-*

beta(2)-25), and (β-,16-) individuals had none of these copies (S3C and S3F Figs).

cidA and *cidB* variants are always located within tandems

Previous investigations describing the *cid* repertoire of a given *Culex* line were done by acquiring the

sequences of *cidA* and *cidB* variants separately [22,27,39–41]. Here, we also amplified and Nanopore

sequenced a region including both *cidA* and *cidB* variable regions in a single amplicon (S5 Fig). We

were thus able to reconstruct the *cidA-cidB* tandems associated in the genome, resulting in the

complete repertoires presented in Fig 3. In order to know if all *cid* variants were associated in

tandems or if they could stand alone in the genome, we compared their relative coverage when

amplified and sequenced alone *vs* amplified and sequenced in *cidA-cidB* tandems acquisitions. These

relative coverages being similar, we considered that all *cidA* and *cidB* variants were always located

within tandems. Thus, (β+,16+) and (β+,16-) sublines have 5 *cidAB* tandems while (β-,16-) lines have 3

cidAB tandems (Fig 3). Individuals of all sublines have *Wolbachia* with two copies of the *cidA-III-*

alpha(5)-25_cidB-III-ae3 tandem and one copy of the *cidA-III-gamma(3)-12_cidB-III-ag1* tandem.

(β+,16+) and (β+,16-) individuals have in addition one *cidA-III-beta(2)-25_cidB-III-ae3* tandem; but

they differ by their fifth tandem: *Wolbachia* in (β+,16+) individuals have a specific *cidA-III-beta(2)-*

16_cidB-III-ae3 while those in (β+,16-) individuals have one *cidA-III-gamma(3)-12_cidB-III-ae3* instead

(Fig 3). Our results show that the *cidB* variant *"cidB-III-ae3"* associates in tandem with four different

cidA variants (Fig. 3).

Discussion

The evolution of *Wolbachia* genomes has already been documented at large phylogenetic

scales, with multiple horizontal transfers and gene losses (e.g. [31,32,37,42–45]). This is especially

true for the *cif* genes family: numerous horizontal transfers have been described, resulting in a strong

incongruence between *cif* genes phylogeny and *Wolbachia* phylogeny [23,24,28]. However, until

now, a single set of studies had linked *Wolbachia* genotype with rapid changes in phenotypes: the

Octomom gene region in *w*Mel and *w*MelPop genomes [35–37]. While rapid shifts in CI patterns

were previously described in *Culex* mosquitoes infected with *w*Pip [38], the *cif* genes were not

described at that time, preventing to study the genetic bases of those phenotypic changes.

Furthermore, the impact of contamination in such shifts could not be ruled out. Here, we could link

changes in CI patterns with their *Wolbachia* genomic bases for the first time.

We detected a recent shift in CI pattern between Slab (*w*Pip-III) and Ist (*w*Pip-IV) lines: while they

were compatible for 12 years (from 2005 to 2017), a quarter of the crosses were fully incompatible in

2021. We looked for underlying genomic variations in their *cid* repertoires. This shift in CI could have

resulted either from a variation in Ist males' *cidB* (toxins) repertoires males or from a variation in Slab

females' *cidA* (antidotes) repertoires. We found that *cid* repertoires did not vary in Ist, but that three

different *cidA* repertoires coexisted in the same Slab isofemale line: two distinct repertoires with 5

cidA copies, named (β+,16+) and (β+,16-), which were able to rescue the modification induced by

*w*Pip-Ist, and a repertoire with three *cidA* copies, named (β-,16-), unable to rescue these

modifications. These results indicate that some *w*Pip genomes within the Slab isofemale line have

lost *cidA* variants. Indeed, screening of mosquitoes from 2005 revealed that the (β-,16-) repertoire

did not exist at that time, showing that some *cidA* variants were lost in some *w*Pip-Slab genomes

between 2017, when all crosses where compatible, and 2021, when the shift in CI was detected.

In studying the genomic bases of phenotypic changes, we showed that *cid* genes repertoires truly

evolved in *w*Pip within 4 years. While previous studies showed *cid* genes are highly variable within

and between *w*Pip groups (with up to 30 distinct variants described so far for *cidA* and 20 for *cidB*,

and different strains have different *cid* repertoires [22,27,40,41]), we found that the new (β-,16-)

repertoire is composed of *cidA* and *cidB* variants present in ancestral repertoires, associated in

identical *cidA-cidB* tandems architecture. We conclude that the shift in CI pattern results from

repertoire evolution and not from any type of contamination.

Rapid evolution of *cid* repertoires through gene losses

In 2017, approximately a quarter of the females had a total of three *cidA-cidB* tandems, showing that

two *cid* tandems had been lost. *cid* copy number distributions are bimodal, with two peaks at three

and 5 copies, and very rare intermediate 4-copy repertoire. These rare intermediates are likely due

to qPCR variability. In this case, it could either mean that (i) 4-copy repertoires were a transient state

that disappeared quickly, due to e.g. a lower fitness or drift; or (ii) that 4-copy repertoires did not

exist, meaning that two tandems were lost simultaneously.

Here, we found that all the observed repertoires were mutually compatible. This means that

mosquitoes infected by putative 4-copy repertoires would also have been compatible with all

surrounding mosquitoes, making its quick disappearance unlikely. The most plausible hypothesis is

that two tandems were lost simultaneously, making it likely that those two tandems are located next

to each other.

Literature on CI evolution suggests that *mod* factors or toxins (*cifB* variants) could degenerate first, as

their evolution is neutral for maternally-transmitted *Wolbachia* [46,47]. It is only once *cifB* have lost

its function that *cifA* (*resc* factors, or antidotes) loss would be possible [48]. In such conceptual

framework, losing a *cifA* without losing a *cifB* would lead to self-incompatibility, being an immediate

evolutionary dead-end. Independently from CI, the presence of *cidB* without an associated *cidA*

might be toxic for the host, as shown recently in insect cells [4]. Such predictions have been

corroborated by a large-scale phylogenetic study of *cif* genes, indicating that *cifB* genes were often

disrupted, while *cifA* were not [24]. However here, at a microevolutionary scale, we observed that it

was possible to decrease qualitatively the diversity in the *cidA* variants without changing qualitatively

the *cidB* variants: all (β+,16+), (β+,16-) and (β-,16-) repertoires described share the same *cidB*

variants (*cidB-III-ag1* and *cidB-III-ae3*), while they have respectively four, three or two distinct *cidA*

variants (Fig 3). Such an apparent discrepancy between predictions and our results is explained by

the fact that *w*Pip genomes contain several *cid* genes. Unlike the other systems studied to date,

where loss of *cifA* can lead to a loss of auto-rescuing ability, *w*Pip displays several *cidA* variants,

making possible to lose one and still maintain compatibility with surrounding mosquitoes. Loss of

cidA variants required for intra-line compatibility may also occur, but mosquitoes having lost such

key antidotes would be quickly wiped out by selection, an hypothesis corroborated by the mutual

compatibility between all newly-established isofemale lines, regardless of their *cid* repertoire.

Laboratory conditions could potentially facilitate such evolution by qualitative reduction of the *cidA*

repertoire. Indeed, in natural populations, multiple *cid* repertoires coexist (as shown by sequencing

and revealed by numerous crosses [22,27,41,49]), meaning that individuals are exposed to a wide

range of toxins. A greater diversity in the *cidA* repertoire enables to rescue a wider range of males

and will thus be under positive selection. While *cidA* diversity is advantageous in natural populations,

in a cage with an isofemale line, females are exposed to a restricted and fixed set of toxins. Loss of

cidA variants which are not strictly required to rescue these toxins will not be counter-selected.

Furthermore, numerous gene copies can be costly (e.g. [50]), loss of *cidAB* tandems could thus confer

a higher fitness to the *Wolbachia* with lower numbers of copies and participate to the fixation of new

Wolbachia genomes in mosquitoes, explaining the rapid *Wolbachia* turnover. It is conceivable that

the 3-copy repertoire of (β-,16-) *Wolbachia*, containing two distinct variants of *cidA* and two distinct

variants of *cidB*, is the minimum repertoire viable configuration, containing only pairs of matching

toxins and antidotes.

No lasting coinfection is involved in changes in CI patterns

in *w*Pip

A plausible explanation for the observed shift in CI could have been that Slab is co-infected by two

strains of *Wolbachia* with different compatibility types and different *cid* repertoires which are usually

co-transmitted, and that some individuals recently lost one of those strains, as observed for instance

in *Aedes albopictus* [51,52]. At first, using common MLST markers, authors found no variability at all

among *w*Pip strains [53], showing that if coinfection occurs, it must be between highly similar strains.

The development of a *w*Pip-specific MLST enabled to uncover *w*Pip polymorphism and to describe

five monophyletic groups within *w*Pip [11]. Yet, while these markers highlighted *w*Pip polymorphism,

no sign of coinfection was found either, every mosquito clearly hosting a single one of the five groups

[11]. Here, working on *cid* genes, highly variable even within a *w*Pip group, we found no clear signs of

coinfections either. We quantified several *cid* markers, and analyzed their values distribution. Had

there been coinfections of two *Wolbachia* strains with respectively three and five *cid* copies,

quantifications outcomes would have varied depending on the proportion of each *Wolbachia* in each

mosquito, leading to a unimodal distribution centered on an intermediate value. For the three

distinct *cid* markers analyzed in 89 females, quantifications displayed a bimodal distribution (S3D-F

Figs), with few intermediate values certainly due to technical variation in qPCR measures. While

coinfection had to happen at some point (all the *Wolbachia* within an individual cannot be wiped out

and replaced by another *Wolbachia* instantly), it had to be brief with no lasting coinfection within a

single host individual. This could be explained by two alternative hypotheses: (i) an exclusion

between the two distinct genotypes (3 and 5-copies), for any non-CI related reason, or (ii) a small

effective population of *Wolbachia* (number of *Wolbachia* transmitted from mother to offspring) in

*w*Pip.

Variability in hatching rates in compatible crosses is not

explained by females' *cid* repertoires

CI has long been described as a 'yes or no' phenomenon in *Cx pipiens*, meaning that crosses were

either compatible or fully incompatible [9]. Yet, having a closer look at hatching rates, we recently

showed that they could be highly variable in so-called compatible crosses [40]. We further showed

that reduced hatching rates were associated with cytological signs of canonical CI [40]. These

phenomena were suggested to be linked with *cid* repertoires, as males infected with *w*Pip-IV, having

distinct *cidB* variants from other *w*Pip groups, induced more intermediate CI than other males.

Here, studying the cross between Slab females (*w*Pip-III) and Ist males (*w*Pip-IV), we also found

intermediate HRs in 'compatible' crosses. While on/off compatibility was clearly explained by the

presence/absence of specific *cidA* variants, these intermediate hatching rates did not correlate with

any of the *cidA* markers studied, nor with *Wolbachia* infection rate.

Yet, once again, intermediate HRs were found in crosses involving males infected with *w*Pip-IV. By

comparing *cidA* and *cidB* sequences from different *w*Pip groups, we showed that the second

interaction interface differs between CidB belonging to *w*Pip-IV and CidB from other groups, with no

variation in the interface II of CidA variants (S6 Fig). This could mean that CidB variants belonging to

*w*Pip group IV never perfectly bind CidA variants. Such lack of match at interface II could result in a

more labile CidA-CidB interaction leading to intermediate HR values.

Recombination seems key to the emergence of new CI pattern

A specific *cidB* region, named *cidB-IV-2*, was previously correlated with strong CI-inducing abilities in males [22,27]. This region is present in the *w*Pip infecting the Ist males, studied here. We showed that this specific region is issued from a recombination between the two main groups of *cidB* downstream regions which are shared by all sequenced *w*Pip groups (Fig 2B).

Furthermore, it was previously shown that the only females able to rescue the modifications induced by Ist males were females infected with *w*Pip-IV and Slab females (*w*Pip-III, studied here) [22,27]. We found that the rescuing ability of Slab females depended on the presence of a specific *cidA* upstream region, *cidA-beta*, all females having this region being able to rescue Ist males, while females having lost this region were not. This variant was also generated by a recombination between two main groups of upstream regions shared by all *w*Pip groups. This time, there were two types of recombinants, the *cidA-beta* regions and the *cidA-IV-delta* region (Fig 2B). When looking at the repertoires found in females infected with *w*Pip-IV, which can all rescue Ist males, we found that all of them had one recombinant region in their repertoire (either *cidA-beta* or *cidA-IV-delta*). This means that all females able to rescue the toxicity of the recombinant region *cidB-IV-2* have a recombinant antidote in their repertoire. Since the upstream *cidA* region (e.g. *beta* or *delta*) interacts with the downstream *cidB* region (e.g. *cidB-IV-2*) in the interaction interface I [21], this finding suggests a recombinant-for-recombinant binding (Fig 2), in which recombinant regions can only bind similarly recombinant regions. New CI phenotypes could thus emerge through the recombination of existing variants.

In conclusion, we found that CI evolved in less than four years within a *Culex* isofemale line. Such evolution is linked to the loss of some *cidA* (antidotes) variants in their *cid* repertoire. Our results suggest that incompatibilities could emerge as a two-step process: local changes in CI repertoires,

while maintaining compatibility with surrounding mosquitoes, then migration and secondary contact

with incompatible lines. Indeed, here, we showed that all repertoires present within a cage were

mutually compatible: changes in repertoires led to a change in CI pattern only when crossed with an

external line. We demonstrated that new compatibility types can emerge rapidly, as a result of the

quick evolution of *Wolbachia* genomes. A common point in the few studies addressing *Wolbachia*

microevolution is that both Octomom in *w*Mel [35–37], and *cid* genes in *w*Pip are amplified genomic

regions, suggesting a strong role of gene amplification in the speed of *Wolbachia* genome evolution.

Such rapid evolution has to be considered when releasing *Wolbachia*-infected mosquitoes, inducing

CI, for vector control purposes.

Materials and methods

444 All DNA extractions in this study were done following the cetrimonium bromide (CTAB) protocol [54].

Isofemale lines used in this study

All the mosquito lines used were isofemale lines, *i.e.* lines created by rearing the offspring resulting

from a single egg-raft, and thus from a single female. We used the following lines: Slab, infected with

Wolbachia from the phylogenetic group *w*Pip-III, and Istanbul (Ist) infected with *Wolbachia* from the

phylogenetic group *w*Pip-IV. Both lines have been established at the lab in 2005.

We also used 6 newly-established isofemale (sub)lines: 2 "(β+,16+)" lines, bearing both the *cidA-III-*

beta and the *cidA-III-16* regions; 2 "(β+,16-)" lines bearing only the *cidA-III-beta* marker, and 2 "(β-

,16-)" lines bearing none of those markers. Briefly, for isofemale line establishment, a cage

containing the Slab isofemale line was blood fed. After five days, females were isolated in glasses for

egg-laying. After egg laying, DNA was extracted and females were genotyped for *cidA-III-beta* and

cidA-III-16 (S5 Table). For each genotype, the 2 most numerous progenies were kept to establish two

separate isofemale lines. All isofemale lines were reared in 65 dm^3 screened cages, in a single room

maintained at 26°C, under a 12h light/ 12h dark cycle. Larvae were fed with a mixture of shrimp

powder and rabbit pellets, and adults were fed on honey solution. Females were fed with turkey

blood, using a Hemotek membrane feeding system (Discovery Workshops, UK), to enable them to lay

eggs.

Crosses

For each cross performed, around 100 virgin females were put in a cage with around 50 virgin males.

After five days in cages, females were fed a blood meal and five days later, individual females were

isolated for egg-laying. This enabled to have a precise correspondence between a female and its egg-

raft. Egg-rafts were deposited into 24-well plates, a picture of the eggs was taken using a Leica

camera, and eggs were counted using ImageJ. Two days after egg laying, larvae issued from each egg

raft were counted manually. HR was calculated as the ratio between the total number of larvae and

the total number of eggs. Egg rafts with a null HR were mounted on a slide and checked with a

binocular magnifier to ensure that eggs were fertilized, and that null HR resulted from CI rather than

unfertilization.

Repertoire acquisition through Nanopore sequencing of

PCR products

We acquired the *cidA* and *cidB* repertoires of 8 individuals from the Slab isofemale line, 4 individuals

from the Ist isofemale line and of 2 individuals per newly-established isofemale line, following the

pipeline described in [39]. We also acquired *cidAB* combinations for all newly established isofemale

lines. Briefly, DNA was extracted from the individuals. The target sequence was amplified using the

adequate primer set (S5 Table, S6 Fig). PCR products were then sequenced by MGX Platform

(Montpellier) by Nanopore sequencing, using a MinION (Oxford Nanopore Technologies). Repertoires

were then acquired following the pipeline described in [39]. For *cidAB* combinations, the pipeline was

adapted: *cidA* and *cidB* repertoires were first obtained, then combinations between all *cidA* and *cidB*

were artificially created and used as a reference base to run the pipeline.

Wolbachia genotyping

Classic PCR

To genotype individuals, DNA was extracted. Presence / absence of both *cidA-III-16* and *cidA-III-beta*

were screened using specific PCRs (S5 Table). For individuals negative for both markers, a PK1 PCR

(S5 Table) was used to control DNA quality. This was done either on adult females or fourth instar

larvae.

Quantitative PCR

Real-time quantitative PCR were run using the LightCycler 480 system (Roche). All DNA samples were

analyzed in triplicates, for each quantification used. A total of 89 females were used for qPCR assays.

The total number of *cidA* copies, along with the number of *cidA-III-beta* and *cidA-III-16* regions copies

were quantified using relative quantitative PCR, corrected by *wsp,* a single-copy *Wolbachia* gene

(following [55], primers in S5 Table). Since *wsp* is present in a single copy per haploid genome, the

ratio of the target and *wsp* signal could be used to estimate the number of target copies per

Wolbachia genome, correcting for *Wolbachia* infection level and DNA quality. For *cidA-III-beta* qPCR,

quantifications were normalized on three distinct individuals with a single *cidA-III-beta* copy.

The infection level *i.e.* the number of *Wolbachia* bacteria per female was assessed by qPCR using a

wsp PCR normalized on the *Culex*-specific single-copy *ace-2* locus [56]. The ratio between *wsp* and

ace-2 signals can be used to estimate the relative number of *Wolbachia* genomes per *Culex* genome.

For *cidA-III-16/wsp* and *cidA/wsp*, the following protocol was used: 3 min at 95°C, 45 cycles of 95°C

for 10 sec, 60°C for 20 sec, and 72°C for 20 sec. For *cidA-III-beta/wsp* and *ace2/wsp*, the protocol was

3 min at 95°C, 45 cycles of 95°C for 3 sec, 62°C for 10 sec, and 72°C for 15 sec. Primers concentration

were respectively 0.6 μM for *cidA-III-16* and *cidA,* 0.4 μM for *wsp* and *ace2* and 1 μM for *cidA-III-beta*.

Variant copy number analysis

511   Variant copy numbers were obtained using both the *cidA* qPCR data and the relative coverage of

512   Nanopore sequencing. Knowing the total number of *cidA* copies, we deduced each variant copy

513   number. For instance, if there are three *cidA* copies and if, out of the two variants present, the

514   coverage of the first one is twice the coverage of the second, it is concluded that there are two

515   copies of the first one and one copy of the second.

Grouping analysis of upstream/downstream regions of variants

519   Phylogenetic analysis of variants was made using all the variants already published (in [22,27,40,41]).

520   Network phylogenies of the two regions of interest (the upstream variable region of *cidA* and the

521   downstream region of *cidB;* corresponding to nucleotides 1-800 for *cidA* and 1201-end for *cidB*) were

522   made using Splitstree [57].

Nomenclature of *cid* variants

The nomenclature established by [27] was updated, since it could result in identical sequences being

called differently in different *w*Pip groups. The basics of the previous nomenclature were kept:

variants names are composed of the name of their upstream region (in Greek letters for *cidA* and in

Latin letters for *cidB*) and of that of their downstream region (digits). A fixed limit was established

between the upstream and downstream regions for both genes (downstream regions starting at

nucleotides 801 and 1201 for *cidA* and *cidB* respectively). A phylogenetic network showed there were

4 clusters of *cidA* upstream regions (Fig 2B), regions within a cluster differing by at most 3 SNPs. We

renamed regions so that all regions belonging to the same cluster were named by the same Greek

letter (e.g. all named alpha). When SNPs were present, to distinguish sequences, we introduced a

number between brackets [e.g. alpha(1)]. The same was done for *cidB* downstream regions. Similarly,

cidA downstream regions and *cidB* upstream regions were renamed so that distinct regions had

different names, and identical regions had identical names, regardless of the *w*Pip-group in which

they are found.

In all variants names, the group in which the variant is found is mentioned [e.g. cidA-III-alpha(5)-25]

is found in a *w*Pip-III].

The correspondence between previous names and names with this new nomenclature is given in S6

Table.

Statistical analyses and graphical representation

544 All statistical analyses were run using R version 4.2.0 [58]. Correlation analyses were done using a

545 Pearson correlation test. All boxplots and histograms were done using ggplot2 [59], and sequence

546 representation was done using the function msaPrettyPrint, from the msa package [60].

Data availability statement

549 The nucleotide sequences of the new *cidA* and *cidB* variants were deposited in GenBank and

550 accession numbers are provided in S7 Table. The authors declare that all other data supporting the

551 findings of this study are available within the article and Supporting information.

Acknowledgments

We thank Michael Turelli and Nicole Pasteur for thorough reading and helpful comments on the manuscript; and Wei Wang and Mark Hochstrasser for inputs on interface residuals. Sequencing data were generated on the Montpellier GénomiX platform.

References

1. Hilgenboecker K, Hammerstein P, Schlattmann P, Telschow A, Werren JH. How many species are infected with *Wolbachia*? - A statistical analysis of current data. FEMS Microbiol Lett. 2008;281: 215–220. doi:10.1111/j.1574-6968.2008.01110.x

2. Zug R, Hammerstein P. Still a host of hosts for *Wolbachia*: Analysis of recent data suggests that 40% of terrestrial arthropod species are infected. PLoS One. 2012;7: 7–9. doi:10.1371/journal.pone.0038544

3. Weinert LA, Araujo-Jnr E V., Ahmed MZ, Welch JJ. The incidence of bacterial endosymbionts in terrestrial arthropods. Proc R Soc B Biol Sci. 2015;282: 3–8. doi:10.1098/rspb.2015.0249

4. Horard B, Terretaz K, Gosselin-Grenet AS, Sobry H, Sicard M, Landmann F, et al. Paternal transmission of the *Wolbachia* CidB toxin underlies cytoplasmic incompatibility. Curr Biol. 2022;32: 1319-1331.e5. doi:10.1016/j.cub.2022.01.052

5. Kaur R, Leigh BA, Ritchie IT, Bordenstein SR. The Cif proteins from *Wolbachia* prophage WO modify sperm genome integrity to establish cytoplasmic incompatibility. PLOS Biol. 2022;20: e3001584. doi:10.1371/journal.pbio.3001584

6. Duron O, Lagnel J, Raymond M, Bourtzis K, Fort P, Weill M. Transposable element polymorphism of *Wolbachia* in the mosquito *Culex pipiens*: evidence of genetic diversity, superinfection and recombination. Mol Ecol. 2005;14: 1561–1573. doi:10.1111/j.1365-294X.2005.02495.x

7. Laven H. Crossing experiments with *Culex* strains. Evolution (N Y). 1951;5: 370–375.

doi:10.2307/2405682

8.   Laven H. Speciation and evolution in *Culex pipiens*. In: Wright J, Pal R, editors. Genetics of

Insect Vectors of Disease. Elsevier: Amsterdam; 1967. pp. 251–275.

9.   Atyame CM, Labbé P, Dumas E, Milesi P, Charlat S, Fort P, et al. *Wolbachia* divergence and the

evolution of cytoplasmic incompatibility in *Culex pipiens*. PLoS One. 2014;9: e87336.

doi:10.1371/journal.pone.0087336

10.   Duron O, Bernard C, Unal S, Berthomieu A, Berticat C, Weill M. Tracking factors modulating

cytoplasmic incompatibilities in the mosquito *Culex pipiens*. Mol Ecol. 2006;15: 3061–3071.

doi:10.1111/j.1365-294X.2006.02996.x

11.   Atyame CM, Delsuc F, Pasteur N, Weill M, Duron O. Diversification of *Wolbachia*

endosymbiont in the *Culex pipiens* mosquito. Mol Biol Evol. 2011;28: 2761–2772.

doi:10.1093/molbev/msr083

12.   Beckmann JF, Fallon AM. Detection of the *Wolbachia* protein WPIP0282 in mosquito

spermathecae: Implications for cytoplasmic incompatibility. Insect Biochem Mol Biol. 2013;43:

867–878. doi:10.1016/j.ibmb.2013.07.002

13.   Beckmann JF, Ronau JA, Hochstrasser M. A *Wolbachia* deubiquitylating enzyme induces

cytoplasmic incompatibility. Nat Microbiol. 2017;2: 1–7. doi:10.1038/nmicrobiol.2017.7

14.   LePage DP, Metcalf JA, Bordenstein SR, On J, Perlmutter JI, Shropshire JD, et al. Prophage WO

genes recapitulate and enhance *Wolbachia*-induced cytoplasmic incompatibility. Nature.

2017;543: 243–247. doi:10.1038/nature21391

15.   Beckmann JF, Bonneau M, Chen H, Hochstrasser M, Poinsot D, Merçot H, et al. The Toxin–

Antidote Model of Cytoplasmic Incompatibility: Genetics and Evolutionary Implications.

Trends Genet. 2019;35: 175–185. doi:10.1016/j.tig.2018.12.004

16.   Namias A, Sicard M, Weill M, Charlat S. From *Wolbachia* genomics to phenotype: molecular

models of cytoplasmic incompatibility must account for the multiplicity of compatibility types.

Curr Opin Insect Sci. 2022;49: 78–84. doi:10.1016/j.cois.2021.12.005

17. Shropshire JD, Leigh B, Bordenstein SR, Duplouy A, Riegler M, Brownlie JC, et al. Models and

Nomenclature for Cytoplasmic Incompatibility: Caution over Premature Conclusions – A

Response to Beckmann et al. Trends Genet. 2019;35: 397–399. Available:

https://doi.org/10.1016/j.tig.2019.03.004

18. Shropshire JD, Bordenstein SR. Two-By-One model of cytoplasmic incompatibility: Synthetic

recapitulation by transgenic expression of *cifA* and *cifB* in *Drosophila*. PLoS Genet. 2019;15: 1–

20. doi:10.1371/journal.pgen.1008221

19. Beckmann JF, Bonneau M, Chen H, Hochstrasser M, Poinsot D, Merçot H, et al. Caution Does

Not Preclude Predictive and Testable Models of Cytoplasmic Incompatibility: A Reply to

Shropshire et al. Trends Genet. 2019;35: 399–400. doi:10.1016/j.tig.2019.03.002

20. Xiao Y, Chen H, Wang H, Zhang M, Chen X, Berk JM, et al. Structural and mechanistic insights

into the complexes formed by *Wolbachia* cytoplasmic incompatibility factors. Proc Natl Acad

Sci U S A. 2021;118: 1–10. doi:10.1073/pnas.2107699118

21. Wang H, Xiao Y, Chen X, Zhang M, Sun G, Wang F, et al. Crystal Structures of *Wolbachia* CidA

and CidB Reveal Determinants of Bacteria-induced Cytoplasmic Incompatibility and Rescue.

Nat Commun. 2022;13: 1–11. doi:10.1038/s41467-022-29273-w

22. Bonneau M, Caputo B, Ligier A, Caparros R, Unal S, Perriat-Sanguinet M, et al. Variation in

Wolbachia cidB gene, but not *cidA*, is associated with cytoplasmic incompatibility *mod*

phenotype diversity in *Culex pipiens*. Mol Ecol. 2019;28: 4725–4736. doi:10.1111/mec.15252

23. Lindsey ARI, Rice DW, Bordenstein SR, Brooks AW, Bordenstein SR, Newton ILG. Evolutionary

Genetics of Cytoplasmic Incompatibility Genes *cifA* and *cifB* in Prophage WO of *Wolbachia*.

Genome Biol Evol. 2018;10: 434–451. doi:10.1093/gbe/evy012

24. Martinez J, Klasson L, Welch JJ, Jiggins FM. Life and Death of Selfish Genes: Comparative

Genomics Reveals the Dynamic Evolution of Cytoplasmic Incompatibility. Mol Biol Evol.

2021;38: 2–15. doi:10.1093/molbev/msaa209

25. Bing XL, Zhao DS, Sun JT, Zhang KJ, Hong XY, Sloan D. Genomic Analysis of *Wolbachia* from

17. Shropshire JD, Leigh B, Bordenstein SR, Duplouy A, Riegler M, Brownlie JC, et al. Models and

Nomenclature for Cytoplasmic Incompatibility: Caution over Premature Conclusions – A

Response to Beckmann et al. Trends Genet. 2019;35: 397–399. Available:

https://doi.org/10.1016/j.tig.2019.03.004

18. Shropshire JD, Bordenstein SR. Two-By-One model of cytoplasmic incompatibility: Synthetic

recapitulation by transgenic expression of *cifA* and *cifB* in *Drosophila*. PLoS Genet. 2019;15: 1–

20. doi:10.1371/journal.pgen.1008221

19. Beckmann JF, Bonneau M, Chen H, Hochstrasser M, Poinsot D, Merçot H, et al. Caution Does

Not Preclude Predictive and Testable Models of Cytoplasmic Incompatibility: A Reply to

Shropshire et al. Trends Genet. 2019;35: 399–400. doi:10.1016/j.tig.2019.03.002

20. Xiao Y, Chen H, Wang H, Zhang M, Chen X, Berk JM, et al. Structural and mechanistic insights

into the complexes formed by *Wolbachia* cytoplasmic incompatibility factors. Proc Natl Acad

Sci U S A. 2021;118: 1–10. doi:10.1073/pnas.2107699118

21. Wang H, Xiao Y, Chen X, Zhang M, Sun G, Wang F, et al. Crystal Structures of *Wolbachia* CidA

and CidB Reveal Determinants of Bacteria-induced Cytoplasmic Incompatibility and Rescue.

Nat Commun. 2022;13: 1–11. doi:10.1038/s41467-022-29273-w

22. Bonneau M, Caputo B, Ligier A, Caparros R, Unal S, Perriat-Sanguinet M, et al. Variation in

Wolbachia cidB gene, but not *cidA*, is associated with cytoplasmic incompatibility *mod*

phenotype diversity in *Culex pipiens*. Mol Ecol. 2019;28: 4725–4736. doi:10.1111/mec.15252

23. Lindsey ARI, Rice DW, Bordenstein SR, Brooks AW, Bordenstein SR, Newton ILG. Evolutionary

Genetics of Cytoplasmic Incompatibility Genes *cifA* and *cifB* in Prophage WO of *Wolbachia*.

Genome Biol Evol. 2018;10: 434–451. doi:10.1093/gbe/evy012

24. Martinez J, Klasson L, Welch JJ, Jiggins FM. Life and Death of Selfish Genes: Comparative

Genomics Reveals the Dynamic Evolution of Cytoplasmic Incompatibility. Mol Biol Evol.

2021;38: 2–15. doi:10.1093/molbev/msaa209

25. Bing XL, Zhao DS, Sun JT, Zhang KJ, Hong XY, Sloan D. Genomic Analysis of *Wolbachia* from

Laodelphax striatellus (Delphacidae, Hemiptera) Reveals Insights into Its "Jekyll and Hyde"

Mode of Infection Pattern. Genome Biol Evol. 2020;12: 3818–3831. doi:10.1093/gbe/evaa006

26. Nor I, Engelstädter J, Duron O, Reuter M, Sagot MF, Charlat S. On the genetic architecture of

cytoplasmic incompatibility: Inference from phenotypic data. Am Nat. 2013;182: E15–E24.

doi:10.1086/670612

27. Bonneau M, Atyame CM, Beji M, Justy F, Cohen-Gonsaud M, Sicard M, et al. *Culex pipiens*

crossing type diversity is governed by an amplified and polymorphic operon of *Wolbachia*. Nat

Commun. 2018;9: 319. doi:10.1038/s41467-017-02749-w

28. Cooper BS, Vanderpool D, Conner WR, Matute DR, Turelli M. *Wolbachia* acquisition by

Drosophila yakuba-clade hosts and transfer of incompatibility loci between distantly related

Wolbachia. Genetics. 2019;212: 1399–1419.

29. Madhav M, Parry R, Morgan JAT, James P. *Wolbachia* Endosymbiont of the Horn Fly

(*Haematobia irritans*): a Supergroup A Strain with Multiple Horizontally Acquired Cytoplasmic

Incompatibility Genes. Appl Environ Microbiol. 2020;86: e02589-19.

30. Bordenstein SR, Wernegreen JJ. Bacteriophage flux in endosymbionts (*Wolbachia*): Infection

frequency, lateral transfer, and recombination rates. Mol Biol Evol. 2004;21: 1981–1991.

doi:10.1093/molbev/msh211

31. Wang N, Jia S, Xu H, Liu Y, Huang D. Multiple Horizontal Transfers of Bacteriophage WO and

Host *Wolbachia* in Fig Wasps in a Closed Community. Front Microbiol. 2016;7: 1–10.

doi:10.3389/fmicb.2016.00136

32. Hill T, Unckless RL, Perlmutter JI. Positive Selection and Horizontal Gene Transfer in the

Genome of a Male-Killing *Wolbachia*. Mol Biol Evol. 2022;39: 1–17.

doi:10.1093/molbev/msab303

33. Gao S, Ren YS, Su CY, Zhu DH. High Levels of Multiple Phage WO Infections and Its

Evolutionary Dynamics Associated With *Wolbachia*-Infected Butterflies. Front Microbiol.

2022;13: 1–12. doi:10.3389/fmicb.2022.865227

34.   Le Clec'h W, Dittmer J, Raimond M, Bouchon D, Sicard M. Phenotypic shift in *Wolbachia*

657   virulence towards its native host across serial horizontal passages. Proc R Soc B Biol Sci.

658   2017;284. doi:10.1098/rspb.2017.1076

659   35.   Chrostek E, Marialva MSP, Esteves SS, Weinert LA, Martinez J, Jiggins FM, et al. *Wolbachia*

660   Variants Induce Differential Protection to Viruses in *Drosophila melanogaster*: A Phenotypic

661   and Phylogenomic Analysis. PLoS Genet. 2013;9. doi:10.1371/journal.pgen.1003896

662   36.   Chrostek E, Teixeira L. Mutualism Breakdown by Amplification of *Wolbachia* Genes. PLoS Biol.

663   2015;13: 1–22. doi:10.1371/journal.pbio.1002065

664   37.   Duarte EH, Carvalho A, López-Madrigal S, Costa J, Teixeira L. Forward genetics in *Wolbachia*:

665   Regulation of *Wolbachia* proliferation by the amplification and deletion of an additive

666   genomic island. PLoS Genetics. 2021. doi:10.1371/journal.pgen.1009612

667   38.   Duron O, Bernard J, Atyame CM, Dumas E, Weill M. Rapid evolution of *Wolbachia*

668   incompatibility types. Proc R Soc B Biol Sci. 2012;279: 4473–4480.

669   doi:10.1098/rspb.2012.1368

670   39.   Namias A, Sahlin K, Makoundou P, Bonnici I, Sicard M. Nanopore sequencing enables

671   multigenic family reconstruction despite highly frequent PCR-induced recombination. bioRxiv.

672   2022; 2022.11.14.516392. doi:10.1101/2022.11.14.516392

673   40.   Sicard M, Namias A, Perriat-Sanguinet M, Carron E, Unal S, Altinli M, et al. Cytoplasmic

674   incompatibility variations in relation with *Wolbachia cid* genes divergence in *Culex pipiens*.

675   MBio. 2021;12: e02797-20. doi:10.1128/mBio.02797-20

676   41.   Bonneau M, Landmann F, Labbé P, Justy F, Weill M, Sicard M. The cellular phenotype of

677   cytoplasmic incompatibility in *Culex pipiens* in the light of *cidB* diversity. PLoS Pathog.

678   2018;14: e1007364. doi:10.1371/journal.ppat.1007364

679   42.   Scholz M, Albanese D, Tuohy K, Donati C, Segata N, Rota-Stabelli O. Large scale genome

680   reconstructions illuminate *Wolbachia* evolution. Nat Commun. 2020;11. doi:10.1038/s41467-

681   020-19016-0

43.   Foster J, Ganatra M, Kamal I, Ware J, Makarova K, Ivanova N, et al. The *Wolbachia* genome of

Brugia malayi: Endosymbiont evolution within a human pathogenic nematode. PLoS Biol.

2005;3: 0599–0614. doi:10.1371/journal.pbio.0030121

44.   Newton ILG, Clark ME, Kent BN, Bordenstein SR, Qu J, Richards S, et al. Comparative genomics

of two closely related *Wolbachia* with different reproductive effects on hosts. Genome Biol

Evol. 2016;8: 1526–1542. doi:10.1093/gbe/evw096

45.   Wang GH, Sun BF, Xiong TL, Wang YK, Murfin KE, Xiao JH, et al. Bacteriophage WO can

mediate horizontal gene transfer in endosymbiotic *Wolbachia* genomes. Front Microbiol.

2016;7: 1–16. doi:10.3389/fmicb.2016.01867

46.   Prout T. Some evolutionary possibilities for a microbe which causes incompatibility in its host.

Evolution (N Y). 1994;48: 909–911.

47.   Turelli M. Evolution of incompatibility-inducing microbes and their hosts. Evolution (N Y).

1994;48: 1500–1513. doi:10.1111/j.1558-5646.1994.tb02192.x

48.   Charlat S, Calme C, Merçot H. On the *mod resc* model and the evolution of *Wolbachia*

compatibility types. Genetics. 2001;159: 1415–1422. doi:10.1093/genetics/159.4.1415

49.   Atyame CM, Duron O, Tortosa P, Pasteur N, Fort P, Weill M. Multiple *Wolbachia* determinants

control the evolution of cytoplasmic incompatibilities in *Culex pipiens* mosquito populations.

Mol Ecol. 2011;20: 286–298. doi:10.1111/j.1365-294X.2010.04937.x

50.   Assogba BS, Milesi P, Djogbénou LS, Berthomieu A, Makoundou P, Baba-Moussa LS, et al. The

ace-1 Locus Is Amplified in All Resistant *Anopheles gambiae* Mosquitoes: Fitness

Consequences of Homogeneous and Heterogeneous Duplications. PLoS Biol. 2016;14: 1–26.

doi:10.1371/journal.pbio.2000618

51.   Kambhampati S, Rai KS, Burgun SJ. Unidirectional Cytoplasmic Incompatibility in the

Mosquito, *Aedes albopictus*. Evolution. 1993. p. 673. doi:10.2307/2410079

52.   Sinkins SP, Braig HR, O'Neill SL. *Wolbachia pipientis*: Bacterial density and unidirectional

cytoplasmic incompatibility between infected populations of *Aedes albopictus*. Exp Parasitol.

1995;81: 284–291. doi:10.1006/expr.1995.1119

53.   Baldo L, Hotopp JCD, Jolley KA, Bordenstein SR, Biber SA, Choudhury RR, et al. Multilocus

sequence typing system for the endosymbiont *Wolbachia pipientis*. Appl Environ Microbiol.

2006;72: 7098–7110. doi:10.1128/AEM.00731-06

54.   Rogers SO, Bendich AJ. Extraction of total cellular DNA from plants, algae and fungi. Plant

Molecular Biology Manual. Springer Netherlands; 1994. pp. 183–190. doi:10.1007/978-94-

011-0511-8_12

55.   Berticat C, Rousset F, Raymond M, Berthomieu A, Weill M. High *Wolbachia* density in

insecticide-resistant mosquitoes. Proc R Soc B Biol Sci. 2002;269: 1413–1416.

doi:10.1098/rspb.2002.2022

56.   Weill M, Berticat C, Raymond M, Chevillon C. Quantitative polymerase chain reaction to

estimate the number of amplified esterase genes in insecticide-resistant mosquitoes. Anal

Biochem. 2000;285: 267–270. doi:10.1006/abio.2000.4781

57.   Bryant D, Moulton V. Neighbor-Net: An Agglomerative Method for the Construction of

Phylogenetic Networks. Mol Biol Evol. 2004;21: 255–265. doi:10.1093/molbev/msh018

58.   R Core Team. R: A language and environment for statistical computing. http://www.R-

project.org/. In: R Foundation for Statistical Computing, Vienna, Austria. 2017.

59.   Wickham H. Elegant Graphics for Data Analysis: ggplot2. Applied Spatial Data Analysis with R.

2008.

60.   Bodenhofer U, Bonatesta E, Horejš-Kainrath C, Hochreiter S. Msa: An R package for multiple

sequence alignment. Bioinformatics. 2015;31: 3997–3999. doi:10.1093/bioinformatics/btv494

Supporting information

S1 Fig. *cidA* and *cidB* **variants observed in the Slab isofemale line, obtained by Nanopore**

sequencing. For each variant, the two previously described variable regions are shown: the upstream

region, named with a Greek letter for *cidA* and with a Latin letter for *cidB*, and the downstream

region, named with a digit for both genes. *cidA* regions whose presence differ among repertoires are

represented in green. *cidB* repertoires are common to all repertoires. Females infected with (β+, 16+)

or a (β+, 16-) *Wolbachia* are able to rescue males infected with *w*Pip-Ist (compatible), while females

infected with (β-, 16-) *Wolbachia* cannot (incompatible).

S2 Fig. Variations of hatching rates in compatible crosses between Slab females and Ist males, in

relation to variations in: A) *Wolbachia* infection level; and numbers of copies of B) total *cidA*; C)

***cidA-III-beta(2)*; and D) *cidA-III-16*.** None of these variations are significantly correlated with hatching

rates (Spearman correlation coefficient, P = 0.49, 0.28, 0.94 and 0.16 for A, B, C and D respectively).

S3 Fig. Variations of *cidA* copy numbers in *Wolbachia* genomes of Slab females. A), B) and C) show

the numbers of copies of total *cidA, cidA-III-16* and *cidA-III-beta(2)* regions, respectively in females

with repertoires (β-,16-), (β+,16-), and (β+,16+). D), E) and F) show the distribution of *cidA, cidA-III-16*

and *cidA-III-beta(2)* copy numbers. All histograms show either bi or trimodal distributions. All copy

numbers were quantified using quantitative PCR relative to the single copy *Wolbachia* gene *wsp*.

S4 Fig. Variations of *cidB* copy numbers in *Wolbachia* genomes of Slab females. *cidB* copy numbers

equal *cidA* copy numbers, (β-,16-) *w*Pip having around 3 *cidB* copies, while (β+,16-) and (β+,16+) have

around 5 *cidB* copies. All copy numbers were quantified using quantitative PCR relative to the single

copy *Wolbachia* gene *wsp*.

S5 Fig. PCR primers used to analyze sequence variation in *cid* genes. PCRs amplifying the full *cidA*

gene (burgundy arrows), variable regions of the *cidB* gene (green arrows) and the *cidAB* tandem

(purple arrows) were used for nanopore sequencing of repertoires. PCRs specific of *cidA-III-beta(2)*

(yellow) and *cidA-III-16* (orange) were used both in classic PCR and qPCR to determine the

presence/absence of these regions.

S6 Fig. Sequence variation at the CidA-CidB interaction interface II. Protein sequences are shown for

the span of the interaction interface II (position of the first and last aminoacids of this interface were

obtained thanks to Wang & Hochstrasser, pers. com.). Polymorphic positions are shown with a grey

background.

This interface is polymorphic in CidB, the *w*Pip-IV CidB showing a different sequence from all other

groups (sequences shown here are representative of all sequenced variants). In contrast, CidA

variants are identical in this interaction interface in all sequenced variants whatever their groups.

S1 Table. *cid* repertoires of the Istanbul isofemale line. Nanopore sequencing of *cid* PCR products

gave identical repertoires for 4 distinct Ist individuals.

S2 Table. Crosses between Slab females and Ist males. For each female, we determined the hatching

rate (HR) (i.e. ratio between hatched larvae and eggs). The specific upstream *cidA-III-beta(2)* and

downstream *cidA-III-16* regions were quantified in each female and normalized over the single copy

Wolbachia gene *wsp*. The total *cidA* and *cidB* copy number was also quantified and normalized over

wsp. Infection rates were assessed by quantifying the single copy gene *wsp* and normalizing it on the

Cx. pipiens single copy gene *ace2*.

S3 Table. Crosses between Slab females from each of the 6 newly established isofemale sublines [2

(β+,16+), 2 (β+,16-) and 2 (β-,16-)] and Istanbul males. Hatching rates and number of hatched larvae

are shown for each female. Eggs were only counted in compatible crosses, involving (β+,16+) and

(β+,16-) females. All females were screened for the presence/absence of two regions of the

polymorphic gene *cidA* (*cidA-III-beta(2)* and *cidA-III-16*) to confirm that they had the same repertoire

as the female from which the isofemale subline was founded. NF = non fertilized

S4 Table. Reciprocal crosses between a (β+,16+) and a (β-,16-) isofemale subline. Number of eggs

and larvae and resulting hatching rates (HR). NF = non fertilized

S5 Table. PCR primers and programs used in this study. qPCR programs are described in the main

text.

S6 Table. Correspondence between variants names in the old (Bonneau et al, Nat Com 2018) and in

current nomenclature.

S7 Table. GenBank accession numbers of new *cidA* and *cidB* variants used in this study.