Swimming ability and flagellar motility of sperm packets of the volvocine green alga *Pleodorina starrii*

Short Title: Flagellar motility of sperm packets of *Pleodorina starrii*

Azusa Kage[1]*, Kohei Takahashi[2], Hisayoshi Nozaki[2], Tetsuya Higashiyama[2], Shoji A. Baba[3], Takayuki Nishizaka[1]

[1]Department of Physics, Gakushuin University, 1-5-1 Mejiro, Toshima-ku, Tokyo 171-8588, Japan

[2]Department of Biological Sciences, Graduate School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

[3]Department of Biology, Ochanomizu University, 2-1-1 Otsuka, Bunkyo-ku, Tokyo 112-8610, Japan

*Corresponding author. azusa.kage@gakushuin.ac.jp

Keywords

Eukaryotic flagella, Volvocales, low Reynolds number, microswimmer, collective motion

Abstract

Eukaryotic flagella collectively form metachronal waves that facilitate the ability to cause flow or

swim. Among such flagellated and planktonic swimmers, large volvocine genera such as *Eudorina*,

Pleodorina and *Volvox* form bundles of small male gametes (sperm) called "sperm packets" for

sexual reproduction. Although these sperm packets reportedly have flagella and the ability to swim,

previous studies on volvocine motility have focused on asexual forms and the swimming

characteristics of sperm packets remain unknown. However, it is important to quantify the motility of

sperm packets and sperm in order to gain insights into the significance of motility in the sexual

reproduction of planktonic algae. In this study, we quantitatively described the behavior of three

flagellated forms of a male strain of *Pleodorina starrii*—asexual colonies, sperm packets, and single

dissociated sperm—with emphasis on comparison of the two multicellular forms. Despite being

smaller, sperm packets swam approximately 1.4 times faster than the asexual colonies of the same

male strain. Body length was approximately 0.5 times smaller in the sperm packets than in asexual

colonies. The flagella from sperm packets and asexual colonies showed asymmetric waveforms,

whereas those from dissociated single sperm showed symmetric waveforms, suggesting the

presence of a switching mechanism between sperm packets and dissociated sperm. Flagella from

sperm packets were approximately 0.5 times shorter and had a beat period approximately twice as

long as those from asexual colonies. The flagella of sperm packets were densely distributed over

the anterior part of the body, whereas the flagella of asexual colonies were sparse and evenly

distributed. The distribution of flagella, but not the number of flagella, appear to illustrate a

significant difference in the speeds of sperm packets and asexual colonies. Our findings reveal

novel aspects of the regulation of eukaryotic flagella and shed light on the role of flagellar motility in

sexual reproduction of planktonic algae.

Introduction

Collective motion is common in nature, such as that observed in schools of fish [1], flocks of birds [2] or herds of sheep [3]. Movements in fluids such as swimming of fish and flying of birds can be characterized by the Reynolds number *Re*, which is defined as $Re = \rho v L / \mu$, where ρ is the density of fluid, *v* is the speed, *L* is the characteristic length (e.g., body length), and μ is the viscosity of the fluid. The Reynolds number represents how inertia and viscosity dominate the movement; higher *Re* indicates the movement is more affected by inertia than by viscosity and vice versa. For low *Re* (<<1), where the spatial scale is small (< 1 mm), inertia does not contribute, and viscosity dominates the movement. Striking examples of low-*Re* movement are microbial swimming and flagellar/ciliary motion [4,5]. Metachronal waves of eukaryotic flagella or cilia [6,7] are a form of collective motion of many flagella/cilia, and are important for microbial swimming and flow generation inside the animal body. Understanding such collective motion on a small scale would lead to better understanding of the ecology of planktonic organisms that swim with flagella/cilia and manipulation of small-scale movements such as those in drug delivery systems.

Volvocine green algae, ranging from unicellular *Chlamydomonas* [8] and four-celled *Tetrabaena* [9] to thousands-celled *Volvox* [10,11], are planktonic swimmers with flagella that have been used as model organisms to investigate the dynamics of multiple flagella. They have two equal-length flagella in one cell, and flagella in the entire colony have been shown to move collectively in many cases. Previous studies on the motility of multicellular volvocines focused on asexual colonies that are in their usual form in culture. Although they have been known to sexually reproduce [12] and the sex-determining locus have been identified in several species (*Pleodorina starrii* [13], *Volvox carteri* [14], *Gonium pectorale* [15], *Yamagishiella unicocca* and *Eudorina* sp. [16]), the function of motility in sexual reproduction in multicellular volvocines remains unclear.

In the process of sexual reproduction, bundles of small male gametes or sperm called "sperm packets" have been reported in large multicellular volvocine genera such as *Eudorina* [17], *Pleodorina* [18,19], and *Volvox* [20,21]. Male gametes or sperm in these genera are characterized as biflagellated cells with elongated cell bodies, which differs from typical animal sperm with a single

flagellum [22,23]. Sperm are bundled as packets in male colonies, and these sperm packets swim

into female colonies [18,19]. Sperm packets subsequently dissociate into single sperm after arrival

at the female colonies and then fertilization occurs between dissociated single sperm and female

gametes held by female colonies. However, these processes have only been described

qualitatively, and no study has quantitatively characterized the movement of flagella in sperm

packets or sperm in volvocines.

In this study, we focused on the motility of the three flagellated forms of a male strain of the

volvocine alga *Pleodorina starrii* (Figure 1)—asexual colonies, sperm packets, and single

dissociated sperm—with emphasis on the comparison of the two multicellular forms (asexual

colonies and sperm packets). *P. starrii* has three sexes: male, female, and hermaphrodite [24]. We

found that the swimming speed of sperm packets was approximately 1.4 times faster than that of

asexual colonies, whereas the body size in sperm packets was approximately 0.5 times smaller

than that in asexual colonies. A comparison with a previous computational study [25] suggested that

the distribution of flagella over the body illustrates the difference in swimming speeds between

sperm packets and asexual colonies. Furthermore, we found that flagella from single dissociated

sperm showed symmetric motion, whereas flagella from sperm packets showed asymmetric motion,

suggesting a waveform switching mechanism in sperm cells. Our findings provide insights into the

function of motility in the sexual reproduction of volvocines, which had previously been overlooked.

Fig. 1. Light microscopic features of *Pleodorina starrii*, based on Takahashi et al. [24]. (A)

Schematic representation of the life cycle. Colonies proliferate asexually. Each colony consists of

32–64 cells with equal-length flagella. The cycle of sexual reproduction begins under a lack of

nitrogen, and consequently, male colonies with sperm packets—bundles of male gametes (sperm)

with two equal-length flagella—are formed in male and bisexual phonotypes. The sperm packet

swims to a female colony produced in female or bisexual phenotypes, and dissociates into sperm

that fertilize with female gametes within the female colony. The body length of asexual colonies and

sperm packets (*2a*) was measured as described in the Materials and Methods section. The flagellar

lengths of asexual colonies, sperm packets, and dissociated single sperm (*L*) were measured as the

length between the tip and base of the flagellum (see Materials and Methods). For details of the

three sex phenotypes (male, female, and bisexual), see Takahashi et al. [24]. (B) 64-celled mature

asexual colony. (C) Sexually induced male colony with sperm packets (sp). (D) Sperm packet. (E)

Dissociated single sperm. All photographs and illustrations are original. For materials and methods

of light micrographs and illustrations, see Takahashi et al. [24].

Materials and Methods

Culture

Pleodorina starrii was maintained at 20 °C in the artificial freshwater-6 (AF6) medium (pH

6.6) [26] under a 10:14 light/dark (LD) cycle. Asexual colonies were transferred into the Volvox

thiamine acetate (VTAC) medium (pH 7.5) [26] and incubated for 2–3 days at 25 °C under a 12:12

LD cycle. Sperm packets were induced by transferring 3-day colonies from the VTAC medium into a

mating medium that lacked a nitrogen source [18,24]. The unisex male strain P7 [24] (NIES-4480

[27]) was used unless otherwise stated. The dissociation process of the sperm packet was

observed using the unisex male strain P7 × P10 F1–14, which is the offspring of P7. The

experiments were performed at a temperature of approximately 23–25 °C during the subjective

daytime.

Recording of free swimming

Swimming activity was recorded using an upright microscope BX53 (Olympus, Tokyo,

Japan) with a halogen lamp (Olympus, Tokyo, Japan), dry dark-field condenser (U-DCD, Olympus,

Tokyo, Japan), and 620-nm long-pass filter to prevent phototaxis (SCF-50S-62R, OptSigma, Tokyo,

Japan). Although phototaxis was not examined in the present study, the wavelength was chosen

based on studies on *Chlamydomonas* phototaxis, in which it did not induce phototaxis [28]. *P. starrii*

was confined between a glass slide (S1126, Matsunami, Osaka, Japan) and coverslip (18 × 18 mm,

Matsunami, Osaka, Japan) with an ~10 × 10 mm square hole in a 0.5 mm silicone rubber spacer

(Kogugo, Tokyo, Japan). Movies were captured at 14.3 fps using a CMOS camera (UI-3250-CP,

IDS, Obersulm, Germany) to monitor swimming and at 1,000 fps using a high-speed CMOS camera

(VCC-H1540, Digimo, Tokyo, Japan) to monitor flagellar motility. A 10× objective lens (UPlanXApo,

NA 0.40, Olympus, Tokyo, Japan) was used to observe sperm packets, whereas a 4× objective lens

(UPlanFl, NA 0.13, Olympus, Tokyo, Japan) was used to observe asexual colonies. Swimming

trajectories and body lengths were quantified using the TrackMate plugin [29] in ImageJ (Fiji). The

swimming speed of each individual was calculated as the mean swimming speed along the

trajectory. The TrackMate plugin estimates the diameter of objects (bright spots); here, the

minimum value of the estimated diameter in the track was considered as the body length or twice

the radius a [25] because the estimated diameter was the minimum when the object was in focus.

Flagellar motility during swimming was visualized using the same setup (objective 10×) except that

a mercury lamp (Olympus, Tokyo, Japan) without a long-pass filter (i.e., white light) was used in

order to ensure sufficient light to visualize the movement of flagella, of which diameter is ~250 nm

[30] and beating frequency is tens of Hz [31], which require high shutter speed (i.e., short exposure

time).

Recording of flagellar motility of sperm packets

To visualize flagellar movement, sperm packets were captured using a glass micropipette

[32]. Glass pipettes with a tip diameter of ~20 μm were fabricated with a puller (PC-100, Narishige,

Tokyo, Japan) and microforge (MF2, Narishige, Tokyo, Japan) from the capillary (G-1, Narishige,

Tokyo, Japan). The glass pipette was set to the injector (IM-11-2, Narishige, Tokyo, Japan) on an

inverted microscope (IX71, Olympus, Tokyo, Japan); 0.5–2 μL droplets of the culture medium

containing sperm packets were placed on a glass-bottom dish (P50G-0-30-F, Mattek, Ashland, MA,

US), and the object inside the droplet was suctioned with the pipette. The droplets were covered

with mineral oil (8042-47-5; ICN Biomedicals, Aurora, OH, US) to prevent evaporation during

observation. Movies were captured at 1,000 fps using a high-speed CMOS camera (VCC-H1540,

Digimo, Tokyo, Japan), long-working distance phase-contrast optics (Olympus, Tokyo, Japan) and a

40× objective lens (UPlanFl, NA 0.75, Olympus, Tokyo, Japan) with a halogen lamp (TH4-100,

Olympus, Tokyo, Japan) and a 620-nm long-pass filter (SCF-50S-62R, OptSigma, Tokyo, Japan).

Analysis of flagellar motility

The flagellar waveform was analyzed using the lab-made software Bohboh [33]. Semi-auto-tracking of flagella was applied to the dark-field images of asexual colonies and dissociated single sperm. For pipette-captured sperm packets, phase-contrast images of the flagella were tracked manually because of poor contrast. From the time series of flagellar traces, we obtained space–time plots (kymographs) of curvature, along with flagellar length and time [34]. Flagellar beat periods were visually estimated from the space–time plots, and flagellar lengths were calculated from traces on Bohboh. As the length fluctuated along the beat period owing to the three-dimensionality of the waveform, the mean length for each flagellum (50–100 frames) was adopted as the flagellar length.

Immunofluorescence microscopy

To visualize the distribution of flagella, indirect immunofluorescence microscopy was performed according to Matsuura et al. [35], using the same method as that used for staining *Chlamydomonas* flagella. The samples were fixed in methanol. Antiacetylated α-tubulin (mouse monoclonal 6-11B-1, ab24610, Abcam, Cambridge, UK) diluted in blocking buffer (containing 10 mM phosphate buffer pH 7.2, 5% normal goat serum, 5% glycerol, 1% cold fish gelatin, 0.004% Na azide) to 1:500 was used as the primary antibody, and samples were incubated with the primary antibody at 37 °C for 1 hour. DyLight® 488-conjugated goat anti-mouse IgG (ab96871, Abcam, Cambridge, UK) diluted in blocking buffer to 1:500 was used as the secondary antibody, and samples were incubated with the secondary antibody at 37 °C for 1 hour. Nuclei were stained with 50 µg/mL Hoechst 33342 (Nacalai Tesque, Kyoto, Japan) in PBS for 10 minutes before immunofluorescence staining. Confocal images were captured using an FV-3000 microscope (Olympus, Tokyo, Japan) equipped with a 100× oil-immersion objective lens (UPlanApo, NA 1.50, Olympus, Tokyo, Japan) at the Department of Life Sciences, Gakushuin University. To grasp the three-dimensional structure of sperm packets and asexual colonies, Z-stacks were overlaid using the FV-3000 software (Olympus, Tokyo, Japan).

Results

Sperm packets swam faster than asexual colonies

To characterize the swimming of sperm packets, we recorded and compared the swimming of sperm packets with asexual colonies of the same male strain (P7) at low magnification. Figure 2A shows the relationship between swimming speed and body length of sperm packets (n = 185) and asexual colonies (n = 313). Hereafter, "body length of a sperm packet" indicates the size of the whole sperm packet, not a single cell (sperm). Sperm packets had smaller body lengths and higher swimming speeds than asexual colonies. The top and right panels of Figure 2A show the distribution of the body lengths and swimming speeds of sperm packets and asexual colonies. The body lengths of sperm packets and asexual colonies were 24.7 ± 5.9 μm (mean ± standard deviation [S.D.], n = 185) and 54.7 ± 14.9 μm (mean ± S.D., n = 313), respectively. The swimming speeds of sperm packets and asexual colonies were 308 ± 73 μm/s (mean ± S.D., n = 185) and 215 ± 63 μm/s (mean ± S.D., n = 313), respectively, indicating that sperm packets swam approximately 1.4 times faster than asexual colonies. Both the speed and the body length were significantly different between sperm packets and asexual colonies (speed: $p < 10^{-36}$ and body length: $p < 10^{-74}$ in Mann–Whitney U-test). The speed and body length showed no correlation in sperm packets (R = 0.07, p = 0.34 in Pearson's correlation test) and a weak but significant negative correlation in asexual colonies (R = −0.31, $p < 10^{-7}$). Figures 2B and 2C show the swimming trajectories of the sperm packets and asexual colonies, respectively. Both exhibited nearly straight trajectories within an observation field of less than 1 mm.

Fig. 2. Swimming of sperm packets and asexual colonies. (A) Swimming speed plotted against body

length (diameter). The top panel shows a histogram of body length; the right panel shows a

histogram of swimming speed (navy: sperm packets, green: asexual colonies). (B) Swimming

trajectories of sperm packets for 50 randomly chosen trajectories. The duration of tracking was 1.30

± 0.69 sec (mean ± S.D.). (C) Swimming trajectories of asexual colonies for 50 randomly chosen

trajectories. The duration of tracking was 4.21 ± 3.33 sec (mean ± S.D.).

Visualization of flagella in dark-field microscopy

To clarify the reason for the difference in speed between sperm packets and asexual

colonies, we visualized their flagellar movements. Figure 3 and Supplementary Movie S1 show a

swimming sperm packet viewed from the side using a dark-field microscope. The sperm packet

swam with flagella anterior (i.e., sperm packets are puller-type swimmers [36]) and metachronal

waves of the flagella traveling from the top region to the periphery of the packet, although it was

difficult to determine the relationship between the direction of the wave and the power stroke of the

flagella because of the dense distribution of flagella. The flagella of asexual colonies (Figure 4,

Supplementary Movie S2) were evenly distributed over the body, and the metachronal waves were

symplectic (i.e., waves propagated in the same direction as the power stroke of the flagella [37]), as

observed in *Volvox* [38].

Fig. 3. Image of a sperm packet (seen from the side) under dark-field microscopy. The sperm

packet moved using anterior flagella (i.e., puller-type swimming). The metachronal wave propagated

from the center (see Supplementary Movie 1). White triangles indicate flagella and the orange arrow

indicates the swimming direction.

Fig. 4. An asexual colony in dark-field microscopy. (A) Images of an asexual colony. Flagella beat

with asymmetric waveform. Registered images of a rolling colony near the wall are shown every 5

ms. The image of the silicone rubber spacer (wall) was deleted (black triangle in the lower right) for

proper registration (see Supplementary Methods). The blue arrow indicates the direction of the

effective stroke; the broken red line indicates where the kymograph (B) was drawn. Time [ms]

corresponds to the time axis in (B). (B) Kymograph of the movement of flagella drawn from the

broken red line in (A). The broken blue arrow indicates the propagation of metachronal waves. (C)

Flagella showing symplectic metachronal waves. Both the effective stroke of the flagella and the

metachronal wave propagated from the left to the right side of the image.

Sperm packets disassemble into individual sperm after arrival in female colonies [19]. Figure

5 and Supplementary Movie S3 show the process of disassembling a sperm packet, as recorded by

dark-field microscopy. The packet disassembled gradually as the sperm moved randomly. To date,

the trigger for this dissociation has not been identified. Figure 6 and Supplementary Movie S4 show

a single sperm after disassembly of a packet; the two flagella of the sperm beat alternately, and

individual sperm barely propel themselves. Flagellar movement was symmetric (i.e., flagella bent in

two directions [39]), similar to human or sea urchin sperm.

Fig. 5. Dissociation of a sperm packet viewed using dark-field microscopy. The sperm packet

gradually collapsed, and the dissociated sperm jumped out of the packet. The white triangle in 90

sec indicates dissociated sperm.

Fig. 6. Dark-field microscopy images of a single dissociated sperm. The two flagella beat with a

symmetric waveform and did not synchronize. One beat period was approximately 25 ms. Images

are shown every 2 ms.

Visualization of flagella of sperm packets held on the micropipette

257   As it was difficult to analyze flagellar motility from freely moving sperm packets (Figure 3),

we suctioned the sperm packets into a glass micropipette. Figure 7 and Supplementary Movie S5

show the flagellar motility of a sperm packet held on a micropipette, using phase-contrast

microscopy. We found that the flagella beat with an asymmetric waveform (i.e., flagella bent in only

one direction). The effective stroke was directed from the top (anterior) region of the body to the

periphery, and the metachronal waves were also directed from the top to the periphery (Figure 7B,

C). Thus, the metachronal waves in the sperm packets were symplectic, similar to those in asexual

colonies.

Fig. 7. Sperm packet captured by a micropipette in phase-contrast microscopy. (A) Image of the

sperm packet. The flagella beat with an asymmetric waveform. Images are shown every 5 ms. Blue

arrows indicate the direction of effective stroke, and the broken red line indicates where the

kymograph (B) was drawn. Time [ms] corresponds to the time axis in (B). Contrast enhanced. (B)

Kymograph of the movement of flagella drawn from the red broken line in (A). Blue broken arrows

indicate the propagation of metachronal waves. (C) Traces of flagella showing metachronal waves

(purple: waves to the right side, orange: waves to the left side). Only a fraction of flagella were

traced manually. Time [ms] corresponds to the time axis in (B).

Analysis of flagellar waveforms

We tracked and quantified flagellar movement. Figures 8A, B, and C show a time series of

flagellar waveforms traced from an asexual colony (Figure 4), dissociated single sperm (Figure 6),

and sperm packets (Figure 7), respectively. The flagellum from the asexual colonies (Figure 8A)

showed asymmetric waveforms, bending in one direction, similar to the colonies of other Volvocales

[11]. The flagellum from the dissociated single sperm (Figure 8B) showed symmetric waveforms

with principal and reverse bends [39]. The flagellum from sperm packets (Figure 8C) showed

asymmetric waveforms with a usually larger period. When we calculated the mean curvature over a

beat period at the midpoint of the flagella, the asexual colony (Figure 8A) showed a curvature of

0.11 rad/μm (at 10 μm from the base), the dissociated single sperm (Figure 8B) showed 0.03

rad/μm (at 7 μm from the base), and the sperm packet (Figure 8C) showed 0.08 rad/μm (at 5 μm

from the base). The flagellum appeared shorter in sperm packets than in single sperm or colonies.

As shown in Figure 8D–F, space–time plots of curvature along with time and flagellar length

supported these views; Figures 8D and 8F indicate that the flagella from the asexual colony and the

sperm packet bent in only one direction (i.e., asymmetric waveform), whereas Figure 8E shows that

the flagella from the single dissociated sperm bent in two directions (i.e., symmetric waveform).

Using waveform analysis (Figure 8, Supplementary Figure S1 and S2), we estimated the

beat period and length of the flagella (Supplementary Table 1). The beat period was 36 ± 13 ms in

asexual colonies (mean ± S.D., n = 4), 30 ± 10 ms in single dissociated sperm (n = 3), and 69 ± 18

ms in sperm packets (n = 3). The flagellar length was 22.1 ± 7.8 μm in asexual colonies (n = 4),

16.7 ± 1.2 μm in single dissociated sperm (n = 3), and 11.0 ± 0.5 μm in sperm packets (n = 3).

Fig. 8. Flagellar waveform of (A) an asexual colony, (B) single dissociated sperm, and (C) a sperm

packet. (A) and (B) show 20 frames at 1,000 fps, and (C) shows 60 frames every 3 frames. (D)–(F)

Space–time plots of curvature along with time and flagellar length obtained from (A)–(C).

Asymmetric waveforms in (D) and (F) show stripes with curvatures of positive (red) and near-zero

(green) values, indicating a one-directional bend, whereas the symmetric waveform in (E) shows

stripes of positive (red) and negative (blue) curvature, indicating a two-directional bend.

Visualization of the distribution of flagella

306   Indirect immunofluorescence microscopy was used to clearly visualize the distribution of

307   flagella. Figures 9A and 9B show the distribution of flagella (green) in sperm packets and asexual

308   colonies, respectively. The flagella of sperm packets were densely distributed over the anterior part

309   of the body, whereas the flagella of asexual colonies were sparse and evenly distributed.

Fig. 9. Distribution of flagella in immunofluorescence microscopy. (A) Sperm packet; (B) asexual colonies. Green structures represent acetylated α-tubulin (flagella) and blue structures represent nuclei. Z projections of confocal images are shown. Orange arrows indicate the swimming direction. Note that flagella were densely distributed in the sperm packet, whereas the asexual colony had sparse distribution of flagella.

Discussion

Mechanisms of straight swimming in asexual colonies and sperm packets

Both asexual colonies and sperm packets showed nearly straight swimming in a field of view of less than 1 mm (Figure 2B, C). Although both showed straight trajectories, the mechanisms are thought to be different, because the distribution (Figure 9) and metachronism (Figure 4, Figure 7) of flagella differ between asexual colonies and sperm packets. In asexual colonies, the distribution of flagella is nearly uniform, and each pair of flagella is nearly identical over a spherical body, although there might be anterior–posterior differences as observed in *Volvox* [11]. Therefore, the time-averaged force distribution generated by flagella is supposedly uniform over the asexual colony (i.e., neutral swimming), and this results in straight swimming. In other neutral swimmers [40] such as the ciliates *Paramecium* [41] and *Tetrahymena* [42], helical trajectories are frequently observed possibly because the metachronal waves are oblique against the body axis [41,43] or the cilia around the mouth (oral groove) move differently from the cilia in the other parts [41]; however, these were not the case in the asexual colonies of *P. starrii* in the present study. In contrast, sperm packets were pullers with tens of flagella in the anterior part (Supplementary Movie S1). The forces generated by flagella are stronger in the anterior part even if averaged over time. Symplectic metachronal waves are generated from the anterior to the posterior part of the packet (Figure 7, Supplementary Movie S5), resulting in straight swimming in the a–p axis direction.

The normalized body length is similar, but the normalized speed is dramatically different

between sperm packets and asexual colonies

336
337

Table 1 summarizes the swimming parameters of the sperm packets and asexual colonies.

We found that, on average, sperm packets had approximately 0.5 times smaller body length, 1.4

times higher speed, 0.5 times shorter flagellar length, and twice longer beat period than sperm

packets. Asexual colonies contained 32–64 cells [19] whereas the sperm packets contained 20–50

cells (n = 3). Thus, it is reasonable to assume that the number of flagella is comparable between the

asexual colonies and sperm packets. From the numerical simulation of ciliated spherical bodies,

Omori et al. [25] showed that the normalized swimming speed $\bar{U}/(\frac{L}{T})$, where $\bar{U}$ is the swimming

speed, L is flagellar length, and T is flagellar beat period, decreased by a power of -2 of the

normalized body length $2a/L$ (where a is the radius of the body) if the number of the flagella N was

fixed for a normalized body length ranging from 10 to 30. If we consider the mean flagellar length of

each category, sperm packets and asexual colonies had normalized body lengths of 2.2 and 2.5,

respectively (i.e., quite similar). However, the normalized swimming speeds were dramatically

different between the two categories. If the normalized body size and direction of the metachronal

waves (symplectic or antiplectic) were the same, the number of flagella would cause a difference in

swimming speed (i.e., a smaller number of flagella results in a slower speed) [25]. However, the

number of flagella would not differ in our case. Thus, in our study, the difference in speed could be

attributed to the difference in the distribution of flagella (Figure 9), which was not considered by

Omori et al. [25]. As the cell shape is elongated in sperm packets but spherical in asexual colonies

[18,19,24], the spacing of the flagella would be smaller in sperm packets. Cell elongation is not

directly related to the radius of the body a, because of the different configuration of cells between

asexual colonies and sperm packets (Figure 9). Elongated cells in sperm packets are

hemispherically aligned with all flagella anterior, whereas spherical cells in asexual colonies are

spherically aligned with all flagella out and with a hollow center.

Table 1. Swimming parameters of asexual colonies, sperm packets, and dissociated single sperm.

	Flagellar length, L [μm]	Flagellar period, T [ms]	L/T [μm/s]	Body length, $2a$ [μm]	Swimming speed, $\bar{U}$ [μm/s]	Normalized body length, $2a/L$	Normalized speed, $\bar{U}/(\frac{L}{T})$
Asexual colonies	22.1 ± 7.8 (n = 4)	36 ± 13 (n = 4)	622 ± 59 (n = 4)	54.7 ± 14.9 (n = 313)	215 ± 63 (n = 313)	2.5	0.3
Sperm packets	11.0 ± 0.5 (n = 3)	69 ± 18 (n = 3)	167 ± 44 (n = 3)	24.7 ± 5.9 (n = 185)	308 ± 73 (n = 185)	2.2	1.8
Single sperm	16.7 ± 1.2 (n = 3)	30 ± 10 (n = 3)	594 ± 198 (n = 3)	-	-	-	-

Flagellar length L and Flagellar period T were measured from the tracking data of flagella (Figure 8

and Supplementary Figure S1). Body length $2a$ was obtained as estimated diameter by TrackMate

(see Materials and Methods) and swimming speed $\bar{U}$ was calculated from each track (Figure 2). The

body length of a sperm packet indicates the size of the whole packet, not a single sperm.

Although the beat period of sperm packets was longer than that of asexual colonies (i.e., the

flagellar beat was slower in sperm packets), swimming was faster in sperm packets than in asexual

colonies. This could be attributed to the differences in size and distribution of flagella between

sperm packets and asexual colonies as well as the difference in the waveform and collective motion

of the flagella. The effect of size difference (i.e., sperm packets were smaller in size and thus had

smaller drag than asexual colonies) might be stronger than the effect of the beat period difference.

Another reason might be the differences in flagellar motion. Although flagellar movements of both

sperm packets and asexual colonies were categorized as asymmetric motion, they showed subtle

differences in actual waveforms (Figure 8A and C) and spatiotemporal dynamics in curvatures

(Figure 8D and F). These differences, as well as the differences in metachronal wave dynamics

(Figure 4B, Figure 7B) and hydrodynamic interactions between flagella, would result in faster

swimming in sperm packets.

Switching mechanism of symmetric and asymmetric waveforms

We found that the flagellar waveform changed in sperm packets and dissociated single

sperm. Sperm packets showed an asymmetric waveform, whereas dissociated single sperm

showed a symmetric waveform. Wild-type *Chlamydomonas* flagella usually show an asymmetric

waveform, and sometimes the waveform becomes symmetric, depending on the Ca^{2+} concentration

inside the flagella [44,45]. A similar mechanism can be considered for waveform switching in the

sperm of *P. starrii*: the Ca^{2+} concentration is constantly higher in the flagella of dissociated single

sperm than in the sperm packets. To test this hypothesis, an experiment using demembranated

models of both sperm packets and dissociated single sperm is desirable. In animal sperm, Ca^{2+}

plays an important role in changing flagellar waveform (and, thus, trajectory) in response to

chemoattractants [46]. It is possible that Ca^{2+} may also play a role in the reproduction of volvocine

green algae.

Flagellar lengths in dissociated single sperm and sperm packets

From the tracking data of flagella (Figure 8), we found differences in the lengths of flagella

for sperm packets and dissociated single sperm. The length was 16.7 ± 1.2 µm in single dissociated

sperm (n = 3) and 11.0 ± 0.5 µm in sperm packets (n = 3). As the sample size was too small to

perform statistical tests, we only discuss these data from the viewpoint of descriptive statistics. The

mean length was approximately 6 µm longer in single dissociated sperm than in sperm packets.

Although the difference might be partly attributed to the three-dimensionality of flagellar waveform

especially in sperm packets, the flagella of the fixed sperm packet also looked short (Figure 9A).

The flagella might be elongated before dissociation of sperm packets; that is, sperm might undergo

a process of maturation during which flagella might grow. In *Chlamydomonas*, the growth rate of

flagella after deflagellation is reported to be approximately 9 µm/hour [47]. Thus, it would be

possible for the *Pleodorina* sperm flagella to grow approximately 6 µm on a timescale of 1 hour

before dissociation. Currently, it is difficult to control the process of dissociation (i.e., the age of

sperm packets), but future study should test the maturation hypothesis of sperm before sperm

packet dissociation.

Significance of faster swimming speed in sperm packets

The sperm packets swam faster than the asexual colonies. To date, we have not measured

the swimming speed of female sexual colonies, which also have flagella. However, it would be

reasonable to assume that the speed of female sexual colonies would not be significantly higher

than that of asexual colonies because the morphology does not differ significantly between sexual

and asexual colonies of females [24]. A previous study reported that female colonies of *Eudorina*

elegans, a close relative to *Pleodorina starrii*, were immobile [48]. Taking the safe side, if we

consider that the speed of female sexual colonies is not significantly different from that of asexual

colonies of male strains, the faster swimming of sperm packets would enable them to catch up with

female colonies during sexual reproduction. However, it remains unclear how the sperm packets

approach female colonies. Three possibilities can be considered: accidental proximity,

hydrodynamic attraction as discussed in Drescher et al. [49], and chemical attraction as occurs in

some other taxa. Although chemoattractants that induce alterations in sperm flagellar waveforms

have been identified in some species of animals [46] and brown algae [50], such an attractant is

unknown in volvocines. Examining chemotaxis and determining chemoattratants, if any, in *P. starrii*

will be the focus of a future study.

Conclusions

In this study, we quantitatively characterized the swimming and flagellar motility of three

forms of the volvocine green alga *Pleodorina starrii*, with emphasis on comparison between asexual

colonies and sperm packets. Two important changes in the flagellar properties during male

gametogenesis and gamete fusion were confirmed. The first is the change in flagellar distribution

during sperm packet formation, which allows the packets to approach the female sexual colony at

high speed. This change in flagellar distribution is accompanied by a drastic change in swimming

mode, from neutral swimmers (asexual colonies) to pullers (sperm packets). The second is the

change from asymmetric to symmetric waveform for effective fusion with the female gametes.

These findings revealed diverse, developmentally- and physiologically-controlled movements under

low Reynolds number conditions even in a male strain of one species. Further clarification of such

diverse movements will lead to the control of motion and cargo transport under low Reynolds

number conditions. Further biophysical, biochemical and bioinformatics studies should clarify the

phototaxis and chemotaxis of sperm packets, the mechanism of switching flagellar waveforms, and

the entire process of sexual reproduction in the context of swimming.

Acknowledgements

This study was supported by JSPS KAKENHI #21K20661 (A.K.) and #22H05689 (A.K.).

We thank Dr. Masafumi Hirono (Hosei University) for the immunofluorescence staining protocol, Ms.

Tomoka Sakamoto (Narishige) for technical help with the suction experiments, Dr. Toshihiro Omori

(Tohoku University) for discussions on fluid mechanics, Dr. Masato S. Abe (Doshisha University) for

discussions on statistical methods, and Dr. Muneyoshi Ichikawa (Fudan University) for advice on

revision of figures.

Author contributions

A.K. conceived the study. K.T., H.N., and T.H. prepared the samples. T.N. constructed the

microscopy system. S.A.B. provided the software and developed the image analysis methods. A.K.

performed the experiments. A.K. and S.A.B. analyzed the data. A.K. drafted the manuscript, and

A.K., K.T., H.N., and T.N. contributed to the final manuscript.

References

1.   Parrish JK, Viscido SV, Grünbaum D. Self-organized fish schools: an examination of emergent

properties. Biol Bull. 2002;202: 296–305. doi: 10.2307/1543482

2.   Attanasi A, Cavagna A, Del Castello L, Giardina I, Jelic A, Melillo S, et al. Emergence of

collective changes in travel direction of starling flocks from individual birds' fluctuations. J R Soc

Interface. 2015;12: 20150319. doi: 10.1098/rsif.2015.0319

3.   Azaïs M, Blanco S, Bon R, Fournier R, Pillot M-H, Gautrais J. Traveling pulse emerges from

coupled intermittent walks: A case study in sheep. PLoS One. 2018;13: e0206817. doi:

10.1371/journal.pone.0206817

4.   Purcell EM. Life at low Reynolds number. Am J Phys. 1977;45: 3–11. doi: 10.1119/1.10903

5.   Wan KY, Poon RN. Mechanisms and functions of multiciliary coordination. Curr Opin Cell Biol.

2023;86: 102286. doi: 10.1016/j.ceb.2023.102286

6.   Aiello E, Sleigh MA. The metachronal wave of lateral cilia of Mytilus edulis. J Cell Biol. 1972;54:

493–506. doi: 10.1083/jcb.54.3.493

7.   Narematsu N, Quek R, Chiam K-H, Iwadate Y. Ciliary metachronal wave propagation on the

compliant surface of Paramecium cells. Cytoskeleton. 2015;72: 633–646. doi:

10.1002/cm.21266

8.   Wan KY, Goldstein RE. Coordinated beating of algal flagella is mediated by basal coupling.

Proc Natl Acad Sci U S A. 2016;113: E2784-93. doi: 10.1073/pnas.1518527113

9.   Tanno A, Tokutsu R, Arakaki Y, Ueki N, Minagawa J, Yoshimura K, et al. The four-celled

Volvocales green alga Tetrabaena socialis exhibits weak photobehavior and high-

photoprotection ability. PLoS One. 2021;16: e0259138. doi: 10.1371/journal.pone.0259138

10. Brumley DR, Wan KY, Polin M, Goldstein RE. Flagellar synchronization through direct

hydrodynamic interactions. Elife. 2014;3: e02750. doi: 10.7554/eLife.02750

11. Ueki N, Wakabayashi K-I. Detergent-extracted Volvox model exhibits an anterior-posterior

gradient in flagellar Ca2+ sensitivity. Proc Natl Acad Sci U S A. 2018;115: E1061–E1068. doi:

10.1073/pnas.1715489115

12. Nozaki H. Morphology and evolution of sexual reproduction in the Volvocaceae (Chlorophyta).

J Plant Res. 1996;109: 353–361. doi: 10.1007/bf02344484

13. Nozaki H, Mori T, Misumi O, Matsunaga S, Kuroiwa T. Males evolved from the dominant

isogametic mating type. Curr Biol. 2006;16: R1018-20. doi: 10.1016/j.cub.2006.11.019

14. Ferris P, Olson BJSC, De Hoff PL, Douglass S, Casero D, Prochnik S, et al. Evolution of an

expanded sex-determining locus in Volvox. Science. 2010;328: 351–354. doi:

10.1126/science.1186222

15. Hamaji T, Mogi Y, Ferris PJ, Mori T, Miyagishima S, Kabeya Y, et al. Sequence of the Gonium

pectorale mating locus reveals a complex and dynamic history of changes in volvocine algal

mating haplotypes. G3 (Bethesda). 2016;6: 1179–1189. doi: 10.1534/g3.115.026229

16. Hamaji T, Kawai-Toyooka H, Uchimura H, Suzuki M, Noguchi H, Minakuchi Y, et al. Anisogamy

evolved with a reduced sex-determining region in volvocine green algae. Commun Biol. 2018;1:

17. doi: 10.1038/s42003-018-0019-5

17. Goldstein M. Speciation and mating behavior inEudorina. J Protozool. 1964;11: 317–344. doi:

10.1111/j.1550-7408.1964.tb01762.x

18. Nozaki H, Kuroiwa H, Mita T, Kuroiwa T. Pleodorina japonica sp. nov. (Volvocales,

Chlorophyta) with bacteria-like endosymbionts. Phycologia. 1989;28: 252–267. doi:

10.2216/i0031-8884-28-2-252.1

19. Nozaki H, Ott FD, Coleman AW. MORPHOLOGY, MOLECULAR PHYLOGENY AND

TAXONOMY OF TWO NEW SPECIES OF PLEODORINA (VOLVOCEAE,

CHLOROPHYCEAE)1. J Phycol. 2006;42: 1072–1080. doi:10.1111/j.1529-8817.2006.00255.x

20. Darden WH Jr. Sexual differentiation in Volvox aureus. J Protozool. 1966;13: 239–255. doi:

10.1111/j.1550-7408.1966.tb01901.x

21. Kochert G. Differentiation of reproductive cells in Volvox carteri. J Protozool. 1968;15: 438–452.

doi: 10.1111/j.1550-7408.1968.tb02154.x

22. Fawcett DW. The mammalian spermatozoon. Dev Biol. 1975;44: 394–436. doi: 10.1016/0012-

1606(75)90411-x

23. Vacquier VD, Hirohashi N. Sea urchin spermatozoa. Methods Cell Biol. 2004;74: 523–544. doi:

10.1016/s0091-679x(04)74021-2

24. Takahashi K, Kawai-Toyooka H, Ootsuki R, Hamaji T, Tsuchikane Y, Sekimoto H, et al. Three

sex phenotypes in a haploid algal species give insights into the evolutionary transition to a self-

compatible mating system. Evolution. 2021;75: 2984–2993. doi: 10.1111/evo.14306

25. Omori T, Ito H, Ishikawa T. Swimming microorganisms acquire optimal efficiency with multiple

cilia. Proc Natl Acad Sci U S A. 2020;117: 30201–30207. doi: 10.1073/pnas.2011146117

26. Nozaki H, Mori F, Tanaka Y, Matsuzaki R, Yamaguchi H, Kawachi M. Cryopreservation of

vegetative cells and zygotes of the multicellular volvocine green alga Gonium pectorale. BMC

Microbiol. 2022;22: 103. doi: 10.1186/s12866-022-02519-9

27. Kawachi M, Ishimoto M, Mori F, Yumoto K, Sato M, M-H N. MCC-NIES. List of Strains, 9th

Edition [DVD]. Tsukuba: National Institute for Environmental Studies; 2013.

28. Nultsch W, Throm G, Rimscha I. Phototactic investigations in Chlamydomonas reinhardii

dangeard in homocontinuous culture. Arch Microbiol. 1971;80: 351–369. doi:

10.1007/bf00406222

29. Tinevez J-Y, Perry N, Schindelin J, Hoopes GM, Reynolds GD, Laplantine E, et al. TrackMate:

An open and extensible platform for single-particle tracking. Methods. 2017;115: 80–90. doi:

10.1016/j.ymeth.2016.09.016

30. Bui KH, Pigino G, Ishikawa T. Three-dimensional structural analysis of eukaryotic flagella/cilia

by electron cryo-tomography. J Synchrotron Radiat. 2011;18: 2–5. doi:

10.1107/S0909049510036812

31. Gilpin W, Bull MS, Prakash M. The multiscale physics of cilia and flagella. Nat Rev Phys.

2020;2: 74–88. doi: 10.1038/s42254-019-0129-0

32. de Maleprade H, Moisy F, Ishikawa T, Goldstein RE. Motility and phototaxis of Gonium, the

simplest differentiated colonial alga. Phys Rev E. 2020;101: 022416. doi:

10.1103/PhysRevE.101.022416

33. Shiba K, Mogami Y, Baba SA. Ciliary movement of sea-urchin Embryos. Nat Sci Rep

Ochanomizu Univ. 2002;53: 49–54.

34. Iida T, Iwata Y, Mohri T, Baba SA, Hirohashi N. A coordinated sequence of distinct flagellar

waveforms enables a sharp flagellar turn mediated by squid sperm pH-taxis. Sci Rep. 2017;7:

12938. doi: 10.1038/s41598-017-13406-z

35. Matsuura K, Lefebvre PA, Kamiya R, Hirono M. Bld10p, a novel protein essential for basal body

assembly in Chlamydomonas: localization to the cartwheel, the first ninefold symmetrical

structure appearing during assembly. J Cell Biol. 2004;165: 663–671. doi:

10.1083/jcb.200402022

36. Ishikawa T, Omori T, Kikuchi K. Bacterial biomechanics-From individual behaviors to biofilm

and the gut flora. APL Bioeng. 2020;4: 041504. doi: 10.1063/5.0026953

37. Knight-Jones EW. Relations between Metachronism and the Direction of Ciliary Beat in

Metazoa. J Cell Sci. 1954;s3-95: 503–521.

38. Brumley DR, Polin M, Pedley TJ, Goldstein RE. Metachronal waves in the flagellar beating of

Volvox and their hydrodynamic origin. J R Soc Interface. 2015;12: 20141358. doi:

10.1098/rsif.2014.1358

39. Gibbons BH, Gibbons IR. Properties of flagellar "rigor waves" formed by abrupt removal of
adenosine triphosphate from actively swimming sea urchin sperm. J Cell Biol. 1974;63: 970–
985. doi: 10.1083/jcb.63.3.970

40. Ohmura T, Nishigami Y, Ichikawa M. Simple dynamics underlying the survival behaviors of
ciliates. Biophys Physicobiol. 2022;19: e190026. doi: 10.2142/biophysico.bppb-v19.0026

41. Jennings HS. Behavior of the lower organisms, by H. S. Jennings. New York,: Columbia
University Press, The Macmillan company, 1906. doi: 10.5962/bhl.title.4123

42. Marumo A, Yamagishi M, Yajima J. Three-dimensional tracking of the ciliate Tetrahymena
reveals the mechanism of ciliary stroke-driven helical swimming. Commun Biol. 2021;4: 1209.
doi: 10.1038/s42003-021-02756-0

43. Ito H, Omori T, Ishikawa T. Swimming mediated by ciliary beating: comparison with a squirmer
model. J Fluid Mech. 2019;874: 774–796. doi: 10.1017/jfm.2019.490

44. Schmidt JA, Eckert R. Calcium couples flagellar reversal to photostimulation in
Chlamydomonas reinhardtii. Nature. 1976;262: 713–715. doi: 10.1038/262713a0

45. Hyams JS, Borisy GG. Isolated flagellar apparatus of Chlamydomonas: characterization of
forward swimming and alteration of waveform and reversal of motion by calcium ions in vitro. J
Cell Sci. 1978;33: 235–253. doi: 10.1242/jcs.33.1.235

46. Yoshida M, Yoshida K. Sperm chemotaxis and regulation of flagellar movement by Ca2+. Mol
Hum Reprod. 2011;17: 457–465. doi: 10.1093/molehr/gar041

47. Lefebvre PA, Nordstrom SA, Moulder JE, Rosenbaum JL. Flagellar elongation and shortening
in Chlamydomonas. IV. Effects of flagellar detachment, regeneration, and resorption on the
induction of flagellar protein synthesis. J Cell Biol. 1978;78: 8–27. doi: 10.1083/jcb.78.1.8

48. Nozaki H. Sexual reproduction in Eudorina elegans (chlorophyta, volvocales). Bot Mag Tokyo.
1983;96: 103–110. doi: 10.1007/bf02491095

49. Drescher K, Leptos KC, Tuval I, Ishikawa T, Pedley TJ, Goldstein RE. Dancing volvox:

hydrodynamic bound states of swimming algae. Phys Rev Lett. 2009;102: 168101. doi:

10.1103/PhysRevLett.102.168101

50. Kinoshita N, Nagasato C, Motomura T. Chemotactic movement in sperm of the oogamous

brown algae, Saccharina japonica and Fucus distichus. Protoplasma. 2017;254: 547–555. doi:

10.1007/s00709-016-0974-y

Supporting Information Files

S1 File. Supplementary methods, figures, and tables.

S1 Movie. Supplementary Movie S1. Sperm packets viewed from the side using dark-field

microscopy (objective ×10). The sperm packets swam along the anterior flagella. Playback speed,

1/50.

S2 Movie. Supplementary Movie S2. Movement of flagella in an asexual colony in dark-field

microscopy (objective ×10). A movie of a rolling colony near the wall was recorded, as described in

the Supplementary Methods, and the registered colony is shown. Playback speed, 1/50.

S3 Movie. Supplementary Movie S3. Dissociation of sperm packets observed using dark-field

microscopy (objective ×10). Playback speed, ×2.

S4 Movie. Supplementary Movie S4. Single dissociated sperm observed using dark-field

microscopy (objective ×10). Playback speed, 1/50.

S5 Movie. Supplementary Movie S5. Sperm packets were held with a micropipette and observed

using phase-contrast microscopy (objective ×40). Contrast enhanced. Playback speed, 1/50.