Updating the sulcal landscape of the human lateral parieto-occipital junction provides anatomical, functional, and cognitive insights

Ethan H. Willbrand[1,2*], Yi-Heng Tsai[3*], Thomas Gagnant[4], Kevin S. Weiner[1,2**]

[1]*Department of Psychology, University of California, Berkeley, Berkeley, CA, USA*
[2]*Helen Wills Neuroscience Institute, University of California, Berkeley, Berkeley, CA, USA*
[3]*Department of Psychology, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA*
[4]*Medical Science Faculty, University of Bordeaux, Bordeaux, France*
*Co-first authors
**Corresponding author: Kevin S. Weiner (kweiner@berkeley.edu)

Abstract

Understanding the tripartite relationship among neuroanatomy, brain function, and cognition is of major interest across neurobiological subdisciplines. Recent advances in neuroimaging have uncovered neuroanatomical structures in evolutionarily-expanded portions of the brain that are related to individual differences in brain function and cognition. Here, we explored this relationship in lateral parietal cortex (LPC), a region crucial for many higher-level cognitive abilities. To do so, we manually defined 2176 individual sulci across 144 hemispheres. We identified four small and shallow indentations of the cerebral cortex (sulci) that were previously unidentified in the lateral parieto-occipital junction (LPOJ) and LPC. One of these sulci (ventral supralateral occipital sulcus, slocs-v) is present in nearly every hemisphere, and is morphologically, architecturally, and functionally dissociable from neighboring sulci. Implementing a data-driven, model-based approach relating sulcal depth to behavior identified different relationships of ventral and dorsal LPC/LPOJ sulcal networks contributing to the perception of spatial orientation. The model identified the slocs-v, further indicating the importance of this new neuroanatomical structure. Our findings build on classic neuroanatomical theories and identify new neuroanatomical targets for future "precision imaging" studies exploring the relationship among brain structure, brain function, and cognitive abilities at the level of the individual participant.

Keywords

Cortical folding, Functional neuroanatomy, Magnetic resonance imaging (MRI), Occipital cortex, Parietal cortex, Spatial orientation

Introduction

A fundamental goal in neuroscience is to understand the complex relationship between brain

structure and brain function, as well as how that relationship provides a scaffold for efficient

cognition and behavior. Of all the neuroanatomical features to target, recent work shows that

morphological features of hominoid-specific, shallow indentations, or sulci, of the cerebral cortex

are not only functionally and cognitively meaningful, but also are particularly impacted by

multiple brain-related disorders and aging [1–20]. The combination of these findings provides

growing support for a classic theory proposing that the late emergence of these structures in

gestation within association cortices, as well as their prolonged development, may co-occur with

specific functional and microstructural features that could support specific cognitive abilities that

also have a protracted development [21]. Nevertheless, despite the developmental,

evolutionary, functional, cognitive, and theoretical relevance of these findings, they have mainly

been restricted to only a subset of association cortices such as the prefrontal, cingulate, and

ventral occipitotemporal cortices [1–20]. Thus, examining the relationship among these

structures (also known as tertiary sulci) relative to architectonic and functional features of the

cerebral cortex, as well as relative to cognition, remains uncharted in other association cortices

such as the lateral parietal cortex (LPC).

As LPC is a cortical extent that has expanded extensively throughout evolution [22,23],

there is great interest in the structure and function of LPC in development, aging, across

species, and in different patient populations. Yet, key gaps in knowledge relating individual

differences in the structure of LPC to individual differences in the functional organization of LPC

and cognitive performance remain for at least four main reasons. First, one line of recent work

shows that LPC displays a much more complex sulcal patterning than previously thought [24–

27], while a second line of work shows that LPC is tiled with many maps and discrete functional

regions spanning modalities and functions such as vision, memory, attention, action, haptics,

and multisensory integration in addition to theory of mind, cognitive control, and subdivisions of

the default mode network [28–34]. Second, a majority of the time, the two lines of work are

conducted independently from one another and the majority of human neuroimaging studies of

LPC implement group analyses on average brain templates—which causes LPC sulci to

disappear (**figure 1**). Third, despite the recently identified complexity of LPC sulcal patterning,

recent studies have also uncovered previously overlooked tertiary sulci in association cortices

(for example, in the posterior cingulate cortex [13,15]). Thus, fourth, it is unknown if additional

LPC tertiary sulci are waiting to be discovered and if so, could improve our understanding of the

structural-functional organization of LPC with potential cognitive insights as in other association

cortices. Critically, while such findings would have developmental, evolutionary, functional,

cognitive, and theoretical implications for addressing novel questions in future studies, they

would also have translational applications as sulci serve as biomarkers in neurodevelopmental

disorders [3–6] and "corridors" for neurosurgery [35,36].

In the present study, we first manually defined LPC sulci in 144 young adult

hemispheres using the most recent definitions of LPC sulci [24]. By manually labeling over

2,000 sulci, we uncovered four previously undefined (Supplemental Methods and Supplemental

figures 1-4 for historical details) sulci in the cortical expanse between the caudal branches of the

superior temporal sulcus (cSTS) and two parts of the intraparietal sulcus (IPS)—a cortical

expanse recently referenced as containing sensory "bridge" regions of the temporal-parietal-

occipital junction [37]—which we term the supralateral occipital sulci (ventral: slocs-v; dorsal:

slocs-d) and posterior angular sulci (ventral: pAngs-d; dorsal: pAngs-d). We then utilized

morphological (depth and surface area), architectural (gray matter thickness and myelination),

and functional (resting-state functional connectivity) data available in each participant to assess

whether the most common of these structures (slocs-v) was dissociable from surrounding sulci.

Finally, we assessed whether the updated view of the LPC/LPOJ sulcal landscape provided

cognitive insights using a model-based, data-driven approach [17] relating sulcal morphology to

behavior on tasks known to activate regions within this cortical expanse (for example, reasoning

and spatial orientation [38–41]).

Materials and Methods

(a) Participants

Data for the young adult human cohort analyzed in the present study were from the Human

Connectome Project (HCP) database (https://www.humanconnectome.org/study/hcp-young-

adult/overview). Here, we used 72 participants (50% female, 22-36 years old, and 90% right-

handed; there was no effect of handedness on our behavioral tasks; Supplemental materials)

that were also analyzed in several prior studies [11–16].

(b) Neuroimaging data acquisition

Anatomical T1-weighted (T1-w) MRI scans (0.8 mm voxel resolution) were obtained in native

space from the HCP database. Reconstructions of the cortical surfaces of each participant were

generated using FreeSurfer (v6.0.0), a software package used for processing and analyzing

human brain MRI images (surfer.nmr.mgh.harvard.edu) [42,43]. All subsequent sulcal labeling

and extraction of anatomical metrics were calculated from these reconstructions generated

through the HCP's version of the FreeSurfer pipeline [44].

(c) Behavioral data:

In addition to structural and functional neuroimaging data, the HCP also includes a wide range

of behavioral metrics from the NIH toolbox [45]. To relate LPC/LPOJ sulcal morphology to

behavior, we leveraged behavioral data related to spatial orientation (Variable Short Penn Line

Orientation Test), relational reasoning (Penn Progressive Matrices Test), and processing speed

(Pattern Completion Processing Speed Test; Supplemental Methods for task details). We

selected these tasks as previous functional neuroimaging studies have shown the crucial role of

LPC/LPOJ in relational reasoning and spatial orientation [38–41], while our previous work

relating sulcal morphology to cognition uses processing speed performance as a control

behavioral task [17,19].

(d) <u>Anatomical analyses:</u>

(i) *Manual labeling of LPC sulci*

Sulci were manually defined in 72 participants (144 hemispheres) guided by the most recent

atlas by Petrides [24], as well as recent empirical studies [25–27], which together offer a

comprehensive definition of cerebral sulcal patterns, including tertiary sulci. For a historical

analysis of sulci in this cortical expanse, please refer to Segal & Petrides [25] and Zlatkina &

Petrides [26]. Our cortical expanse of interest was bounded by the following sulci and gyri: (i)

the postcentral sulcus (PoCS) served as the anterior boundary, (ii) the superior temporal sulcus

(STS) served as the inferior boundary, (iii) the superior parietal lobule (SPL) served as the

superior boundary, and (iv) the medial and lateral transverse occipital sulci (mTOS and lTOS)

served as the posterior boundary. We also considered the following sulci within this cortical

expanse: the three different branches of the caudal superior temporal sulcus (posterior to

anterior: cSTS3, 2, 1), the supramarginal sulcus (SmgS), posterior intermediate parietal sulcus

(pips), sulcus of Brissaud (sB), anterior intermediate parietal sulcus of Jensen (aipsJ),

paroccipital intraparietal sulcus (IPS-PO), intraparietal sulcus (IPS), and the superior parietal

sulcus (SPS). Of note, the IPS-PO is the portion of the IPS extending ventrally into the occipital

lobe. The IPS-PO was first identified as the paroccipital sulcus by Wilder (1886). There is often

an annectant gyrus separating the horizontal portion of the IPS proper from the IPS-PO [26,46].

Additionally, we identified as many as four previously uncharted and variable tertiary

LPC/LPOJ sulci for the first time: the supralateral occipital sulcus (slocs; composed of ventral

(slocs-v) and dorsal (slocs-d) components) and the posterior angular sulcus (pAngs; composed

of ventral (pAngs-v) and dorsal (pAngs-d) components). In the Supplemental Methods and

Supplemental Figures 1-4, we discuss the slocs and pAngs within the context of modern and

historical sources.

For each participant in each hemisphere, the location of each sulcus was confirmed by

trained independent raters (E.H.W., Y.T., and T.G.) and finalized by a neuroanatomist (K.S.W.).

All LPC sulci were then manually defined in FreeSurfer using tksurfer tools, as in previous work

[11–20], from which morphological and anatomical features were extracted. For four example

hemispheres with these 13-17 sulci identified, see **figure 1*a*** (Supplemental figure 5 for all

hemispheres). The specific criteria to identify the slocs and pAngs are outlined in **figure 1*b***. We

also generated sulcal probability maps for each sulcus (Supplemental Methods and

Supplemental figure 6).

To test whether the incidence rates of the slocs and pAngs components were statistically

different, we implemented a binomial logistic regression GLM with sulcus (slocs-v, slocs-d,

pAngs-v, and pAngs-d) and hemisphere (left and right), as well as their interaction, as predictors

for sulcal presence [0 (absent), 1 (present)]. GLMs were carried out with the glm function from

the built-in stats R package. ANOVA χ2 tests were applied to each GLM with the Anova function

from the car R package, from which results were reported.

Figure 1. Four previously undefined small and shallow sulci in the lateral parieto-occipital junction (LPOJ). *(a)* Four example inflated (top) and pial (bottom) left hemisphere cortical surfaces displaying the 13-17 sulci manually identified in the present study. Each hemisphere contains 1-4 of the previously undefined and variable LOC/LPOJ sulci (slocs and pAngs). Each sulcus is numbered according to the legend. *(b)* Criteria for defining slocs and pAngs components. (i) Slocs-v is the cortical indentation between the cSTS3 and ITOS. (ii) Slocs-d is the indentation between cSTS3/cSTS2 and IPS-PO. (iii) pAngs-v is the indentation between the cSTS2 and pips. (iv) pAngs-d is the indentation between cSTS2/cSTS1 and IPS. *(c)* The variability of the slocs and pAng components can cause them to disappear when individual surfaces are averaged together. Left to right: (i) 10 HCP participants, (ii) 20 HCP participants, (iii) 100 HCP participants, and iv) 650 HCP participants. The disappearance of these sulci on average surfaces, which are often used for group analyses in neuroimaging research, emphasizes the importance of defining these structures in individual hemispheres.

(ii) *Extracting and comparing the morphological and architectural features from sulcal*

labels

Morphologically, we compared sulcal depth and surface area across sulci, as these are two of

the primary morphological features used to define and characterize sulci [8,10–13,16–18,21,47–

51]. As in our prior work [17,18], mean sulcal depth values (in standard FreeSurfer units) were

computed in native space from the .sulc file generated in FreeSurfer [42] with custom Python

code [17]. Briefly, depth values are calculated based on how far removed a vertex is from what

is referred to as a "mid-surface," which is determined computationally so that the mean of the

displacements around this "mid-surface" is zero. Thus, generally, gyri have negative values,

while sulci have positive values. Each depth value was also normalized by the deepest point in

the given hemisphere. Surface area (mm^2) was calculated with the FreeSurfer

mris_anatomical_stats   function

(https://surfer.nmr.mgh.harvard.edu/fswiki/mris_anatomical_stats). The morphological features

of all LPC/LPOJ sulci are documented in Supplemental figure 7.

Architecturally, we compared cortical thickness and myelination, as in our prior work in

other cortical expanses [12,13,16,17]. Mean gray matter cortical thickness (mm) was extracted

using the FreeSurfer mris_anatomical_stats function. To quantify myelin content, we used the

T1-w/T2-w maps for each hemisphere, an in vivo myelination proxy [52]. To generate the T1-

w/T2-w maps, two T1-w and T2-w structural MR scans from each participant were registered

together and averaged as part of the HCP processing pipeline [44]. The averaging helps to

reduce motion-related effects or blurring. Additionally, and as described by Glasser and

colleagues [44], the T1-w/T2-w images were bias-corrected for distortion effects using field

maps. We then extracted the average T1-w/T2-w ratio values across each vertex for each

sulcus using custom Python code [12]. The architectural features of all LPC/LPOJ sulci are

documented in Supplemental figure 7.

To assess whether these four metrics differed between the slocs-v and surrounding sulci

(cSTS3 and ITOS), we ran a repeated measure analysis of variance (rm-ANOVA) with the

within-participant effects of sulcus (slocs-v, cSTS3, and ITOS), metric (surface area, depth,

cortical thickness, and myelination), and hemisphere (left and right). Rm-ANOVAs (including

sphericity correction) were implemented with the aov_ez function from the afex R package.

Effect sizes for the ANOVAs are reported with the partial eta-squared metric ($\eta2$). Post-hoc

analyses were computed with the emmeans function from the emmeans R package (*p*-values

corrected with Tukey's method). We also repeated these analyses for the three cSTS

components [24,25] and the two intermediate parietal sulcal components (ips: aipsJ and pips

[24,26]; detailed in the Supplemental Results and Supplemental Figure 8) as these components,

to our knowledge, have not been quantitatively compared in previous work.

(f) <u>Functional analyses</u>

To determine if the slocs-v is functionally distinct from surrounding sulci, we generated

functional connectivity profiles using recently developed analyses [12,13,53]. First, we used

resting-state network parcellations for each individual participant from Kong and colleagues [54],

who generated individual network definitions by applying a hierarchical Bayesian network

algorithm to produce maps for each of the 17 networks in individual HCP participants.

Importantly, this parcellation was conducted blind to both cortical folding and our sulcal

definitions. Next, we resampled the network profiles for each participant onto the fsaverage

cortical   surface,   and   then   to   each   native   surface   using   CBIG   tools

(https://github.com/ThomasYeoLab/CBIG). We then calculated the spatial overlap between a

sulcus and each of the 17 individual resting-state networks via the Dice coefficient (Equation 1):

$$(1) \quad Dice\,(\square, \square) \;=\; \frac{2\,|\square \cap \square|}{|\square| + |\square|}$$

This process of calculating the overlap between each sulcus and the 17-network

parcellation generated a "connectivity fingerprint" for each sulcus in each hemisphere of each

participant. We then ran an rm-ANOVA with within-participant factors of sulcus (slocs-v, cSTS3,

and ITOS), network (17 networks), and hemisphere (left and right) to determine if the network

profiles (i.e., the Dice coefficient overlap with each network) of the slocs-v was differentiable

from the surrounding sulci (i.e., cSTS3 and ITOS). Here we discuss effects related to networks

that at least showed minor overlap with one sulcus (i.e., Dice ≥ .10). As in the prior analysis, we

also repeated these analyses for the three cSTS components and the two intermediate parietal

sulcal components (Supplemental Results and Supplemental Figure 8).

(g) Behavioral analyses

(i) *Model selection*

The analysis relating sulcal morphology to spatial orientation and/or reasoning consisted of

using a cross-validated least absolute shrinkage and selection operator (LASSO) regression to

select the sulci that explained the most variance in the data and determined how much variance

is explained by sulcal depth as a predictor of behavior, as implemented in our previous work

[16–18]. The depths of all LPC/LPOJ sulci were included as predictors in the LASSO regression

model (Supplemental Methods for details on demographic control variables). As the shrinkage

parameter (alpha) increases, it decreases the coefficient of each of the sulci to zero except for

those with the strongest association. Therefore, this technique highlights the sulci whose

morphology was most closely related to behavior. We used cross-validation to optimize the

shrinking parameter for the LASSO regression. Conventionally, we selected the model

parameters that minimized the cross-validated mean squared error (MSE_{cv}) [55]. The

optimization was performed with the GridSearchCV function sklearn in Python, which allowed us

to determine the model parameters minimizing the MSE_{cv}. To evaluate the performance of any

model selected by the LASSO regression, as in prior work [16–18], we measured the model

performance for the relevant behavioral task using nested model comparison. With leave-one-

out cross-validation (LooCV), we compared the LASSO-selected model with the predictors to a

model with all left hemisphere sulci as predictors.

(ii) *Assessing morphological and behavioral specificity*

To assess whether our findings generalized to other anatomical features, we considered cortical

thickness, which is consistently studied in cognitive neuroscience studies relating morphology to

cognition [16–18,56,57]. To do so, we replaced sulcal depth with cortical thickness as the

predictive metric in our LASSO-selected model. As with depth, the model was fit to the data with

LooCV. To compare the thickness model to the depth model, we used the Akaike Information

Criterion (AIC), which provides an estimate of in-sample prediction error and is suitable for non-

nested model comparison. By comparing AIC scores, we are able to assess the relative

performance of the two models. If the ΔAIC is > 2, it suggests an interpretable difference

between models. If the ΔAIC is > 10, it suggests a strong difference between models, with the

lower AIC value indicating the preferred model [58]. To also ascertain whether the relationship

between LPC/LPOJ sulcal depth and cognition is specific to spatial orientation performance, or

transferable to other general measures of cognitive processing, we investigated the

generalizability of the sulcal-behavior relationship to another widely used measure of cognitive

functioning: processing speed [59]. Specifically, we used LooCV to predict processing speed

instead of spatial orientation score. As with thickness, we compared the two models with the

AIC.

Results

(a) Four previously undefined small and shallow sulci in the lateral parieto-occipital

junction (LPOJ)

In addition to defining the 13 sulci previously described within the LPC/LPOJ, as well as the

posterior superior temporal cortex (**Materials and Methods**) [24] in individual participants, we

could also identify as many as four small and shallow sulci situated within the LPC/LPOJ that

were more variable across individuals and uncharted until now (Supplemental Methods and

Supplemental figures 1-4). Macroanatomically, we could identify two sulci between the cSTS3

and the IPS-PO/ITOS ventrally and two sulci between the cSTS2 and the pips/IPS dorsally.

Ventrally, we refer to these sulci as ventral (slocs-v; sulcus 5 in **figure 1**) and dorsal

(slocs-d; sulcus 6 in **figure 1**) components of the supralateral occipital sulcus. The slocs-v,

located between the posterior cSTS3 and ITOS, was present in 98.6% of hemispheres (left

hemisphere: N = 71/72; right hemisphere: N = 71/72; **figure 1**). Conversely, the slocs-d, located

between the cSTS3 and IPS-PO, was present 68.0% of the time (left hemisphere: N = 50/72;

right hemisphere: N = 48/72; **figure 1**). Dorsally, we refer to the other newly identified sulci as

the ventral (pAngs-v; sulcus 7 in **figure 1**) and dorsal (pAngs-d; sulcus 8 in **figure 1**)

components of the posterior angular sulcus. The pAng components were more rare than the

slocs components. Specifically, pAngs-v, located between cSTS2 and pips, was identifiable

31.3% of the time (19 left and 26 right hemispheres; **figure 1**). Located between cSTS2 and the

IPS, pAngs-d was identifiable only 13.2% of the time (8 left and 11 right hemispheres; **figure 1**).

Though we characterize these sulci in this paper for the first time, the location of these four sulci

is consistent with the presence of variable "accessory sulci" in this cortical expanse mentioned

in prior modern and classic studies (Supplemental Methods).

These incidence rates were significantly different (GLM, main effect of sulcus:

$\chi 2(3)\square=\square166.53$, $p < .0001$; no hemispheric effects: $ps > .68$). The slocs-v was more common

than the other three sulci ($ps < .0001$), slocs-d was more common than the pAngs components

($ps < .0001$), and pAngs-v was more common than pAngs-d ($p = .002$). We could further identify

these sulci in post-mortem hemispheres (Supplemental figures 2, 3). Finally, to help guide future

research on these newly- and previously-classified LPC/LPOJ sulci, we generated probabilistic

maps of each of these 17 sulci and share them with the field with the publication of this paper

(Supplemental figure 6; **Data accessibility statement**).

(b) The slocs-v is morphologically, architecturally, and functionally dissociable from

nearby sulci

Given that the slocs-v was present in the majority of participants (98.6% across hemispheres),

we focused our analyses on this stable sulcal feature of the LPOJ. To do so, we first tested

whether the slocs-v was morphologically (depth and surface area) and architecturally (gray

matter thickness and myelination) distinct from the two sulci surrounding it: the cSTS3 and ITOS

(**figure 1**; Supplemental figure 7 for these metrics in all 17 sulci examined). An rm-ANOVA

(within-participant factors: sulcus, metric, and hemisphere for standardized metric units)

revealed a sulcus x metric interaction ($F(4, 276.19) = 179.15$, $\eta2 = 0.38$, $p < .001$). Post hoc

tests showed four main differences: (i) the slocs-v was shallower than cSTS3 ($p < .001$) but not

ITOS ($p = .60$), (ii) the slocs-v was smaller than both the cSTS3 and ITOS ($ps < .001$), (iii) the

slocs-v was thicker than both the cSTS3 and ITOS ($ps < .001$), and iv) the slocs-v was less

myelinated than both the cSTS and ITOS ($ps < .001$; **figure 2a**). There was also a sulcus x

metric x hemisphere interaction ($F(4.20, 289.81) = 4.16$, $\eta 2 = 0.01$, $p = .002$; Supplemental

Results).

We then tested whether the slocs-v was also functionally distinct from the cSTS3 and

ITOS by leveraging resting-state network parcellations for each individual participant to quantify

"connectivity fingerprints" for each sulcus in each hemisphere of each participant (**Materials**

and Methods) [54]. An rm-ANOVA (within-participant factors: sulcus, network, and hemisphere

for Dice coefficient overlap) revealed a sulcus x network interaction ($F(32, 2144) = 80.18$, $\eta 2 =$

0.55, $p < .001$). Post hoc tests showed that this interaction was driven by four effects: (i) the

cSTS3 overlapped more with the Default A subnetwork than both the slocs-v and ITOS ($ps <$

$.001$), (ii) the slocs-v overlapped more with the Default C subnetwork than the ITOS ($p < .001$)

and marginally than the cSTS3 ($p = .077$), (iii) the slocs-v overlapped more with the Dorsal

Attention A subnetwork than both the cSTS3 and ITOS ($ps < .001$), and iv) the ITOS overlapped

more with the Visual A and Visual B subnetworks than both the cSTS3 and slocs-v ($ps < .004$;

figure 2b). There was also a sulcus x network x hemisphere interaction ($F(32, 2144) = 3.99$, $\eta 2$

$= 0.06$, $p < .001$; Supplemental Results). Together, these results indicate that the slocs-v is a

morphologically, architecturally, and functionally distinct structure from its sulcal neighbors, and

thus, deserves a distinct neuroanatomical definition.

We further found that the three caudal STS rami [24,25] and intermediate parietal sulci

(aipsJ and pips) [24,26] are morphologically, architecturally, and functionally distinct structures

for the first time (to our knowledge), which empirically supports their distinctions with separate

sulcal labels (Supplemental Results and Supplemental figure 8).

Figure 2. The slocs-v is morphologically, architecturally, and functionally dissociable from nearby sulci. *(a)* Radial plot displaying the morphological (upper metrics: depth, surface area) and architectural (lower metrics: cortical thickness, myelination) features of the slocs-v (gray), cSTS3 (blue), and ITOS (green). Each dot and solid line represent the mean. The dashed lines indicate ± standard error. These features are colored by sulcus (legend). Metrics are in standardized units. *(b)* Radial plot displaying the connectivity fingerprints of these three sulci: the Dice Coefficient overlap (values from 0-1) between each component and individual-level functional connectivity parcellations [54].

(c) The morphology of LPC/LPOJ sulci, including the slocs-v, is related to cognitive performance

Finally, leveraging a data-driven approach of cross-validated LASSO feature selection, we sought to determine whether sulcal depth, a main defining feature of sulci, related to cognitive performance (**Materials and Methods**). To do so, we primarily focused on spatial orientation and reasoning given that these abilities recruit multiple subregions of lateral parietal and/or occipital cortices [38–41]. As in prior work [16–18], we chose the model at the alpha that minimized MSE_{cv}. Participants with a slocs-v in both hemispheres and all behavioral metrics were included (N = 69). Due to their rarity (being in less than 70% of hemispheres at most), we did not include the slocs-d or pAng components in this analysis.

This method revealed an association between spatial orientation scores and normalized sulcal depth in the left hemisphere (MSE_{cv} = 25.63, alpha = 0.05; **figure 3*a***), but not in the right hemisphere (MSE_{cv} = 26.41, alpha = 0.3). Further, we found that no LPC/LPOJ sulci were selected for reasoning in either hemisphere (right*:* alpha = 0.3, MSE = 24.01; left: alpha = 0.3, MSE = 24.01). Six left hemisphere LPC/LPOJ sulci were related to spatial orientation task

performance **figure 3a, *b***). Four of these sulci were positioned ventrally: cSTS3 (β = -9.77),

slocs-v (β = -3.36), lTOS (β = -4.91), and mTOS (β = -0.06), whereas two were positioned

dorsally: pips (β = 5.02), and SPS (β = 4.30; **figure 3a, *b***). Using LooCV to construct models

that predict behavior, the LASSO-selected model explained variation in spatial orientation score

(R^2_{cv} = 0.06, MSE_{cv} = 23.99) above and beyond a model with all left hemisphere sulci (R^2_{cv} <

0.01, MSE_{cv} = 27.12). This model also showed a moderate correspondence (r_s = 0.29, p = .01;

figure 3*c*) between predicted and actual measured scores. We then tested for anatomical and

behavioral specificity using the AIC, which revealed two primary findings. First, we found that

the LASSO-selected sulcal depth model outperformed a model using the cortical thickness of

the six LASSO-selected sulci (R^2_{cv} < .01, MSE_{cv} = 26.02, $AIC_{cortical\ thickness}$ − $AIC_{sulcal\ depth}$ = 2.19).

This model also showed task specificity as these sulci outperformed a model with processing

speed (R^2_{cv} < .01, MSE_{cv} = 254.65, $AIC_{processing\ speed}$ − $AIC_{spatial\ orientation}$ = 63.57). Thus, our data-

driven model explains a significant amount of variance on a spatial orientation task and shows

behavioral and morphological specificity.

Figure 3. The morphology of LPC/LPOJ sulci, including the slocs-v, is related to cognitive performance. *(a)* Beta-coefficients for each left hemisphere LPC/LPOJ sulcus at a range of shrinking parameter values [alpha (α)]. Highlighted gray bar indicates coefficients at the chosen α-level. Bottom: Cross-validated mean-squared error

393 (MSE$_{CV}$) at each α level. By convention, we selected the α that minimized the MSE$_{CV}$ (dotted line). *(b)*
394 Inflated left hemisphere cortical surface from an example participant highlighting the two groups of sulci—
395 *dorsal positive* (orange) and *ventral negative* (green)—related to spatial orientation performance. *(c)*
396 Spearman's correlation (r$_s$) between the measured and the predicted spatial orientation scores from the
397 LASSO-selected model is shown in *(a)*.
398

Discussion

(a) <u>Overview</u>

In the present study, we examined the relationship between LPC/LPOJ sulcal morphology,

functional connectivity fingerprints, and cognition. We report five main findings. First, while

manually defining sulci in LPC/LPOJ across 144 hemispheres, we uncovered four new small

and shallow sulci that are not included in present or classic neuroanatomy atlases or

neuroimaging software packages. Second, we found that the most common of these structures

(the slocs-v; identifiable 98.6% of the time) was morphologically, architecturally, and functionally

differentiable from nearby sulci. Third, using a model-based, data-driven approach quantifying

the relationship between sulcal morphology and cognition, we found a relationship between the

depths of six LPC/LPOJ sulci and performance on a spatial orientation processing task. Fourth,

the model identified distinct dorsal and ventral sulcal networks in LPC/LPOJ: ventral sulci had

negative weights while dorsal sulci had positive weights (**figure 3*b***). These findings are

consistent with previous neuroimaging work from Gur et al. [41] who demonstrated separate

functional activations in dorsal parietal and the more ventrally situated occipital-parietal cortices

for the judgment of line orientation task used in the present study. Fifth, the model identified that

the slocs-v is cognitively relevant, further indicating the importance of this new neuroanatomical

structure. In the sections below, we discuss (i) the slocs-v relative to modern functional and

cytoarchitectonic parcellations in the LPC/LPOJ, as well as anatomical connectivity to other

parts of the brain, (ii) underlying anatomical mechanisms relating sulcal morphology and

behavior more broadly, and (iii) limitations of the present study. Implications for future studies

are distributed throughout each section.

(b) <u>The slocs-v relative to modern functional and cytoarchitectonic parcellations in the</u>

<u>LPC/LPOJ, as well as anatomical connectivity to other parts of the brain</u>

To lay the foundation for future studies relating the newly-identified slocs-v to different

anatomical and functional organizational features of LPC/LPOJ, we situate probabilistic

predictions of slocs-v relative to probabilistic cortical areas identified using multiple modalities.

For example, when examining the correspondence between the slocs-v and modern multimodal

(HCP-MMP [37]) and observer-independent cytoarchitectural (Julich-Brain atlas [60]) areas

(Supplemental Methods), the slocs-v is located within distinct areas. In particular, the slocs-v

aligns with the multimodally- and cytoarchitecturally-defined area PGp bilaterally and

cytoarchitecturally-defined hIP4 in the right hemisphere (**figure 4**). In classic neuroanatomical

terms [61], this indicates that the slocs-v is a putative "axial sulcus" for these regions, which

future work can assess with analyses in individual participants.

Figure 4. The slocs-v relative to modern functional and cytoarchitectonic parcellations in LPC/LPOJ. *(a)* Top: Left (LH) and right (RH) hemispheres of the inflated fsaverage surface with two areas from the modern HCP multimodal parcellation (HCP-MMP; blue) [37] relative to an MPM of the slocs-v (warm colors indicate areas with at least 20% overlap across participants; Supplemental figure 6). Bottom: Same as top, except for two observer-independent cytoarchitectonic regions from the Julich-Brain Atlas [60]. *(b)* Overlap between the slocs-v and each area (Supplemental Methods). Each dot and solid line represents the mean. The dashed lines indicate ± standard error (left: gray; right: white).

Aside from recent multimodal and observer-independent cytoarchitectonic parcellations, an immediate question is: What is the relationship between the slocs-v and other functional regions at this junction between the occipital and parietal lobes, as well as potential anatomical connectivity? For example, there are over a dozen visual field maps in the cortical expanse spanning the TOS, IPS-PO, and the IPS proper (see (i), (ii), and (iii), respectively in **figure 5a**) [33]. When projecting probabilistic locations of retinotopic maps from over 50 individuals from Wang and colleagues [62] (Supplemental Methods), the slocs-v is likely located outside of visual field maps extending into this cortical expanse (**figure 5a**). Nevertheless, when also projecting the map of the mean R^2 metric from the HCP retinotopy dataset from 181 participants shared by Benson and colleagues [63] (Supplemental Methods), the slocs-v is in a cortical expanse that explains a significant amount of variance (left hemisphere: $R^2_{mean} = 19.29$, $R^2_{max} = 41.73$; right hemisphere: $R^2_{mean} = 21.17$, $R^2_{max} = 44.23$; **figure 5b**).

Figure 5. The slocs-v relative to retinotopy. *(a)* Top: Left (LH) and right (RH) hemispheres of the
inflated fsaverage surface displaying the probabilistic locations of retinotopic maps from over 50
individuals from Wang and colleagues [62] (black outlines). The predicted slocs-v location from the MPMs
is overlaid in orange (as in figure 4). (i), (ii), and (iii) point out the retinotopic maps in the cortical expanse
spanning the TOS, IPS-PO, and IPS, respectively. *(b)* Same format as in *(a)*, but with a map of the mean
R^2 metric from the HCP retinotopy dataset [63] overlayed on the fsaverage surfaces (thresholded
between R^2 values of 10% and 90%). This metric measures how well the fMRI time series at each vertex
is explained by a population receptive field (pRF) model. The mean and max R^2 values for the slocs-v
MPM in each hemisphere are included below each surface.

In terms of anatomical connectivity, as the slocs-v co-localizes with cytoarchitectonically

defined PGp (**figure 4**) and previous studies have examined the anatomical connectivity of the

probabilistically defined PGp, we can glean insight regarding the anatomical connectivity of

slocs-v from these previous studies [64,65]. This prior work showed that PGp was anatomically

connected to temporooccipital regions, other regions in the temporal lobe, middle and superior

frontal cortex, as well as the inferior frontal cortex and insula [64,65]. Of course, the location of

the slocs-v relative to multimodal, cytoarchitectonic, and retinotopic areas, as well as the

anatomical connectivity of the slocs-v, would need to be examined in individual participants, but

the present work makes clear predictions for future studies as fleshed out here. To conclude this

section, as the multimodal area PGp (**figure 4**) was recently proposed as a "transitional area"

by Glasser and colleagues [37] (Supplemental table 1), future studies can also further

functionally and anatomically test the transitional properties of slocs-v.

(c) Underlying anatomical mechanisms relating sulcal morphology and behavior

In this section, we discuss potential anatomical mechanisms contributing to the relationship

between sulcal depth and behavior in two main ways. First, long-range white matter fibers have

a gyral bias, while short-range white matter fibers have a sulcal bias in which some fibers

project directly from the deepest points of a sulcus [66–70]. As such, recent work hypothesized

a close link between sulcal depth and short-range white matter properties [16–18,71,72]: deeper

sulci would reflect even shorter short-range white matter fibers, which would result in faster

communication between local, cortical regions and in turn, contribute to improved cognitive

performance. This increased neural efficiency could underlie individual differences in cognitive

performance. Ongoing work is testing this hypothesis which can be further explored in future

studies incorporating anatomical, functional, and behavioral measures, as well as computational

modeling.

Second, our model-based approach identified separate dorsal and ventral sulcal

networks in which deeper sulci dorsally and shallower sulci ventrally contributed to the most

explained variance on the spatial orientation task. A similar finding was identified by our

previous work in the lateral prefrontal cortex [18]. These previous and present findings may be

explained by the classic anatomical compensation theory, which proposes that the size and

depth of a sulcus counterbalance those of the neighboring sulci [23,48,73]. Thus, a larger,

deeper sulcus would be surrounded by sulci that are smaller and shallower, rendering the

overall degree of cortical folding within a given region approximately equal [23,48,73]. Future

work can incorporate underlying white matter architecture into the compensation theory, as well

as a recent modification that proposed to also incorporate local morphological features such as

the deepest sulcal point (e.g., sulcal pit or sulcal root [74]), which has recently been shown to be

related to different functional features of the cerebral cortex [50,51,75]. Altogether, these and

recent findings begin to build a multimodal mechanistic neuroanatomical understanding

underlying the complex relationship between sulcal depth and cognition relative to other

anatomical features.

(d) Limitations

The main limitation of our study is that presently, the most accurate methodology to define sulci

—especially the small, shallow, and variable tertiary sulci—requires researchers to manually

trace each structure on the cortical surface reconstructions. This method is arduous and time-

consuming, which, on the one hand, limits the sample size in terms of number of participants,

while on the other, results in thousands of precisely defined sulci - a push-pull relationship

reflecting a conversation in the broader human brain mapping and cognitive neuroscience fields

between a balance of large N studies and "precision imaging" studies in individual participants

[76–79]. Though our sample size is comparable to other studies that produced reliable results

relating sulcal morphology to brain function and cognition (e.g., [3,4,8,10,12,13,17–19,46]),

ongoing work that uses deep learning algorithms to automatically define sulci should result in

much larger sample sizes in future studies [80,81]. Finally, the time-consuming manual

definitions of primary, secondary, and tertiary sulci also limit the cortical expanse explored in

each study, thus, restricting the present study to LPC/LPOJ.

(e) <u>Conclusion</u>

In conclusion, we uncovered four previously-undefined sulci in LPC/LPOJ and quantitatively

showed that the slocs-v is a stable sulcal landmark that is morphologically, architecturally, and

functionally differentiable from surrounding sulci. We further used a data-driven, model-based

approach relating sulcal morphology to behavior, which identified different relationships of

ventral and dorsal LPC/LPOJ sulcal networks contributing to the perception of spatial

orientation. The model identified the slocs-v, further indicating the importance of this new

neuroanatomical structure. Altogether, this work provides a scaffolding for future "precision

imaging" studies interested in understanding how anatomical and functional features of

LPC/LPOJ relate to cognitive performance at the individual level.

Ethics statement

Human participants: HCP consortium data were previously acquired using protocols approved by the Washington University Institutional Review Board (Mapping the Human Connectome: Structure, Function, and Heritability; IRB # 201204036). Informed consent was obtained from all participants. The data are treated according to the WU-Minn HCP Consortium data use terms and the terms of use for the restricted data.

Competing interests statement

The authors declare no competing financial interests.

Data accessibility statement

The processed and anonymized HCP neuroimaging data required to perform all statistical analyses and reproduce all figures used for this project are available on GitHub (https://github.com/cnl-berkeley/stable_projects/tree/main/Updating_LPOJ_sulci) and Dryad (DOI: doi:10.5061/dryad.rbnzs7hh9) repositories. The analysis pipelines are available on Open Science Framework (https://osf.io/7fwqk/).

Acknowledgments

We thank Jacob Miller, Benjamin Parker, and Willa Voorhies for helping develop the analysis pipelines implemented in this project. We also thank the HCP researchers for participant recruitment and data collection and sharing, as well as the participants who participated in the study.

Funding information

This research was supported by NSF CAREER Award 2042251 (PI Weiner). Neuroimaging and

560   behavioral data were provided by the HCP, WU-Minn Consortium (PIs: David Van Essen and

561   Kamil Ugurbil; NIH Grant 1U54-MH-091657) funded by the 16 NIH Institutes and Centers that

562   support the NIH Blueprint for Neuroscience Research, and the McDonnell Center for Systems

563   Neuroscience at Washington University.

References

1. Amiez C, Wilson CRE, Procyk E. 2018 Variations of cingulate sulcal organization and link with cognitive performance. *Sci. Rep.* **8**, 1–13. (doi:10.1038/s41598-018-32088-9)

2. Amiez C, Sallet J, Hopkins WD, Meguerditchian A, Hadj-Bouziane F, Ben Hamed S, Wilson CRE, Procyk E, Petrides M. 2019 Sulcal organization in the medial frontal cortex provides insights into primate brain evolution. *Nat. Commun.* **10**, 1–14. (doi:10.1038/s41467-019-11347-x)

3. Cachia A, Borst G, Jardri R, Raznahan A, Murray GK, Mangin J-F, Plaze M. 2021 Towards Deciphering the Fetal Foundation of Normal Cognition and Cognitive Symptoms From Sulcation of the Cortex. *Front. Neuroanat.* **15**, 712862. (doi:10.3389/fnana.2021.712862)

4. Garrison JR, Fernyhough C, McCarthy-Jones S, Haggard M, Australian Schizophrenia Research Bank, Simons JS. 2015 Paracingulate sulcus morphology is associated with hallucinations in the human brain. *Nat. Commun.* **6**, 8956. (doi:10.1038/ncomms9956)

5. Nakamura M, Nestor PG, Shenton ME. 2020 Orbitofrontal Sulcogyral Pattern as a Transdiagnostic Trait Marker of Early Neurodevelopment in the Social Brain. *Clin. EEG Neurosci.* **51**, 275–284. (doi:10.1177/1550059420904180)

6. Ammons CJ, Winslett M-E, Bice J, Patel P, May KE, Kana RK. 2021 The mid-fusiform sulcus in autism spectrum disorder: Establishing a novel anatomical landmark related to face processing. *Autism Res.* **14**, 53–64. (doi:10.1002/aur.2425)

7. Fornito A, Yücel M, Wood S, Stuart GW, Buchanan J-A, Proffitt T, Anderson V, Velakoulis D, Pantelis C. 2004 Individual differences in anterior cingulate/paracingulate morphology are related to executive functions in healthy males. *Cereb. Cortex* **14**, 424–431. (doi:10.1093/cercor/bhh004)

8. Lopez-Persem A, Verhagen L, Amiez C, Petrides M, Sallet J. 2019 The Human Ventromedial Prefrontal Cortex: Sulcal Morphology and Its Influence on Functional Organization. *J. Neurosci.* **39**, 3627–3639. (doi:10.1523/JNEUROSCI.2060-18.2019)

9. Harper L *et al.* 2022 Prenatal Gyrification Pattern Affects Age at Onset in Frontotemporal Dementia. *Cereb. Cortex* (doi:10.1093/cercor/bhab457)

10. Weiner KS. 2019 The Mid-Fusiform Sulcus (sulcus sagittalis gyri fusiformis). *Anat. Rec.* **302**, 1491–1503. (doi:10.1002/ar.24041)

11. Miller JA, Voorhies WI, Li X, Raghuram I, Palomero-Gallagher N, Zilles K, Sherwood CC, Hopkins WD, Weiner KS. 2020 Sulcal morphology of ventral temporal cortex is shared between humans and other hominoids. *Sci. Rep.* **10**, 17132. (doi:10.1038/s41598-020-73213-x)

12. Miller JA, Voorhies WI, Lurie DJ, D'Esposito M, Weiner KS. 2021 Overlooked Tertiary Sulci Serve as a Meso-Scale Link between Microstructural and Functional Properties of Human Lateral Prefrontal Cortex. *J. Neurosci.* **41**, 2229–2244. (doi:10.1523/JNEUROSCI.2362-20.2021)

13. Willbrand EH *et al.* 2022 Uncovering a tripartite landmark in posterior cingulate cortex.
Science Advances **8**, eabn9516. (doi:10.1126/sciadv.abn9516)

14. Hathaway CB, Voorhies WI, Sathishkumar N, Mittal C, Yao JK, Miller JA, Parker BJ,
Weiner KS. 2023 Defining putative tertiary sulci in lateral prefrontal cortex in chimpanzees
using human predictions. *Brain Struct. Funct.* (doi:10.1007/s00429-023-02638-7)

15. Willbrand EH, Maboudian SA, Kelly JP, Parker BJ, Foster BL, Weiner KS. 2023 Sulcal
morphology of posteromedial cortex substantially differs between humans and
chimpanzees. *Communications Biology* **6**, 1–14. (doi:10.1038/s42003-023-04953-5)

16. Willbrand EH, Ferrer E, Bunge SA, Weiner KS. 2023 Development of human lateral
prefrontal sulcal morphology and its relation to reasoning performance. *J. Neurosci.*
(doi:10.1523/JNEUROSCI.1745-22.2023)

17. Voorhies WI, Miller JA, Yao JK, Bunge SA, Weiner KS. 2021 Cognitive insights from tertiary
sulci in prefrontal cortex. *Nat. Commun.* **12**, 5122. (doi:10.1038/s41467-021-25162-w)

18. Yao JK, Voorhies WI, Miller JA, Bunge SA, Weiner KS. 2022 Sulcal depth in prefrontal
cortex: a novel predictor of working memory performance. *Cereb. Cortex* , bhac173.
(doi:10.1093/cercor/bhac173)

19. Willbrand EH, Voorhies WI, Yao JK, Weiner KS, Bunge SA. 2022 Presence or absence of a
prefrontal sulcus is linked to reasoning performance during child development. *Brain Struct.*
Funct. **227**, 2543–2551. (doi:10.1007/s00429-022-02539-1)

20. Parker BJ *et al.* 2023 Hominoid-specific sulcal variability is related to face perception ability.
Brain Struct. Funct. (doi:10.1007/s00429-023-02611-4)

21. Sanides F. 1964 Structure and function of the human frontal lobe. *Neuropsychologia* **2**,
209–219. (doi:10.1016/0028-3932(64)90005-3)

22. Van Essen DC, Donahue CJ, Glasser MF. 2018 Development and Evolution of Cerebral
and Cerebellar Cortex. *Brain Behav. Evol.* **91**, 158–169. (doi:10.1159/000489943)

23. Zilles K, Palomero-Gallagher N, Amunts K. 2013 Development of cortical folding during
evolution and ontogeny. *Trends Neurosci.* **36**, 275–284. (doi:10.1016/j.tins.2013.01.006)

24. Petrides M. 2019 *Atlas of the Morphology of the Human Cerebral Cortex on the Average*
MNI Brain. Academic Press. See
https://play.google.com/store/books/details?id=qeWcBAAAQBAJ.

25. Segal E, Petrides M. 2012 The morphology and variability of the caudal rami of the superior
temporal sulcus. *Eur. J. Neurosci.* **36**, 2035–2053. (doi:10.1111/j.1460-9568.2012.08109.x)

26. Zlatkina V, Petrides M. 2014 Morphological patterns of the intraparietal sulcus and the
anterior intermediate parietal sulcus of Jensen in the human brain. *Proc. Biol. Sci.* **281**.
(doi:10.1098/rspb.2014.1493)

27. Drudik K, Zlatkina V, Petrides M. 2022 Morphological patterns and spatial probability maps
of the superior parietal sulcus in the human brain. *Cereb. Cortex*

639  (doi:10.1093/cercor/bhac132)

640 28. Goodale MA, Milner AD. 1992 Separate visual pathways for perception and action. *Trends*
641   *Neurosci.* **15**, 20–25. (doi:10.1016/0166-2236(92)90344-8)

642 29. Humphreys GF, Tibon R. 2023 Dual-axes of functional organisation across lateral parietal
643   cortex: the angular gyrus forms part of a multi-modal buffering system. *Brain Struct. Funct.*
644   **228**, 341–352. (doi:10.1007/s00429-022-02510-0)

645 30. Harvey BM, Klein BP, Petridou N, Dumoulin SO. 2013 Topographic representation of
646   numerosity in the human parietal cortex. *Science* **341**, 1123–1126.
647   (doi:10.1126/science.1239052)

648 31. Harvey BM, Fracasso A, Petridou N, Dumoulin SO. 2015 Topographic representations of
649   object size and relationships with numerosity reveal generalized quantity processing in
650   human parietal cortex. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 13525–13530.
651   (doi:10.1073/pnas.1515414112)

652 32. Konen CS, Kastner S. 2008 Representation of eye movements and stimulus motion in
653   topographically organized areas of human posterior parietal cortex. *J. Neurosci.* **28**, 8361–
654   8375. (doi:10.1523/JNEUROSCI.1930-08.2008)

655 33. Mackey WE, Winawer J, Curtis CE. 2017 Visual field map clusters in human frontoparietal
656   cortex. *Elife* **6**. (doi:10.7554/eLife.22974)

657 34. Schurz M, Tholen MG, Perner J, Mars RB, Sallet J. 2017 Specifying the brain anatomy
658   underlying temporo-parietal junction activations for theory of mind: A review using
659   probabilistic atlases from different imaging modalities. *Hum. Brain Mapp.* **38**, 4788–4805.
660   (doi:10.1002/hbm.23675)

661 35. Tomaiuolo F, Giordano F. 2016 Cerebal sulci and gyri are intrinsic landmarks for brain
662   navigation in individual subjects: an instrument to assist neurosurgeons in preserving
663   cognitive function in brain tumour surgery (Commentary on Zlatkina et al.). *Eur. J. Neurosci.*
664   **43**, 1266–1267. (doi:10.1111/ejn.13072)

665 36. Tomaiuolo F, Raffa G, Morelli A, Rizzo V, Germanó A, Petrides M. 2022 Sulci and gyri are
666   topological cerebral landmarks in individual subjects: a study of brain navigation during
667   tumour resection. *Eur. J. Neurosci.* **55**, 2037–2046. (doi:10.1111/ejn.15668)

668 37. Glasser MF *et al.* 2016 A multi-modal parcellation of human cerebral cortex. *Nature* **536**,
669   171–178. (doi:10.1038/nature18933)

670 38. Vendetti MS, Bunge SA. 2014 Evolutionary and developmental changes in the lateral
671   frontoparietal network: a little goes a long way for higher-level cognition. *Neuron* **84**, 906–
672   917. (doi:10.1016/j.neuron.2014.09.035)

673 39. Wendelken C. 2014 Meta-analysis: how does posterior parietal cortex contribute to
674   reasoning? *Front. Hum. Neurosci.* **8**, 1042. (doi:10.3389/fnhum.2014.01042)

675 40. Karnath HO. 1997 Spatial orientation and the representation of space with parietal lobe
676   lesions. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **352**, 1411–1419.

(doi:10.1098/rstb.1997.0127)

41. Gur RC *et al.* 2000 An fMRI study of sex differences in regional activation to a verbal and a
spatial task. *Brain Lang.* **74**, 157–170. (doi:10.1006/brln.2000.2325)

42. Dale AM, Fischl B, Sereno MI. 1999 Cortical surface-based analysis. I. Segmentation and
surface reconstruction. *Neuroimage* **9**, 179–194. (doi:10.1006/nimg.1998.0395)

43. Fischl B, Sereno MI, Dale AM. 1999 Cortical surface-based analysis. II: Inflation, flattening,
and a surface-based coordinate system. *Neuroimage* **9**, 195–207.
(doi:10.1006/nimg.1998.0396)

44. Glasser MF *et al.* 2013 The minimal preprocessing pipelines for the Human Connectome
Project. *Neuroimage* **80**, 105–124. (doi:10.1016/j.neuroimage.2013.04.127)

45. Barch DM *et al.* 2013 Function in the human connectome: task-fMRI and individual
differences in behavior. *Neuroimage* **80**, 169–189. (doi:10.1016/j.neuroimage.2013.05.033)

46. Roell M, Cachia A, Matejko AA, Houdé O, Ansari D, Borst G. 2021 Sulcation of the
intraparietal sulcus is related to symbolic but not non-symbolic number skills. *Dev. Cogn.*
Neurosci. **51**, 100998. (doi:10.1016/j.dcn.2021.100998)

47. Welker W. 1990 Why Does Cerebral Cortex Fissure and Fold? In *Cerebral Cortex:*
Comparative Structure and Evolution of Cerebral Cortex, Part II (eds EG Jones, A Peters),
pp. 3–136. Boston, MA: Springer US. (doi:10.1007/978-1-4615-3824-0_1)

48. Armstrong E, Schleicher A, Omran H, Curtis M, Zilles K. 1995 The ontogeny of human
gyrification. *Cereb. Cortex* **5**, 56–63. (doi:10.1093/cercor/5.1.56)

49. Chi JG, Dooling EC, Gilles FH. 1977 Gyral development of the human brain. *Ann. Neurol.*
1, 86–93. (doi:10.1002/ana.410010109)

50. Natu VS, Arcaro MJ, Barnett MA, Gomez J, Livingstone M, Grill-Spector K, Weiner KS.
2021 Sulcal Depth in the Medial Ventral Temporal Cortex Predicts the Location of a Place-
Selective Region in Macaques, Children, and Adults. *Cereb. Cortex* **31**, 48–61.
(doi:10.1093/cercor/bhaa203)

51. Leroy F *et al.* 2015 New human-specific brain landmark: the depth asymmetry of superior
temporal sulcus. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 1208–1213.
(doi:10.1073/pnas.1412389112)

52. Glasser MF, Van Essen DC. 2011 Mapping human cortical areas in vivo based on myelin
content as revealed by T1- and T2-weighted MRI. *J. Neurosci.* **31**, 11597–11616.
(doi:10.1523/JNEUROSCI.2180-11.2011)

53. Willbrand EH, Bunge SA, Weiner KS. 2023 Neuroanatomical and functional dissociations
between variably present anterior lateral prefrontal sulci. *bioRxiv.* , 2023.05.25.542301.
(doi:10.1101/2023.05.25.542301)

54. Kong R *et al.* 2019 Spatial Topography of Individual-Specific Cortical Networks Predicts
Human Cognition, Personality, and Emotion. *Cereb. Cortex* **29**, 2533–2551.

(doi:10.1093/cercor/bhy123)

55. Heinze G, Wallisch C, Dunkler D. 2018 Variable selection - A review and recommendations
for the practicing statistician. *Biom. J.* **60**, 431–449. (doi:10.1002/bimj.201700067)

56. Dickerson BC *et al.* 2008 Detection of cortical thickness correlates of cognitive
performance: Reliability across MRI scan sessions, scanners, and field strengths.
Neuroimage **39**, 10–18. (doi:10.1016/j.neuroimage.2007.08.042)

57. Gogtay N *et al.* 2004 Dynamic mapping of human cortical development during childhood
through early adulthood. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 8174–8179.
(doi:10.1073/pnas.0402680101)

58. Wagenmakers E-J, Farrell S. 2004 AIC model selection using Akaike weights. *Psychon.*
Bull. Rev. **11**, 192–196. (doi:10.3758/bf03206482)

59. Kail R, Salthouse TA. 1994 Processing speed as a mental capacity. *Acta Psychol.* **86**,
199–225. (doi:10.1016/0001-6918(94)90003-5)

60. Amunts K, Mohlberg H, Bludau S, Zilles K. 2020 Julich-Brain: A 3D probabilistic atlas of the
human brain's cytoarchitecture. *Science* **369**, 988–992. (doi:10.1126/science.abb4588)

61. Cunningham DJ. 1892 *Contribution to the Surface Anatomy of the Cerebral Hemispheres*.
Academy House. See https://play.google.com/store/books/details?id=t4VCAQAAIAAJ.

62. Wang L, Mruczek REB, Arcaro MJ, Kastner S. 2015 Probabilistic Maps of Visual
Topography in Human Cortex. *Cereb. Cortex* **25**, 3911–3931. (doi:10.1093/cercor/bhu277)

63. Benson NC *et al.* 2018 The Human Connectome Project 7 Tesla retinotopy dataset:
Description and population receptive field analysis. *J. Vis.* **18**, 23. (doi:10.1167/18.13.23)

64. Caspers S, Eickhoff SB, Rick T, von Kapri A, Kuhlen T, Huang R, Shah NJ, Zilles K. 2011
Probabilistic fibre tract analysis of cytoarchitectonically defined human inferior parietal
lobule areas reveals similarities to macaques. *Neuroimage* **58**, 362–380.
(doi:10.1016/j.neuroimage.2011.06.027)

65. Wang J, Fan L, Zhang Y, Liu Y, Jiang D, Zhang Y, Yu C, Jiang T. 2012 Tractography-
based parcellation of the human left inferior parietal lobule. *Neuroimage* **63**, 641–652.
(doi:10.1016/j.neuroimage.2012.07.045)

66. Reveley C, Seth AK, Pierpaoli C, Silva AC, Yu D, Saunders RC, Leopold DA, Ye FQ. 2015
Superficial white matter fiber systems impede detection of long-range cortical connections
in diffusion MR tractography. *Proc. Natl. Acad. Sci. U. S. A.* **112**, E2820–8.
(doi:10.1073/pnas.1418198112)

67. Van Essen DC, Jbabdi S, Sotiropoulos SN, Chen C, Dikranian K, Coalson T, Harwell J,
Behrens TEJ, Glasser MF. 2014 Mapping connections in humans and non-human
primates. In *Diffusion MRI*, pp. 337–358. Elsevier. (doi:10.1016/b978-0-12-396460-
1.00016-0)

68. Cottaar M, Bastiani M, Boddu N, Glasser MF, Haber S, van Essen DC, Sotiropoulos SN,

Jbabdi S. 2021 Modelling white matter in gyral blades as a continuous vector field. *Neuroimage* **227**, 117693. (doi:10.1016/j.neuroimage.2020.117693)

69. Schilling K, Gao Y, Janve V, Stepniewska I, Landman BA, Anderson AW. 2018 Confirmation of a gyral bias in diffusion MRI fiber tractography. *Hum. Brain Mapp.* **39**, 1449–1466. (doi:10.1002/hbm.23936)

70. Schilling KG *et al.* 2023 Superficial white matter across development, young adulthood, and aging: volume, thickness, and relationship with cortical features. *Brain Struct. Funct.* **228**, 1019–1031. (doi:10.1007/s00429-023-02642-x)

71. Pron A, Deruelle C, Coulon O. 2021 U-shape short-range extrinsic connectivity organisation around the human central sulcus. *Brain Struct. Funct.* **226**, 179–193. (doi:10.1007/s00429-020-02177-5)

72. Bodin C, Pron A, Le Mao M, Régis J, Belin P, Coulon O. 2021 Plis de passage in the superior temporal sulcus: Morphology and local connectivity. *Neuroimage* **225**, 117513. (doi:10.1016/j.neuroimage.2020.117513)

73. Connolly CJ. 1950 *External morphology of the primate brain*. CC Thomas.

74. Régis J, Mangin J-F, Ochiai T, Frouin V, Riviére D, Cachia A, Tamura M, Samson Y. 2005 'Sulcal Root' Generic Model: a Hypothesis to Overcome the Variability of the Human Cortex Folding Patterns. *Neurol. Med. Chir.* **45**, 1–17. (doi:10.2176/nmc.45.1)

75. Bodin C, Takerkart S, Belin P, Coulon O. 2018 Anatomo-functional correspondence in the superior temporal sulcus. *Brain Struct. Funct.* **223**, 221–232. (doi:10.1007/s00429-017-1483-2)

76. Naselaris T, Allen E, Kay K. 2021 Extensive sampling for complete models of individual brains. *Current Opinion in Behavioral Sciences* **40**, 45–51. (doi:10.1016/j.cobeha.2020.12.008)

77. Gratton C, Nelson SM, Gordon EM. 2022 Brain-behavior correlations: Two paths toward reliability. *Neuron.* **110**, 1446–1449. (doi:10.1016/j.neuron.2022.04.018)

78. Rosenberg MD, Finn ES. 2022 How to establish robust brain-behavior relationships without thousands of individuals. *Nat. Neurosci.* **25**, 835–837. (doi:10.1038/s41593-022-01110-9)

79. Allen EJ *et al.* 2022 A massive 7T fMRI dataset to bridge cognitive neuroscience and artificial intelligence. *Nat. Neurosci.* **25**, 116–126. (doi:10.1038/s41593-021-00962-x)

80. Borne L, Rivière D, Mancip M, Mangin J-F. 2020 Automatic labeling of cortical sulci using patch- or CNN-based segmentation techniques combined with bottom-up geometric constraints. *Med. Image Anal.* **62**, 101651. (doi:10.1016/j.media.2020.101651)

81. Lyu I *et al.* 2021 Labeling lateral prefrontal sulci using spherical data augmentation and context-aware training. *NeuroImage.* **229**, 117758. (doi:10.1016/j.neuroimage.2021.117758)