The involvement of the human prefrontal cortex in the emergence of visual awareness

Ze- peng Fang[1], Yuan-yuan Dang[2], Zhi-pei Ling[2], Yong- zheng Han[3], Hu- lin Zhao[2*], Xin Xu[2*], Ming- sha Zhang[1*]

[1] *State Key Laboratory of Cognitive Neuroscience and Learning and IDG/McGovern Institute for Brain Research, Division of Psychology, Beijing Normal University, Beijing 100875, China*

[2] *Department of Neurosurgery, Chinese PLA General Hospital, Beijing 100853, China*

[3] *Department of Anesthesiology, Peking University Third Hospital, Beijing, China*

[*] Corresponding authors

zhaohulin_90@sohu.com

xuxinmm@hotmail.com

mingsha.zhang@bnu.edu.cn

Abstract

Exploring the neural mechanisms of awareness is a fundamental task of cognitive neuroscience. There is an ongoing dispute regarding the role of the prefrontal cortex (PFC) in the emergence of awareness, which is partially raised by the confound between report- and awareness-related activity. To address this problem, we designed a visual awareness task that can minimize report-related confounding. Our results show that saccadic latency is significantly shorter in the aware trials than in the unaware trials. Local field potential (LFP) data from 6 patients consistently show early (200-300 ms) awareness-related activity in the PFC, including event-related potential and high-gamma activity. Moreover, the awareness state can be reliably decoded by the neural activity in the PFC since the early stage, and the neural pattern is dynamically changed rather than being stable during the representation of awareness. Furthermore, the enhancement of dynamic functional connectivity, through the phase modulation at low frequency, between the PFC and other brain regions in the early stage of the awareness trials may explain the mechanism of conscious access. And the spatiotemporal overlap between the awareness-related activity and the inter-regional connectivity in PFC suggested that conscious access and phenomenal awareness may be closely coupled. These results indicate that the PFC is critically involved in the emergence of awareness.

Keywords: visual consciousness, local field potential (LFP), saccade, neural correlates of consciousness (NCC), stereoelectroencephalography (sEEG).

Introduction

A conscious state means that a person has some kind of subjective experiences, such as seeing a painting, hearing a sound, thinking about a problem, or feeling an emotion [1]. The neural mechanism of consciousness is one of the most fundamental, exciting and challenging pursuits in

the 21st-century brain science [2]. Among the various aspects of consciousness, visual awareness, as a core of consciousness, has attracted enormous attention in consciousness research. Visual awareness seems quite simple: we see an object and seem to immediately know its shape, color and other properties. However, the neural processes behind the emergence of visual awareness are quite sophisticated and largely unknown [3].

The typical method to study the neural mechanism of visual awareness is to present external stimuli with similar physical properties but to elicit different conscious experiences, such as using near perceptual threshold stimuli, visual masking, or attentional blinks [4,5]. Then, the aware and unaware states are determined through the subjective reports of participants, and the awareness-related neural activity is dissociated by the brain activation difference between the two states. Using this approach, some studies have found that activities in broad brain regions are correlated with visual awareness. For example, an fMRI study of lexical visual masking found that only consciously perceived and reported words can induce activation of higher-order cortical regions such as the frontal and parietal lobes [6]. Another study using a visual masking task found changes in the amplitude of intracranial electroencephalography (iEEG) signals and high-gamma power in occipital-temporal regions, which indicated that these regions played an important role in the emergence of awareness [7].

However, there is still much debate about the origin of conscious awareness in the brain. Some hypotheses (global neuronal workspace theory, higher-order theory, etc.) argued that visual awareness arose from the frontoparietal network, in which the prefrontal cortex played a key role, and they argued that the earliest biomarker of awareness in EEG is the P3b wave onset approximately 300 ms after stimulus onset [8,9]. For example, based on the electrophysiological study of a visual awareness-guided saccade selection paradigm in macaques, by comparing the action potential firing characteristics of neurons in the dorsolateral prefrontal cortex (dlPFC), V4 and V1, only dlPFC neurons were closely associated with visual awareness states. Therefore, the authors argued that stimuli must elicit a threshold level of prefrontal activity for conscious detection to occur [10]. In contrast, other hypotheses (integrated information theory, recurrent processing theory, etc.) proposed that the "hot zone" of the posterior cortex of the brain, including the parietal lobe, occipital lobe, and temporal lobe, is sufficient to generate visual awareness, and its biomarkers on EEG are the relatively earlier visual awareness negativity (VAN) at approximately 200 ms after stimulus onset [1,11-13], whereas the PFC is not necessary for the emergence of conscious experience.

This controversy is partially caused by the confounding effect of report-related activity, such as motor planning and execution, on awareness-related activity. The reason is that the study of awareness usually requires subjective reports to determine different awareness states, while the prefrontal cortex is closely related to such report-related functions [14]. To solve this problem, some researchers use the no-report paradigm to rule out report-related impacts. For instance, a functional magnetic resonance imaging (fMRI) study employing human binocular rivalry paradigms found that when subjects need to manually report their awareness states, the frontal, parietal, and occipital lobes are all involved in the process of binocular rivalry, but when a report is not needed, rivalry-related activation disappears in the frontal lobe but remains in the occipital and parietal lobes [15]. However, the no-report paradigm may overestimate unconscious processing because it cannot directly measure the awareness state but infers the awareness state through other relevant physiological indicators, such as optokinetic nystagmus and pupil size [16]. Therefore, neither the typical report paradigm nor the no-report paradigm is ideal for visual awareness research; thus, it is

difficult to solve the problem of whether the PFC participates in conscious perception.

In addition, when studying the role of the PFC in visual awareness, the noninvasive
technologies mostly used in previous studies, i.e., Functional magnetic resonance imaging (fMRI)
and M/EEG have limitations. Although fMRI research can detect the whole brain activation pattern,
it is difficult to describe the dynamic changes and sequences of different brain regions in the rapid
process of visual awareness due to its limited time resolution (approximately several seconds). For
example, even though a recent fMRI study using the no-reported paradigm found that there are still
visual awareness-related activities in the PFC [17], it remains unclear when these activities begin and
whether PFC activity occurs earlier or later than other brain regions. At the same time, although
EEG and MEG have a very high temporal resolution, their low spatial resolution and low signal-to-
noise ratio limit their interpretation of awareness-related activities, especially high-frequency
activities (highly correlated with local population neuronal activities). For example, although
increasing evidence shows that the earliest biomarker of visual awareness in EEG is more likely to
be the VAN, which starts at approximately 180-300 ms (may be delayed to 300-460 ms for low-
visible stimulus) and peaks at the parieto-occipital region [18-20], rather than the P3b wave onset at
approximately 300-500 ms which peaks in the central frontoparietal region, it is still unclear whether
there are local visual awareness-related activities in the PFC during the VAN period. In addition, a
few recent studies have suggested that awareness may be related to the activities of brain networks,
including the PFC, rather than that of a single brain region. However, most of these brain network
studies used noninvasive research methods [17,21], and due to the limitations mentioned above, it is
difficult to detect the dynamic changes in brain networks in a more precise spatial-temporal profile
during the generation of visual awareness.

To address the questions mentioned above, we first designed a novel visual awareness task
(Figure 1). The contrast of the cue stimulus (grating) was roughly divided into three levels: well
above the visual perceptual threshold, near the visual perceptual threshold and zero contrast (no
stimulus). During each experiment, in nearly 80% of trials, the contrast of the grating was close to
the subject's visual perceptual threshold so that the subjects could sometimes see the grating and
sometimes could not. Subjects were required to choose a saccade direction according to the visual
awareness state (seeing the grating or not seeing the grating) and the color of the fixation point.
Notably, subjects were unable to choose the saccade direction or effectively prepare for saccades
until the color of the fixation point was changed. Moreover, the subjects needed to choose the
saccade direction, both in the awareness and unawareness states, according to the color of the
fixation point so that the report behaviors were matched between the two awareness states. Such a
paradigm can effectively dissociate awareness-related activity from report-related activity in terms
of time (the fixation point changes color 650 ms after the grating onset) and report behavior, thereby
being report-independent of the awareness states [22], thus minimizing the impact of report-related
factors on visual awareness. In addition, saccade parameters (such as reaction time, etc.) under
different awareness states can be compared to explore the influence of visual awareness on saccade
behavior. Then, we applied the above behavioral paradigm to 6 clinical patients with implanted
electrodes (stereoelectroencephalography, sEEG) in prefrontal and other cortices and 10 healthy
subjects. The behavior and local field potential (LFP) data of 6 patients were recorded while they
performed the task, and we also collected the behavioral data of 10 healthy subjects for comparison
with the patients' behavioral data.

Figure 1. Schematic diagram of the visual awareness task. A trial started when a fixation point (0.5°x0.5°, white cross) appeared at the center of the screen (radius of eye position check window is 4°, the dotted circle). After the subject fixated on the fixation point for 600 ms, a cue stimulus (Gabor grating, 2x2° circle) was presented for 50 ms at a fixed position (7°) on the left (or right, see Methods) side of the screen for all participants. In 70% of the trials, the grating contrast was maintained near the subject's perceptual threshold by a staircase method; in 10% of the trials, the stimulus contrast was well above the threshold; and in the other 20% of the trials, the stimulus contrast was 0, namely, no stimulus appeared. After another 600 ms delay, the color of the fixation point turned red or green, and two saccade targets (1x1°, white square) appeared at fixed positions (10°) on the left and right sides of the screen. If the grating was seen, the green fixation point was required to make a saccade to the right target, while the red fixation point was required to make a saccade to the left target. If the grating was not seen, the rule of saccadic direction was inverted.

Results

Behavioral Results

To detect the effect of cue stimulus (grating) contrast on the emergence of visual awareness, we first analyzed the proportion of participants being aware of the grating as a function of its contrast. Figure 2A shows exemplified sessions from a patient and healthy subject, and Figure 2B shows the population results of the two participant groups. The results showed that with increasing grating contrast, the ratio of reporting 'awareness' of the grating gradually increased for all participants (including patients and healthy subjects), showing a classical psychometric curve. Such results showed that the grating contrast level had a direct impact on the emergence of visual awareness and thus is a reliable way to study visual awareness. In addition, in trials with no grating and grating contrast well above the perceptual threshold, subjects showed high accuracy (patients 94.75% ± 2.65; healthy subjects, 96.80% ± 0.71, mean ± SEM), indicating that the subjects understood and performed the task well following the task rules.

To explore whether visual awareness affects behavioral performance, we divided the trials
into 4 groups according to the level of grating contrast and the participants' reported awareness
state, i.e., high-contrast aware (HA), near-threshold-contrast aware (NA), near-threshold-contrast
unaware (NU), and low-contrast unaware (LU) (Figure 2A, also see Methods). We compared the
saccadic parameters between the awareness and unawareness states in near-threshold contrast
levels (NA vs. NU), as well as in high and low contrast levels (HA vs. LU), as a control. While
saccadic reaction time showed a significant difference between the two awareness states (Figure
2B), the other parameters did not. Figure 2B shows the mean saccadic reaction time of patients
and healthy subjects under different awareness states and contrast levels. The results showed that
the average saccadic reaction time was significantly shorter under the awareness state than under
the unawareness state at both the near-threshold contrast level and the high-low contrast level ($p <$
0.05, Wilcoxon signed rank test), and it was also explicit and consistent at the individual level.
The results showed that the saccadic reaction time in the aware trials was systematically shorter
than that in the unaware trials. Such results demonstrate that visual awareness significantly affects
the speed of information processing in the brain.

A

Patient

Healthy Subjects

B

C

Figure 2. Behavioral performance and electrode location

A. Psychometric detection curve in a single session. Left panels from one patient and right from one healthy subject. Each black point in the graph represents the aware percent in a contrast level, and the black curve represents the fitted psychometric function. The contrast level resulted in an awareness percentage greater than 25%, and less than 75% was defined as near-threshold, whereas an awareness percentage less than 25% was low and greater than 75% was high.

B. Psychometric detection curves for all participants. The same as Panel A except the contrast is aligned to the individual subject's perceptual threshold, i.e., the contrast 0 represents each subject's perceptual threshold. (N, number of subjects; R^2, coefficient of determination).

C. Saccadic reaction time in the aware and unaware trials under high+low- and near-threshold-level contrasts of gratings. Bars show the mean values of saccadic reaction time. In the near-threshold condition, the mean reaction

times of patients and healthy subjects in the aware trials were 676.55±42.38 and 556.16 ± 28.45 ms, respectively.
The mean reaction times in the unaware trials were 892.57 ± 43.28 and 702.30 ± 30.93 ms, respectively, and the p
values were 0.03 and 9.77 x10 $^{-4,}$ respectively, Wilcoxon signed-rank test. In the high+low condition, the mean
reaction times for patients and healthy subjects under the aware trials were 577.78 ± 55.28 and 524.08 ± 31.06 ms,
and the mean reaction times of the unaware trials were 898.64 ± 57.21 and 687.59 ± 49.07 ms, respectively, with p
values of 0.002 and 9.77x10$^{-4,}$ respectively, Wilcoxon signed-rank test. Each black dot represents one participant.
The gray solid line represents the paired dots. Error bars represent the standard error of the mean (SEM).
D. Left, right and top views of all recording sites projected on an MNI brain template. Each color represents a
participant.

Local Field Potential Results

189   While the patients were performing our visual awareness task, we recorded the local field
190 potential (LFP) from 901 recording sites (Figure 2D, projected on Montreal Neurological Institute
191 (MNI) brain template ICBM152), in which 245 sites were located in the PFC. The table in Figure
192 2D shows a summary of the proportions of recording sites in different lobes and structures.

Early visual awareness-related iERP activity in the PFC

While looking through the event-related response of the LFP data, a striking phenomenon in
all patients was that the amplitude of LFP showed remarkably different patterns between awareness
and unawareness states. We show the grand average (left) and single-trial data (right) of LFP activity
of recording sites in the PFC from each patient in Figure 3A, which represents the typical awareness-
related neural activity of the PFC. There was a vigorous visual evoked response under the HA
condition (light red line) compared to the LU (light blue line) condition. However, in the near-
threshold condition, although the contrast of the grating was similar (see Results below), the LFP
amplitude of the NA (red line) and NU (blue line) trials started to differ significantly at
approximately 200-300 ms (thick black line in panels, $p < 0.01$ corrected, by independent t-test,
details see Methods). This difference is also very explicit and consistent at the single-trial (>180
trials for NA and NU conditions in individual participants, see Methods) and single-subject level.
The divergence onset time (DOT, see Methods) of different recording sites in the PFC is consistently
within the period of 200~300 ms after grating onset, i.e., the time window of VAN in scalp EEG
mentioned above.
As proposed above, the different activities between NA and NU are regarded as awareness-
related activities. We found 89 sites with awareness-related iERP activity in the PFC (323 for all
recording sites). Figure 3B shows the spatial-temporal dynamics of visual awareness-related
activities in all recording sites during the 0-650 ms period (for a clearer display, projected to the
cortex, see the Supplementary Materials for the complete video). Visual awareness-related activities
clearly appeared in the PFC during 200-300 ms, and these early activities were mainly concentrated
in the middle lateral prefrontal cortex (LPFC) and then spread to other brain areas of the PFC. We
further calculated the start time of awareness-related activity, i.e., DOT, of all recording sites. Figure
3C shows the spatial distribution of all recording sites DOT (standardized and convenient for visual
display), and it can be found that the DOT of the PFC is not explicitly later than those in posterior
brain regions, including the posterior parietal lobe, temporal lobe, occipital lobe, etc., and the earliest
region in PFC is still located in the middle lateral prefrontal cortex.

Considering the possibility that the brain region with earlier awareness-related activity might
be more important for visual awareness, we tried to detect awareness-related activity in the PFC
during the 'early' period, i.e., the VAN time window mentioned above. We focused on the sites in
PFC with earlier DOTs. Because of the low-visible stimulus used in this experiment, the onset time
of VAN may be delayed to 300-350 ms, and we took 350 ms as the cutoff value and divided DOT
into early (n=43, 48.31%) and late (n=46, 51.69%) phases. We focused on the early DOT sites and
calculated the normalized average iERP response of the early DOT sites (Figure 3D). At the
population level, visual awareness-related activities in the PFC were also significant and started at
approximately 200-300 ms, which is located in the VAN time window. Importantly, there were sites
with DOT<350 ms from each patient and these sites from different patients were uniformly
concentrated in the middle lateral prefrontal cortex (Figure 3D, right). That is, the early visual
awareness-related activities in the prefrontal cortex are robust and consistent not only at the group
level but also at the individual level.

233
234
235

Figure 3. Visual awareness-related iERP activities in the PFC.

A. LFP activity of example sites from each patient. Each panel represents one example site from one patient. In each

panel, the left plot shows the grand average of LFP activity in different conditions. The pink line represents data

under HA conditions, the red line represents data under NA conditions, the blue line represents data under NU

conditions, and the light blue line represents data under LU conditions. The shaded area of the curve represents the

SEM. The two black dotted lines at 0 ms and 650 ms represent the time when the grating and fixation point change

color (the appearance of saccade targets), and the black thick solid line area represents the period when the LFP

amplitude is significantly different under NA and NU conditions (p<0.01 corrected, independent sample t-test). The

right figure in each panel shows the single-trial data in the NA (upper) and NU (lower) conditions. The color

represents the voltage.

B. The spatial-temporal distribution of awareness-related ERP activities after the appearance of the grating. Each

brain image showed a significant (p<0.01 corrected, see Methods) difference in local field potential in NA and NU

trials at all visual awareness-related sites at a specific time point (lower left corner) after the grating appeared. The

color represents the standardized voltage difference (see Methods).

C. Spatial distribution of divergence onset time. The color represents the normalized DOT value.

D. Population results of early phase ERP response in the prefrontal cortex. Different lines represent different

conditions, as shown in Panel A. The right table shows the number of recording sites for different patients. The lower

right is the location of these sites. The dots with different colors represent different subjects, as shown in Figure 2D.

Early visual awareness-related high-gamma activity in the PFC

Although the above iERP results indicate that there are early visual awareness-related neural

activities in the prefrontal cortex, it is difficult to explain whether this response is the result of local

processing in the PFC or transmission from other brain regions. Therefore, we calculated the LFP

response in the high-gamma (HG) band, which is more representative of the local processing, of the

prefrontal cortex. Figure 4A shows the exemplified high-gamma activity in the prefrontal cortex

from each patient at the grand average (left), spectrogram (middle), and single-trial level (right). It

is similar to the iERP results (Figure 3) except the data are the HG power and there was a vigorous

visual evoked power increase under the HA condition compared to the LU condition. However, in

the near-threshold condition, while the contrast of the grating was similar, the high-gamma power

of the NA and NU trials started to differ significantly after ~ 230 ms (thick black line in panels, p <

0.01 corrected by independent t-test; for details, see Methods). This difference is also very explicit

and consistent at the single-trial and single-subject levels. The HG activity divergence onset time

(DOT) of different recording sites in the PFC was also consistent within the early stage, which was

similar to the above iERP result.

We found 31 sites with awareness-related HG activity in the PFC (74 for all recording sites).
Figure 4B shows the spatial-temporal dynamics of visual awareness-related HG activities in all
recording sites during the 0-650 ms period (for a clearer display, projected to the cortex, see the
Supplementary Materials for the complete video). Visual awareness-related HG activities also
clearly appeared in the PFC during 200-300 ms, while they were rarely seen in lower-level visual
areas (occipital and temporal lobe), and these early activities were mainly concentrated in the middle
lateral prefrontal cortex (LPFC) and then spread to other brain areas of the PFC. We further
calculated the start time of awareness-related HG activity, i.e., DOT, of all recording sites. Figure
4C shows the spatial distribution of all recording sites DOT (standardized and convenient for visual
display), and it can be found that the DOT of the PFC is not explicitly later than those in posterior
brain regions, and the earliest region is still located in the middle lateral prefrontal cortex.

We continue to take 350 ms as the cutoff value and divide the HG DOT into early (n=18,
58.06%) and late (13, 41.94%?). We focused on the early DOT sites and calculated the normalized
mean response (Figure 4D) and spectrum of the early DOT sites. At the population level, early visual
awareness-related HG activities of the prefrontal sites still exist, and the start time is approximately
200 ms, which is within the VAN time window. Importantly, there are sites with HG DOT<350 ms
for each patient except one, and these sites are also uniformly concentrated in the middle lateral
prefrontal cortex. In other words, this effect is significant and consistent at both the group and
individual levels. It is worth noting that in the PFC recording sites, the early sites with a high-gamma
visual response (7.35%) were explicitly less than those with ERP visual response (17.55%), which
may indicate that in the process of visual awareness generation, there may also be a variety of other
bands activity for information interaction between multiple brain regions.

294
295

Figure 4. Visual awareness-related high-gamma activities in the PFC.

A. High-gamma activity of example sites from each patient. Each panel represents one example site from one patient. In each panel, the left figure shows the grand average of high-gamma power in different conditions. The pink line represents data under HA conditions, the red line represents data under NA conditions, the blue line represents data under NU conditions, and the light blue line represents data under LU conditions. The shaded area of the curve represents the SEM. The two black dotted lines at 0 ms and 650 ms represent the time when the grating and fixation point change color (the appearance of saccade targets), and the black thick solid line area represents the period when

the high-gamma power is significantly different under NA and NU conditions (p<0.01 corrected, independent sample
t-test). The right figure in each panel shows the single-trial data in the NA (upper) and NU (lower) conditions. The
color represents the normalized power.
B. The spatial-temporal distribution of awareness-related high-gamma activities after the appearance of the grating.
Each brain image showed a significant (p<0.01 corrected, see Methods) difference in high-gamma activity in NA
and NU trials at all visual awareness-related sites at a specific time point (lower left corner) after the grating appeared.
The color represents the standardized power difference.
C. Spatial distribution of divergence onset time for high-gamma activity. The color represents the normalized DOT
value.
D. Population results of early phase high-gamma response in the prefrontal cortex. Different lines represent different
conditions, as shown in Panel A. The middle panel shows the spectrogram in NA (upper) and NU (lower) trials. The
right table shows the number of recording sites for different patients. The lower right is the location of these sites.
The dots with different colors represent different subjects, as shown in Figure 2D.

Decoding awareness state through ERP and HG activity in the PFC

In addition to the univariate analysis above, we also adopted the machine learning decoding method based on multivariate pattern analysis (MVPA) to test the ability of broadband and high-gamma activities in the PFC to predict the awareness state at the single-trial level. We applied a linear discriminant analysis (LDA) to classify aware versus unaware trials based on broadband and high-gamma LFP activities in the PFC.

Figure 5A shows the decoding performance through the ERP activity in the PFC. We found that the ERP activity of the PFC can accurately predict the state of visual awareness at the single-trial level in the delayed period after grating onset, and this result is also relatively consistent at the individual level. Moreover, this decoding accuracy has begun to be much higher than the chance level in 200-300 ms, consistent with our above univariate analysis results. Furthermore, we applied the temporal generalization method to test the predictive ability across time. The results (Figure 5B) show that the decoding performance of population ERP is very limited in temporal generalization and cannot be well generalized over a long period. Importantly, this phenomenon is also very consistent at the individual level, which indicates that the representation of visual awareness via broadband activities in the PFC is relatively dynamic rather than static. Moreover, we also applied temporal generalization across the control conditions, i.e., train on NA/NU trials but test on HA/LU trials to test the predictive ability across conditions. The classifier trained on the NA/NU trials can be well generalized to the HA/LU trials, and the generalization across time is also very consistent. These results further indicated the robustness of the above decoding results.

Figure 5C-D shows the results of a similar analysis of HG activity in the prefrontal cortex. Figure 5C shows the decoding performance through HG activity in the PFC. We found that the HG activity of the PFC can accurately predict the state of visual awareness at the single-trial level in the delayed period, and this result is also relatively consistent at the individual level. Moreover, this decoding accuracy began to be significantly higher than chance at approximately 200-300 ms, which is also consistent with our above univariate analysis results. Furthermore, we applied the temporal generalization method to the HG activity. Although the group results still show that the decoding performance of HG activity is limited in time generalization, the generalization results are not consistent at the individual level, which may be attributed to the variability of electrode location in different patients. This may suggest that the representation of visual awareness-related information by population neurons at different regions in the PFC is inconsistent, i.e., some are relatively stable,

350 and others are dynamic. Similarly, we applied temporal generalization across the HA/LU conditions,
351 and the result was also very consistent.

Figure 5. Decoding the awareness state through broadband and HG activities.

A. Decoding the awareness state through broadband LFP activity. The population result (left) is the average result

of six individual subjects (right) after decoding analysis.

B. Time generalization of broadband activity decoding. The left panel represents generalization across time. The

right panel represents generalization across different conditions, which means training on NA&NU data and testing

on HA&LU data. In each panel, the population result (left) is the average result of six subjects (right) after decoding

analysis.

C. The awareness state is decoded by HG activity. Same as A, except that the HG magnitude is used for decoding.

D. Time generalization of HG activity decoding. Same as B, except that the HG magnitude is used for decoding.

Dynamic functional connectivity changes in the PFC associated with visual awareness

We further explored the dynamic functional connectivity between the prefrontal cortex and
other brain regions in the generation of visual awareness. First, we calculated the time-across phase-
locking value (PLV) [23] in NA and NU conditions at the sensor level (baseline removed, see Methods
for details). Figure 6A-B shows the example results of two exemplified patients (see SM for other
patients). We found that in the low-frequency band (1-8 Hz), overall, there was a significant
difference in the PLV between the NA and NU trials. In the first 150 ms after the appearance of the
grating, there was no significant difference in the PLV between these two conditions. However, in
the period of approximately 200-300 ms, the PLV between many recoding sites in the NA trials
began to increase explicitly and lasted until after 600 ms. In contrast, there was no such phenomenon
in the NU trials. Importantly, this phenomenon is strikingly explicit and consistent in each patient.

Furthermore, we grouped the recording sites by brain regions according to the DKT brain atlas
[24] and averaged the PLV between different sites and different subjects according to brain regions
(Figure 6C). The functional connectivity between the prefrontal cortex and subcortical structures,
such as the thalamus, and the posterior cortex, such as the posterior parietal cortex, was significantly
enhanced in the NA trials compared with the PLV in NU trials. In addition, the functional
connectivity between the brain regions within the PFC was also significantly enhanced. For clearer
visualization, we show the dynamics of PLV in the NA and NU trials on a topographic map (Figure
6D).

Interestingly, we found that at the single-subject level, the sites in the prefrontal cortex that
showed early awareness-related PLV activity coincided with the above sites with early awareness-
related HG activity. Figure 6E shows the averaged PLV between these sites (n=18) with early visual
awareness-related HG activity in the PFC and other sites in the NA and NU trials. The averaged
PLV also showed significant visual awareness-related activities. Moreover, at the population level,
the start time of awareness-related PLV activity at these sites also started at approximately 200 ms,
which is very similar to that of HG activity.

B

P2 (n=93 sites)

Time from cue onset (ms)

C

Time from cue onset (ms)

D

N=6

Time from cue onset (ms)

Threshold by 85% for visualization

Figure 6. Functional connectivity analysis results.

A-B. Phase-locking value (PLV) changes at the sensor level (NxN) in two example patients. From left to right, the

changes in PLV between each recording site at the five time points of 0/150/300/450/600 ms after the appearance of

the grating in the near-threshold aware (upper) and near-threshold unaware (lower) trials are displayed (baseline

removed, see the Methods). The color represents PLV.

C. The population results of functional connectivity analysis averaged according to brain regions. From left to right,

PLV between different brain regions at the five time points of 0/150/300/450/600 ms after the appearance of the

grating in the near-threshold aware (upper) and near-threshold unaware (lower) trials (baseline removed, see the

Methods). The color represents the PLV.

D. For the same data as C, only the strongest 15% PLV is displayed on the brain template for clear visualization.

E. PLV under different conditions at early-phase HG awareness-related sites in the PFC. The red line represents the

average PLV in the NA trials, and the blue line represents the average PLV in the NU trials. The faded line represents

the PLV of each site.

The different neural activity between NA and NU trials is not caused by the grating contrast

Since there may be a slight difference in the grating contrast under the NA and NU trials, the

aforementioned different activity between the NA and NU conditions may be caused by the

different contrast of grating. To rule out this possibility, we compared the distribution of grating

contrast in the NA and NU conditions for each patient (Figure 7A). Overall, we find that the

distributions of grating contrasts largely overlap with each other (red and blue curves). The

Kolmogorov–Smirnov test resulted in no significant difference ($p > 0.05$) between the two contrast

distributions in all patients except patient 2. Although intuitively the trials with higher contrast

contributed more to the NA condition and the trials with lower contrast contributed more to the

NU condition, the contrast difference between the NA and NU conditions were not significantly

different perhaps because the grating contrast was designed to be near the perceptual threshold

(the 'aware' percent is 50%) in most trials. Moreover, for patient 2, the difference between the

mean grating contrast in the NA and NU trials was also subtle (0.15%), much less than the

difference between NA and HA (17.35%) or NU and LU (0.84%). However, it is clear in the

results above that the different activities between NA and NU were much stronger than those

between the same awareness states, i.e., NA vs. HA or NU vs. LU. Therefore, the difference in

neural activity between aware and unaware trials cannot be explained by the contrast differences

in the grating.

Moreover, we generalized the classifier trained under the NA/NU condition to the NU/LU
condition. If the decoding result under the NA/NU condition is based on the difference in the
grating contrast rather than on the awareness state, then this classifier should also be able to
generalize across the NU/LU condition because the difference in the grating contrast between the
NU/LU conditions is greater than that under the NA/NU condition, while the awareness state
remains the same (both being unaware of the grating). Figure 7B shows the results of
generalization decoding results of ERP (left) and HG (right) data. The classifier trained between
NA/NU cannot be well generalized to NU/LU conditions for broadband or HG data. Therefore, the
above decoding results also cannot be attributed to the difference in grating contrast.

A

B

Figure 7. Grating contrast in NA and NU conditions
A. Distribution of grating contrast in the NA and NU conditions for each patient. Each panel
represents one patient. The red and blue lines represent the distribution of grating contrast in NA
and NU conditions, respectively.
B. Temporal generalization of broadband and HG activity across the NA&NU condition and
NU&LU condition. The left panel represents the generalization of ERP activity across NA&NU
and NU&LU conditions, which means trained on NA&NU data and tested on NU&LU data. The
right panel is the same analysis except for high-gamma activity.

Discussion

We employed a novel visual awareness task and found that, first, the saccadic reaction time of

all subjects was shorter when they were aware of grating than when they were not aware. Second,
there are ERP and HG activities related to visual awareness in the prefrontal LFP, and these activities
begin at the early stage (200-300 ms). Then, we found that the awareness state can be reliably
decoded from the broadband and HG activities of the PFC at approximately 200-300 ms after
stimulus onset. Finally, the functional connectivity between the PFC and other brain regions in the
low-frequency band (1-8 Hz) also appeared awareness-related changes, and the initial time and
location of this change are consistent with the above HG results. The results above are not caused
by the difference in the physical properties of external stimuli. Therefore, we propose that the
prefrontal cortex also plays a key role in the early stage of visual awareness generation even if motor
effects are excluded.

Visual awareness states affect saccadic reaction time

Many factors affect saccadic reaction time, including the characteristics of external stimuli
and the internal state of the brain. The physical properties of external visual stimuli have a direct
impact on the performance of saccades; for example, saccadic reaction time decreases with
increasing visual stimulus contrast [25,26]. A widely accepted view is that the greater contrast of
visual stimuli will automatically attract more bottom-up attention and speed up information
processing in the brain, thereby shortening the saccadic reaction time [27,28]. Other studies found that
endogenous factors, such as top-down attention, can also significantly shorten the saccadic
reaction time [28,29].

Consciousness, as a high-level cognitive function of the brain, should have some similar effects
as other cognitive functions on behavior (for example, saccadic reaction time). With this question
in mind, we carefully searched the literature about the relationship between consciousness and
behavior; surprisingly, we failed to find any relevant literature. One study briefly reported that the
average reaction time (button press) for word semantic classification in the awareness state was
approximately 300 ms faster than in the unawareness state, but the statistical test did not reach a
significant level [30]. In this study, we found that the saccadic reaction time of all subjects in the aware
trials was significantly smaller than that in the unaware trials (Figure 2C). More critically, the grating
contrast near the threshold was not significantly different between the aware and unaware trials
(Figure 7A); thus, such results cannot be caused by the contrast difference in the grating. Here, for
the first time, we showed that awareness states significantly affect saccadic reaction time, which
supports our previous hypothesis.

Visual awareness-related activities appeared in the PFC at an early stage

The long debate about the neural mechanism of consciousness focuses on the location, i.e.,
'front' vs. 'back', and the time, i.e., 'early' vs. 'late', of its origin in the brain [14]. Concerning the
dispute of location, our results show that the prefrontal cortex still displays visual awareness-related
activities even after eliminating the influence of the confounding variables related to subjective
reports such as motion preparation, which indicates that the prefrontal cortex does participate in the
information processing of visual awareness. Regarding the time dispute, our results show that at
200-300 ms after stimulus onset, the VAN time window, the prefrontal cortex has activities related
to visual awareness, and these activities include not only ERP activities but also HG activities, which
are highly related to the firing of local population neurons [7]. Notably, the recording sites with early
ERP and HG visual awareness-related activities are mostly concentrated in the middle lateral
prefrontal cortex. This shows that the prefrontal cortex, especially the middle lateral prefrontal

cortex, started visual awareness-related activities, including local activities, at a relatively early stage.

Some recent studies utilizing the no-report paradigm supported our results. For example, a recent electrophysiological study of rhesus monkeys found that even if monkeys do not need to report their conscious content under the binocular competition paradigm, there were also visual awareness-related spiking activities in the prefrontal cortex [31]. In addition, some previous human intracranial studies also reported ERP, HG, and single-neuron activities related to visual perception in the higher visual areas of the temporal lobe, such as the superior temporal gyrus [7,32]. However, due to the limitations of the experimental paradigm, these studies cannot completely exclude the impact of subjective reports. Moreover, most of the electrodes in previous studies were located in the temporal lobe, and electrodes in the prefrontal cortex are rare. As our data show that the proportion of sites with earlier ERP and HG responses in the prefrontal cortex is low and relatively concentrated, it is not hard to explain that earlier visual awareness-related activities in the prefrontal cortex were not found in previous iEEG studies, especially in the HG band. Our results, on the one hand, support that the prefrontal cortex may play an important role in the emergence of visual awareness and thus support the 'front of the brain' theories of consciousness in the origin brain regions. However, our data also support the prediction of the early emergence of consciousness proposed by the 'back of the brain' theories. This means that our result also challenges some predictions of these mainstream theories.

Awareness state can be effectively decoded from PFC activities from an early stage

Compared with traditional univariate analysis, MVPA-based machine learning decoding may be more sensitive to the information contained in brain activities [33], so it has gradually become a powerful tool for the study of neural correlates of consciousness (NCCs). Therefore, we used linear discriminant analysis (LDA) to decode the awareness states at the single-trial level through broadband and HG activities in the prefrontal cortex. Our decoding results show that whether through broadband or HG activities, the decoding performance began well above the chance level at approximately 200-300 ms. This shows that the prefrontal cortex has begun to encode visual awareness-related information in 200-300 ms, which is consistent with the above univariate analysis results. In addition, a recent electrophysiological study of rhesus monkeys using the no-report paradigm also found that even without the requirement of explicit reporting, awareness-related neurons can still be found in the lateral prefrontal cortex of rhesus monkeys, and conscious content can be accurately decoded by the MVPA method [31]. Another human fMRI study also found similar results [17]. These results supported our experimental results.

In addition, we also applied the temporal generalization method during decoding. Temporal generalization analysis shows how different brain activity patterns change and transform over time [34]. Our results show that the representation pattern of neural activity in the prefrontal cortex for visual awareness-related information is dynamic, not static, and this result is very consistent at the individual level and across the control conditions. Such a result may be consistent with the nature of conscious perception. The process of conscious perception (visual awareness) includes a series of processes from phenomenal consciousness ('qulia') to conscious access. Therefore, at the level of large, high-order brain regions, the representation of this complex process in the brain should be continuously transmitted and transformed, that is, dynamically changed.

Evaluate the role of the prefrontal cortex in visual awareness at the brain network level

There is growing evidence showing that conscious experience may be represented at the brain
network level [17,21]. Thanks to the simultaneously recorded LFP data in multiple brain regions with
a high signal-to-noise ratio and high spatial-temporal resolution, we can analyze the dynamic
functional connectivity changes under the aware and unaware conditions at the millisecond scale.
Since the signals in the low-frequency band are more involved in the information interaction
between different brain regions and the activities in the high-frequency band are more representative
of the local neural activities, it is easy to understand the functional connectivity changes in the low-
frequency band between relatively distant brain regions. Some previous studies have found that the
phase synchronization of neural activities in beta and gamma bands is related to visual awareness,
but because most of these studies are based on fMRI, it is difficult to describe the dynamic changes
in such brain networks on a more precise time scale. The only human iEEG study [30] reported that
the phase synchronization of the beta band in the aware condition also occurred relatively late (>300
ms). Moreover, most of these studies did not rule out the influence of report-related cognitive
processing, and these advanced cognitive processing processes are often associated with changes in
the functional connectivity of multiple brain regions. To the best of our knowledge, we are the first
to report dynamic changes in functional connectivity on the millisecond time scale in the low-
frequency band (1-8 Hz) associated with visual awareness, and this change is not caused by
reported-related confounding factors, such as motion preparation. Moreover, the early sites of PLV
and HG overlapped each other to a large extent. Such converging results may further suggest that
the middle lateral prefrontal cortex, the brain regions where these sites are located, plays a crucial
role in the emergence of visual awareness.

Our results may contribute to some theoretical concepts related to consciousness.

Compared with the unaware trials, the high-frequency activity in the PFC and the functional
connectivity between many brain regions in the aware trials increased. These results may confirm
some theoretical concepts, such as 'ignition' and conscious access. Global neuronal workspace
theory (GNWT) emphasizes ignition and conscious access [5,8]. Ignition means nonlinear brain
activation to small stimulus changes (such as perceptual threshold stimulation). For the near-
threshold stimulus used in the current study, while the sensory information in the brain passes
through a certain threshold in a specific brain area, such as the lateral prefrontal cortex, the
population of neurons discharges (HG activity of local field potential), that is, "ignition". This is
consistent with many previous human intracranial studies [7,32]. However, due to clinical limitations,
these studies are largely confined to the temporal lobe. To the best of our knowledge, our study
detected 'ignition' in the human PFC for the first time.

Conscious access means that conscious contents can be shared by many other local information
processors in the brain [5,8]. For example, when you see the grating, you can either report it orally or
draw it down. Such a function may require the participation and coordination of many related brain
regions or networks, but there is no clear biomarker for how these brain regions coordinate. The
explicit increase in the PLV, specifically appearing in the aware trials, in the low-frequency band
found in the current study may provide potential biological meaning for conscious access; that is,
many brain regions interact with each other through phase synchronization in the low-frequency
band to accomplish information sharing of conscious content.

Interestingly, the beginning time of prefrontal visual awareness-related activities found in ERP
and HG activities is very consistent with the onset time of dynamic functional connectivity
enhancement, both of which start at 200-300 ms. In addition, the dynamic functional connectivity

of the sites with early HG visual awareness-related activity is usually the earliest to enhance compared to other sites, and the two have good consistency. And this consistency suggested that conscious access and phenomenal awareness may be closely coupled, occurring initially in the lateral PFC at the early stage , and indicate a potential process of conscious perception: subtle change in the external stimulus or internal brain state - population neuron discharge in specific brain areas (ignition, phenomenal consciousness appeared) - phase synchronization of various brain areas induced by specific brain areas (conscious access). Further research is needed to regulate the activity of the middle lateral prefrontal cortex in the high-gamma band or the phase synchronization in the low-frequency band to explore whether the prefrontal cortex plays a causal role in the process of conscious perception.

Methods

Data acquisition

Six adult patients with drug-resistant epilepsy (6 males, 32.33±4.75 years old, mean±SEM) and 10 healthy subjects (4 males, 6 females, age 27.80 ± 3.92, mean ± SEM) participated in this study. All subjects had normal or corrected-to-normal vision. Electrophysiological signals (LFPs) were obtained from 6 patients. Stereotaxic EEG depth electrodes (SINOVATION MEDICAL TECHNOLOGY CO., LTD. Ltd., Beijing, China) containing 8 to 20 sites were implanted in patients in the Department of Neurosurgery, PLA General Hospital. Each site was 0.8 mm in diameter and 2 mm in length, with 1.5 mm spacing between adjacent sites. A few electrodes were segmented, and the distance between the two segments was 10 mm. Electrode placement was based on clinical requirements only. Recordings were referenced to a site in white matter. The sEEG signal was sampled at a rate of 1 kHz, filtered between 1 and 250 Hz, and notched at 50 Hz (NEURACLE Technology Co., Ltd., Beijing, China). Stimulus-triggered electrical pulses were recorded simultaneously with sEEG signals to precisely synchronize the stimulus with electrophysiological signals.

Ten healthy subjects performed the experiment in a quiet dark room. Stimuli were presented on a 27-inch LED screen (BenQ, refresh rate of 120 Hz, resolution of 1920 x 1080), and eye movement data were recorded by an infrared eye tracker (EyeLink 1000 Desktop; SR Research) with a sampling rate of 1 kHz. Patients performed the experiment in a quiet dark environment. The stimuli were presented on a 24-inch LED screen (Admiral Overseas Corporation, refresh rate of 144 Hz, resolution of 1920 x 1080), and eye position data were obtained by an infrared eye tracker (Jsmz EM2000C, Beijing Jasmine Science & Technology, Co. Ltd), sampling at 1 kHz. The experimental paradigm was presented by MATLAB (The MathWorks) and the toolbox Psychtoolbox-3 (PTB-3; Brainard and Pelli). Patients were regular recipients of standard medication for epilepsy treatment during the experimental period.

All subjects provided informed consent to participate in this study. The Ethics Committee of Beijing Normal University approved the experimental procedures.

Electrode localization

Electrode locations were determined by registering postoperative computed tomography (CT) scans and preoperative T1 MRI using a standardized mutual information method in SPM12 [35]. Cortical reconstruction and parcellation (Desikan-Killiany Atlas) were conducted by FreeSurfer [24].

619 To demonstrate the MNI coordinates of recording sites, a nonlinear surface registration was
620 performed using a nonlinear registration in SPM12, aligning to the MNI ICBM152 template.

Experimental task

A trial started when a fixation point (0.5°x0.5°, white cross) appeared at the center of the screen
(radius of eye position check window is 4°, the dotted circle). After the subject fixated on the fixation
point for 600 ms, a cue stimulus (Gabor grating, 2x2° circle) was presented for 50 ms at a fixed
position (7°) on the left (or right) side of the screen for all participants. The side where the grating
appears (left or right) is opposite to the hemisphere where the patients' electrodes are implanted (left
or right). For example, if the patient's electrodes are implanted on the right side, the grating would
appear on the left side of the screen. For patients 1, 5, and 6, the electrodes were implanted in both
hemispheres, and the grating is set on the right side. In 70% of the trials, the grating contrast (weber
contrast, see [10]) was maintained near the subject's perceptual threshold by a 1 up/1 down staircase
method, and the step was either 0.39% or 0.78% for each participant. In 10% of the trials, the
stimulus contrast was well above the threshold, and in the other 20% of the trials, the stimulus
contrast was 0, namely, no stimulus appeared. After another 600 ms delay, the color of the fixation
point turned red or green, and two saccade targets (1x1°, white square) appeared at fixed positions
(10°) on the left and right sides of the screen. If the grating was seen, the green fixation point was
required to make a saccade to the right target, while the red fixation point was required to make a
saccade to the left target. If the grating was not seen, the rule of saccadic direction was inverted.
Gratings with high contrast (well above the perceptual threshold) and zero contrast (grating absent)
served as control conditions to evaluate the understanding and performance of the task. Before
starting to collect data, subjects accepted 1-2 training sessions, and the contrast perceptual threshold
was determined for each subject. In the formal experiments, each session consisted of 180 trials,
and the intertrial interval (ITI) was 800 ms.

Data analysis

Each healthy subject completed 2 sessions, and each patient completed 5-7 sessions. Each
session contained 180 trials in total. We excluded trials in which subjects broke fixation during the
fixation period (eye position exceeded 4x4° check window by more than 40 ms) or the saccadic
latency exceeded max response duration (2000 ms for most patients, 5000 ms for patient 6 according
to the patient's performance) after the target appeared. For all subjects, more than 85% of the trials
were available for further analysis.

In each session, according to the subjects' reports of being aware and unaware of the existence
of grating, we divided the trials into awareness and unawareness states. According to the percentage
of subjects reporting 'awareness', we further divided grating contrast into 3 levels: high contrast
(aware percentage > 75%), near-threshold contrast (25% <= aware percentage <= 75%), and low
contrast (aware ratio<25%). Due to the few trials of low-contrast-aware and high-contrast-unaware
trials, we next analyzed only trials under four conditions of awareness: low-contrast-unaware (LU),
near-threshold-unaware (NU), near-threshold-aware (NA) and high contrast-awareness (HA). In
total, under LU/NU/NA/HA conditions, there were 243/353/343/160 trials (22.1/32.1/31.2/15.6%,
averaged across patients). Since we were more concerned with the differences in subjects' behavior
and electrophysiological responses during the physical stimulus, we focused on the results under
NA and NU conditions in the subsequent analysis. The results under LU and HA conditions were
classified as the control group and were only used to verify and check the results during calculation.

Behavioral data analysis

Psychophysical curve fitting uses the following formula:

664
$$AP = a + \frac{b}{1 + exp(-d * (SC - c))}$$

AP, ratio of reported awareness; SC, stimulus contrast; a-d, individual fitting parameters;

For population psychometric curve fitting, adjacent contrast levels were averaged, resulting in
8 contrast levels in each session.

The saccadic latency was calculated according to a custom program. Eye position and saccade
velocity were smoothed with 10 ms sliding windows, and each step was 1 ms. Saccade onset time
was calculated as the time when the saccade velocity first exceeded 30°/s and lasted more than 20
ms. Saccadic latency was defined as the time interval between the onset of the saccadic target and
the onset of the saccade.

LFP data analysis

Data preprocessing

Data preprocessing was performed by the software package EEGLAB [36] in MATLAB. Data
were filtered between 1-250 Hz and notch-filtered at 50, 100, 150, 200, and 250 Hz with an FIR
filter. Epochs were extracted from -490 ms before grating onset to 1299 ms after grating onset. Bad
recording sites were discarded for analysis based on visual inspection and power spectral density
(PSD) analysis. each recording site was applied a bipolar reference, i.e., rereferenced to its direct
neighbor.

Spectral analyses

Time-frequency decomposition in high-gamma bands (60-150 Hz) was achieved by Morlet
wavelets in the Brainstorm toolbox [37]. Before computation, the mean ERP in each condition was
removed. For visualization, we used a prestimulus baseline correction over -200 to 0 ms, and the
results in Figure 4 show the HG activity increases or decreases relative to this baseline. To avoid
windowing artifacts, we only reported results of -200 to 800 ms.

Quantitative definition of visual awareness-related activity in recording sites

We first identified visual awareness-related sites. For all recording sites, we compared the
broadband amplitude (for ERP analysis) or high-gamma magnitude (for HG analysis) of LFP
between the NA and NU conditions within 650 ms after grating onset by statistical tests for each
time point. We defined the site where there was a significant difference (p < 0.01 FDR corrected for
time points and channels, independent t-test) in the LFP between NA and NU conditions and
persisted for more than 20 ms as an awareness-related site. We also defined the start time of
divergence as the divergence onset time (DOT) and defined the mean of the LFP amplitude
difference in the first divergence period as the divergence amplitude (DA).

Normalized DOT:

697
$$T_d = T_{target} - DOT$$

698
$$normDOT = \frac{T_d - T_{dmin}}{T_{dmax} - T_{dmin}}$$

DOT, divergence onset time; T_{target}, time from target onset to grating onset, 650 ms; T_d, time from
DOT to T_{target}; T_{dmin}, the minimum value of T_d in all awareness-related sites within one patient; T_{dmax},

the maximum value of T_d of all awareness-related sites within one patient; normDOT, normalized

DOT value, a larger normDOT value means a shorter DOT;

For population analysis of recording sites, we conducted baseline normalization for each site.

Baseline normalization for ERP amplitude:

707
$$x_{norm} = \frac{x - \mu}{\mu} * 100$$

708   x_{norm}, normalized value; x, LFP amplitude; μ, mean LFP amplitude in the baseline period (-200 to

709   0 ms).

711   Baseline normalization for HG magnitude:

712
$$x_{norm} = \frac{x - \mu}{\sigma} * 100$$

713   x_{norm}, normalized value; x, high-gamma magnitude or power; μ, mean of high-gamma magnitude

714   or power in the baseline period (-200 to 0 ms); σ, standard error of high-gamma magnitude or

715   power in the baseline period (-200 to 0 ms).

717   Baseline normalization for spectrogram:

718
$$x_{norm} = 10 * log10(x/\mu)$$

x_{norm}, normalized value; x, power value; μ, mean power in the baseline period (-200 to 0 ms).

Topographic display of ERP and HG data

To visualize the topography of the ERP and HG activities, we used the Brainstorm [37] package

of MATLAB to map the results on the ICBM152 cortical surface. It is a simple 3D interpolation

with Shepard's method (weights decreasing with the square of the distance), similar to what is done

when projecting subject sources to an anatomy template, where vertices of the cortical surface that

are further than 15 mm from the center of the sEEG contact are ignored.

Decoding Analysis

We implemented a machine learning framework for trial-by-trial classification using broadband

amplitude and high-gamma magnitude of LFP activities in the PFC via the MVPA-Light toolbox

[38]. We applied the LDA classification and temporal generalization on each participant separately,

and all trials from each class were undersampled to balance the unequal trial number between

different conditions. Then, the trials were randomly assigned into 10 folds, and all trials within

each fold were subaveraged across every 4 trials to increase the signal-to-noise ratio. Decoding

was then followed with a leave-one-out cross-validation procedure on the subaveraged trials. This

procedure was repeated 10 times so that each fold in the dataset was used exactly once for testing,

and at least once for training, but never at the same time. And the fold assignment was repeated 5

times, which resulted in total 50 iterations for the decoding analysis of each patient.

Functional Connectivity Analysis

The time-across phase-locking value (PLV) was calculated in the brainstorm toolbox. Time-

frequency transformation was conducted via the Hilbert transform method. The time-across PLV

measures if the phase difference between two signals is consistent across trials with respect to the

event that defines the trials (i.e., the output is a mean across trials per time point)[39]. PLV values range between 0 (for totally random) and 1 (for perfect phase locking). For clear demonstration here, the mean PLV in the baseline period (-200 to 0 ms) of each pair of recording sites was subtracted. The group analysis of PLV was performed according to [40,41]. The topographic display of the connectivity was conducted by the BrainNet Viewer toolbox [42].

Data and software availability

Electrophysiological data were analyzed using MATLAB in conjunction with the toolboxes mentioned above. Data and customized MATLAB analysis scripts are available upon reasonable request from Ming-sha Zhang (mingsha.zhang@bnu.edu.cn).

ACKNOWLEDGMENTS

We thank the participants for volunteering to take part in the study. This study was funded by STI2030-Major Projects 2021ZD0204300 and National Natural Science Foundation of China (32061143004, 32030045 and 31871078).

Reference

1. Koch, C., Massimini, M., Boly, M., and Tononi, G. (2016). Neural correlates of consciousness: progress and problems. Nat Rev Neurosci *17*, 307-321. 10.1038/nrn.2016.22.

2. Mashour, G.A. (2018). The controversial correlates of consciousness. Science *360*, 493-494. 10.1126/science.aat5616.

3. Dehaene, S., Lau, H., and Kouider, S. (2017). What is consciousness, and could machines have it? Science *358*, 486-492. 10.1126/science.aan8871.

4. Kim, C.Y., and Blake, R. (2005). Psychophysical magic: rendering the visible 'invisible'. Trends Cogn Sci *9*, 381-388. 10.1016/j.tics.2005.06.012.

5. Dehaene, S., and Changeux, J.P. (2011). Experimental and theoretical approaches to conscious processing. Neuron *70*, 200-227. 10.1016/j.neuron.2011.03.018.

6. Dehaene, S., Naccache, L., Cohen, L., Bihan, D.L., Mangin, J.F., Poline, J.B., and Riviere, D. (2001). Cerebral mechanisms of word masking and unconscious repetition priming. Nat Neurosci *4*, 752-758. 10.1038/89551.

7. Fisch, L., Privman, E., Ramot, M., Harel, M., Nir, Y., Kipervasser, S., Andelman, F., Neufeld, M.Y., Kramer, U., Fried, I., and Malach, R. (2009). Neural "ignition": enhanced activation linked to perceptual awareness in human ventral stream visual cortex. Neuron *64*, 562-574. 10.1016/j.neuron.2009.11.001.

8. Mashour, G.A., Roelfsema, P., Changeux, J.P., and Dehaene, S. (2020). Conscious Processing and the Global Neuronal Workspace Hypothesis. Neuron *105*, 776-798. 10.1016/j.neuron.2020.01.026.

9. Brown, R., Lau, H., and LeDoux, J.E. (2019). Understanding the Higher-Order Approach to Consciousness. Trends Cogn Sci *23*, 754-768. 10.1016/j.tics.2019.06.009.

10. van Vugt, B., Dagnino, B., Vartak, D., Safaai, H., Panzeri, S., Dehaene, S., and Roelfsema, P.R. (2018). The threshold for conscious report: Signal loss and response bias in

visual and frontal cortex. Science *360*, 537-542.

11.   Koch, C. (2018). What Is Consciousness? Nature *557*, S8-S12. 10.1038/d41586-018-05097-x.

12.   Lamme, V.A.F. (2018). Challenges for theories of consciousness: seeing or knowing, the missing ingredient and how to deal with panpsychism. Philos Trans R Soc Lond B Biol Sci *373*. 10.1098/rstb.2017.0344.

13.   Dembski, C., Koch, C., and Pitts, M. (2021). Perceptual awareness negativity: a physiological correlate of sensory consciousness. Trends in Cognitive Sciences *25*, 660-670. 10.1016/j.tics.2021.05.009.

14.   Seth, A.K., and Bayne, T. (2022). Theories of consciousness. Nat Rev Neurosci *23*, 439-452. 10.1038/s41583-022-00587-4.

15.   Frassle, S., Sommer, J., Jansen, A., Naber, M., and Einhauser, W. (2014). Binocular rivalry: frontal activity relates to introspection and action but not to perception. J Neurosci *34*, 1738-1747. 10.1523/JNEUROSCI.4403-13.2014.

16.   Tsuchiya, N., Wilke, M., Frassle, S., and Lamme, V.A.F. (2015). No-Report Paradigms: Extracting the True Neural Correlates of Consciousness. Trends Cogn Sci *19*, 757-770. 10.1016/j.tics.2015.10.002.

17.   Kronemer, S.I., Aksen, M., Ding, J.Z., Ryu, J.H., Xin, Q.L., Ding, Z.X., Prince, J.S., Kwon, H., Khalaf, A., Forman, S., et al. (2022). Human visual consciousness involves large scale cortical and subcortical networks independent of task report and eye movement activity. Nat Commun *13*. ARTN 7342

10.1038/s41467-022-35117-4.

18.   Northoff, G., and Lamme, V. (2020). Neural signs and mechanisms of consciousness: Is there a potential convergence of theories of consciousness in sight? Neurosci Biobehav Rev *118*, 568-587. 10.1016/j.neubiorev.2020.07.019.

19.   Koivisto, M., and Revonsuo, A. (2010). Event-related brain potential correlates of visual awareness. Neurosci Biobehav Rev *34*, 922-934. 10.1016/j.neubiorev.2009.12.002.

20.   Koivisto, M., Lahteenmaki, M., Sorensen, T.A., Vangkilde, S., Overgaard, M., and Revonsuo, A. (2008). The earliest electrophysiological correlate of visual awareness? Brain Cogn *66*, 91-103. 10.1016/j.bandc.2007.05.010.

21.   Huang, Z., Mashour, G.A., and Hudetz, A.G. (2023). Functional geometry of the cortex encodes dimensions of consciousness. Nat Commun *14*, 72. 10.1038/s41467-022-35764-7.

22.   Merten, K., and Nieder, A. (2012). Active encoding of decisions about stimulus absence in primate prefrontal cortex neurons. P Natl Acad Sci USA *109*, 6289-6294.

23.   Tass, P., Rosenblum, M.G., Weule, J., Kurths, J., Pikovsky, A., Volkmann, J., Schnitzler, A., and Freund, H.J. (1998). Detection of $\mathit{n}\!:\!\mathit{m}$ Phase Locking from Noisy Data: Application to Magnetoencephalography. Physical Review Letters *81*, 3291-3294. 10.1103/PhysRevLett.81.3291.

24.   Fischl,   B.   (2012).   FreeSurfer.   Neuroimage   *62*,   774-781. 10.1016/j.neuroimage.2012.01.021.

25.   Allik, J., and Kreegipuu, K. (1998). Multiple visual latency. Psychol. Sci. *9*, 135-138. 10.1111/1467-9280.00025.

26.   Deplancke, A., Madelain, L., Chauvin, A., Cardoso-Leite, P., Gorea, A., and Coello,

Y. (2010). Influence of near threshold visual distractors on perceptual detection and reaching movements. J Neurophysiol *104*, 2249-2256. 10.1152/jn.01123.2009.

27. Posner, M.I., and Petersen, S.E. (1990). The Attention System of the Human Brain. Annual Review of Neuroscience *13*, 25-42. DOI 10.1146/annurev.ne.13.030190.000325.

28. Corbetta, M., and Shulman, G.L. (2002). Control of goal-directed and stimulus-driven attention in the brain. Nat Rev Neurosci *3*, 201-215. 10.1038/nrn755.

29. Posner, M.I. (1980). Orienting of Attention. Q J Exp Psychol *32*, 3-25. Doi 10.1080/00335558008248231.

30. Gaillard, R., Dehaene, S., Adam, C., Clemenceau, S., Hasboun, D., Baulac, M., Cohen, L., and Naccache, L. (2009). Converging intracranial markers of conscious access. PLoS Biol *7*, e61. 10.1371/journal.pbio.1000061.

31. Kapoor, V., Dwarakanath, A., Safavi, S., Werner, J., Besserve, M., Panagiotaropoulos, T.I., and Logothetis, N.K. (2022). Decoding internally generated transitions of conscious contents in the prefrontal cortex without subjective reports. Nat Commun *13*. ARTN 1535

10.1038/s41467-022-28897-2.

32. Reber, T.P., Faber, J., Niediek, J., Bostrom, J., Elger, C.E., and Mormann, F. (2017). Single-Neuron Correlates of Conscious Perception in the Human Medial Temporal Lobe. Curr Biol *27*, 2991-+. 10.1016/j.cub.2017.08.025.

33. Grootswagers, T., Wardle, S.G., and Carlson, T.A. (2017). Decoding Dynamic Brain Patterns from Evoked Responses: A Tutorial on Multivariate Pattern Analysis Applied to Time Series Neuroimaging Data. J Cogn Neurosci *29*, 677-697. 10.1162/jocn_a_01068.

34. King, J.R., and Dehaene, S. (2014). Characterizing the dynamics of mental representations: the temporal generalization method. Trends Cogn Sci *18*, 203-210. 10.1016/j.tics.2014.01.002.

35. Ashburner, J., and Friston, K. (1997). Spatial transformation of images. spm course notes.

36. Delorme, A., and Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics. Journal of Neuroscience Methods *134*, 9-21.

37. Tadel, F., Baillet, S., Mosher, J.C., Pantazis, D., and Leahy, R.M. (2011). Brainstorm: a user-friendly application for MEG/EEG analysis. Comput Intell Neurosci *2011*, 879716. 10.1155/2011/879716.

38. Guggenmos, M., Sterzer, P., and Cichy, R.M. (2018). Multivariate pattern analysis for MEG: A comparison of dissimilarity measures. Neuroimage *173*, 434-447. 10.1016/j.neuroimage.2018.02.044.

39. Mercier, M.R., Dubarry, A.S., Tadel, F., Avanzini, P., Axmacher, N., Cellier, D., Vecchio, M.D., Hamilton, L.S., Hermes, D., Kahana, M.J., et al. (2022). Advances in human intracranial electroencephalography research, guidelines and good practices. Neuroimage *260*, 119438. 10.1016/j.neuroimage.2022.119438.

40. Betzel, R.F., Medaglia, J.D., Kahn, A.E., Soffer, J., Schonhaut, D.R., and Bassett, D.S. (2019). Structural, geometric and genetic factors predict interregional brain connectivity patterns probed by electrocorticography. Nat Biomed Eng *3*, 902-916. 10.1038/s41551-019-0404-5.

41. Sadaghiani, S., Brookes, M.J., and Baillet, S. (2022). Connectomics of human

868 electrophysiology. Neuroimage *247*. ARTN 118788

869 10.1016/j.neuroimage.2021.118788.

870 42. Xia, M., Wang, J., and He, Y. (2013). BrainNet Viewer: a network visualization tool

871 for human brain connectomics. PLoS One *8*, e68910. 10.1371/journal.pone.0068910.

Supplementary Information

SI Figures and Videos

C

D

Figure S1. Functional connectivity analysis results of Patient 3-6.

A-D. Phase-Locking Value (PLV) changes at sensor level (NxN) in four patients P3-P6. From left to right, the

changes of PLV between each recording site in the five time points of 0/150/300/450/600ms after the appearance of

the grating in the near-threshold aware (upper) and near-threshold unaware (lower) trials are displayed respectively

(baseline removed, see the Method). The color represents PLV.

Video S1. The spatial-temporal dynamics of visual awareness-related activities (event-related potential (ERP))

in all recording sites during the 180-650 ms period (for a clearer display, projected to the cortex). The curves in the

upper left panel represent the significant (p<0.01 corrected, see Methods) difference (normalized) in averaged

amplitude of local field potential in NA and NU trials at all visual awareness-related ERP sites. The brain image

showed the difference projected on the surface at a specific time point after the grating appeared (top, left and right

views were showed at upper right, lower left and lower right panels, respectively). The color represents the

normalized voltage difference (see Methods).

Video S2. The spatial-temporal dynamics of visual awareness-related activities (high-gamma (HG) activity) in

all recording sites during the 180-650 ms period (for a clearer display, projected to the cortex). The same as Video

S1 but for high-gamma activity. The curves in the upper left panel represent the significant (p<0.01 corrected, see

Methods) difference (normalized) in averaged magnitude of high-gamma activity in NA and NU trials at all visual

awareness-related HG sites. The rain image showed the difference projected on the surface at a specific time point
after the grating appeared (top, left and right views were showed at upper right, lower left and lower right panels,
respectively). The color represents the normalized magnitude difference (see Methods).
901
902