Graphical Abstract

Parthenogenesis, sexual conflict and the evolution of oogamy

Xiaoyuan Liu, Jon W. Pitchford, George W.A. Constable

Highlights

Parthenogenesis, sexual conflict and the evolution of oogamy

Xiaoyuan Liu, Jon W. Pitchford, George W.A. Constable

- Adapting the classic Parker-Baker-Smith model for the evolution of anisogamy (gametic sex cells of differing sizes), we provide the first analysis of the co-evolution of fertilization rate and gamete cell mass in organisms that can reproduce both sexually and asexually through parthenogenesis.

- We identify a novel mechanism by which oogamy (motile microgametes and sessile macrogametes) can be selected for, arising from sexual conflict between microgametes (sperm) and macrogametes (eggs).

- We identify a novel mechanism by which a state of isogamy (equal sized gametes) can be stabilized under evolution, resulting from a bet-hedging strategy in switching environments.

- For all of the above insights, we develop a mathematical analysis with predictions that are supported by detailed computational simulations.

Parthenogenesis, sexual conflict and the evolution of oogamy

Xiaoyuan Liu[a], Jon W. Pitchford[a,b], George W.A. Constable[a]

[a]*Department of Mathematics, University of York, York, YO10 5DD, Yorkshire, U.K.*
[b]*Department of Biology, University of York, York, YO10 5DD, Yorkshire, U.K.*

Abstract

Organisms with external fertilization exhibit a variety of reproductive modes, from simple parthenogenesis to isogamy, anisogamy and oogamy. Here, we develop a mathematical model that helps to explain the evolution of these modes through the co-evolution of cell size and fertilization rate. By assuming that gametes can develop parthenogenetically should they fail to fertilize, and that survival of a propagule (zygote/unfertilized gamete) depends on size, we find that an isogamous population can evolve to anisogamy through evolutionary branching. Oogamy can then evolve from an anisogamous population under sexual conflict. Furthermore, we derive analytic results on the model parameters required to arrest evolution on this isogamy-oogamy trajectory. Low fertilization rates stabilise isogamy, while low fertilization costs stabilise anisogamy. Additionally we show using adaptive dynamics that isogamy can be maintained as a bet-hedging strategy in a stochastically switching environment.

Keywords:

Anisogamy, Oogamy, Fertilization Kinetics, Adaptive Dynamics, Evolutionary Branching, Switching environments
2000 MSC: 92D15, 92D25

1. Introduction

The origin of the sexes lies in the types of sex cells they produce; males produce small microgametes while females produce larger macrogametes [1]. This gamete size dimorphism is referred to as a state of anisogamy [2]. Should the microgamete be motile (sperm) and the macrogamete sessile (egg), the population is said to be oogamous [3]. While anisogamy is the most commonly observed mode of sexual reproduction in eukaryotic organisms [4], having evolved several times in evolutionary history [5], it is now widely accepted to be derived from isogamy (equal gamete sizes) [6]. While rarer, isogamous species include study organisms such as the yeast *Saccharomyces cerevisiae* [7] and the green algae *Chlamydomonas reinhardtii* [8], where self-incompatible mating types play the role of ancestral versions of the true sexes [9]. Indeed, the volvocine algae (of which *C. reinhardtii* is a member), provide neat empirical examples of these transitions, with phylogenetic analysis indicating that numerous independent lineages have undergone the transition from isogamy, to anisogamy, and finally to oogamy [3, 10]. Explaining the evolutionary mechanisms behind these transitions has been the focus of much work in evolutionary theory.

While theoretical investigations of the evolution of anisogamy date back to the 1930's [11] and were developed into the 1960's [12, 13], it was arguably the Parker-Baker-Smith (PBS) model [14] that synthesised these ideas into a complete evolutionary model that is now widely accepted as providing an explanation for the evolution of anisogamy [15, 16, 17]. They assumed that a fixed mass or energy budget is allocated by individuals to gamete production, such that microgametes can be produced in larger quantities than macrogametes. However, while microgametes may be more numerous, they contribute a lower fitness than macrogametes to a fertilized zygote due to their low cytoplasmic volume [18]. In this way the PBS model was able to show how anisogamy was the result of a quality-quantity trade-off, with disruptive selection acting on gamete size [15].

The key elements of the PBS model have since been set in a game theoretic [19] and population genetic [20, 21] context, as well as extended to account for more general reproductive modes such as hermaphrodism [22]. Models using adaptive dynamics [23] in particular have been useful. It has been shown analytically that even in the absence of mating types, anisogamy can evolve from isogamy through evolutionary branching in mass [24]. Meanwhile, ac-

counting for self-incompatibility of mating types and investigating the effect of varying fertilization rate, [25] showed that anisogamy can evolve from isogamy through both gamete competition and gamete limitation. Altogether, the results described above suggest that fertilization rate is a crucial factor that may impact gamete survival, and selection is likely to act on the fertilization rate between microgametes and macrogametes.

As addressed, oogamy, the loss of motility in eggs and specialization for motility in sperm, is often seen as the "last step in the evolution of the egg–sperm dichotomy" [26], a view supported by empirical analyses that suggest oogamy is derived from anisogamy [27]. However the theory of this transition is comparatively less studied than the earlier transition from isogamy to anisogamy. Ghiselin [28] provided an argument based on the physiological division of labour between macro and microgametes, with females specialising in provisioning and males in motility. Most other work has considered the evolution of oogamy as a strategy to maximise gamete encounter rate. This can be achieved by having a population of pheromone emitters and receivers [29, 30] and by having a large stationary target egg and small motile sperm [31, 32]. Although the assumption of an inverse speed-size relationship in [32] has justification in some gametic systems, it is also worth noting that positive speed-size relationship have been observed in *C. reinhardtii*, due to larger cells having greater propulsive forces; at scales such as these the precise speed-size relationship is complicated by details of cell morphology [33]. Lastly, the prior evolution of internal fertilization has been proposed as a mechanism that could generate selection for oogamy [34], consistent with empirical evidence from *volvox* (external fertilization and anisogamous) and its sister lineage *platydorina* (internal fertilization and oogamous) [35].

The work described above all assumes obligate sexual reproduction (unfertilized gametes die at the end of each generation) and a fixed fertilization rate (the rate at which gametes encounter one another and fertilize to form zygotes) [19, 25, 26]. Recently, however, inspired by the life-histories of green and brown algae such as *Blidingia minima* (isogamous [36]), *Urospora neglecta* (anisogamous [37]), and *Saccharina japonica* (oogamous [38]), whose gametes can develop parthenogenetically should they fail to find a mate (see [36, 37, 39], respectively), the first of these assumptions was relaxed in two theoretical papers [40, 41]. In [40], extra survival costs were incurred by gametes developing parthenogenetically, while in [41] extra survival costs on either the parthenogenetic or the sexual reproductive route were considered. They found that in the presence of two self-incompatible mating types, isogamy can be stabilised under low costs to parthenogenesis, while anisogamy is the evolutionary stable state when fertilization is favoured.

The studies described above on parthenogenesis and the evolution of anisogamy lead to a natural question; if the sexual route (via fertilization) and asexual route (via parthenogenesis) carry different survival costs, how should the fertilization rate evolve to account for this? Should this rate increase (to minimise the number of unfertilized gametes taking a potentially perilous route to survival) or should it decrease (to avoid potential costs incurred during cell fusion)? The modelling of fertilization kinetics is an interesting topic in its own right [42, 43], and plays a role in models for the evolution of anisogamy from isogamy [25]. However fertilization kinetics are also clearly a key element of the evolution of oogamy, with selection on sperm to increase their encounter rate with eggs and selection on eggs to remain sessile.

In this paper, we modify the adaptive dynamics models of [24, 40, 41] to study the co-evolution of gamete size and fertilization rate in species capable of parthenogenesis under external fertilization. For simplicity, we assume an absence of self-incompatible mating types [6, 24]. We show that under such assumptions, anisogamy can evolve from an initial state of isogamy, followed by the subsequent evolution to oogamy under sexual conflict between microgametes and macrogametes. In Section 2, we introduce the model, describing its key behaviour in Section 3. These behaviours include the evolution of oogamy (Section 3.2), conditions that stabilise isogamy (Section 3.3), conditions that stabilise anisogamy (Section 3.4), and the emergence of isogamy as a bet-hedging strategy in switching environments (Section 3.5).

2. Model

In this section we describe the specifics of the models we use, paying careful attention to the various time scales involved. We begin by considering the evolutionary model in a fixed environment (Section 2.1), before generalising to the case of a switching environment under which bet-hedging strategies can evolve (Section 2.2).

2.1. Model dynamics in a fixed environment

The evolutionary dynamics of the model are built from a hierarchy of timescales, which are particularly important to keep in mind once environmental

switching is introduced in later sections. The shortest timescale is the generational timescale. The intermediate timescale is that over which the invasion of a rare mutant (taking place over many generations) can occur. The longest timescale is the evolutionary timescale, representing the cumulative effect of multiple mutations and invasions.

Dynamics within each generation

At the start of each discrete generation, a number of adults with haploid mass (or energy budget) M produce gametes (mass m), such that mass/energy budget is conserved (i.e. each adult produces M/m gametes). Note that this implicitly assumes, for simplicity, that a continuous (rather than discrete) number of gametes is possible. Gametes then enter a pool and the fertilization process takes place. After a finite time window, the resultant cells face a round of survival dependent on their mass. The surviving cells form the basis of the next generation, completing the generational cycle, as illustrated in Figure 1.

Fertilization Kinetics For simplicity, we assume that all gametes may fertilize each other (i.e. there are no self-incompatible mating types). Given a total of A adults, the population is initially comprised of $N = AM/m$ single gametes produced by adults through meiosis. We assume fertilization is external, with cells fertilizing according to mass action dynamics at rate α, such that the number of single (unfertilized) cells, N, is given by the solution to

$$\frac{\mathrm{d}N}{\mathrm{d}t} = -\alpha N^2, \qquad N(0) = \frac{AM}{m} \qquad (1)$$

(see also [44]). At the end of the fertilization window or duration T there are then $N(T)$ single cells remaining, and $(N(0) - N(T))/2$ fertilized cells.

The parameter α is variously referred to as the fertilization rate, the collision rate [42], the "'aptitude' for union" [13], or the "bimolecular reaction constant" [45]. We will refer to α as the fertilization rate, and treat it as a trait subject to evolution. Overall, α captures the compound effect of the propensity for fertilization between cells encountering each other, as well as additional mechanisms to enhance cell encounter rate such as increased emission or response to cheomoattractants [30] or cell motility [46]. In practice there is a likely upper-bound on α, brought about by energy trade-offs, diffusion in aquatic environments, or dispersal in terrestrial environments. To account for this we introduce a ceiling on α, and restrict its evolutionary dynamics to the range $\alpha_{\max} \geq \alpha \geq 0$.

Figure 1: Schematic of dynamics within each generation. Mature cells (adults) produce gametes at the start of a generation. All the gametes are given a fixed time period T in which to complete the fertilization process. At the end of the fertilization period, there will be a pool of fertilized zygotes and unfertilized gametes, both of which are capable of maturation. Each cell survives according to its independent survival functions ($(1 - C_z)S(\beta, m_1 + m_2)$ and $(1 - C_p)S(\beta, m_i)$ respectively) to produce a number of mature cells in the subsequent generation. The pool of gametes consist of resident (blue) and mutant gametes (green), where the mutation occurs in either the mass m or fertilization rate α.

Survival Probability

At the end of a finite fertilization window at time T, the population will consist of both fertilized and unfertilized gametes. Fertilised gametes produce diploid zygotes, while unfertilized gametes can develop as haploids parthenogenetically (e.g. parthenosporophytes [47]). We assume that the probability that either of these cell types survives is given by the Vance survival function [18], which is a common assumption in the literature [19, 25, 48]. Given a cell size m_c (for either fertilized or unfertilized cells), the survival probability is given by

$$S(\beta, m_c) = \exp\left(-\frac{\beta}{m_c}\right). \qquad (2)$$

Note that this is an increasing function of cell size, and we do not account for gamete mortality during the fertilization period. Thus, although both fertilized and unfertilized gametes are exposed to the same survival function, fertilized cells (with a mass around twice the size of unfertilized cells in a monomorphic isogamous population) will have a greater survival probability than unfertilized cells. Meanwhile for a given mass m_c, increasing β will decrease the survival probability. We therefore refer to β as the resistance to survival, with high β corresponding to harsh environments in which survival is difficult, and low β corresponding to more benign environments in which even cells of modest mass have a high probability of surviving.

In addition to the benefits explained in the introduction, fertilization may also carry risks: generally these include cell-fusion failure [49], selfish extra-genomic elements in the cytoplasm [50] and cytoplasmic con-

flict [51, 52]. We therefore introduce an additional fixed cost, $1 \geq C_z \geq 0$, applied to fertilized zygotes independent of their mass. Similarly, there can be costs associated to parthenogenetic development such as reduced fitness due to lack of genetic diversity [53] and the possibility of failure of parthenogenetic development [40]. The final probability of survival for a zygote formed from the fertilization of two gametes of sizes m_1 and m_2 is then given by $(1 - C_z) \exp\left[-\beta/(m_1 + m_2)\right]$, while the probability of survival for an unfertilized cell of size m is $(1 - C_p) \exp\left[-\beta/m\right]$.

Invasion Dynamics

We assume that haploid gametes are characterised by two genetically determined non-recombining traits; their mass, m, and their fertilization rate, α. We next consider a monomorphic resident population to which a mutant individual is introduced at rate μ. This mutant may produce gametes of a different mass to its ancestor, $m \pm \delta m$, where δm represents the size of a mutational step. Under this scenario the mutant may produce more or fewer gametes than its ancestor (see Appendix A.1), but the survival probability of its unfertilized cells (mass $m \pm \delta m$), mutant-resident fertilized cells (mass $2m \pm \delta m$), and mutant-mutant fertilized cells (mass $2(m \pm \delta m)$), will also be simultaneously decreased or increased (see Eq. (2) and Appendix A.3). The cumulative effect of this quality-quantity trade-off will either lead to selection for or against the mutant over subsequent generations.

Alternatively, the mutant may engage in an increased or decreased fertilization rate relative to its ancestor, at a rate $\alpha \pm \delta\alpha$ (see Appendix A.2). Under this scenario the mutant fertilizes with residents at their average fertilization rate $(2\alpha \pm \delta\alpha)/2$ and other mutants at a rate $\alpha \pm \delta\alpha$. Mutants will either contribute to more or fewer fertilized cells and depending on the resistance to survival, β, the costs to fertilization, C_z, and the costs to parthenogenesis, C_p, may experience a selective advantage over the resident by devoting more of its gametes to one of the reproductive routes (see Appendix A.4).

In order to mathematically characterise the invasion dynamics (which occur over discrete generations), we assume that δm and $\delta\alpha$ are small, so that we can approximate the dynamics continuously. Denoting by $\hat{f}_m$ the frequency of mutants of size $m \pm \delta m$ in the population, and t_g the number of generations, we find (see Appendix B.1)

$$\frac{\mathrm{d}\hat{f}_m}{\mathrm{d}t_g} = h_m(m, \alpha, \beta, C_z, C_p)\hat{f}_m(1 - \hat{f}_m), \tag{3}$$

where $h_m(m, \alpha, \beta, C_z, C_p)$ is a constant selective pressure that depends on the parameters $m, \alpha, \beta, C_z, C_p$ (see Eq. (B.1)). Similarly, denoting by $\hat{f}_\alpha$ the frequency of mutants with fertilization rate $\alpha \pm \delta\alpha$ in the population (see Appendix B.2), we find

$$\frac{\mathrm{d}\hat{f}_\alpha}{\mathrm{d}t_g} = h_\alpha(m, \alpha, \beta, C_z, C_p)\hat{f}_\alpha(1 - \hat{f}_\alpha) \tag{4}$$

where $h_\alpha(m, \alpha, \beta, C_z, C_p)$ is a constant that depends on the parameters $m, \alpha, \beta, C_z, C_p$ (see Eq. (B.2)). We thus see that in the case of a single mutant, we have frequency-independent selection for both traits, with the direction of selection determined by the resistance to survival β, the cost to fertilization and parthenogenesis C_z and C_p, and the traits of the resident population, m and α. We assume for the remainder of this section that mutants encounter a strictly monomorphic resident population, allowing us infer a simple trait substitution process that considerably simplifies the evolutionary analysis. However we will return to the possibility of mutants coexisting with residents in Section 3.2 of the Results.

Evolutionary Dynamics

Following standard approaches in adaptive dynamics [54] (see also Appendix C), we construct the evolutionary equations for the gamete mass, m, and the fertilization rate, α. Denoting by τ the evolutionary timescale over which mutations appear and trait substitutions occur, we find

$$\begin{aligned}
\frac{\mathrm{d}m}{\mathrm{d}\tau} &= H_m(m, \alpha; \beta, C_z, C_p) \\
&= -\frac{4(1 - C_p)m(m - \beta) + AM\alpha T(1 - C_z)e^{\frac{\beta}{2m}}(4m - \beta)}{4m^2((1 - C_p)m + AM\alpha T(1 - C_z)e^{\frac{\beta}{2m}})} \\
\frac{\mathrm{d}\alpha}{\mathrm{d}\tau} &= H_\alpha(m, \alpha; \beta, C_z, C_p) \\
&= -m\frac{((1 - C_p) - (1 - C_z)e^{\frac{\beta}{2m}})\ln(1 + \frac{AM\alpha T}{m})}{2\alpha((1 - C_p)m + AM\alpha T(1 - C_z)e^{\frac{\beta}{2m}})}.
\end{aligned} \tag{5}$$

Note that these equations involve highly nonlinear functions, with some evident non-physical behaviour near the boundaries (e.g. when cell masses are zero). This feature is common in simplified models of anisogamy evolution [41].

2.2. Evolutionary dynamics in switching environments

We now wish to consider the case of a population characterised by gamete mass (m) and fertilization rate (α) traits, evolving subject to changing environmental

conditions. Explicitly, we allow the resistance to survival to alternate between two values β_1 and β_2; recall from Eq. (2) that if $\beta_1 > \beta_2$ then β_1 represents a comparatively "harsh environment", where cells have a lower survival probability than in environment β_2.

Switching between these two environments is modelled as a discrete stochastic telegraph process [55, 56]; the time spent in each environment is distributed geometrically (a discrete analogue of the exponential distribution), spending an average period $\tau_1 \approx 1/\lambda_{1\to2}$ in environment 1 and $\tau_2 \approx 1/\lambda_{2\to1}$ in environment 2, where $\lambda_{i\to j}$ is the transition rate from environment i to j. We must carefully consider the magnitude of these timescales in comparison with the other timescales at work in the model (see Section 2.1).

First consider the case where the environmental switching timescales, τ_1 and τ_2, are larger than the generational timescale (t_g), but much smaller than the invasion timescale (characterised by the inverse of the strength of selection, proportional to $1/\delta m$ and $1/\delta\alpha$) and the mutational timescale ($1/\mu$). We call this the 'fast relative to invasion' switching regime (FRTI). In this scenario, the population does not switch environments during a single round of fertilization kinetics, but typically switches between the two environmental states many times before an invasion has time to complete. When switching occurs this frequently, we can approximate the dynamics mathematically by observing that the population experiences the weighted average of the selection pressures in the two environments [55]. Denoting by P_1 and P_2 the probability of finding the population in either of the respective environments, we have

$$P_1 = \frac{\tau_1}{\tau_1 + \tau_2}, \quad P_2 = \frac{\tau_2}{\tau_1 + \tau_2} = (1 - P_1). \quad (6)$$

The effective selection pressure on mutants with mass $m + \delta m$ in a resident population of mass m during an invasion is then given by

$$\left[P_1 h_m(m, \alpha, \beta_1, C_z, C_p) + P_2 h_m(m, \alpha, \beta_2, C_z, C_p) \right] \quad (7)$$

which can be contrasted against the selection pressure $h_m(m, \alpha, \beta, C_z, C_p)$ in Eq. (3). An analogous approach allows us to approximate the invasion dynamics for mutants with a different fertilization rate to their ancestors in this FRTI regime (see Appendix F.1).

With equations for $\mathrm{d}\hat{f}_m/\mathrm{d}t_g$ and $\mathrm{d}\hat{f}_\alpha/\mathrm{d}t_g$ in hand, we can proceed to apply the same standard techniques from adaptive dynamics as were used to derive Eq. (5) from Eqs. (3-4) (see Appendix F). We obtain the effective evolutionary dynamics

$$\frac{\mathrm{d}m}{\mathrm{d}\tau} = P_1 H_m(m, \alpha; \beta_1, C_z, C_p) + P_2 H_m(m, \alpha; \beta_2, C_z, C_p)$$
$$\frac{\mathrm{d}\alpha}{\mathrm{d}\tau} = P_1 H_\alpha(m, \alpha; \beta_1, C_z, C_p) + P_2 H_\alpha(m, \alpha; \beta_2, C_z, C_p)$$
$$(8)$$

where $H_m(m, \alpha; \beta_1, C_z, C_p)$ and $H_\alpha(m, \alpha; \beta_1, C_z, C_p)$ retain the functional forms given in Eq. (5), and P_1 and P_2 are taken from Eq. (6).

In contrast to the FRTI regime discussed above, we can also investigate the regime in which environmental switching occurs on a comparable or slower timescale than invasion, but still occurs fast relative to the evolutionary timescale. In this scenario, which we call the "fast relative to evolution" switching regime (FRTE), environmental switching occurs on a similar rate to that at which new mutations are introduced, but much faster than the combined effect of mutation and selection (e.g. $\lambda_{i\to j} \ll \mu \times \delta m$), such that only a small number of mutations can fixate in either environment before the population switches to the alternate environment. Although we shall show via simulations in Section 3.5 that this FRTE regime leads to quantitatively different evolutionary trajectories compared to the FRTI regime, we show mathematically in Appendix F.2 that the evolutionary dynamics can be approximated by the same equations (see Eq. (8)).

3. Results

In this section we proceed to analyse the evolutionary dynamics derived in Sections 2.1-2.2 and compare our results to numerical simulations of the full stochastic simulations.

3.1. Initial evolution of fertilization rate

In Figure 2, we see two potential evolutionary outcomes for the co-evolutionary dynamics of m and α in a single fixed environment that are dependent on the initial conditions and parameters; the population can either evolve to large fertilization rates (limited only by $\alpha_{\max}$) or to zero fertilization rates.

When the cost to fertilization, C_z, is low (Figure 2, panel (a)), there exists a smaller region of initial conditions that drive α towards zero (pink shaded region). When $\alpha = 0$ within this region, selection on gamete mass, m, drives the population towards the point $m = \beta$ (red dot, see also Appendix C.2). As this point exists outside the region in which $\mathrm{d}\alpha/\mathrm{d}\tau < 0$, selection for increased α can again manifest along the evolutionary

Figure 2: Phase portraits for the co-evolutionary dynamics in a fixed environment (see Eq. (5)). High fertilization rates are the only evolutionary outcome in panel (a), while high and zero fertilization rate are both evolutionary outcomes in panel (b) (see Eq. (9)). The red region shows trajectories leading to points on the $\alpha = 0$ boundary for which evolution selects for decreasing fertilization rate ($d\alpha/d\tau < 0$) (the point at which $d\alpha/d\tau = 0$ is marked by a red arrow). Red filled (stable) and unfilled (unstable) circles mark a fixed point in the evolutionary dynamics of m ($m^* = \beta$, see Eq. (C.2)). Blue circles and arrows illustrate the high fertilization rate attractor ($(m^*, \alpha^*) \to (\beta/4, \infty)$, see Eq. (C.4)). Average population trait trajectories, $(\langle m \rangle(t), \langle \alpha \rangle(t))$, from simulation of the stochastic model (see Appendix C.3) are plotted in light gray, and their mean over multiple realisations are given dark gray. The cost to fertilization is $C_z = 0.3$ (panel (a)) and $C_z = 0.6$ (panel (b)). In both panels $\beta = 1$ and $C_p = 0$. Remaining model parameters are given in Appendix G.

trajectory. Thus when costs are sufficiently low, high fertilization rates are the only evolutionary outcome.

Conversely when costs to fertilization, C_z, are intermediate (Figure 2, panel (b)), there exists a larger region of initial conditions that drive α towards zero (pink shaded region). The point $m = \beta$ (red dot), towards which the population evolves when $\alpha = 0$, is now contained within this region in which $d\alpha/d\tau < 0$, and so is a stable fixed point. Thus when costs are sufficiently high, there are two evolutionary outcomes, depending on the initial conditions; either high fertilization rates or zero fertilization rates.

In Appendix C.2 we conduct a mathematical and numerical analysis to formalise the arguments above. In summary, the possible early evolutionary attractors, (m^*, α^*), are given by:

$$
\begin{aligned}
\text{If} \quad & 1 - \frac{1 - C_p}{\sqrt{e}} > C_z \geq 0, \quad (m^*, \alpha^*) \to (\beta/4, \alpha_{\max}) \\
\text{If} \quad & 1 \gtrsim C_z > 1 - \frac{1 - C_p}{\sqrt{e}}, \quad (m^*, \alpha^*) \to (\beta/4, \alpha_{\max}) \\
& \qquad\qquad\text{or} \quad (m^*, \alpha^*) \to (\beta, 0) \\
\text{If} \quad & C_z \approx 1, \qquad\qquad (m^*, \alpha^*) \to (\beta, 0)
\end{aligned}
$$
(9)

where we note that in the absence of parthenogenesis costs i.e $C_p = 0$, the condition in which a high fertilization rate is the only evolutionary outcome is $0 \leq C_z < 1 - 1/\sqrt{e} \approx 0.39$ (see Figure 2). Since situations in which the fertilization rate initially tends to zero are not of interest in the current analysis, we work in the remainder of this paper in the regime in which increasing fertilization rate is selected for; that is, $1 \gtrsim C_z$ and with initial conditions that do not lead to a stable $\alpha = 0$

fixed point (see Figure 2, unshaded regions).

3.2. Evolutionary branching can lead to anisogamy, followed by oogamy

In Figure 2 we see that the approximation obtained for the co-evolutionary dynamics of m and α in a single fixed environment, Eq. (5), accurately captures the dynamics of the full model realised via numerical simulation at early times. One point of departure is that at long times as α increases along the $m \approx \beta/4$ manifold, we see the mean mass trait value from simulations increasing to higher values than those predicted analytically. In Figure 3, we show that this is a result of evolutionary branching in the simulations.

Should the evolutionary trajectories reach a sufficiently large fertilization rate, α, along the $m \approx \beta/4$ manifold, anisogamy can evolve through evolutionary branching in mass (see Figure 3), which gives rise to a dimorphic gamete population that contains one large gamete (macrogamete) with $m_{macro} \approx \beta - \delta m$ and one small gamete (microgamete) with $m_{micro} = \delta m$. Analytically, this is shown to be due to disruptive selection in mass in the vicinity of the $m \approx \beta/4$ manifold at high values of α. In Appendix D, we show that this region is an approximate evolutionary singularity. This initial branching has been noted in other models that also do not account for self-incompatible mating types but that consider the case of fixed α and obligate sexual reproduction [24].

As the mass of the macrogamete becomes larger (and that of the microgamete smaller), a branching in fertilization rates can also occur (see Figure 3 (b)). Near this mass $m_{macro} \approx \beta - \delta m$, the macrogamete has a high survival rate under parthenogenesis, and the benefits of fer-

Figure 3: Numerical illustration of evolutionary branching in Figure 2(b). Panel (a): Analytic predictions for the early evolutionary dynamics (as Figure 2(b)) overlaid with trajectories $(m_i(t), \alpha_i(t))$ for each i^{th} trait. Evolutionary branching is observed along the $m \approx \beta/4$ manifold, indicated by the blue star. Green arrows show the temporal progression of the branching. Panels (b) and (c): The temporal trajectories of the traits $\alpha_i(t)$ and $m_i(t)$ respectively, showing that the evolutionary trajectory passes from isogamy to oogamy. Parameters used are $A = 100$, $M = 1$, $T = 1$, $C_z = 0.6$, $C_p = 0$, $\beta = 1$, $\delta = 0.01$, $\mu = 10^{-3}$, $f_0 = 2 \times 10^{-3}$ and simulation run for 3.5×10^6 generations.

tilization (in particular with microgametes of very small mass) can be outweighed by the costs of fertilization. Selection can thus act to lower the fertilization rate of macrogametes, α_{macro}, towards zero. This in turn leads to an increased selection pressure for the microgametes to increase their fertilization rate, with $\alpha_{\text{micro}} \to \alpha_{\text{max}}$, to increase the probability of microgametes fertilizing macrogametes (averting the low survival probabilities of microgametes under parthenogenesis). This is a situation in which macrogametes are still fertilized by microgametes (at a rate $\alpha_{\text{micro}}/2$) but do not fertilize themselves. Biologically, this situation can be interpreted as the evolution of oogamy, with $\alpha_{\text{micro}} \to \alpha_{\text{max}}$ and obligate sexual reproduction in the limit $\alpha_{\text{max}} \to \infty$.

Above we have shown that when costs to fertilization are accounted for, a continuous evolutionary trajectory can exist that takes the population from a state of isogamy to oogamy. In the following sections we demonstrate how each of these transitions can be arrested under various parameter regimes.

3.3. A low ceiling on the fertilization rate can stabilize isogamy

If the maximum possible fertilization rate α_{max} is limited to a low value, for instance due to energetic constraints on motility or environmental constraints such as turbulence, then it is possible to prevent the transition from isogamy to anisogamy. In Appendix C.4, we derive the minimum value of α_{max} at which evolutionary branching in mass can occur, and find that if

$$\frac{48(1 - C_p)\beta}{35AMT(1 - C_z)e^{\frac{7}{6}}} > \alpha_{\text{max}} \tag{10}$$

evolutionary branching in mass cannot occur. These results are illustrated in Figure 4. Should α_{max} lie below this threshold (equivalent in approximately 70% of cells gametes fertilized in Figure 4, see Figure A.8) the population is held in a state of isogamy.

3.4. High costs of parthenogenesis relative to zygote formation can stabilize anisogamy

Analytically, we show in Appendix C.4 that when costs for zygote formation are relatively low and costs for parthenogenesis relatively high, true oogamy (in which the macrogamete's fertilization rate evolves towards zero, $\alpha_{\text{macro}} \to 0$) does not evolve following the transition to anisogamy. The costs to fertilization must instead be sufficiently high, such that

$$C_z > 1 - (1 - C_p) \exp\left(-\frac{\delta m}{\beta - \delta m}\right) \tag{11}$$

for true oogamy to evolve. This is illustrated in Figure 5. Essentially if the costs to the macrogamete of fertilization are not sufficiently high, then there is no longer a selective pressure (as described in Section 3.2) for the macrogamete to avoid fertilizations by decreasing its fertilization rate. While the population does not evolve towards oogamy, its ultimate state is dependent on the maximum fertilization rate, α_{max}.

Suppose we initially place no limit on the maximum fertilization rate, $\alpha_{\text{max}} \to \infty$, and that the costs C_p and C_z are such that there is no longer selection for the fertilization rate of the macrogamete to decrease (i.e. the inequality in Eq. (11) does not hold). While there may be a selective pressure for the fertilization rate of the macrogamete to increase, this selection pressure is weaker for the parthenogenetic macrogamete than the microgamete, as the microgamete relies more heavily on the fertilization pathway for its survival. Thus although the fertilization rates of both macrogametes and microgametes, α_{macro} and α_{micro}, evolve to increase α indefinitely the macrogamete does so at a considerably slower rate. Under this scenario, we therefore observe pseudoogamy [27], where the macrogamete fertilization rate does not evolve toward zero but is still lower

Figure 4: Analytical prediction (blue) and numerical illustration of the range of α_{max} in which branching to anisogamy is possible. Black markers represent the mass of each gamete genotype within the population after 2.5×10^6 generations. Once branching to anisogamy has occurred, a dimorphic gamete population, characterised by the presence of two genotypes would be present. One genotype where $m \approx \beta - \delta m$ and one where $m = \delta m$. Vertical line represents the analytically predicted α_{max} above which branching can occur in m Eq. (10). The blue curve left of this line is the numerical solution to $dm/d\tau|_{\alpha=\alpha_{max}} = 0$ in Eq. (C.1) and the blue lines towards the right contains the theoretical masses of the macrogamete ($m = \beta - \delta m$) and microgametes ($m = \delta m$). Parameters are $A = 100$, $M = 1$, $T = 0.1$, $C_z = 0.3$, $C_p = 0$, $\beta = 1$, $\delta = 0.01$, $\mu = 10^{-3}$ and $f_0 = 2 \times 10^{-3}$.

than the microgamete fertilization rate. While this situation is only observed when $\alpha_{max} \to \infty$, in the case of large but finite α_{max}, pseudooogamy is a prolonged transient state (see Figures 5 and E.13). Lastly, should α_{max} be smaller, such that the increasing fertilization rate of macrogametes can "catch up" with that of microgametes, the population can return to anisogamy following a transient period of oogamy.

3.5. In a switching environment, the population can evolve a bet-hedging strategy that stabilizes isogamy

In Figure 6, we see that the analysis for the evolutionary dynamics in the case of a switching environment (see Eq. (8)), provides a good approximation to the dynamics of the full model (which accounts for multiple traits coexisting under a mutation-selection balance) realised via numerical simulation. We now see three broad evolutionary outcomes.

We begin by considering the intuitive limit of $\tau_1 \gg \tau_2$. In this scenario the population spends almost all of the time in environment 1, and a comparatively insignificant amount of time in environment 2 (i.e. $P_1 \approx 1$ and $P_2 \approx 0$). Consequently, the population evolves approximately as if it were simply in a fixed environment with $\beta = \beta_1$, and the conditions given for the fixed environment, Eq. (9), can be used to infer the evolutionary outcome. An analogous argument holds for the dynamics when $\tau_2 \gg \tau_1$, but with $\beta = \beta_2$ in Eq. (9). Once on the

Figure 5: Numerical illustration of the ratio of macrogamete to microgamete fertilization rate, $\alpha_{macro}/\alpha_{micro}$. Oogamy is favoured over anisogamy above a sufficiently high fertilization cost C_z, predicted analytically by the vertical line (see Eq. (11)). Here $C_p = 0.5$, $(m(0), \alpha(0)) = (0.25, 0.02)$ and $\alpha_{max} = 1.3$. All other parameters are as in Figure 2 except $f_0 = 0.02$ and the simulation is run for 6×10^6 generations. Back arrows point in the direction towards which $\alpha_{macro}/\alpha_{micro}$ evolves in infinite time.

$m = \beta_i/4$ manifold towards the high-fertilization rate attractor, evolutionary branching towards anisogamy and oogamy can occur as described in Section 3.2. When the time spent in each environment is of the a comparable order however (e.g when τ_1, and τ_2 have not entirely dissimilar magnitudes), we find the emergence of bet-hedging strategies (see Figures 6, 7 and Eq. (14)).

Under a range of parameter conditions, we find analogous evolutionary attractors for the fertilization rate as in the fixed-environment case (see Eq. (9) and Figure 6), but with bet-hedging strategies for the gamete mass;

$$
(m^*, \alpha^*) \to (m^*_{\alpha=0}, 0)
$$
$$
(m^*, \alpha^*) \to (m^*_{\alpha\to\infty}, \alpha_{max})
$$
(12)

with

$$
m^*_{\alpha=0} = P_1\beta_1 + (1 - P_1)\beta_2
$$
$$
m^*_{\alpha\to\infty} = \frac{1}{4}(P_1\beta_1 + (1 - P_1)\beta_2).
$$
(13)

The population thus initially evolves either to high fertilization rates or zero fertilization rates, but with a mass which is the weighted average of the optimal strategy in either environment.

Under a more restricted set of parameter conditions however we find that a bet-hedging strategy for the fertilization rate can also evolve; essentially a switching-induced fixed point can manifest, as illustrated in Figures 6 and 7. Here the tension between the evolutionary dynamics in the two environments (which can select for high fertilization rate and large gametes in one environment, and zero fertilization rate and small gametes in the other) can lead to the population being held in a state at which intermediate finite fertilization rate and isogamy form the evolutionarily stable bet-hedging

Figure 6: Phase portraits for the co-evolutionary dynamics in a switching environment (see Eq. (8)). In addition to qualitatively similar dynamics as in the fixed environment (see Figure 2), two new evolutionary scenarios are now possible, including populations in which stable intermediate fertilization rates (filled blue circle) are the only evolutionarily stable state (see panel (a)) and populations in which there is an additional zero fertilization stable state (filled red circle, see panel (b)). Open orange circles represent the (now unstable) states to which the population can be attracted in either environment 1 or 2 (where $\beta = \beta_1$ or $\beta = \beta_2$). Average population trait trajectories, $(\langle m \rangle(t), \langle \alpha \rangle(t))$, from simulation of the full stochastic model in the FRTI regime are overlaid in gray. In panel (c) we plot the time-series in the FRTI regime (gray, as in panel (a)) alongside those in the FRTE regime (purple) and our analytic predictions (black-dashed). In all panels, $A = 100$, $M = 10$, $T = 0.1$ and $f_0 = 2 \times 10^{-3}$. Under the FRTI regime (overlaid in gray in panels (a), (b) and (c)), $\mu = 3.5 \times 10^{-4}$, $\delta = 7 \times 10^{-3}$ and the simulation is run for 10^7 generations. Under the FRTE regime (overlaid in purple in panel (c)), $\mu = 5 \times 10^{-3}$, $\delta = 5 \times 10^{-3}$ and the simulation is run for 7×10^6 generations. In panel (a) (and panel (c) gray), $C_z = 0.35$, $\beta_1 = 3$, $P_1 = 0.335$, $(m(0), \alpha(0)) = (2, 0.4)$, $\lambda_{1 \to 2} = 0.250$ and $\lambda_{2 \to 1} = 0.126$. In panel (b), $C_z = 0.7$, $\beta_1 = 4$, $P_1 = 0.74$, $(m(0), \alpha(0)) = (2, 0.6)$, $\lambda_{1 \to 2} = 5.86 \times 0.01$ and $\lambda_{2 \to 1} = 0.167$. In panels (c) and (d), the switching rates for the FRTE regime (purple) are $\lambda_{1 \to 2} = 2.93 \times 10^{-5}$ and $\lambda_{2 \to 1} = 8.34 \times 10^{-5}$. Switching rates and mutation rates μ are all measured in units of (number of generations)$^{-1}$ and $C_p = 0$ in all panels.

strategy. In Appendix F.3, we show that the switching-induced fixed point is given by

$$m^* = \frac{1}{4} \left[P_1 \beta_1 + (1 - P_1) \beta_2 \right]$$

$$\alpha^* = \frac{(1 - C_p) m^*}{AMT(1 - C_z)}$$

$$\times \left(\frac{(1 - C_z) \left(P_1 e^{\frac{\beta_1}{2m^*}} + (1 - P_1) e^{\frac{\beta_2}{2m^*}} \right) - (1 - C_p)}{(1 - C_p)(P_1 e^{\frac{\beta_2}{2m^*}} + (1 - P_1) e^{\frac{\beta_1}{2m^*}}) - (1 - C_z) e^{\frac{\beta_1 + \beta_2}{2m^*}}} \right)$$

(14)

In Figure 6(c-d) we see that this switching induced fixed point is observed in the evolutionary simulations in which multiple traits can be held in the population under a mutation-selection balance. In the FRTI regime (gray lines), in which the environment changes more quickly, the population is held at the predicted mass m, while more variability between simulations is seen around the predicted fertilization rate α as a result of the weaker selection on this trait. In the slower-switching FRTE regime (purple lines), a greater quantitative difference between the results of simulations and the analytic predictions is observed. However in both regimes, the key prediction of finite fertilization rate (Figures 6 and 7)

and isogamy is indeed captured (see Figures F.15 and F.16).

4. Conclusions

In this paper we have extended classic results on the evolution of anisogamy [14, 25] to account for both parthenogenetic development [40, 41] and the co-evolution of fertilization rate and gamete cell mass. In doing so we have demonstrated the possibility of a continuous evolutionary trajectory from an initial state of isogamy to anisogamy followed by oogamy, an evolutionary trajectory observed throughout the eukaryotes [3, 10].

Consistent with earlier theoretical results on the evolution of anisogamy that neglected the pre-existence of self-incompatible mating types [24], we have shown that anisogamy can evolve from isogamy through evolutionary branching. However our model also shows that when the possibility of parthenogenetic development is accounted for, a subsequent branching to oogamy in the population is possible. Importantly, this transition is possible even without explicitly accounting for pheromone-receptor systems [29, 30], speed-mass relationships [32], mechanistic gamete encounter rates [31]

Figure 7: Analytic theory for the fertilization rate at the switching-induced isogamous fixed point (blue line, see Eq. (14)) against simulations (dots). The black vertical line is the analytic prediction for when this fixed point vanishes, with parameters left of the line corresponding to destabilised isogamy. Simulations are obtained by averaging over 25 realisations of $\langle \alpha(t) \rangle$ corresponding to stochastic evolutionary trajectories. Parameters are $\delta = 0.01$, $\mu = 5 \times 10^{-4}$, $f_0 = 2 \times 10^{-3}$, $\tau_1 = 0.25$ and simulation run for 1.2×10^7 generations. All other parameters are the same as in the FRTI regime of Figure 6 (a). Red markers are for $(m(0), \alpha(0)) = (0.29, 0.075)$, green for $(m(0), \alpha(0)) = (0.29, 0.15)$ and blue for $(m(0), \alpha(0)) = (0.29, 0.3)$.

or costly motility [57], as in previous work. Instead in our model, oogamy can naturally evolve from anisogamy via sexual conflict [58], following the evolution of anisogamy from isogamy. In particular, microgametes receive a survival advantage by increasing their fertilization rate with macrogametes, while macrogametes evolve to zero fertilization rate to overcome fertilization costs. This requirement of fertilization costs for oogamy to evolve under parthenogenesis provides a complementary result to [27], where instead high motility costs were required for oogamy.

We have shown analytically that various conditions exist that can arrest the population at different stages of the isogamy-anisogamy-oogamy evolutionary trajectory. In particular our model suggests that isogamy can be stabilised under greater energetic constraints or in highly turbulent environments. We have also shown that isogamy can be stabilised as a bet-hedging strategy under switching environmental conditions. Here, in organisms that reproduce parthenogenetically, microgametes that fail to find a partner face low survival prospects under harsh environmental conditions, and thus the transition to anisogamy is frustrated. Such dynamics may be at play in isogamous species that reproduce parthenogenetically, such as those among the *ectocarpus* [59] and *bildingia minima* [36].

The results described above are dependent on the capacity for parthenogenetic development amongst gametes that fail to find a partner to fertilize with. Such a capacity is widespread in the brown algae [47] and can also be found in the green algae [40]. We have

also shown that for the evolution of oogamy in our model, we require costs of fertilization to exceed those of parthenogenetic development. While empirically parthenosporophytes have lower survival probabilities than zygotes [40], disentangling the costs of development pathways (parthenogenesis or fertilization) from those of increased mass is challenging [41]. By invoking costs to zygote formation arising from cytoplasmic conflict [50], our work is reminiscent of alternative hypotheses for the evolution of anisogamy [51, 52], that suggest that anisogamy reduces the potential for such conflict by limiting cytoplasmic contributions from the microgamete. However many other costs to fertilization could be present, including the possibility of failure during cell fusion [49], and the energetic and time costs of sexual reproduction [60]. Unpacking and quantifying these costs remains an interesting area of empirical research.

Our primary reason for neglecting the existence of mating types in our model was for mathematical simplicity [24]. Including such types in a model would be a natural next step, and based on the biological reasoning above, such a model should demonstrate similar qualitative behaviours. However it is worthwhile noting that many models of the evolution of sexual reproduction in early eukaryotes suppose the existence of a "unisexual" early ancestor that mated indiscriminately [61]. Thus our model, which has neglected the existence of mating types, could be seen as reflecting such a system. In this context, the evolution we have seen of incompatibility between macrogametes (that fuse with each other at rate zero in the oogamous scenario) is interesting, as the origins of self-incompatible mating types are still debated [62, 63]. This mechanism would represent an "anisogamy consequence" model for the evolution of mating types that manages to identify the conditions under which fusion between large macrogametes is disadvantageous [64]. However as such "anisogamy consequence" models are not consistent with most empirical observations (that suggest mating types preceded anisogamy), this interpretation should be treated with caution.

As with any model, there are obviously numerous other omissions from our formulation. These include the mortality of gametes (known to stabilise isogamy [25]) the discrete nature of cell divisions leading to gametes [40], and non-local trait mutations. While these would be interesting additions to our model, the key insights derived from the PBS model have remained remarkably robust to such generalisations, and so the inclusion of these additional considerations to our model may lead primarily to quantitative,

rather than qualitative, changes in results. More generally, extending our mathematical approach leveraging adaptive dynamics to switching environments in other facultatatively sexual populations might prove particularly fruitful [65, 66].

In this paper, we have extended the models of [24, 67] in several ways; by allowing the fertilization rate to evolve, accounting for the possibility for unfertilized gametes to develop parthenogenetically should they fail to locate a partner, and subjecting the system to switching environments. In doing so, we have shown its capacity to parsimoniously capture continuous evolutionary trajectories from isogamy to oogamy in parthenogens. Moreover, our models emphasise the importance of investigating the coevolutionary dynamics for a range of evolutionary parameters and its implications for the evolution of oogamy.

Appendix A. Within generation dynamics

At the start of each generation, we assume a total of A organisms of mature cell size (or energy budget) M divide to form gametes. We further assume that the frequency of mutants in this adult population is given by $1 \geq \hat{f}_i \geq 0$, with $i = m$ for mutants that change the mass of gametes and $i = \alpha$ for mutants that change the fertilization rate of gametes.

Appendix A.1. Fertilization kinetics: mutant with different mass

If a mutation occurs changing the size of gametes produced by the mutant ($\hat{m} = m + \delta m$ for resident gamete mass m), the fertilization kinetics themselves (see Eq. (1)) are unaffected by the change in mass. However the number of gametes produced by the mutant will change. Denoting by N the number of resident gametes and $\hat{N}$ the number of mutant gametes, we have

$$\frac{\mathrm{d}N}{\mathrm{d}t} = -\alpha(N^2 + N\hat{N}); \qquad N(0) = \frac{AM(1 - \hat{f}_m)}{m}$$

$$\frac{\mathrm{d}\hat{N}}{\mathrm{d}t} = -\alpha(\hat{N}^2 + N\hat{N}); \qquad \hat{N}(0) = \frac{AM\hat{f}_m}{\hat{m}},$$

(A.1)

which has a solution

$$N(t) = \frac{N(0)}{1 + \left(N(0) + \hat{N}(0)\right)\alpha t}$$

$$\hat{N}(t) = \frac{\hat{N}(0)}{1 + \left(N(0) + \hat{N}(0)\right)\alpha t}.$$

(A.2)

This allows us to determine the number of unfertilized cells of each type at the end of the fertilization window at $U = N(T)$ and $\hat{U} = \hat{N}(T)$, where we recall that T is the length of the fertilization window.

We also need to determine the number of fertilized cells formed from the fertilization of two residents, F_N^N, two mutants, $F_{\hat{N}}^{\hat{N}}$, and a mutant and a resident, $F_N^{\hat{N}}$. For this we need to solve

$$\frac{\mathrm{d}F_N^N}{\mathrm{d}t} = \frac{1}{2}\alpha N^2; \qquad F_N^N(0) = 0$$

$$\frac{\mathrm{d}F_N^{\hat{N}}}{\mathrm{d}t} = \alpha N\hat{N}; \qquad F_N^{\hat{N}}(0) = 0$$

$$\frac{\mathrm{d}F_{\hat{N}}^{\hat{N}}}{\mathrm{d}t} = \frac{1}{2}\alpha \hat{N}^2; \qquad F_{\hat{N}}^{\hat{N}}(0) = 0,$$

(A.3)

which can be solved by substituting for N and $\hat{N}$ from Eq. (A.2) to yield

$$F_N^N(t) = \frac{1}{2}\frac{N(0)^2 t}{1 + \left(N(0) + \hat{N}(0)\right)\alpha t}$$

$$F_N^{\hat{N}}(t) = \frac{N(0)\hat{N}(0)t}{1 + \left(N(0) + \hat{N}(0)\right)\alpha t}$$

(A.4)

$$F_{\hat{N}}^{\hat{N}}(t) = \frac{1}{2}\frac{\hat{N}(0)^2 t}{1 + \left(N(0) + \hat{N}(0)\right)\alpha t}.$$

Figure A.8: Illustration of the total proportion of cells that are fertilized at the end of a fertilization period (length $T = 1$) in a monomorphic isogamous population (without branching) as a function of trait variables m and α. Parameters used are the same as those in Figure 2. The vertical black dashed line gives the location of the manifold $(\beta/4, \alpha)$, along which the population is attracted to when approaching the high α attractor.

Appendix A.2. Fertilization kinetics: mutant with different fertilization rate

If a mutation occurs changing the fertilization rate of gametes produced by the mutant ($\hat{\alpha} = \alpha + \delta\alpha$ for residents with fertilization rate α), the fertilization kinetics themselves are altered relative to Eq. (1). We assume for simplicity that the fertilization rate between resident-pairs is the mean of the fertilization rate between the two types in isolation, such that

$$\frac{\mathrm{d}N}{\mathrm{d}t} = -\alpha N^2 - \left(\frac{\alpha + \hat{\alpha}}{2}\right)N\hat{N}; \qquad N(0) = \frac{AM(1 - \hat{f}_\alpha)}{m}$$

$$\frac{\mathrm{d}\hat{N}}{\mathrm{d}t} = -\hat{\alpha}\hat{N}^2 - \left(\frac{\alpha + \hat{\alpha}}{2}\right)N\hat{N}; \qquad \hat{N}(0) = \frac{AM\hat{f}_\alpha}{m}.$$

(A.5)

Solving this equation is slightly less straightforward than solving Eq. (A.1). However we can make analytic progress by making a change of variables and applying an approximation based on small mutational step size $\delta\alpha$.

We introduce the transformed variables N_{tot} and $\hat{r} = \frac{\hat{N}}{N+\hat{N}}$, representing the total number of unfertilized cells and the frequency of unfertilized mutant cells respectively. Eq. (A.5) then becomes

$$\frac{\mathrm{d}N_{\text{tot}}}{\mathrm{d}t} = -N_{\text{tot}}^2(\alpha + \delta\alpha\,\hat{r})$$
$$\frac{\mathrm{d}\hat{r}}{\mathrm{d}t} = -\frac{1}{2}\delta\alpha\,N_{\text{tot}}\hat{r}(1-\hat{r})\,. \tag{A.6}$$

We now see that although this equation is also intractable, the leading order dynamics of N_{tot} are governed by α. Therefore when $\alpha \gg \delta\alpha$, we make the approximation $(\mathrm{d}N_{\text{tot}}/\mathrm{d}t) \approx -\alpha N_{\text{tot}}^2$. We then obtain

$$N_{\text{tot}}(t) \approx \frac{N_{\text{tot}}(0)}{1 + N_{\text{tot}}(0)\alpha t}\,, \tag{A.7}$$

and substituting this into our equation for $\hat{r}$ in Eq. (A.6), we can solve to obtain

$$\hat{r}(t) = \frac{\hat{r}(0)}{\hat{r}(0) + (1-\hat{r}(0))(1 + N_{\text{tot}}(0)\alpha t)^{\frac{\delta\alpha}{2\alpha}}} \tag{A.8}$$

Inverting the transformation, we then arrive at

$$N(t) = N_{\text{tot}}(t)(1 - \hat{r}(t))$$
$$\hat{N}(t) = N_{\text{tot}}(t)\hat{r}(t)\,, \tag{A.9}$$

with N_{tot} and $\hat{r}(t)$ taken from Eqs. (A.7-A.8). We shall see in Appendix A.3 that calculating F_N^N, $F_{\hat{N}}^N$, and $F_N^{\hat{N}}$ explicitly is in fact unnecessary, and these expressions for $N(t)$ and $\hat{N}(t)$ are sufficient for analytical progress.

Appendix A.3. Change in mutant frequency over a generation: mutant with different mass

We begin by calculating the fitnesses of the resident and a mutant that changes the mass of gametes, w_m and $\hat{w}_m$ respectively, which are simply give by the total number of cells of each type at the end of a generation. Recalling that fertilized and unfertilized gametes both survive with a probability governed by the parameter β and the mass of the cell m_c (see Eq. (2)), and that fertilized cells survive with an additional probability $(1 - C_z)$, we have

$$w_m = (1 - C_z)\big(2F_N^N(T)S(\beta, 2m)$$
$$+ F_N^{\hat{N}}(T)S(\beta, m+\hat{m})\big) + N(T)S(\beta, m)(1 - C_p)$$
$$\hat{w}_m = (1 - C_z)\big(2F_{\hat{N}}^{\hat{N}}(T)S(\beta, 2\hat{m})$$
$$+ F_N^{\hat{N}}(T)S(\beta, m+\hat{m})\big) + \hat{N}(T)S(\beta, \hat{m})(1 - C_p)$$
$$\tag{A.10}$$

where $F_N^N(T)$, $F_{\hat{N}}^N(T)$, and $F_N^{\hat{N}}(T)$ are taken from Eq. (A.4), and $N(t)$ and $\hat{N}(t)$ are taken from Eq. (A.2). These expressions can be used to calculate the frequency at the end of the generation, $\hat{f}_m'$, of a mutant that changes the mass of gametes as

$$\tilde{\hat{f}}_m = \frac{\hat{w}_m}{w_m + \hat{w}_m}\,. \tag{A.11}$$

The change is the frequency of the mutant over the course of a generation is then $\hat{f}_m' - \hat{f}_m$.

Appendix A.4. Change in mutant frequency over a generation: mutant with different fertilization rate

Taking an analogous approach to Appendix A.3, we begin by calculating the fitnesses of the resident and a mutant that changes the fertilization rate, w_α and $\hat{w}_\alpha$ respectively. We obtain

$$\begin{aligned}
w_\alpha &= (1 - C_z)\big(2F_N^N(T)S(\beta, 2m)\\
&\quad + F_N^{\hat{N}}(T)S(\beta, 2m)\big) + (1 - C_p)N(T)S(\beta, m)\\
&= (1 - C_z)S(\beta, 2m)(N(0) - N(T)) + (1 - C_p)N(T)S(\beta, m)\\
\hat{w}_\alpha &= (1 - C_z)\big(2F_{\hat{N}}^{\hat{N}}(T)S(\beta, 2m)\\
&\quad + F_N^{\hat{N}}(T)S(\beta, 2m)\big) + (1 - C_p)\hat{N}(T)S(\beta, m)\\
&= (1 - C_z)S(\beta, 2m)\big(\hat{N}(0) - \hat{N}(T)\big) + (1 - C_p)\hat{N}(T)S(\beta, m)
\end{aligned} \tag{A.12}$$

where $N(t)$ and $\hat{N}(t)$ are now taken from Eq. (A.9). Here we have used the fact that since the survival function for fertilized cells, $S(\beta, 2m)$, is independent of the composition of the fertilized cells (mutations here only affect fertilization rate) the number of cell-types contributing to the fertilized cells can be inferred under cell conservation during the fertilization period (e.g. for resident cell types $\mathrm{d}/\mathrm{d}t\big[2F_N^N(t) + F_N^{\hat{N}}(t) + N(t)\big] = 0$, and similarly for mutant cell types).

These expressions can be used to calculate the frequency at the end of the generation, $\hat{f}_\alpha'$, of a mutant that changes the fertilization rate as

$$\tilde{\hat{f}}_\alpha = \frac{\hat{w}_\alpha}{w_\alpha + \hat{w}_\alpha}\,. \tag{A.13}$$

The change is the frequency of the mutant over the course of a generation is then $\hat{f}_\alpha' - \hat{f}_\alpha$.

Appendix B. Invasion dynamics

Our aim is to derive the dynamics of the frequency of a mutant with a small mutation in either of the traits, m or α, over multiple generations. We introduce t_g as a measure of the number of discrete generations.

Figure B.9: Invasion dynamics for a mutant with mass $m + \delta m$. Blue - analytical prediction using Eq. (B.1), black - numerical simulation. The initial condition is $(m(0), \alpha(0)) = (0.4, 0.1)$. Parameters are $\delta m = -0.005$, $f_0 = 0.002$, $G = 4 \times 10^3$, $A = 100$, $M = 1$, $T = 1$, $C_z = 0.6$, $C_p = 0$ and $\beta = 1$.

Figure B.10: Invasion dynamics for a mutant with fertilization rate $\alpha + \delta \alpha$. Blue - analytical prediction using Eq. (B.2), black - numerical simulation. The initial conditions and parameters are the same as in Figure B.9 except $\delta \alpha = 1/200$, $\delta m = 0$ and $G = 1.5 \times 10^4$.

Appendix B.1. Invasion ODE: mutant with different mass

We begin by deriving the dynamics for the frequency of a mutant that changes the mass of gametes. We begin by assuming that the mutational step size, δm, is small. Under these conditions, the frequency of mutants changes only by a small amount over the course of one generation, and we can approximate the frequency of mutants at the end of the generation (see Appendix A.3) by $\hat{f}'_m = \hat{f}_m + \delta m \, (\mathrm{d}\hat{f}'_m / \mathrm{d}\,\delta m)|_{\delta m = 0}$, where $\hat{f}_m$ is the frequency of the mutants at the beginning of the generation. The dynamics of the mutant frequency over an invasion can then be approximated by

$$
\begin{aligned}
\frac{\mathrm{d}\hat{f}_m}{\mathrm{d}t_g} &= \left. \left(\frac{\hat{f}'_m - \hat{f}_m}{t_g} \right) \right|_{t_g \to 1} \\
&= \delta m \left[\frac{\mathrm{d}\hat{f}'_m}{\mathrm{d}\,\delta m} \right]_{\delta m = 0} \\
&= \delta m \left[\frac{AM\alpha T(1 - C_z)e^{\frac{\beta}{2m}}(\beta - 4m) - 4m(m - \beta)(1 - C_p)}{4m^2 \left(AM\alpha T(1 - C_z)e^{\frac{\beta}{2m}} + m(1 - C_p) \right)} \right] \\
&\quad \times \hat{f}_m(1 - \hat{f}_m) \\
&\equiv h_m(m, \alpha, \beta, C_z, C_p)\hat{f}_m(1 - \hat{f}_m),
\end{aligned}
\tag{B.1}
$$

(see also Eq. (3)), where $\mathrm{d}\hat{f}'_m / \mathrm{d}\,\delta m$ is derived from Eq. (A.11). We show in Figure B.9 that as expected, this is a good approximation for the dynamics when δm is small.

Appendix B.2. Invasion ODE: mutant with different fertilization rate

We now derive the dynamics for the frequency of a mutant that changes the fertilization rate of gametes. Taking an analogous approach to Appendix B.1, we assume the mutational step size for mutation, $\delta \alpha$, is small. We can then approximate the frequency of mutants at the end of the generation (see Appendix A.4) by $\hat{f}'_\alpha = \hat{f}_\alpha + \delta \alpha \, (\mathrm{d}\hat{f}'_\alpha / \mathrm{d}\,\delta \alpha)|_{\delta \alpha = 0}$, where $\hat{f}_\alpha$ is the frequency of the mutants at the beginning of the generation. The dynamics of the mutant frequency over an invasion can then be approximated by

$$
\begin{aligned}
\frac{\mathrm{d}\hat{f}_\alpha}{\mathrm{d}t_g} &= \left. \left(\frac{\hat{f}'_\alpha - \hat{f}_\alpha}{t_g} \right) \right|_{t_g \to 1} \\
&= \delta \alpha \left[\frac{\mathrm{d}\hat{f}'_\alpha}{\mathrm{d}\,\delta \alpha} \right]_{\delta \alpha = 0} \\
&= \delta \alpha \left[\frac{m \left((1 - C_z)e^{\frac{\beta}{2m}} - (1 - C_p) \right) \ln \left(\frac{\alpha AMT}{m} + 1 \right)}{2\alpha \left(\alpha A(1 - C_z)MT e^{\frac{\beta}{2m}} + m(1 - C_p) \right)} \right] \\
&\quad \times \hat{f}_\alpha(1 - \hat{f}_\alpha) \\
&\equiv h_\alpha(m, \alpha, \beta, C_z, C_p)\hat{f}_\alpha(1 - \hat{f}_\alpha),
\end{aligned}
\tag{B.2}
$$

(see also Eq. (4)) where $\mathrm{d}\hat{f}'_\alpha / \mathrm{d}\,\delta \alpha$ is derived from Eq. (A.13). We show in Figure B.10 that as expected, this is a good approximation for the dynamics when $\delta \alpha$ is small.

Appendix C. Evolutionary dynamics: Fixed environment

Appendix C.1. Derivation of evolutionary ODEs

We begin by noting that the functional form of $\mathrm{d}\hat{f}_m/\mathrm{d}t_g$ (see Eq. (B.1)) and $\mathrm{d}\hat{f}_\alpha/\mathrm{d}t_g$ (see Eq. (B.2)) is $\hat{f}_i(1 - \hat{f}_i)$, which implies a situation of trait substitution [54]; mutants are either driven to fixation or extinction, and polymorphic equilibria are not possible. This simplifies the subsequent analysis considerably.

Taking a classic adaptive dynamics approach [54] and define the invasion fitness of the mutants as their per-capita rate of reproduction upon arising in the population (i.e. when $\hat{f}_m \approx 0$ and $\hat{f}_\alpha \approx 0$). Under the standard assumptions of adaptive dynamics (i.e. that mutations are of small effect, $1 \gg \delta m, \delta \alpha$, and occur sufficiently rarely that each mutation can fixate before a new mutation occurs), the evolutionary dynamics are given by

$$
\begin{aligned}
\frac{\mathrm{d}m}{\mathrm{d}\tau} &= \frac{\mathrm{d}}{\mathrm{d}\hat{f}_m}\frac{\mathrm{d}\hat{f}_m}{\mathrm{d}t_g}\bigg|_{\hat{f}_m=0} = h_m(m, \alpha, \beta, C_z, C_p) \\
&= H_m(m, \alpha, \beta, C_z, C_p) \\
\frac{\mathrm{d}\alpha}{\mathrm{d}\tau} &= \frac{\mathrm{d}}{\mathrm{d}\hat{f}_\alpha}\frac{\mathrm{d}\hat{f}_\alpha}{\mathrm{d}t_g}\bigg|_{\hat{f}_\alpha=0} = h_\alpha(m, \alpha, \beta, C_z, C_p) \\
&= H_\alpha(m, \alpha, \beta, C_z, C_p) \, .
\end{aligned}
\tag{C.1}
$$

Substituting for $\mathrm{d}\hat{f}_m/\mathrm{d}t_g$ from Eq. (B.1), and $\mathrm{d}\hat{f}_\alpha/\mathrm{d}t_g$ from Eq. (B.2), we obtain Eq. (5) in the main text.

Appendix C.2. Analysis of ODEs

In this section we aim to analytically characterise the long term evolutionary behaviour of the population in a fixed environment, with dynamics given by Eq. (5), as illustrated in Figure 2.

We begin by calculating the evolutionary behaviour of m when the fertilization rate is fixed to zero ($\alpha = 0$). Solving $H_m(m, 0, \beta, C_z, C_p) = 0$, for $m = m^*_{\alpha=0}$ we obtain

$$
\delta m \frac{(\beta - m)}{m^2} = 0
$$
$$
\implies m^*_{\alpha=0} = \beta \, .
\tag{C.2}
$$

We now turn to the evolutionary behaviour of m and α. For $(m^*_{\alpha=0}, 0)$ identified above to remain stable if evolution on α is allowed requires that $[\mathrm{d}\alpha/\mathrm{d}\tau]|_{(m=\beta,\alpha=0)} < 0$ (i.e. that evolution selects against increases in α). Stating this condition in full, we have

$$
0 > \delta \alpha \left[\frac{m\left((1-C_z)e^{\frac{\beta}{2m}} - (1-C_p)\right)\ln\left(\frac{\alpha AMT}{m} + 1\right)}{2\alpha\left(\alpha A(1-C_z)MT e^{\frac{\beta}{2m}} + m(1-C_p)\right)} \right]\Bigg|_{(m=\beta,\alpha=0)}
$$
$$
\implies 0 > (1-C_z)e^{\frac{1}{2}} - (1-C_p)
$$
$$
\implies C_z > 1 - (1-C_p)e^{-\frac{1}{2}} \, .
\tag{C.3}
$$

If fertilization costs exceed this this value, evolutionary trajectories starting at $(m^*_{\alpha=0}, 0)$ will remain there (see Figure 2, panels (b-c)). Conversely if fertilization costs do not exceed this value, evolutionary trajectories starting at $(m^*_{\alpha=0}, 0)$ will initially experience a selective pressure for increasing α, due to the fact $[\mathrm{d}\alpha/\mathrm{d}\tau]|_{m=\beta} > 0$ under these conditions (see Figure 2, panel (a)); the dynamics are then pulled to another attractor, which we characterise below.

When $C_z > 1 - (1 - C_p)\exp(-1/2)$ (but less than another critical value, yet to be determined), only a subset of initial conditions fall within the basin of attraction of this attractor described above (see Figure 2, panel (b)), with remaining initial conditions leading to an evolutionary state at which $\alpha^* \to \infty$. Conversely, when $C_z < 1 - (1 - C_p)\exp(-1/2)$, *all* initial conditions lead to this attractor at which $\alpha^* \to \infty$ (see Figure 2, panel a). We now calculate the mass to which the population evolves at this second attractor. Taking the limit $\alpha \to \infty$ in the evolutionary dynamics for m (see Eq. (5)) and solving for zero;

$$
\begin{aligned}
0 &= H_m(m, \alpha, \beta, C_z, C_p)|_{\alpha\to\infty} \\
&= \delta m \frac{(\beta - 4m)}{4m^2} \\
\implies m^* &= \frac{\beta}{4} \, .
\end{aligned}
\tag{C.4}
$$

Therefore the second early evolutionary attractor is at $(m, \alpha) \to (\beta/4, \infty)$.

For even greater costs of fertilization, C_z, our mathematical analysis (which assumes monomorphic resident populations and no evolutionary branching) suggests that $(m, \alpha) \to (m^*_{\alpha=0}, 0)$ becomes the only attractor, with $(m, \alpha) \to (\beta/4, \infty)$ ceasing to be an attractor. To determine the critical cost at which this occurs, we take $[\mathrm{d}\alpha/\mathrm{d}\tau]_{m=\beta/4}$ and calculate the conditions under which this is negative when α is large (i.e. when $(m, \alpha) \to (\beta/4, \infty)$ is no longer attracting, but repelling). Expanding $[\mathrm{d}\alpha/\mathrm{d}\tau]_{m=\beta/4}$ in small $1/\alpha$, we find that to leading order we must have;

$$0 > \delta\alpha \left[\frac{m\left((1 - C_z)e^2 - (1 - C_p)\right)\ln\left(\frac{\alpha AMT}{\beta}\right)}{8\alpha^2 A(1 - C_z)MTe^2} \right]$$

$$\implies 0 > (1 - C_z)e^2 - (1 - C_p)$$

$$\implies C_z > 1 - (1 - C_p)e^{-2}$$

$$\implies C_z \gtrapprox 0.86 , \qquad \text{for } C_p = 0$$

(C.5)

However, while this accurately captures the short-term evolutionary dynamics, we see evolutionary branching (which our model Eq. (5) does not account for) should trajectories approach the $m \approx \beta/4$ manifold. Unless costs are exceedingly high ($C_z \approx 1$), this eventually leads to anisogamy followed by oogamy (see Figure C.11). We discuss this more in Appendix D.

Appendix C.3. Implementation of simulations

Here we detail the process of numerical simulation (see *Code availability statement*). We employ a multigenotype model whereby a mutation occurs at rate μ. This rate is the inverse of the expected number of fertilization processes (i.e. generations) until the next mutation event. Before each successive mutation event, the number of fertilization processes until the next mutation event is determined by generating a number from a $Geo(\mu)$ distribution and taking the inverse of that number. In an S genotype model, if a given genotype i has a trait value of (m_i, α_i) and has frequency f_i, then the mean population trait value is given by

$$\langle m \rangle = \sum_{i=1}^{S} m_i f_i \quad , \quad \langle \alpha \rangle = \sum_{i=1}^{S} \alpha_i f_i$$

(C.6)

The simulations in Figure 2 are repeated for 5500 mutation events. Below, we detail how we simulate the dynamics on each timescale.

Fertilization Kinetics

We construct our model for fertilization kinetics assuming that each genotype, characterised by their unique (m_i, α_i), can fertilize with one another. The input parameters of this function are the trait values of each genotype m and α, their frequency in the preceding adult generation f, and the parameters A, M and T. The fertilization kinetics simulations are then run according to

$$\frac{dN_i}{dt} = -\frac{1}{2}\sum_{j=1}^{S}(\alpha_i + \alpha_j)N_i N_j$$

(C.7)

$$\frac{dF_{ij}}{dt} = \frac{1}{2}(\alpha_i + \alpha_j)N_i N_j$$

(C.8)

where F_{ij} is the number of fertilized cells formed from genotypes i and j, and initial conditions

$$N_i(0) = \frac{AMf_i}{m_i}$$

(C.9)

$$F_{ij}(0) = 0$$

(C.10)

The fertilization process is run for a fixed time period T. At the end of this time period, the function outputs the number of cells of each type. These include the number of unfertilized cells with each trait pair, $N_i(T)$, and fertilized cells of each type $F_{ij}(T)$.

Single Generation Dynamics

We simulate the frequency of each genotype after the end of each generation taking into account the survival probability of each genotype. Upon maturation, only a fraction of unfertilized and fertilized cells survive to adulthood. We use the outputs of the fertilization kinetic function along with the Vance survival functions Eq. (2) to calculate the probability that each progeny survives into adulthood. The single generation dynamics are run according to

$$f_i' = \frac{w_i}{\sum_{j=1}^{S} w_j}$$

(C.11)

$$w_i = (1 - C_p)N_i(T)S(\beta, m_i)$$

(C.12)

$$+ (1 - C_z)\left(\sum_{j=1, j \neq i}^{S} F_{ij}(T)S(\beta, m_i + m_j) \right.$$

$$+ \left. 2F_{ii}(T)S(\beta, 2m_i) \right)$$

where f_i' is the frequency of genotype i in the subsequent generation and w_i is its absolute fitness. The input parameters of this function are $N_i(T)$ and $F_{ij}(T)$, m, α, β, C_z and C_p whilst the output is f, the frequency of all genotypes at the end of the generation.

Invasion Dynamics

The invasion dynamics are run for approximately one unit of τ (i.e. until the next randomly chosen mutation event after G generations). As outlined in the opening paragraph of Appendix C.3, G is generated from a $Geo(\mu)$ distribution following each mutation event. The inputs of this function are f, m, α, A, M, T, C_z, C_p, G and β and the output is the frequency of each genotype f after one unit of τ (i.e after G fertilization processes). Please note we have adopted one single notation for the frequency of all genotypes.

Evolutionary Dynamics

The evolutionary dynamics have the input parameters δ, μ, $\boldsymbol{f}$, NEVOL, f_0, A, M, T, C_z and C_p where δ is the mutational stepsize, the initial frequency of a newly introduced mutant f_0 is chosen to be small (equal to 0.002 in our model). We initialise the simulation with two genotypes, where each genotype is characterised by a unique pair of trait values e.g. (m_i, α_i) for genotype i. A mutant is introduced into the population after a random number of fertilization processes G (generated from a $Geo(\mu)$ distribution) at frequency f_0. The mutation is chosen to occur in either α or m with equal probability $1/2$. The mutation also acts to increase/decrease the trait value each with probability $1/2$. Upon introduction of this mutant, the invasion dynamics of the population is run for G fertilization processes. In the meantime, the mean mass and fertilization rate of the population is recorded using Eq. (C.6). Next, we repeat the process of introducing a new mutant into the population. Since the population now has more than two genotypes, the genotype that mutates is chosen with probability weighted by the frequency of each genotype. There is now the possibility of back mutation to one of the existing genotypes. In this case, if f_k is frequency of this genotype before the introduction of the genotypically identical mutant, then the frequency of this genotype becomes $(1-f_0)f_k + f_0$ after its introduction. Furthermore, a genotype is thought to be extinct if its frequency falls below 10^{-3}, in which case we remove that genotype.

Appendix C.4. Evolutionary Branching in Gamete Mass and fertilization rate

In this section we present additional numerical results investigating the evolutionary branching that occurs in simulations on the manifold $m \approx \beta/4$ (along which $d\alpha/dt \approx 0$).

In a fixed environment, when C_z is small relative to C_p, branching in fertilization rate still occurs but no longer gives rise to oogamy

When the cost to fertilization, C_z is small compared to C_p, we find that although branching in both gamete mass and fertilization rate still occurs, the branching in α no longer acts to decrease the fertilization rate of macrogametes. Therefore, we observe pseudooogamy but not oogamy. This behaviour is illustrated in Figures 5 and E.13. We can estimate the critical value of C_z at which this occurs by comparing the survival probability of a macrogamete that does not fertilize with a microgamete, which has cost C_p with that of a macrogamete that does fertilize with a microgamete at cost C_z; if this first probability exceeds the second, there should be no evolutionary pressure for oogamy to evolve. We find

$$(1-C_p)\exp\left(-\frac{\beta}{m_{\mathrm{macro}}}\right) > (1-C_z)\exp\left(-\frac{\beta}{m_{\mathrm{macro}} + m_{\mathrm{micro}}}\right)$$
(C.13)

or, noting that m_{macro} approaches $\beta - \delta m$ and m_{micro} approaches δm

$$(1 - C_p)\exp\left(-\frac{\beta}{\beta - \delta m}\right) > (1 - C_z)\exp(-1) \qquad \text{(C.14)}$$

$$\implies 1 - (1 - C_p)\exp\left(-\frac{\delta m}{\beta - \delta m}\right) > C_z \geq 0. \qquad \text{(C.15)}$$

In a fixed environment, when C_z is very large, branching in mass and fertilization rates can still occur, with oogamy possible (dependent on initial conditions)

Our analysis of the dynamics of Eq. (5) in Appendix C.2 suggested that zero fertilization rates is the only evolutionary attractor when costs to fertilization are high (see Eq. (C.5)). However, our analysis in Appendix D along with our simulations reveal that although $d\alpha/dt < 0$ along the $m \approx \beta/4$ manifold, any trajectory that approaches this manifold can experience branching in gamete mass, unless $C_z = 1$. Once branching in gamete mass occurs, the smaller gametes (e.g. microgametes) again experience a strong selective pressure to increase their fertilization rate, despite the high costs imposed by fertilization; sexual conflict drives the population towards obligate oogamy imposed by motile microgametes. Thus as for $C_z < 1$, there is always a subset of initial conditions that lead to trajectories along the $m \approx \beta/4$ manifold (most obviously initial conditions on the manifold itself) and thus oogamy remains one of the two evolutionary outcomes, albeit requiring an increasingly small and biologically unrealistic set of initial conditions (see Figure C.11 below).

Figure C.11: Numerical illustration of evolutionary branching for the case where $C_z = 0.9$ and $C_p = 0$. All other parameters the same as Figure 2, except $(m(0), \alpha(0)) = (0.6, 0.5)$ and run for $3000/\mu$ generations.

Appendix D. $\alpha_{\mathbf{max}}$ below which branching to anisogamy can be arrested

The two necessary conditions for evolutionary branching to occur is for the fitness gradient of an invading mutant to be zero (i.e. $dm/d\tau$ in Eq. (C.1) to equal zero if the mutation occurs in mass, this is also known as an evolutionary singularity) and the second derivative of Eq. (A.11) with respect to the mutational stepsize δm to be positive [68, 69] when $\delta m = 0$. Using Eq. (C.1), we see that the vertical manifold $m \approx \beta/4$ is an approximate evolutionary singularity since $[dm/d\tau]|_{m=\beta/4} \approx 0$. The condition for selection in mass to be disruptive about an evolutionary singularity is

$$\frac{d}{d\hat{f}}\left[\frac{\mathrm{d}^2\hat{f}'}{\mathrm{d}\,\delta m^2}\right]_{\delta m=0}\Bigg|_{\hat{f}=0} > 0 \qquad (D.1)$$

where $\hat{f}'$ is the frequency of the mutant in the subsequent generation, given by Eq. (A.11). The explicit expression for the left hand side of Eq. (D.1) is given by

$$\frac{d}{d\hat{f}}\left[\frac{\mathrm{d}^2\hat{f}'}{\mathrm{d}\,\delta m^2}\right]_{\delta m=0}\Bigg|_{\hat{f}=0} = U/V \qquad (D.2)$$

where

$$
\begin{aligned}
U &= A(-1+C_z)e^{\frac{\beta}{2m}}MT\alpha\left(32m^2 - 12m\beta + \beta^2\right)\\
&\quad +16(-1+C_p)m\left(2m^2 - 4m\beta + \beta^2\right)\\
V &= 16m^4\left((-1+C_p)m + A(-1+C_z)e^{\frac{\beta}{2m}}MT\alpha\right)
\end{aligned}
$$

The first condition for branching to occur in mass i.e $dm/d\tau = 0$ can be calculated straightforwardly by setting Eq. (C.1) to zero and solving for α, which gives

$$\alpha = \frac{4m(C_p - 1)(m - \beta)e^{-\frac{\beta}{2m}}}{AMT(C_z - 1)(4m - \beta)} \qquad (D.3)$$

We then substitute Eq. (D.3) into Eq. (D.2) to get

$$\frac{\partial}{\partial\hat{f}}\left[\frac{\mathrm{d}^2\hat{f}'}{\mathrm{d}\,\delta m^2}\right]_{\delta m=0}\Bigg|_{\hat{f}=0} = \frac{(3\beta - 7m)(\beta - 4m)\beta}{12\beta m^4} \qquad (D.4)$$

which needs to be positive for selection to be disruptive. Eq. (D.4) equals 0 if $m = 3\beta/7$ or $m = \beta/4$ and is positive for $\beta/4 < m < 3\beta/7$. By substituting the boundaries of this interval into Eq. (D.3), we find that it corresponds to

$$\frac{48(1 - C_p)\beta}{35AMT(1 - C_z)e^{\frac{7}{6}}} < \alpha_{\mathrm{max}} < \infty \qquad (D.5)$$

and thus the value of α_{max} above which branching will occur in mass is the value given on the left hand side of the inequality Eq. (D.5). In other words, Eq. (D.5) is the interval in α_{max} in which branching is expected to occur. The vertical line in Figure 4 corresponds to the lower limit of this interval. In Figure D.12 below, we provide a numerical example showing how isogamy can be stabilised below a sufficiently low α_{max}.

Figure D.12: Numerical illustration of the stabilisation of isogamy below a sufficiently low α_{max}. System parameters are $A = 100$, $M = 1$, $T = 0.1$, $C_z = 0.6$, $C_p = 0$, $\beta = 1$, $\alpha_{max} = 0.1$ and simulation parameters are $\delta = 5 \times 10^{-3}$, $f_0 = 2 \times 10^{-3}$ and run for 6×10^6 generations. Using Eq. (D.5) we can calculate that branching would occur if $\alpha_{max} \gtrsim 0.1068$ for these parameter values.

Appendix E. The stabilisation of Anisogamy under high costs of parthenogenesis relative to fertilization

In Figure E.13, we provide an example of a parameter regime where pseudooogamy occurs. When the parameters are close to the boundary where we observe the transition between oogamy and anisogamy (i.e close to where the inequality in Eq. (C.15) becomes an equality), we observe pseudooogamy, where there is a considerably stronger selection pressure for microgametes to increase their fertilization rate than macrogametes. Under pseudooogamy, macrogametes are still motile but are considerably less motile than microgametes.

Figure E.13: Numerical illustration of the evolution of pseudo-oogamy in fixed environment. Parameters are $C_z = 0$, $C_p = 0$, $\delta m = 0.005$, $(m(0), \alpha(0)) = (0.25, 0)$ and simulation run for 1.6×10^7 generations. Remaining parameters are as given in Appendix G.

Appendix F. Evolutionary dynamics: switching environments with bet-hedging

We first tackle the derivation of the approximate dynamics in the FRTI (fast relative to invasion) switching regime in Appendix F.1, before verifying the qualitative robustness of these results in the FRTE switching regime in Appendix F.1.

Appendix F.1. Derivation of evolutionary ODEs: FRTI

$$
\begin{aligned}
\frac{\mathrm{d}\hat{f}_m}{\mathrm{d}t_g} &\approx \Big[P_1 h_m(m, \alpha, \beta_1, C_z, C_p) \\
&\quad + P_2 h_m(m, \alpha, \beta_2, C_z, C_p) \Big] \hat{f}_m (1 - \hat{f}_m) \\
\frac{\mathrm{d}\hat{f}_\alpha}{\mathrm{d}t_g} &\approx \Big[P_1 h_\alpha(m, \alpha, \beta_1, C_z, C_p) \\
&\quad + P_2 h_\alpha(m, \alpha, \beta_2, C_z, C_p) \Big] \hat{f}_\alpha (1 - \hat{f}_\alpha)
\end{aligned} \tag{F.1}
$$

where $h_m(m, \alpha, \beta, C_z, C_p)$ and $h_\alpha(m, \alpha, \beta, C_z, C_p)$ are given in Eq. (B.1) and Eq. (B.2) respectively. In Figure F.14 we show that this indeed is a good approximation of the dynamics when δm and $\delta \alpha$ are small and when the residency times in each environment are small relative to the invasion time. Note that as in the case of the fixed environment, the functional form of f_i in these equations implies that trait substitution occurs for independent mutations on the gamete mass and fertilization rate.

We now apply the same approach as in Appendix C.1 (see Eq. (C.1)) to obtain

$$
\begin{aligned}
\frac{\mathrm{d}m}{\mathrm{d}\tau} &= \Big[P_1 h_m(m, \alpha, \beta_1, C_z, C_p) \\
&\quad + P_2 h_m(m, \alpha, \beta_2, C_z, C_p) \Big] \frac{\mathrm{d}}{\mathrm{d}\hat{f}_m} \left(\hat{f}_m (1 - \hat{f}_m) \right) \Big|_{\hat{f}_m = 0} \\
&= P_1 H_m(m, \alpha, \beta_1, C_z, C_p) + P_2 H_m(m, \alpha, \beta_2, C_z, C_p) \\
\frac{\mathrm{d}\alpha}{\mathrm{d}\tau} &= \Big[P_1 h_\alpha(m, \alpha, \beta_1, C_z, C_p) \\
&\quad + P_2 h_\alpha(m, \alpha, \beta_2, C_z, C_p) \Big] \frac{\mathrm{d}}{\mathrm{d}\hat{f}_\alpha} \left(\hat{f}_\alpha (1 - \hat{f}_\alpha) \right) \Big|_{\hat{f}_\alpha = 0} \\
&= P_1 H_\alpha(m, \alpha, \beta_1, C_z, C_p) + P_2 H_\alpha(m, \alpha, \beta_2, C_z, C_p).
\end{aligned} \tag{F.2}
$$

We see in Figure 6 that these also provide a good approximation of the evolutionary dynamics.

Appendix F.2. Derivation of evolutionary ODEs: FRTE

In the FRTI scenario in the previous section, we supposed that switching between the environments was

Figure F.14: Invasion dynamics for a population undergoing bet-hedging when the environment switches FRTI. Blue curve is the analytical approximation using Eq. (F.1) and jagged curve is the numerical simulation. Mutation occurred in mass with $\delta = 0.005$, $f_0 = 0.002$, $(m(0), \alpha(0)) = (0.3, 0.1)$, $\lambda_{1 \to 2} = 13/222$, $\lambda_{2 \to 1} = 1/6$. All other parameters the same as Figure 6.

happening sufficiently regularly relative to the timescale of invasion that the effective invasion dynamics could be described by a weighted mean of the invasion dynamics in both environments (see Eq. (F.1)). Applying an analogous logic, we now assume in the FRTE scenario that switching between the environments occurs regularly relative to the timescale of evolution (the arrival rate of new mutations) that the effective evolutionary dynamics can be described by a weighted mean of the evolutionary dynamics in both environments; that is

$$
\begin{aligned}
\frac{\mathrm{d}m}{\mathrm{d}\tau} &= P_1 H_m(m, \alpha, \beta_1, C_z, C_p) + P_2 H_m(m, \alpha, \beta_2, C_z, C_p) \\
\frac{\mathrm{d}\alpha}{\mathrm{d}\tau} &= P_1 H_\alpha(m, \alpha, \beta_1, C_z, C_p) + P_2 H_\alpha(m, \alpha, \beta_2, C_z, C_p).
\end{aligned} \tag{F.3}
$$

We note that these are in fact exactly the same evolutionary dynamics as derived in the FRTI scenario (see Eq. (F.2)). In Figure 6 we show that these do indeed remain a good approximation for the dynamics in the FRTE scenario. The difference between the dynamics in both regimes is quantitative, rather than qualitative. In the very-fast switching FRTI regime, the populations follow the effective dynamics very closely. In the comparatively slower FRTE regime, although the populations no longer follow the dynamics as well, the qualitative picture of the dynamics is still captured by (D.3). In particular, we still observe an evolutionary stable state of intermediate fertilization rate (see Figure 6).

We begin, as in Appendix C.2, by considering the evolutionary behaviour of m when the fertilization rate is fixed to zero ($\alpha = 0$). We recall that if the population was fixed in environment 1 or 2 with $\alpha = 0$, the mass of gametes would evolve to $m = \beta_1$ and $m = \beta_2$ respectively. In the switching environment we instead find the bet-hedging strategy

$$m^*_{\alpha=0} = P_1\beta_1 + (1 - P_1)\beta_2 \,. \tag{F.4}$$

Meanwhile the region of the boundary $\alpha = 0$ over which reduced fertilization rates are selected for is given by $[d\alpha/d\tau]|_{\alpha=0} < 0$, or

$$0 > (1 - C_z)\left[P_1 e^{\frac{\beta_1}{2m}} + (1 - P_1)e^{\frac{\beta_2}{2m}}\right] - (1 - C_p)$$

$$\implies C_z > (1 - C_p) - \frac{1}{P_1 e^{\frac{\beta_1}{2m}} + (1 - P_1)e^{\frac{\beta_2}{2m}}} \,. \tag{F.5}$$

If this condition holds for $m = m^*_{\alpha=0}$, then the switching-induced fixed point $(m, \alpha) = (m^*_{\alpha=0}, 0)$ is stable. We next turn to the high fertilization rate attractor, for which $\alpha \to \infty$. In a similar manner to Eq. (C.4) (albeit with $H_m(m, \alpha, \beta, C_z, C_p)$ replaced with $[P_1 H_m(m, \alpha, \beta_1, C_z, C_p) + P_2 H_m(m, \alpha, \beta_2, C_z, C_p)]$), we find the bet-hedging strategy

$$m^* = \frac{1}{4}\left[P_1\beta_1 + (1 - P_1)\beta_2\right] \,. \tag{F.6}$$

However, unlike in the fixed environment case, $(m, \alpha) = (m^*, \infty)$ is not always an evolutionary attractor.

In the switching environment, a third evolutionary attractor can emerge, brought about by a balance between selection for high fertilization rates in one environment and zero fertilization rates in the other. Solving $P_1 H_\alpha(m, \alpha, \beta_1, C_z, C_p) + P_2 H_\alpha(m, \alpha, \beta_2, C_z, C_p) = 0$ for α we obtain

$$\alpha^* = \frac{m^*(1 - C_p)}{AMT(1 - C_z)}$$

$$\times \left(\frac{(1 - C_z)\left(P_1 e^{\frac{\beta_1}{2m^*}} + (1 - P_1)e^{\frac{\beta_2}{2m^*}}\right) - (1 - C_p)}{(1 - C_p)(P_1 e^{\frac{\beta_2}{2m^*}} + (1 - P_1)e^{\frac{\beta_1}{2m^*}}) - (1 - C_z)e^{\frac{\beta_1+\beta_2}{2m^*}}}\right) \tag{F.7}$$

A good approximation for m^* in this equation can be deduced by noting that the fixed point sits on a vertical manifold of trajectories along which m is approximately held constant as $\alpha \to \infty$; thus we substitute $\approx m^*$ from Eq. (F.6) to obtain the approximate expression for the

Figure F.15: Numerical simulation showing an absence of branching for a system undergoing bet-hedging in an environment that switches FRTI. Parameters are same as Figure 6 (a) and system run for $4000/\mu$ generations.

attractor (m^*, α^*) given in Eq. (14). In Appendix F.5, we verify from simulations that evolutionary branching does not occur at this switching-induced fixed point, and that the population is instead held in a state of isogamy.

Appendix F.4. Implementation of simulations

In the bet-hedging scenario, we simulate the evolutionary dynamics by implementing a Gillespie algorithm. In particular, we introduce mutations and environmental switching events randomly with geometrically distributed waiting times, where the waiting time is measured in units of number of fertilization processes t_g. To simulate the FRTI regime, we set the environmental switching rates to larger values than the mutation rate i.e. $\lambda_{2\to1}, \lambda_{1\to2} \gg \mu$. Likewise, in the FRTE regime, we set $\lambda_{2\to1}$ and $\lambda_{1\to2}$ to smaller values than μ. The mutation rate in the numerical simulation of the FRTI regime overlaid in Figure 6 (a) is $\mu = 3.5 \times 10^{-4}$ and the switching rates are $\lambda_{2\to1} = 67/532$ and $\lambda_{1\to2} = 1/4$. For Figure 6 (b) we have $\mu = 3.5 \times 10^{-4}$ and the switching rates are $\lambda_{2\to1} = 1/6$ and $\lambda_{1\to2} = 13/222$. For the FRTE regime, $\mu = 5 \times 10^{-4}$ and the switching rates are $\lambda_{2\to1} = (67\mu)/532$ and $\lambda_{1\to2} = \mu/4$ in Figure 6 (a) and $\lambda_{2\to1} = \mu/6$ and $\lambda_{1\to2} = (13\mu)/222$ in Figure 6 (b).

Appendix F.5. Absence of Evolutionary Branching at Switching-Induced fixed point

In this section we present additional numerical results that confirm that evolutionary branching does not occur at the switching-induced fixed point calculated in Eq. (14). We see that under both FRTI and FRTE, the population is held in a state of isogamy. This behaviour is illustrated in Figure F.15.

Figure F.16: Numerical illustration showing an absence of evolutionary branching for a system undergoing bet-hedging in an environment that switches FRTE. Parameters the same as Figure 6 (c) and the system is run for $4000/\mu$ generations.

Appendix G. Model parameters in figures

Appendix G.1. Figure 2

The initial points of the trajectories are $(m(0), \alpha(0)) = (1.5, 0.6)$ and $(m(0), \alpha(0)) = (2, 0.1)$. Remaining model parameters are $A = 100$, $M = 1$, $T = 1$ and $\beta = 1$. Simulation parameters: Initial frequency of novel mutant genotype $f_0 = 2 \times 10^{-3}$, mutation rate $\mu = 5 \times 10^{-4}$ (number of generations)$^{-1}$, run for 1.1×10^7 generations in panel (a) and 1.24×10^7 generations in panel (b). The mutational stepsize is $\delta = 5 \times 10^{-3}$. Throughout the paper, we use δm and $\delta \alpha$ to denote mutational stepsize in m and α respectively, however in simulations involving co-evolution of m and α, the same stepsize is used for both traits. For convenience, we thus denote mutational stepsize as δ in these simulations.

Acknowledgements

Ph.D. studentship (X.L.) is funded by the University of York though the EPSRC DTP, Grant No. EP/V52010X/1.

Code availability statement

Code for running all simulations in the manuscript is provided at https://github.com/phil-liu1020/ParthEvolOogamy/tree/main/Github_submission JTB, and is described in Appendix C.3 and Appendix F.4

References

[1] R. F. Hoekstra, The evolution of sexes, in: S. C. Stearns (Ed.), The Evolution of Sex and Its Consequences, Birkhaeuser Verlag, 1987.

[2] J. Umen, S. Coelho, Algal sex determination and the evolution of anisogamy, Annual review of microbiology 73 (2019) 267–291.

[3] D. L. Kirk, Oogamy: inventing the sexes, Current Biology 16 (24) (2006) R1028–R1030.

[4] H. Matsuda, P. A. Abrams, Why are equally sized gametes so rare? the instability of isogamy and the cost of anisogamyunopened xml element'f'closed, Evolutionary Ecology Research 1 (7) (1999) 769–784.

[5] E. R. Hanschen, M. D. Herron, J. J. Wiens, H. Nozaki, R. E. Michod, Multicellularity drives the evolution of sexual traits, The American Naturalist 192 (3) (2018) E93–E105.

[6] G. Bell, The evolution of anisogamy, Journal of Theoretical Biology 73 (2) (1978) 247–270.

[7] B. P. Nieuwenhuis, D. Aanen, Sexual selection in fungi, Journal of evolutionary biology 25 (12) (2012) 2397–2411.

[8] R. Innami, S. Miyamura, T. Okoshi, T. Nagumo, K. Ichihara, T. Yamazaki, S. Kawano, Gamete dimorphism of the isogamous green alga (chlamydomonas reinhardtii), is regulated by the mating type-determining gene, mid, Communications Biology 5 (1) (2022) 1333.

[9] Y. Krumbeck, G. W. A. Constable, T. Rogers, Fitness differences suppress the number of mating types in evolving isogamous species, Royal Society Open Science 7 (2020) 192126.

[10] T. Hamaji, Y. Mogi, P. J. Ferris, T. Mori, S. Miyagishima, Y. Kabeya, Y. Nishimura, A. Toyoda, H. Noguchi, A. Fujiyama, et al., Sequence of the gonium pectorale mating locus reveals a complex and dynamic history of changes in volvocine algal mating haplotypes, G3: Genes, Genomes, Genetics 6 (5) (2016) 1179–1189.

[11] H. Kalmus, Über den erhaltungswert der phänotypischen (morphologischen) anisogamie und die entstehung der ersten geschlechtsunterschiede, Biol Zentralbl 52 (1932) 716–736.

[12] H. Kalmus, C. Smith, Evolutionary origin of sexual differentiation and the sex-ratio, Nature 186 (1960) 1004–1006.

[13] F. M. Scudo, The adaptive value of sexual dimorphism: I, anisogamy, Evolution (1967) 285–291.

[14] G. A. Parker, R. R. Baker, V. Smith, The origin and evolution of gamete dimorphism and the male-female phenomenon, Journal of Theoretical Biology 36 (3) (1972) 529–553.

[15] T. Togashi, J. Bartelt, J. Yoshimura, K. Tainaka, P. Cox, Evolutionary trajectories explain the diversified evolution of isogamy and anisogamy in marine green algae, Proc. Natl. Acad. Sci. 109 (2012) 13692–13697.

[16] J. Lehtonen, The legacy of parker, baker and smith 1972: Gamete competition, the evolution of anisogamy and model robustness, Cells 10 (3) (2021) 573.

[17] G. A. Parker, How soon hath time... a history of two "seminal" publications, Cells 10 (2) (2021) 287.

[18] R. R. Vance, On reproductive strategies in marine benthic invertebrates, The American Naturalist 107 (955) (1973) 339–352.

[19] M. G. Bulmer, G. A. Parker, The evolution of anisogamy: a game-theoretic approach, Proc. R. Soc. Lond. B. 269 (2002) 2381–2388.

[20] B. Charlesworth, The population genetics of anisogamy, Journal of theoretical biology 73 (2) (1978) 347–357.

[21] P. Iyer, J. Roughgarden, Gametic conflict versus contact in the evolution of anisogamy, Theoretical population biology 73 (4) (2008) 461–472.

[22] J. M. Henshaw, M. Bittlingmaier, L. Schärer, Hermaphroditic origins of anisogamy, Philosophical Transactions of the Royal Society B 378 (1876) (2023) 20220283.

[23] É. Kisdi, A. Stefan, H. Geritz, Adaptive dynamics: a framework to model evolution in the ecological theatre, Journal of mathematical biology 61 (1) (2010) 165.

[24] N. Maire, M. Ackermann1, M. Doebeli, Evolutionary branching and the evolution of anisogamy, Selection 2 (1-2) (2002) 119–131.

[25] J. Lehtonen, H. Kokko, Two roads to two sexes: unifying gamete competition and gamete limitation in a single model of anisogamy evolution, Behav. Ecol. Sociobiol. 65 (2011) 445–459. doi:10.1007/s00265-010-1116-8.
URL https://doi.org/10.1007/s00265-010-1116-8

[26] C. M. Lessells, R. R. Snook, D. J. Hosken, The evolutionary origin and maintenance of sperm: selection for a small, motile gamete mating type, in: Sperm biology, Elsevier, 2009, pp. 43–67.

[27] J. da Silva, The evolution of sexes: A specific test of the disruptive selection theory, Ecology and evolution 8 (1) (2018) 207–219.

[28] M. T. Ghiselin, The economy of nature and the evolution of sex, Journal of the History of Biology 9 (2) (1976).

[29] R. F. Hoekstra, Evolution of gamete motility differences ii. interaction with the evolution of anisogamy, Journal of theoretical Biology 107 (1) (1984) 71–83.

[30] Z. Hadjivasiliou, Y. Iwasa, A. Pomiankowski, Cell–cell signalling in sexual chemotaxis: a basis for gametic differentiation, mating types and sexes, Journal of the Royal Society Interface 12 (109) (2015) 20150342.

[31] D. B. Dusenbery, Selection for high gamete encounter rates explains the success of male and female mating types, Journal of Theoretical Biology 202 (1) (2000) 1–10.

[32] P. A. Cox, J. A. Sethian, Gamete motion, search, and the evolution of anisogamy, oogamy, and chemotaxis, The American Naturalist 125 (1) (1985) 74–101.

[33] C. D. Soulsbury, S. Humphries, Biophysical determinants and constraints on sperm swimming velocity, Cells 11 (21) (2022) 3360.

[34] J. Lehtonen, G. A. Parker, Evolution of the two sexes under internal fertilization and alternative evolutionary pathways, The American Naturalist 193 (5) (2019) 702–716.

[35] H. Nozaki, T. K. Yamada, F. Takahashi, R. Matsuzaki, T. Nakada, New "missing link" genus of the colonial volvocine green algae gives insights into the evolution of oogamy, BMC Evolutionary Biology 14 (1) (2014) 1–11.

[36] M. Tatewaki, M. Lima, Life histories of blidingia minima (chlorophyceae), especially sexual reproduction 1, Journal of phycology 20 (3) (1984) 368–376.

[37] G. Lokhorst, B. Trask, Taxonomic studies on urospora (acrosiphoniales, chlorophyceae) in western europe, Acta botanica neerlandica 30 (5/6) (1981) 353–431.

[38] N. Kinoshita, C. Nagasato, T. Motomura, Chemotactic movement in sperm of the oogamous brown algae, saccharina japonica and fucus distichus, Protoplasma 254 (1) (2017) 547–555.

[39] J. Li, S. Pang, F. Liu, T. Shan, S. Gao, Spermatozoid life-span of two brown seaweeds, saccharina japonica and undaria pinnatifida, as measured by fertilization efficiency, Chinese journal of oceanology and limnology 31 (4) (2013) 774–781.

[40] J. Lehtonen, Y. Horinouchi, T. Togashi, G. A. Parker, Evolution of anisogamy in organisms with parthenogenetic gametes, The American Naturalist 198 (3) (2021) 360–378.

[41] G. W. Constable, H. Kokko, Parthenogenesis and the evolution of anisogamy, Cells 10 (9) (2021) 2467.

[42] J. Lehtonen, Models of fertilization kinetics, Royal Society Open Science 2 (9) (2015) 150175.

[43] J. Lehtonen, L. Dardare, Mathematical models of fertilization—an eco-evolutionary perspective, The Quarterly Review of Biology 94 (2) (2019) 177–208.

[44] J. Lehtonen, Models of fertilization kinetics, Royal Society Open Science 2 (9) (2015) 150175. doi:10.1098/rsos.150175.

[45] H. Vogel, G. Czihak, P. Chang, W. Wolf, Fertilization kinetics of sea urchin eggs, Mathematical Biosciences 58 (2) (1982) 189–216.

[46] H. Van den Ende, F. Klis, A. Musgrave, The role of flagella in sexual reproduction of chlamydomonas eugametos, Acta botanica neerlandica 37 (3) (1988) 327–350.

[47] L. Mignerot, K. Avia, R. Luthringer, A. P. Lipinska, A. F. Peters, J. M. Cock, S. M. Coelho, A key role for sex chromosomes in the regulation of parthenogenesis in the brown alga ectocarpus, PLoS genetics 15 (6) (2019) e1008211.

[48] D. R. Levitan, Optimal egg size in marine invertebrates: theory and phylogenetic analysis of the critical relationship between egg size and development time in echinoids, The American Naturalist 156 (2) (2000) 175–192.

[49] A. E. Hall, M. D. Rose, Cell fusion in yeast is negatively regulated by components of the cell wall integrity pathway, Molecular biology of the cell 30 (4) (2019) 441–452.

[50] E. Harrison, R. MacLean, V. Koufopanou, A. Burt, Sex drives intracellular conflict in yeast, Journal of Evolutionary Biology 27 (8) (2014) 1757–1763.

[51] L. D. Hurst, W. D. Hamilton, Cytoplasmic fusion and the nature of sexes, Proceedings of the Royal Society of London. Series B: Biological Sciences 247 (1320) (1992) 189–194.

[52] V. Hutson, R. Law, Four steps to two sexes, Proceedings of the Royal Society of London. Series B: Biological Sciences 253 (1336) (1993) 43–51.

[53] L. Adams, K. Lyons, J. Monday, E. Larkin, J. Wyffels, Costs of parthenogenesis on growth and longevity in ex situ zebra sharks stegostoma tigrinum, Endangered Species Research 50 (2023) 81–91.

[54] J. A. Metz, S. A. Geritz, G. Meszéna, F. J. Jacobs, J. S. Van Heerwaarden, Adaptive dynamics: a geometrical study of the consequences of nearly faithful reproduction (1995).

[55] P. G. Hufton, Y. T. Lin, T. Galla, A. J. McKane, Intrinsic noise in systems with switching environments, Physical Review E 93 (5) (2016) 052119.

[56] E. Berríos-Caro, T. Galla, G. W. A. Constable, Switching environments, synchronous sex, and the evolution of mating types, Theoretical population biology 138 (2021) 28–42.

[57] J. da Silva, V. L. Drysdale, Isogamy in large and complex volvocine algae is consistent with the gamete competition theory of the evolution of anisogamy, Proceedings of the Royal Society B 285 (1890) (2018) 20181954.

[58] N. W. Burke, R. Bonduriansky, Sexual conflict, facultative asexuality, and the true paradox of sex, Trends in Ecology & Evolution 32 (9) (2017) 646–652.

[59] J. Lehtonen, H. Kokko, G. A. Parker, What do isogamous organisms teach us about sex and the two sexes?, Phil. Trans. R. Soc. B 371 (2016) 20150532.

[60] J. Lehtonen, M. D. Jennions, H. Kokko, The many costs of sex, Trends in ecology & evolution 27 (3) (2012) 172–178.

[61] J. Heitman, Evolution of sexual reproduction: A view from the fungal kingdom supports an evolutionary epoch with sex before sexes, Fungal Biology Reviews 29 (3) (2015) 108 – 117, special Issue: Fungal sex and mushrooms – A credit to Lorna Casselton. doi:https://doi.org/10.1016/j.fbr.2015.08.002.
URL http://www.sciencedirect.com/science/article/pii/S1749461315000391

[62] Z. Hadjivasiliou, N. Lane, R. Seymour, A. Pomiankowski, Dynamics of mitochondrial inheritance in the evolution of binary mating types and two sexes, Proc. Biol. Sci. 280 (1769) (2013) 20131920.

[63] S. Billiard, M. López-Villavicencio, M. Hood, T. Giraud, Sex, outcrossing and mating types: unsolved questions in fungi and beyond, Journal of evolutionary biology 25 (6) (2012) 1020–1038.

[64] S. Billiard, M. López-Villavicencio, B. Devier, M. E. Hood,

C. Fairhead, T. Giraud, Having sex, yes, but with whom? inferences from fungi on the evolution of anisogamy and mating types, Biological reviews 86 (2) (2011) 421–442.

[65] G. W. A. Constable, H. Kokko, The rate of facultative sex governs the number of expected mating types in isogamous species, Nat. Ecol. Evol. 2 (7) (2018) 1168–1175.

[66] P. Czuppon, G. W. A. Constable, Invasion and extinction dynamics of mating types under facultative sexual reproduction, Genetics 213 (2) (2019) 567–580.

[67] G. A. Parker, R. Baker, V. G. Smith, The origin and evolution of gamete dimorphism and the male-female phenomenon, J. Theor. Biol. 36 (3) (1972) 529–553.

[68] F. Dercole, F. Della Rossa, P. Landi, The transition from evolutionary stability to branching: A catastrophic evolutionary shift, Scientific reports 6 (1) (2016) 1–8.

[69] S. A. Geritz, E. Kisdi, G. Meszéna, J. A. Metz, Evolutionary singular strategies and the adaptive growth and branching of the evolutionary tree (1996).

23