1 *Alternative splicing of a potato disease resistance gene maintains homeostasis between development*

2 *and immunity*

4 Biying Sun[1,2], Jie Huang[1,2,6], Liang Kong[1,2], Chuyun Gao[1,2], Fei Zhao[1], Jiayong Shen[1], Tian Wang[1],

5 kangping Li[1], Luyao Wang[1,5], Yuanchao Wang[1,2], Dennis A. Halterman[3,4,*], Suomeng Dong[1,2,*]

[1]Department of Plant Pathology, Nanjing Agricultural University, Nanjing 210095, China.

[2]The Key Laboratory of Integrated Management of Crop Diseases and Pests (Ministry of Education),

Nanjing Agricultural University, Nanjing 210095, China.

[3]Department of Plant Pathology, University of Wisconsin-Madison, Madison, Wisconsin, United States of

America.

[4]U.S. Department of Agriculture-Agricultural Research Service, Vegetable Crops Research Unit, Madison,

Wisconsin, United States of America

[5]Shenzhen Branch, Guangdong Laboratory of Lingnan Modern Agriculture, Genome Analysis Laboratory

of the Ministry of Agriculture and Rural Affairs, Agricultural Genomics Institute at Shenzhen, Chinese

Academy of Agricultural Sciences, Shenzhen, Guangdong, 518120, China.

[6]Plant Chemetics Laboratory, Department of Biology, University of Oxford, Oxford, UK

*Address correspondence to: smdong@njau.edu.cn, dennis.halterman@usda.gov

Short title:

Resistance gene alternative splicing regulates development and immunity homeostasis

One-sentence summary:

Potato resistance gene *RB* balances plant growth and immunity through AS (alternative splicing), while

pathogen-secreted effector IPI-O1 mediates AS of *RB* by targeting the conserved splicing factor

StCWC15, further increasing the *RB_CDS* expression level to activate immunity.

Abstract

Plants possess a robust and sophisticated innate immune system against pathogens. The intracellular receptors with nucleotide-binding, leucine-rich repeat (NLR) motifs recognize pathogen-derived effector proteins to trigger the immune response. To balance plant growth and rapid pathogen detection, NLR expression is precisely controlled in multifaceted ways. The role of post-transcriptional processing of NLRs, particularly alternative splicing (AS) of introns, in response to infection is recurrently observed but poorly understood. Here we report that the potato NLR gene *RB* undergoes AS of its intron, resulting in two transcriptional isoforms, which coordinately regulate plant immunity and growth homeostasis. During normal growth, *RB* predominantly exists as intron-retained isoform *RB_IR*, encoding a truncated protein containing only the N-terminus of the NLR. Upon late blight infection, the causal pathogen *Phytophthora infestans* induces intron splicing of *RB*, increasing the abundance of *RB_CDS*, which encodes a full-length, and active R protein. By deploying the *RB* splicing isoforms fused with a *luciferase* reporter system, we identified IPI-O1 (also known as Avrblb1), the RB cognate effector, as a facilitator of *RB* AS. Importantly, IPI-O1 directly interacts with potato splicing factor StCWC15 to promote *RB* splicing for activation of *RB*-mediated resistance. Thus, our study reveals that StCWC15 serves as a surveillance facilitator sensing the pathogen-secreted effector, and regulates the trade-off between *RB*-mediated plant immunity and growth, expanding our understanding of molecular plant-microbe interactions.

Keywords: Potato late blight; resistance gene; RB; IPI-O1; CWC15

Introduction

Upon pathogen infection, plants have evolved two major mechanisms for recognition of pathogen effectors and elicitation of plant defense (Jones and Dangl, 2006). One utilizes plasma membrane-localized pattern-recognition receptors (PRRs) to perceive pathogen-associated molecular patterns (PAMPs), leading to PAMP-triggered immunity (PTI) (Couto and Zipfel, 2016; Yu et al., 2017; Zhou and Zhang, 2020). The other uses intracellular NLR immune receptors to detect cytoplasmic effectors introduced by the pathogen, thereby activating effector-triggered immunity (ETI; Jones and Dangl, 2006; Dodds and Rathjen, 2010; Cui et al., 2015). During ETI, NLRs oligomerize to form "resistosome" structures (Wang et al., 2019; Ofir et al., 2021) and the response culminates in programmed cell death,

known as the hypersensitive response (HR), to prevent pathogen spread. The enormous diversity of *NLR* genes in plants demonstrates their important role in plant immunity (Li et al., 2015; Van de Weyer et al., 2019).

Late blight, caused by oomycete *Phytophthora infestans*, is the most devastating disease in potato production (Haverkort et al., 2009; Kamoun et al., 2015; Sun et al., 2022). Deployment of host resistance (*R*) genes is a crucial strategy for controlling the disease and many potato cultivars have been bred to introduce resistance genes against *P. infestans* (*Rpi* genes) that originate from potato wild relatives (Kamoun and Smart, 2005; Karki et al., 2021). More than 70 *Rpi* genes have been identified and mapped in 32 *Solanum* species (Paluchowska et al., 2022). Among these *R* genes, *RB* was cloned from the wild diploid potato species *Solanum bulbocastanum*, which is known for its durable late blight resistance (Song et al., 2003; van der Vossen et al., 2003). Like most plant *R* genes, *RB* encodes an NLR receptor which recognizes *P. infestans* effector IPI-O1 (also known as Avrblb1) to activate immune responses (Vleeshouwers et al., 2008; Champouret et al., 2009). Previous studies hypothesized that IPI-O1 might change the conformation of RB to facilitate the formation a functional multimeric complex, thereby activating *RB*-mediated disease resistance (Chen et al., 2012; Zhao and Song, 2021). Consequently, for *RB* with a durable-resistance properties, the functions of *RB*-mediated plant immunity need to be further determined. Despite their usefulness in activating resistance, constitutive expression of *NLR* genes causes auto-immunity, leading to a negative effect on plant growth and development (Cheng et al., 2011; Deng et al., 2019; Cui et al., 2020). Therefore, the expression of *NLR* genes needs to be precisely regulated to maintain homeostasis between growth and immunity. The regulatory mechanisms for the expression of NLRs are mainly via transcription regulation, RNA processing, post-translational modification and proteasome-mediated degradation (Li et al., 2015). Intron splicing is common in post-transcriptional RNA processing and is executed by a supramolecular ribonucleoprotein (RNP) complex known as the spliceosome (Reddy, 2007). Intron retention during RNA processing is a form of alternative splicing (AS) and can lead to changes in the translated protein. AS contributes to transcriptome and proteome diversity in animals and plants and it has been shown to play an important role in regulating plant immunity (Reddy et al., 2013a; Wu et al., 2016; Bazin et al., 2020; Zhang et al., 2022). Several *NLR* genes have been demonstrated to undergo AS, suggesting that this may be an important mechanism for regulation of this important class of immunity-related genes (Dinesh-Kumar and Baker, 2000; Halterman et al., 2003; Halterman and Wise, 2006; Xu et al., 2012; Sanchez-Martin et al., 2021). However, the molecular

mechanisms that control *NLR* gene AS and the significance of AS in maintaining plant growth and
immunity homeostasis is incompletely understood.

In this work, we report that the potato resistance gene *RB* undergoes alternative splicing, resulting in
generation of two transcripts *RB_IR* and *RB_CDS*. Removal of the intron results in intact RB protein
(RB_CDS) that confers resistance to *P. infestans*, while intron retention results in a truncated RB protein
(RB_IR) that cannot protect against the pathogen. Upon late blight infection, *P. infestans* is able to
modulate the AS of *RB*, increasing the abundance of *RB_CDS* to activate *RB*-mediated immunity.
Interaction of *P. infestans* effector IPI-O1 with potato splicing factor CWC15 in the nucleus leads to an
increase of *RB* intron splicing during infection. This process balances plant growth and defense and
provides a unique molecular basis for NLR recognition of pathogen effectors.

Results
The AS of *RB* is specifically modulated by *P. infestans* infection
The *RB* gene encompasses a single 679-nucleotide (nt) intron initiating 428 nt downstream of the
translation start codon and terminating at 1107 nt (Fig. 1A). We performed rapid amplification of cDNA
ends (5' RACE) using total RNA from non-inoculated *RB* transgenic potato cv. Desiree expressing a
single genomic copy of *RB* with its native promoter (Zhu et al., 2015), the transcriptional isoforms were
identified that correspond to the transcripts with the intron spliced (*RB_CDS*) or retained (*RB-IR*) (Fig.
1A). The existence of two isoforms, *RB_CDS* and *RB_IR*, were further validated by reverse transcription
PCR (RT-PCR) using *RB* transgenic potato and *Nicotiana benthamiana* (Fig. 1B). Furthermore, the two
transcripts caused by alternative intron splicing were also detected in RNA from the diploid potato species
Solanum bulbocastanum clone PT29, where *RB* was identified (Fig. 1B) (Song et al., 2003). *Rpi-sto1*, an
RB-homologous gene from *S. stoloniferum*, also showed the same two transcripts detected by RT-PCR in
Rpi-sto1 transgenic potato (Fig. 1B) (van der Vossen et al., 2003; Wang et al., 2008). Splicing of introns
in pre-mRNA is directed by spliceosome targeting of particular sequences at the intron/exon junctions
called splice sites (Wan et al., 2020), which are commonly found as a conserved GT dinucleotide at the 5'
splice site and a conserved AG dinucleotide at the 3' splice site (Sheth et al., 2006). We generated *RB*
splicing mutant *RB_IVS* (IVS=intervening sequence), with synonymously mutated splice sites in *RB* from
GT to GA (5') and from AG to CG (3') (Fig. 1A), and transformed it into potato and *N. benthamiana*. We
found that *RB_IVS* is unable to undergo intron splicing in both transgenic potato and *N. benthamiana* (Fig.

1B). Among the two transcripts of *RB*, *RB_CDS* is predicted to encode the entire NLR protein, while *RB_IR* encodes a truncated protein containing only the N-terminus of the NLR protein, due to a premature termination codon (PTC) in the intron (Fig. 1A). As expected, immunoblot analysis showed that RB fused with green florescent protein at the N-terminus (GFP-RB) produced two bands corresponding to RB_CDS and RB_IR, respectively (Fig. 1C). Together, the data support that *RB* undergoes AS to produce two different transcriptional isoforms that each encode different size proteins.

To investigate the dynamic change of *RB* splicing during pathogen infection, we inoculated *RB* transgenic potato with *P. infestans* strain JH19 and measured the expression levels of different *RB* isoforms compared to levels following water (mock) inoculation. Quantitative RT-PCR data demonstrated that the splicing ratio (transcript level of the spliced isoform divided by the transcript level of the unspliced isoform) of *RB (RB_CDS/RB_IR)* significantly increased to 1.66 at 1 dpi and 2.78 at 2 dpi following *P. infestans* inoculation, indicating an increase in *RB_CDS* expression and a decrease of *RB_IR* expression during *P. infestans* infection (Fig. 1D and Supplemental Fig. S1A). Conversely, inoculation of *RB* transgenic potato with zoospores of *Phytophthora capsici* isolate LT1534 had no significant impact on *RB* splicing ratio alteration (Fig. 1D and Supplemental Fig. S1A). Following these experiments, the total gene expression of *RB* did not change significantly in either *P. infestans*-, *P. capsici*-, or mock-inoculated plants (Supplemental Fig. S1B). Previously, our results revealed that both the intron-spliced *RB_CDS* and intron-retained *RB_IR* can encode proteins when transiently expressed in *N. benthamiana* (Fig. 1C). A change in abundance of the two proteins was also observed during *P. infestans* infection. The abundance of the RB_CDS protein increased during *P. infestans* infection compared to mock treatment, whereas the accumulation of the truncated RB_IR decreased. However, no considerable change in protein expression level was noted during *P. capsici* infection (Supplemental Fig. S1C). These results indicate that the *RB* gene produces more RB_CDS due to increased intron splicing subsequent to *P. infestans* inoculation. To further verify the effect of *P. infestans* on *RB* AS, we tested multiple *P. infestans* genotypes. Genotype T30-4 and HB1501 had been sequenced and the presence of IPI-O1 had been reported (Thilliez et al., 2019; Zhang et al., 2021). The presence of IPI-O1 in genotype JH19 was also verified by PCR. The RT-PCR results show that the relative expression intensity of the *RB_IR* isoform was reduced at 1 and 2 dpi, whereas the relative expression intensity of the *RB_CDS* isoform increased during the same timeframe, regardless of the pathogen genotype (Fig. 1E). The expression intensity remained unchanged in *P. capsici* inoculated plants and mock treatment (Fig. 1E). Collectively, these data supplemented that the *RB* gene

produces more *RB_CDS* transcript due to increased intron splicing during *P. infestans* infection.

Given that *P. infestans* affects the expression level of *RB_CDS* via modulation of *RB* AS, we

hypothesized that impeding the splicing process would impact *RB_CDS* expression. To test this, we

assessed the *RB* splicing ratio change in splicing mutant *RB_IVS* transgenic potato during *P. infestans*

infection. We found that the splicing ratio in this transgenic plant remained low and did not change

significantly during infection, indicating that *RB* is maintained with the intron present (Fig. 1F and

Supplemental Fig. S1D). In eukaryotes, the splicing process is executed by the spliceosome, a conserved

ribonucleoprotein complex dynamically composed of various proteins (Chen and Manley, 2009). Among

these components, U1 small nuclear ribonucleoprotein particle 70k (U1-70k) is crucial for binding the

5'splice site in pre-mRNA (Cho et al., 2011), and splicing factor 3b subunit 1(SF3b1) as the core

component of the U2 snRNP, plays a pivotal role in branch site recognition (Will and Lührmann, 2011).

Thus, we silenced these targets to verify the effect on splicing of *RB* transcripts. Additionally, as a control,

we silenced the RNA-binding protein *GRP7*, known to regulate RNA processing, thereby affecting the

PTI response and specifically influencing NLR protein Gpa2/Rx1-mediated immunity (Wang et al., 2020;

Sukarta et al., 2022). The quantitative PCR results indicated that silencing of *U1-70k* and *SF3B1* both

notably suppressed up-regulation of the *RB* splicing ratio following *P. infestans* infection compared to

non-inoculated samples, whereas *GRP7* silencing did not affect the induction of *RB* splicing following

infection (Fig. 1G and Supplemental Fig. S1, E and F). We also conducted a chemical splicing inhibitor

assay using Pladienolide B (PB), a splicing modulator that targets SF3B1 and disrupts its proper

recognition of the intron (Kotake et al., 2007; Cretu et al., 2018). The transcript expression analysis

indicated that 30 μM PB effectively suppressed the splicing ratio increase during infection (Fig. 1H and

Supplemental Fig. S1G). In conclusion, our results adequately demonstrated that the *RB_CDS* transcript

was most prevalent following *P. infestans* infection, which is the consequence of an increase in intron

splicing.

Intron splicing of *RB* is necessary for the activation of resistance and functions independent of ETI

In light of the observed change in the *RB* splicing ratio during *P. infestans* infection, we aimed to

characterize the function of different transcripts of *RB* in *RB*-mediated resistance. We transformed

RB_CDS, *RB_IR*, and the splicing mutant *RB_IVS* into potato cultivar Favorita. The wild-type *RB* gene

was also transformed into Favorita as a control. Multiple potato plants expressing the intron-spliced

transcript *RB_CDS* displayed resistance to *P. infestans*, similar to the wild-type *RB* transgenic potato. In

contrast, plants expressing the intron-retained transcript *RB_IR* or splicing mutant *RB_IVS* showed severe

disease symptoms (Fig. 2A). The results of detached leaf assays of the transgenic potatoes were consistent

with the whole plant phenotypes (Fig. 2, B and C). Additionally, we generated *RB*, *RB_CDS*, *RB_IR*, and

RB_IVS transgenic *N. benthamiana* for our assays. Notably, only *RB_CDS* limited *P. infestans* lesion

development (Supplemental Fig. S2, A to C). Furthermore, *RB_CDS*, neither *RB_IR* nor *RB_IVS*, induced

HR in *N. benthamiana* in presence of its cognate effector IPI-O1 (Supplemental Fig. S2, D and E), further

supporting that *RB_CDS* is required for *RB*-mediated immune responses. Of note, overexpression of

RB_CDS and RB_IR did not induce HR in *N. benthamiana* (Supplemental Fig. S2, F to H).

Given that only the protein encoded by the intron spliced transcript *RB_CDS* can activate immunity,

we investigated whether interdiction of the splicing process might affect *RB*-mediated resistance to *P.*

infestans. Silencing of *U1-70k, SF3B1*, and *GRP7* in *RB* transgenic *N. benthamiana* rendered *RB*

transgenic *N. benthamiana* susceptible to *P. infestans* compared to the EV control, while silencing *GRP7*

did not affect *RB*-mediated resistance (Fig. 2, D to F). Moreover, treatment with 30 μM PB significantly

reduced *RB*-mediated resistance in *RB* transgenic *N. benthamiana*, but this treatment did not affect the

resistance activation conferred by RB_CDS (Fig. 2, G to I). These results could adequately explain that

the modulation of *RB* pre-mRNA intron splicing is essential for initiating *RB* resistance activation against

P. infestans.

Most *R* genes function by recognizing their corresponding effectors and triggering a HR to prevent

spreading of disease. To test whether disruption of *RB* splicing also interfered with elicitation of the HR,

we co-expressed IPI-O1 and *RB* in *N. benthamiana* in which spliceosome components were silenced.

Additionally, we introduced the intron-containing *R* gene *Rpi-blb2* (van der Vossen et al., 2005) and the

intron-lacking *R* gene *R3a* (Huang et al., 2005) with their corresponding effector genes for HR testing.

Following the silencing of *U1-70k* and *SF3B1*, co-infiltration of the intron-containing potato *NLR* genes

RB or *Rpi-blb2* with their corresponding effectors *IPI-O1* and *Avrblb2*, respectively, resulted in a

significantly weakened HR (Supplemental Fig. S3, A and B). Conversely, co-infiltration of the intronless

NLR genes *RB_CDS* or *R3a* with the corresponding effectors *IPI-O1* and *Avr3a* induced a robust HR

following the silencing of *U1-70k* and *SF3B1* (Supplemental Fig. S3, A and B). Silencing of *GRP7* had no

effect on suppression the HR following co-infiltration of any tested NLRs with their corresponding

effectors (Supplemental Fig. S3, A and B). Furthermore, treatment with 30 μM PB significantly reduced

the incidence of HR upon IPI-O1 infiltration in *RB* transgenic *N. benthamiana*. However, this treatment

did not affect the HR incidence conferred by RB_CDS (Supplemental Fig. S3C). These results suggest

that the splicing plays a vital role in the activity of intron containing NLRs, including *RB* and *Rpi-blb2*.

IPI-O1 mediates *RB* AS during the infection process

To determine the proteins secreted by *P. infestans* that regulate *RB* intron splicing, we designed an

RB alternative splicing reporter system that can directly reflect the extent of *RB* intron splicing. This

system allows the detection of intron splicing modulation without ETI activation. The reporter system is

made up of the AS region of *RB* genomic sequences (from 1 nt to 1987 nt) fused with the *luciferase* (*LUC*)

reporter gene, which is predicted to produce two transcripts (Fig. 3A): the first is *RB_IR-LUC*, which

retains the intron, introduces a premature termination codon, and does not produce a LUC signal; the

second is *RB_CDS-LUC*, in which the intron is spliced and translates a complete RB CC-NB protein and

functional LUC protein. To verify intron splicing changes via the *RB* AS reporter system following *P.*

infestans inoculation, we expressed *RB-LUC* via agrobacterium in *N. benthamiana* and subsequently

inoculated with *P. infestans* JH19. Luciferase activity of *RB-LUC* was measured at 1 dpi and showed a

significant increase following *P. infestans* treatment compared to the water-inoculated (mock). The

luciferase activity of *RB_CDS-LUC* and *LUC* constructs remained unaltered following *P. infestans*

inoculation (Supplemental Fig. S4). This verified the modulation of *RB* intron splicing by *P. infestans*

using the *RB* AS reporter system, demonstrating the feasibility of this system. This supported the use of

the *RB* AS reporter system in screening for *P. infestans* effectors that facilitate *RB* splicing. Indeed, in a

previous study, we identified several splicing regulatory effectors (SRE) using another splicing reporter

system (Huang et al., 2020a). To assess the impact of SRE proteins on gene *RB* AS, we co-expressed

several SRE proteins with the *RB-LUC* reporter system in *N. benthamiana*. Screening results show that

IPI-O1 (SRE8) significantly induced *RB* intron splicing compared to the mock-infiltrated control (Fig.

3B). Infiltration of SRE proteins with *RB_CDS-LUC*, which lacks the intron, did not affect LUC activity

(Fig. 3, B and C).

Classified within the *ipiO* locus in *P. infestans*, IPI-O1 is one of 16 variants in this family. Among

them, IPI-O2 is highly similar to IPI-O1 and also induces HR with *RB*. The most divergent variant IPI-O4,

differs from IPI-O1 by 20 amino acids and evades *RB* recognition (Champouret et al., 2009; Halterman et

al., 2010). To test the ability of other IPI-O family members to modulate *RB* AS, we overexpressed these

proteins in *RB* transgenic *N. benthamiana*. The followed RT-PCR of *RB* illustrated that, like IPI-O1, IPI-O2 induced *RB* intron splicing, but IPI-O4 does not (Fig. 3, D and E). Consistent with screening results by the *RB* AS reporter system, RT-PCR also revealed overexpression of other SRE proteins had no effect on *RB* splicing changes (Supplemental Fig. S5). Overall, it appears that IPI-O1 and its closely related family members can promote *RB* intron splicing, and this modulation correlates with RB-mediated immunity induction.

NLS motif of IPI-O1 is required for inducing *RB* splicing change and immunity activation.

IPI-O1 is a nuclear-localized effector, and has been demonstrated to modulate mRNA splicing (Wang et al., 2018; Huang et al., 2020b). To investigate the correlation between nuclear localization of IPI-O1 and its role in *RB* splicing, we utilized cNLS Mapper to predict the presence of an internal nuclear localization signal (NLS) motif within IPI-O1. Using this information, we generated an IPI-O1 mutant, IPI-O1^{nls}, in which a total of seven charged amino acids in NLS motif, K (lysine) and R (arginine), were mutated to A (alanine), and monitored the subcellular localization of proteins by confocal microscopy (Fig. 4A). We found that IPI-O1^{nls} failed to accumulate in the nucleoplasm and nucleolus like IPI-O1, and instead is localized primarily in the cytoplasm (Fig. 4, B-D). To examine the AS ability of IPI-O1 and the mutant, we expressed IPI-O1 and IPI-O1^{nls} in the *RB-LUC* reporter system. Monitoring luciferase activity showed that IPI-O1, but not IPI-O1^{nls} can significantly increase the luciferase activity (Fig. 4E). Further evaluation of *RB* splicing ratio alterations through RT-qPCR demonstrated that IPI-O1 increased the *RB_CDS*/*RB_IR* splicing ratio, indicating elevated *RB_CDS* expression level and reduced *RB_IR* expression. Conversely, overexpression of IPI-O1^{nls} did not elicit this alteration (Fig. 4F and Supplemental Fig. S6A). To detect the IPI-O1 and IPI-O1^{nls} modulation of RB protein expression levels, we co-expressed IPI-O1 and IPI-O1^{nls} respectively with GFP-RB in *N. benthamiana* for immuno-blot analysis. The data demonstrated an increase in RB_CDS and a decrease in RB_IR only in the presence of IPI-O1, not IPI-O1^{nls}, when compared to GFP-RB expressed alone (Fig. 4G). These results confirmed that the IPI-O1 mutant IPI-O1^{nls} has lost the ability to modulate *RB* AS changes. To further validate the function of IPI-O1 nuclear localization on *RB* AS modulation, we complemented the mutant of the NLS in IPI-O1^{nls} by fusion of a NES or NLS motif to generate NES-IPI-O1^{nls} and IPI-O1^{nls}-NLS mutants (Supplemental Fig. S6B). Microscopy showed that NES-IPI-O1^{nls} and IPI-O1^{nls}-NLS are localized to the cytoplasm and nucleus, respectively (Supplemental Fig. S6C). Nevertheless, overexpression of these

mutants in the *RB* AS reporter system showed that, like IPI-O1[nls], neither NES-IPI-O1[nls] nor IPI-O1[nls]-NLS can promote *RB* intron splicing or increase abundance of *RB_CDS* (Supplemental Fig. S6D). In conclusion, not only is the nuclear localization of IPI-O1 crucial for *RB* AS modulation, but also the 7 amino acids in the NLS motif.

IPI-O1, as the corresponding effector of *RB*, has the capability to induce *RB* intron splicing and activate the HR. To test the impact of IPI-O1[nls] on HR induction with *RB*, we expressed all IPI-O1 mutants in *RB* transgenic *N. benthamiana*. The results showed that IPI-O1 induced the HR in both *RB* and *RB_CDS* transgenic *N. benthamiana*, but not in *RB_IR* plants (Supplemental Fig. S6E). However, the IPI-O1[nls] mutant and its localization complement mutants were unable to induce the HR in *RB* transgenic *N. benthamiana*. Nevertheless, they were still able to elicit the HR in *RB_CDS* transgenic *N. benthamiana*, regardless of their localization to the nucleus or cytoplasm (Supplemental Fig. S6, E and F). These findings collectively suggest that the arginine and lysine residues within the IPI-O1 nuclear localization motif are critical for promoting *RB* intron splicing, which is necessary for immunity activation.

IPI-O1 interacts physically with splicing factor CWC15 in the nucleus

To identify the potential splicing regulators of *RB*, we performed an ULTImate (Hybrigenics) yeast-two hybrid (Y2H) screen to identify IPI-O1-interacting proteins within a *N. benthamiana* cDNA library. A total of 81.6 million interactions were analyzed and 254 clones were processed by sequencing of the prey inserts. Of these 254 clones, 28 contained an insert with homology to the *CWC15* gene from *N. tomentosiformis* (LOC104090615). CWC15 is a core component of the Nineteen complex within the spliceosome (Will and Lührmann, 2011). Y2H assays have verified the interaction of two CWC15 homologs in *N. benthamiana* with IPI-O1 (Supplemental Fig. S7A). Analysis of *CWC15* in potato we found two *CWC15*-related gene sequences in the double-monoploid potato DM-1-3: *Soltu.DM.10G000900.1* (*StCWC15a*) and *Soltu.DM.07G027590.1* (*StCWC15b*), encoding 231 aa (amino acid) and 228 aa proteins respectively. StCWC15a and StCWC15b share 93% similarity at the amino acid levels (Supplemental Fig. S7B). Further analysis of CWC15 sequence variation in multiple wild and cultivated potatoes determined that CWC15 is well conserved (Supplemental Fig. S7C). We performed co-immunoprecipitation in planta to explore the interaction between IPI-O1 and mutant IPI-O1[nls] with CWC15a/b in potato. The results showed that both StCWC15a and StCWC15b interacted with IPI-O1,

but the nuclear localization mutant IPI-O1[nls] did not interact with either protein (Fig. 5A). No interaction was observed between StCWC15 proteins and *P. infestans* effector Pi04314, another RxLR effector that interacts with three host protein phosphatase 1 catalytic isoforms, causing their re-localization from the nucleolus to the nucleoplasm (Boevink et al., 2016). The interactions were verified *in vitro* by GST-pull down and Y2H, aligning with the co-IP results (Fig. 5B and Supplemental Fig. S8A). Interaction tests between NES-IPI-O1[nls] and IPI-O1[nls]-NLS with StCWC15 indicated that neither mutant interacted with StCWC15 in vivo or in vitro (Supplemental Fig. S8, B and C). These data demonstrated that interaction between IPI-O1 and StCWC15 proteins is abolished when the 7 amino acids within the nuclear localization motif of IPI-O1 are mutated, regardless of its subcellular localization to the nucleus or cytoplasm. This underscores the essential role of these 7 amino acids in facilitating the interaction between IPI-O1 and CWC15.

We performed bimolecular fluorescence complementation (BiFC) to investigate the subcellular localization of the interaction between IPI-O1 and StCWC15. The results demonstrated that IPI-O1 interacted with StCWC15 proteins in nuclear speckles and the nucleolus (Fig. 5C). We did not observe fluorescence when testing the interaction between StCWC15 proteins and IPI-O1[nls] or Pi04314 (Fig. 5C). It is noteworthy that both the nucleolus and nuclear speckles are cellular structures highly correlated with RNA splicing (Liao and Regev, 2021). While expression of fusion proteins of StCWC15a and StCWC15b with the red fluorescence protein (RFP) demonstrated that StCWC15a primarily localized to the nucleoplasm (91%), the protein was occasionally found in the nucleolus (5%) and nuclear speckles (4%). StCWC15b was distributed similarly and was localized in the nucleoplasm (90%), nucleolus (5%), and nuclear speckles (5%) (Supplemental Fig. S9A). To investigate potential changes in StCWC15 subcellular localization influenced by the IPI-O1 effector, we co-expressed StCWC15a/b-RFP fusion proteins with GFP-IPI-O1 in *N. benthamiana*, with GFP-IPI-O1[nls] and GFP protein alone as controls. The results showed that StCWC15a protein was re-localized into speckles (63%) or the nucleolus (26%), and that StCWC15b protein was re-localized into speckles (60%) or the nucleolus (25%) when co-expressed with IPI-O1, but not with IPI-O1[nls] or GFP (Fig. 5D and Supplemental Fig. S9B). To further investigate the re-localization of StCWC15 during *P. infestans* infection, *N. benthamiana* expressing GFP–CWC15a/b was inoculated with zoospores of transgenic *P. infestans* isolate 88069 expressing the fluorescent protein RFP. The number of cells with speckle-localized CWC15 near the infected hyphae increased more than in cells distant from the infected hyphae (Supplemental Fig. S10). This effect was not observed in cells

330 expressing the GFP control during *P. infestans* infection (Supplemental Fig. S10). These findings suggest

331 that IPI-O1 interacts with StCWC15 proteins in the nucleus and induces their re-localization from the

332 nucleoplasm into speckles and the nucleolus.

CWC15 positively regulates *RB*-mediated HR and plant resistance against *P. infestans* infection

To determine the effect of CWC15 on *RB*-mediated immunity, we transiently silenced *CWC15*,

transcription-related factor transcription initiation factor TFIID subunit 12b-like (*PPS7*) (Katou et al.,

2005), splicing-related genes serine/Arginine-rich protein 30 (*SR30*), and arginine/serine-rich zinc

knuckle-containing protein 32 (*RS2Z32*) (Reddy, 2004) in *RB* transgenic *N. benthamiana* followed by

transient expression of IPI-O1. Infiltrated areas were assayed for HR elicitation and we found that

silencing of *CWC15* resulted in a reduced incidence of HR compared to the EV control, while *PPS7*,

SR30, or *RS2Z32* silencing had no effect on *RB* recognition of IPI-O1 (Supplemental Fig. S11A). *SGT1*

silencing, known to be required for *RB* function (Bhaskar et al., 2008), also reduced HR incidence

compared to the EV control (Supplemental Fig. S11A). The effect of CWC15 on *R* gene function

appeared to be specific for *RB*, as silencing of *CWC15* had no effect on the elicitation of HR following

co-infiltration of *R1*, *Rpiblb2*, or *R3a* with their corresponding effectors, or following infiltration of

elicitor *INF* (Supplemental Fig. S11B).

Based on the transient screening results, we further verified the function of *CWC15* by virus-induced

gene silencing (VIGS). The TRV-*NbCWC15* silenced plant exhibited small and short morphology

compared to wildtype (Supplemental Fig. S12, A and B). Similar to the transiently silenced results, TRV-

NbCWC15 silenced plants disturbed the recognition of IPI-O1 by *RB* (Supplemental Fig. S12, C-E).

Together, these indicated CWC15 as an interactor of IPI-O1 involved in *RB* activation. To verify the

function of potato *CWC15* in *RB*-mediated resistance, we stably silenced *StCWC15* by introducing a

hairpin-silencing construct into *RB*-transgenic potato by transformation. We obtained three independent

transgenic lines in which *StCWC15* transcription was reduced and a susceptible phenotype was exhibited

following inoculation with *P. infestans* when compared to the *RB* transgenic parent potato (Fig. 6, A to D).

CWC15 is able to regulate mRNA alternative splicing in *Arabidopsis* (Slane et al., 2020). To test whether

StCWC15 could play a role in *P. infestans*-induced AS of *RB*, we detected *RB* transcript levels in the

highest silencing efficiency line *RB-StCWC15* RNAi#2 following inoculation with *P. infestans* isolate

JH19. We found that the *RB_CDS/RB_IR* splicing ratio remained constant in the *StCWC15* knockdown

plant, while the ratio increased in the *RB* transgenic control plant (Fig. 6E). These data suggest that

CWC15 can modulate *RB* splicing, thereby influencing *RB*-mediated immunity.

To detect the possibility of CWC15 involvement in the recognition of IPI-O1 by RB_CDS as a

tradition guardee protein. We assayed for *P. infestans* resistance and HR induction following silencing

CWC15 in *RB*, *RB_CDS* and *Rpi-blb2* transgenic *N. benthamiana*. Silencing of *CWC15* significantly

impaired IPI-O1 induced HR and resistance to *P. infestans* in the *RB* line, but had no effect on the

resistance in *RB_CDS* and *Rpi-blb2* transgenic lines (Fig. 6, F to H). Additionally, *CWC15* silencing had

no effect on the HR induced by RB_CDS/IPI-O1, Avr3a/R3a or INF1 (Supplemental Fig. S13, A and B).

This effect does not appear to be due to interaction between CWC15 and RB, as no interaction was found

with RB_CDS or RB_IR proteins (Supplemental Fig. S13C). These data highlight that splicing factor

CWC15 is specifically involved in regulating *RB* splicing changes during *P. infestans* infection, effecting

RB-mediated immunity in potatoes, but does not affect *RB* function once the intron is removed. In further

experiments, we complemented *StCWC15* in the TRV-*NbCWC15* silenced *RB* transgenic *N. benthamiana*

using silencing-resistant *CWC15* synthetic constructs to test CWC15 function. These synthetic genes were

constructed with shuffled synonymous codons and encode full-length StCWC15a/b (Supplemental Fig.

S14A). Expression of StCWC15a^{syn} and StCWC15b^{syn} rescued the impaired HR and *RB* resistance in

TRV silenced-*CWC15* transgenic plants (Fig. 7, I to K; Supplemental Fig. S14B). Immunoblot analysis

revealed that expression of StCWC15a^{syn} and StCWC15b^{syn} recovered the expression of CWC15 in

silenced *N. benthamiana* (Supplemental Fig. S14, C and D). Collectively, our results indicate that IPI-O1

interacts with the splicing factor CWC15 to mediate the AS of *RB*, leading to increased expression of

resistant transcript *RB_CDS* and activation of *RB*-mediated immunity.

Regulation of *RB* intron splicing is critical for balancing plant growth and immunity

Our prior findings indicated that the *RB* intron is primarily unspliced in the absence of *P. infestans*,

leading to the production of mainly truncated protein. The presence of *P. infestans* effector IPI-O1 in the

nucleus leads to increased intron splicing and increased translation of RB_CDS protein. This discovery

suggests that the plant maintains resistance gene *RB* in a non-spliced state during normal growth and may

avoid inappropriate activation of immunity responses that could impact growth and development. To test

this, we assayed *RB*, *RB_CDS*, and *RB_IR* transgenic lines for their growth characteristics. Positive

transgenic lines were screened from multiple differentiated clustered buds using PCR verification. To

ensure consistency, three lines were selected from 67 *RB*-transformed seedlings, 34 RB_CDS transformed seedlings, and 29 RB_IR transformed seedlings, all displaying similar *RB* transcription levels and an estimated single copy of the transgene. (Supplemental Fig. S15A and Table S1). *RB_CDS* transgenic lines exhibited shorter stem height and root length than non-transformed Favorita in the seedling stage, while *RB* and *RB_IR* transgenic lines were not significantly different (Fig. 7, A-C). Furthermore, *RB_CDS* transgenic potato also displayed slower growth and dwarf phenotypes compared to the wild-type *RB* transgenic potatoes when grown in pots (Supplemental Fig. S15, B and C). The total tuber quality per plant and the individual tuber quality were significantly reduced in plants expressing *RB_CDS* compared with Favorita, while total tuber quality and individual tuber quality in *RB* and *RB_IR* transgenic potatoes were not significantly altered (Fig. 7, D and E, Supplemental Fig. S 15D). Studies have shown that accumulation of salicylic acid (SA) in response to biotic stress affects the growth of plants (Ning et al., 2017; Gao et al., 2021a; Gao et al., 2021b). We compared SA concentrations in *RB* and *RB_CDS* transgenic potatoes with non-transformed wild-type Favorita potatoes and found that SA accumulation in the *RB_CDS* transgenic line was higher than in wild-type and *RB* transgenic potatoes (Fig. 7F). Additionally, the transcription of *SAR Deficient 1* (*SARD1*), a key regulator of SA synthesis and plant immunity (Zhang et al., 2010), and the SA-regulated gene *pathogenesis-related gene 1* (*PR1*) were up-regulated in *RB_CDS* transgenic potatoes compared to wild-type Favorita and *RB* transgenic potatoes (Fig. 7G and H). These results indicate that the sustained expression of *RB_CDS* leads to continuous activation of SA-mediated defense responses, which likely affects plant growth and tuber production.

Previous reports have shown that truncated NLR proteins are expressed and have a potential role in plant immunity (Nishimura et al., 2017a; Liang et al., 2019). Especially the C-terminally truncated protein, which are encoded CC or TIR of NLR, their self-association is critical for their activation of downstream immunity (Rairdan et al., 2008; Gutierrez et al., 2010; Maekawa et al., 2011). N terminal CC domain of RB also displayed self-interaction in vivo and in vitro, and is vital for the immunity activation (Chen et al., 2012; Zhao and Song, 2021). In this work, we identified RB encoded a truncated NLR protein RB_IR during normal growth stage, which encompassed the N-terminal of RB but cannot activate immunity. To deeply explore the interplay of truncated and full-length RB proteins, we detected their interaction by co-IP. The result showed that RB_IR can interact with RB_CDS, and especially existed interaction with N terminal CC domain of RB_CDS (Fig. 8A). This led us to hypothesize that overexpression RB_IR could affect the RB CC motif self-association and thereby impact immunity activation. To test this, we

overexpressed RB_IR in *RB_CDS* transgenic *N. benthamiana*. The results show that overexpression

RB_IR compromised RB_CDS resistance compared with GFP control, whereas overexpression of the

protein R2-CC, the N-terminus of R2, had no effect on RB_CDS resistance (Fig.8, B to D).

Overexpression of RB_IR also disrupted the HR induced by IPI-O1 in *RB_CDS* transgenic *N.*

benthamiana (Supplemental Fig. S16). Together, these results suggest that truncated protein RB_IR can

interact with CC domain of RB_CDS, and it overexpression affects RB_CDS mediated resistance during

infection. This is likely due to the disruption of RB-CC oligomerization by RB_IR, as we observed that

overexpression of RB_IR could disrupt the self-association of RB-CC through RB_IR heteromeric

binding (Figure. 8E). Whereas another N-terminal protein, R2-CC, had no effect on either RB_CDS

immunity activation or RB-CC self-association (Figure. 8E). These findings suggest that the

overexpression of RB_IR could potentially compete with N-terminal self-association of full-length RB

protein, consequently disrupting immunity activation.

Discussion

Earlier studies indicate interaction between IPI-O1 and the RB CC domain in vivo (Chen et al.,

2012). Further evidence indicates that IPI-O1 accumulates in the nucleus and it induces the HR with RB

protein in the cytoplasm (Wang et al., 2018; Zhao and Song, 2021). Our current work builds on previous

data mainly uncovered a new model of NLR RB mediated plant immunity responses: a pathogen effector

(IPI-O1) targets a component of the spliceosome (CWC15), modulating the intron splicing of a plant

NLR gene (*RB*). This results in increased expression of the functional transcript RB_CDS that can

provide resistance to pathogen. This study implicates the spliceosome as a sensor to surveil pathogen

effector activity, and activate plant immunity through proper splicing of an NLR. In this model, the plant

utilizes alternative intron splicing to regulate expression of the NLR to prevent inappropriate stress-

induced responses during normal growth and timely activate plant immunity response upon the presence

of pathogen (Fig. 8F). Here, we highlighted the presence of IPI-O1 in the nucleus interacted with splicing

factor StCWC15 forming a regulatory hub, the intron of RB participates in the regulatory model to form

resistance under biotic stress.

Pathogens and plants are eternally entrenched in molecular warfare, utilizing an arsenal of microbial

effector proteins that manipulate the host to cause disease and an equally diverse set of host protein

receptors that recognize the presence of pathogen effectors to activate immune responses. Given the

complexity of possible interactions between host and pathogen molecules, it is no surprise that multiple mechanisms exist to mediate effector recognition and activate resistance. Classical activation models of NLR proteins and effectors include direct recognition, indirect recognition, integrated decoy, and activation traps (Duxbury et al., 2021). The simplest mechanism is direct interaction between NLR proteins and their corresponding effector, leading to changes in protein structure or location that results in resistance activation (Dodds et al., 2006). The indirect recognition mechanisms are divided into the guardee and the decoy models. In the guardee model, the effectors target a defense-related protein, either through direct binding or protein modification, that is monitored by an NLR protein. When changes in the guarded protein are recognized, this leads to activation of the NLR protein and elicitation of immune responses (Dangl and Jones, 2001; Axtell and Staskawicz, 2003). The decoy model is derived from the guardee model, but decoy proteins have no function in host defense or susceptibility in the absence of the NLR protein (van der Hoorn and Kamoun, 2008; Zhou and Chai, 2008). Our results demonstrate an additional mechanism of effector modulation of *NLR* expression in plants. IPI-O1 is one of multiple *P. infestans* effectors capable of reprogramming genome-wide alternative intron splicing of the host (Huang et al., 2020b). The proposed model involves the spliceosome component CWC15 protein, which is translocated to the nucleolus in the presence of IPI-O1, leading to splicing of the intron within the *RB* gene. *RB* intron splicing is necessary for the expression of a full-length protein that can activate immunity. The outcome of this model is similar to TAL effector activation traps that result in increased transcription of multiple executor *R* genes (Bogdanove et al., 2010). However, in the case of IPI-O1, we found that total *RB* transcription is not significantly induced during infection. This is inconsistent with the results of previous studies that demonstrated an increase in *RB* transcription during *P. infestans* infection in potato (Lara C. Kramer, 2009), but is consistent with reports that *RB* is transcribed at a constitutive level (James M. Bradeen, 2009). These contrasting observations are most likely due to differences in host and pathogen genotypes used in our experiments, which could affect IPI-O expression and *RB* transcript accumulation. The primers and methods used for detection of RB transcripts was also different between studies. Regardless, our results indicate that *RB* gene expression is likely regulated at the co-transcriptional or post-transcriptional level, and does not follow a transcription activation model through promoter binding as it is reported in other pathosystems such as with TAL effectors.

The prevalence of alternative splicing (AS) in plants contributes significantly to transcriptome and proteome diversity. It is estimated that between 42% and 61% of all gene transcripts are alternatively

spliced (Filichkin et al., 2010; Lu et al., 2010b; Marquez et al., 2012; Reddy et al., 2013b; Klepikova et al., 2016). AS serves as a key player in gene regulation at the RNA level and has been recognized for its role in regulating plant immunity (Reddy, 2007; Huang et al., 2017; Reddy et al., 2020). Several NLR genes undergo AS in the coding sequence or untranslated regions (UTRs; Yang et al., 2014; Lai and Eulgem, 2018). Our data indicate that the *RB* gene produces two different transcripts that encode either a full-length NLR protein that can confer resistance to *P. infestans*, or a truncated protein only containing the N-terminally domain of NLR. Premature termination codon (PTC)-containing transcripts that arise as a result of AS are often subjected to a nonsense-mediated mRNA decay (NMD) process that is intended to prevent their translation (Brogna and Wen, 2009). An analysis of transcripts produced during bacterial infection in Arabidopsis showed that the majority of CC (coiled coil)-NLR (65.1%) and TIR (Toll-like, Interleukin 1)-NLR (81.2%) transcripts exhibit evidence of NMD (Jung et al., 2020). However, transcripts containing PTCs can be used to express the truncated protein or have functional relevance (Liu et al., 2017; Nishimura et al., 2017b). Our results indicate that the *RB* gene transcripts that retain the intron are still detectable by RT-PCR and the truncated RB_IR protein is detectable in plants (Fig. 1C), suggesting that the short transcripts are not immediately degraded. We also demonstrated that this truncated protein could have an impact on immunity activation through the interaction with full-length RB protein. Previous evidence indicates that the CC domain of RB can self-oligomerize and this oligomerization is important in activation of immunity (Chen and Halterman, 2017; Zhao and Song, 2021). In this paper, we preliminarily demonstrated that overexpression of truncated RB_IR can disrupt the self-association of RB. The impact of this interaction on formation of a resistosome structure is therefore likely, but requires further confirmation. Such an investigation would help elucidate whether an RB_IR and RB_CDS hybrid structure is formed to prevent autoactivation due to RB_CDS self-oligomerization, thus avoiding inappropriate immune responses during regular plant growth.

In addition to directly affecting the expression of *NLR* genes, AS plays a more global role in regulating plant disease resistance responses. *Arabidopsis snc1* mutants constitutively activate defense responses and have increased resistance to pathogens (Zhang and Li, 2005). A screen for suppressors of *snc1* revealed two *MOS* (modifier of *snc1*) genes, *MOS4* and *MOS14*, that are involved in mRNA splicing (Palma et al., 2007; Xu et al., 2011). The MOS14 protein is similar to transportin-SR, which is necessary for nuclear import of serine-arginine (SR) rich proteins. The *mos14-1* in *Arabidopsis* affects splicing of *NLR* genes *SNC1* and *RPS4* and compromises RPS4-dependent immunity and basal resistance to

Pseudomonas syringae DC3000 (Xu et al., 2011). In *Arabidopsis*, *P. syringae* introduces small RNAs that silence *SWP73A*, a chromatin-remodeling gene that regulates expression of several *NLRs* through promoter binding and alternative splicing (Huang et al., 2021). *CDC5* has been reported to regulate alternative splicing of *RPS4*, which is important for RPS4 activity (Palma et al., 2007; Zhang and Gassmann, 2007). In this work, the IPI-O1 effector interacts with CWC15 to regulate splicing of the *RB* gene. Further work It seems unlikely that *RB* is only regulated by IPI-O1. More likely, the CWC15 and RB intron composed plant decoy can been regulated by more proteins, especially effectors. The impact of other effectors on regulation of *RB*-mediated plant defense responses will be a focus of future research.

For plants, homeostasis between growth and defense is vital for survival and reproduction in nature (He et al., 2022a, b). Research on the role of *NLR* expression in regulating growth versus defense trade-offs in plants has been reported previously (Li et al., 2012; Tsuchiya and Eulgem, 2013; Cui et al., 2020; Huang et al., 2021; He et al., 2022b). NLR proteins are the intracellular immune receptors and activators in plants. The transition between autoinhibited and activated states of NLRs is fine-tuned by intra- and inter-molecular interactions. Over-accumulation of R proteins could lead to autoimmunity (Dinh et al., 2022), whereas insufficient NLR expression could lead to pathogen susceptibility. We found that the intron within the coding sequence of *RB* is alternatively spliced and, under normal growth conditions, transcripts are maintained with the intron retained, leading to a truncated protein. Presumably, this is a strategy to reduce elicitation of immune responses in the absence of an invading pathogen by regulating expression of the full-length RB protein in the plant. This is supported by our results that overexpression of an *RB* gene with the intron removed leads to constitutive expression of defense-related genes, smaller plants, and lowered tuber yield. Following *P. infestans* inoculation to introduce the effector IPI-O1, *RB* produces more intron spliced *RB_CDS*, which encodes a full-length NLR protein to provide resistance. Our findings suggest a strategy to reduce elicitation of immune responses in the absence of an invading pathogen by regulating expression of the functional plant *R* gene.

Materials and methods

RT-qPCR analysis of *RB* splicing ratio and gene expression

Total RNA was extracted using E.Z.N.A.® Plant RNA Kit (R6827-01, OMEGA) following the manufacturer's instructions and treated with RNase-Free DNase (M6101, Promega) to remove genomic DNA contamination. Approximately 700 ng RNA was used for reverse transcription with HiScript III 1st

Strand cDNA Synthesis Kit (R312-01, Vazyme). RT-qPCR was performed with 2μL of 3-fold-diluted

cDNA in a total reaction volume of 20μL in technical duplicates using ChamQ Universal SYBR qPCR

Master Mix (Q711-02, Vazyme Biotech) with an ABI 7500 Fast Real-time PCR system according to the

manufacturer's instructions. PCR conditions were as follows: 95°C for the 30s, followed by 40 cycles of

95°C for 5 s, 60°C for 34 s, and then 95°C for 15 s, 60°C for 1 min, 95°C for 30 s, 60□ for 15s for the

melt curve. Relative transcript amount differences were calculated using the $2^{-\Delta\Delta CT}$ method in ABI 7500

System Sequence Detection Software. Target-specific amplification efficiencies are given in

Supplemental Data Set S4. Three biological replicates were performed and the results showed similar

trends.

Gene silencing and complementation in *N. benthamiana*

Virus-induced gene silencing (VIGS) was performed in *N. benthamiana* as previously described

(Stam et al., 1997). For VIGS target design, the specific DNA fragments with 300 base pairs (bp) from

GFP, *NbCWC15* (Niben101Scf04847g02012.1/Ni-ben101Scf00693g01009.1) predicted by VIGS tool

(https://vigs.solgenomics.net/) were cloned into Tobacco Rattle Virus (TRV) construct. The sequences of

the targets are listed in Supplemental Data Set S1. The constructs were introduced into *A. tumefaciens*

GV3101 by electroporation. Suspensions of *A. tumefaciens* strain GV3101 harboring TRV1 and TRV2

constructs were mixed in a 1:1 ratio in infiltration buffer (10 mM MES, 10 mM MgCl$_2$ pH 5.6, and 150

uM acetosyringone) to a final OD$_{600}$ of 0.4, and then left at room temperature for 2h. TRV2-EV was used

as a negative control. Two-week-old *N. benthamiana* plants were infiltrated with *A. tumefaciens* for VIGS

assays, and their upper leaves were used 2-3 weeks later. The silencing efficiency was confirmed by RT-

qPCR. For the complementation assays, the shuffled synonymous synthetic *StCWC15* sequences were

generated by Sangon Biotech and cloned into expression constructs pK7WGF2 with a C-terminal GFP tag.

Synthetic proteins were detected by western blot.

The 300-bp fragment of *NbPPS7a/b* (*Niben101Scf07798g01001.1/Niben101Scf01573g01003.1*),

NbSR30 (*Niben101Scf05298g04002.1*), *NbRS2Z32* (*Niben101Scf07227g00011.1*), *NbSGT1*

(*Niben101Scf03241g00015.1*), *NbU1-70ka/b* (*NbS00011208g0006.1/ NbS00015731g0004.1*),

NbSF3b1a/b (Niben101Scf06716g01016.1/ Niben101Scf06128g00004.1), *NbGRP7*

(Niben101Scf00215g03004.1) were cloned and ligated into pK7GWIWG2D(II) to generate a hairpin

silencing construct. The sequences of the targets are listed in Supplemental Data Set S1. The

pK7GWIWG2D(II) empty vector was used as a control. Six- to seven-week *N. benthamiana* leaves were

infiltrated with GV3101 *A. tumefaciens* strains containing either the silencing construct or empty vector

control diluted to an OD_{600}=0.5.

Growth of plant material and microbial strains

Solanum tuberosum cv. Favorita and transgenic plants were grown on Murashige and Skoog (MS)

medium in a growth chamber under 16 h light/8 h dark conditions at a constant temperature of 24 °C and

light intensity of 100 μM m^{-2} s^{-1}. Plants in soil were growth with 12 h light/12 h dark conditions at a

constant temperature of 22°C and light intensity of 153 μM m^{-2} s^{-1}.

N. benthamiana plants were grown and maintained in the greenhouse with 16 h light at 25°C and 8 h

dark at 20°C throughout the experiments. *P. infestans* strains JH19 (Howard et al., 2013), and 88069[td],

expressing the red fluorescent marker tandem dimer RFP (tdTomato) (Dagdas et al., 2016) were grown

on rye sucrose agar (RSA) at 18°C. *P. capsici* strain LT1534 was grown on V8 medium at 25□.

Escherichia coli strains DH5α, JM109, and BL21 and *A. tumefaciens* strain GV3101 were grown on

Luria-Bertani (LB) medium at 37 °C and 30 °C, respectively.

S. bulbocastanum accession PT29 (PI 243510) was originally obtained from the United States Potato

Genebank (NRSP-6) in Sturgeon Bay, WI. The accession sto389-4 containing *Rpi-sto1*, blb8005

containing *RB* were kindly provided by Prof. Jack H. Vossen. Binary plasmids pBINPLUS: *Rpi-sto1*,

pBINPLUS: *RB* were constructed in house and transformed into cv Desiree using Agrobacterium

mediated transformation (Zhu et al., 2014).

Plasmid construction

All primers and plasmids used for cloning and experiments are listed in Supplemental Data Set S2.

All the templates used for PCR amplification come from genomic DNA (gDNA) or complementary DNA

(cDNA) of *P. infestans* (strain T30-4), *N. benthamiana*, and *Solanum tuberosum* cv. Favorita using 2 ×

Phanta Max Master Mix (P515-01, Vazyme Biotech). FastPure Plasmid Mini Kit, FastPure Gel DNA

Extraction Mini Kit and ClonExpress Ultra One Step Cloning Kit (Cat No. DC201-01, DC301-01 and

C115-01) were used for nucleic acid purification, digestion, and ligation.

Co-immunoprecipitation assays and GST pull-down

The *IPI-O1* gene without the signal peptide was inserted into pK7WGF2 with N-terminal RFP tag

using ClonExpress Ultra One Step Cloning Kit (C115-01, Vazyme Biotech), and *StCWC15a/b* were

inserted into pK7WGF2 with C-terminal GFP for expression in *N. benthamiana.* Leaves of 6-week-old *N.*

benthamiana plants were co-infiltrated with RFP-tagged *IPI-O1*, GFP-tagged *StCWC15a/b*, and the P19

silencing suppressor. Two days after agro-infiltration, the leaves were frozen in liquid nitrogen and

ground to powder using a mortar and pestle. Proteins were extracted with extraction buffer (10% [vol/vol]

glycerol, 25mM Tris pH=7.5, 1mM EDTA, 150mM NaCl, 2% [wt/vol] PVPP, 10mM DTT, 1×Protease

Inhibitor Cocktail (Sigma-Aldrich), 0.1% Tween 20). The samples were centrifuged at 4°C for 20 min at

$12,000 \times g$, and the supernatant was transferred to a new tube. For GFP-IP, 1 mL of supernatant was

incubated at 4°C for 2 h with GFP-Trap_A beads (Chromotek, Planegg-Martinsried, Germany). The beads

were collected by centrifugation at $800 \times g$ and washed four times in 1 mL washing buffer (25 mM Tris

pH=7.5, 150 mM NaCl and 1mM EDTA, 10% [vol/vol] glycerol, 0.1% Tween 20). The beads were boiled

for 10 min at 95°C. SDS-PAGE was performed with the supernatant, and co-IP of StCWC15a/b with IPI-

O1 was detected using anti-GFP or anti-RFP (Chromotek, Planegg-Martinsried, Germany).

For *in vitro* GST pull-down assays, GST-StCWC15a/b and His-IPI-O1 were expressed in *E. coli* strain

BL21. Co-precipitation of StCWC15a/b with IPI-O1 was examined by protein blotting before (input) and

after affinity purification (pull-down) using glutathione agarose beads (GE Healthcare). Interactions were

detected using anti-GST and anti-His (Abmart).

***P. infestans* and *P. capsici* infection assays**

P. infestans strains were grown on RSA medium plate at 18°C for 14 d. The sporangia were scraped

into cold, distilled water and incubated at 4°C for 1-2 h until zoospore release. The zoospore suspension

was adjusted to 100 zoospores/μL before inoculation. Ten microliter drops of *P. infestans* zoospores were

placed on the abaxial side of detached leaves. The inoculated leaves were kept at 18°C for 4-7 days. The

whole potato infection assays used a zoospore suspension sprayed on the plant followed by incubation in

a controlled environment chamber at 20°C for 6 days. The infected *N. benthamiana* leaves were stained

with trypan blue. Leaves were soaked in trypan blue solution (10 mL 85% lactic acid aqueous, 10 mL

water-saturated phenol, 10 mL 98% glycerol, 10 mL distilled water, 15 mg trypan blue) overnight.

Stained leaves were then transferred into ethanol and incubated at 37°C for 24 h before observation. *P.*

capsici strains were grown on V8 medium plate at 25□ for 3 d. Ten pieces of mycelium with a size of

about 2 mm^2 were cultured in V8 liquid medium for 4 days. The culture medium was discarded, the

mycelium was washed with distilled water three times, and distilled water was added with just enough to

cover the mycelium. The culture was incubated at 25□ for 12 hours until sporangia were observed, then

was placed at 4°C for 20 minutes to stimulate zoospores release.

Agroinfiltrations

Construct plasmids were transformed into *A. tumefaciens* strain GV3101 and grown overnight in

Luria-Bertani (LB) culture medium with the antibiotic kanamycin and rifampicin at 30°C. Bacteria were

harvested by centrifugation at 5000g for 4 min and then resuspended and diluted in an agroinfiltration

buffer (10 mM $MgCl_2$, 150 mM acetosyringone, 10 mM MES) to an optical density of $OD_{600} = 0.01$-1.0

depending on the experiments. After 2-4 h of incubation at room temperature, they were infiltrated with a

needleless syringe into the abaxial side of leaves from 4–5-week-old *N. benthamiana* plants. For the co-

expression of multiple constructs, suspensions carrying each construct were thoroughly mixed before

infiltration.

Confocal microscopy and bimolecular fluorescence complementation (BiFC) assays

Confocal microscopy was performed on *N. benthamiana* leaf patches post infiltration by LSM 980

laser-scanning microscope with 20×/0.8 or 60×/0.95 objective lens (ZEISS LSM 980 with Airyscan2).

Parameters that are used for fluorescence detection are GFP/YFP, (excitation 488 nm, emission 505–540

nm), RFP (excitation 514 nm, emission 518–582 nm), DAPI (excitation 405 nm, emission 430–480 nm).

Leaves were stained with 2 µg/mL of 4', 6-diamidino-2-phenylindole (DAPI) for 5 min to indicate nuclei.

ZEISS ZEN Microscope Software was used to process the images.

pGTQL1221YC and pGTQL1211YN (nYFP/cYFP) vectors were used for bimolecular fluorescence

complementation (BiFC) assays (Lu et al., 2010a). Coding regions of *IPI-O1* and *IPI-O1^{nls}* without a

signal peptide and *StCWC15a/b* were amplified with primers containing attB tails and BP-cloned into

pDONR207 (Invitrogen). To generate BiFC constructs, constructs were integrated into destination vectors

pGTQL1221YC or pGTQL1211YN by LR recombination using Gateway LR clonase II (Invitrogen). The

constructs were transformed into *A. tumefaciens* strain GV3101. Two days after agroinfiltration, the Zeiss

980 laser scanning confocal microscope was used for YFP fluorescence detection.

Transformation of potato and RNAi in potato

For generating transgenic potato, the entire *RB* gene sequence (3592 bp), *RB_CDS* transcript (2913

bp), *RB_IR* transcript (507 bp) were cloned into pBinGFP2 binary vector containing the *CaMV35S*

promoter. The constructs were transformed into the *A. tumefaciens* strain GV3101. To generate stable

transgenic potato plants, stems were used for transformation. Stem segments of *Solanum tuberosum* cv.

Favorita were incubated in MS-medium (Murashige and Skoog, 1962). Agrobacterium containing each

plasmid was grown and diluted to a cell concentration of OD_{600}=0.5. Stem segments were added and

incubated in the dark for 45 mins. After co-incubation, stem segments were transferred to MS solid

medium supplemented with 2.5 mg/L zeantinribose, 0.2 mg/L NAA, 0.02 mg/L GA_3 for 48 hours. After

that, stems were washed with 300mg/L Timentin and transferred to MS solid medium supplemented with

2.5 mg/L zeantinribose, 0.2 mg/L NAA, 0.02 mg/L GA_3 and 200 mg/L Timentin for 2 weeks. For callus

induction, stems segments were kept on MS solid medium with 2 mg/L zeantinribose, 0.02 mg/L NAA,

0.02 mg/L GA_3, 300 mg/L Timentin and 50 mg/L kanamycin. Every 14 days, stem segments or

developing calli were placed onto new medium. After bud initiation formation (following 4-5 transfers),

shoots were cut off from the callus and transferred to MS-medium containing 300 mg/L Timentin and 50

mg/L kanamycin.

For hairpin RNA interference (RNAi), the chalcone synthase (CHSA) intron was amplified from

plasmid pFGC5941. The PCR-amplified 3'StCWC15 cDNA fragments overlapped the 5' and 3' end of the

CHSA intron from two directions. The overlapped fragment was inserted in the p1300 vector and

transformed into *A. tumefaciens* strain GV3101 before potato transformation. Transformation is the same

as potato stable overexpression transformation. *StCWC15s* silencing transgenic lines were confirmed by

RT-qPCR in buds using the primers indicated in Supplemental Data Set S2.

5' RACE

Rapid amplification of cDNA ends (5' RACE) was used to determine the transcripts of the *RB* gene

using the SMARTer RACE cDNA Amplification Kit (Clontech) according to the protocol. For synthesis

of first-strand cDNA, we designed the *RB* specific primer (Supplemental Data Set S2). Next, we used 10

μM *RB* specific primer and the provided UPM primer in the kit for RACE PCR. Amplification was

performed using a touchdown PCR protocol with five cycles of 94°C for 30 s, 72°C for 4 min, then five

cycles of 94°C for 30 s, 70°C for 30s, 72°C for 4 min, final 25 cycles of 94°C for 30 s, 68°C for 30 s,

72°C for 4 min. The obtained 5'RACE products were gel excised, introduced into the pUC19 vector using

in-fusion cloning, and sequenced by Sanger sequencing to detect the transcripts.

LUC reporter assays

For LUC reporter system assays in *N. benthamiana* leaves, a 1986-bp *RB* fragment including the

intron structure was amplified from *RB* transgenic potato (SP951) genomic DNA and fused with *LUC*

gene ligated into pCAMBIA1300. Four- or five-week-old *N. benthamiana* leaves were infiltrated with

Agrobacterium strain GV3101 harboring the indicated constructs resuspended in infiltration buffer for 48

h, after which the leaves were sprayed with 1 mM D-luciferin substrate and incubated for 5 min in the

dark. The sprayed leaves were photographed using a Tanon-5200 Multi Chemiluminescent Imaging

System (Tanon, China). For quantification of LUC activity, *N. benthamiana* leaf discs were collected at

48 h after infiltration (hpi) and incubated with 1 mM D-luciferin in a 96-well plate, and LUC activity was

quantified with the Promega GloMax Navigator microplate luminometer (Promega, USA).

Yeast two-hybrid screening

To identify interacting partners with the *P. infestans* effector IPI-O1, a yeast two-hybrid interaction

screen was performed by Hybrigenics Services (Évry-Courouronnes, France) using IPI-O1 as a bait in an

ULTImate Y2H library constructed from *N. benthamiana*. IPI-O1, encoding amino acids 2-132, was

cloned into the inducible pB35 vector as a fusion with GAL4 at the amino terminus. The prey library was

prepared in the proprietary pP6 Hybrigenics vector. A total of 81.6 million interactions were analyzed and

254 clones were processed by sequencing of the prey inserts. These prey fragments were fused out-of-

frame with the Gal4 activation domain (AD) in pP6. However, *Saccharaomyces cerevisiae* are able to

translate a percentage of the correct fusion proteins using ribosomal frameshift (Fromont-Racine et al.,

1997; Vidal and Legrain, 1999). For this reason, it is possible to identify true interacting proteins from

out-of-frame clones.

Hypersensitive response phenotype and ion leakage

717   Hypersensitive cell death during the ETI process was scored at 4 dpi and was quantified using an ion

718   leakage assay (Mittler et al., 1999). Membrane ion leakage was determined by measuring electrolytes

719   leaked from leaf discs. For each measurement, five 9 mm leaf discs were immersed in 5 mL distilled

720   water at room temperature (22-25°C) for 3h. After which, the initial conductivity was measured with

Multi-parameter tester (Mettler Toledo S400-K), referred to as value A. Total conductivity was determined after boiling at 95°C for 25 min and cooling to room temperature; this is referred to as value B. The ion leakage was expressed as the percentage of the initial conductivity versus the total conductivity (value A/value B) ×100%.

Salicylic Acid Quantification

For SA quantification, three replicates of each genotype transgenic potato were collected. Approximately 100 mg (fresh weight) of leaf tissues were frozen in liquid nitrogen, ground and extracted with 1 mL 80% methanol (Anaqua™ Chemicals Supply, CAS: 67-56-1) containing 1% formic acid (Anaqua™ Chemicals Supply, CAS: 64-18-6) mixture. The resulting solution was then vortexed and placed in a shaker at 4□ for 24 h. Samples were centrifuged 10 min at 15,000 rpm, and supernatant was transferred to new tubes. Add 1 mL mixture and 0.3 g PVPP in the precipitate to shaker at 4□ for 24 h again. Mix two times supernatant and drying most of the liquid with a vacuum pump. Using 200 μL of 80% methanol for redissolve. Supernatants were filtered through 0.2 mm PTFE membrane (Millipore, Bedford, MA) and transferred to auto sampler vials. Quantification was performed by HPLC with the ACQUITY UPLC HSS C18 column.

Inhibitor assay

Pladienolide B (CAS:445493-23-2) was purchased from Adooq. PB, dissolved in DMSO, was infiltrated into *RB* transgenic *N. benthamiana*. The exact concentration of DMSO was used as control infiltrated in *N. benthamiana*.

Lanthanum (III) chloride (LaCl$_3$) (Sigma, CAS: 10099-58-8) was dissolved in water and infiltrated at a concentration of 2 mM.

ACKNOWLEDGMENTS.

We thank Sebastian Schornack (University of Cambridge), Jiamu Du, Zhe Wu (Southern University of Science and Technology), Jiming Jiang (Michigan State University), Yan Wang (Nanjing Agriculture University) for helpful discussions.

This work was supported by the Chinese National Science Funds Grants (32130088, 32102234) and the China Agriculture Research System-potato (CARS-potato, CARS-09-P20). This research was supported

751   in part by the United States Department of Agriculture (USDA) Agricultural Research Service (ARS).

752   Funding was provided through USDA National Institute of Food and Agriculture (NIFA) Plant-

753   Associated Microbes and Plant-Microbe Interactions award number 2014-67013-21593. The findings and

754   conclusions in this publication are those of the author(s) and should not be construed to represent any

755   official USDA or United States Government determination or policy.

References

Axtell, M.J., and Staskawicz, B.J. (2003). Initiation of RPS2-Specified Disease Resistance in Arabidopsis Is Coupled to the AvrRpt2-Directed Elimination of RIN4. CELL **112,** 369–377.

Bazin, J., Mariappan, K., Jiang, Y., Blein, T., Voelz, R., Crespi, M., and Hirt, H. (2020). Role of MPK4 in pathogen-associated molecular pattern-triggered alternative splicing in Arabidopsis. PLoS Pathog **16,** e1008401.

Bhaskar, P.B., Raasch, J.A., Kramer, L.C., Neumann, P., Wielgus, S.M., Austin-Phillips, S., and Jiang, J. (2008). Sgt1, but not Rar1, is essential for the RB-mediated broad-spectrum resistance to potato late blight. BMC Plant Biol **8,** 8.

Boevink, P.C., Wang, X., McLellan, H., He, Q., Naqvi, S., Armstrong, M.R., Zhang, W., Hein, I., Gilroy, E.M., Tian, Z., and Birch, P.R.J. (2016). A Phytophthora infestans RXLR effector targets plant PP1c isoforms that promote late blight disease. Nat Commun **7,** 10311.

Bogdanove, A.J., Schornack, S., and Lahaye, T. (2010). TAL effectors: finding plant genes for disease and defense. Current Opinion in Plant Biology **13,** 394-401.

Brogna, S., and Wen, J. (2009). Nonsense-mediated mRNA decay (NMD) mechanisms. Nature Structural & Molecular Biology **16,** 107-113.

Champouret, N., Bouwmeester, K., Rietman, H., Lee, T.v.d., Maliepaard, C., Heupink, A., Vondervoort, P.J.I.v.d., Jacobsen, E., Visser, R.G.F., Vossen, E.A.G.v.d., Govers, F., and Vleeshouwers, V.G.A.A. (2009). *Phytophthora infestans* Isolates Lacking Class I *ipiO* Variants Are Virulent on *Rpi-blb1* Potato. MPMI **22,** 1535-1545.

Chen, M., and Manley, J.L. (2009). Mechanisms of alternative splicing regulation: insights from molecular and genomics approaches. Nat Rev Mol Cell Biol **10,** 741-754.

Chen, Y., and Halterman, D.A. (2017). Phytophthora infestans Effectors IPI-O1 and IPI-O4 Each Contribute to Pathogen Virulence. Phytopathology **107,** 600-606.

Chen, Y., Liu, Z., and Halterman, D.A. (2012). Molecular determinants of resistance activation and suppression by Phytophthora infestans effector IPI-O. PLoS Pathogen **8,** e1002595.

Cheng, Y.T., Li, Y., Huang, S., Huang, Y., Dong, X., Zhang, Y., and Li, X. (2011). Stability of plant immune-receptor resistance proteins is controlled by SKP1-CULLIN1-F-box (SCF)-mediated protein degradation. Proc Natl Acad Sci U S A **108,** 14694-14699.

Cho, S., Hoang, A., Sinha, R., Zhong, X.-Y., Fu, X.-D., Krainer, A.R., and Ghosh, G. (2011). Interaction between the RNA binding domains of Ser-Arg splicing factor 1 and U1-70K snRNP protein determines early spliceosome assembly. Proceedings of the National Academy of Sciences **108,** 8233-8238.

Couto, D., and Zipfel, C. (2016). Regulation of pattern recognition receptor signalling in plants. Nat Rev Immunol **16,** 537-552.

Cretu, C., Agrawal, A.A., Cook, A., Will, C.L., Fekkes, P., Smith, P.G., Luhrmann, R., Larsen, N., Buonamici, S., and Pena, V. (2018). Structural Basis of Splicing Modulation by Antitumor Macrolide Compounds. Mol Cell **70,** 265-273 e268.

Cui, C., Wang, J.-J., Zhao, J.-H., Fang, Y.-Y., He, X.-F., Guo, H.-S., and Duan, C.-G. (2020). A Brassica miRNA regulates plant growth and immunity through distinct modes of action. Molecular plant **13,** 231-245.

Cui, H., Tsuda, K., and Parker, J.E. (2015). Effector-Triggered Immunity: From Pathogen Perception to

Robust Defense. Annual Review of Plant Biology **66,** 487-511.

Dagdas, Y.F., Belhaj, K., Maqbool, A., Chaparro-Garcia, A., Pandey, P., Petre, B., Tabassum, N., Cruz-Mireles, N., Hughes, R.K., Sklenar, J., Win, J., Menke, F., Findlay, K., Banfield, M.J., Kamoun, S., and Bozkurt, T.O. (2016). An effector of the Irish potato famine pathogen antagonizes a host autophagy cargo receptor. Elife **5**.

Dangl, J.L., and Jones, J.D.G. (2001). Plant pathogens and integrated defence responses to infection. Nature **411,** 826-833.

Deng, Y., Zhai, K., Xie, Z., Yang, D., Zhu, X., Liu, J., Wang, X., Qin, P., Yang, Y., Zhang, G., Li, Q., Zhang, J., Wu, S., Milazzo, J., Mao, B., Wang, E., Xie, H., Tharreau, D., and He, Z. (2019). Epigenetic regulation of antagonistic receptors confers rice blast resistance with yield balance. Science **355,** 962-965.

Dinesh-Kumar, S.P., and Baker, B.J. (2000). Alternatively spliced N resistance gene transcripts: Their possible role in tobacco mosaic virus resistance. Proc. Natl Acad. Sci. USA **97,** 1908–1913.

Dinh, H.X., Singh, D., Gomez de la Cruz, D., Hensel, G., Kumlehn, J., Mascher, M., Stein, N., Perovic, D., Ayliffe, M., Moscou, M.J., Park, R.F., and Pourkheirandish, M. (2022). The barley leaf rust resistance gene *Rph3* encodes a predicted membrane protein and is induced upon infection by avirulent pathotypes of *Puccinia hordei*. Nature Communications **13,** 2386.

Dodds, P.N., and Rathjen, J.P. (2010). Plant immunity: towards an integrated view of plant-pathogen interactions. Nature reviews. Genetics **11,** 539-548.

Dodds, P.N., Lawrence, G.J., Catanzariti, A.-M., Teh, T., Wang, C.-I.A., Ayliffe, M.A., Kobe, B., and Ellis, J.G. (2006). Direct protein interaction underlies gene-for-gene specificity and coevolution of the flax resistance genes and flax rust avirulence genes. Proceedings of the National Academy of Sciences **103,** 8888-8893.

Duxbury, Z., Wu, C.H., and Ding, P. (2021). A Comparative Overview of the Intracellular Guardians of Plants and Animals: NLRs in Innate Immunity and Beyond. Annu Rev Plant Biol **72,** 155-184.

Filichkin, S.A., Priest, H.D., Givan, S.A., Shen, R., Bryant, D.W., Fox, S.E., Wong, W.-K., and Mockler, T.C. (2010). Genome-wide mapping of alternative splicing in Arabidopsis thaliana. Genome Research **20,** 45-58.

Fromont-Racine, M., Rain, J.-C., and Legrain, P. (1997). Toward a functional analysis of the yeast genome through exhaustive two-hybrid screens. Nature Genetics **16,** 277-282.

Gao, M., He, Y., Yin, X., Zhong, X., Yan, B., Wu, Y., Chen, J., Li, X., Zhai, K., Huang, Y., Gong, X., Chang, H., Xie, S., Liu, J., Yue, J., Xu, J., Zhang, G., Deng, Y., Wang, E., Tharreau, D., Wang, G.L., Yang, W., and He, Z. (2021a). Ca(2+) sensor-mediated ROS scavenging suppresses rice immunity and is exploited by a fungal effector. Cell **184,** 5391-5404 e5317.

Gao, W., Liu, Y., Huang, J., Chen, Y., Chen, C., Lu, L., Zhao, H., Men, S., and Zhang, X. (2021b). MES7 Modulates Seed Germination via Regulating Salicylic Acid Content in Arabidopsis. Plants (Basel) **10**.

Gutierrez, J.R., Balmuth, A.L., Ntoukakis, V., Mucyn, T.S., Gimenez-Ibanez, S., Jones, A.M.E., and Rathjen, J.P. (2010). Prf immune complexes of tomato are oligomeric and contain multiple Pto-like kinases that diversify effector recognition. The Plant Journal **61,** 507-518.

Halterman, D., Wei, F., and Wise, R. (2003). Powdery mildew induced *Mla* mRNAs are alternatively spliced and contain multiple upstream open reading frames. Plant Physiol **131,** 558-567.

Halterman, D.A., and Wise, R.P. (2006). Upstream open reading frames of the barley Mla13 powdery mildew resistance gene function co-operatively to down-regulate translation. Molecular Plant

Pathology **7,** 167-176.

Halterman, D.A., Chen, Y., Sopee, J., Berduo-Sandoval, J., and Sanchez-Perez, A. (2010).
Competition between Phytophthora infestans effectors leads to increased aggressiveness on plants
containing broad-spectrum late blight resistance. PLoS One **5,** e10536.

Haverkort, A.J., Struik, P.C., Visser, R.G.F., and Jacobsen, E. (2009). Applied Biotechnology to
Combat Late Blight in Potato Caused by Phytophthora Infestans. Potato Research **52,** 249-264.

He, Z., Webster, S., and He, S.Y. (2022a). Growth-defense trade-offs in plants. Current Biology **32,**
R634-R639.

He, Z., Webster, S., and He, S.Y. (2022b). Growth-defense trade-offs in plants. Curr Biol **32,** R634-
R639.

Howard, B.E., Hu, Q., Babaoglu, A.C., Chandra, M., Borghi, M., Tan, X., He, L., Winter-Sederoff,
H., Gassmann, W., Veronese, P., and Heber, S. (2013). High-Throughput RNA Sequencing of
Pseudomonas-Infected Arabidopsis Reveals Hidden Transcriptome Complexity and Novel Splice
Variants. Plos One **8**.

Huang, C.Y., Rangel, D.S., Qin, X., Bui, C., Li, R., Jia, Z., Cui, X., and Jin, H. (2021). The
chromatin-remodeling protein BAF60/SWP73A regulates the plant immune receptor NLRs. Cell
Host Microbe **29,** 425-434 e424.

Huang, J., Lu, X., Wu, H., Xie, Y., Peng, Q., Gu, L., Wu, J., Wang, Y., Reddy, A.S.N., and Dong, S.
(2020a). *Phytophthora* effectors modulate genome-wide alternative splicing of host mRNAs to
reprogram plant immunity. Molecular plant **13,** 1470-1484.

Huang, J., Lu, X., Wu, H., Xie, Y., Peng, Q., Gu, L., Wu, J., Wang, Y., Reddy, A.S.N., and Dong, S.
(2020b). Phytophthora Effectors Modulate Genome-wide Alternative Splicing of Host mRNAs to
Reprogram Plant Immunity. Mol Plant.

Huang, J., Gu, L., Zhang, Y., Yan, T., Kong, G., Kong, L., Guo, B., Qiu, M., Wang, Y., Jing, M.,
Xing, W., Ye, W., Wu, Z., Zhang, Z., Zheng, X., Gijzen, M., Wang, Y., and Dong, S. (2017).
An oomycete plant pathogen reprograms host pre-mRNA splicing to subvert immunity. Nature
Communications **8**.

Huang, S., Van Der Vossen, E.A.G., Kuang, H., Vleeshouwers, V.G.A.A., Zhang, N., Borm, T.J.A.,
Van Eck, H.J., Baker, B., Jacobsen, E., and Visser, R.G.F. (2005). Comparative genomics
enabled the isolation of the R3a late blight resistance gene in potato. The Plant Journal **42,** 251-
261.

James M. Bradeen, M.I., Dimitre S. Mollov, John Raasch, Lara Colton Kramer, Benjamin P .
Millett, Sandra Austin-Phillips, Jiming Jiang, and Domenico Carputo. (2009). Higher Copy
Numbers of the Potato *RB* Transgene Correspond to Enhanced Transcript and Late Blight
Resistance Levels. MPMI **22,** 437–446.

Jones, J.D., and Dangl, J.L. (2006). The plant immune system. Nature **444,** 323-329.

Jung, H.W., Panigrahi, G.K., Jung, G.Y., Lee, Y.J., Shin, K.H., Sahoo, A., Choi, E.S., Lee, E., Man
Kim, K., Yang, S.H., Jeon, J.S., Lee, S.C., and Kim, S.H. (2020). Pathogen-associated
molecular pattern-triggered immunity involves proteolytic degradation of core nonsense-mediated
mRNA decay factors during the early defense response. Plant Cell **32,** 1081-1101.

Kamoun, S., and Smart, C.D. (2005). Late Blight of Potato and Tomato in the Genomics Era. Plant
Disease **89,** 692-699.

Kamoun, S., Furzer, O., Jones, J.D.G., Judelson, H.S., Ali, G.S., Dalio, R.J.D., Roy, S.G., Schena, L.,
Zambounis, A., Panabieres, F., Cahill, D., Ruocco, M., Figueiredo, A., Chen, X.R., Hulvey, J.,

Stam, R., Lamour, K., Gijzen, M., Tyler, B.M., Grunwald, N.J., Mukhtar, M.S., Tome, D.F.A., Tor, M., Van den Ackerveken, G., McDowell, J., Daayf, F., Fry, W.E., Lindqvist-Kreuze, H., Meijer, H.J.G., Petre, B., Ristaino, J., Yoshida, K., Birch, P.R.J., and Govers, F. (2015). The Top 10 oomycete pathogens in molecular plant pathology. Molecular Plant Pathology **16,** 413-434.

Karki, H.S., Jansky, S.H., and Halterman, D.A. (2021). Screening of Wild Potatoes Identifies New Sources of Late Blight Resistance. Plant Disease **105,** 368-376.

Katou, S., Yoshioka, H., Kawakita, K., Rowland, O., Jones, J.D.G., Mori, H., and Doke, N. (2005). Involvement of PPS3 Phosphorylated by Elicitor-Responsive Mitogen-Activated Protein Kinases in the Regulation of Plant Cell Death. Plant Physiology **139,** 1914-1926.

Klepikova, A.V., Kasianov, A.S., Gerasimov, E.S., Logacheva, M.D., and Penin, A.A. (2016). A high resolution map of the Arabidopsis thaliana developmental transcriptome based on RNA-seq profiling. The Plant Journal **88,** 1058-1070.

Kotake, Y., Sagane, K., Owa, T., Mimori-Kiyosue, Y., Shimizu, H., Uesugi, M., Ishihama, Y., Iwata, M., and Mizui, Y. (2007). Splicing factor SF3b as a target of the antitumor natural product pladienolide. Nat Chem Biol **3,** 570-575.

Lai, Y., and Eulgem, T. (2018). Transcript-level expression control of plant NLR genes. Mol Plant Pathol **19,** 1267-1281.

Lara C. Kramer, M.J.C., Susan M. Wielgus, Pudota B. Bhaskar, and Jiming Jiang. (2009). Correlation Between Transcript Abundance of the RB Gene and the Level of the RB-Mediated Late Blight Resistance in Potato. MPMI.

Li, F., Pignatta, D., Bendix, C., Brunkard, J.O., Cohn, M.M., Tung, J., Sun, H., Kumar, P., and Baker, B. (2012). MicroRNA regulation of plant innate immune receptors. Proceedings of the National Academy of Sciences USA **109,** 1790-1795.

Li, X., Kapos, P., and Zhang, Y. (2015). NLRs in plants. Curr Opin Immunol **32,** 114-121.

Liang, W., van Wersch, S., Tong, M., and Li, X. (2019). TIR-NB-LRR immune receptor SOC3 pairs with truncated TIR-NB protein CHS1 or TN2 to monitor the homeostasis of E3 ligase SAUL1. New Phytologist **221,** 2054-2066.

Liao, S.E., and Regev, O. (2021). Splicing at the phase-separated nuclear speckle interface: a model. Nucleic Acids Res **49,** 636-645.

Liu, N., Hake, K., Wang, W., Zhao, T., Romeis, T., and Tang, D. (2017). CALCIUM-DEPENDENT PROTEIN KINASE5 Associates with the Truncated NLR Protein TIR-NBS2 to Contribute to exo70B1-Mediated Immunity. Plant Cell **29,** 746-759.

Lu, Q., Tang, X., Tian, G., Wang, F., Liu, K., Nguyen, V., Kohalmi, S.E., Keller, W.A., Tsang, E.W.T., Harada, J.J., Rothstein, S.J., and Cui, Y. (2010a). Arabidopsis homolog of the yeast TREX-2 mRNA export complex: components and anchoring nucleoporin. The Plant Journal **61,** 259-270.

Lu, T., Lu, G., Fan, D., Zhu, C., Li, W., Zhao, Q., Feng, Q., Zhao, Y., Guo, Y., Li, W., Huang, X., and Han, B. (2010b). Function annotation of the rice transcriptome at single-nucleotide resolution by RNA-seq. Genome Research **20,** 1238-1249.

Maekawa, T., Cheng, W., Spiridon, L.N., Toller, A., Lukasik, E., Saijo, Y., Liu, P., Shen, Q.H., Micluta, M.A., Somssich, I.E., Takken, F.L.W., Petrescu, A.J., Chai, J., and Schulze-Lefert, P. (2011). Coiled-coil domain-dependent homodimerization of intracellular barley immune receptors defines a minimal functional module for triggering cell death. Cell Host Microbe **9,** 187-199.

Marquez, Y., Brown, J.W.S., Simpson, C., Barta, A., and Kalyna, M. (2012). Transcriptome survey reveals increased complexity of the alternative splicing landscape in Arabidopsis. Genome

Research **22,** 1184-1195.

Mittler, R., Herr, E.H., Orvar, B.L., van Camp, W., Willekens, H., Inzé, D., and Ellis, B.E. (1999).
Transgenic tobacco plants with reduced capability to detoxify reactive oxygen intermediates are
hyperresponsive to pathogen infection. Proceedings of the National Academy of Sciences **96,**
14165-14170.

Ning, Y., Liu, W., and Wang, G.L. (2017). Balancing Immunity and Yield in Crop Plants. Trends Plant
Sci **22,** 1069-1079.

Nishimura, M.T., Anderson, R.G., Cherkis, K.A., Law, T.F., Liu, Q.L., Machius, M., Nimchuk, Z.L.,
Yang, L., Chung, E.H., El Kasmi, F., Hyunh, M., Osborne Nishimura, E., Sondek, J.E., and
Dangl, J.L. (2017a). TIR-only protein RBA1 recognizes a pathogen effector to regulate cell death
in Arabidopsis. Proc Natl Acad Sci U S A **114,** E2053-E2062.

Nishimura, M.T., Anderson, R.G., Cherkis, K.A., Law, T.F., Liu, Q.L., Machius, M., Nimchuk, Z.L.,
Yang, L., Chung, E.H., El Kasmi, F., Hyunh, M., Osborne Nishimura, E., Sondek, J.E., and
Dangl, J.L. (2017b). TIR-only protein RBA1 recognizes a pathogen effector to regulate cell death
in Arabidopsis. Proceedings of the National Academy of Sciences USA **114,** E2053-E2062.

Ofir, G., Herbst, E., Baroz, M., Cohen, D., Millman, A., Doron, S., Tal, N., Malheiro, D.B.A.,
Malitsky, S., Amitai, G., and Sorek, R. (2021). Antiviral activity of bacterial TIR domains via
immune signalling molecules. Nature **600,** 116-120.

Palma, K., Zhao, Q., Cheng, Y.T., Bi, D., Monaghan, J., Cheng, W., Zhang, Y., and Li, X. (2007).
Regulation of plant innate immunity by three proteins in a complex conserved across the plant and
animal kingdoms. Genes & Development **21,** 1484-1493.

Paluchowska, P., Sliwka, J., and Yin, Z. (2022). Late blight resistance genes in potato breeding. Planta
255, 127.

Rairdan, G.J., Collier, S.M., Sacco, M.A., Baldwin, T.T., Boettrich, T., and Moffett, P. (2008). The
Coiled-Coil and Nucleotide Binding Domains of the Potato Rx Disease Resistance Protein
Function in Pathogen Recognition and Signaling The Plant Cell **20,** 739-751.

Reddy, A.S. (2007). Alternative splicing of pre-messenger RNAs in plants in the genomic era. Annu Rev
Plant Biol **58,** 267-294.

Reddy, A.S.N. (2004). Plant serine/arginine-rich proteins and their role in pre-mRNA splicing. Trends in
Plant Science **9,** 541-547.

Reddy, A.S.N., Marquez, Y., Kalyna, M., and Barta, A. (2013a). Complexity of the Alternative
Splicing Landscape in Plants. . The Plant Cell **25,** 3657-3683.

Reddy, A.S.N., Marquez, Y., Kalyna, M., and Barta, A. (2013b). Complexity of the Alternative
Splicing Landscape in Plants. The Plant Cell **25,** 3657-3683.

Reddy, A.S.N., Huang, J., Syed, N.H., Ben-Hur, A., Dong, S., and Gu, L. (2020). Decoding co-/post-
transcriptional complexities of plant transcriptomes and epitranscriptome using next-generation
sequencing technologies. Biochemical Society Transactions **48,** 2399-2414.

Sanchez-Martin, J., Widrig, V., Herren, G., Wicker, T., Zbinden, H., Gronnier, J., Sporri, L., Praz,
C.R., Heuberger, M., Kolodziej, M.C., Isaksson, J., Steuernagel, B., Karafiatova, M., Dolezel,
J., Zipfel, C., and Keller, B. (2021). Wheat Pm4 resistance to powdery mildew is controlled by
alternative splice variants encoding chimeric proteins. Nat Plants **7,** 327-341.

Sheth, N., Roca, X., Hastings, M.L., Roeder, T., Krainer, A.R., and Sachidanandam, R. (2006).
Comprehensive splice-site analysis using comparative genomics. Nucleic Acids Res **34,** 3955-
3967.

Slane, D., Lee, C.H., Kolb, M., Dent, C., Miao, Y., Franz-Wachtel, M., Lau, S., Macek, B., Balasubramanian, S., Bayer, M., and Jurgens, G. (2020). The integral spliceosomal component CWC15 is required for development in Arabidopsis. Sci Rep **10,** 13336.

Song, J., Bradeen, J.M., Naess, S.K., Raasch, J.A., Wielgus, S.M., Haberlach, G.T., Liu, J., Kuang, H., Austin-Phillips, S., Buell, C.R., Helgeson, J.P., and Jiang, J. (2003). Gene RB cloned from Solanum bulbocastanum confers broad spectrum resistance to potato late blight. Proc Natl Acad Sci U S A **100,** 9128-9133.

Stam, M., Mol, J.N.M., and Kooter, J.M. (1997). The silence of genes in transgenic plants. Annals of Botany **79,** 3-12.

Sukarta, O.C.A., Zheng, Q., Slootweg, E.J., Mekken, M., Mendel, M., Putker, V., Bertran, A., Brand, A., Overmars, H., Pomp, R., Roosien, J., Boeren, S., Smant, G., and Goverse, A. (2022). GLYCINE-RICH RNA-BINDING PROTEIN 7 potentiates effector-triggered immunity through an RNA recognition motif. Plant Physiol **189,** 972-987.

Sun, H., Jiao, W.B., Krause, K., Campoy, J.A., Goel, M., Folz-Donahue, K., Kukat, C., Huettel, B., and Schneeberger, K. (2022). Chromosome-scale and haplotype-resolved genome assembly of a tetraploid potato cultivar. Nat Genet **54,** 342-348.

Thilliez, G.J.A., Armstrong, M.R., Lim, T.-Y., Baker, K., Jouet, A., Ward, B., van Oosterhout, C., Jones, J.D.G., Huitema, E., Birch, P.R.J., and Hein, I. (2019). Pathogen enrichment sequencing (PenSeq) enables population genomic studies in oomycetes. New Phytologist **221,** 1634-1648.

Tsuchiya, T., and Eulgem, T. (2013). An alternative polyadenylation mechanism coopted to the Arabidopsis RPP7 gene through intronic retrotransposon domestication. Proceedings of the National Academy of Sciences USA **110,** E3535-3543.

Van de Weyer, A.L., Monteiro, F., Furzer, O.J., Nishimura, M.T., Cevik, V., Witek, K., Jones, J.D.G., Dangl, J.L., Weigel, D., and Bemm, F. (2019). A Species-Wide Inventory of NLR Genes and Alleles in Arabidopsis thaliana. Cell **178,** 1260-1272 e1214.

van der Hoorn, R.A., and Kamoun, S. (2008). From Guard to Decoy: a new model for perception of plant pathogen effectors. Plant Cell **20,** 2009-2017.

van der Vossen, E., Sikkema, A., Hekkert, B., Gros, J., Stevens, P., Muskens, M., Wouters, D., Pereira, A., Stiekema, W., and Allefs, S. (2003). An ancient R gene from the wild potato species Solanum bulbocastanum confers broad-spectrum resistance to Phytophthora infestans in cultivated potato and tomato. Plant J **36,** 867-882.

van der Vossen, E.A., Gros, J., Sikkema, A., Muskens, M., Wouters, D., Wolters, P., Pereira, A., and Allefs, S. (2005). The Rpi-blb2 gene from Solanum bulbocastanum is an Mi-1 gene homolog conferring broad-spectrum late blight resistance in potato. Plant J **44,** 208-222.

Vidal, M., and Legrain, P. (1999). Yeast forward and reverse 'n'-hybrid systems. Nucleic Acids Research **27,** 919-929.

Vleeshouwers, V.G., Rietman, H., Krenek, P., Champouret, N., Young, C., Oh, S.K., Wang, M., Bouwmeester, K., Vosman, B., Visser, R.G., Jacobsen, E., Govers, F., Kamoun, S., and Van der Vossen, E.A. (2008). Effector genomics accelerates discovery and functional profiling of potato disease resistance and phytophthora infestans avirulence genes. PLoS One **3,** e2875.

Wan, R., Bai, R., Zhan, X., and Shi, Y. (2020). How Is Precursor Messenger RNA Spliced by the Spliceosome? Annu Rev Biochem **89,** 333-358.

Wang, J., Hu, M., Wang, J., Qi, J., Han, Z., Wang, G., Qi, Y., Wang, H.W., Zhou, J.M., and Chai, J. (2019). Reconstitution and structure of a plant NLR resistosome conferring immunity. Science

364.

Wang, L., Yang, T., Wang, B., Lin, Q., and Yu, F. (2020). RALF1-FERONIA complex affects splicing
dynamics to modulate stress responses and growth in plants. Science Advances **6,** eaaz1622.

Wang, M., Allefs, S., van den Berg, R.G., Vleeshouwers, V.G., van der Vossen, E.A., and Vosman, B.
(2008). Allele mining in Solanum: conserved homologues of Rpi-blb1 are identified in Solanum
stoloniferum. Theor Appl Genet **116,** 933-943.

Wang, S., McLellan, H., Bukharova, T., He, Q., Murphy, F., Shi, J., Sun, S., van Weymers, P., Ren,
Y., Thilliez, G., Wang, H., Chen, X., Engelhardt, S., Vleeshouwers, V., Gilroy, E.M., Whisson,
S.C., Hein, I., Wang, X., Tian, Z., Birch, P.R.J., and Boevink, P.C. (2018). Phytophthora
infestans RXLR effectors act in concert at diverse subcellular locations to enhance host
colonization. Journal of Experimental Botany **70,** 343-356.

Will, C.L., and Lührmann, R. (2011). Spliceosome structure and function. Cold Spring Harb Perspect
Biol **3**.

Wu, Z., Zhu, D., Lin, X., Miao, J., Gu, L., Deng, X., Yang, Q., Sun, K., Zhu, D., Cao, X., Tsuge, T.,
Dean, C., Aoyama, T., Gu, H., and Qu, L.J. (2016). RNA Binding Proteins RZ-1B and RZ-1C
Play Critical Roles in Regulating Pre-mRNA Splicing and Gene Expression during Development
in Arabidopsis. The Plant Cell **28,** 55-73.

Xu, F., Xu, S., Wiermer, M., Zhang, Y., and Li, X. (2012). The cyclin L homolog MOS12 and the
MOS4-associated complex are required for the proper splicing of plant resistance genes. The plant
jouranl **70,** 916-928.

Xu, S., Zhang, Z., Jing, B., Gannon, P., Ding, J., Xu, F., Li, X., and Zhang, Y. (2011). Transportin-SR
is required for proper splicing of resistance genes and plant immunity. PLOS Genetics **7,**
e1002159.

Yang, S., Tang, F., and Zhu, H. (2014). Alternative splicing in plant immunity. International journal of
molecular sciences **15,** 10424-10445.

Yu, X., Feng, B., He, P., and Shan, L. (2017). From Chaos to Harmony: Responses and Signaling upon
Microbial Pattern Recognition. Annual Review of Phytopathology **55,** 109-137.

Zhang, F., Fang, H., Wang, M., He, F., Tao, H., Wang, R., Long, J., Wang, J., Wang, G.L., and Ning,
Y. (2022). APIP5 functions as a transcription factor and an RNA-binding protein to modulate cell
death and immunity in rice. Nucleic Acids Res **50,** 5064-5079.

Zhang, F., Chen, H., Zhang, X., Gao, C., Huang, J., Lü, L., Shen, D., Wang, L., Huang, C., Ye, W.,
Zheng, X., Wang, Y., Vossen, J.H., and Dong, S. (2021). Genome Analysis of Two Newly
Emerged Potato Late Blight Isolates Sheds Light on Pathogen Adaptation and Provides Tools for
Disease Management. Phytopathology® **111,** 96-107.

Zhang, X.-C., and Gassmann, W. (2007). Alternative splicing and mRNA levels of the disease
resistance gene *RPS4* are induced during defense responses. Plant Physiology **145,** 1577-1587.

Zhang, Y., and Li, X. (2005). A putative nucleoporin 96 is required for both basal defense and
constitutive resistance responses mediated by *suppressor of npr1-1, constitutive 1*. The Plant Cell
17, 1306-1316.

Zhang, Y., Xu, S., Ding, P., Wang, D., Cheng, Y.T., He, J., Gao, M., Xu, F., Li, Y., Zhu, Z., Li, X., and
Zhang, Y. (2010). Control of salicylic acid synthesis and systemic acquired resistance by two
members of a plant-specific family of transcription factors. Proc Natl Acad Sci U S A **107,** 18220-
18225.

Zhao, J., and Song, J. (2021). NLR immune receptor RB is differentially targeted by two homologous

but functionally distinct effector proteins. Plant Communications **2**.

Zhou, J.-M., and Chai, J. (2008). Plant pathogenic bacterial type III effectors subdue host responses.
Current Opinion in Microbiology **11,** 179-185.

Zhou, J.-M., and Zhang, Y. (2020). Plant Immunity: Danger Perception and Signaling. Cell **181,** 978-
989.

Zhu, S., Vossen, J.H., Bergervoet, M., Nijenhuis, M., Kodde, L., Kessel, G.J.T., Vleeshouwers, V.,
Visser, R.G.F., and Jacobsen, E. (2014). An updated conventional- and a novel GM potato late
blight R gene differential set for virulence monitoring of Phytophthora infestans. Euphytica **202,**
219-234.

Zhu, S., Vossen, J.H., Bergervoet, M., Nijenhuis, M., Kodde, L., Kessel, G.J.T., Vleeshouwers, V.,
Visser, R.G.F., and Jacobsen, E. (2015). An updated conventional- and a novel GM potato late
blight R gene differential set for virulence monitoring of Phytophthora infestans. Euphytica **202,**
219-234.

Figure 1. The AS of *RB* is specifically modulated by *P. infestans* infection.
(A) Gene model of *RB* and splicing mutant *RB_IVS*. The arrows indicate the primers used in (B). Red letters in *RB_IVS* denote mutations to change the intron splicing recognition sites. The upper is schematic of two transcripts of *RB* amplified using 5′-rapid amplification of cDNA ends (5′-RACE). UPM primer is Universal Primer A Mix, GSP primer is gene specific primer of *RB*. The white vertical lines represent single nucleotide polymorphisms that appeared in the sequencing results (Left panel). *RB* alternative splicing produced splicing isoforms. The asterisk shows the location of a premature termination codon. Diagrams showing the *RB* encoded predicted proteins and with domains indicated by color (Right panel). **(B)** RT-PCR analysis of *RB_IR* and *RB_CDS* in wildtype (WT), *RB* and *RB_IVS* transgenic *S. tuberosum* and *N. benthamiana*, PT29 (*S. bulbocastanum*); and AS events of *Rpi-sto1* in *Rpi-sto1* transgenic potato (*S. tuberosum*). *RB_IR* is 995 bp, and *RB_CDS* is 316 bp. *StEF1α* and *NbEF1α* were used as internal controls. **(C)** Expression of N terminal GFP-tagged fusions of RB, RB_CDS, and RB_IR in agroinfiltrated *N. benthamiana* was

confirmed by protein blot using an anti-GFP antibody. The asterisks indicate the predicted sizes of RB_CDS and RB_IR. Equal protein loading is visualized by Ponceau S staining. **(D)** Splicing ratio of *RB* in transgenic potato following inoculation with water (mock), *P. infestans* JH19, or *P. capsici* LT1534. Transcript levels were detected by RT-qPCR and normalized to *RB_CDS* following mock treatment at 0 dpi. Bars represent the mean ± SD from three independent replicates. **(E)** RT-PCR of *RB_IR* or *RB_CDS* variants in *RB* transgenic potato following inoculation with different *P. infestans* strains (JH19, HB1501, T30-4), water (mock) and *P. capsici* strain LT1534. *StEF1α* was used as the internal control. The intensities under the PCR bands of each product were quantified by ImageJ software. **(F)** Splicing ratio of *RB* isoforms in *RB_IVS* transgenic potato following inoculation with *P. infestans* zoospores. RNA was extracted 0, 1, 2 days post inoculation. Splicing ratio levels were detected by RT-qPCR. Bars represent the mean ± SD in three independent replicates. **(G)** Splicing ratio of *RB* in *RB* transgenic *N. benthamiana* following agroinfiltration of RNAi-hairpin constructs then inoculation with *P. infestans* zoospores 2 day later. RNA was extracted 0 day and 2 days after pathogen inoculation. Splicing ratio levels were detected by RT-qPCR. Bars represent the mean ± SD in three independent replicates. **(H)** Splicing ratio of *RB* in *RB* transgenic *N. benthamiana* following treatment with DMSO (control), 1 μM PB, or 30 μM PB followed by inoculation with *P. infestans* zoospores 6 hours later. RNA was extracted at 0 day and 2 days post inoculation (dpi). Splicing ratio levels were detected by RT-qPCR and compared to *RB* splicing ratio at 0 dpi. ns, no significant difference. One asterisk denotes a significant difference at P<0.05, and three asterisks denotes a significant difference at P<0.001 using a two-way ANOVA with Tukey's test. The above experiments were repeated 3 times with similar results.

Figure 2. Intron retention in *RB* results in susceptibility to *P. infestans* infection.
(A) Zoospore suspension of *P. infestans* strain JH19 was used to inoculate *RB*, *RB_CDS*, *RB_IR* and *RB_IVS* transgenic potato plants at a concentration of 15,000 zoospores/mL. Plants were photographed at 5 dpi. For every transgenic potato, each line has nine repeats (n=9). The numbers represent the number of resistant plants compared to all inoculated plants. Bar=5 cm. **(B)** Zoospore suspension of *P. infestans* strain JH19 was drop-inoculated on detached leaves of *RB* transgenic potato expressing the indicated gene variants. The infected leaves were photographed at 4 dpi. **(C)** Infected areas (%) of inoculated leaves in (B). The infected area was measured in ImageJ. The data represent the mean ± SD from 12 replicates. Different letters denote a significant difference at P<0.001 using a two-way ANOVA with Dunn's test. **(D)** Trypan blue staining of *RB* transgenic *N. benthamiana* following agroinfiltration of RNAi-hairpin constructs and inoculation with *P. infestans* zoospores. *EV* = empty vector control. Leaves were photographed at 5 dpi. **(E)** Lesion areas

(cm^2) of inoculated leaves in (D). Data represent the mean ± SD from 8 replicates. **(F)** Relative biomass of *P. infestans* was determined by qPCR of *P. infestans* genomic DNA normalized to potato genomic DNA at 5 days post inoculation (dpi). Bars represent the mean ± SD from three independent replicates. Different letters above bars denote a significant difference at P<0.001 using a one-way ANOVA with Dunn's test. **(G)** Zoospores of *P. infestans* strain JH19 were inoculated on *RB* or *RB_CDS* transgenic *N. benthamiana* leaves pretreated with DMSO or 30 μM Pladienolide B (PB). Leaves were stained with trypan blue and photographed at 5 dpi. **(H)** Lesion areas (cm^2) of inoculated leaves in (D). Data represent the mean ± SD from 14 replicates. **(I)** Relative biomass of *P. infestans* was determined by qPCR of *P. infestans* genomic DNA normalized to potato genomic DNA at 5 days post inoculation (dpi). Bars represent the mean ± SD from three independent replicates. ns = no significant difference and three asterisks denotes a statistical difference at P<0.001 using a two-way ANOVA with Tukey's test. The above experiments were repeated 3 times with similar results.

Figure 3. IPI-O1 induces *RB* intron splicing.

(A) Schematic diagram of the *RB_LUC* alternative splicing reporter system. The genomic sequence of *RB* was fused with the *luciferase* (*LUC*) gene to construct *35S::RB-LUC*. Intron retention introduces a premature termination codon (PTC) and no production of the luciferase reporter. Intron splicing leads to an entire RB-LUC protein with luciferase activity. The *35S::RB_CDS-LUC* construct with the intron removed produces a full-length RB-LUC protein with luciferase activity regardless of intron splicing activity. **(B)** Results of co-expression of the reporters with multiple splicing regulatory effectors (SRE) that we previously identified (Huang et al., 2020). LUC activity was quantified at 48 hpi. Relative LUC activity was normalized to the activity of the *35S::RB_CDS-LUC* construct with mock treatment. Data represent the mean ± SD from 24 replicates. Different letters above each bar represent a significant difference at P<0.001 using a two-way ANOVA with Dunn's test. **(C)** GFP-tagged SRE fusion protein expression in *N. benthamiana* was confirmed by western blotting using an anti-GFP antibody. Equal protein loading was visualized by Ponceau S staining. **(D)** Transcript levels of *RB_IR* and *RB_CDS* were determined by RT-

PCR following agroinfiltration of IPI-O1, IPI-O2 and IPI-O4 into leaves of *RB* transgenic *N. benthamiana*. *NbEF1α* was used as the internal control. **(E)** Bars represent relative intensity of RT-PCR products in (D) and normalized with internal control *NbEF1α*. Intensities of the bands were quantified by ImageJ software, and the relative abundances of *RB* isoforms are relative to *RB_CDS* at 0 dpi. Bars represent the mean $\pm$ SD from three independent replicates. ns = no significant difference, one asterisk denotes a significant difference at $P<0.05$, two asterisks denotes a significant difference at $P<0.01$ and three asterisks denotes a significant difference at $P<0.001$ using a two-way ANOVA with Tukey's test. The above experiments were repeated 3 times with similar results.

Figure 4. NLS motif of IPI-O1 is necessary for *RB* splicing modulation.
(A) Schematic diagrams at the top show the location of alanine replacements within the nuclear localization signal of IPI-O1. **(B)** GFP-IPI-O1 and GFP-IPI-O1^{nls} were expressed in *N. benthamiana* leaves for 48 h before imaging, Scale bar = 20 μm. **(C)** Bars represent the number of times that GFP was detected in the nucleoplasm or nucleolus following infiltration of GFP-IPI-O1 and GFP-IPI-O1^{nls}. Data represent the mean ± SD of three replicates. **(D)** Expression of GFP-tagged fusions of IPI-O1 and IPI-O1^{nls} in *N. benthamiana* was confirmed by protein blotting using an anti-GFP antibody. Equal protein loading was visualized by Ponceau S staining. **(E)** Results of co-expression of RB LUC reporters with mock, GFP, IPI-O1 and IPI-O1^{nls} in *N. benthamiana* leaves by agroinfiltration. LUC activity was quantified at 2 dpi. Relative LUC activity was normalized to the activity of the *LUC* construct with mock treatment. Data represent the mean ± SD of 30 replicates. For all graphs, ns = no significant difference and three asterisks denote a statistical

difference at P<0.001 using a two-way ANOVA with Tukey's test. **(F)** Splicing ratio of *RB* in *RB* transgenic *N. benthamiana* following agroinfiltration of IPI-O1 and IPI-O1^{nls} at 0, 1 and 2 days by RT-qPCR. Bars represent the mean ± SD of three independent replicates. **(G)** The image on the left shows protein blot analysis of the protein accumulation of RB_CDS and RB_IR following transient expression of the indicated genes in *N. benthamiana* leaves. On the right, RFP tag fusions with IPI-O1 and IPI-O1^{nls} were confirmed by protein blotting using an anti-RFP antibody. Equal protein loading is visualized by Ponceau S staining. The above experiments were repeated 3 times with similar results.

Figure 5. IPI-O1 physically interacts with splicing factor CWC15.
(A) Immunoprecipitation (IP) of StCWC15a/b-GFP with RFP-IPI-O1, RFP-IPI-O1^{nls}, or RFP-Pi04314 following agroinfiltration in *N. benthamiana* leaves. Expression of constructs in the leaves is indicated by +. Markers are indicated on the right, and the Ponceau S staining was used to show levels of protein loading. **(B)** *In vitro* immunoprecipitation (IP) using the GST-tagged StCWC15a/b and IPI-O1 or IPI-O1^{nls}. Input and GST-pull-down proteins were visualized by protein blotting using anti-GST and anti-His antibodies. Asterisks (*) indicate the protein bands that correspond to the predicted size of GST-CWC15a/b. **(C)** IPI-O1 interacts with StCWC15 in the nucleolus and nuclear speckles. Split-YFP/BiFC of IPI-O1 or IPI-O1^{nls} and Pi04314 with StCWC15a and b using fluorescence imaging. Samples were stained with 2 ug/mL DAPI to indicate the nucleus. Numbers represent the frequency of cells showing YFP fluorescence compared to the total number of cells in the 25 mm^2 discs from three biological replicates. Bar = 5 μm. **(D)** IPI-O1 co-localizes with StCWC15a/b in nuclear speckles and the nucleolus. GFP-IPI-O1, GFP-IPI-O1^{nls}, and GFP alone were transiently co-expressed with StCWC15a/b-RFP by agroinfiltration in *N. benthamiana*. The left image is from the GFP channel, the middle image is from the RFP channel, and the right image is the overlay of the GFP and RFP channels. Numbers represent the frequency of cells showing indicated StCWC15 subcellular localization compared to the total StCWC15-positive cells in 25mm^2 discs from three biological replicates. Graphs show the fluorescence intensity profiles across the arrows in the merged images. Bar = 5μm.

Figure 6. Silencing *StCWC15* disrupts *RB*-mediated resistance.
(A) Whole plants of *RB* and *RB-StCWC15* RNAi transgenic were spray-inoculated with zoospores of *P. infestans* strain JH19 at a concentration of 15,000 zoospores/mL. Plants were photographed at 5 dpi. For every transgenic potato, each line has nine repeats (n=9). Bar = 3 cm. The numbers represent resistant plants compared to all inoculated plants. **(B)** Zoospore suspension of *P. infestans* strain JH19 was drop-inoculated on detached leaves of *RB* and *RB-StCWC15* RNAi transgenic potato lines. The infected leaves were photographed at 4 dpi. The numbers represent the number of resistant leaves compared to all inoculated leaves. **(C)** Relative expression of *StCWC15a/b* in three RNAi transformed lines compared to *RB* transgenic potato. Bars represent the mean ± SD of three independent replicates. Three asterisks designate a statistical difference from the *RB* control at P<0.001. **(D)** Infected areas (%) of inoculated leaves in (B). The infected area was measured in ImageJ. The data represent the mean ± SD from 10 replicates. Different letters denote a significant difference at P<0.001 using a two-way ANOVA with Dunn's test. **(E)** Splicing ratio of *RB* in *RB-StCWC15* RNAi transgenic potato line#2 following inoculation with *P. infestans* zoospores at 2 dpi by RT-qPCR. Bars represent the mean ± SD from three independent replicates. ns=no significant difference and three asterisks denote a statistical difference at P<0.001. **(F)** *NbCWC15* was silenced by VIGS in *RB*, *RB_CDS*, and *Rpi_blb2* transgenic *N. benthamiana* and inoculated with *P. infestans* JH19 on detached leaves. Leaves were stained with trypan blue and photographed at 5 dpi. **(G)** Lesion areas(cm^2) of inoculated leaves in (F) were calculated using ImageJ. Data represent the mean ± SD from 18 replicates. **(H)** Relative biomass of *P. infestans* was determined by qPCR of *P. infestans* genomic DNA normalized to potato genomic DNA at 5 days post inoculation (dpi). Bars represent the mean ± SD from three independent replicates. Three asterisks designate a statistical difference from the control at P<0.001 using a two-way ANOVA with Tukey's test. **(I)** Complementation of *RB* function caused by *CWC15* silencing in *N. benthamiana* via synthetic StCWC15 constructs (StCWC15a/b^{syn}). Agrobacterium expressing StCWC15a/b^{syn} constructs were infiltrated into *NbCWC15* silenced *RB* transgenic *N. benthamiana*, then inoculated with *P. infestans* JH19 zoospores. Leaves were stained with trypan blue and photographed at 5 dpi. **(J)** Lesion areas(cm^2) of inoculated leaves in (I) were calculated using ImageJ. Data represent the mean ± SD from 8 replicates. **(K)** Relative biomass of *P. infestans* was determined by qPCR of *P. infestans* genomic DNA normalized to potato genomic DNA at 5 days post inoculation (dpi). Bars represent the mean± SD from three independent replicates. Different letters denote a significance difference at P<0.001 using one-way ANOVA with Dunn's test.

Figure 7. Alternative intron splicing of *RB* balances plant growth and immunity.
(A) Photos of 21-day-old wild-type (WT) Favorita and *RB*, *RB_CDS*, and *RB_IR* transgenic tissue culture plants. Bar = 1 cm. **(B)** Average stem height and **(C)** root length of Favorita (WT) and 21-day-old transgenic potato tissue culture plants. Bars represent the mean ± SD from 5 tissue culture plants of 3 independent

lines for each genotype. **(D)** Total tuber mass and **(E)** individual tuber mass of tubers from greenhouse grown plants. Measurements were taken at 8 to 10 weeks after tissue culture plants were transplanted. Bars represent the mean ± SD from 5 replicates of 3 independent lines for each genotype. **(F)** Total salicylic acid (SA) content in Favorita (WT) and transgenic *RB* and *RB_CDS* lines without pathogen inoculation. FW, fresh weight. Bars represent the mean ± SD from three independent replicates. **(G-H)** Relative expression of *SARD1* (G) and *PR1* (H) in the indicated transgenic potato as determined by RT-qPCR. Bars represent the mean ± SD from three independent replicates. For all graphs, different letters represent a statistical difference at $P<0.001$ using one-way ANOVA with Dunn's test.

Fig.8 Overexpression RB_IR disrupts RB_CDS self-association and immunity activation.
(A) Immunoprecipitation (IP) of RB_CDS-GFP and RB-CC-GFP with RB_IR-RFP, R2-CC-RFP following agroinfiltration in *N. benthamiana* leaves. Expression of constructs in the leaves is indicated by +. The Ponceau S staining was used to show levels of protein loading. **(B)** Zoospores of *P. infestans* strain JH19 were inoculated on *RB_CDS* transgenic *N. benthamiana* leaves expressing with the indicated proteins by

agroinfiltration. Leaves were stained with trypan blue and photographed at 5 dpi. **(C)** Lesion areas(cm^2) of inoculated leaves in (B) were calculated using ImageJ. Data represent the mean ± SD from 10 replicates from three independent experiments. **(D)** Relative biomass of *P. infestans* was determined by qPCR of *P. infestans* genomic DNA normalized to *N. benthamiana* genomic DNA at 5 dpi. Bars represent the mean ± SD from three independent replicates. **(E)** Agrobacterium carrying RB-CC-GFP and RB-CC-RFP (OD$_{600}$ = 0.3) were co-expressed in *N. benthamiana* in the presence of GFP-10×Myc or RB_IR-10×Myc or R2-CC-10×Myc (OD$_{600}$ = 0.6) by agroinfiltration. Total protein was incubated with GFP-Trap beads for co-IP assays, and immunoprecipitated proteins were analyzed using anti-GFP, anti-RFP or anti-10×Myc antibodies. Ponceau S staining was used to show levels of protein loading. These experiments were repeated three times with similar results.

Fig. 9 Schematic model of the P. infestans host pathosystem related to resistance gene RB and effector IPI-O1.

During normal growth (left), CWC15 proteins (grey triangles) are maintained in the nucleoplasm and *RB* transcripts exist primarily with the intron retained (*RB_IR*). This results in the production of a truncated, non-functional RB protein that supports growth over immunity. Following *P. infestans* infection, effector IPI-O1 is introduced into the plant cell and translocated to the nucleus, leading to a re-localization of CWC15 from the nucleoplasm to the nucleolus and speckles. This results in increased splicing of the *RB* intron (*RB_CDS*), resulting in production of a full-length RB protein that can elicit immunity.