HES1 and *ID3* temporally regulate p27 levels and endothelial cell flow-mediated quiescence depth

Natalie T Tanke[a], Ziqing Liu[b], Tianji Yu[b], Morgan Oatley[b], Pauline Bougaran[b], Kaitlyn Quigley[b], Jeanette Gowen Cook[c], and Victoria L Bautch[a,b,d,#]

[a]Curriculum in Cell Biology and Physiology, The University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599, USA
[b]Department of Biology, The University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599, USA
[c]Department of Biochemistry and Biophysics, The University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599, USA
[d]McAllister Heart Institute, The University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599, USA

[#]Corresponding author: Victoria L Bautch, PhD
                       Professor of Biology
                       Department of Biology, CB#3280
                       University of North Carolina at Chapel Hill
                       Chapel Hill, NC 27599 USA
                       Email: bautch@med.unc.edu

Running head: Endothelial cell quiescence depth varies with time

Words: 8,293 (main text and references)
Abstract: 197 words
Figures: 6
Supplemental Figures: 6
Supplemental Tables: 6
References: 68

Keywords: Endothelial cell, laminar shear stress, quiescence depth, p27, cell cycle, *HES1*, *ID3*

48 **GRAPHICAL ABSTRACT**
HIGHLIGHTS

• Different quiescence stimuli lead to distinct transcriptional and functional

quiescence profiles in endothelial cells

• Flow-mediated endothelial quiescence depth changes from deep to shallow over

time and correlates with p27 levels

• *HES1* and *ID3*, flow-upregulated repressors of p27 expression, are required for

the temporal shift in quiescence depth under flow

ABSTRACT
86

Vascular endothelial cells regulate their cell cycle as blood vessels remodel and mature,

and as they transition from active angiogenesis to quiescence. Mechanical cues

provided by fluid shear stress orchestrate this transition, and laminar blood flow

instigates a quiescent (G_0) state and homeostasis. However, how flow-mediated

quiescence is set up and maintained is poorly understood. We found that flow-mediated

endothelial cell quiescence has unique properties and temporal regulation of

quiescence depth. Flow-exposed endothelial cells had a distinct transcriptome, and

quiescent endothelial cells re-entered the cell cycle more rapidly after extended flow

exposure compared to contact inhibition, indicating a shallow quiescence depth. The

cell cycle inhibitor *CDKN1B* (p27) was required for endothelial cell flow-mediated

quiescence but was not significantly expressed after extended flow exposure. Rather,

flow-exposed endothelial cells first established a deep quiescence that subsequently

became shallow, and p27 levels positively correlated with these distinct quiescent

states. *HES1* and *ID3*, transcriptional repressors of p27 downstream of flow-regulated

Notch and BMP signaling, were required for flow-mediated quiescence depth changes

and the reduced p27 levels associated with shallow quiescence. These findings are

consistent with a model whereby flow-mediated endothelial cell quiescence depth is

temporally regulated downstream of transcriptional regulation of p27.

105
106

107   **INTRODUCTION**
108

Blood vessel network expansion via sprouting angiogenesis during development

is followed by vascular homeostasis; this transition depends on the appropriate

responses of endothelial cells to incoming signals, including laminar shear stress

provided by blood flow. Laminar flow leads to endothelial cell cytoskeletal realignment

and polarization relative to the flow vector and dramatically reduces proliferation [1–6].

Transcriptional profiles reveal numerous changes with flow exposure [7,8], and signaling

pathways such as Notch and BMP are upregulated to promote alignment and vascular

homeostasis [9,10]. Once vascular homeostasis is achieved, endothelial cells rarely divide

[7,11,12]; rather, they are arrested in a quiescent (G_0) state but capable of re-entering the

cell cycle when stimulated by angiogenic growth factors such as VEGF-A [13–15]. How

flow-mediated vascular quiescence is established and maintained is poorly understood,

yet it is critical to blood vessel function.

In non-endothelial cells, quiescence is influenced by cell type and the quiescence

stimulus, and its establishment and maintenance has been extensively studied in

fibroblasts and transformed cells [16–18]. Quiescence stimuli include growth factor

deprivation, contact inhibition, loss of cell adhesion or cell cycle inhibition. Different

quiescence stimuli applied to the same cells lead to transcriptional profiles that are

partially overlapping but have strong unique signatures, suggesting that the quiescence

stimulus qualitatively influences the transcriptome at steady state [16,17]; however, how

these differences affect quiescence properties is poorly understood.

The cell cycle is regulated by cyclins and cyclin-dependent kinases (CDKs) that

promote and CDK inhibitors that block cell cycle progression [19]. In cycling cells, proteins

are post-translationally modified and/or carry destabilization sequences that lead to

rapid gain and loss of function required for passage through the cell cycle [20,21]. In

contrast, quiescence is characterized by transcriptional repression of cell cycle

activators and often by upregulation of cell cycle inhibitors [16,19]. Thus, upon removal of a

quiescence stimulus cell cycle re-entry is delayed, and re-entry time is an indicator of

quiescence depth [22,23]. Quiescence depth is variable; for example, quiescent fibroblasts

have delayed cell cycle re-entry that positively correlates with temporal exposure to the

quiescence stimulus [22,23]. Non-homogeneous quiescence depth is also documented in

muscle stem cells that respond to injury-induced circulating cues by entering a shallow

quiescence called "G_0 alert" that allows for prompt proliferation in response to a

subsequent injury [24]**.** Moreover, spontaneous quiescence has been described recently,

whereby epithelial cells enter quiescence in the absence of obvious environmental or

pharmacological cues, suggesting that quiescence entry has a stochastic component

and/or complex inputs [17,25]. Thus, how quiescence is established, maintained, and

exited is complex.

We examined properties of flow-induced endothelial cell quiescence, and found

that laminar flow-induced endothelial cell quiescence is transcriptionally distinct from

contact-mediated quiescence and functionally exhibits dynamic temporal regulation of

quiescence depth. Prolonged laminar flow leads to an initial stage of deep quiescence

followed by transition to a shallow quiescence depth. Temporal changes in endothelial

cell quiescence depth positively correlate with expression of the cell cycle inhibitor p27.

p27 repression and the shift from deep to shallow quiescence depth requires the

transcription factors *HES1* and *ID3*, targets of the Notch and BMP signaling pathways,

respectively, that are up-regulated by laminar flow and repress p27 levels. These

findings show that endothelial cells respond to laminar flow-mediated quiescence

signals dynamically in a way that correlates with flow-mediated transcriptional changes

affecting p27 expression, and suggest complex regulation of quiescence depth that may

protect from senescence and facilitate the robust angiogenic response required in

physiological settings such as wound healing.

MATERIALS AND METHODS

Cell culture

HUVEC (Lonza #C2519A) were cultured according to the manufacturer's

recommendations in EBM2 (CC-3162, Lonza) supplemented with added growth factors

(Bullet kit, CC-3162, Lonza, referred to as EGM2) at 37°C and 5% CO_2 and used

between passages 2-4, unless otherwise noted. Critical experiments were replicated

with multiple lots of HUVEC.

Microscopy

Imaging was performed using confocal microscopy (Olympus Fluoview FV3000,

IX83) using 405nm, 488nm, 561nm, and 640nm lasers and a UPlanSApo 40x silicone-

immersion objective (NA 1.25), UPlanSApo 60x oil-immersion objective (NA 1.40), or

UPlanSApo 100x oil-immersion objective (NA 1.40). Imaging was performed at RT and

images were acquired with the Fluoview FV31S-SW software. All imaging analysis was

completed in FIJI [26]. For each replicate within an experiment, images were acquired at

the same settings.

Immunofluorescence

Endothelial cells were fixed in 4% PFA (15713 (100504-940), VWR) at 37°C for

10 min and permeabilized in 0.1% Triton (T8787-100ML, Sigma) in PBS for 10 min at

RT. Following permeabilization, HUVEC were blocked for 1 hr at RT in 5% NBCS

(Gibco, #16010-159), antibiotic-antimycotic (Gibco, #15240062), 0.1% sodium azide

(Sigma s2002-100G). Following 3X washes in PBS, cells were incubated in primary

antibody diluted in blocking solution and incubated for 24 hr at 4°C (**Supp. Table S2**,

primary antibody information). Cells were washed 3X with PBS before adding

secondary antibody diluted in blocking solution and incubated for 3 hr at RT in the dark

(**Supp. Table S3**, secondary antibody information). Slides were mounted with coverslips

using Prolong Diamond Antifade mounting media (P36961, Life Technology), sealed

with nail polish, and stored at 4°C. Glass-bottom Ibidi slides or well dishes were stored

in 1X PBS to prevent desiccation.

Endothelial Cell Transfection

HUVEC were grown to subconfluency and treated with siRNAs **(Supp. Table S4,**

siRNA information**)** diluted in OptiMem (Gibco, #11058021) and Lipofectamine 3000

(ThermoFisher, #L3000015) according to manufacturer's protocol

(https://www.thermofisher.com/us/en/home/references/protocols/cell-

[culture/transfection-protocol/lipofectamine-2000.html](https://www.thermofisher.com/us/en/home/references/protocols/cell-culture/transfection-protocol/lipofectamine-2000.html)). Briefly, siRNA at 0.48 µM in Opti-

MEM (31985-070, Gibco) and a 1:20 dilution of Lipofectamine in Opti-MEM were

incubated separately at RT for 5 min, then combined and incubated at RT for 15 min.

HUVEC were transfected at 80% confluency with siRNA at 37°C for 24 hr. After 24 hr,

media was replaced with 10 mL of fresh EGM-2. All experiments were initiated 48 hr

following exposure to siRNA.

Endothelial Cell Flow and Quiescence Depth

Endothelial cells were exposed to laminar flow for 72 hr at 15dyn/cm^2 using an

Ibidi pump system (CC-1090, Ibidi). Static (non-flow) control slides were seeded at

lower densities and incubated for 48 hr prior to fixation with 4% PFA, and subconfluence

status was verified visually prior to fixation and collection. All HUVEC for flow

experiments were cultured in EBM2 supplemented with 2% FBS, 1% antibiotic-

antimycotic (Gibco, #15240062), and 1% Nystatin (Sigma, #N1638-20ML). To prevent

cell shearing, cells were first exposed to 5 d/cm^2 flow for 30 min, 10 d/cm^2 for 30 min,

and then 15 d/cm^2 for the remaining 16 hr (flow establishment (Flow-E)) or 72 hr (flow

maintenance (Flow-M)). For orbital flow experiments, an orbital shaker (Hoefer Red

Rotor Mixer Platform Shaker PR70-115V) was used as previously described [27].

Flow-mediated quiescence depth experiments were performed as follows. After

16 or 72 hr of exposure to laminar flow, cells were pulse labeled for 30 min with EdU

and either fixed (0 hr) or fixed 2, 5, or 8 hr post-flow in 4% PFA for 10 min at RT, then

visualized for EdU and/or stained for antibodies as described (**Supp. Tables S2, S3**).

Contact Inhibition and Quiescence Depth

Endothelial cells were seeded in a 6-well plate at 1.2 million cells/well (high

density) or at 200,000 cells/well (low density) and incubated for 24 hr. A p1000 pipette

tip was used to scrape the cell monolayer and create a gap as previously described [28].

Following the scratch, cells were pulse labeled with EdU for 30 min and fixed (0 hr) or

fixed 2, 5, or 8 hr post-scratching in 4% PFA for 10 min at RT, then visualized for EdU

and/or stained for antibodies as described (**Supp. Table S2, S3**). Cells within 1000 μm

of the scratch edge were imaged **(Supp. Fig. 2a)**.

EdU Labeling

EdU labeling was performed according to the Click-IT EdU 488, 594, or 647

protocol (Invitrogen, C10337). For all experiments with the exception of quiescence

depth experiments, cells were incubated in EdU for 1 hr at 37°C in 5% CO_2 and fixed

with 4% PFA for 10 min at RT. For quiescence depth experiments, cells were incubated

in EdU for 30 min at 37°C in 5% CO_2 and fixed at 4% PFA for 10 min at RT.

Quantification of Positive Endothelial Cells

The DAPI channel was used as a nuclear mask, and nuclear fluorescence

intensity was measured for each cell per image field. Positive cells were defined as

those above the sensitivity threshold. For each image acquired, the percentage of

positive cells was quantified.

Quantification of Cell Axis Ratio

Cell shape and alignment were measured as described previously [7]. Cells were

stained for VE-cadherin upon fixation, and measured at the longest axis of cell divided

by shortest axis to calculate the cell axis ratio, with the longest cell axis being the

direction of flow.

Western Blot Analysis

Western blot analysis was performed according to [29] with modifications. Briefly,

cells were scraped into PBS, centrifuged at 13000 rpm at 4°C for 20 min, then

resuspended in RIPA buffer with protease/phosphatase inhibitor (5872S, Cell

Signaling). The cell pellet was resuspended in sample loading buffer and dithiothreitol

(R0861, Thermo Fisher) and boiled for 10 min. 10μg of sample was run on a 10% SDS-

PAGE gel (161-0183, BioRad) then transferred onto a membrane for 1.5 hr on ice.

Membranes were incubated in primary antibodies overnight at 4°C (**Supp. Table S2**).

Membranes were washed 3X in PBST and stained with secondary antibody diluted in

One-Block at RT for 1 hr (**Supp. Table S3**). Immobilon Forte HRP Substrate

(WBLUF0100, 769 Millipore Sigma) was added for 30 sec. Blots were exposed for 2 sec

ChemiDoc XRS with Chemi High Resolution setting.

RT-qPCR

RNA was extracted using TRIzol (15596018, Invitrogen). cDNA was generated

from 1ug RNA using iScript reverse transcription kit (Bio-Rad, #1708891) and diluted

1:3 in water. qRT–PCR was performed using iTaq Universal SYBR Green SuperMix

(Bio-Rad, #1725121). SYBR Green real-time PCR was performed in triplicate on the

Applied Biosystems QuantStudio 6 Flex Real-Time PCR System (**Supp. Table S5**). For

quantification, relative expression of each gene to β-actin in each sample was

calculated by $2\wedge(CT$ of gene$-CT$ of β-actin). Statistical significance was determined by

unpaired Student's T-test.

Bulk RNA Sequencing and Analysis

RNA was extracted using TRIzol (15596018, Invitrogen). For each condition 3

Ibidi slides were pooled. Stranded libraries were prepared using KAPA mRNA

HyperPrep Kit (7961901001, Roche) and sequenced using NovaSeq S1 at the UNC

Sequencing Core. Data was analyzed as described previously [30]. Briefly, 2–3×10^7 50 bp

paired-end reads per sample were obtained and mapped to the human genome

GRCh38 with STAR using default settings [31]. Quality of sequencing reads was

confirmed with FastQC before mapping. Mapping rate was over 87% for all samples

(**Supp. Table S6**), and gene expression was determined with Htseq-count using the

union mode [32]. Genes with low expression were filtered out (total raw counts in all

samples < 10), differential expression analysis was performed with DESeq2 [33] using

default settings in R, and lists of differentially expressed genes were obtained (p

adjusted <0.1). Gene ontology analysis was performed using enrichGO function in the R

package clusterProfiler. All gene ontology terms shown in this study have a corrected P

value < 0.1.

Quiescence Score

The endothelial quiescence score was calculated using the following formula: QS

$= 1/n\sum_{i=1}^{n} QSUi - 1/m\sum_{j=1}^{m} QSDj$ (QS = Quiescence Score, QSU = QS of upregulated

genes under quiescence (total n genes), QSD = QS of downregulated genes under

quiescence (total m genes)). Selected genes were upregulated or downregulated

relative to control, in this case static (non-flow) samples. All genes were normalized for

sequencing depth and scaled for expression as previously described [8]. To generate the

gene list used for the endothelial quiescence score we utilized two datasets (**Supp.**

Table S1). First, S/G$_2$/M phase associated genes from the CellCycleScoring function of

the R package Seurat were tested in our scRNAseq HUVEC dataset

(https://satijalab.org/seurat/articles/cell_cycle_vignette.html). Genes that were down-

regulated under flow were included in the score. Second, genes that were consistently

upregulated in response to different quiescence stimuli in non-endothelial cells in a

previous report were tested in our scRNA dataset, and if upregulated in response to

flow they were included [16]. The scRNAseq HUVEC dataset was previously published

and is available in NCBI GEO database (accession code GSE151867[8]). The epithelial

quiescence score was developed by the Barr lab as previously described, and utilized

an independent gene list generated using non-transformed human epithelial cell

data[17,34].

Mouse scRNAseq data

scRNAseq data of aged mouse (the *Tabula Muris Senis* dataset) was

downloaded from figshare (https://doi.org/10.6084/m9.figshare.8273102.v2) and

analyzed using Seurat [35]. Endothelial cells were extracted from the original gene count

matrix and down sampled for the 1-, 3- and 18-month time points. Endothelial cells

below the sensitivity level for *Cdkn1b* expression were excluded from analysis.

For mouse ear scRNAseq, a previously generated dataset (Liu et al, in

preparation) using enriched endothelial cells from the ear at P8 (postnatal day 8) or

adult (2-2.5 months) was analyzed. A total of ten vascular endothelial cell populations

were identified by dimension reduction and clustering analysis with Seurat. The

percentage of *Cdkn1b-expressing* cells in the endothelial cells of 9 of the 10

subpopulations (2 samples for each stage) was calculated (one subpopulation had < 10

cells in adult stage and was excluded from analysis).

Statistics

Unpaired Students T-Test was used to determine statistical significance in

situations with two experimental groups, and One-Way ANOVA with Tukey correction

was used for experimental groups of 3 or more. All statistical tests and graphs were

made using the Prism 9.4.1 software (GraphPad Software). *$p \leq 0.05$, ** $p \leq 0.01$, *** p

≤ 0.001, ****$p \leq 0.0001$, ns = not significant*. Statistical methods and significance

parameters are also described in the Figure Legends.

Data availability

Bulk RNA Sequencing data has been deposited in GEO under accession code

GSE213323. Data available upon request.

340   **RESULTS**
341
Laminar flow-exposed endothelial cells have a distinct transcriptional profile

Endothelial cells exposed to laminar shear stress align relative to the flow vector

and significantly reduce their proliferation rate, hallmarks of vascular homeostasis or

quiescence [1,2,5]. To determine whether endothelial cells exposed to laminar flow were

transcriptionally quiescent relative to proliferating cells (static, non-flow), we first applied

a published quiescence score generated in non-endothelial epithelial cells [34] to a

scRNAseq dataset that we previously generated in primary endothelial cells (HUVEC,

human umbilical vein endothelial cells) exposed to laminar flow [8]. Application of the

epithelial quiescence score to this data binned by flow status showed that endothelial

cells exposed to homeostatic laminar flow (here called Flow-Maintenance ($15d/cm^2$, 72

hr)) had a significantly higher score than non-flowed (static condition, 48hr post-seeding

and subconfluent) control endothelial cells **(Fig 1a)**. Previous UMAP analysis of this

dataset revealed 9 groups that primarily separated by flow condition, and the cluster

containing the majority of cells exposed to flow (Flow Cluster 1) had a significantly

higher epithelial quiescence score compared to the equivalent cluster in the static group

(Static Cluster 1), indicating that this published quiescence score positively correlates

with flow-exposed endothelial cells that are predicted to be quiescent. Quiescence

score comparisons among other flow clusters with fewer cells showed heterogeneity

relative to Flow Cluster 1, and a similar comparison of static clusters also revealed

heterogeneity relative to Static Cluster 1, indicating that these distinct clusters exhibit

quiescence heterogeneity. This trend was especially noted for Flow Cluster 2, Static

Cluster 2, and Mix Cluster (contains endothelial cells from both flow and static

conditions) that were previously defined as proliferative populations and had a

significantly reduced quiescence score **(Fig. 1a)**[8].

We next used the same dataset to generate an independent endothelial

quiescence score. Since a hallmark of cellular quiescence is downregulation of

proliferation markers and upregulation of cell cycle inhibitors [16], this score was primarily

generated using expression levels of cell cycle genes that were up- or downregulated in

endothelial cells exposed to extended laminar flow (Flow Maintenance (Flow-M),

15dyn/cm^2/72hr) (**Supp. Table 1**). The endothelial quiescence score was then applied

to the dataset to verify that the score predicted the relationships. Consistent with the

epithelial score, flow cluster 1 had a significantly higher quiescence score compared to

static cluster 1, and the clusters associated with proliferation had significantly lower

scores (**Fig. 1b**). These findings indicate that endothelial cells exposed to homeostatic

laminar flow (Flow-M) are in a transcriptionally quiescent cell cycle state.

We next applied the endothelial quiescence score to an independently generated

dataset to validate the score and further examine endothelial cell quiescence

relationships. Both fibroblasts and epithelial cells exposed to different quiescence

stimuli have unique transcriptional profiles [16,17], so we asked whether the mode of

quiescence induction affected the quiescence score of endothelial cells exposed to

static vs. extended laminar flow (Flow-M) or low vs. high density static (non-flow)

conditions. Application of the endothelial quiescence score to HUVEC bulk RNA seq

data showed significant increases under flow and high density compared to static and

low density conditions, respectively **(Fig. 1c)**, and these relationships held when the

epithelial quiescence score was applied in the same manner **(Supp. Fig. 1a)**. Venn

diagrams revealed that extended laminar flow and contact inhibition have both shared

and distinct transcriptional changes in endothelial cells, with approximately one quarter

of the regulated genes shared for up- and down-regulated categories (upregulated:

21.7% (494/2272) flow and 24.3% (494/2030) density shared genes; down-regulated

25.1% (573/2282) flow and 23.7% (573/2415) density shared genes) (**Supp. Fig. 1b**).

Further analysis of highly differentially expressed genes revealed only one overlapping

gene (*CCDC190*) that was downregulated between flow and high density conditions,

and no genes were highly upregulated with both flow and high density conditions,

suggesting that while there is some overall overlap, the quiescence profiles of

endothelial cells exposed to extended laminar flow or high density are largely distinct

(Supp. Fig. 1c-f). Gene Ontogeny (GO) analysis revealed several terms linked to

migration and development shared by endothelial cells under extended laminar flow and

high density (**Supp. Fig. 1g-j**). Although endothelial cell quiescence is sometimes

characterized as an inactive state, migration of quiescent cells has been documented[36]

and contributes to developmental and pathological angiogenesis [37–39]. However, more

prominent were numerous categories unique to either flow exposed or high density

endothelial cells; for example, flow exposed cells had adhesion terms upregulated and

protein ubiquitination terms downregulated, while contact inhibited endothelial cells had

TGF-β signaling upregulated and mitochondrial gene expression downregulated (**Supp.**

Fig. 1g-j), consistent with the largely unique transcriptional profiles for different modes

of endothelial cell quiescence.

We next examined how expression of specific cell cycle regulatory genes varied

under extended flow exposure or high density, and found that expression of cell cycle

activators was down-regulated regardless of quiescence induction stimulus, while

transcription of cell cycle inhibitors showed an expected increased trend under high

density conditions but not under flow exposure **(Fig. 1d-e).** Thus, endothelial cells

respond to extended laminar flow exposure with a distinct quiescence transcriptional

profile compared to high density-induced quiescence that includes differences in how

cell cycle inhibitors are regulated, suggesting unique properties of flow-mediated

endothelial cell quiescence.

Shallow quiescence depth characterizes endothelial cells exposed to extended

laminar flow

Quiescence depth varies in fibroblasts and stem cells, as defined by the time to

re-enter the cell cycle upon removal of the quiescence stimulus [22–24]. The lack of cell

cycle inhibitor RNA upregulation in endothelial cells exposed to laminar flow led us to

hypothesize that laminar flow-exposed cells are in a relatively shallow quiescence state.

To measure endothelial cell quiescence depth, we assessed cell cycle re-entry time via

EdU pulse labeling and pRB (phospho-retinoblastoma protein) expression at defined

times after removal of the quiescence stimulus (**Supp. Fig. 2a**) [40,41]. The timepoints for

analysis were selected based on the doubling time of HUVEC (Lonza Manufacturer).

EdU labels cells in S-phase, and phosphorylation releases RB from E2F transcription

factors, allowing for progression through the G_1-S checkpoint and proliferation [41,42].

Endothelial cells exposed to either extended laminar flow (Flow-M) or contact inhibition

had significantly reduced percentages of EdU-labeled **(Fig. 2a-b, d-e)** and pRB positive

cells **(Fig. 2a, c, d, f)** at the end of the flow or contact inhibition period, consistent with

previously published work [7,12,43], as well as decreased pRB nuclear intensity **(Supp.**

Fig. 2b-c). Endothelial cells released from extended laminar (Flow-M) significantly

increased EdU-labeled and pRB positive cells within 2 hr of flow cessation, while

endothelial cells released from contact inhibition did not show significant increases in

EdU-labeling and pRB reactivity until 8 hr post-release **(Fig. 2a-f, Supp. Fig. 2b-c).**

These findings show that endothelial cells exposed to homeostatic laminar flow (Flow-

M) are less deeply quiescent compared to endothelial cells that are contact inhibited.

Cell cycle inhibitor p27 expression correlates with quiescence depth and is

required for flow-mediated quiescence

Although several cell cycle inhibitors participate in cell cycle progression, p27

(*CDKN1B*) is often upregulated with quiescence in non-endothelial cells [44–46]. However,

after exposure to extended laminar flow (Flow-M), RT-qPCR analysis revealed a

significant decrease in *CDKN1B* RNA expression, and protein analysis showed a highly

significant decrease in the percentage of p27+ endothelial cells relative to static controls

(static 47% vs. flow 2%) **(Fig. 3a-c; Supp. Fig 3a,c).** In contrast, p27 protein levels

increase in contact inhibited endothelial cells [43], and we confirmed that both RNA levels

and the percentage of p27+ endothelial cells significantly increased with contact

inhibition compared to low density controls (14% low vs. 99% high density) **(Fig. 3d-f;**

Supp. Fig. 3b-c). It is striking that endothelial cells exposed to extended laminar flow

(Flow-M) and with shallow quiescence depth have 2% p27+ cells, while contact inhibited

endothelial cells with more profound quiescence depth have 99% p27+ cells. Taken

together, these findings show that p27 expression positively correlates with quiescence

depth in endothelial cells.

459
460
We next asked whether p27 is functionally required for flow-mediated quiescence

induction in endothelial cells by depleting p27 via siRNA knockdown (KD). p27-depleted

cells exposed to extended flow (Flow-M) had a significantly increased percentage of

cells positive for EdU-labeling and staining for Ki67, a marker of cells in S, G_2, or M

phase [47], compared to non-targeting (NT) siRNA controls (**Fig. 4a-b; Supp. Fig. 3d-f**),

indicating that, despite relatively low expression at the end of flow exposure, p27 is

required for flow-mediated endothelial cell quiescence.

To better understand how p27 affects endothelial cell quiescence, we

transcriptionally profiled quiescent endothelial cells depleted for p27 under extended

laminar flow and high density conditions. Consistent with differences in p27 expression

with different quiescence stimuli, there was little overlap in the profiles of genes up- or

down-regulated in each condition upon p27 depletion (upregulated: 9.0% (12/133) flow

and 8.6% (12/139) density shared genes; down-regulated 13.4% (18/134) flow and

0.4% (18/313) density shared genes) **(Supp. Fig. 4a)**. Further analysis of highly

differentially expressed genes revealed no overlapping genes between control and p27

depleted cells under extended laminar flow or contact inhibition, and comparison of

gene ontology (GO) terms of p27 depleted cells to controls under extended laminar flow

or contact inhibition revealed no shared terms **(Supp. Fig. 4b-i)**, suggesting that p27

affects endothelial cell transcriptional quiescence programs differently depending on the

quiescence stimulus.

Flow-mediated quiescence depth is temporally regulated in endothelial cells

p27 is functionally required to mediate flow-mediated endothelial cell repression

of proliferation despite having significantly reduced expression by 72 hr of laminar flow

exposure (Flow-M). We hypothesized that p27 expression levels fluctuated during the

flow period, and temporal analysis showed that the percentage of p27+ cells

significantly increased 16 hr after flow initiation compared to the 0 hr levels (time of flow

cessation), consistent with another study showing elevated p27 levels after 16 hr of flow

[48], and percent p27+ cells then decreased during the remaining time and were almost

undetectable at 72 hr **(Fig. 4c-d, Supp. Fig. 5a).** Levels of p21, another member of the

CIP-KIP cell cycle inhibitor family, showed a similar decrease in levels even earlier in

the time course **(Fig. 4c, e, Supp. Fig. 5b)**, and Ki67+ cells also decreased to

undetectable levels over flow time **(Fig. 4c,f; Supp. Fig 5c)**, consistent with the

conclusion that endothelial cells leave the cell cycle and become quiescent by 16-24hr

of laminar flow, then subsequently down-regulate p27 levels while maintaining

quiescence.

We then asked whether the quiescence depth of endothelial cells under laminar

flow was temporally regulated, by assessing time to cell cycle re-entry upon flow

cessation after 16 hr of flow (flow establishment (Flow-E)), a timepoint showing elevated

levels of p27. After this relatively short exposure to laminar flow, endothelial cells did not

exhibit significant EdU-labeling and pRB reactivity until 8 hr post-flow, similar to

endothelial cells released from contact inhibition and longer than the 2 hr time to cell

cycle re-entry exhibited by endothelial cells exposed to extended laminar flow (Flow-M)

(Fig. 4g-i; Supp. Fig. 5d). These findings correlate with the temporal regulation of p27

levels over the flow period, and they suggest that endothelial cells initially respond to

laminar flow by p27 upregulation and establishment of a relatively "deep" quiescence

(Flow-E) that over time becomes shallower, along with reduced levels of cell cycle

inhibitors p27 and p21. We further characterized the flow response by assessment of

endothelial cell alignment, which was not established at 16 hr of flow (Flow-E) but was

established by 72 hr of flow (Flow-M) in our model (**Supp. Fig. 5e-f).** Thus, endothelial

cells under laminar flow transition from a deep to a shallower quiescence over time,

coincident with flow-mediated cell alignment and reduced levels of p27.

Transcriptional targets of flow-mediated endothelial cell signaling regulate p27

levels under laminar flow

518
To better understand how endothelial cell p27 expression decreases over time

with laminar flow, we examined genes whose expression is upregulated by laminar flow

and whose encoded proteins transcriptionally repress p27. It is well-established that

signaling pathways such as Notch and BMP are upregulated with laminar flow [2,9,10], and

this in turn leads to upregulation of targets of these pathways. *ID3* and *HES1* are

downstream targets of the Notch and BMP pathways, respectively, and both

transcriptionally repress *CDKN1B* expression via direct promoter interactions [49,50]. Upon

exposure to extended laminar flow (Flow-M), we observed an accumulation of *HES1*

and *ID3* RNA via RT-qPCR compared to static conditions (**Supp. Fig. 6a-b**). We directly

assessed the effects of these transcription factors on p27 levels in endothelial cells via

siRNA KD and found that depletion of either *HES1* or *ID3* RNA prevented the normal

decrease in the percent of p27+ cells under extended laminar flow (Flow-M) **(Fig. 5a-b;**

Supp. Fig. 6c), indicating that these repressors are required for flow-mediated

reduction of p27 levels over time.

We hypothesized that *HES1* and *ID3* are required for the temporal changes to

endothelial cell quiescence depth over time, and we predicted that *HES1* or *ID3*

depletion would prevent the transition from deep to shallower quiescence depth with

time. We performed cell cycle re-entry experiments upon cessation of extended laminar

flow (Flow-M) and found that either *HES1* or *ID3* depletion significantly extended the

time required to see significant increase in percent EdU+ and pRB+ endothelial cells

compared to controls (**Fig. 5c-h, Supp. Fig. 6d-e),** suggesting that the normal change

to a shallow quiescence depth over time was prevented when these repressors were

depleted. These findings are consistent with the idea that endothelial quiescence depth

is temporally regulated from deep to shallow over time in conjunction with laminar flow-

mediated changes, and that p27 regulates this transition downstream of the

transcriptional repressors *HES1* and *ID3*.

546
547
548   **Endothelial cell p27 expression decreases with age *in vivo***
549

To begin to determine whether the transition from deep to shallow quiescence

documented here was mirrored *in vivo*, we used endothelial cell p27 expression as a

surrogate for quiescence depth and re-analyzed a published scRNAseq dataset

collected from mice of different ages[35]. We predicted higher levels *of Cdkn1b* in

endothelial cells of young mice with expanding vessel networks that are not yet

established, and we found that *Cdkn1b* expression was significantly higher in

endothelial cells from 1-month old mice compared to endothelial cells of older mice (3 or

18 months) **(Fig. 6a)**. Next, we analyzed temporal scRNAseq data we generated and

found that percent *Cdkn1b*+ endothelial cells in ear skin trended lower (p=0.16) with age

from neonatal (postnatal day 8) to adult (2-2.5 months) mice **(Fig. 6b)**. These findings

are consistent with the idea that endothelial cell quiescence depth changes with

developmental time, and that establishment initiates a deep quiescence that becomes

shallower over time as endothelial cells are exposed to extended laminar flow that

upregulates repressors of p27 expression (**Fig. 6c**).

DISCUSSION

Endothelial cell quiescence is important for vascular homeostasis and proper

barrier function, but how this quiescence is set up and maintained is not well-

understood. Here we show that quiescence depth is shallow in endothelial cells

exposed to extended laminar flow under maintenance conditions compared to contact-

inhibited endothelial cell quiescence depth in the absence of flow, and that flow-

mediated quiescence depth moves from a deeper to a shallower state as the

establishment phase ends and the maintenance phase commences. This discovery is

unexpected because we originally presumed that cells exit the cell cycle to a shallow

quiescent state that becomes deeper over time rather than the opposite. This surprising

finding correlates with changes in expression of the cell cycle inhibitor p27 and requires

transcriptional repressors normally upregulated by laminar flow, and it suggests a new

critical control point for vascular homeostasis that is linked to aging [51–53].

To better understand how endothelial cell quiescence varies with stimulus and

conditions, we developed an endothelial quiescence score using a cohort of genes

primarily chosen because the encoded proteins are flow-regulated and involved in the

cell cycle [8]. Application of this score to bulk RNA seq data not used to generate the

score validated that it was higher in endothelial cells predicted to be quiescent.

Moreover, application of a unique quiescence score developed by the Barr lab for

epithelial cells found similar relationships [54], suggesting that some aspects of vascular

quiescence are shared among cell types and conditions. Both quiescence scores were

higher in endothelial cells made quiescent via extended laminar flow or contact

inhibition compared to controls, and further transcriptome analysis revealed that both

conditions were characterized by down-regulation of cell cycle activators while only

contact inhibited endothelial cells showed increased expression of cell cycle inhibitor

genes. Further analysis of bulk RNAseq data indicated that although exposure to

extended flow or contact inhibition led to about a quarter of differentially regulated

genes being shared, most were unique to the quiescence stimulus, and comparison of

the most highly regulated genes and GO categories supported that the transcriptional

profiles are largely unique to flow or contact inhibition.

The global differences in cell cycle inhibitor expression in extended flow- vs.

contact inhibition-mediated quiescence correlated with expression of the cell cycle

inhibitor p27, and quiescence depth experiments revealed that endothelial cells under

extended laminar flow reentered the cell cycle soon after flow cessation while released

contact inhibited cells took significantly longer, indicating a deeper quiescence state.

This was surprising, as intuitively vascular homeostasis associated with extended

laminar flow was thought to be accompanied by relatively deep quiescence, and

endothelial cells exposed to laminar flow are in a high density monolayer. However,

these findings positively correlated with p27 levels, and reanalysis of several RNA seq

datasets showed a trend of reduced endothelial cell p27 levels with age. This is in line

with a recent study that found high levels of interstitial flow associated with shallower

quiescence in fibroblasts [55]. In another study, quiescence depth was invoked to explain

in vivo findings, as muscle stem cells entered a shallow quiescence state in response to

injury, and the shallow quiescence was proposed to be advantageous in wound

resolution [24].

Reduced p27 expression in endothelial cells exposed to extended flow was

nevertheless accompanied by a functional requirement for p27 to establish quiescence,

suggesting different requirements for the establishment vs. maintenance of vascular

quiescence. Direct analysis showed that quiescence depth changed with exposure time

to laminar flow, starting with a deep quiescence during the establishment phase (Flow-

E) when endothelial cells were not yet aligned to the flow vector, and becoming shallow

over extended exposure (Flow-M) and cell alignment to flow, consistent with our model

invoking distinct requirements for establishment vs. maintenance of laminar flow-

mediated endothelial quiescence. This idea is also consistent with our finding that p27

levels vary temporally during flow-mediated quiescence, increasing over the first 16-24

hours of laminar flow exposure and decreasing to very low levels with further flow.

Taken together, these data suggest that flow-mediated quiescence exhibits different

requirements for establishment vs. maintenance, and that the functional requirement for

p27 is related to establishment but not maintenance of flow-mediated quiescence. This

phenomenon is unique to flow, as both we and others have found that under high

density contact-inhibited endothelial cells accumulate p27 [43].

p27 expression is regulated by transcriptional repressors that are in turn

positively regulated by laminar flow, suggesting how endothelial cell p27 levels are

dynamically regulated in response to laminar flow **(Fig 6c)**. Both Notch and BMP

signaling are flow-regulated [9,10], and *HES1* and *ID3* are known transcriptional targets of

Notch and BMP, respectively, that are upregulated with laminar flow [56–58]. We found that

both repressors are functionally required non-redundantly to dampen p27 expression as

endothelial cells move from the establishment phase to the maintenance phase of

quiescence and vascular homeostasis. Endothelial cell quiescence depth is more

profound in the absence of either repressor, indicating that increased Notch and BMP

signaling during quiescence establishment, and subsequent upregulation of these

repressors leads to reduced p27 expression levels and a change from a deep to shallow

endothelial quiescence depth. This model is consistent with a defined role for p27-

mediated cell cycle regulation during angiogenesis to promote arterio-venous

differentiation important for early stages of angiogenesis and remodeling [48,59]. It is also

consistent with our analysis of RNA seq data showing decreased endothelial cell p27

levels with age, since age is predicted to result in more endothelial cells in the

maintenance phase of vascular quiescence.

Our work shows that endothelial cells have different responses to quiescence-

inducing stimuli. Quiescence depth changes over time from deep to shallow with

laminar flow, and this temporal regulation requires upregulation of p27 repressors and

reduced levels of p27 protein. Why might there be an advantage to endothelial cells

supporting a shallower quiescence state under flow maintenance (homeostasis)

conditions? One possibility is protection from premature senescence. Senescent cells

are permanently arrested [60], and fibroblasts that remain in quiescence for extended

periods of time eventually undergo senescence [61]. Senescent endothelial cells often

express a senescence-associated secretory phenotype (SASP) [62] that includes

upregulation of inflammatory cytokines that contribute to endothelial dysfunction,

leading to cardiovascular disease [63–65]. Thus, endothelial cells in a shallow quiescence

state at steady state may be protected from inappropriate senescence. *HES1*

expression in fibroblasts prevents premature senescence [66], supporting the concept that

shallow quiescence may protect from senescence.

A shallower endothelial cell quiescence depth may also help promote

physiological angiogenesis. A seminal study showed that endothelial cells *in vivo* are

non-proliferative but can respond to angiogenic/growth signals and re-enter the cell

cycle relatively quickly [67]. It was posited that these properties set up vascular

homeostasis while allowing for endothelial cells to rapidly respond to physiological or

pathological signals for angiogenesis. More recent work provides evidence that some

endothelial cells in the dorsal aorta respond to clamp injury with clonal expansion to

cover the denuded area in a relatively short time period [68]. Thus, the need to

simultaneously maintain homeostasis yet be primed for prompt injury response may be

facilitated by a relatively shallow endothelial cell quiescence depth. Conversely, shallow

quiescence depth may also contribute to pathological neo-angiogenesis associated with

diseases such as cancer.

ACKNOWLEDGMENTS

We thank members of the Bautch and Cook labs for their support and feedback, and the

UNC High Throughput Sequencing Core for technical help with bulk RNAseq sample

preparation and sequencing. We thank BioRender for figure model generation.

AUTHOR CONTRIBUTIONS

Natalie T Tanke (NTT), Jeanette Gowen Cook (JGC), Victoria L Bautch (VLB)

conceptualized the work; NTT, Ziqing Liu (ZL), Morgan Oatley (MO), Tianji Yu (TY),

Pauline Bougaran (PB), and Kaitlyn Quigley (KQ) performed and analyzed experiments.

NTT and VLB wrote and edited the manuscript; VLB and JGC provided study

supervision and oversight.

FUNDING

This work was supported by grants from the National Institutes of Health (R35

HL139950 to VLB), the MiBioX T32 Training Grant (T32 GM 119999-03, NTT), a Ruth

L. Kirschstein Predoctoral Fellowship (1F31HL156527-01 to NTT), and an American

Heart Association Postdoctoral Fellowship (AHA829371 to MO).

DECLARATION OF INTERESTS: none

FIGURE LEGENDS

Figure 1. Laminar flow-mediated quiescence is transcriptionally distinct.

(a) Quantification of epithelial quiescence score in HUVEC scRNA dataset by cluster.

Clusters were previously defined[8]. Flow-M, flow maintenance (laminar flow

($15d/cm^2/72hr$)). Static cells were visually checked for subconfluence and fixed 48 hr

post-seeding. **(b)** Quantification of endothelial quiescence score in HUVEC scRNA

dataset by cluster. **(c)** Endothelial quiescence score on bulk RNA seq data of HUVEC

exposed to indicated stimuli, n= 3 replicates. **(d-e)** Heatmaps showing relative

expression of cell cycle proliferation markers (green font) and inhibitors (red font) plotted

using bulk RNAseq data of HUVEC under indicated conditions, n=3 replicates.

Statistics, one-way ANOVA with Tukey's multiple comparison test; *** $p \leq 0.01$, **** $p \leq$

*0.001, *****$p \leq 0.0001$, ns = not significant*.

Figure 2. Endothelial cell flow maintenance quiescence stimulus leads to shallow

quiescence depth.

(a) Representative images of HUVEC under static (non-flow) or Flow-M (flow

maintenance) conditions with EdU incorporation and fixation at indicated times post

Flow-M release. Cultures stained for DAPI (white, nuclear mask) and EdU (red, s-

phase) or pRB (blue, interphase). Scale bar, 50 μm. White arrow, flow direction. **(b)**

Quantification of percent EdU+ cells with indicated conditions. n=3 replicates, 5 images

per condition per replicate. **(c)** Quantification of percent pRB+ cells with indicated

conditions. n=3 replicates, 5 images per condition per replicate. **(d)** Representative

images of HUVEC under low or high density conditions with EdU incorporation and

fixation at indicated times post high-density release. Cultures stained for DAPI (white,

nuclear mask) and EdU (red, s-phase) or pRB (blue, interphase). Scale bar, 50 μm. **(e)**

Quantification of percent EdU+ cells with indicated conditions. n=3 replicates, 6 images

per condition per replicate. **(f)** Quantification of percent pRB+ cells with indicated

conditions. n=3 replicates, 6 images per condition per replicate. Statistics, one-way

ANOVA with Tukey's multiple comparisons test; *$p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$,

****$p \leq 0.0001$, ns = not significant.

731
732
Figure 3. Cell cycle inhibitor p27 expression levels vary with endothelial

quiescence stimulus.

(a) RT-qPCR for *CDKN1B* levels in indicated conditions. n=3 replicates. **(b)**

Representative images of HUVEC under indicated conditions stained for p27 (green)

and DAPI (white, nuclear mask). Scale bar, 50 μm. White arrow, flow direction. **(c)**

Quantification of HUVEC p27+ cells under indicated conditions, n=3 replicates, 5

images per condition per replicate. **(d)** RT-qPCR for *CDKN1B* levels in indicated

conditions. n=3 replicates. **(e)** Representative images of HUVEC under indicated

conditions stained for p27 (green) and DAPI (white, nuclear mask). Scale bar, 50 μm. **(f)**

Quantification of HUVEC p27+ cells under indicated conditions. n=3 replicates, 5

images per condition per replicate. Statistics, student's two-tailed *t*-test; ** $p \leq 0.01$,

****$p \leq 0.0001$.

Figure 4. Endothelial quiescence depth varies with laminar flow exposure time

and positively correlates with p27 expression levels

(a) Representative images of HUVEC with indicated siRNA treatments and conditions.

after EdU incorporation (red, S-phase) and staining with DAPI (white, nuclear mask).

Scale bar, 50 µm. White arrow, flow direction. **(b)** Quantification of EdU+ cells with

indicated conditions. n=3 replicates, 5 images per condition per replicate. **(c)**

Representative images of HUVEC with indicated conditions and stained for p27

(inhibitor), p21 (inhibitor), Ki67 (proliferation marker), and DAPI (nucleus) in white. Scale

bar, 50 µm. White arrow, flow direction. **(d-f)** Quantification of percent p27+ (d), p21+

cells (e), or Ki67+ cells (f). n=3 replicates, 5 images per condition per replicate, **(g)**

Representative images of HUVEC under indicted conditions and after EdU incorporation

(red, S-phase) and stained for pRB (blue, interphase) and with DAPI (white, nuclear

mask), Flow-E, flow establishment; Flow-M, flow maintenance. Scale bar, 50 µm. White

arrow, flow direction. **(h)** Quantification of EdU+ cells in indicated conditions. n=3

replicates, 5 images per condition per replicate. **(i)** Quantification of pRB+ cells in

indicated conditions. n=3 replicates, 5 images per condition per replicate. Statistics,

one-way ANOVA with Tukey's multiple comparisons test; *$p \leq 0.05$, ****$p \leq 0.0001$, ns =*

not significant.

Figure 5. BMP and NOTCH regulated p27 repressors *HES1* and *ID3* regulate p27

levels and flow-mediated quiescence depth

(a) Representative images of HUVEC under indicated conditions and with indicated

siRNA treatment. Cells stained for p27 (green) and DAPI (white, nuclear mask). Scale

bar, 50 µm. White arrow, flow direction. **(b)** Quantification of p27+ cells under indicated

conditions and treatments. n=3 replicates, 5 images per condition per replicate. **(c)**

Representative images of HUVEC with indicated conditions and siRNA treatments.

Cells were pulse labeled with EdU (red, S-phase) and stained for pRB (blue, interphase)

and DAPI (white, nuclear mask). Scale bar, 50 µm. White arrow, flow direction. **(d)**

Representative images of HUVEC with indicated conditions and siRNA treatments.

Cells were pulse labeled with EdU (red, S-phase) and stained for pRB (blue, interphase)

and DAPI (white, nuclear mask). Scale bar, 50 µm. White arrow, flow direction. **(e)**

Quantification of EdU+ cells with indicated conditions. n=3 replicates, 5 images per

condition per replicate. **(f)** Quantification of pRB+ cells with indicated conditions. n=3

replicates, 5 images per condition per replicate. **(g)** Quantification of EdU+ cells with

indicated conditions. n=3 replicates, 5 images per condition per replicate. **(h)**

Quantification of pRB+ positive cells with indicated conditions. n=3 replicates, 5 images

per condition per replicate. Statistics, one-way ANOVA with Tukey's multiple

comparisons test. ** *p ≤ 0.01, *** p ≤ 0.001, ****p ≤ 0.0001, ns = not significant.*

786
787

Figure 6. *Cdkn1b* expression does not accumulate with age *in vivo*

(a) scRNAseq analysis of normalized *Cdkn1b* expression in endothelial cells isolated

from mice of indicated ages. See Methods and text for details of dataset reanalysis.

Endothelial cells with *Cdkn1b* expression at 0 were excluded from analysis. *****p ≤*

0.0001, ns = not significant by one-way ANOVA with Tukey's multiple comparisons test.

(b) scRNAseq analysis of percent *Cdkn1b*+ endothelial cells isolated from neonatal (P8)

or adult (2-2.5m) murine ears. Each dot represents one endothelial subpopulation in

one sample. Statistics, student's two-tailed *t*-test **(c)** Proposed model for dynamic

regulation of endothelial cell flow-mediated quiescence depth with flow exposure time.

In the first 16-24 hr (Flow-E, flow establishment), a deep quiescence is established

accompanied by high levels of cell cycle inhibitor p27. With time, transcription factors

that are targets of flow-mediated Notch and BMP signaling are upregulated, and these

p27 transcriptional repressors prevent ongoing p27 synthesis and lead to significantly

reduced p27 levels. The regulators are required to transition to a shallow quiescence

depth that characterizes extended exposure to laminar flow (Flow-M, flow

maintenance).

804
805

806   **SUPPLEMENTARY FIGURE LEGENDS**
807
Supplemental Figure 1. Endothelial bulk RNAseq-derived quiescence

transcriptional profiles are distinct with different stimuli.

(a) Quantification of an epithelial quiescence score [54] of HUVEC bulk RNAseq dataset

from different conditions. Statistics, *$p \leq 0.05$ using One-Way ANOVA with Tukey's

multiple comparison test, n=3 replicates. Flow-M, flow maintenance. **(b)** Venn diagrams

showing bulk RNA seq analysis of genes up- and down-regulated in HUVEC under

indicated conditions. **(c-f)** Heatmaps showing relative expression levels of genes (top 50

by fold change and p-value) in bulk RNA seq from HUVEC under indicated conditions.

(g-j) GO term analysis from bulk RNA-seq data with indicated comparisons.

Representative biological processes GO terms significantly enriched (P adjusted < 0.1)

in differentially regulated genes are shown.

Supplemental Figure 2. pRB nuclear fluorescence intensity under indicated

conditions.

(a) Schematic showing areas of scratch wound used for imaging post-scratch for

contact inhibition quiescence depth experiments. **(b)** Quantification of pRB nuclear

fluorescence intensity under indicated conditions. n=3 replicates, 5 images averaged

per condition per replicate. **(c)** Quantification of nuclear fluorescence intensity of pRB in

low vs. high density release timepoints. n=3 replicates, n=5 averaged images per

condition per replicate. Statistics, one-way ANOVA with Tukey's multiple comparisons

test *$p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, ****$p \leq 0.0001$, ns= not significant.

Supplemental Figure 3. Cell cycle inhibitor p27 expression differs with

quiescence stimulus.

(a) Quantification of p27 nuclear fluorescence intensity in indicated conditions. n=3

replicates, 5 images averaged per condition per replicate. **(b)** Quantification of p27

nuclear fluorescence intensity in indicated conditions. n=3 replicates, 5 images

averaged per condition per replicate. **(c)** Western blot of p27 expression under indicated

conditions, and with indicated antibodies. p27 Ab #1 (Cell Signaling) and p27 Ab #2

(Santa Cruz). **(d)** Representative images of HUVEC with indicated siRNA and

treatments. Endothelial cells stained with Ki67 (yellow, proliferation marker) and DAPI

(gray, nucleus mask). Scale bar 50 µm. White arrow, flow direction. **(e)** Quantification of

percent Ki67+ cells. n=3 replicates, n= 5 images per condition per replicate. **(f)**

Quantification of nuclear fluorescence intensity of Ki67. n=3 replicates, n=5 averaged

images per condition per replicate. Statistics, one-way ANOVA with Tukey's multiple

comparisons test; *** $p \leq 0.01$, ****$p \leq 0.0001$, ns= not significant.*

Supplemental Figure 4. p27 depletion leads to distinct transcriptional changes

dependent on quiescence stimulus.

(a) Venn diagrams showing overlap of HUVEC genes differentially regulated in

indicated conditions and for *CDKN1B* KD compared to NT. **(b-e)** Heatmaps showing

relative expression levels of genes differentially regulated (top 50 by fold change and p-

value) in response to indicated conditions. **(f-i)** GO analysis performed on differentially

expressed genes from bulk RNA-seq data under comparing indicated

conditions/treatments. Representative biological processes GO terms significantly

enriched (P adjusted < 0.1) in differentially regulated genes are shown.

855
856
Supplemental Figure 5. Cell cycle inhibitor and Ki67 expression intensity varies

with flow exposure and correlates with flow-mediated endothelial cell alignment.

(a) Quantification of p27 nuclear fluorescence intensity with indicated conditions. n=3

replicates, 5 averaged images per condition per replicate. **(b)** Quantification of p21

nuclear fluorescence intensity with indicated conditions. n=3 replicates, 5 averaged

images per condition per replicate. **(c)** Quantification of Ki67 nuclear fluorescence

intensity with indicated conditions. n=3 replicates, 5 averaged images per condition per

replicate. **(d)** Quantification of pRB nuclear fluorescence intensity under indicated

conditions. n=3 replicates, 5 averaged images per condition per replicate. **(e)**

Representative images of HUVEC stained with VE-cadherin (white, junction marker)

and DAPI (blue, nucleus) under indicated conditions. Scale bar, 50 μm. White arrow,

flow direction. **(f)** Cell axis ratio quantification in indicated conditions. n=3 replicates, 5

images per condition per replicate. Statistics, one-way ANOVA with Tukey's multiple

comparisons test; *$p \leq 0.05$, **$p \leq 0.01$, *** $p \leq 0.001$, ****$p \leq 0.0001$, ns = not*

significant.

Supplemental Figure 6. Nuclear fluorescence intensity of markers and changes

with *HES1* or *ID3* depletion.

(a) RT-qPCR for *HES1* expression under indicated conditions. n=3 replicates. Statistics,

*** *$p \leq 0.001$* with student's two-tailed *t*-test. **(b)** RT-qPCR for *ID3* expression under

indicated conditions, n=3 replicates. Statistics, * $p \leq 0.05$ with student's two-tailed *t*-test.

(c) Quantification of p27 nuclear fluorescence intensity with indicated siRNA treatments

and conditions. n=3 replicates, 5 averaged images per condition per replicate. Statistics,

**$p \leq 0.01$, ****$p \leq 0.0001$ by one-way ANOVA with Tukey's multiple comparisons test.

(d) Quantification of pRB nuclear fluorescence intensity with indicated siRNA treatments

and conditions. n=3 replicates, 5 averaged images per condition per replicate. Statistics,

****$p \leq 0.0001$, *ns= not significant* by one-way ANOVA with Tukey's multiple comparisons

test. **(e)** Quantification of pRB nuclear fluorescence intensity with indicated siRNA

treatments and conditions. n=3 replicates, 5 averaged images per condition per

replicate. Statistics, **$p \leq 0.01$, ****$p \leq 0.0001$, *ns= not significant* by one-way ANOVA

with Tukey's multiple comparisons test.

REFERENCES

1. Privratsky JR, Newman PJ. PECAM-1: regulator of endothelial junctional integrity.
Cell Tissue Res. 2014;355(3):607-619. doi:10.1007/s00441-013-1779-3

2. Baeyens N, Schwartz MA. Biomechanics of vascular mechanosensation and
remodeling. *Mol Biol Cell*. 2016;27(1):7-11. doi:10.1091/mbc.E14-11-1522

3. Sluysmans S, Vasileva E, Spadaro D, Shah J, Rouaud F, Citi S. The role of apical
cell-cell junctions and associated cytoskeleton in mechanotransduction. *Biol Cell*.
2017;109(4):139-161. doi:10.1111/boc.201600075

4. Levesque MJ, Sprague EA, Schwartz CJ, Nerem RM. The Influence of Shear
Stress on Cultured Vascular Endothelial Cells: The Stress Response of an
Anchorage-Dependent Mammalian Cell. *Biotechnol Prog*. 1989;5(1):1-8.
doi:10.1002/btpr.5420050105

5. Levesque MJ, Nerem RM, Sprague EA. Vascular endothelial cell proliferation in
culture and the influence of flow. *Biomaterials*. 1990;11(9):702-707.
doi:10.1016/0142-9612(90)90031-k

6. Ziegler T, Nerem RM. Effect of flow on the process of endothelial cell division.
Arterioscler Thromb J Vasc Biol. 1994;14(4):636-643. doi:10.1161/01.atv.14.4.636

7. Ruter DL, Liu Z, Ngo KM, X S, Marvin A, Buglak DB, Kidder EJ, Bautch VL.
SMAD6 transduces endothelial cell flow responses required for blood vessel
homeostasis. *Angiogenesis*. 2021;24(2):387-398. doi:10.1007/s10456-021-09777-7

8. Liu Z, Ruter DL, Quigley K, Tanke NT, Jiang Y, Bautch VL. Single-Cell RNA
Sequencing Reveals Endothelial Cell Transcriptome Heterogeneity Under
Homeostatic Laminar Flow. *Arterioscler Thromb Vasc Biol*. 2021;41(10):2575-
2584. doi:10.1161/ATVBAHA.121.316797

9. Yang J, Li X, Li Y, Southwood M, Ye L, Long L, Al-Lamki RS, Morrell NW. Id
proteins are critical downstream effectors of BMP signaling in human pulmonary
arterial smooth muscle cells. *Am J Physiol - Lung Cell Mol Physiol*.
2013;305(4):L312-L321. doi:10.1152/ajplung.00054.2013

10. Jouve C, Palmeirim I, Henrique D, Beckers J, Gossler A, Ish-Horowicz D, Pourquie
O. Notch signalling is required for cyclic expression of the hairy-like gene HES1 in
the presomitic mesoderm. *Development*. 2000;127(7):1421-1429.
doi:10.1242/dev.127.7.1421

11. Caplan BA, Schwartz CJ. Increased endothelial cell turnover in areas of in vivo
Evans Blue uptake in the pig aorta. *Atherosclerosis*. 1973;17(3):401-417.
doi:10.1016/0021-9150(73)90031-2

12. Schlereth K, Weichenhan D, Bauer T, Heumann T, Giannakouri E, Lipka D, Jaeger
S, Schlesner M, Aloy P, Eils R, Plass C, Augustin HG. The transcriptomic and
epigenetic map of vascular quiescence in the continuous lung endothelium. *eLife*.
2018;7:e34423. doi:10.7554/eLife.34423

13. Millauer B, Wizigmann-Voos S, Schnürch H, Martinez R, Møller NPH, Risau W,
Ullrich A. High affinity VEGF binding and developmental expression suggest Flk-1
as a major regulator of vasculogenesis and angiogenesis. *Cell*. 1993;72(6):835-
846. doi:10.1016/0092-8674(93)90573-9

14. Gerhardt H, Golding M, Fruttiger M, Ruhrberg C, Lundkvist A, Abramsson A,
Jeltsch M, Mitchell C, Alitalo K, Shima D, Betsholtz C. VEGF guides angiogenic
sprouting utilizing endothelial tip cell filopodia. *J Cell Biol*. 2003;161(6):1163-1177.
doi:10.1083/jcb.200302047

15. Bernatchez PN, Soker S, Sirois MG. Vascular Endothelial Growth Factor Effect on
Endothelial Cell Proliferation, Migration, and Platelet-activating Factor Synthesis Is
Flk-1-dependent *. *J Biol Chem*. 1999;274(43):31047-31054.
doi:10.1074/jbc.274.43.31047

16. Coller HA, Sang L, Roberts JM. A New Description of Cellular Quiescence. *PLOS
Biol*. 2006;4(3):e83. doi:10.1371/journal.pbio.0040083

17. Min M, Spencer SL. Spontaneously slow-cycling subpopulations of human cells
originate from activation of stress-response pathways. *PLoS Biol*.
2019;17(3):e3000178. doi:10.1371/journal.pbio.3000178

18. Johnson MS, Cook JG. Cell cycle exits and U-turns: Quiescence as multiple
reversible forms of arrest. *Fac Rev*. 2023;12:5. doi:10.12703/r/12-5

19. Hartwell LH, Culotti J, Pringle JR, Reid BJ. Genetic Control of the Cell Division
Cycle in Yeast. *Science*. 1974;183(4120):46-51. doi:10.1126/science.183.4120.46

20. Tarn WY, Lai MC. Translational control of cyclins. *Cell Div*. 2011;6:5.
doi:10.1186/1747-1028-6-5

21. MINSHULL J, PINES J, GOLSTEYN R, STANDART N, MACKIE S, COLMAN A,
BLOW J, RUDERMAN JV, WU M, HUNT T. The role of cyclin synthesis,
modification and destruction in the control of cell division. *J Cell Sci*.
1989;1989(Supplement_12):77-97. doi:10.1242/jcs.1989.Supplement_12.8

22. Kwon JS, Everetts NJ, Wang X, Wang W, Croce KD, Xing J, Yao G. Controlling
Depth of Cellular Quiescence by an Rb-E2F Network Switch. *Cell Rep*.
2017;20(13):3223-3235. doi:10.1016/j.celrep.2017.09.007

23. Fujimaki K, Li R, Chen H, Croce KD, Zhang HH, Xing J, Bai F, Yao G. Graded
regulation of cellular quiescence depth between proliferation and senescence by a

lysosomal dimmer switch. *Proc Natl Acad Sci*. Published online October 21, 2019.
doi:10.1073/pnas.1915905116

24. Rodgers JT, King KY, Brett JO, Cromie MJ, Charville GW, Maguire KK, Brunson C,
Mastey N, Liu L, Tsai CR, Goodell MA, Rando TA. mTORC1 controls the adaptive
transition of quiescent stem cells from G0 to GAlert. *Nature*. 2014;510(7505):393-
396. doi:10.1038/nature13255

25. Min M, Rong Y, Tian C, Spencer SL. Temporal integration of mitogen history in
mother cells controls proliferation of daughter cells. *Science*. 2020;368(6496):1261-
1265. doi:10.1126/science.aay8241

26. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T,
Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein
V, Eliceiri K, Tomancak P, Cardona A. Fiji: an open-source platform for biological-
image analysis. *Nat Methods*. 2012;9(7):676-682. doi:10.1038/nmeth.2019

27. Dardik A, Chen L, Frattini J, Asada H, Aziz F, Kudo FA, Sumpio BE. Differential
effects of orbital and laminar shear stress on endothelial cells. *J Vasc Surg*.
2005;41(5):869-880. doi:10.1016/j.jvs.2005.01.020

28. Liang D, Chang JR, Chin AJ, Smith A, Kelly C, Weinberg ES, Ge R. The role of
vascular endothelial growth factor (VEGF) in vasculogenesis, angiogenesis, and
hematopoiesis in zebrafish development. *Mech Dev*. 2001;108(1):29-43.
doi:10.1016/S0925-4773(01)00468-3

29. Mahmood T, Yang PC. Western Blot: Technique, Theory, and Trouble Shooting.
North Am J Med Sci. 2012;4(9):429-434. doi:10.4103/1947-2714.100998

30. Buglak DB, Bougaran P, Kulikauskas MR, Liu Z, Monaghan-Benson E, Gold AL,
Marvin AP, Burciu A, Tanke NT, Oatley M, Ricketts SN, Kinghorn K, Johnson BN,
Shiau CE, Rogers S, Guilluy C, Bautch VL. Nuclear SUN1 stabilizes endothelial
cell junctions via microtubules to regulate blood vessel formation. Koh GY,
Akhmanova A, eds. *eLife*. 2023;12:e83652. doi:10.7554/eLife.83652

31. A D, Ca D, F S, J D, C Z, S J, P B, M C, Tr G. STAR: ultrafast universal RNA-seq
aligner. *Bioinforma Oxf Engl*. 2013;29(1). doi:10.1093/bioinformatics/bts635

32. Putri GH, Anders S, Pyl PT, Pimanda JE, Zanini F. Analysing high-throughput
sequencing data in Python with HTSeq 2.0. *Bioinforma Oxf Engl*.
2022;38(10):2943-2945. doi:10.1093/bioinformatics/btac166

33. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol*. 2014;15(12):550.
doi:10.1186/s13059-014-0550-8

34. Wiecek AJ, Cutty SJ, Kornai D, Parreno-Centeno M, Gourmet LE, Tagliazucchi
GM, Jacobson DH, Zhang P, Xiong L, Bond GL, Barr AR, Secrier M. Genomic

hallmarks and therapeutic implications of G0 cell cycle arrest in cancer. Published
online April 15, 2023:2021.11.12.468410. doi:10.1101/2021.11.12.468410

35.   Almanzar N, Antony J, Baghel AS, Bakerman I, Bansal I, Barres BA, Beachy PA,
Berdnik D, Bilen B, Brownfield D, Cain C, Chan CKF, Chen MB, Clarke MF, Conley
SD, Darmanis S, Demers A, Demir K, de Morree A, Divita T, du Bois H, Ebadi H,
Espinoza FH, Fish M, Gan Q, George BM, Gillich A, Gòmez-Sjöberg R, Green F,
Genetiano G, Gu X, Gulati GS, Hahn O, Haney MS, Hang Y, Harris L, He M,
Hosseinzadeh S, Huang A, Huang KC, Iram T, Isobe T, Ives F, Jones RC, Kao KS,
Karkanias J, Karnam G, Keller A, Kershner AM, Khoury N, Kim SK, Kiss BM, Kong
W, Krasnow MA, Kumar ME, Kuo CS, Lam J, Lee DP, Lee SE, Lehallier B,
Leventhal O, Li G, Li Q, Liu L, Lo A, Lu WJ, Lugo-Fagundo MF, Manjunath A, May
AP, Maynard A, McGeever A, McKay M, McNerney MW, Merrill B, Metzger RJ,
Mignardi M, Min D, Nabhan AN, Neff NF, Ng KM, Nguyen PK, Noh J, Nusse R,
Pálovics R, Patkar R, Peng WC, Penland L, Pisco AO, Pollard K, Puccinelli R, Qi
Z, Quake SR, Rando TA, Rulifson EJ, Schaum N, Segal JM, Sikandar SS, Sinha
R, et al. A single-cell transcriptomic atlas characterizes ageing tissues in the
mouse. *Nature*. 2020;583(7817):590-595. doi:10.1038/s41586-020-2496-1

36.   Chen BR, Cheng HH, Lin WC, Wang KH, Liou JY, Chen PF, Wu KK. Quiescent
Fibroblasts Are More Active in Mounting Robust Inflammatory Responses Than
Proliferative Fibroblasts. *PLOS ONE*. 2012;7(11):e49232.
doi:10.1371/journal.pone.0049232

37.   Hayer A, Shao L, Chung M, Joubert LM, Yang HW, Tsai FC, Bisaria A, Betzig E,
Meyer T. Engulfed cadherin fingers are polarized junctional structures between
collectively migrating endothelial cells. *Nat Cell Biol*. 2016;18(12):1311-1323.
doi:10.1038/ncb3438

38.   Lee HW, Xu Y, He L, Choi W, Gonzalez D, Jin SW, Simons M. Role of Venous
Endothelial Cells in Developmental and Pathologic Angiogenesis. *Circulation*.
2021;144(16):1308-1322. doi:10.1161/CIRCULATIONAHA.121.054071

39.   Lee HW, Shin JH, Simons M. Flow goes forward and cells step backward:
endothelial migration. *Exp Mol Med*. 2022;54(6):711-719. doi:10.1038/s12276-022-
00785-1

40.   Pereira PD, Serra-Caetano A, Cabrita M, Bekman E, Braga J, Rino J, Santus R,
Filipe PL, Sousa AE, Ferreira JA. Quantification of cell cycle kinetics by EdU (5-
ethynyl-2′-deoxyuridine)-coupled-fluorescence-intensity analysis. *Oncotarget*.
2017;8(25):40514-40532. doi:10.18632/oncotarget.17121

41.   Goodrich DW, Wang NP, Qian YW, Lee EY, Lee WH. The retinoblastoma gene
product regulates progression through the G1 phase of the cell cycle. *Cell*.
1991;67(2):293-302. doi:10.1016/0092-8674(91)90181-w

42.   Buck SB, Bradford J, Gee KR, Agnew BJ, Clarke ST, Salic A. Detection of S-phase
cell cycle progression using 5-ethynyl-2′-deoxyuridine incorporation with click

chemistry, an alternative to using 5-bromo-2′-deoxyuridine antibodies.
BioTechniques. 2008;44(7):927-929. doi:10.2144/000112812

43.   Noseda M, Chang L, McLean G, Grim JE, Clurman BE, Smith LL, Karsan A. Notch
1041   activation induces endothelial cell cycle arrest and participates in contact inhibition:
1042   role of p21Cip1 repression. *Mol Cell Biol*. 2004;24(20):8813-8822.
1043   doi:10.1128/MCB.24.20.8813-8822.2004

1044   44.   Oesterle EC, Chien WM, Campbell S, Nellimarla P, Fero ML. p27(Kip1) is required
1045   to maintain proliferative quiescence in the adult cochlea and pituitary. *Cell Cycle*
1046   *Georget Tex*. 2011;10(8):1237-1248. doi:10.4161/cc.10.8.15301

1047   45.   Besson A, Gurian-West M, Chen X, Kelly-Spratt KS, Kemp CJ, Roberts JM. A
1048   pathway in quiescent cells that controls p27Kip1 stability, subcellular localization,
1049   and tumor suppression. *Genes Dev*. 2006;20(1):47-64. doi:10.1101/gad.1384406

1050   46.   Levenberg S, Yarden A, Kam Z, Geiger B. p27 is involved in N-cadherin-mediated
1051   contact inhibition of cell growth and S-phase entry. *Oncogene*. 1999;18(4):869-876.
1052   doi:10.1038/sj.onc.1202396

1053   47.   Miller I, Min M, Yang C, Tian C, Gookin S, Carter D, Spencer SL. Ki67 is a Graded
1054   Rather than a Binary Marker of Proliferation versus Quiescence. *Cell Rep*.
1055   2018;24(5):1105-1112.e5. doi:10.1016/j.celrep.2018.06.110

1056   48.   Fang JS, Coon BG, Gillis N, Chen Z, Qiu J, Chittenden TW, Burt JM, Schwartz MA,
1057   Hirschi KK. Shear-induced Notch-Cx37-p27 axis arrests endothelial cell cycle to
1058   enable arterial specification. *Nat Commun*. 2017;8. doi:10.1038/s41467-017-
1059   01742-7

1060   49.   Chassot AA, Turchi L, Virolle T, Fitsialos G, Batoz M, Deckert M, Dulic V,
1061   Meneguzzi G, Buscà R, Ponzio G. Id3 is a novel regulator of p27kip1 mRNA in
1062   early G1 phase and is required for cell-cycle progression. *Oncogene*.
1063   2007;26(39):5772-5783. doi:10.1038/sj.onc.1210386

1064   50.   Murata K, Hattori M, Hirai N, Shinozuka Y, Hirata H, Kageyama R, Sakai T, Minato
1065   N. Hes1 directly controls cell proliferation through the transcriptional repression of
1066   p27Kip1. *Mol Cell Biol*. 2005;25(10):4262-4271. doi:10.1128/MCB.25.10.4262-
1067   4271.2005

1068   51.   Herrera MD, Mingorance C, Rodríguez-Rodríguez R, Alvarez de Sotomayor M.
1069   Endothelial dysfunction and aging: an update. *Ageing Res Rev*. 2010;9(2):142-152.
1070   doi:10.1016/j.arr.2009.07.002

1071   52.   SEALS DR, JABLONSKI KL, DONATO AJ. Aging and vascular endothelial function
1072   in humans. *Clin Sci Lond Engl 1979*. 2011;120(9):357-375.
1073   doi:10.1042/CS20100476

53.　Donato AJ, Machin DR, Lesniewski LA. Mechanisms of Dysfunction in the Aging
Vasculature and Role in Age-Related Disease. *Circ Res*. 2018;123(7):825-848.
doi:10.1161/CIRCRESAHA.118.312563

54.　Wiecek AJ, Cutty SJ, Kornai D, Parreno-Centeno M, Gourmet LE, Tagliazucchi
GM, Jacobson DH, Zhang P, Xiong L, Bond GL, Barr AR, Secrier M. Genomic
hallmarks and therapeutic implications of cancer cell quiescence. *bioRxiv*.
Published online January 1, 2022:2021.11.12.468410.
doi:10.1101/2021.11.12.468410

55.　Liu B, Wang X, Jiang L, Xu J, Zohar Y, Yao G. Extracellular Fluid Flow Induces
Shallow Quiescence Through Physical and Biochemical Cues. *Front Cell Dev Biol*.
2022;10:792719. doi:10.3389/fcell.2022.792719

56.　Mack JJ, Mosqueiro TS, Archer BJ, Jones WM, Sunshine H, Faas GC, Briot A,
Aragón RL, Su T, Romay MC, McDonald AI, Kuo CH, Lizama CO, Lane TF, Zovein
AC, Fang Y, Tarling EJ, Vallim TQ de A, Navab M, Fogelman AM, Bouchard LS,
Iruela-Arispe ML. NOTCH1 is a mechanosensor in adult arteries. *Nat Commun*.
2017;8(1):1620. doi:10.1038/s41467-017-01741-8

57.　Helle E, Ampuja M, Antola L, Kivelä R. Flow-Induced Transcriptomic Remodeling of
Endothelial Cells Derived From Human Induced Pluripotent Stem Cells. *Front
Physiol*. 2020;11. Accessed April 7, 2023.
https://www.frontiersin.org/articles/10.3389/fphys.2020.591450

58.　Zhang MJ, Pisco AO, Darmanis S, Zou J. Mouse aging cell atlas analysis reveals
global and cell type-specific aging signatures. Han JDJ, Tyler JK, Hou L, eds. *eLife*.
2021;10:e62293. doi:10.7554/eLife.62293

59.　Acharya BR, Fang JS, Jeffery ED, Chavkin NW, Genet G, Vasavada H, Nelson EA,
Sheynkman GM, Humphries MJ, Hirschi KK. Connexin 37 sequestering of
activated-ERK in the cytoplasm promotes p27-mediated endothelial cell cycle
arrest. *Life Sci Alliance*. 2023;6(8). doi:10.26508/lsa.202201685

60.　Hayflick L, Moorhead PS. The serial cultivation of human diploid cell strains. *Exp
Cell Res*. 1961;25(3):585-621. doi:10.1016/0014-4827(61)90192-6

61.　Marthandan S, Priebe S, Hemmerich P, Klement K, Diekmann S. Long-Term
Quiescent Fibroblast Cells Transit into Senescence. *PLOS ONE*.
2014;9(12):e115597. doi:10.1371/journal.pone.0115597

62.　Yin K, Patten D, Gough S, de Barros Gonçalves S, Chan A, Olan I, Cassidy L,
Poblocka M, Zhu H, Lun A, Schuijs M, Young A, Martinez-Jimenez C, Halim TYF,
Shetty S, Narita M, Hoare M. Senescence-induced endothelial phenotypes
underpin immune-mediated senescence surveillance. *Genes Dev*. 2022;36(9-
10):533-549. doi:10.1101/gad.349585.122

63. Miyauchi H, Minamino T, Tateno K, Kunieda T, Toko H, Komuro I. Akt negatively
regulates the in vitro lifespan of human endothelial cells via a p53/p21-dependent
pathway. *EMBO J.* 2004;23(1):212-220. doi:10.1038/sj.emboj.7600045

64. Chen Y, Liu Z, Chen H, Huang X, Huang X, Lei Y, Liang Q, Wei J, Zhang Q, Guo
X, Huang Q. p53 SUMOylation Mediates AOPP-Induced Endothelial Senescence
and Apoptosis Evasion. *Front Cardiovasc Med*. 2022;8. Accessed April 4, 2023.
https://www.frontiersin.org/articles/10.3389/fcvm.2021.795747

65. Minamino T, Miyauchi H, Yoshida T, Ishida Y, Yoshida H, Komuro I. Endothelial
Cell Senescence in Human Atherosclerosis. *Circulation*. 2002;105(13):1541-1544.
doi:10.1161/01.CIR.0000013836.85741.17

66. Sang L, Coller HA, Roberts JM. Control of the reversibility of cellular quiescence by
the transcriptional repressor HES1. *Science*. 2008;321(5892):1095-1100.
doi:10.1126/science.1155998

67. Ausprunk DH, Folkman J. Migration and proliferation of endothelial cells in
preformed and newly formed blood vessels during tumor angiogenesis. *Microvasc
Res*. 1977;14(1):53-65. doi:10.1016/0026-2862(77)90141-8

68. McDonald AI, Shirali AS, Aragón R, Ma F, Hernandez G, Vaughn DA, Mack JJ, Lim
TY, Sunshine H, Zhao P, Kalinichenko V, Hai T, Pelegrini M, Ardehali R, Iruela-
Arispe ML. Endothelial Regeneration of Large Vessels Is a Biphasic Process
Driven by Local Cells with Distinct Proliferative Capacities. *Cell Stem Cell*.
2018;23(2):210-225.e6. doi:10.1016/j.stem.2018.07.011

1132
1133

1134   **Nonstandard Abbreviations and Acronyms**
1135
HUVEC: Human Umbilical Vein Endothelial Cells
Flow-M: Flow maintenance; 72 hr 15d/cm^2
Flow-E: Flow establishment; 72 hr 15d/cm^2
siRNA: Small interfering RNA
KD: Knockdown
A.U.: Arbitrary unit
pRB: Phospho retinoblastoma protein
EdU: 5-Ethynyl-2'-deoxyuridine
HES1: Hes Family BHLH Transcription Factor 1
ID3: Inhibitor of DNA Binding 3
1146

Figure 1. Laminar flow-mediated quiescence is transcriptionally distinct.
(a) Quantification of epithelial quiescence score in HUVEC scRNA dataset by cluster.
Clusters were previously defined[8]. Flow-M, flow maintenance (laminar flow
(15d/cm^2/72hr)). Static cells were visually checked for subconfluence and fixed 48 hr
post-seeding. **(b)** Quantification of endothelial quiescence score in HUVEC scRNA
dataset by cluster. **(c)** Endothelial quiescence score on bulk RNA seq data of HUVEC
exposed to indicated stimuli, n= 3 replicates. **(d-e)** Heatmaps showing relative
expression of cell cycle proliferation markers (green font) and inhibitors (red font) plotted
using bulk RNAseq data of HUVEC under indicated conditions, n=3 replicates.
Statistics, one-way ANOVA with Tukey's multiple comparison test; ** $p \leq 0.01$, *** $p \leq$
*0.001, ****$p \leq 0.0001$, ns = not significant*.

Figure 2. Endothelial cell flow maintenance quiescence stimulus leads to shallow quiescence depth.
(a) Representative images of HUVEC under static (non-flow) or Flow-M (flow maintenance) conditions with EdU incorporation and fixation at indicated times post Flow-M release. Cultures stained for DAPI (white, nuclear mask) and EdU (red, s-phase) or pRB (blue, interphase). Scale bar, 50 µm. White arrow, flow direction. **(b)** Quantification of percent EdU+ cells with indicated conditions. n=3 replicates, 5 images per condition per replicate. **(c)** Quantification of percent pRB+ cells with indicated conditions. n=3 replicates, 5 images per condition per replicate. **(d)** Representative images of HUVEC under low or high density conditions with EdU incorporation and fixation at indicated times post high-density release. Cultures stained for DAPI (white, nuclear mask) and EdU (red, s-phase) or pRB (blue, interphase). Scale bar, 50 µm. **(e)** Quantification of percent EdU+ cells with indicated conditions. n=3 replicates, 6 images per condition per replicate. **(f)** Quantification of percent pRB+ cells with indicated conditions. n=3 replicates, 6 images per condition per replicate. Statistics, one-way ANOVA with Tukey's multiple comparisons test; *$p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, ****$p \leq 0.0001$, ns = not significant*.

Figure 3. Cell cycle inhibitor p27 expression levels vary with endothelial quiescence stimulus.
(a) RT-qPCR for *CDKN1B* levels in indicated conditions. n=3 replicates. (b) Representative images of HUVEC under indicated conditions stained for p27 (green) and DAPI (white, nuclear mask). Scale bar, 50 μm. White arrow, flow direction. (c) Quantification of HUVEC p27+ cells under indicated conditions, n=3 replicates, 5 images per condition per replicate. (d) RT-qPCR for *CDKN1B* levels in indicated conditions. n=3 replicates. (e) Representative images of HUVEC under indicated conditions stained for p27 (green) and DAPI (white, nuclear mask). Scale bar, 50 μm. (f) Quantification of HUVEC p27+ cells under indicated conditions. n=3 replicates, 5 images per condition per replicate. Statistics, student's two-tailed *t*-test; ** $p \leq 0.01$, ****$p \leq 0.0001$.

Figure 4. Endothelial quiescence depth varies with laminar flow exposure time and positively correlates with p27 expression levels
(a) Representative images of HUVEC with indicated siRNA treatments and conditions. after EdU incorporation (red, S-phase) and staining with DAPI (white, nuclear mask). Scale bar, 50 µm. White arrow, flow direction. **(b)** Quantification of EdU+ cells with indicated conditions. n=3 replicates, 5 images per condition per replicate. **(c)** Representative images of HUVEC with indicated conditions and stained for p27 (inhibitor), p21 (inhibitor), Ki67 (proliferation marker), and DAPI (nucleus) in white. Scale bar, 50 µm. White arrow, flow direction. **(d-f)** Quantification of percent p27+ (d), p21+ cells (e), or Ki67+ cells (f). n=3 replicates, 5 images per condition per replicate, **(g)** Representative images of HUVEC under indicted conditions and after EdU incorporation (red, S-phase) and stained for pRB (blue, interphase) and with DAPI (white, nuclear mask), Flow-E, flow establishment; Flow-M, flow maintenance. Scale bar, 50 µm. White arrow, flow direction. **(h)** Quantification of EdU+ cells in indicated conditions. n=3 replicates, 5 images per condition per replicate. **(i)** Quantification of pRB+ cells in indicated conditions. n=3 replicates, 5 images per condition per replicate. Statistics, one-way ANOVA with Tukey's multiple comparisons test; *$p \leq 0.05$, ****$p \leq 0.0001$, ns = not significant*.

Figure 5. BMP and NOTCH regulated p27 repressors *HES1* and *ID3* regulate p27
levels and flow-mediated quiescence depth
(a) Representative images of HUVEC under indicated conditions and with indicated
siRNA treatment. Cells stained for p27 (green) and DAPI (white, nuclear mask). Scale
bar, 50 μm. White arrow, flow direction. **(b)** Quantification of p27+ cells under indicated
conditions and treatments. n=3 replicates, 5 images per condition per replicate. **(c)**
Representative images of HUVEC with indicated conditions and siRNA treatments.
Cells were pulse labeled with EdU (red, S-phase) and stained for pRB (blue, interphase)
and DAPI (white, nuclear mask). Scale bar, 50 μm. White arrow, flow direction. **(d)**
Representative images of HUVEC with indicated conditions and siRNA treatments.
Cells were pulse labeled with EdU (red, S-phase) and stained for pRB (blue, interphase)
and DAPI (white, nuclear mask). Scale bar, 50 μm. White arrow, flow direction. **(e)**
Quantification of EdU+ cells with indicated conditions. n=3 replicates, 5 images per
condition per replicate. **(f)** Quantification of pRB+ cells with indicated conditions. n=3
replicates, 5 images per condition per replicate. **(g)** Quantification of EdU+ cells with
indicated conditions. n=3 replicates, 5 images per condition per replicate. **(h)**
Quantification of pRB+ positive cells with indicated conditions. n=3 replicates, 5 images
per condition per replicate. Statistics, one-way ANOVA with Tukey's multiple
comparisons test. *** $p \leq 0.01$, **** $p \leq 0.001$, ***** $p \leq 0.0001$, ns = not significant*.

Figure 6. *Cdkn1b* expression does not accumulate with age *in vivo*
(a) scRNAseq analysis of normalized *Cdkn1b* expression in endothelial cells isolated
from mice of indicated ages. See Methods and text for details of dataset reanalysis.
Endothelial cells with *Cdkn1b* expression at 0 were excluded from analysis. ****$p \leq$
0.0001, ns = not significant by one-way ANOVA with Tukey's multiple comparisons test.
(b) scRNAseq analysis of percent *Cdkn1b+* endothelial cells isolated from neonatal (P8)
or adult (2-2.5m) murine ears. Each dot represents one endothelial subpopulation in
one sample. Statistics, student's two-tailed *t*-test **(c)** Proposed model for dynamic
regulation of endothelial cell flow-mediated quiescence depth with flow exposure time.
In the first 16-24 hr (Flow-E, flow establishment), a deep quiescence is established
accompanied by high levels of cell cycle inhibitor p27. With time, transcription factors
that are targets of flow-mediated Notch and BMP signaling are upregulated, and these
p27 transcriptional repressors prevent ongoing p27 synthesis and lead to significantly
reduced p27 levels. The regulators are required to transition to a shallow quiescence
depth that characterizes extended exposure to laminar flow (Flow-M, flow
maintenance).
1452