Paradoxical anti-arrhythmic effects of Zim3, a CRISPRi effector, in human iPSC-cardiomyocytes

Julie L. Han, Yuli W. Heinson and Emilia Entcheva
Department of Biomedical Engineering, The George Washington University

Abstract:
We show that Zim3, when used as Zim3-KRAB-dCas9 effector in interference CRISPR, without any guide RNAs, paradoxically upregulates key cardiac ion channel genes in human induced pluripotent stem-cell-derived cardiomyocytes, iPSC-CMs, responsible for healthy resting membrane potential, repolarization of the action potential and electrical transmission of signals. These were found to yield expected functional enhancements consistent with a more mature iPSC-CM phenotype, with anti-arrhythmic properties.

Zinc Finger imprinted 3 (Zim3 or ZNF657/264) is a protein-coding gene in humans, found at low levels in testis, brain, skeletal muscle and skin[1]. As other ZNF-KRAB (Krueppel associated box) domains, which bind to DNA and through complexing with KAP1 (KRAB-associated protein 1) affect chromatin organization[2,3], Zim3-KRAB may be involved in the regulation of transcription at multiple genomic loci. Recently, Zim3-KRAB has been identified as a potent effector (repressor) in combination with dCas9 for interference CRISPR (CRISPRi) gene regulation studies[4]. Indeed, using Zim3-KRAB-dCas9 adenoviral delivery in combination with suitable gRNAs in post-differentiated induced pluripotent stem cell-derived cardiomyocytes (iPSC-CMs), we have shown robust suppression of the expression of select ion channel genes, e.g. KCNH2, with concomitant functional consequences[5]. While performing control experiments, we observed unexpected effects of Zim3-KRAB-dCas9 alone on cardiac electrophysiology without the addition of any gRNAs or with the use of scrambled/control gRNA. Namely, we found Zim3-KRAB-dCas9 expression to significantly upregulate the mRNA expression of several key cardiac ion channels: KCNJ2, encoding for the Kir2.1 protein – inward rectifier K+ channel, contributing to the maintenance of healthy negative resting membrane potential; KCNH2, encoding for the main repolarizing K+ channel in cardiomyocytes, the rapid delayed rectifier that controls action potential duration, APD; and GJA1, encoding for Cx43 – the main gap-junctional protein in ventricular cardiomyocytes; for Cx43 this was also confirmed at the protein level, **Figure 1**. Using all-optical electrophysiology[6-9], **Figure 1a**, we pursued functional studies, **Figure 1c-e**, which indeed corroborated physiologically relevant functional effects, expected for upregulated KCNH2 (APD shortening in paced conditions), **Figure 1c, f**; upregulated KCNJ2 (reduced or eliminated spontaneous firing), **Figure 1d, f**; and upregulated GJA1 (conduction velocity increase), **Figure 1e-h**. In combination, these electrophysiological changes are perceived as desirable and a signature of mature iPSC-CMs with anti-arrhythmic properties[10]. These gene expression changes and functional effects cannot be attributed to the expression of dCas9, as we have shown that regular KRAB-dCas9[11] without gRNAs does not alter these genes. The fluorescent reporter, mCherry, also does not have such effects. Related earlier-generation effector construct for CRISPRi – Mec2p-KRAB-dCa9[12] caused similar, but much milder changes when delivered in the same type of viral vector - adenovirus with an SFFV (spleen focus-forming virus) promoter and an mCherry reporter, at the same multiplicity of infection, MOI 1000, as Zim3-KRAB-dCas9. In fact, we are unaware of any pharmacological or other single perturbation acting as a potent agonist for all three genes. These findings should be taken into consideration when performing CRISPRi studies in iPSC-CMs. Zim3 also needs to be investigated further as a potential therapeutic tool for iPSC-CMs maturation for their use in drug screening, disease modeling and regenerative applications[13-15]; furthermore, it would be interesting to explore if Zim3 may exert similar effects *in vivo* in the adult human heart as an anti-arrhythmic therapy.

Figure 1. Effects of dCas9/Zim3 expression in human iPSC-CMs: RNA, protein, and electrophysiological consequences. a. Schematic illustrating the experimental workflow with all-optical electrophysiology and molecular characterization. b. Validation of virally-mediated delivery of mCherry tagged constructs (mCherry, Mec2p-KRAB-dCa9-mCherry, Zim3-KRAB-dCa9-mCherry) in human iPSC-CMs. Scale bar 10 μmc. Functional changes in action potential duration (APD) and calcium transient duration (CTD) under 0.5 Hz pacing upon expression of the various constructs (n=9-34 biologically independent samples; one-way ANOVA with post-hoc Tukey correction, **$p<0.01$). d. Changes on spontaneous beat frequency in dCas9/Zim3 expressing cells compared to non-infected control. (n=10 biologically independent samples; unpaired t-test, **$p<0.01$). e. Activation maps of hiPSC-CMs expressing dCas9/MECP2 and dCas9/Zim3 compared to non-infected controls; representative images and summary of

normalized conduction velocity (CV_{norm}) changes in the differentially treated groups at 0.75 Hz and 1.0 Hz pacing frequencies (n=2-9 biologically independent samples). Data are represented as mean ± S.E.M. f. qPCR analysis of KCNH2, GJA1, and KCHJ2 of pristine samples expressing dCas9/Zim3 compared to mCherry control (n=4-7 biologically independent samples, n=3 technical replicates; unpaired t-test, **$p<0.01$). g. qPCR analysis of KCNH2 and GJA1 mRNA changes upon expression of dCas9/MECP2 and dCas9/Zim3 compared to non-infected control post functional experiments (n=4-16 biologically independent samples, n=3 technical replicates; one-way ANOVA, **$p<0.01$). h. Western blot (Wes by ProteinSimple) protein quantification of Cx43 upon the different treatments post all-optical electrophysiology (n=6-13 biologically independent samples; one-way ANOVA, **$p<0.01$). Biorender was used for some of the panels.

References

Kim, J., Bergmann, A., Wehri, E., Lu, X. & Stubbs, L. Imprinting and Evolution of Two Kruppel-type Zinc-Finger Genes, ZIM3 and ZNF264, Located in the PEG3/USP29 Imprinted Domain. *Genomics* **77**, 91-98, doi:https://doi.org/10.1006/geno.2001.6621 (2001).

Lupo, A. *et al.* KRAB-Zinc Finger Proteins: A Repressor Family Displaying Multiple Biological Functions. *Curr Genomics* **14**, 268-278, doi:10.2174/13892029113149990002 (2013).

Stoll, G. A., Pandiloski, N., Douse, C. H. & Modis, Y. Structure and functional mapping of the KRAB-KAP1 repressor complex. *The EMBO Journal* **41**, e111179, doi:https://doi.org/10.15252/embj.2022111179 (2022).

Alerasool, N., Segal, D., Lee, H. & Taipale, M. An efficient KRAB domain for CRISPRi applications in human cells. *Nat Methods* **17**, 1093-1096, doi:10.1038/s41592-020-0966-x (2020).

Han, J. L., Heinson, Y. W., Chua, C. J., Liu, W. & Entcheva, E. CRISPRi Gene Modulation and All-Optical Electrophysiology in Post-Differentiated Human iPSC-Cardiomyocytes. *bioRxiv*, 2023.2005.2007.539756, doi:10.1101/2023.05.07.539756 (2023).

Klimas, A., Ortiz, G., Boggess, S. C., Miller, E. W. & Entcheva, E. Multimodal on-axis platform for all-optical electrophysiology with near-infrared probes in human stem-cell-derived cardiomyocytes. *Prog Biophys Mol Biol* **154**, 62-70, doi:10.1016/j.pbiomolbio.2019.02.004 (2020).

Klimas, A. *et al.* OptoDyCE as an automated system for high-throughput all-optical dynamic cardiac electrophysiology. *Nature communications* **7**, 11542, doi:10.1038/ncomms11542 (2016).

Heinson, Y. W., Han, J. L. & Entcheva, E. Portable low-cost macroscopic mapping system for all-optical cardiac electrophysiology. *J Biomed Opt* **28**, 016001, doi:10.1117/1.Jbo.28.1.016001 (2023).

Liu, W., Han, J. L., Tomek, J., Bub, G. & Entcheva, E. Simultaneous Widefield Voltage and Dye-Free Optical Mapping Quantifies Electromechanical Waves in Human Induced Pluripotent Stem Cell-Derived Cardiomyocytes. *ACS Photonics* **10**, 1070-1083, doi:10.1021/acsphotonics.2c01644 (2023).

Karbassi, E. *et al.* Cardiomyocyte maturation: advances in knowledge and implications for regenerative medicine. *Nature reviews. Cardiology* **17**, 341-359, doi:10.1038/s41569-019-0331-x (2020).

Gilbert, L. A. *et al.* CRISPR-mediated modular RNA-guided regulation of transcription in eukaryotes. *Cell* **154**, 442-451, doi:10.1016/j.cell.2013.06.044 (2013).

Yeo, N. C. *et al.* An enhanced CRISPR repressor for targeted mammalian gene regulation. *Nat Methods* **15**, 611-616, doi:10.1038/s41592-018-0048-5 (2018).

Gintant, G. *et al.* Use of Human Induced Pluripotent Stem Cell-Derived Cardiomyocytes in Preclinical Cancer Drug Cardiotoxicity Testing: A Scientific Statement From the American Heart Association. *Circ Res* **125**, e75-e92, doi:10.1161/res.0000000000000291 (2019).

Yang, X., Ribeiro, A. J. S., Pang, L. & Strauss, D. G. Use of Human iPSC-CMs in Nonclinical Regulatory Studies for Cardiac Safety Assessment. *Toxicological sciences : an official journal of the Society of Toxicology* **190**, 117-126, doi:10.1093/toxsci/kfac095 (2022).

Barker, R. A. *et al.* Lessons learnt, and still to learn, in first in human stem cell trials. *Stem cell reports*, doi:10.1016/j.stemcr.2022.11.019 (2022).