Evolutionary history of Calcium-sensing receptors sheds light into hyper/hypocalcemia-

causing mutations

Aylin Bircan[1], Nurdan Kuru[1], Onur Dereli[1], Ogün Adebali[1,2*],

1 Faculty of Engineering and Natural Sciences, Sabanci University, İstanbul, Türkiye

2 TÜBİTAK Research Institute for Fundamental Sciences, Gebze, Türkiye

* oadebali@sabanciuniv.edu

Abstract

The Calcium Sensing Receptor (CaSR) is very important in controlling the levels of calcium in

the body by interacting with different types of G-protein. This receptor is highly conserved

among other G-protein coupled receptors (GPCRs) and has been linked to disorders affecting

the balance of calcium in the body, such as hypercalcemia and hypocalcemia. Although there

has been progress in understanding the structure and function of CaSR, there is still a lack of

knowledge about which specific residues are important for their function and how it differs from

other receptors in the same class. In this study, we used phylogeny-based methods to identify

functionally-equivalent orthologs of CaSR, predict the importance of each residue, and calculate

specificity-determining position (SDP) scores to uncover the evolutionary basis of its function.

Our results showed that the CaSR subfamily is highly conserved, with higher SDP scores than

its closest receptor subfamilies. Residues with high SDP scores are likely to be critical in

receptor activation and pathogenicity. We applied gradient-boosting trees with evolutionary

metrics as inputs to predict the functional consequences of each substitution, and discriminate

between gain and loss-of-function mutations those causing hypo- and hypercalcemia,

respectively. Our study provides insight into the evolutionary fine-tuning of CaSR, which can

help understand its role in calcium balance and related disorders.

Introduction

Calcium sensing receptor (CaSR) is a class C G-protein-coupled receptor (GPCR) that

maintains extracellular Ca2+ homeostasis by sensing calcium ions in the blood and regulating

parathyroid hormone release and urinary calcium[1, 2] . The CaSR is activated by Ca2+ and L-

amino acids such as L-Phe and L-Trp as well as polyamines and polypeptides[3-5]. Ligands

bind to the extracellular Venus flytrap (VFT) domain of the receptor-like the other class C

GPCRs such as metabotropic glutamate receptors [6].

Class C GPCRs are obligatory dimers, forming either homo or heterodimers [6]. CaSR forms a

homodimer where each subunit is composed of an extracellular domain (ECD), comprising a

bilobed (LB1, LB2) VFT and a cysteine-rich domain (CRD) connected to a heptahelical

transmembrane (7TM) domain[3, 5].

Crystal structures of the ECD [4, 7] and cryo-electron microscopy structures of the full-length

CaSR[3, 5, 8-10] reveal the structural basis for activation mechanisms and ligand binding sites.

L-amino acid binding site at the interdomain cleft of LB1-LB2[3-5, 11-13] and multiple Ca2+

amino acid binding sites on the VFT domain are shown in the literature[3-5]. While Ca2+ is the

composite agonist to the CaSR, L-amino acids promote the receptor activation along with the

Ca2+, but they are not able to activate the receptor alone[14]. Even though Ca2+ alone

activates the receptor in functional assays[14], whether it activates the CaSR in the absence of

L-amino acid is still controversial[3, 5].

Variants in CaSR may cause malfunctions that result in Ca2+ homeostasis diseases in humans.

More than 400 germline loss/gain-of-function mutations cause hypercalcaemic disorders,

neonatal severe hyperparathyroidism (NSHPT) and the milder familial hypocalciuric

hypercalcemia type-1 (FHH1) and autosomal dominant hypocalcemia type-1 (ADH1)

respectively[2]. Many more CaSR variants are anticipated to be identified as more population-

level genetic data become available[2]. Understanding the role of each residue in receptor

structure and activation mechanisms could provide additional information about the likelihood of

variant pathogenicity and CaSR signaling. The role of each residue in a receptor can be

revealed by comparison of receptors in a family and between different families; however, the

structure and complete activation mechanisms of many families in class C GPCRs are still

unknown, especially G-protein coupled receptor family C group 6 member A (GPRC6A) and

type 1 taste receptors (TAS1Rs; members 1,2 and 3) that are the closest subfamilies to CaSR.

While all subfamily receptors of class C GPCRs share common domains and structural features,

details of responding to different ligands and activating signaling pathways are diverse between

even closely related receptors[6]. Gene duplication is the main mechanism that generates new

protein functions across GPCRs. Protein families are evolved by speciation events following a

gene duplication[15, 16]; thus sequence comparisons of members within a subfamily and

between subfamilies can show the evolutionarily conserved domains as well as diverged protein

sites that distinguish one subfamily from others. One challenge with this analysis is that

excessive gene duplication events complicate the identification of functionally identical orthologs

in a subfamily. Moreover, the conservation patterns in paralogs and distant homologs may help

inferring the specific roles of a single residue in protein function. Because the evolutionary

pressure on the paralogs and close orthologues are not the same, allowed substitutions on

paralogs may not be acceptable in close orthologues. Thus, using functionally identical

orthologs in sequence comparisons is crucial to infer the role of each residue in a protein family.

Here, we show the importance of each residue in CaSR by comparing it with the closely related

subfamilies, GPRC6A and TAS1Rs. We identified all orthologues sequences in each subfamily

by phylogenetic tree analysis. To obtain orthologues without requiring computationally

expensive phylogenetic tree step, subfamily-specific profile HMMs are generated from the true

orthologues in subfamilies that we determined by the phylogenetic tree analysis. We calculated

a specificity score for each residue in a subfamily by calculating scores based on a modified

version of PHACT[17] scores which considers independent evolutionary events on the

phylogenetic tree while scoring the acceptability of an amino acid substitution. We predicted the

functional consequence of each substitution in CaSR by using the gradient boosting trees

machine learning approach.

Results

Evolutionary History of Class C GPCRs

To reveal the evolutionary constraints on protein families, we developed to develop a strategy to

precisely define a protein subfamily. Precise subfamily definition can be precisely accomplished

by revealing the evolutionary history of the superfamily. Evolutionary history of gene families

can only be established by reconstructing high-quality phylogenetic trees, which can be used to

pinpoint gene duplication events. Discrimination between gene duplication and speciation nodes

enabled us to define of the paralogous and orthologous protein sequences. We further analyzed

the phylogenetic trees to classify the orthologous sequences that are likely equivalent in

function. We used functionally-equivalent orthologs in comparative analyses between

subfamilies, which eventually yielded subfamily-specific signatures that can be used to define

that particular subfamily and its function. Finally, the association between the signature and

function would enable a better understanding of specific molecular mechanisms and the effects

of variants, particularly for the protein subfamily of interest. Here, we aim to reveal the

signatures of the CaSR subfamily, that is implied in the specific function of calcium-sensing and

downstream signaling.

We have retrieved the complete proteomes of 478 species from the NCBI database. To identify

proteins that belong to class C GPCR family, we performed searched profile HMM of the seven

transmembrane domain profile (Pfam: 7tm_3) (Fig 1) against the proteomes. While this search

allowed us to retrieve the entire class C GPCRs hitting the hmm profile, it does not yield

subfamily (22 human GPCRs) annotations. We performed scan profile HMM of the PfamA

profile against to Class C GPCR to select canonical isoforms. We used seven-transmembrane

domain only to assign subfamilies and trimmed the N-terminus region of this domain for further

homology steps. To generate a general HMM profile for each subfamily, we first applied a Blast

search using each human class-C as a query. For each subject, we blasted them against the

human proteome. We retrieved the bidirectional best hits (Core subfamily assignment).

Figure 1: **Summary of the Methodological Framework**. *478 complete proteomes were retrieved from NCBI*

database. Each sequence was searched by HMMsearch against Pfam 7tm_3 domain profile to retrieve all class C

GPCRs. Domain architectures of class C GPCRs were determined by HMMscan against PFAM.A profile to identify

canonical isoforms. Species-specific BLAST databases from TM domains of the canonical isoforms were built. Bi-

directional mutual best hits were detected by blasting each canonical sequence against the species databases (Core

subfamily assignment). Core subfamily sequences were aligned and ML trees were built to make subfamily profile

HMMs. By HMMsearch against subfamily profile HMMs other sequences in the subfamilies were found (Subfamily

extension). Sequences in each subfamily were aligned and ML trees were built. Based on the ML trees paralogs were

filtered and functionally identical groups were identified (FIG).

For proteins that did not have bidirectional mutual best hits, we assigned them to a subfamily

based on their homology search against the HMM profiles generated in the previous step

(Subfamily Extension). We produced maximum likelihood (ML) trees of extended subfamilies

and filtered paralogous sequences to obtain functionally identical groups (FIGs).

The CaSR subfamily produced over five thousand hits, which included vomeronasal and

olfactory receptors that have never been shown to sense calcium. Previous research has shown

that CaSR is classified in the pheromone/olfactory cluster of class C GPCRs[18] (18). In species

that had multiple proteins assigned to the CaSR subfamily, we constructed a maximum

likelihood tree using these hits and other human class C GPCR protein sequences. These trees

revealed that a significant number of duplication events occurred in the species after the clade

diverged from CaSR. As a result, we defined this diverged clade as a new subfamily named

CaSR-likes. The sequences in this subfamily is unlikely to maintain calcium homeostasis, and

therefore should not be annotated as calcium-sensing receptors.

We selected representative sequences from different species for each subfamily of 22 different

receptor subfamilies and 264 CaSR-like sequences and built a ML tree (Fig 1A). Also, we built

the ML trees of all proteins from CaSR, GPRC6A, taste receptors and merged these trees to the

representative tree of class C GPCRs (Fig 2 A). The resulting phylogeny shows that are five

major clades: CaSR-related, GABA, mGluR, Orphans, and retinoic acid induced (RAIG).

Orphan receptors, GPR158 and GPR179, formed a clade that was diverged from other

receptors consistent with previous trees[19] and with 0.95 transfer bootstrap (TB) value. γ-

aminobutyric acidB receptors (GABBR1 and GABBR2) formed another clade diverged from

GPR156 with 0.97 TB. γ-aminobutyric acid$_B$ receptors evolved earliest that have a common

ancestor with the highest taxonomic rank (33213–Bilateria) compared to other subfamilies.

CaSR group (CaSR, CaSR-likes, GPRC6A and taste receptors) was diverged from

metabotropic glutamate receptors (GRM1-8) and RAIG receptors (GPRC5A, GPRC5B,

GPRC5C, GPRC5D) with 1 and 0.98 TB values respectively. Within the CaSR group clade

CaSRs and CaSR-likes were diverged from GPRC6A and taste receptors with 1 TB. Except

TAS1R1 and TAS1R2, all CaSR group subfamilies have a common ancestor from taxonomy

clade 7776-Gnathostomata. TAS1R1 and TAS1R2 were more specific than other CaSR group

subfamilies that were evolved from 117571-Euteleostomi. Comparison analysis of branch

lengths[20] among common species between CaSR, GPRC6A and taste receptors shows that

CaSR subfamily is significantly more conserved than its closest subfamilies (Fig 2 B)

The higher diversity of CaSR-likes relative to CaSRs is reflected in the ML tree (Fig 2A). Branch

lengths of CaSR-likes are longer in contrast to shorter branch lengths in CaSR. Longer branch

lengths show that more variation, and thus divergence occurred in the CaSR-like clade.

Moreover, extensive gene duplication events occurred in this clade. For instance, rodents such

as Dipodomys ordii (taxid:10020),Octodon degus**(**taxid:10160**)** and snakes such as

Notechis scutatus(taxid: 8663) have more than a hundred receptors that match to CaSR profile.

However, these matches include type 2 vomeronasal receptors (V2R) and type 2 vomeronasal

receptor likes. Among mammals, V2R genes exhibit significant variation. While dogs, cows, and

primates except prosimians do not have functional V2Rs**,** rodents, reptiles and fish have

multiple intact V2Rs[21].Since these receptors do not have functional orthologs in mammals,

separating them from functionally-equivalent CaSRs is crucial**.**

*Figure 2: **Evolution of Class C GPCRs**. (A) The maximum likelihood phylogenetic tree of Class C GPCRs,*
spanning representative species from each subfamily is shown. Subfamilies are represented as circular layers around
the ML tree. All twenty-two Class C GPCR subfamilies are shown in the inner circle. In addition to these subfamilies,
vomeronasal and other orphan receptors are represented as CaSR-like receptors. All proteins in CaSR, GPRC6A
and TAS1Rs are merged to this representative species tree. (B) Branch lengths from leaf to the root of the common
species that exist in all CaSR, GPRC6A and TAS1Rs are taken from the subfamily trees. Welch's t-Test by using
ggstatsplot package results are shown on the graph.

Subfamily-specific Profile HMMs to Obtain Orthologs

In the class C GPCR family, gene duplication events give rise to new specificity, and each duplicated gene with a new function is evolved by further speciation events and produce a set of orthologous sequences[15, 16]. Each subfamily of class C shares relatively conserved membrane-spanning region as well as a degree of variability underling functional differences. At the molecular level, residues that are responsible for certain functional characteristics such as ligand and coupling selectivity are called specificity-determining residues[15]. Conservation analysis from multiple sequence alignments can be used to find residues that are conserved in all subfamilies through evolution as well as specificity-determining residues that are only conserved in a subfamily and differ in other subfamilies. However, the success of this method depends on the sequences that are used to build alignments. Hence, it is vital to use functionally identical orthologs in the analysis.

The seven-transmembrane domains of class C GPCRs are used to build a class-specific general profile for this family (Pfam:7tm_3). However, this domain cannot be used to differentiate subfamilies further.

Moreover, excessive gene duplication events as seen in the CaSR-like clade requires precise phylogenetic analysis to differentiate CaSR and CaSR-like sequences. Also, subfamily specific profile HMMs are shown to be promising methods to detect protein sequences belong to a protein subfamily, as well as separation of homologs and non-homologs [22, 23].Therefore, we built subfamily-specific profile HMMs that match with all orthologs of a subfamily while excluding closely related like sequences (Fig 3).

*Figure 3: **Subfamily Specific HMM Models.** Subfamily specific HMM model method. Based on the phylogenetic tree*
the target, the closest and the rest groups are determined. Representative amino acids in each group are selected,
and their scores are calculated. Weights to scale the emission probability are calculated.

We define the target family, its closest family (phylogenetic neighboring clade), and the rest
based on the phylogenetic tree. We weighted the identity score of each amino acid to calculate
the emission probabilities. The highest weight is given to the residues which are only conserved
in the target subfamily; hence they differentiate one subfamily from the others. Minimum weight
is given to the residues which are conserved both in the target subfamily and its closest clade.
We tested our subfamily-specific profile HMMs' performance on an independent sequences
retrieved UNIPROT dataset[24] and not seen during the training process. We assigned
sequences to their corresponding subfamilies by following the same steps as NCBI dataset[25]
used to build these models. We selected new taxons that were not in the NBCI dataset as test
sequences. Our subfamily-specific profile HMMs correctly hit all members of a subfamily while
they do not hit any protein from another subfamily (Table 1).

Table 1: Subfamily Specific Profile HMM's Performance

Subfamily HMM	Test Cases	Hits	Missed
CaSR	81	81	-
GPRC6A	62	62	-
TAS1R1	75	75	-
TAS1R2	21	21	-
TAS1R3	74	74	-

Specificity Determining Residues

CaSR is distinguished from other subfamilies of class C GPCRs by its oversensitivity to many

substitutions that are caused either gain or loss of function mutations, because it maintains

systemic calcium homeostasis and highly sensitive to a very slight change in extracellular Ca2+

concentrations[26]. Since CaSR is the most conserved and ancestral subfamily among the

CaSR-likes, GPRC6A and TAS1Rs, it is reasonable to expect in some positions can be under

the relaxation of existing purifying selection in any CaSR-likes, GPCR6A or TAS1Rs, but not in

CaSR. On the other hand, at some positions the same amino acid remains functionally

important in both subfamilies, and at others a position remains important in each subfamily but a

different amino acid is favored in each duplicate.

To identify and order residues that differentiate a subfamily from its closest relatives, we

employed multiple sequence alignment- and phylogenetic tree-based approaches. Specificity-

determining residues that are conserved in a subfamily, but differ from its sister clade can be

predicted by directly comparing ancestral family sequences and calculating their divergence

scores (26). However, using multiple sequence alignments only does not discriminate between

the number of substitution event. For example, a single substitution event in the common

ancestor of bony fish clade of CaSR subfamily can be inherited to multiple descendants'

sequences. Assessing this single event as independent events result in overcounting of these

changes as if they are independent. Hence, the position is considered (i) to tolerate that

particular amino acid and (ii) functionally less important. In contrast, a single evolutionary event

might have been compensated by other substitutions in the same evolutionary node. Such a

substitution might not be tolerated in the other clades of the subfamily.

Another consideration to identify and order specificity-determining residues is treating

substitution events on the phylogenetic tree unequally. When an amino acid in CaSR remains

the same but can differ in the nearby subfamily, CaSR-likes, it indicates that the amino acid has

a unique purpose for CaSR. The SDP score of such an amino acid must be high. If an amino

acid is conserved in both CaSR and remote subfamilies like taste receptors but likely to be

substituted in CaSR-likes, it suggests that the amino acid plays a common functional role in

both CaSR and other subfamilies. For such an amino acid, the SDP score must be low, since it

is not a specific position for CaSR.

For CaSR group(CaSR, GPRC6A and TAS1Rs), we identified and order residues by specificity

which differentiate a subfamily from others by using an adaption of functionally divergent

residues method[27] along with an adaption of PHACT method[17]. We calculated probability of

each amino acid at each node of the CaSR-group phylogenetic tree by ancestral sequence

reconstruction (Fig 4 A). Starting from the root of the tree, we identified each substitution event

and at which subfamily node that event happened. Counting the number of independent

substitution events in a subfamily clade and comparing the probability of the same substitution

in other subfamily clades, we ordered the specificity-determining residues. We assumed that if

an amino acid is allowed to change on sister subfamily nodes and poorly conserved in sister

subfamily nodes while it is highly conserved on the target subfamily node, it is a specific residue

to the target subfamily only. If a substitution event is observed on a clade close to the target

node, we consider that event to increase specificity of a residue because it diverges the target

group from its closest, sister clade. The details and the algorithm are given in materials and

methods (Algorithm 3).

*Figure 4:***Specificity Determining Position Scores. (A)** *Calculation of SDP scores uses the phylogenetic tree,*

*probability distribution of amino acids at each ancestral node. **(B)** SDP score distributions of each subfamily are*

*shown. **(C)**Cryo-EM structure of human CaSR bound with Ca²⁺and L-Trp (PDB:7DTV) and homology models of*

We calculated specificity scores for each CaSR, GPRC6A and TAS1Rs. Specificity score

distributions show that CaSR subfamily have more specific residues compared to other

subfamilies (Fig 4 B). On the VFT domain, specific residues are clustered different regions. We

found a cluster of specific residues on the interdomain cleft between LB1-LB2 that is the L-

amino acid binding site in other class C GPCRs[3]. It suggests that this region is the primary

Ca2+ binding site in CaSR consistent with[14]. We found two different clusters of specific

residues on the ECD. First cluster was on the LB1 domain and on the LB1-LB1 dimer interface.

LB1 domain plays a role in anchoring ligands and initiating domain twisting by conformational

changes at the interface between LB1 regions[3, 5]. The second cluster was found at the

cytosolic side of the LB2 and at the interface between LB2-CRD where Ca2+ ions are bind[3-5].

Interaction between LB2 subunits are required for CaSR activation that propagates to large-

scale transitions of the 7TMDs[3, 5]. Specific residues on the LB1 domain, LB1-LB1 dimer

interface and LB2-CRD interface indicate that they provide the structural conformational

changes upon ligand binding to the interdomain cleft. Mutations located on these regions are

associated with loss and gain of function mutations (Fig 6)[2]. Other specific residues are found

on the CR, ECL2 and TM domains. On the ECL2 acidic residues D758 and E759 are specific to

CaSR. The intersubunit electrostatic repulsion between the ECL2 regions could facilitate the

activation of CaSR[3, 5]. In the agonist+PAM bound state the ECL2 is moved by the interaction

among E759, W590, and K601. Deletion of D758 and E759, and single mutations of K601E and

W590E disrupts the CaSR activity, however Δ758–759 mutant was expressed at the cell surface

with the comparable levels to that of WT, while W590E and K601E mutants were expressed on

the cell surface lower than the WT level[3]. We found that residues W590 and K601 are not

specific to CaSR. The TM domains of two protomers of CaSR come into close proximity upon

receptor activation[5].

The orientation of the TM5-TM6 dimer in the CaSR distinguishes it from other Class C receptors

such as mGluR and GABA$_B$ receptors, which results in its inactive conformation[9]. The

interaction between TM4-5 of each subunit in the inactive state is essential[14], while the

interaction between TM6-TM6 is crucial for the active state[3, 8, 14]. The structural findings and

the presence of CaSR specific residues on each TM domain suggest that CaSR is specialized

in both dimerization and ligand binding. Specific residues on TM domain guarantees the correct

orientation for activation upon ligand binding and inactive conformation otherwise. Interactions

between the domains and ligand-receptor are quite sensitive that slight changes cause

malfunctions in the receptor. On the other hand, GPRC6A and taste receptors are more prone to

acceptable substitutions and they are not very specialized to respond a single ion. GPRC6A and

taste receptors are activated by a broad spectrum of ligands[28, 29]. Even though the ligand of

GPRC6A is controversial in the literature, multiple ligands such as osteocalcin (Ocn),

testosterone, basic amino acids and cations such as L-Arg, L-Lys, L-Orn, calcium, magnesium,

and zinc are suggested to bind GPRC6A[29]. Taste receptors bind to different ligands including

sugar, L- and D-amino acids, sweet proteins, and artificial sweeteners[30].

On the TM region we also find CaSR specific cholesterol recognition/interaction amino acid

consensus (CRAC) motif (L783,F789,S820) that is defined by the consensus (L/V)X1–5YX1–

5(R/K) and is often present at junctions between membrane- and cytosol-exposed domains and

shown in GRM2 receptor[31]. Phylogenetic analysis shows that TAS1R3 evolved earliest (7776

Gnathostomata) among TAS1Rs, TAS1R1and TAS1R2 subfamilies have common ancestor

117571 Euteleostomi. TAS1R3 forms heterodimers with TAS1R1 and TAS1R2[28, 30, 32].

Interactions between the cytosolic terminus of the extracellular CRD is needed for T1R3

dimerization. TAS1R1 and TAS1R2 recognize a broad spectrum of L-amino acids that bind to

the intercleft between LB1-LB2 and induce the positional shift of the CRD regions, however

T1R3 loses the corresponding function[32]. Our analysis showed that TAS1R1 have specific

residues on LB1, LB2 and extracellular loop regions. Also, TAS1R2 has specific residues on

LB1,LB2 and CR domains. On the other hand, in TAS1R3, we found specific residues only on

the LB1 and one on the CR domain. Since LB1-LB2 domains create a cavity for ligand binding,

specific residues on LB1-LB2 domains of TAS1R1 and TAS1R2 may contribute to domain

transformation upon ligand binding. However, the number and distribution of specific amino

acids suggest that taste receptors are not under selective pressure as CaSR.

Gradient Boosting Trees Machine Learning Approach to Predict the Mutation Types in

CaSR Because CaSR is a highly conserved subfamily, any substitution on the receptor disrupts

the function of the receptor and causes either gain or loss of function mutations (Fig 3C).

However, predicting the functional consequence of a substitution is challenging. Evolutionary

conservation of a residue among subfamilies might reflect the common structural constraints,

but it does not distinguish between loss and gain of function mutations (i.e., LoF and GoF,

respectively). In addition, at some positions substitution to different amino acids causes either

loss or gain of function mutations[21]. We hypothesized that "activating" mutations are more

likely to be tolerated in the neighboring clades such as GPRC6A and TAS1Rs and not in CaSR

whereas, in general, loss-of-function (inactivating) mutations are not tolerated in the larger clade

of these receptor subfamilies. To test this hypothesis whether we can discriminate between GoF

and LoF mutations in CaSR, we applied a tree boosting machine learning algorithm,

XGBoost[33] that linked multiple features such as conservation scores, physico-chemical

properties of amino acids and domain information.

We used sequence-based features, identity scores from multiple sequence alignments, physico-

chemical properties of amino acids, and domain information as input features to train our model

(Fig 5 A). Since we calculated our feature values from the multiple sequence alignments, we

divided our dataset into training, validation and test datasets before we created feature matrices

to prevent information leakage. We performed 50 replications with different random splitting of

datasets to obtain a more robust model performance.

*Figure 5: **Gradient Boosting Trees Machine Learning Approach to Predict the Mutation Types in CaSR.***

***(A)**Model architecture. We used MSA of CaSR,CaSR-likes, GPRC6A and TAS1Rs to generate features as well as*

*amino acid physico-chemical features and domain information. We performed 50 replications. **(B)** The performance*

and feature importance of XGBoost algorithm. AUROC and AUPR values of 50 replications are shown. Average AUC

levels of 50 replications are 0.83 and 0.78 for the train and test respectively. Average AUPR levels of 50 replications

are 0.93 and 0.9 for the train and test respectively. Contributions of Shapley values for type of pathogenicity

classification to the model output for XGBoost. aa0:the amino acid found in the human CaSR,aa1:substituted amino

Table 2: Model's predictions for the new CASR gain and loss of function mutations from the literature. The correct

predictions are indicated by a star symbol (*) next to them.

Mutation	Cause	Prediction
p.I857S[34]	hypocalcemia	gain-of-function*
p.Y825F [35]	hypocalcemia	gain-of-function*
p.P393R [36]	hypercalcemia	loss-of-function*
p.C60G[37]	hypercalcemia	loss-of-function*
p.D99N[38]	hypercalcemia	loss-of-function*
p.T186N[39]	hypocalcemia	loss-of-function
p.A840V[24]	hypocalcemia	gain-of-function*
p.S448P[40]	hypercalcemia	loss-of-function*
p.L696V[41]	hypocalcemia	gain-of-function*
p.D433Y[42]	hypercalcemia	loss-of-function*
p.S147L[43]	hypercalcemia	loss-of-function*
p.D398N[44]	hypercalcemia	loss-of-function*
p.K805R[45]	hypercalcemia	gain-of-function
p.C60Y[46]	hypercalcemia	loss-of-function*
p.S820N[47]	hypocalcemia	loss-of-function
p.L606P[48]	hypercalcemia	loss-of-function*
p.H41R[49]	hypercalcemia	gain-of-function
p.A110D[50]	hypercalcemia	gain-of-function
p.I139T[51]	hypocalcemia	gain-of-function*
p.Q164R[52]	hypercalcemia	loss-of-function*
p.T699N[53]	hypercalcemia	gain-of-function
p.R701G[53]	hypercalcemia	loss-of-function*
p.T808P[53]	hypercalcemia	loss-of-function*

The ROC and PR curves are used to understand the performance of a binary classifier that assigns each element of data into two groups. ROC curve is a graphical plot that shows the false positive rate versus the true positive rate for different threshold values between 0.0 and 1. A PR curve is a plot of the precision and the recall for different threshold values and it is useful for imbalance datasets. We used the areas under the ROC and PR curves (i.e., AUC and AUPR, respectively) to compare the performances of the model on the train and test datasets for 50 replications. Higher AUC and AUPR values are associated with better performance. AUC and AUPR over all replications were shown in (Fig 5 B). Our average AUC values for training and test among 50 replications are 0.83 and 0.78 (Fig 5 B). Our average main AUPR values for training and test among 50 replications are 0.93 and 0.9 respectively (Fig 5 B). After we reported our algorithm performance, we trained our algorithm with the whole dataset. We tested our algorithm with new test cases from literature (Table 2). Additionally, we categorized amino acids that are observed in the CaSR MSA as neutrals. To date, no pathogenic substitution has been reported in the literature for these amino acids that we identified as neutral. We visualized all predictions in the form of a heatmap for every other amino acids at each position until the disordered region (position 892) of the human CaSR (Fig 6 A). We mapped known CaSR loss and gain of function mutations on the cryo-EM structure of human CaSR bound with Ca^{2+} and L-Trp (PDB:7DTV (3)) (Fig 6 B). There is a tendency that loss-of-function mutations are on the outer-core regions, while gain-of-function mutations are on the inner-core regions. In the heatmap we observed a similar prediction pattern that gain-of-function predictions are mostly in the inner-core regions. SHAP (SHapley Additive exPlanations) values provide a way to decode the inner workings of a machine learning model like XGBoost. These values calculate the average contribution of each feature to the overall prediction, taking into account any interactions between the features. Based on the SHAP values, the conservation scores of human CaSR amino acids in other subfamilies play a significant role in the model's prediction, as shown in Figure 5B. If the amino acid is also conserved in GPRC6A and taste receptors (in fact conservation score in TAS1R3 has the highest contribution), the model predicts a substitution of that amino acid as loss-of-function. Another important feature is the domain of the amino acid. Our findings indicate that if the amino acid is located in the TM domain, a substitution would result in a gain-of-function mutation.It is known that the majority of gain of function mutations are located in the TM domain, as shown in Figure 6B. The presence of certain amino acids on the TM domain of CaSR suggests that they play a crucial role in its activation mechanism. Even though substituting those amino acids might be acceptable in

GPRC6A and taste receptors, they might lead to the lock of TM domains and result in the

overactivation of CaSR.

*Figure 6: **All Possible Amino Acid Substitution Predictions. (A)Visualizing** the precision of our XGBoost model.*

The heatmap displays the XGBoost model's predictions for each of the 20 amino acids at every position except

*disordered regions (892-1078) within the human CaSR. **(B)** Mutations on human CaSR structure. Loss and gain of*

function associates mutations are shown on the cryo-EM structure of human CaSR bound with Ca2+ and L-Trp

(PDB:7DTV) as blue and red spheres respectively.

Discussion

In this study, we showed the evolution of CaSR by developing a methodology in precisely

defining functionally equivalent orthologous sequences across species and therefore

subfamilies. We built a high-quality phylogenetic tree of CaSR with its closest subfamilies,

GPRC6A and TAS1Rs. Statistical analysis of branch length distances from this phylogenetic

tree showed that CaSR is evolutionarily more conserved compared to GPRC6A and TAS1Rs.

While GPRC6A and taste receptors can bind to a diverse range of ligands and able to tolerate

substitutions at most of the positions, CaSR requires a delicate balance for proper functioning.

High evolutionarily conservation and specificity of CaSR in contrast to closest subfamilies are

reflected in specificity determining position (SDP) score analysis. CaSR has specific residues

clustered on different regions of the receptor. They are located on Ca2+ and L-Trp binding sites

on the VFT, as well as on the dimerization sites between two sub-units of the homodimer.

Specific residues on the dimer interfaces indicate that dimerization maintained by interactions

between different subunits is required for ligand binding and correct activation of the CaSR.

Ca2+ ions binding and interactions between LB2-CR domains and conformational changes in

LB1 domain were suggested that they are required to activate CaSR[3-5]. Mutational analysis at

some positions on LB1 domain have been shown to reduce the effect of Ca2+-stimulated

intracellular Ca^{2+} mobilization in cells[3, 5]. In contrast, substitutions caused negative charge

neutralization on the ECL2 result in prompting the activation of CaSR[5]. Our results suggest

that residues with low SDP scores on any domain are required for common activation

mechanism since they are conserved across functionally different receptor subfamilies.

However, residues with high SDP scores cause malfunctions in the CaSR. Any substitution in a

residue with high SDP score might either cause over or less activation. Deep mutational

scanning approaches or new methods that simultaneously profile variant libraries[54] are

needed to provide further evidence to functionally assay all possible missense mutants.

To predict the functional consequence of a mutation in human CaSR, we used Extreme

Gradient Boosting (XGBoost) method. XGBoost is able to perform well on small datasets by

incorporating variety of regularization methods to control the model complexity, which helps to

prevent overfitting. We have a small and unbalanced dataset in that the number of gain-of-

function mutations was very low, therefore it is prone to overfit. To prevent overfitting while

achieving high predictive performance, we used a simple method along with regularization

parameters. Moreover, we tried to keep the ratio between the number of loss-of-function and the

number of gain-of-function mutations for training and test sets as close as possible. To get a

robust performance, we repeated the train-validation-test splitting procedure fifty times. To

increase the predictive performance, we could use a more complex methods such as deep

learning, however they require larger datasets. Studies that used deep learning or ensemble

methods for similar assessment are different in terms of prediction in which they predict the type

of a mutation as only pathogenic or neutral [55-58]. Even though there are number of mutations

of human CaSR in the Clinvar, the functional consequences of most of them are not known.

Given the constraints of the small dataset and limited additional data, we carefully selected and

processed the features for our model's training. Features that are used to train a machine

learning model heavily determine the performance of the model. The more features we use, the

more information the model has to learn from, which can lead to improved predictive

performance. However, having too many features can also lead to overfitting. Moreover, the

quality of the features is more important than the quantity. One important evolutionary process

that can affect the functional consequence of a substitution is co-evolution. From the multiple

sequence alignment of CaSR proteins, we manually selected six

positions,p.180,p.212,p.228,p.241,p.557 and p.883, that are in our dataset and co-evolved. We

masked the co-evolved amino acids from the MSA and performed train-validation-test splitting

procedure fifty times again. Our average AUC values for training and test among 50 replications

were 0.83 and 0.77 respectively, and average AUPR values were 0.93 and 0.89. Despite not

experiencing an improvement in performance, we found that the amino acid changes p.I212T,

p.F180C, and p.I212S were now predicted to cause loss of function, contrary to their previous

prediction of causing gain of function. We cannot accurately assess the impact of co-evolution

on performance because there is a lack of effective tools for identifying co-evolved positions and

our dataset contains only a limited number of co-evolved positions, but we anticipate that it is an

important feature to differentiate gain and loss of function.

We built subfamily-specific profile HMMs to get all functionally-equivalent orthologs while

excluding other proteins. To generate these HMM models, we manually decided target, closest

and rest groups based on the phylogenetic tree of CaSR group. Based on the nature of a

phylogenetic tree, selection of these groups is changed, so that this process can be further

automated. We did not anticipate our specific models to match any receptor from other classes

of GPCRs, since they are evolutionarily more distant to CaSR group. We expect that our

subfamily specific profile HMMs can be used to obtain orthologs in different protein families for

the upcoming genomes. They can be particularly useful for studying protein families with many

duplications and orphan protein families, where it can be difficult to identify true members.

These models are particularly important to avoid computationally expensive and expertise-

required phylogenetic tree reconstruction and analysis.

Materials and methods

Class C Proteins and Their Domain Architectures

478 complete eukaryotic proteomes were downloaded from NCBI genomes website(https : //f

tp.ncbi.nlm.nih.gov/genomes/archive/old ref seq/) in 2018. hmmsearch of HMMER

software[59](http://hmmer.org/) was run for each proteome against Pfam 7TM3 profile[60].

Sequences with significant 7TM3 hit based on hmmsearch results (above the default threshold)

were compiled from proteomes. hmmscan of HMMER software[59](http://hmmer.org/) was run

for these sequences against Pfam-A 32.0 database[60]. Based on the results of hmmscan, the

longest isoform was taken and saved in a separate file named by taxonomic id, however

canonical sequences were obtained for human (based on given canonical proteins in UniProt

website[61]). Because plants do not have GPCRs, plants were eliminated from the analysis. For

single isoform sequences of each proteome a BLAST database was built[62].

Subfamily Definition and Subfamily Specific Models

Each protein sequence of each taxon was queried through BLASTP against each prepared

BLAST database[62]. reciprocal mutual best hits of each human class C GPCR were collected

in a file named gene id. reciprocal mutual best hits of each class C GPCR and remaining human

class C GPCRs were collected and 7TM domains of these sequences were taken based on

hmmscan results (Longest sequence which hit the 7TM3). Sequences were aligned using

MAFFT v7.221 E-INS-I algorithm with default parameters[63]. Maximum likelihood based

phylogenetic tree (ML tree) of each subfamily of class C GPCR was built using RAxML version

8.2.12 with automatic protein substitution model selection (PROTGAMMAAUTO) and 100 rapid

bootstrapping parameters[64]. Most common lowest taxonomic level was added to the

phylogenetic tree with ETE toolkit[65]. Based on the phylogenetic tree, sequences belong to the

corresponding subfamily were taken and an profile HMM was built. Subfamily Assignment The

process begins by scanning each sequence with a 7TM3 domain against profile Hidden Markov

Models (profile HMMs). After the sequence is scanned, the subfamily is determined based on

three conditions: (1)The maximum score value of the hmmscan must belong to the given

subfamily. (2) E-value is a measure of the significance of a match in a database search and the

lower the E-value, the more significant the match is. The E-value of the sequence must be the

lowest. (3)The sequence must belong to the most common highest taxonomic level of the given

subfamily. Taxonomic level refers to the classification of an organism within a biological

classification system. If a sequence meets these three conditions, it is assigned to the

corresponding subfamily. After this, the full length sequences of each subfamily were then

aligned using the MAFFT v7.221 algorithm[63] and trimmed using the gappy-out method of the

trimAl tool[66].

Paralog Filter

There were a number of duplications in CaSR subfamily. For example, Dipodomys ordii has 116

CaSR sequences. To reduce the number of seqeunces,human CaSR and other human class C

GPCR proteins sequences compiled with CaSR seqeunces of each taxon, and and aligned with

MAFFT v7.221 auto algortihm[63], and the gappy-out method of the trimAl tool was used to trim

the multiple sequence alignments (MSA)[66]. ML tree was built using RAxML-NG v0.9.0 with ML

tree search and bootstrapping (Felsenstein Bootstrap and Transfer Bootstrap) parameters[67].

Based on the ML tree, proteins that were diverged from the common ancestor of the human

CaSR clade were classified as CaSR-likes. Proteins that were clustered with the human CaSR

were accepted as CaSRs. After we assigned all proteins to their subfamilies, we built final ML

trees for CaSR,GPRC6A, and TAS1Rs. We added human CaSR sequence was added to

GPRC6A and TAS1Rs subfamilies, and human GPRC6A sequence was added to CaSR

subfamily as an outgroup. We aligned each subfamily sequences with MAFFT v7.221 einsi

algortihm[63] and built the ML trees by using RAxML-NG v0.9.0 with FTT model parameter[67].

We labeled the duplications at each node on the ML trees. Based on the duplications, we

manually checked the trees and removed a clade that was a subset of its sister clade by using

ETE toolkit[65]. We took each branch and node length from leaf to root of the tree by using

common species in all CaSR, GPRC6A and taste receptor trees to calculate subfamily

conservation by using Welch's t-Test by using ggstatsplot package[20].

Subfamily Specific Profile HMMs

[63].After we took all receptors from CaSR, CaSR-like, GPRC6A, and taste receptors, we

aligned them by using MAFFT v7.221 auto algortihm[63]. For each subfamily we removed the

positions from the multiple sequence alignment (MSA) that correspond to a gap in the human

receptor. Then, we divided the MSA into subfamily alignments. We generated a HMM from the

gap removed alignment of each subfamily, and we added weight to the emission probabilities of

the HMMs. To calculate emission probability weights, based on the maximum likelihood

phylogenetic tree (ML tree) we defined the target, the closest and the rest groups. We took the

closest node as the closest group and other nodes as the rest. According to that we have five

different scenarios:

• CaSR is the target group, CaSR-likes are the closest group, and GPRC6A and taste
receptors, (TAS1Rs) are the rest.
• GPRC6A is the target group, TAS1Rs are the closest group, CaSR and CaSR-likes are
the rest.
• TAS1R1 is the target group, TAS1R2 is the closest group and TAS1R3 is the rest.
• TAS1R2 is the target group, TAS1R1 is the closest group and TAS1R3 is the rest.
• TAS1R3 is the target group, TAS1R1 and TAS1R2 are the closest group and GPRC6A
is the rest.

ALGORITHM 1: REPRESENTATIVE AMINO ACID AND INITIAL SCORE FOR POSITION "K"

Input: Representative amino acid of target subfamily, R_T; the frequency of R_T in the target, S_T; the most frequent amino acid of subfamily i, (i=1,...,N) and its frequency, a_i,

F_i, respectively; the number of subfamilies in close and rest groups, n_c and n_r, respectively; conservation threshold for target and close/rest groups, thr_1 and thr_2; the threshold for Blosum scores, thr_{bls}.

STEP 1: *Choose representative amino acid and related frequency for each group*

1   *for $j \in \{c, r\}$*

2   *if $n_j = 1$*

3   *$R_j = a_k$ where k is the subfamily in group j*

4   *$S_j = F_k$*

5   *else*

6   *if $R_T \in \{a_j, j = 1, \ldots, n_j\}$*

7   *$R_j = R_T$*

8   *$S_j = F_k$ where k is the subfamily with the most frequent amino acid is R_T*

9   *else*

10   *$R_j = a_k$ where k is group with highest frequency*

11   *$S_j = \frac{\sum_{s=1}^{n_j} F_s}{n_j}$*

STEP 2: *Assign position type and initial score to position "k"*

Category 1

12 *if $R_c = R_r$ and they are gap*

13   *if R_T is gap*

14   *$type_k = II$*

15   *$score_k = \frac{1}{\sum_{i \in \{T,c,r\}} S_i}$*

16   *else*

17   *$type_k = I$*

18   *$score_k = \sum_{i \in \{T,c,r\}} S_i$*

Category 2

19 *else if R_T is gap*

20   *if R_c is gap or R_r is gap*

21 $type_k = II$

22 $score_k = \dfrac{1}{\sum_{i \in \{T,c,r\}} S_i}$

23   *else*

24 $type_k = I$

25 $score_k = \sum_{i \in \{T,c,r\}} S_i$

Category 3

26   *else if $R_T \neq R_c \neq R_r$*

27   *if R_T, R_r and R_c are not gaps*

28   *if $S_T \geq thr_1$ and $S_c, S_r \geq thr_2$*

29 $type_k = I$

30 $score_k = \sum_{i \in \{T,c,r\}} S_i$

31   *else if $Blosum(R_T, R_c) \leq thr_{bls}$ and $Blosum(R_T, R_r) \leq thr_{bls}$*

32 $type_k = I$

33 $score_k = \sum_{i \in \{T,c,r\}} S_i$

34   *else*

35 $type_k = IV$

36 $score_k = \sum_{i \in \{T,c,r\}} S_i$

37   *else if R_c is gap*

38   *if $S_T \geq thr_1$ and $S_r \geq thr_2$*

39 $type_k = I$

40 $score_k = \sum_{i \in \{T,c,r\}} S_i$

41   *else if $Blosum(R_T, R_r) \leq thr_{bls}$*

42 $type_k = I$

43 $score_k = \sum_{i \in \{T,c,r\}} S_i$

44   *else*

45 $type_k = IV$

46 $score_k = \sum_{i \in \{T,c,r\}} S_i$

47   else if R_r is gap

48   if $S_T \geq thr_1$ and $S_c \geq thr_2$

49 $type_k = I$

50 $score_k = \sum_{i \in \{T,c,r\}} S_i$

51   else if $Blosum(R_T, R_c) \leq thr_{bls}$

52 $type_k = I$

53 $score_k = \sum_{i \in \{T,c,r\}} S_i$

54   else

55 $type_k = IV$

56 $score_k = \sum_{i \in \{T,c,r\}} S_i$

Category 4

57   else if $R_T = R_c$

58 $type_k = II$

59 $score_k = \dfrac{1}{\sum_{i \in \{T,c,r\}} S_i}$

Category 5

60   else if $R_T \neq R_c$ and $R_T = R_r$

61   if R_c is gap

62 $type_k = III$

63 $score_k = \sum_{i \in \{T,c,r\}} S_i$

64   else

65   if $Blosum(R_T, R_c) \leq thr_{bls}$ and $S_c \geq thr_2$

66 $type_k = III$

67 $score_k = \sum_{i \in \{T,c,r\}} S_i$

68   else

69 $type_k = II$

70 $\qquad score_k = \dfrac{1}{\sum_{i \in \{T,c,r\}} S_i}$

Category 6

71 else if $R_T \neq R_c$ and $R_c = R_r$

72 $\qquad$ if $S_T \geq thr_1$ and $S_c, S_r \geq thr_2$

73 $\qquad\qquad type_k = I$

74 $\qquad\qquad score_k = \sum_{i \in \{T,c,r\}} S_i$

75 $\qquad$ else if $Blosum(R_T, R_c) \leq thr_{bls}$

76 $\qquad\qquad type_k = I$

77 $\qquad\qquad score_k = \sum_{i \in \{T,c,r\}} S_i$

78 $\qquad$ else

79 $\qquad\qquad type_k = II$

80 $\qquad\qquad score_k = \dfrac{1}{\sum_{i \in \{T,c,r\}} S_i}$

ALGORITHM 2: COMPUTE WEIGHT FOR ALL POSITIONS

Input: Types for each position k (k=1,…,K), $type_k$; initial score for each position k of type t, $score_k^t$; number of type i positions, n_i where $n_1 + n_2 + n_3 + n_4 = K$; a predefined constant value as max weight of Type II positions, c_2.

Weight of Type I positions

1 for $p_1 = 1:n_1$

2 $\qquad weight_{p_1} = \dfrac{score_{p_1}^1}{\min_{l=1,\dots,n_1}(score_l^1)}$

Weight of Type IV positions

3 for $p_2 = 1:n_2$

4 $\qquad weight_{p_2} = \dfrac{score_{p_2}^2}{\max_{l=1,\dots,n_2}(score_l^2)} mean\left(weight_{p_1}\right)_{p_1 \in \{1,\dots,n_1\}}$

Weight of Type III positions

5 for $p_3 = 1:n_3$

6 $\qquad$ if target is CaSR

7 $$c_1 = \frac{\min\left(weight_{p_2}\right)_{p_2 \in \{1,\dots,n_2\}}}{2}$$

8   *else*

9 $$c_1 = \min\left(weight_{p_2}\right)_{p_2 \in \{1,\dots,n_2\}}$$

4 $$weight_{p_3} = \frac{score_{p_3}^3}{\max_{l=1,\dots,n_3}\left(score_l^3\right)} c_1$$

Weight of Type II positions

7   *for* $p_4 = 1 : n_4$

8 $$weight_{p_4} = \frac{score_{p_4}^4}{\max_{l=1,\dots,n_4}\left(score_l^4\right)} c_2$$

Subfamily Specific Position Scores

From the alignment we used to make subfamily specific profile HMMs, we randomly selected

264 CaSR like sequences (same number of sequences as CaSRs) and took all CaSR (264

proteins), GPRC6A (242 proteins) and TAS1Rs (TAS1R1 has 210, TAS1R2 has 173 and

TAS1R3 has 273 proteins). We built an ML tree by using IQ-TREE multicore version 2.0.6[68]

with automatic model selection[69] (-m MFP) and ultrafast bootstrap[70] (-bb 1000) parameters.

For CaSR, GPRC6A, and TAS1Rs, we removed the positions from the multiple sequence

alignment that correspond to a gap in the human receptor respectively. By using gap removed

alignments and the ML tree, we did ancestral sequence reconstructions for each subfamily with

IQ-TREE multicore version 2.0.6 with -m JTT+R10 model parameter[68]. We showed specific

residues that have a SDP score higher than 5, on the structures. We used cyro-EM structure of

CaSR (PDB:7DTV) and Swiss models[71] for GPRC6A and taste receptors since they do not

have experimental structures. To visualize structures and residues we used UCSF Chimera

tool[72].

We calculated SDP scores by a method extended from[27] by considering the phylogenetic

trees and a phylogeny-based scoring approach, adjPHACT, based on the methodology of

PHACT algorithm. The details of how we compute SDP score for any position k can be found in

Algorithm 3. PHACT computes the tolerance for each amino acid for the query specie which is

human by using a tree traversal approach. By checking the probability differences, PHACT

detects the location of amino acid substitutions and compute weighted summation of positive

probability differences based on the distance between the node of change and human. On the

other hand, here we aim to determine the acceptability of each amino acid per subfamily. To

achieve this, we modify PHACT by starting the tree traversal from the root node and eliminating

the node weighting approach. At the end, we have a probability distribution per position for each

subfamily which is computed by considering the independent events. Again, we determine the

representative amino acid for target subfamily by picking the most frequently observed amino

acid and its adjPHACT score. For the remaining subfamilies, we keep the adjPHACT score of

the representative amino acid of the target. Then, similar to [27] we check whether the same

amino acid is conserved across all subfamilies. On the other hand, our approach differentiates

from[27] in terms of considering multiple subfamilies and using adjPHACT scores which employ

phylogenetic trees and ancestral reconstruction probabilities. In our approach, we compute the

contribution of each subfamily to the SDP score by checking whether the representative amino

acid of target has a high adjPHACT score in that subfamily (line 1). In the final SDP score for

any position k is computed by considering the distance between target and other subfamilies

(which is computed by considering the distance between root nodes), the conservation level of

the target subfamily in terms of independent amino acid alterations and the individual score

coming from each subfamily (line 3).

ALGORITHM 3: SDP SCORE FOR POSITION "K"

Input: Amino acid with the highest adjPHACT score in the target group, aa; the adjPHACT score of aa for target, P_{aa}^T; adjPHACT score of aa for other subfamilies i=1,…,n, P_{aa}^i; distance between target subfamily and subfamily i, d_i.

1   *Compute score for each subfamily i,*

$$S_i = \exp(1) - \exp\left(P_{aa}^i\right).$$

2   *The overall weight,*

$$\omega = 1 - \max\left(P_{aa}^i\right).$$

3   *The SDP score for position k,*

$$SDP = P_{aa}^T + \omega \left(\sum_{i=1}^{n} \frac{1}{D_i} S_i\right)^{P_{aa}^T}$$

Evolution of Class C GPCRs

567   We selected representative sequences from different taxonomic levels for each subfamily and

568   264 CaSR-like sequences. We aligned them with MAFFT v7.221 einsi algorithm[63]. We built

569   the ML tree by RAxML-NG-0.9.0 with the model JTT and transfer bootstrap expectation –bs-

570 metric fbp, tbe parameters[67]. We merged the ML trees of CaSR, GPRC6A and taste receptors

571 by checking clades by using ETE toolkit [65].

Machine Learning

Dataset and Feature Preparation

To predict the consequence of a substitution in human CaSR, we used a gradient boosting-

based machine learning algorithm, XGBoost[33]. We used XGBoost library for R[73] to train our

model. We selected total of 337 loss and gain-of-function mutations from the literature[2] to train

our model. Since we used conservation scores as features to train our model, we divided

subfamily alignments and mutations randomly as 80% training and the remaining 20% test data

before creating feature matrices to prevent information leakage. 25% of the training data was

randomly picked as the validation data five times for cross validation. For each dataset split we

used the sklearn train test split model with stratify option to keep loss-of-function to gain-of-

function ratio almost the same in the datasets[74]. We calculated the conservation score of the

reference amino acid and the substituted amino acid in human CaSR in each subfamily. The

reference and the substituted amino acids were represented BLOSUM62 encoded matrices.

Amino acid physico-chemical feature values Zimmerman polarity[75], average flexibility[76],

Dayhoff[77], average buried area[78], Doolittle hydropathicity[79], atomic weight ratio[80],

molecular weight, and bulkiness[75] from ProtScale database[81]; and domain information of

the reference amino acid were used as other features. We normalized the physico-chemical

feature values prior to model training. We repeated the whole random dataset splitting and

feature preparation procedure 50 times to obtain more robust results.

Model Selection and Parameter Tuning

We picked the model parameters for each replication by applying a 5-fold cross-validation

technique on the training set. We tuned the model parameters step-by-step using the same

validation sets for each parameter to decrease the time complexity. We used the following order

of model parameters, so that the parameter that has the highest impact on model outcome was

tuned first: Eta and nrounds, gamma, maxdepth, subsample, colsample bytree, min child

weight, lambda, alpha. We selected the maxdepth as 2, the minimum maxdepth value to

prevent overfitting. We chose eta, gamma, colsample bytree, subsample, min child weight from

the sets 0.00001,0.00002,..., 0.001,0,0.1,0.2,...,0.5, 0.5,0.55,...,1, 0.5,0.55,...,1, 1,2,...,6

respectively. We selected regularization parameters lambda and alpha from the set 0, 1e-4, 1e-

3, 1e-2, 1e-1, 1, 10, 100. We set nrounds parameter as 200.

Performance Metrics

We used the area under the receiver operating characteristic curve (AUROC) and the area under the precision-recall curve (AUPR) to evaluate the performance of our prediction model. AUROC and AUPR are performance measures that are widely used to evaluate the performance of binary classification problems. The higher the AUROC and AUPR, the better the model distinguishes classes. To understand how our model makes predictions, we used SHAP (SHapley Additive exPlanations) values. SHAP values provide an estimate of the contribution of each feature to the prediction made by the model[82]. We calculated SHAP values for our final model trained by all samples by using R shapviz package[83].

Predictive Performance

After we evaluated the performance of our machine learning algorithm over 50 replications, we used the whole dataset to train the model that we used to make predictions for every possible mutation in human CaSR. We selected model parameters by using 5-fold cross-validation technique on the whole dataset. To create a new test dataset, we took subfamily alignments of the species from the new Uniprot dataset that did not exist in the training data. We eliminated amino acids that are observed in the CaSR alignment as neutral. In each position we predicted the gain or loss-of -function class for any substitution. We did a literature search to find new clinical cases that cause either gain or loss of function mutations. We reported our predictions in the table.

Acknowledgments

This study was supported by EMBO Installation Grant no:4163 funded by TÜBİTAK (to OA). Ogun Adebali is supported by the BAGEP program of the Science Academy - Türkiye, and the TÜBA-GEBİP program of the Turkish Academy of Sciences.

References

1. Cook, A.E., et al., *Biased allosteric modulation at the CaS receptor engendered by structurally diverse calcimimetics.* British journal of pharmacology, 2015. **172**(1): p. 185-200.

2. Gorvin, C.M., *Molecular and clinical insights from studies of calcium-sensing receptor mutations.* J Mol Endocrinol, 2019. **63**(2): p. R1-R16.

3. Chen, X., et al., *Structural insights into the activation of human calcium-sensing receptor.* Elife, 2021. **10**.

4. Geng, Y., et al., *Structural mechanism of ligand activation in human calcium-sensing receptor.* Elife, 2016. **5**.

5.   Ling, S., et al., *Structural mechanism of cooperative activation of the human calcium-*
sensing receptor by Ca(2+) ions and L-tryptophan. Cell Res, 2021. **31**(4): p. 383-394.

6.   Wootten, D., et al., *Mechanisms of signalling and biased agonism in G protein-coupled*
receptors. Nat Rev Mol Cell Biol, 2018. **19**(10): p. 638-653.

7.   Zhang, C., et al., *Structural basis for regulation of human calcium-sensing receptor by*
magnesium ions and an unexpected tryptophan derivative co-agonist. Sci Adv, 2016.
2(5): p. e1600241.

8.   Gao, Y., et al., *Asymmetric activation of the calcium-sensing receptor homodimer.*
Nature, 2021. **595**(7867): p. 455-459.

9.   Park, J., et al., *Symmetric activation and modulation of the human calcium-sensing*
receptor. Proc Natl Acad Sci U S A, 2021. **118**(51).

10.   Wen, T., et al., *Structural basis for activation and allosteric modulation of full-length*
calcium-sensing receptor. Science Advances, 2021. **7**(23): p. eabg1483.

11.   Mun, H.-C., et al., *A double mutation in the extracellular Ca2+-sensing receptor's venus*
flytrap domain that selectively disables L-amino acid sensing. Journal of Biological
Chemistry, 2005. **280**(32): p. 29067-29072.

12.   Zhang, C., et al., *Identification of an L-phenylalanine binding site enhancing the*
cooperative responses of the calcium-sensing receptor to calcium. Journal of Biological
Chemistry, 2014. **289**(8): p. 5296-5309.

13.   Zhang, Z., et al., *Three adjacent serines in the extracellular domains of the CaR are*
required for L-amino acid-mediated potentiation of receptor function. Journal of
Biological Chemistry, 2002. **277**(37): p. 33727-33735.

14.   Liu, H., et al., *Illuminating the allosteric modulation of the calcium-sensing receptor.*
Proceedings of the National Academy of Sciences, 2020. **117**(35): p. 21711-21722.

15.   Chagoyen, M., J.A. Garcia-Martin, and F. Pazos, *Practical analysis of specificity-*
determining residues in protein families. Brief Bioinform, 2016. **17**(2): p. 255-61.

16.   Studer, R.A., B.H. Dessailly, and C.A. Orengo, *Residue mutations and their impact on*
protein structure and function: detecting beneficial and pathogenic changes. Biochemical
journal, 2013. **449**(3): p. 581-594.

17.   Kuru, N., et al., *PHACT: Phylogeny-Aware Computing of Tolerance for Missense*
Mutations. Mol Biol Evol, 2022. **39**(6).

18.   Pin, J.P., T. Galvez, and L. Prezeau, *Evolution, structure, and activation mechanism of*
family 3/C G-protein-coupled receptors. Pharmacol Ther, 2003. **98**(3): p. 325-54.

19.   Harpsoe, K., et al., *Structural insight to mutation effects uncover a common allosteric*
site in class C GPCRs. Bioinformatics, 2017. **33**(8): p. 1116-1120.
20.   Patil, I., *Visualizations with statistical details: The'ggstatsplot'approach.* Journal of Open
Source Software, 2021. **6**(61): p. 3167.
21.   Goes van Naters, W. and C. Mucignat-Caretta, *Frontiers in Neuroscience Drosophila*
Pheromones: From Reception to Perception. Neurobiology of Chemical Communication.
Boca Raton (FL): CRC Press/Taylor & Francis (c), 2014.
22.   Brown, D., et al., *Subfamily hmms in functional genomics.* Pac Symp Biocomput, 2005:
p. 322-33.
23.   Srivastava, P.K., et al., *HMM-ModE--improved classification using profile hidden Markov*
models by optimising the discrimination threshold and modifying emission probabilities
with negative training sequences. BMC Bioinformatics, 2007. **8**: p. 104.
24.   Roberts, M.S., et al., *Treatment of Autosomal Dominant Hypocalcemia Type 1 With the*
Calcilytic NPSP795 (SHP635). J Bone Miner Res, 2019. **34**(9): p. 1609-1618.
25.   Pruitt, K.D., T. Tatusova, and D.R. Maglott, *NCBI reference sequences (RefSeq): a*
curated non-redundant sequence database of genomes, transcripts and proteins.
Nucleic Acids Res, 2007. **35**(Database issue): p. D61-5.
26.   Gorvin, C.M., *Calcium-sensing receptor signaling—How human disease informs biology.*
Current opinion in endocrine and metabolic research, 2021. **16**: p. 10-18.
27.   Bradley, D. and P. Beltrao, *Evolution of protein kinase substrate recognition at the active*
site. PLoS biology, 2019. **17**(6): p. e3000341.
28.   Chun, L., W.H. Zhang, and J.F. Liu, *Structure and ligand recognition of class C GPCRs.*
Acta Pharmacol Sin, 2012. **33**(3): p. 312-23.
29.   Pi, M., S.K. Nishimoto, and L.D. Quarles, *GPRC6A: Jack of all metabolism (or master of*
none). Mol Metab, 2017. **6**(2): p. 185-193.
30.   Nango, E., et al., *Taste substance binding elicits conformational change of taste receptor*
T1r heterodimer extracellular domains. Sci Rep, 2016. **6**: p. 25745.
31.   Kumari, R., C. Castillo, and A. Francesconi, *Agonist-dependent signaling by group I*
metabotropic glutamate receptors is regulated by association with lipid domains. J Biol
Chem, 2013. **288**(44): p. 32004-19.
32.   Nuemket, N., et al., *Structural basis for perception of diverse chemical substances by*
T1r taste receptors. Nat Commun, 2017. **8**: p. 15530.

33.   Chen, T. and C. Guestrin. *Xgboost: A scalable tree boosting system*. in *Proceedings of*
the 22nd acm sigkdd international conference on knowledge discovery and data mining.
2016.

34.   Chou, K.J., et al., *A new missense mutation of calcium sensing receptor with isoleucine*
replaced by serine at codon 857 leading to type V Bartter syndrome. Exp Cell Res,
2022. **414**(1): p. 113080.

35.   Moon, J.E., et al., *A cell function study on calcium regulation of a novel calcium-sensing*
receptor mutation (p.Tyr825Phe). Ann Pediatr Endocrinol Metab, 2021. **26**(1): p. 24-30.

36.   Palmieri, S., et al., *Case Report: Unusual Presentations of Loss-of-Function Mutations of*
the Calcium-Sensing Receptor. Front Med (Lausanne), 2021. **8**: p. 809067.

37.   Li, N., et al., *A novel homozygous mutation of the calcium-sensing receptor gene*
associated with apparent autosomal recessive inheritance of familial hypocalciuric
hypercalcemia. Chin Med J (Engl), 2021. **134**(15): p. 1869-1871.

38.   Hao, Y., et al., *Radiofrequency Ablation of Parathyroid Glands to Treat a Patient With*
Hypercalcemia Caused by a Novel Inactivating Mutation in CaSR. Front Endocrinol
(Lausanne), 2021. **12**: p. 743517.

39.   Tsuji, T., et al., *Autosomal Dominant Hypocalcemia With Atypical Urine Findings*
Accompanied by Novel CaSR Gene Mutation and VitD Deficiency. J Endocr Soc, 2021.
5(3): p. bvaa190.

40.   Dharmaraj, P., et al., *Neonatal Hypocalcemic Seizures in Offspring of a Mother With*
Familial Hypocalciuric Hypercalcemia Type 1 (FHH1). J Clin Endocrinol Metab, 2020.
105(5).

41.   Gomes, V., et al., *Autosomal dominant hypocalcaemia: identification of two novel*
variants of CASR gene. BMJ Case Rep, 2020. **13**(6).

42.   Magno, A.L., et al., *Functional Analysis of Calcium-Sensing Receptor Variants Identified*
in Families Provisionally Diagnosed with Familial Hypocalciuric Hypercalcaemia. Calcif
Tissue Int, 2020. **107**(3): p. 230-239.

43.   Majumdar, S.K., et al., *A Novel Variant in the Calcium-Sensing Receptor Associated with*
Familial Hypocalciuric Hypercalcemia and Low-to-Normal PTH. Case Rep Endocrinol,
2020. **2020**: p. 8752610.

44.   Zajickova, K., et al., *Familial hypocalciuric hypercalcemia in an index male: grey zones*
of the differential diagnosis from primary hyperparathyroidism in a 13-year clinical follow
up. Physiol Res, 2020. **69**(Suppl 2): p. S321-S328.

45.   Sagi, S.V., et al., *A novel CASR variant in a family with familial hypocalciuric*
734   *hypercalcaemia and primary hyperparathyroidism.* Endocrinol Diabetes Metab Case
735   Rep, 2020. **2020**.

736   46.   Dong, Q., et al., *[Clinical and genetic analysis of a child with neonatal severe*
737   *parathyroidism].* Zhonghua Yi Xue Yi Chuan Xue Za Zhi, 2020. **37**(11): p. 1247-1249.

738   47.   Hawkes, C.P., D.I. Shulman, and M.A. Levine, *Recombinant human parathyroid*
739   *hormone (1-84) is effective in CASR-associated hypoparathyroidism.* Eur J Endocrinol,
740   2020. **183**(6): p. K13-K21.

741   48.   Wejaphikul, K., et al., *Subtotal parathyroidectomy successfully controls calcium levels of*
742   *patients with neonatal severe hyperparathyroidism carrying a novel CASR mutation.*
743   Hormone Research in Paediatrics, 2023: p. 1-7.

744   49.   Courtney, A., et al., *Familial hypocalciuric hypercalcaemia type 1 caused by a novel*
745   *heterozygous missense variant in the CaSR gene, p (His41Arg): two case reports.* BMC
746   Endocrine Disorders, 2022. **22**(1): p. 324.

747   50.   Bletsis, P., et al., *A Novel missense CASR gene sequence variation resulting in familial*
748   *hypocalciuric hypercalcemia.* AACE Clinical Case Reports, 2022. **8**(5): p. 194-198.

749   51.   Wu, Y., et al., *Autosomal dominant hypocalcemia with a novel CASR mutation: a case*
750   *study and literature review.* Journal of International Medical Research, 2022. **50**(7): p.
751   03000605221110489.

752   52.   Coughlan, A., F. Khan, and M. Brassill, *A Novel Genetic Variant Resulting in Familial*
753   *Hypocalciuric Hypercalcaemia.* Irish Medical Journal, 2022. **115**(2): p. 545-545.

754   53.   Mullin, B.H., et al., *Functional assessment of calcium-sensing receptor variants confirms*
755   *familial hypocalciuric hypercalcemia.* Journal of the Endocrine Society, 2022. **6**(5): p.
756   bvac025.

757   54.   Jones, E.M., et al., *Structural and functional characterization of G protein–coupled*
758   *receptors with deep mutational scanning.* Elife, 2020. **9**: p. e54895.

759   55.   Alirezaie, N., et al., *ClinPred: prediction tool to identify disease-relevant nonsynonymous*
760   *single-nucleotide variants.* The American Journal of Human Genetics, 2018. **103**(4): p.
761   474-483.

762   56.   Ioannidis, N.M., et al., *REVEL: an ensemble method for predicting the pathogenicity of*
763   *rare missense variants.* The American Journal of Human Genetics, 2016. **99**(4): p. 877-
764   885.

765   57.   Rentzsch, P., et al., *CADD: predicting the deleteriousness of variants throughout the*
766   *human genome.* Nucleic acids research, 2019. **47**(D1): p. D886-D894.

58. Rogers, M.F., et al., *FATHMM-XF: accurate prediction of pathogenic point mutations via*
extended features. Bioinformatics, 2018. **34**(3): p. 511-513.

59. Eddy, S.R., *Accelerated profile HMM searches.* PLoS computational biology, 2011.
7(10): p. e1002195.

60. Finn, R.D., et al., *The Pfam protein families database: towards a more sustainable*
future. Nucleic acids research, 2016. **44**(D1): p. D279-D285.

61. *UniProt: the universal protein knowledgebase.* Nucleic acids research, 2017. **45**(D1): p.
D158-D169.

62. Altschul, S.F., et al., *Basic local alignment search tool.* Journal of molecular biology,
1990. **215**(3): p. 403-410.

63. Yamada, K.D., K. Tomii, and K. Katoh, *Application of the MAFFT sequence alignment*
program to large data—reexamination of the usefulness of chained guide trees.
Bioinformatics, 2016. **32**(21): p. 3246-3251.

64. Stamatakis, A., *RAxML version 8: a tool for phylogenetic analysis and post-analysis of*
large phylogenies. Bioinformatics, 2014. **30**(9): p. 1312-1313.

65. Huerta-Cepas, J., F. Serra, and P. Bork, *ETE 3: reconstruction, analysis, and*
visualization of phylogenomic data. Molecular biology and evolution, 2016. **33**(6): p.
1635-1638.

66. Capella-Gutierrez, S., J.M. Silla-Martinez, and T. Gabaldon, *trimAl: a tool for automated*
alignment trimming in large-scale phylogenetic analyses. Bioinformatics, 2009. **25**(15): p.
1972-3.

67. Kozlov, A.M., et al., *RAxML-NG: a fast, scalable and user-friendly tool for maximum*
likelihood phylogenetic inference. Bioinformatics, 2019. **35**(21): p. 4453-4455.

68. Nguyen, L.-T., et al., *IQ-TREE: a fast and effective stochastic algorithm for estimating*
maximum-likelihood phylogenies. Molecular biology and evolution, 2015. **32**(1): p. 268-
274.

69. Kalyaanamoorthy, S., et al., *ModelFinder: fast model selection for accurate phylogenetic*
estimates. Nature methods, 2017. **14**(6): p. 587-589.

70. Hoang, D.T., et al., *UFBoot2: improving the ultrafast bootstrap approximation.* Molecular
biology and evolution, 2018. **35**(2): p. 518-522.

71. Waterhouse, A., et al., *SWISS-MODEL: homology modelling of protein structures and*
complexes. Nucleic acids research, 2018. **46**(W1): p. W296-W303.

72. Pettersen, E.F., et al., *UCSF Chimera--a visualization system for exploratory research*
and analysis. J Comput Chem, 2004. **25**(13): p. 1605-12.

73. Chen, T., et al., *Xgboost: extreme gradient boosting. R package version 04-2.* OS
Independent, 2015.
74. Pedregosa, F., et al., *Scikit-learn: Machine learning in Python.* the Journal of machine
Learning research, 2011. **12**: p. 2825-2830.
75. Zimmerman, J., N. Eliezer, and R. Simha, *The characterization of amino acid sequences*
in proteins by statistical methods. Journal of theoretical biology, 1968. **21**(2): p. 170-201.
76. Bhaskaran, R. and P. Ponnuswamy, *Positional flexibilities of amino acid residues in*
globular proteins. International Journal of Peptide and Protein Research, 1988. **32**(4): p.
241-255.
77. Barker, W.C., L.K. Ketcham, and M.O. Dayhoff, *A comprehensive examination of protein*
sequences for evidence of internal gene duplication. Journal of Molecular Evolution,
1978. **10**: p. 265-281.
78. Rose, G.D., et al., *Hydrophobicity of amino acid residues in globular proteins.* Science,
1985. **229**(4716): p. 834-838.
79. Kyte, J. and R.F. Doolittle, *A simple method for displaying the hydropathic character of a*
protein. Journal of molecular biology, 1982. **157**(1): p. 105-132.
80. Grantham, R., *Amino acid difference formula to help explain protein evolution.* science,
1974. **185**(4154): p. 862-864.
81. Walker, J.M., *The proteomics protocols handbook.* 2005: Springer.
82. Lundberg, S.M. and S.-I. Lee, *A unified approach to interpreting model predictions.*
Advances in neural information processing systems, 2017. **30**.
83. Mayer, M. *shapviz: SHAP Visualizations. R package version 0.9.0.* 2023; Available from:
https://github.com/ModelOriented/shapviz.