Hepatic lipid-associated macrophages mediate the beneficial effects of bariatric surgery against MASH

Gavin Fredrickson[1], Kira Florczak[1], Fanta Barrow[1], Katrina Dietsche[1], Haiguang Wang[1], Preethy Parthiban[1], Rawan Almutlaq[1], Oyedele Adeyi[2], Adam Herman[3], Alessandro Bartolomucci[1], Christopher Staley[3], Cyrus Jahansouz[3], Jesse Williams[1,4], Douglas G. Mashek[5], Sayeed Ikramuddin[3*], and Xavier S. Revelo[1,4,6*]

[1]Department of Integrative Biology & Physiology, University of Minnesota, Minneapolis, MN 55455, USA.
[2]Department of Laboratory Medicine and Pathology, University of Minnesota, Minneapolis, MN 55455, USA.
[3]Department of Surgery, University of Minnesota, Minneapolis, MN 55455, USA.
[4]Center for Immunology, University of Minnesota, Minneapolis, MN 55455, USA.
[5]Department of Biochemistry, Molecular Biology, and Biophysics, University of Minnesota, Minneapolis, MN 55455, USA; Division of Diabetes, Endocrinology, and Metabolism, Department of Medicine, University of Minnesota, Minneapolis, MN, USA.
[6]Lead contact
*Correspondence: xrevelo@umn.edu or ikram001@umn.edu

Contact Information
Xavier S. Revelo, Ph.D.
Cancer and Cardiovascular Research Building
2231 6th St SE
Minneapolis, MN 55455
Phone: +1 (612) 301-7688

ABSTRACT

For patients with obesity and metabolic syndrome, bariatric procedures such as vertical sleeve gastrectomy (VSG) have a clear benefit in ameliorating metabolic dysfunction-associated steatohepatitis (MASH). While the effects of bariatric surgeries have been mainly attributed to nutrient restriction and malabsorption, whether immuno-modulatory mechanisms are involved remains unclear. Here we report that VSG ameliorates MASH progression in a weight loss-independent manner. Single-cell RNA sequencing revealed that hepatic lipid-associated macrophages (LAMs) expressing the triggering receptor expressed on myeloid cells 2 (TREM2) increase their lysosomal activity and repress inflammation in response to VSG. Remarkably, TREM2 deficiency in mice ablates the reparative effects of VSG, suggesting that TREM2 is required for MASH resolution. Mechanistically, TREM2 prevents the inflammatory activation of macrophages and is required for their efferocytotic function. Overall, our findings indicate that bariatric surgery improves MASH through a reparative process driven by hepatic LAMs, providing insights into the mechanisms of disease reversal that may result in new therapies and improved surgical interventions.

MAIN

Metabolic dysfunction-associated steatotic liver disease (MASLD) is estimated to affect 30% of the population and is one of the leading causes of abnormal liver function[1]. MASLD is highly associated with obesity and covers a wide spectrum of liver pathology ranging from simple lipid accumulation to the more serious metabolic dysfunction-associated steatohepatitis (MASH), characterized by inflammation, hepatocellular injury, and fibrosis[2, 3]. Hepatic inflammation is primarily driven by liver macrophages and is a critical component in the initiation and progression of MASH[3, 4]. Although lifestyle and dietary intervention can result in moderate weight loss with concomitant improvements in comorbidities, these effects are transient, and weight regain is common[5]. Bariatric procedures such as vertical sleeve gastrectomy (VSG) are the most successful and effective treatment options for obese adults[6] and are effective at ameliorating the progression of MASH[7, 8, 9]. While changes in body weight have a profound metabolic impact, bariatric surgery often results in improved insulin sensitivity before any substantial weight loss, indicative of weight-loss-independent effects[10, 11]. Several mechanisms of weight-loss-independent improvements in metabolism due to bariatric surgery have been proposed including bile acid signaling[12], gut hormones[13], intestinal tissue reprogramming[14], changes in the gut microbiome[15, 16], and HIF2α-mediated iron absorption[17]. However, whether bariatric surgery ameliorates the progression of MASH through immunomodulatory mechanisms has not been investigated.

Recent research has shown that MASH is associated with the emergence of macrophages with a lipid-associated signature, termed hepatic lipid-associated macrophages (LAMs), that express the triggering receptor expressed on myeloid cells-2 (TREM2)[18, 19]. Hepatic LAMs are conserved in mice and humans, correlate with disease severity, preferentially localize to steatotic regions, and are highly responsive to dietary intervention[19, 20]. In MASH, TREM2 expression in LAMs is required for effective efferocytosis of apoptotic hepatocytes[21]. However, prolonged hypernutrition and hepatic inflammation promote the shedding of TREM2 from the cell surface of LAMs leading to an ineffective clearance of dying hepatocytes[21]. Cleavage of TREM2 increases soluble TREM2 (sTREM2) in the circulation, making it a promising biomarker of MASH severity[22]. TREM2 is also required for adequate metabolic coordination between macrophages and hepatocytes, lipid handling, and extracellular matrix remodeling[21, 22, 23]. In a mouse model of hepatotoxic injury, TREM2 deficiency exacerbated inflammation-associated injury through enhanced toll-like receptor signaling, in agreement with its anti-inflammatory function[24]. Given the key roles of TREM2 in the diseased liver, mounting evidence has shown that TREM2 deficiency worsens most aspects of MASLD and MASH[21, 22, 23, 24, 25]. As TREM2 is a central signaling hub regulating macrophage function, therapeutic efforts have been made to stimulate the TREM2 active domain or block its shedding in neurodegenerative disease[26]. Despite the evidence suggesting that LAMs are protective, the mechanisms promoting their restorative function in MASH are poorly understood.

In the current study, we found that VSG ameliorates the progression of MASH and results in profound effects on the macrophage transcriptomic profile in a weight loss-independent manner. Remarkably, the newly described hepatic lipid-associated macrophages (LAMs) upregulate their reparative, lipid-quenching, and anti-inflammatory programs in response to bariatric surgery. Most notably, TREM2 deficiency prevented VSG-induced MASH reversal, suggesting that TREM2 is essential for the beneficial effects of bariatric surgery in the liver. Mechanistically, TREM2 prevents the production of inflammatory cytokines by macrophages and is required for adequate efferocytosis of apoptotic hepatocytes. Overall, our results indicate that bariatric surgery improves MASH through a reparative process driven by hepatic LAMs.

RESULTS

Vertical sleeve gastrectomy ameliorates MASH progression independent of weight loss

To study the mechanisms by which bariatric surgery ameliorates MASH, we used a validated mouse model of VSG in which approximately 80% of the lateral stomach of mice is clamped by a gastric clip and excised[15]. To induce MASH, we fed mice a high-fat high-carbohydrate (HFHC) diet that recapitulates aspects of human disease such as obesity, hepatic lipid accumulation, inflammation, and fibrosis[27]. Mice were fed the HFHC diet ad libitum for 12 weeks and then assigned to either sham (Sham AL) or VSG surgery (**Fig. 1A**) and remained on the HFHC diet for 5 weeks. Most mice survived the surgery, and we confirmed surgical anatomy by oral gavage of barium and imaging[15]. To determine weight-loss-independent effects, we included a sham group that was pair-fed to the VSG group (Sham PF) to match their caloric intake during the post-surgery period[28]. In addition, a group of mice were fed a normal chow diet (**NCD**) for the duration of the studies. Compared with Sham AL, both VSG and Sham PF mice resulted in a similar decrease in average daily food intake (**Fig. 1B**). Following the surgeries, body weight rapidly decreased in Sham AL, PF, and VSG groups due to the intervention, as previously reported[29] (**Fig. 1C**). However, while the body weight in Sham AL mice recovered, Sham PF and VSG mice maintained their weight loss throughout the study (**Fig. 1C**). Five weeks after the surgeries, both Sham PF and VSG mice showed a similar decrease in liver weight (**Fig. 1D**). Compared with Sham AL controls, only the VSG group showed a reduction in hepatic lipid accumulation (**Fig. 1E and 1F)**, ALT/AST (**Fig. 1G**), fibrosis (**Fig. 1H and Extended Data Fig. 1B**), and NAS score (**Extended Data Fig. 1A**), suggesting that the effects of VSG are partly independent of the reduced caloric intake induced by the surgery. As decreased intestinal lipid absorption has been proposed as a mechanism of VSG actions[30], we measured fecal lipids and found that VSG mice had increased fecal lipids compared with Sham AL mice (**Fig. 1I**). We also determined the effects of VSG on MASH progression 10 weeks after surgeries and found that the VSG-induced reductions in body and liver weight, liver triglycerides, AST, and ALT were maintained at this later timepoint (**Extended Data Fig. 1C-G**). Overall, these findings indicate that VSG results in substantial improvements in MASH in a weight-loss-independent manner.

To determine if the VSG-induced improvements in MASH were associated with alterations in the gut microbiota and its metabolites, we analyzed the fecal microbial composition of NCD, Sham AL, Sham PF, and VSG mice 5 weeks after surgeries. Compared with NCD mice, HFHC feeding induced dramatic remodeling in gut microbial species, including increased abundances of *Akkermansia*, *Erysipelotrichaceae*, *Parasutterella*, *Ruminococceae*, *Clostridium XVIII*, and decreased abundances of *Lactobacillus*, and *Alistipes* (**Fig. 1K and Supplemental Table 1**). While HFHC feeding caused a substantial loss of *Lactobacillus*, regardless of surgical treatment, VSG resulted in a restoration of this genus towards levels found in NCD mice (**Fig. 1J**). In contrast, the HFHC diet increased the abundance of *Clostridium XVIII* whereas it was partially lowered following VSG (**Fig. 1L**). No differences in fecal microbiota composition between Sham AL and Sham PF mice were detected 5 weeks after the interventions (**Supplemental Table 1**). Intestinal products can translocate into the portal vein and reach the liver where they can regulate immune homeostasis and inhibit inflammation[31]. Thus, we measured the concentrations of bile acids (BA) and short-chain fatty acids (SCFA) in the portal vein blood. While HFHC feeding increased the total amount of BA, as observed in Sham AL mice, both Sham PF and VSG mice had decreased BA concentrations to levels similar to NCD controls (**Fig. 1M**). In particular, VSG and Sham PF mice had a similar decrease in taurine-conjugated cholic acid while VSG was more effective at reducing deoxycholic acid and taurodeoxycholic acid (**Fig. 1N**). Independent of surgical intervention, HFHC feeding decreased the levels of cholic acid, chenodeoxycholic acid, and hyodeoxycholic acid and increased several taurine-conjugated BA in all groups without any

145 effects of Sham PF or VSG (**Extended Data Fig. 1H**). HFHC feeding also decreased the
146 concentration of SCFAs in hepatic portal serum and neither VSG nor Sham PF had an impact on
147 their concentration (**Fig. 1O and Supplemental Table 2**). Overall, these data suggest that the
148 VSG-induced improvements in MASH are associated with a partial restoration of specific gut
149 microbial species and BAs.
150
151 **scRNA-seq reveals profound effects of VSG on hepatic LAMs**
152
To explore the impact of VSG on hepatic macrophages, we first determined the abundance of
macrophages and Kupffer cells (KCs) subsets by flow cytometry[32]. We did not detect any
differences in the number of monocyte-derived macrophages (MoMF), embryonic KCs (emKC),
monocyte-derived KCs (moKC), and VSIG4[-] macrophages between Sham and VSG groups at 5
or 10 weeks after surgeries (**Extended Data Fig. 2A**). To determine how VSG influences the
function of hepatic macrophages, we profiled the gene expression of macrophages from Sham
AL, Sham PF, and VSG mice using droplet-based single-cell RNA sequencing (scRNA-seq, **Fig.
2A**). Livers were perfused before immune cell isolation to remove circulatory cells[33]. We
multiplexed samples using cell multiplexing oligos (CMOs) to track the sample of origin. Cells
were loaded into the ports of a 10x-Genomics chip following a single-cell 3' kit. The gene
expression and CMO libraries were sequenced using a Novaseq S4 chip (2x150bp PE). After
quality control, data were normalized and de-multiplexed. Monocytes and macrophages were
identified using the cell ID function of Seurat and re-clustered for analysis. We profiled the gene
expression of 58,904 single cells (avg. read depth 50,000, **Fig. 2B**). Unsupervised graph-based
clustering was performed on the integrated dataset and cells were visualized using uniform
manifold approximation and projection (UMAP). Independent of surgical groups, integrated UMAP
analysis revealed 18 clusters of monocytes and macrophages, whose identity was determined
based on the expression of established marker genes[18, 34] (**Fig. 2C, Extended Data Fig. 2B, and
Supplemental Table 3**).
172
Independent of surgical intervention, clusters 0, 2, 5, 6, 7, 8, 13, and 16 were identified as
monocyte-like cells while clusters 1, 3, 4, 9, 10, 11, 12, and 14 were macrophages (**Fig. 2C**).
Monocyte clusters had heterogeneous gene expression profiles indicative of their progressive
stages of maturation and function. Clusters 0, 2, and 7 had dual monocyte and macrophage
features, such as *Ly6c2* and *H2-Ab1*, suggesting that they were transitioning monocytes on the
trajectory of becoming macrophages. Cells in cluster 6 were identified as classical monocytes
based on their high expression of *Ly6c2* and *Ccr2,* which allows them to migrate in response to
inflammation. Monocytes in cluster 8 were enriched in *Spn* and did not express *Ccr2* and *Ly6c2*
indicating these cells were non-classical monocytes capable of patrolling. Clusters 13 and 16
corresponded to unidentifiable monocyte populations that expressed high levels of *Il2rb* and
Cxcr2, respectively (**Fig. 2C**). We next analyzed the macrophage subsets which can broadly be
divided into Kupffer cells (KCs) and MoMFs. Clusters 3, 9, and 10 were identified as KCs (KC1-
3) based on their expression of *Clec1b*, *Clec4f*, *Vsig4*, and *Folr2*. These clusters of KCs had
minimal expression of *Timd4* suggesting that they are primarily moKCs[32]. Clusters 3 and 9 had a
similar gene expression pattern, although cluster 3 was enriched in genes associated with
efferocytosis such as *Mertk* and *Wdfy3*. Cluster 10 was enriched in *Esam* and *Cd36* and
resembles a recently identified subset of pathogenic KCs termed "KC2"[35]. Among MoMFs, cells
in cluster 1 were enriched for LAM genes including *Trem2*, *Spp1*, *Lipa*, and *Cd36*[36] (**Fig. 2C, 2D
and Supplemental Table 3**). Pathway analysis showed that LAMs have enriched gene programs
associated with lipid metabolism such as "Lipoprotein particle binding" and "Lipase activity" (**Fig.
2E**). Cluster 4 was composed of MoMFs and transitioning monocytes that could not be further
characterized based on their differential gene expression. Cells in cluster 14 were enriched in
hemoglobin genes *Hba-a1* and *Hba-a2* which are highly expressed by erythrophages. Cluster 12

had a high G2M score and was enriched with *Top2a*, typical of proliferating macrophages (**Fig.**
2C, 2D and Supplemental Table 3). To gain insight into the differentiation of macrophage
subsets, we performed a slingshot trajectory analysis[37] of our integrated dataset. Although the
origin (ontogeny) and cellular turnover of hepatic LAMs is unknown, recent work suggests that
recruited MoMFs give rise to hepatic LAMs[22, 38]. Using monocytes as the origin, we found three
primary trajectories by which newly recruited monocytes differentiate into transitioning monocytes
and then either differentiate into LAMs, KC1s, or Trans Mon2s (**Fig. 2F and Extended Data Fig.**
2C). These data suggest that LAMs are derived from classical monocytes and are consistent with
recent studies that have used single cell transcriptomics to reveal a previously unappreciated
heterogeneity in hepatic macrophages in the MASH liver[18, 39].
206
Following cluster identification and trajectory analysis, we demultiplexed the samples based on
their experimental group and determined the relative abundance of each cluster. There were no
differences in the proportion and number of the major myeloid clusters 5 weeks post-VSG (**Fig.**
3A). To investigate the effects of VSG on the transcriptome of hepatic macrophages, we
performed differential gene expression analysis between Sham AL, Sham PF, and VSG groups
and found distinct expression patterns in macrophage clusters analyzed in bulk (**Extended Data**
Fig. 3A) and per cluster (**Supplemental Table 4**). Given the reparative functions of LAMs in
MASH, we focused our subsequent analysis on these cells. First, we quantified the average
single-cell expression of the *Trem2* gene in the LAM cluster and found no differences between
groups (**Fig. 3B**). Despite no effect on the abundance of LAMs and *Trem2* gene expression, VSG
mice had a decreased serum level of sTREM2 (**Fig. 3C**), suggesting a reduced cleavage of
membrane-bound TREM2 in macrophages due to improved inflammation[21]. Next, we performed
differential gene expression analysis between LAMs from VSG, Sham AL, and Sham PF mice.
Compared with LAMs from Sham AL (37 DEGs), and to a lesser extent Sham PF (24 DEGs),
LAMs from VSG mice showed an increased expression of genes involved in lysosomal activity
(*Lyz2*, *Ctsl, Ctss,* and *H2-Eb1*), antigen presentation (*H2-Eb1*), repression of inflammation (*Egr1*),
and fatty acid metabolism (*Lipa*) (**Supplemental Table 4** and **Fig. 3D**). In contrast, several genes
associated with inflammation (*Cd83*, *Nfkb1*, *Junb*, and *Mmp7*) were downregulated in LAMs from
VSG mice (**Fig. 3D**). Pathway analysis revealed an upregulation of pathways associated with
immune activation and lysosomal activity such as "Chemokine signaling", "Cytokine receptor
interaction", and "Phagosome" in LAMs from VSG mice (**Fig. 3E**). Similarly, gene set enrichment
analysis showed that genes involved in "Chemokine signaling", "Lysosome", "Peroxisome" and
"Fatty acid metabolism" had increased expression in VSG LAMs, compared with Sham PF
controls (**Fig. 3G**). Overall, these data highlight lysosomal and metabolic regulatory mechanisms
by which VSG may sustain the protective function of LAMs against MASH in response to VSG.
232
233 **Hepatic TREM2⁺ LAMs mediate the reparative effects of VSG against MASH**
234
Recent work has shown that systemic TREM2 deficiency worsens diet-induced MASH as TREM2
is required for LAM survival, the metabolic coordination between LAMs and hepatocytes, and the
clearance of dying liver cells [21, 22, 23, 40]. To determine if LAMs directly mediate the reparative
process induced by VSG against MASH, we performed sham or VSG surgeries on WT and
TREM2-deficient (TREM2 KO) mice fed an HFHC diet for 12 weeks. Expression of *Trem2* was
not detectable in bone marrow-derived macrophages (BMDM) from TREM2 KO mice (**Extended**
Data Fig. 4A). Five weeks post-surgery, WT and TREM2 KO mice showed a similar degree of
weight loss (**Fig. 4A**) and an increase in fecal lipid excretion (**Fig. 4B**) in response to VSG. In
agreement with our previous experiments, we found that sTREM2 was reduced following VSG in
the serum of WT mice but was not detectable in TREM2 KO mice (**Extended Data Fig. 4B**).

Notably, while VSG ameliorated MASH progression in WT mice, this effect was blunted in TREM2
KO mice (**Fig. 4C-G**). Compared with their Sham controls, VSG failed to decrease the liver weight
(**Fig. 4C**), hepatic steatosis (**Fig. 4D and 4E**), ALT (**Fig. 4F**), and AST (**Fig. 4G**) in TREM2 KO
mice, suggesting that TREM2 is required for the VSG-induced reversal of MASH. To explore the
underlying mechanisms, we first assessed whether TREM2 deficiency alters the hepatic
macrophage populations in mice with MASH before and after VSG. We found no differences in
the number of MoMFs, emKCs, moKCs, and VSIG4- macrophages between WT and TREM2 KO
fed the HFHC diet for 12 weeks without any intervention (**Extended Data Fig. 4C**). Similarly, the
blunted effect of VSG in TREM2 KO mice was not associated with alterations in the number of
hepatic macrophage subsets (**Fig. 4H and Extended Data Fig. 4D**). To determine the potential
mechanisms by which TREM2 KO mice are resistant to the beneficial effects of VSG, we
magnetically sorted and performed bulk RNA sequencing on total macrophages from the livers of
WT and TREM2 KO mice after sham or VSG surgeries. Unsupervised PCA showed a distinct
separation between WT Sham and WT VSG, whereas there was no distinction between TREM2
KO Sham and TREM2 KO VSG macrophages (**Fig. 4I**). Differential gene expression analysis
revealed a more robust response of WT macrophages to VSG (331 DEGs), compared with that
of TREM2 KO cells (44 DEGs) (**Fig. 4J and Extended Data Table 5**). In agreement, unsupervised
clustering of the top 500 most variable genes revealed substantial gene expression differences
and clustering between WT Sham and VSG macrophages but not between TREM2 KO Sham
and VSG cells (**Extended Data Fig. 4E**). We performed pathway and gene ontology (GO)
analyses and found that TREM2 KO macrophages from VSG mice upregulated genes enriched
in inflammatory and immune activation pathways such as "cytokine-cytokine receptor interaction"
and the GO terms "inflammatory response" and "chemotaxis" (**Fig. 4K**), suggesting that TREM2
is required for preventing an inflammatory activation of macrophages. To test this possibility, we
examined the response of BMDMs from WT and TREM2 KO mice with or without stimulation with
palmitate (PA) in vitro to mimic the lipid-rich environment of the MASH liver. Compared with WT
controls, TREM2 KO BMDMs showed a markedly increased expression of *Il1b*, *Il6*, and *Tnf*,
regardless of stimulation (**Fig. 4L**). Next, to determine whether TREM2 facilitates the ability of
macrophages to clear apoptotic cells, we induced apoptosis in AML12 hepatocytes by PA
treatment (**Extended Data Fig. 4F**) and co-cultured them with either WT or TREM2 KO peritoneal
macrophages. Consistent with its key role in efferocytosis, TREM2 was required for macrophages
to perform effective efferocytosis of apoptotic hepatocytes (**Fig. 4M**). Together, these data
suggest that hepatic LAMs mediate the VSG-induced reversal of MASH by repressing
inflammation and facilitating efferocytosis in a TREM2-dependent manner.
279
280 **VSG increases the content of inflammatory lipid species in hepatic macrophages**
281
Our scRNA-seq data shows that hepatic LAMs are uniquely equipped with the enzymatic
machinery to recognize, scavenge, and catabolize lipids[20, 38]. Because our data showed that total
liver TGs decreased in response to VSG while hepatic LAMs upregulate lipid metabolism genes,
we performed metabolic profiling of sorted F4/80[+] macrophages from Sham AL and VSG livers to
determine how bariatric surgery alters their lipid content. We performed metabolite profiling by
liquid chromatography-tandem mass spectrometry (MxP® Quant 500 kit, Biocrates), which
revealed 317 unique detectable metabolites in macrophages. The majority of these were lipid
species, predominantly TGs and phosphatidylcholines, although we also detected amino acids
and bile acids (**Fig. 5A**). Unsupervised principal component analysis (PCA) of detected
metabolites showed a moderate clustering of VSG samples with more separation among Sham
AL specimens (**Fig. 5B**). We found no difference in the concentration of total TGs in the
mcarophages from Sham AL and VSG mice (**Fig. 5C**). However, when we performed chain length
enrichment analysis of all lipid species, we found that macrophages from Sham AL mice were
enriched in species with longer chain lengths while those from VSG mice were enriched in species

of shorter chain length (**Fig. 5D**). Quantification of the major lipid families showed that macrophages from VSG mice had increased total phosphatidylcholines and sphingolipids, but no changes in total, cholesterol esters, fatty acids, glycosylceramides, ceramides, sphingolipids, and diacylglycerols (**Fig. 5E**). To further assess the effects of VSG on the lipid profile of hepatic macrophages, we assessed the composition of the individual lipid species. VSG macrophages had increased levels of phosphatidylcholines particularly species containing 2 acyl-bound (aa), one acyl- and one alkyl-bound (ae), and monounsaturated fatty acids (**Fig. 5F and Extended Data Fig. 5**). The role of phosphatidylcholines in macrophages is unclear with studies reporting both pro-inflammatory[41] and anti-inflammatory[42] responses. VSG macrophages were also enriched in sphingolipids containing long, very long-chain fatty acids, and hydroxyl-free species (**Fig. 5G**), as well as ceramides rich in very long-chain fatty acids (**Fig. 5H**). Given that sphingolipids acylated with fatty acids give rise to ceramides[43], and that excess ceramides have detrimental effects on the liver including steatosis[44], insulin resistance[45], inflammation, and oxidative stress[46], it is possible that VSG macrophages protect the liver from their detrimental effects. Although there were trending increases in VSG macrophages, we were unable to detect significant differences in the content of mono- and poly-unsaturated fatty acids (**Fig. 5I**) and cholesterol esters (**Fig. 5J**), compared with Sham controls. As the liver has decreased steatosis after VSG, the increase of macrophage intracellular lipids suggests an improved ability to clear lipids after the surgery.

Spatial transcriptomic of MASH livers following bariatric surgery reveal an improved metabolic status in the macrophage microenvironment

Given that our scRNA-seq analysis revealed substantial changes in the gene expression profile of hepatic macrophages, we explored the effects of VSG on the hepatic areas surrounding these cells using spatial transcriptomic analysis of liver sections (Nanostring GeoMx). Tissues from NCD, Sham AL, Sham PF, and VSG mice were collected 5 weeks after surgery and stained with a fluorescently labeled antibody against the macrophage marker CD68 to capture gene expression changes in CD68⁻ microanatomic areas (**Fig. 6A**). Following imaging and sequencing, the initial data was subjected to quality checks, filtering, and scaling, leading to a total of 7074 detectable genes and 32-36 regions of interest (ROI) per group (**Fig. 6B**) Unsupervised PCA of the normalized genes showed a substantial separation between CD68-expressing ROIs, and to a lesser extent between experimental groups (**Fig. 6C**). Due to the small size of their ROIs, we were unable to detect meaningful gene expression data from the CD68⁺ macrophage areas. However, differential expression gene analysis of the CD68⁻ ROIs revealed 185, 103, and 139 DEGs (FDR<0.05, FC > 1.5; **Supplemental File 6**) between NCD vs. Sham AL (**Fig. 6D**), Sham AL vs. and VSG (**Fig. 6E**), and Sham PF vs. VSG (**Fig. 6F**), respectively. To better understand the biological meaning of these changes, we performed enrichment and pathway analyses. Gene set enrichment analysis revealed that our analysis covered a wide range of cellular pathways including "metabolism", "immune system", "signal transduction" and "metabolism of proteins" (**Extended Data Fig. 6A**). Compared with the NCD group, pathway analysis showed that Sham AL mice had downregulated genes involved in the "metabolism of steroids" and "respiratory electron transport" and upregulation of "plasma lipoprotein remodeling" and "chemokine receptors bind chemokines" (**Fig. 6G**). Compared with Sham AL, VSG downregulated pathways involved in "biological oxidations", "metabolism of amino acids", and "metabolism of lipids" in agreement with the reduced lipid accumulation observed in VSG mice. On the other hand, VSG upregulated pathways such as "degranulation of neutrophils/platelets" and "complement cascade" indicative of an active immune and tissue remodeling response (**Fig. 6H**). Compared with the Sham PF group, we observed a downregulation of metabolic pathways and an upregulation of complement pathways in the VSG ROIs, similar to the VSG-induced changes relative to the Sham AL group (**Fig. 6I**). Overall, these findings highlight the VSG-induced transcriptomic changes in metabolic

347 and immune pathways in the microenvironment surrounding hepatic macrophages that are
348 associated with MASH reversal.
349
Finally, we determined whether our main findings could be relevant to human patients with MASH
undergoing VSG. We performed spatial transcriptomics on needle biopsy specimens from a
patient with MASH, collected before and one year after VSG (**Extended Data Fig. 6B**) as VSG
resulted in the resolution of MASH (NAS 4 to NAS 0 after 12 months, **Table 1**). We also included
a liver specimen from a donor patient with MASH without any surgical intervention. After staining
with cytokeratin (green) and the macrophage marker CD68 (magenta), 32 ROIs were annotated
to define hepatic zones I, II, and III (**Extended Data Fig. 2C**). Unsupervised PCA of detectable
genes revealed a substantial separation between specimens but small differences between
hepatic zones (**Extended Data Fig. 2D**). Pathway analysis of these comparisons revealed several
downregulated pathways involved in the "metabolism of lipids", "cholesterol biosynthesis", and
"metabolism of steroids" (**Extended Data Fig. 2E**), consistent with the MASH resolution observed
in this patient. To explore if human LAMs are responsive to VSG as in our mouse studies, we
assessed the expression of genes involved in the LAM differentiation program including *Trem2*,
Plin2, *Ctss*, *Cd36*, and *Lipa*[38, 40]. LAM genes were primarily enriched in zone I of the liver
(**Extended Data Fig. 2F**), in agreement with a recently published spatial transcriptomic study[20].
Furthermore, the LAM genes were upregulated in the Pre-VSG and MASH samples but were
lower in the Post-VSG sample (**Extended Data Fig. 2F**). These data suggest that LAMs correlate
with disease progression in human MASH.
368
369 **DISCUSSION**
370
Bariatric surgeries are the most effective options available for the treatment of obesity
complications including MASLD and MASH[7, 8, 9, 47]. Indeed, a recent randomized trial showed that
bariatric surgery is more effective than lifestyle intervention and medical care in the treatment of
patients with MASH, including those with liver fibrosis[48]. The beneficial effects of bariatric
surgeries against obesity-related diseases are usually attributed to the mechanical restriction of
the stomach and malabsorption of nutrients. However, accumulating evidence shows that the
surgeries exert profound effects in the regulation of several physiological functions ranging from
caloric intake[49], respiration[50], intestinal bile acids and lipid absorption[30], iron metabolism[17], and
immune responses[51]. Here we show that VSG induces weight-loss-independent improvements in
MASH in a process that requires TREM2-expressing macrophages. These data implicate the
innate immune system in the control of inflammation as an important mechanism of VSG action.
Previous work has shown that systemic inflammatory mediators associated with obesity gradually
decline following bariatric surgery in humans[52] and mouse models of disease[53, 54]. As TREM2-
deficient mice do not respond to the surgery, our findings further demonstrate that the reduced
inflammation in the liver is not secondary to caloric restriction or weight loss but a contributing
factor.
387
Recent research has shown that MASH is associated with the emergence of LAMs as a
specialized subset of macrophages that are required for adequate metabolic coordination, lipid
handling, extracellular matrix remodeling, and clearance of apoptotic hepatocytes[21, 22, 23, 40, 55].
Notably, metabolic stress and inflammation cause TREM2 shedding in LAMs resulting in the loss
of immune homeostasis and aggravated disease[21]. Here we show that TREM2 is necessary for
the VSG-induced reversal of disease in a process associated with decreased shedding of TREM2.
Mechanistically, we show that TREM2 is required for dampening hepatic inflammation and
effective efferocytosis of lipid-laden hepatocytes by macrophages. Furthermore, these two
processes are likely dependent on each other as efferocytosis has been shown to suppress
inflammation via the liver X receptor[56]. Given that TREM2 is a master regulator of the LAM

phenotype[36], we reasoned that TREM2 would mediate their reparative role induced by VSG.
Indeed, our findings indicate that VSG restores several TREM2-dependent pathways such as
"AKT-PI3K signaling", required for effective suppression of inflammation[57], and "oxidative
phosphorylation", which is typical of macrophages in anti-inflammatory or reparative states[58].
Overall, our data show that VSG induces a restorative function in macrophages in a TREM2-
dependent manner.
404
Given that the local micro-environment has a profound impact on the transcriptional program and
function of macrophages and neighboring cells[58], we used spatial transcriptomics to assess the
effects of VSG on hepatic macrophage-adjacent areas. In agreement with reduced steatosis, lipid
metabolism pathways were suppressed by VSG in the non-macrophage areas. However, one of
the main immune pathways that was upregulated following VSG was the complement system.
This was surprising as increased levels of complement proteins have been associated with MASH
severity in humans[59] and complement has been shown to increase hepatic de novo lipogenesis,
inflammation, and insulin resistance[60]. Additional studies are needed to determine if complement
proteins play a role in the response of the MASH liver to VSG due to the limited biological
implications that can be drawn from transcriptomic data. Nevertheless, hepatic complement
proteins have been shown to promote a reparative or apoptotic cell-clearing phenotype[61]. In
support of this notion, we found that macrophages from VSG mice have an enrichment of several
lipid species at the same time as the liver presents with decreased steatosis and a downregulation
of genes associated with lipid metabolism.
419
Not surprisingly, bariatric surgery induces rapid and profound changes in the gut microbiota likely
due to anatomical and metabolic mechanisms[62, 63]. Increased *Lactobacillus*, considered a
beneficial commensal bacteria[64, 65], has been proposed to be a mechanism by which VSG
improves obesity and glucose tolerance[66]. As the potential ligands of TREM2 include bacterial
products[67], it is possible that gut-derived microbial products may have direct effects on hepatic
LAMs or other macrophage subsets through pattern recognition receptors[68], or gut-derived
metabolites drained via the portal vein[69]. For example, increased intestinal butyrate may be an
additional mechanism by which VSG induces a reparative macrophage phenotype as VSG
restores the levels of butyrate-producing *Lactobacillus*[70].
429
One limitation of our study is that we utilized a mouse with a systemic TREM2 deficiency. Although
TREM2 is primarily expressed by myeloid cells, bacterial and viral infections have been reported
to induce TREM2 expression in CD4 T cells[71, 72]. Additionally, TREM2-expressing macrophages
in the adipose tissue have been shown to prevent obesity and adiposity, which could influence
the development of hepatic steatosis[36]. While several key experiments such as our scRNAseq
and spatial transcriptomics analyses show that hepatic LAMs are responsive to VSG, we cannot
rule out the potential contribution of adipose tissue LAMs to the VSG-induced protective effects
in the liver.
438
In summary, here we show that VSG induces a reversal of MASH independent of weight loss
accompanied by a substantial remodeling of the gut microbiota and hepatic macrophages. Here,
we report that hepatic TREM2[+] LAMs respond to VSG by increasing their expression of lysosomal
and oxidative phosphorylation genes while downregulating inflammatory genes. Notably, TREM2
deficiency ablates the protective effects of VSG, providing causal evidence that LAMs mediate
the beneficial effects of bariatric surgery against MASH.
445
MATERIALS AND METHODS
Animals

Five-week-old C57BL/6J (000664) and TREM2-deficient (TREM2 KO, 027197) male mice were
purchased from The Jackson Laboratory and maintained in a pathogen-free, temperature-
controlled environment. At six weeks of age, mice were fed a high-fat high-carbohydrate (HFHC,
40% kcal palm oil, 20% kcal fructose, and 2% cholesterol supplemented with 23.1 g/L d-fructose
and 18.9 g/L d-glucose in the drinking water) diet ad-libitum for 12 weeks to induce MASH.
Subsequently, they received either a sham operation or VSG surgery. Post-surgery, the mice
were provided a liquid diet for 2 days and the HFHC diet was slowly re-introduced. 7 days post-
surgery the mice were only fed the HFHC diet for the remainder of the study. To account for
weight-loss-independent effects, there was a sham ad-libitum-fed group (Sham AL) and a sham
group that was pair-fed (Sham PF) to the VSG group to match the weight loss post-surgery. The
amount of HFHC diet fed to the Sham PF group was the mean consumption of the VSG group
from the preceding day. Additionally, we included a group of mice that were fed a normal chow
diet (NCD) for the duration of the study. All animal experiments were approved by the University
of Minnesota Institutional Animal Care and Use Committee.
462
Macrophage characterization and isolation
Intrahepatic immune cells were isolated from perfused livers using enzymatic digestion as
previously described[73]. For flow cytometry, single cell suspensions were incubated with Zombie
NIR (1:200, Biolegend) for 20 min at room temperature, TruStain FcX (1:50, Biolgend) for 5 min
at room temperature, and with fluorophore-conjugated primary antibodies (1:100) for 30 min at
4°C. Following the staining, cells were washed and fixed with 100 μL Fixation Buffer (BioLegend)
for 20 min at 4°C. Flow cytometry data were acquired on a Fortessa flow cytometer (BD
Biosciences) and analyzed using Flowjo (TreeStar) software. For macrophage-specific assays,
macrophages were isolated from immune cell suspensions using anti-F4/80 microbeads and a
MACS separator (Miltenyi Biotech).
473
Bone marrow-derived macrophages
Bone marrow-derived macrophages (BMDM) were generated as previously described[74]. Briefly,
bone marrow was isolated from the femurs and tibias of male mice, then seeded in 2mL of
conditioned media with macrophage colony-stimulating factor (mCSF, 25 ng/mL, BioLegend).
Five days later, 1 mL of conditioned media with mCSF (50 ng/mL) was added. Seven days after
seeding, the BMDMs were washed and stimulated with palmitate for 24 hours (200 uM, Cayman
Chemical). Total RNA was extracted from the BMDMs using the RNeasy Plus Mini kit (Qiagen).
cDNA was prepared using the iScript cDNA Synthesis kit (Bio-Rad). Gene expression was
calculated using the 2(-ΔΔCT) method and normalized to GAPDH.
483
Efferocytosis assessment
Peritoneal macrophages were isolated from WT and TREM2 KO mice by lavage with phosphate-
buffered saline (PBS) with 2mM EDTA, as previously described[75]. Macrophages were then plated
on 24 well non-tissue culture-treated plates (Corning) in conditioned RPMI with 10% fetal bovine
serum and 1% penicillin and streptomycin for 1 hour. The media was replaced to remove non-
adherent cells and the cells were cultured for 4 hours. Subsequently, the peritoneal macrophages
were stained with CellTrace Violet (CTV, ThermoFisher) for 20 min at 37°C. AML12 hepatocytes
(ATCC) were treated with PA (1200 μM) for 24 hours to induce apoptosis (**Extended Data Fig.**
4F), then stained with CellTracker Green (CMFDA) for 30 min at 37°C. Following, CTV+
macrophages were cocultured with CMFDA+ apoptotic hepatocytes at a ratio of 1:4, respectively,
for 2 hours. The cells were then lifted, centrifuged, and resuspended for flow cytometry
acquisition. Flow cytometry data were acquired on a Fortessa flow cytometer (BD Biosciences)
and analyzed using Flowjo (TreeStar) software. Efferocytosis capacity was calculated as a
percentage of CTV+, and CMFDA+ cells relative to total CTV+ cells.
498

Fecal lipid extraction

Fecal lipid extraction was performed as previously described[76]. Feces were collected, added to 5 mL of PBS, and vortexed until the fecal pellets disintegrated. Next, 5 mL of chloroform:methanol (2:1, v/v) was added, vortexed, and centrifuged at 1,000g for 10 min. The lower liquid phase was collected, transferred to a glass tube, and evaporated in a fume hood. The remaining lipids were weighed.

MASH phenotyping

Hepatic triglycerides were assessed using a calorimetric assay (Cayman Chemical). Concentrations of serum ALT and AST were assessed by the University of Minnesota Veterinary Medical Center's Clinical Pathology Laboratory. Liver sections were fixed in 10% formalin and hematoxylin & eosin (H&E) and Picro Sirius Red staining was performed by the Biorepository & Laboratory Services at the UMN Clinical and Translational Science Institute. The collagen area was assessed using Picro Sirius Red stained liver sections as previously described[77]. NAS scoring of the H&E-stained liver sections was performed by a blinded pathologist.

Metabolite assessments

Metabolite assays were performed by the University of Minnesota's Center for Metabolomics and Proteomics. Hepatic macrophage metabolites were determined utilizing Biocrates MxP® Quant 500 kit. Lipids and hexoses were measured by flow injection analysis-tandem mass spectrometry (FIA-MS/MS) and small molecules were measured by liquid chromatography-tandem mass spectrometry (LC-MS/MS) using multiple reaction monitoring (MRM) for the detection of analytes utilizing a Shimadzu LC-20AD XR (Shimadzu USA Manufacturing Inc.) coupled to a Sciex QTRAP 5500 mass spectrometer (Sciex,). Hepatic portal vein bile acids and short chain fatty acids were acquired using an Agilent 1290 series HPLC (Agilent,) with an Acquity C18 BEH (2.1mm x 50mm, 1.7 um) column coupled to an Agilent 6495C Triple Quadrupole (Agilent) mass spectrometer in negative ion-mode. Bile acid samples were prepared by combining 100 uL of hepatic portal vein serum, 20 uL of the internal standard, 30 ul of HCL (1 M), and 1 mL of acetonitrile[78]. The acquired data was imported into Agilent MassHunter software for additional analysis. Hepatic portal vein short chain fatty acids samples were prepared by combining 90uL of hepatic portal vein serum 10 uL of the internal standard, and 420 µL of cold methanol. Concentrations of metabolites were calculated in the Biocrates MetIDQ™ software (Biocrates Life Sciences AG). Metabolites missing more than 20% of measurements were excluded from the statistical analysis. Missing values were imputed using the k-nearest neighbors approach as implemented in the *impute* R-package[79]. Hypothesis testing was performed using a Welch's two sided t-test. Lipid set enrichment was performed using LipidSuite software[80].

Bulk RNA sequencing

Total RNA was extracted from hepatic macrophages using the RNeasy Plus Mini kit (Qiagen). Samples were sequenced on a Novaseq 6000 using a 150 PE flow cell at the University of Minnesota Genomics Center. The SMARTer Stranded RNA Pico Mammalian V2 kit (Takara Bio) was used to create Illumina sequencing libraries. Differential gene expression analysis was performed using edgeR (Bioconductor). Gene ontology and pathway analysis were completed using iDEP.96 software[81].

Fecal microbiota assessment

DNA was extracted from single fecal pellets using the PowerSoil DNA isolation kit (QIAGEN) on the automated QIAcube platform using the inhibitory removal technology (IRT) protocol. The V4 hypervariable region of the 16S rRNA gene was amplified and sequenced using the 505F/816R primer set[82]. Paired-end sequencing was done at a length of 301 nucleotides (nt) on the Illumina

MiSeq platform by the University of Minnesota Genomics Center using their previously described
dual-index method[83]. Raw sequence data are deposited in the Sequence Read Archive under
BioProject accession number SRP325945. Sequence data processing and analysis were
performed using Mothur ver. 1.41.1[84]. Sequences were aligned against the SILVA database[85]
(ver. 138) for processing. Clustering of operational taxonomic units was done at 99% similarity
using the furthest-neighbor algorithm and taxonomic classifications were made against the
Ribosomal Database Project (ver. 18)[86]. Compositional data are reported from non-normalized
data. For statistical comparisons, samples were rarefied to 10,000 reads/sample[87]. Ordination of
samples was done using principal coordinate analysis[88], and differences in community
composition were evaluated by analysis of similarity (ANOSIM), with Bonferroni correction for
multiple comparisons[89]. Differences in alpha diversity were determined by ANOVA with Tukey's
posthoc test and differences in genera relative abundances were evaluated by the non-parametric
Kruskal-Wallis test, with pairwise comparisons done using the Steel-Dwass-Critchlow-Fligner
procedure. To determine dissimilarity from baseline composition, SourceTracker2[90] was used with
baseline samples in each group designated as the source. All statistics were evaluated at α=0.05
prior to correction for multiple comparisons.
566
Spatial transcriptomics
Fixed moused liver sections were mounted and stained with DAPI and an anti-CD68 to create
macrophage-free ROIs. Fixed human liver sections were mounted and stained with DAPI/anti-
CD68/anti-panCK antibodies to create zone-specific ROIs. Spatial sequencing was performed as
previously described using the GeoMx platform (NanoString)[91]. Analysis of the sequencing data
was performed on the GeoMx Digital Spatial Profiler (NanoString).
573
Human specimens information
We used human de-identified specimens previously collected by the Biological Materials
Procurement Network (BioNet) at the University of Minnesota. We used specimens collected by
liver biopsy from one patient before and one year VSG and from a patient diagnosed with MASH
without any surgical intervention (**Table 1**). Samples were obtained with institutional review board
approval.
580
581 **Table 1. Human spatial transcriptomics specimen information**

Specimen	Bx Date	NAS Score	Age	Sex	Race	BMI	DM Dx	MS Dx	Alcohol Use
PreVSG	11/13/19	4	42	F	Caucasian	35.3	PreDM	No	None
PostVSG	12/23/20	0	43	F	Caucasian	26.7	None	No	None
NASH	N/A	6	70	F	Caucasian	N/A	N/A	N/A	N/A

582
Single-cell RNA sequencing
Two million CD45[+] and F4/80[+] cells from each sample were individually incubated with cell
multiplexing oligos (CMO) to allow for downstream multiplexing of samples. Cells were washed
with PBS containing 0.04% BSA and centrifuged at 400g for 5 minutes at room temperature. The
supernatant was carefully discarded, and the cell pellet was resuspended in 100 µL of CMO for 5
minutes. Cells were washed with 1.9mL PBS containing 1% BSA and centrifuged at 400g for 5
minutes at 4ºC. Then, cells were washed three times using 2 mL of wash buffer. Following,
samples were combined at a concentration of 1,500 cells/µL and loaded into capture ports of a
10x-Genomics chip following a single cell 3' kit. Separate libraries were created for gene
expression and CMO then sequenced utilizing a Novaseq S4 chip (2 x 150bp PE). Feature
quantification was performed using cellranger (version 3.0, 10X Genomics). All subsequent
analyses were performed using Seurat 3.1.1[92]. After sample demultiplexing of CMOs, we retained

single cells and normalized within each pool using the 4 sctransform method in Seurat[92].
Visualization of different clusters was enabled using Uniform Manifold Approximation and
Projection (UMAP) dimensional reduction[93]. The Seurat cell ID function was utilized to identify
monocytes and macrophages, which were re-clustered for analysis. Differential expression
testing was performed using the default Wilcoxon rank-sum test. To evaluate the developmental
path of macrophages, we inferred trajectories using slingshot[37]. To limit our analyses to the
developmental stages of interest, we first created a data set only containing the major monocyte
and macrophage clusters. Then, we inferred the trajectories with cluster 6 set as the origin and
cluster 1 as the terminal state. To visualize gene expression patterns across developmental
pseudotime and across lineages, we first fit a negative binomial generalized additive model to the
observed gene expression, which was then used to predict smoothed expression values for each
cell and lineage using the tradeseq package[94]. All Seurat, slingshot, and tradeseq analyses were
conducted in R (v.4.1.0). Pathway analysis was completed using iDEP.96[81] and DAVID[95]
software. Gene set enrichment was performed using the GSEA software[96].

609
Statistical analysis
Statistical significance was determined using a one-way ANOVA with Holm-Šídák multiple
comparison test using GraphPad Prism (version 10.0.2). All data analyzed by ANOVA followed a
Gaussian distribution (Shapiro-Wilk test) and had equal variances (Brown-Forsythe). For datasets
that failed the normality test, statistical significance was determined using multiple non-parametric
two-sided Mann-Whitney tests. For in vitro experiments, each data point represents a biological
replicate with cells from one mouse. Mouse studies were repeated in at least two independent
experiments. Data are presented as means ± SEM with statistical significance denoted.

ACKNOWLEDGMENTS
This study was supported by the National Institute of Diabetes and Digestive and Kidney Diseases
(DK122056 to X.S.R.) and the National Heart, Lung, and Blood Institute (R01HL155993 to
X.S.R.). G.F. is supported by a T32 NIH training grant (T32DK083250) We recognize the staff
from the Research Animal Resources, University Flow Cytometry Resource, Center for
Metabolomics and Proteomics, Genomics Center, and the Clinical and Translational Science
Institute at the University of Minnesota for their assistance.

626
AUTHOR CONTRIBUTIONS
X.S.R., G.F., and S.I conceived the study and designed experiments. G.F. and X.S.R interpreted
results, generated figures and tables, and wrote the manuscript. G.F., K.F., F.B, K.D., H.W., P.P.,
C.J., C.S., and R.A. performed experiments. O.A. analyzed and provided data from liver
specimens. A.H. performed the analysis of single-cell RNA sequencing data. A.B., C.J., J.W., and
D.G.M. provided feedback and supervised aspects of the study. X.S.R obtained funding for,
supervised, and led the overall execution of the study.

REFERENCES
636
637 1.   Younossi, Z. *et al.* Global burden of NAFLD and NASH: trends, predictions, risk factors
638   and prevention. *Nature reviews. Gastroenterology & hepatology* **15**, 11-20 (2018).

639
640 2.   Marengo, A., Jouness, R.I. & Bugianesi, E. Progression and Natural History of
641   Nonalcoholic Fatty Liver Disease in Adults. *Clinics in liver disease* **20**, 313-324 (2016).

642
643 3.   Tilg, H. & Moschen, A.R. Evolution of inflammation in nonalcoholic fatty liver disease: the
644   multiple parallel hits hypothesis. *Hepatology (Baltimore, Md.)* **52**, 1836-1846 (2010).

646   4.   Schuster, S., Cabrera, D., Arrese, M. & Feldstein, A.E. Triggering and resolution of
647   inflammation in NASH. *Nat Rev Gastroenterol Hepatol* **15**, 349-364 (2018).
648
649   5.   Yanovski, S.Z. & Yanovski, J.A. Obesity. *The New England journal of medicine* **346**, 591-
650   602 (2002).
651
652   6.   Stefater, M.A., Wilson-Perez, H.E., Chambers, A.P., Sandoval, D.A. & Seeley, R.J. All
653   bariatric surgeries are not created equal: insights from mechanistic comparisons.
654   *Endocrine reviews* **33**, 595-622 (2012).
655
656   7.   Inge, T.H. *et al.* Perioperative outcomes of adolescents undergoing bariatric surgery: the
657   Teen-Longitudinal Assessment of Bariatric Surgery (Teen-LABS) study. *JAMA pediatrics*
658   **168**, 47-53 (2014).
659
660   8.   Cherla, D.V. *et al.* Impact of sleeve gastrectomy and Roux-en-Y gastric bypass on biopsy-
661   proven non-alcoholic fatty liver disease. *Surgical endoscopy* **34**, 2266-2272 (2020).
662
663   9.   Froylich, D. *et al.* Effect of Roux-en-Y gastric bypass and sleeve gastrectomy on
664   nonalcoholic fatty liver disease: a comparative study. *Surgery for obesity and related*
665   *diseases : official journal of the American Society for Bariatric Surgery* **12**, 127-131 (2016).
666
667   10. Wickremesekera, K., Miller, G., Naotunne, T.D., Knowles, G. & Stubbs, R.S. Loss of
668   insulin resistance after Roux-en-Y gastric bypass surgery: a time course study. *Obesity*
669   *surgery* **15**, 474-481 (2005).
670
671   11. Rizzello, M. *et al.* Early postoperative insulin-resistance changes after sleeve gastrectomy.
672   *Obesity surgery* **20**, 50-55 (2010).
673
674   12. Ryan, K.K. *et al.* FXR is a molecular target for the effects of vertical sleeve gastrectomy.
675   *Nature* **509**, 183-188 (2014).
676
677   13. le Roux, C.W. *et al.* Gut hormones as mediators of appetite and weight loss after Roux-
678   en-Y gastric bypass. *Annals of surgery* **246**, 780-785 (2007).
679
680   14. Saeidi, N. *et al.* Reprogramming of intestinal glucose metabolism and glycemic control in
681   rats after gastric bypass. *Science (New York, N.Y.)* **341**, 406-410 (2013).
682
683   15. Jahansouz, C. *et al.* Sleeve gastrectomy drives persistent shifts in the gut microbiome.
684   *Surgery for obesity and related diseases : official journal of the American Society for*
685   *Bariatric Surgery* **13**, 916-924 (2017).
686
687   16. Liou, A.P. *et al.* Conserved shifts in the gut microbiota due to gastric bypass reduce host
688   weight and adiposity. *Science translational medicine* **5**, 178ra141 (2013).
689
690   17. Evers, S.S. *et al.* Gut HIF2α signaling is increased after VSG, and gut activation of HIF2α
691   decreases weight, improves glucose, and increases GLP-1 secretion. *Cell Rep* **38**,
692   110270 (2022).
693
694   18. Remmerie, A. *et al.* Osteopontin Expression Identifies a Subset of Recruited Macrophages
695   Distinct from Kupffer Cells in the Fatty Liver. *Immunity* **53**, 641-657 e614 (2020).
696

697   19.   Xiong, X. *et al.* Landscape of Intercellular Crosstalk in Healthy and NASH Liver Revealed
698   by Single-Cell Secretome Gene Analysis. *Mol Cell* **75**, 644-660 e645 (2019).
699
700   20.   Guilliams, M. *et al.* Spatial proteogenomics reveals distinct and evolutionarily conserved
701   hepatic macrophage niches. *Cell* **185**, 379-396.e338 (2022).
702
703   21.   Wang, X. *et al.* Prolonged hypernutrition impairs TREM2-dependent efferocytosis to
704   license chronic liver inflammation and NASH development. *Immunity* **56**, 58-77.e11
705   (2023).
706
707   22.   Hendrikx, T. *et al.* Soluble TREM2 levels reflect the recruitment and expansion of
708   TREM2(+) macrophages that localize to fibrotic areas and limit NASH. *J Hepatol* (2022).
709
710   23.   Hou, J. *et al.* TREM2 sustains macrophage-hepatocyte metabolic coordination in
711   nonalcoholic fatty liver disease and sepsis. *J Clin Invest* **131**, e135197 (2021).
712
713   24.   Perugorria, M.J. *et al.* Non-parenchymal TREM-2 protects the liver from immune-mediated
714   hepatocellular damage. *Gut* **68**, 533-546 (2019).
715
716   25.   Liebold, I. *et al.* TREM2 Regulates the Removal of Apoptotic Cells and Inflammatory
717   Processes during the Progression of NAFLD. *Cells*; 2023.
718
719   26.   Deczkowska, A., Weiner, A. & Amit, I. The Physiology, Pathology, and Potential
720   Therapeutic Applications of the TREM2 Signaling Pathway. *Cell* **181**, 1207-1217 (2020).
721
722   27.   Asgharpour, A. *et al.* A diet-induced animal model of non-alcoholic fatty liver disease and
723   hepatocellular cancer. *J Hepatol* **65**, 579-588 (2016).
724
725   28.   Ben-Zvi, D. *et al.* Time-Dependent Molecular Responses Differ between Gastric Bypass
726   and Dieting but Are Conserved Across Species. *Cell Metab* **28**, 310-323 e316 (2018).
727
728   29.   Hao, Z. *et al.* RYGB Produces more Sustained Body Weight Loss and Improvement of
729   Glycemic Control Compared with VSG in the Diet-Induced Obese Mouse Model. *Obesity
730   surgery* **27**, 2424-2433 (2017).
731
732   30.   Ding, L. *et al.* Vertical sleeve gastrectomy confers metabolic improvements by reducing
733   intestinal bile acids and lipid absorption in mice. *Proceedings of the National Academy of
734   Sciences* **118**, e2019388118 (2021).
735
736   31.   Godlewska, U., Bulanda, E. & Wypych, T.P. Bile acids in immunity: Bidirectional mediators
737   between the host and the microbiota. *Front Immunol* **13**, 949033 (2022).
738
739   32.   Tran, S. *et al.* Impaired Kupffer Cell Self-Renewal Alters the Liver Response to Lipid
740   Overload during Non-alcoholic Steatohepatitis. *Immunity* **53**, 627-640.e625 (2020).
741
742   33.   Ghazarian, M. *et al.* Type I Interferon Responses Drive Intrahepatic T cells to Promote
743   Metabolic Syndrome. *Science immunology* **2** (2017).
744
745   34.   Barrow, F. *et al.* Microbiota-Driven Activation of Intrahepatic B Cells Aggravates NASH
746   Through Innate and Adaptive Signaling. *Hepatology* **74**, 704-722 (2021).
747

35. Blériot, C. *et al.* A subset of Kupffer cells regulates metabolism through the expression of CD36. *Immunity* **54**, 2101-2116.e2106 (2021).

36. Jaitin, D.A. *et al.* Lipid-Associated Macrophages Control Metabolic Homeostasis in a Trem2-Dependent Manner. *Cell* **178**, 686-698.e614 (2019).

37. Street, K. *et al.* Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics. *BMC Genomics* **19**, 477 (2018).

38. Daemen, S. *et al.* Dynamic Shifts in the Composition of Resident and Recruited Macrophages Influence Tissue Remodeling in NASH. *Cell Reports* **34**, 108626 (2021).

39. Ramachandran, P. *et al.* Resolving the fibrotic niche of human liver cirrhosis at single-cell level. *Nature* **575**, 512-518 (2019).

40. Hendrikx, T. *et al.* Soluble TREM2 levels reflect the recruitment and expansion of TREM2(+) macrophages that localize to fibrotic areas and limit NASH. *J Hepatol* **77**, 1373-1385 (2022).

41. Cauvi, D.M., Hawisher, D., Derunes, J. & De Maio, A. Phosphatidylcholine Liposomes Reprogram Macrophages toward an Inflammatory Phenotype. *Membranes (Basel)* **13** (2023).

42. Treede, I. *et al.* Anti-inflammatory Effects of Phosphatidylcholine*. *Journal of Biological Chemistry* **282**, 27155-27164 (2007).

43. Maceyka, M. & Spiegel, S. Sphingolipid metabolites in inflammatory disease. *Nature* **510**, 58-67 (2014).

44. Xia, J.Y. *et al.* Targeted Induction of Ceramide Degradation Leads to Improved Systemic Metabolism and Reduced Hepatic Steatosis. *Cell Metab* **22**, 266-278 (2015).

45. Blouin, C.M. *et al.* Plasma Membrane Subdomain Compartmentalization Contributes to Distinct Mechanisms of Ceramide Action on Insulin Signaling. *Diabetes* **59**, 600-610 (2009).

46. Yu, X.-D. & Wang, J.-W. Ceramide de novo synthesis in non-alcoholic fatty liver disease: Pathogenic mechanisms and therapeutic perspectives. *Biochemical Pharmacology* **202**, 115157 (2022).

47. Lassailly, G. *et al.* Bariatric Surgery Provides Long-term Resolution of Nonalcoholic Steatohepatitis and Regression of Fibrosis. *Gastroenterology* **159**, 1290-1301.e1295 (2020).

48. Verrastro, O. *et al.* Bariatric-metabolic surgery versus lifestyle intervention plus best medical care in non-alcoholic steatohepatitis (BRAVES): a multicentre, open-label, randomised trial. *Lancet* **401**, 1786-1797 (2023).

49. le Roux, C.W. & Bueter, M. The physiology of altered eating behaviour after Roux-en-Y gastric bypass. *Exp Physiol* **99**, 1128-1132 (2014).

799 50. Arble, D.M., Schwartz, A.R., Polotsky, V.Y., Sandoval, D.A. & Seeley, R.J. Vertical sleeve
800 gastrectomy improves ventilatory drive through a leptin-dependent mechanism. *JCI
801 Insight* **4** (2019).
802

803 51. Gihring, A. *et al.* Influence of bariatric surgery on the peripheral blood immune system of
804 female patients with morbid obesity revealed by high-dimensional mass cytometry. *Front
805 Immunol* **14**, 1131893 (2023).
806

807 52. Biobaku, F., Ghanim, H., Monte, S.V., Caruana, J.A. & Dandona, P. Bariatric Surgery:
808 Remission of Inflammation, Cardiometabolic Benefits, and Common Adverse Effects. *J
809 Endocr Soc* **4**, bvaa049 (2020).
810

811 53. Ahn, C.H. *et al.* Vertical sleeve gastrectomy induces distinctive transcriptomic responses
812 in liver, fat and muscle. *Sci Rep* **11**, 2310 (2021).
813

814 54. Griffin, C. *et al.* Inflammatory responses to dietary and surgical weight loss in male and
815 female mice. *Biology of Sex Differences* **10**, 16 (2019).
816

817 55. Han, H. *et al.* Macrophage-derived Osteopontin (SPP1) Protects From Nonalcoholic
818 Steatohepatitis. *Gastroenterology* (2023).
819

820 56. N, A.G. *et al.* Apoptotic cells promote their own clearance and immune tolerance through
821 activation of the nuclear receptor LXR. *Immunity* **31**, 245-258 (2009).
822

823 57. Wang, Y. *et al.* TREM2 ameliorates neuroinflammatory response and cognitive
824 impairment via PI3K/AKT/FoxO3a signaling pathway in Alzheimer's disease mice. *Aging
825 (Albany NY)* **12**, 20862-20879 (2020).
826

827 58. Rasheed, A. & Rayner, K.J. Macrophage Responses to Environmental Stimuli During
828 Homeostasis and Disease. *Endocr Rev* **42**, 407-435 (2021).
829

830 59. Xu, C. *et al.* Serum complement C3 levels are associated with nonalcoholic fatty liver
831 disease independently of metabolic features in Chinese population. *Scientific Reports* **6**,
832 23279 (2016).
833

834 60. Guo, Z. *et al.* The role of complement in nonalcoholic fatty liver disease. *Frontiers in
835 Immunology* **13** (2022).
836

837 61. Bohlson, S.S., O'Conner, S.D., Hulsebus, H.J., Ho, M.-M. & Fraser, D.A. Complement,
838 C1q, and C1q-Related Molecules Regulate Macrophage Polarization. *Frontiers in
839 Immunology* **5** (2014).
840

841 62. Furet, J.P. *et al.* Differential adaptation of human gut microbiota to bariatric surgery-
842 induced weight loss: links with metabolic and low-grade inflammation markers. *Diabetes*
843 **59**, 3049-3057 (2010).
844

845 63. Kong, L.C. *et al.* Gut microbiota after gastric bypass in human obesity: increased richness
846 and associations of bacterial genera with adipose tissue genes. *Am J Clin Nutr* **98**, 16-24
847 (2013).
848

64. Cani, P.D. *et al.* Changes in gut microbiota control metabolic endotoxemia-induced inflammation in high-fat diet-induced obesity and diabetes in mice. *Diabetes* **57**, 1470-1481 (2008).

65. Caesar, R., Tremaroli, V., Kovatcheva-Datchary, P., Cani, P.D. & Bäckhed, F. Crosstalk between Gut Microbiota and Dietary Lipids Aggravates WAT Inflammation through TLR Signaling. *Cell Metab* **22**, 658-668 (2015).

66. Shao, Y. *et al.* Vertical sleeve gastrectomy increases duodenal Lactobacillus spp. richness associated with the activation of intestinal HIF2α signaling and metabolic benefits. *Mol Metab* **57**, 101432 (2022).

67. Daws, M.R. *et al.* Pattern Recognition by TREM-2: Binding of Anionic Ligands. *The Journal of Immunology* **171**, 594-599 (2003).

68. Weaver, L.K. *et al.* Microbiota-dependent signals are required to sustain TLR-mediated immune responses. *JCI Insight* **4** (2019).

69. Abu-Gazala, S. *et al.* Sleeve Gastrectomy Improves Glycemia Independent of Weight Loss by Restoring Hepatic Insulin Sensitivity. *Diabetes* **67**, 1079-1085 (2018).

70. Ulker, İ. & Yildiran, H. The effects of bariatric surgery on gut microbiota in patients with obesity: a review of the literature. *Biosci Microbiota Food Health* **38**, 3-9 (2019).

71. Wu, Y. *et al.* TREM-2 is a sensor and activator of T cell response in SARS-CoV-2 infection. *Sci Adv* **7**, eabi6802 (2021).

72. Wu, Y. *et al.* TREM-2 promotes Th1 responses by interacting with the CD3ζ-ZAP70 complex following Mycobacterium tuberculosis infection. *The Journal of Clinical Investigation* **131** (2021).

73. Lynch, R.W. *et al.* An efficient method to isolate Kupffer cells eliminating endothelial cell contamination and selective bias. *J Leukoc Biol* **104**, 579-586 (2018).

74. Bailey, J.D. *et al.* Isolation and culture of murine bone marrow-derived macrophages for nitric oxide and redox biology. *Nitric Oxide* **100-101**, 17-29 (2020).

75. Zhang, X., Goncalves, R. & Mosser, D.M. The isolation and characterization of murine macrophages. *Curr Protoc Immunol* **Chapter 14**, 14.11.11-14.11.14 (2008).

76. Kraus, D., Yang, Q. & Kahn, B.B. Lipid Extraction from Mouse Feces. *Bio Protoc* **5** (2015).

77. Fredrickson, G. *et al.* Exercise of high intensity ameliorates hepatic inflammation and the progression of NASH. *Mol Metab* **53**, 101270 (2021).

78. Krautbauer, S. & Liebisch, G. LC-MS/MS Analysis of Bile Acids. In: Giera, M. (ed). *Clinical Metabolomics: Methods and Protocols*. Springer New York: New York, NY, 2018, pp 103-110.

79. Troyanskaya, O. *et al.* Missing value estimation methods for DNA microarrays. *Bioinformatics* **17**, 520-525 (2001).

80.   Mohamed, A. & Hill, M.M. LipidSuite: interactive web server for lipidomics differential and
enrichment analysis. *Nucleic Acids Research* **49**, W346-W351 (2021).

81.   Ge, S.X., Son, E.W. & Yao, R. iDEP: an integrated web application for differential
expression and pathway analysis of RNA-Seq data. *BMC Bioinformatics* **19**, 534 (2018).

82.   Caporaso, J.G. *et al.* Ultra-high-throughput microbial community analysis on the Illumina
HiSeq and MiSeq platforms. *The ISME Journal* **6**, 1621-1624 (2012).

83.   Gohl, D.M. *et al.* Systematic improvement of amplicon marker gene methods for increased
accuracy in microbiome studies. *Nature Biotechnology* **34**, 942-949 (2016).

84.   Schloss Patrick, D. *et al.* Introducing mothur: Open-Source, Platform-Independent,
Community-Supported Software for Describing and Comparing Microbial Communities.
Applied and Environmental Microbiology **75**, 7537-7541 (2009).

85.   Pruesse, E. *et al.* SILVA: a comprehensive online resource for quality checked and aligned
ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res* **35**, 7188-7196
(2007).

86.   Cole, J.R. *et al.* The Ribosomal Database Project: improved alignments and new tools for
rRNA analysis. *Nucleic Acids Research* **37**, D141-D145 (2009).

87.   Gihring, T.M., Green, S.J. & Schadt, C.W. Massively parallel rRNA gene sequencing
exacerbates the potential for biased community diversity comparisons due to variable
library sizes. *Environ Microbiol* **14**, 285-290 (2012).

88.   Anderson, M.J. & Willis, T.J. CANONICAL ANALYSIS OF PRINCIPAL COORDINATES:
A USEFUL METHOD OF CONSTRAINED ORDINATION FOR ECOLOGY. *Ecology* **84**,
511-525 (2003).

89.   Clarke, K.R. Non-parametric multivariate analyses of changes in community structure.
Australian Journal of Ecology **18**, 117-143 (1993).

90.   Knights, D. *et al.* Bayesian community-wide culture-independent microbial source
tracking. *Nature Methods* **8**, 761-763 (2011).

91.   Zollinger, D.R., Lingle, S.E., Sorg, K., Beechem, J.M. & Merritt, C.R. GeoMx™ RNA
Assay: High Multiplex, Digital, Spatial Analysis of RNA in FFPE Tissue. In: Nielsen, B.S.
& Jones, J. (eds). *In Situ Hybridization Protocols.* Springer US: New York, NY, 2020, pp
331-345.

92.   Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-
seq data using regularized negative binomial regression. *Genome Biology* **20**, 296 (2019).

93.   Becht, E. *et al.* Dimensionality reduction for visualizing single-cell data using UMAP.
Nature Biotechnology **37**, 38-44 (2019).

94.   Van den Berge, K. *et al.* Trajectory-based differential expression analysis for single-cell
sequencing data. *Nat Commun* **11**, 1201 (2020).

951
952   95.   Sherman, B.T. *et al.* DAVID: a web server for functional enrichment analysis and functional
953   annotation of gene lists (2021 update). *Nucleic Acids Res* **50**, W216-w221 (2022).
954
955   96.   Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for
956   interpreting genome-wide expression profiles. *Proceedings of the National Academy of*
957   *Sciences* **102**, 15545-15550 (2005).
958

Figure 1. Vertical sleeve gastrectomy (VSG) ameliorates MASH progression independent of weight loss. (**A**) Experimental design, (**B**) Mean daily food intake post-surgery, (**C**) Weekly body weights post-surgery, (**D**) Liver weight, (**E**) Hepatic triglyceride content, (**F**) Representative H&E-stained liver sections, (**G**) Serum ALT (Left) and AST (Right), (**H**) Collagen area in Picro Sirius Red-stained liver sections, (**I**) Fecal lipid content, (**J**) Relative abundance of Lactobacillus, (**K**) Relative abundance of microbiota species, (**L**) Relative abundance of Clostridium XVIII, (**M**) Total portal vein bile acids, (**N**) Concentration of portal vein taurocholic acid (TCA), deoxycholic acid (DCA), and taurodeoxycholic acid (TDCA), and (**O**) Total portal vein short-chain fatty acids (SCFA) in C57BL6/J (WT) mice fed an HFHC diet for 12 weeks before assignment to Sham ad libitum (Sham AL; n = 6-17), Sham pair-fed (Sham PF; n = 6-17), or VSG (n = 6-25) surgeries. Mice were maintained on the HFHC diet for 5 weeks post-surgery. A cohort of mice without any intervention were fed a normal chow diet (NCD; n = 5-25) throughout the study. Data are biological experimental units presented as mean ± standard error of the mean (SEM). Except for the microbiota data, which were analyzed by a non-parametric Kruskal-Wallis test, data were analyzed by one-way ANOVA with Holm-Šídák multiple comparison test.

Figure 2. ScRNA-seq reveals profound effects of VSG on hepatic LAMs. (**A**) Experimental design, (**B**) Proportion (left) and number (right) of sequenced cells per sample, (**C**) Integrated uniform manifold approximation and projection (UMAP) analysis of all monocytes and macrophages and expression of marker genes, (**D**) Heat and dot plot of the expression and coverage of marker genes in all subsets, (**E**) Pathway analysis of upregulated differentially expressed genes (DEG) in LAMs, compared with all other subsets (Gene ontology terms, p-value < 0.1), and (**F**) Slingshot trajectory analysis (top left) and gene expression over pseudotime for trajectories 1 (top right), 2 (bottom left), and 3 (bottom right) from single-cell RNA sequencing (scRNA-seq) of hepatic macrophages and monocytes from Sham AL (n = 8), Sham PF (n = 8), and VSG (n = 8) mice 5-10 weeks post-surgeries. Differential expression testing was performed by a Wilcoxon rank-sum test.

Figure 3. VSG enhances lipid metabolism and lysosomal gene programs in hepatic LAMs.
(**A**) UMAPs (left), relative cluster proportion (middle), and number of cells per cluster (right) and (**B**) Expression of *Trem2* in cluster 1 (LAMs) from the scRNA-seq analysis of monocytes and macrophages from Sham AL, Sham PF, or VSG groups. (**C**) Concentration of soluble TREM2 (sTREM2) in the serum of NCD (n = 8), Sham AL (n = 7), Sham PF (n = 9), and VSG (n = 8) mice 5 weeks post-surgeries. (**D**) Volcano plots showing differentially expressed genes (DEGs) between Sham AL and VSG (left), and Sham PF and VSG (right) LAMs, (**E**) Pathway analysis of DEGs in LAMs from Sham AL vs. VSG (top) and Sham PF vs. VSG (bottom) comparisons, and (**F**) Gene seat enrichment analysis (GSEA) of the DEGs between Sham PF and VSG LAMs in the scRNA-seq data. The cell number and sTREM2 concentration data were analyzed by one-way ANOVA with Holm-Šídák multiple comparison test. Differential expression testing was performed by a Wilcoxon rank-sum test. Pathway analysis was performed by Generally Applicable Gene-set Enrichment Analysis (GAGE, p-value < 0.1). Data are biological experimental units presented as mean ± SEM.

Figure 4. Hepatic TREM2⁺ LAMs mediate the reparative effects of VSG against MASH. (**A**) Body weight change post-surgery, (**B**) Fecal lipid content, (**C**) Liver weight, (**D**) Representative H&E-stained liver sections, (**E**) Hepatic triglyceride content, (**F**) Serum ALT and (**G**) AST, (**H**) Representative flow plot (top) and quantification of monocyte-derived (MoMFs) and F4/80^{hi} CD11b^{int} macrophages, (**I**) Unsupervised PCA of bulk RNA-seq gene expression data from F4/80⁺ sorted macrophages. (**J**) Venn diagram with the number of differentially expressed genes (DEGs) between Sham and VSG in WT and TREM2 KO macrophages, and (**K**) Pathway analysis (KEGG and Gene ontology) of DEGs between WT and TREM2 KO macrophages from sham or VSG mice. C57BL6/J (WT) and TREM2 KO (KO) mice were fed an HFHC diet for 12 weeks, assigned to either sham or VSG surgeries, and analyzed 5 weeks post-surgery (WT Sham, n = 5-8; WT VSG, n = 6-7; KO Sham, n = 5-8; and KO VSG, n = 5-8). (**L**) Gene expression of inflammatory cytokines in bone marrow-derived macrophages from WT (n = 3-7) or TREM2 KO (n = 4-6) mice left unstimulated (unstim, top) or stimulated with palmitate (PA, bottom). (**M**) Representative flow plots (left) and quantification (right) of CellTracker Green (CMFDA)-positive macrophages following coculture of peritoneal macrophages from WT (n = 7) and TREM2 KO (n = 5) mice with CMFDA-labeled apoptotic AML12 hepatocytes. Data from four experimental groups were analyzed by one-way ANOVA with Holm-Šídák multiple comparison test. Pathway analysis was performed by GAGE (p-value < 0.05). Data from two experimental groups were analyzed by Mann Whitney tests. Data are biological experimental units presented as mean ± SEM.

Figure 5. VSG increases the content of inflammatory lipid species in hepatic macrophages. (**A**) Composition of detectable metabolites in macrophages, (**B**) Unsupervised PCA of metabolite data, (**C**) Concentration of triacylglycerols (TG), (**D**) Chain length enrichment analysis of lipid species. The box represents the interquartile range, the middle line is the median, and the top and bottom lines indicate quartile 3 and quartile 1, respectively. Dots indicate outliers and red boxes indicate chain lengths with statistical significance between Sham AL and VSG, (**E**) Concentration of major lipid species including phosphatidylcholines (PC), cholesterol esters (CE), fatty acids (FA), glycosylceramides (HexCer), ceramides (Cer), sphingolipids (SM), and diacylglycerols (DG), (**F**) PC subspecies such as PCs (aa), PCs (ae), monounsaturated fatty acid (MUFA) PCs, polyunsaturated fatty acid (PUFA) PCs, and saturated fatty acid (SFA) PCs, (**G**) SM subspecies including long chain fatty acid (LCFA) SMs, very long chain fatty acid (VLCFA) SMs, and SMs with or without hydroxyl (-OH) groups, (**H**) Cer subspecies including VLCFA-Cer and LCFA-Cer, (**I**) FA subspecies including MUFAs, PUFAs, and omega-3 (w-3) FAs, and (**J**) CE subspecies determined by mass spectrometry and liquid chromatography of hepatic macrophages isolated from the livers of HFHC-fed mice assigned to Sham AL (n = 4) or VSG (n = 4) 5 weeks post-surgery. Data were analyzed by Welch's two-sided t tests. Data are biological experimental units presented as mean ± SEM.

Figure 6. Spatial transcriptomic of MASH livers following bariatric surgery reveals an improved metabolic status in the macrophage microenvironment. (**A**) Representative immunofluorescence images showing an entire section (left) and a magnified field (right) with CD68⁻ (orange) and CD68⁺ regions of interest (ROIs, green) in liver specimens analyzed by spatial transcriptomics (Geomx). (**B**) Steps in data processing (top) and number of genes after processing (bottom), (**C**) Unsupervised PCA of gene expression data from CD68⁺ and CD68⁻ ROIs, Volcano plots showing differentially expressed genes between (**D**) NCD and Sham AL, (**E**) Sham AL and VSG, (**F**) Sham PF and VSG CD68⁻ ROIs. Pathway analysis between (**G**) NCD and Sham AL, (**H**) Sham AL and VSG (**I**) Sham PF and VSG CD68⁻ ROIs. Liver specimens were collected from C57BL6/J (WT) fed either a NCD or an HFHC diet for 12 weeks, assigned to Sham AL, Sham PF, or VSG surgeries, and analyzed 5 weeks post-surgery (n = 4). Data were analyzed utilizing a Mann-Whitney test and corrected with a Benjamini-Hochberg procedure. Pathway analysis was performed by GSEA (p-value < 0.05).

Extended Data Figure 1 (related to Figure 1). (**A**) NAFLD activity score; NCD (n = 8), Sham AL (n = 6), Sham PF (n = 4), VSG (n = 5), and (**B**) Representative Picro Sirius red-stained liver sections 5 weeks post surgeries. (**C**) Percentage change in body weight, (**D**) Liver weight, (**E**) Serum AST, (**F**) Serum ALT, and (**G**) Hepatic triglyceride content in C57BL6/J (WT) mice fed either an NCD (n = 8) for 22 weeks or an HFHC diet for 12 weeks, assigned to Sham AL (n = 6-7), Sham PF (n = 6), or VSG (n = 9) surgeries, and maintained on the HFHC diet for 10 weeks post-surgery. (**H**) Concentration of portal vein bile acids 5 weeks post-surgery: cholic acid (CA), ursodeoxycholic acid (UDCA), chenodeoxycholic acid (CDCA), taurolithocholic acid (TLCA), lithocholic acid (LCA), taurochenodeoxycholic acid (TCDCA), hyodeoxycholic acid (HDCA), taurohyodeoxycholic acid (THDCA), tauroursodeoxycholic acid (TUDCA); NCD (n = 10), Sham AL (n = 6), Sham PF (n = 8), VSG (n = 12). Data were analyzed by one-way ANOVA with Holm-Šídák multiple comparison test. Data are biological experimental units presented as mean ± SEM.

Extended Data Figure 2 (related to Figure 2). (**A**) Representative flow plots (left) and quantification of hepatic macrophage subsets per gram of liver (right) at 5 weeks (top) and 10 weeks post-surgery (bottom); 5 week Sham AL (n = 4), 5 week VSG (n = 5), 10 week Sham AL (n = 6) and 10 week VSG (n = 9). (**B**) Heat map of top genes for each cluster. (**C**) Gene expression over pseudotime for trajectories 1 (top left), 2 (top right) and 3 (bottom). Data were analyzed by Mann Whitney tests. Data are biological experimental units presented as mean ± SEM.

A

Extended Data Figure 3 (related to Figure 3). (**A**) Heat maps of top DEGs between all groups at both time points for total macrophages. All data presented in this figure (n = 4).

Extended Data Figure 4 (related to Figure 4). (**A**) Relative expression *Trem2* in WT and TREM2 KO BMDMs; WT (n = 3) and KO (n = 3). (**B**) Concentration of sTREM2 in the serum 5 weeks post-VSG; WT Sham (n = 8), WT VSG (n = 6), KO Sham (n = 8), and KO VSG (n = 8). (**C**) Representative flow plots (top) and quantification of hepatic macrophage subsets per gram of liver (bottom) after 12 weeks of HFHC feeding; WT (n = 15) and KO (n = 8). (**D**) Quantification of hepatic macrophage subsets per gram of liver 5 weeks post-VSG; WT (n = 5) and KO (n = 5). (**E**) Heat map of the top 500 most variable genes between WT Sham and WT VSG macrophages (left) and KO Sham and KO VSG macrophages (right). (**F**) Representative flow plots of AML12 cells that were either unstimulated or stimulated with PA. Data were analyzed by Mann Whitney tests. Data are biological experimental units presented as mean ± SEM.

Name	Abbreviation	Class	Raw P-val	Abundance Ratio Sham AL / VSG	Abundance CV Sham	Abundance CV VSG
Phosphatidylcholine ae C34:2	PC ae C34:2	Glycerophospholipids	0.0025	0.5881	15.0864	8.4421
Dihexosylceramide(d18:1/18:0)	Hex2Cer(d18:1/18:0)	Glycosylceramides	0.0059	0.5484	19.5180	18.5255
Hydroxysphingomyelin C16:1	SM (OH) C16:1	Sphingolipids	0.0099	0.6133	18.5626	14.1484
Sphingomyelin C16:1	SM C16:1	Sphingolipids	0.0106	0.6695	13.9359	16.2424
Phosphatidylcholine ae C36:3	PC ae C36:3	Glycerophospholipids	0.0119	0.6389	19.1640	6.1530
Phosphatidylcholine ae C32:1	PC ae C32:1	Glycerophospholipids	0.0124	0.6525	17.0621	14.1475
Phosphatidylcholine ae C34:1	PC ae C34:1	Glycerophospholipids	0.0130	0.6809	15.8386	11.1349
Phosphatidylcholine ae C30:2	PC ae C30:2	Glycerophospholipids	0.0137	0.6019	23.7268	18.7709
Phosphatidylcholine aa C28:1	PC aa C28:1	Glycerophospholipids	0.0139	0.7055	6.8880	15.9828
Sphingomyelin C16:0	SM C16:0	Sphingolipids	0.0151	0.6438	17.2193	18.1432
Phosphatidylcholine ae C40:3	PC ae C40:3	Glycerophospholipids	0.0159	0.6162	21.2772	15.6241
Phosphatidylcholine ae C38:4	PC ae C38:4	Glycerophospholipids	0.0209	0.6981	10.5045	18.3192
Phosphatidylcholine aa C30:0	PC aa C30:0	Glycerophospholipids	0.0222	0.6534	16.7497	20.7499
Phosphatidylcholine ae C36:2	PC ae C36:2	Glycerophospholipids	0.0253	0.6661	18.3194	4.1832
Phosphatidylcholine ae C36:1	PC ae C36:1	Glycerophospholipids	0.0258	0.7076	17.9638	10.3597
Hexosylceramide(d18:1/26:0)	HexCer(d18:1/26:0)	Glycosylceramides	0.0264	0.6677	15.0474	20.3008
Trihexosylceramide(d18:1/24:1)	Hex3Cer(d18:1/24:1)	Glycosylceramides	0.0272	0.4968	24.1495	41.8201
Phosphatidylcholine ae C36:4	PC ae C36:4	Glycerophospholipids	0.0275	0.6746	17.7006	20.9567
Phosphatidylcholine ae C38:5	PC ae C38:5	Glycerophospholipids	0.0292	0.7129	14.6775	17.7371
Phosphatidylcholine aa C32:1	PC aa C32:1	Glycerophospholipids	0.0299	0.6131	25.4020	13.5715

Extended Data Figure 5 (related to Figure 5). Table of top metabolites in hepatic macrophages 5 weeks post-VSG.

A

Metabolism

Organelle biogenesis and maintenance

Gene expression (Transcription)

Cell Cycle

Cellular responses to external stimuli

Coverage
0%

100.00%

Transport of small molecules Neuronal System

Signal Transduction

Immune System

Extracellular matrix organization

Developmental Biology

Vesicle-mediated transport

Metabolism of proteins

Hemostasis

Autophagy

DNA Repair

Cell-Cell communication

Chromatin organization

Programmed Cell Death

DNA Replication

Metabolism of RNA

B

MASH

DAPI
CD68
panCK

PreVSG

PostVSG

C

DAPI
CD68
panCK

Zone 1

Zone 2

Zone 3

D

PC3

-10

-20

-30

0   50
PC1

Group
MASH
PostVSG
PreVSG

ID
1
2
3

E

Selenoamino acid metabolism
Metabolism of amino acids and derivatives
Metabolism of RNA
Respiratory electron transport
Platelet degranulation
Bile acid and bile salt metabolism
Fatty acyl-CoA biosynthesis
Metabolism of lipids
Regulation of cholesterol biosynthesis by SREBP
Metabolism of steroids

-5   0   5   10
Zone 1 PreVSG vs PostVSG
Normalized Enrichment

Metabolism of amino acids and derivatives
Metabolism of RNA
Gluconeogenesis
TCA cycle and respiratory electron transport
Regulation of Complement cascade
Metabolism of lipids
Metabolism of steroids
Regulation of cholesterol biosynthesis by SREBP
Activation of gene expression by SREBF

-5   0   5   10
Zone 2 PreVSG vs PostVSG
Normalized Enrichment

F

MASH   PreVSG   PostVSG
I   II   III   I   II   III   I   II   III

Cd63
Cd68
Plin2
Trem2
Glul
Dhrs3
Socs3
Ctss
Pld3
Ctsl
Cd36
Lipa
Itgb5
Blvrb

60   90   120   150

Extended Data Figure 6 (related to Figure 6). (**B-F**) Spatial sequencing of human liver samples pre-VSG and post-VSG. (**A**) Pathway coverage for mouse VSG GeoMx analysis. (**B**) Representative human liver sections used for GeoMx. (**C**) Representative human liver section used for GeoMx with liver zones indicated. (**D**) Unsupervised PCA of all human liver ROIs (n = 1). (**E**) Pathway analysis for Zone 1 PreVSG vs. Zone 1 PostVSG (top) and Zone 2 PreVSG vs. Zone 2 PostVSG (bottom). (**F**) Heatmap of LAM genes.