The orientation of choroidal macrophage polarization significantly influences the development of myopia in murine models

Jing Hou[1,2#], Shin-ichi Ikeda[1,2#], Kiwako Mori[1,2], Heonuk Jeong[1,2], Hidemasa Torii[1,2], Kazuno Negishi[1], Kazuo Tsubota[1,3*], Toshihide Kurihara[1,2*]

[1]Department of Ophthalmology, Keio University School of Medicine, 35 Shinanomachi, Shinjuku-ku, 160-8582, Tokyo, Japan.

[2]Laboratory of Photobiology, Keio University School of Medicine, 35 Shinanomachi, Shinjuku-ku, 160-8582, Tokyo, Japan.

[3]Tsubota Laboratory, Inc., 304 Toshin Shinanomachi-ekimae Bldg., 34 Shinanomachi Shinjuku-ku, 160-0016, Tokyo, Japan.

[#] Those authors contributed equally to the work and request the double first authorship.

*Corresponding authors:

Toshihide Kurihara, MD, PhD

Principal Investigator of Laboratory of Photobiology and Associate Professor of the Department of Ophthalmology, Keio University School of Medicine; 35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan.

Tel: +81-3-5363-3204, Fax: +81-3- 5363-3274

E-mail: kurihara@z8.keio.jp

Kazuo Tsubota, MD, PhD

Professor Emeritus of the Department of Ophthalmology, Keio University School of Medicine, and

the CEO of Tsubota Laboratory, Inc.;304 Toshin Shinanomachi-Eki Mae Bldg., 34 Shinanomachi

Shinjuku-ku, Tokyo 160-0016, Japan.

Tel: +81-3-6384-2866

E-mail: tsubota@tsubota-lab.com

Competing Interest Statement

K.T. reports that he is the CEO of Tsubota Laboratory, Inc., Tokyo, Japan, a company that

develops products for the treatment of myopia. The other authors have declared that they have

no conflicts of interest. Patent-pending (J.H. S.I. T.K. K.T. 2022-72596).

Funding

This work is supported by AMED under Grant Number JP22gm1510007 to T. Kurihara.

Keywords: Macrophage; Myopia; Choroid; Vascular Maintenance; Anti-inflammatory.

Article Type: Research paper

Word count: 199 words for the abstract

Number of table(s) and figure(s): 4 figures, and 3 supplementary figures

Abstract

Myopia is a primary contributor to visual impairment and has emerged as a global public health

concern. Evidence indicates that one of the main structural features of myopia is the

corresponding decrease in choroidal thickness, and choroidal macrophages play an important

role in maintaining the choroidal thickness. Nevertheless, the effect of choroidal macrophages on

myopia remains unclear. Here, we discovered that the continuous intraperitoneal injection of

clodronate liposomes depleted choroidal macrophages and leads to myopia, which confirmed that

the presence of choroidal macrophages plays an important role in myopia development.

Subsequently, based on the phenotypic characteristics of macrophages, experiments were

designed to study the effects of different polarization directions of macrophages on myopia

development. We found that lipopolysaccharides (LPS) injection can induce the polarization of

choroidal M1 macrophages, thinning the choroidal thickness and resulting in myopia. Conversely,

IL-4 or IL-13 injection causes choroidal M2 macrophage polarization, thickens the choroid, and

suppresses the progression of myopia. Additionally, we demonstrated that the opposite effects of

M1 and M2 macrophages on myopia development may be related to their impacts on choroidal

thickness, inflammation, and oxidative stress response. These findings establish that choroidal

macrophages are critically important in the development of myopia and provide new strategies for

the development of myopic therapies.

Main Text

Introduction

Myopia, commonly referred to as "short-sightedness," is the most prevalent vision-related

disease, which is caused by refractive error depending on the elongation of the eyeball along the

visual axis (1-4). Current research has shown that myopia can be influenced by a variety of

factors, including light environmental impacts, dietary factors, and genetics; however, the

pathogenesis of myopia remains unclear (5-8). Numerous studies have suggested that

inflammation plays a role in the development of myopia: increased inflammation in the eye

accelerates the development of myopia, whereas lower inflammation inhibits it (9-12). It is worth

noting that macrophages are essential for the occurrence, maintenance, and resolution of

inflammation (13).

Macrophages are an essential component of the innate immune system that exists in all

mammalian tissues and exhibit great functional diversity. It not only plays a variety of functions

related to inflammation but also plays a crucial role in the development and maintenance of

homeostasis (14, 15). A healthy cornea contains a population of macrophages that maintain

corneal lymphatic homeostasis and promote lymphangiogenesis under inflammatory conditions

(16). The neural retina is populated by microglia, specialized resident macrophages that

contribute to the structural and functional integrity of neuronal synapses (16, 17). Under

homeostatic conditions, there are no microglia in the subretinal space or the choroid and are

restricted to the inner layers of the retina (18-20). Further, the choroid was found to contain a

dense network of resident tissue macrophages as early as the 1990s (21). At the level of

choroidal arteries and arterioles, dendritic resident macrophages were closely aligned along the

vessel wall. In the choroidal terminal choriocapillaris, they are not tightly packed on the vessel

wall, but are widely distributed in the connective tissue between blood vessels (22). These

findings provide a basis for exploring the relationship between choroidal vessels and

macrophages. Subsequently, researchers found that the ablation of a considerable majority of

choroidal macrophages was associated with broad and gradual atrophy of the choroidal

vasculature and thinning of the choroid (23, 24). This finding reveals that choroidal macrophages

play a hitherto underestimated role in the maintenance of the choroidal vascular structure. In the

field of myopia research, the stability of the choroidal vasculature is recognized as the most

important component for maintaining choroidal thickness, and choroidal thickness variation is a

structural characteristic of the development of myopia (25-28). Therefore, we hypothesized that

choroidal macrophages play a crucial role in the progression of myopia.

Macrophage subsets can be classified as classic M1 macrophages and alternative M2

macrophages, according to their pro- or anti-inflammatory properties in the polarized state (29).

M1 macrophages can be polarized by lipopolysaccharide (LPS), leading to a high expression of

inflammatory cytokines, such as tumor necrosis factor-α (TNFA) and interleukin-6 (IL6), as well

as reactive oxygen species. Correspondingly, M2 macrophage polarization induced by IL-4 and

IL-13 diminishes and alleviates inflammation following infection or damage, and inhibits the

production of reactive oxygen species (30-32). Studies have shown that inflammatory diseases

increase the risk of myopia, and inflammation in the eye directly contributes to the progression of

myopia (12, 33), all this underscores the importance of controlling inflammation in inhibiting the

development of myopia. In addition, it has been suggested that the existence of several

supplements which convert M1 into M2 macrophages, but this property has not been studied and

discussed in conjunction with myopia suppression (34-36). Therefore, we hypothesized that

different subsets of macrophages in the choroid have diametrically opposite effects on the

development of myopia according to their polarization direction and characteristics.

In this study, we investigated the depletion and polarization of choroidal macrophages in a mouse

model and combined a lens-induced myopia model (LIM) with the polarization of choroidal

macrophage to investigate the effect of different macrophage polarization directions on myopia

development. We found that choroidal macrophage ablation was associated with myopia

progression. We also discovered that the tendency for myopia development or inhibition is closely

related to the direction of macrophage polarization in the choroid and that macrophages interfere

with the development of myopia based on their effects on inflammation and oxidative stress.

Taken together, our findings suggest that macrophages play a key role in myopia and emphasize

the different characteristics of macrophages as a potential mechanism for developing or inhibiting

myopia. Regulating the phenotype of macrophages in the choroid may have therapeutic effects

on maintaining choroidal vasculature thereby slowing the progression of myopia.

Results

Sustained intraperitoneal injection of clodronate liposome depletes macrophages in the

choroid and can trigger myopia

It has been previously reported that resident macrophages in the choroid positively affect the

choroidal thickness and maintain vascular status (24). In contrast, a series of animal models have

demonstrated that choroidal thinning is an inevitable change in the development of experimental

myopia and is a prominent feature in the development of myopia (26, 28, 37-39). To investigate

the effect of choroidal-resident macrophages on myopia in a mouse model, 3-weeks-old C57BL/6

J mice were randomly divided into 2 groups and were injected intraperitoneally with clodronate or

phosphate-buffered saline (PBS) liposomes. In the clodronate liposome group, after being

engulfed by macrophages, the clodronate liposomes were degraded by lysosomal

phospholipases, which released clodronate into the macrophages and induced apoptosis, thereby

promoting macrophage depletion (40). The injection volume of clodronate liposome was 0.1 ml

per 10 g of mice and the injection frequency was once every two days. Refraction, axial length,

and choroidal thickness were measured using an infrared photorefractor and a spectral domain-

optical coherence tomography (SD-OCT) system, before injection and after 8 days of injections,

respectively (Fig. 1A). Mice injected with clodronate liposome showed a higher refractive shift

(−7.30 ± 3.61 D vs. +1.17 ± 4.40 D, $P < 0.001$), axial length elongation (0.08 ± 0.02 mm vs.

0.05 ± 0.02 mm, $P < 0.01$), and thinner choroid (−1.56 ± 0.89 mm vs. 1.09 ± 0.51 mm, $P < 0.001$)

compared to the PBS liposome injection group (Fig. 1B). Flow cytometry analysis of the

percentage of choroidal resident macrophages demonstrated a corresponding decrease after 8

days of clodronate liposome injection compared to the control group (2.583 ± 0.002% vs. 9.290

± 0.018%, $P < 0.05$) (Fig. 1C). These results indicated that clodronate liposome injection depletes

resident macrophages in the choroid, promotes choroidal thinning, and induces myopia in 3-

week-old mice. Previous studies showed that choroidal thickness did not increase and tended to

be stable from approximately 8 weeks of age in mice (41). However, whether the depletion of

choroidal macrophages in adult mice also contributes to myopia has not been elucidated. To rule

out the possibility that choroidal macrophage depletion can cause myopia independent from the

developmental stage of choroidal thickness, 8-weeks-old adult mice were injected with clodronate

liposomes or PBS liposomes in the same manner, and changes in refraction, axial length, and

choroidal thickness were analyzed 8 days later (Fig. 1D). Intraperitoneal injection of clodronate

liposomes significantly reduced body weight while consuming resident macrophages, which

affects the comparison of changes in the axial length of the eye. To avoid the effects of body

weight changes on the axial length, we used the ratio of the axial length to body weight to

represent the changes in the axial length of the eye. In the clodronate liposome injection group, a

higher refraction shift (−7.52 ± 2.57 D vs. +1.52 ± 4.91 D, $P < 0.01$), a higher axial length/body

weight (0.168 ± 0.015 mm/g vs. 0.155 ± 0.002 mm/g, $P < 0.05$), and thinner choroidal

(−2.39 ± 1.33 mm vs. −0.12 ± 1.48 mm, $P < 0.01$) were observed compared to the PBS liposome

injection group (Fig. 1E), suggesting that clodronate liposome injection causes choroidal resident

macrophage depletion and thinning of the choroid, which leads to myopia regardless of whether

the choroid is developing.

Continuous intraperitoneal injection of LPS stimulates choroidal M1 macrophage

polarization and induces myopia in a murine model

Macrophages are classified into two categories based on their activation state and functions: M1

type (classically activated macrophages) and M2 type (alternatively activated macrophages),

which exhibit pro-inflammatory and anti-inflammatory properties, respectively (31). As the

upregulation of allergic inflammation has been implicated in the progression of myopia in animal

models (33), we investigated whether M1 macrophage polarization can induce myopia in a

mouse model. Mice were divided into LPS (10 mg/ml) and PBS injection groups for daily

intraperitoneal injections. To determine whether myopia had developed, refraction, axial length,

and choroidal thickness were measured using SD-OCT before the injections, 1 week after, and 2

weeks after the injections (Fig. 2A). After 1 week of LPS injection, LPS-injected mice showed

refractive shift (−9.21 ± 5.91 D vs. +2.91 ± 3.91 D, $P < 0.001$), rise in axial length/body weight

(0.28 ± 0.01 mm/g vs. 0.22 ± 0.005 mm/g, $P < 0.001$), and thinning of the choroid (−1.11 ± 0.90

mm vs. 1.36 ± 0.70 mm, $P < 0.001$), compared to mice injected with PBS for a week. The

injections were continued until the end of the 2 weeks, and LPS-injected mice maintained a

significantly higher refractive shift (−7.23 ± 4.36 D vs. +8.41 ± 7.01 D, $P < 0.001$), increase in the

ratio of axial length to mouse body weight (0.20 ± 0.005 mm/g vs. 0.17 ± 0.004 mm/g, $P < 0.01$),

and thinning of the choroid (−1.77 ± 1.02 mm vs. 2.05 ± 0.66 mm, $P < 0.001$), in comparison to

PBS-injected mice (Fig. 2B). Choroidal blood perfusion was measured by optical coherence

tomography angiography (OCTA), and the results showed that the choroidal blood perfusion of

mice injected with LPS for 2 weeks was reduced compared to that of mice injected with PBS

(37.36 ± 3.74 % area vs. 46.79 ± 4.74 % area, $P < 0.001$) (Fig. 2C). These data show that the

intraperitoneal injection of LPS can induce myopia in a mouse model. Choroid samples were

collected for real-time polymerase chain reaction (PCR) analysis to verify the polarization of M1

macrophages in the choroid stimulated by intraperitoneal injection of LPS. The expression levels

of the pro-inflammatory cytokines *Tnf* and *Il6* were significantly upregulated in LPS-injected

choroid samples compared to those in PBS-injected choroid samples (Fig. 2D). These data

indicate that the intraperitoneal injection of LPS polarizes M1 macrophages in the choroid,

promotes the release of inflammatory cytokines, and causes myopia in a mouse model. As

reported in LPS-treated macrophages, pro-inflammatory cytokines (TNFA, and IL6) secretion was

significantly attenuated by eicosapentaenoic acid (EPA), which can inhibit myopia progression

(42, 43). Therefore, an experiment was designed to administer an EPA diet to LPS-induced

myopic mice, and it was found that myopia did not progress (Supplementary Figs. 1A, B, C).

Taken together, these data suggest that the LPS-induced phenotypic switch toward M1

macrophages is associated with the development of myopia.

The constant intraperitoneal injection of IL-4/IL-13 encouraged the polarization of

choroidal M2 macrophages and suppressed myopia progression in a murine model of LIM

We found that intraperitoneal injection of LPS enhanced the expression of pro-inflammatory

cytokines in the choroid and promoted myopia development; contrarily, choroidal M2 macrophage

polarization should inhibit myopia development. Previous studies have demonstrated that IL-4

and IL-13, as anti-inflammatory interleukins, can directly activate M2 macrophages, which can be

identified by the expression of M2 macrophage markers, including mannose receptor 1 (Mrc1),

also referred to as CD206 and CD163 (44). Above all, to investigate the frequency of continuous

IL-4 injection, we compared the proportion of M2 macrophages in the choroid at 0 h, 24 h, and 48

h after being subjected to 0.1 µg/100 µl IL-4 intraperitoneal injection; F4/80 and CD11b-positive

cells were regarded as macrophages and CD206 was used as a marker to identify M2

macrophages. Flow cytometry analysis showed a growth in the ratio of macrophages (6.73 ± 0.22

% vs. 3.78 ± 0.41 %, $P < 0.001$) and M2 macrophages (43.57 ± 4.72 % vs. 30.33 ± 8.35 %, $P <$

0.05) at 48 h after IL-4 intraperitoneal injection, compared to 0 h (Fig. 3A). Real-time PCR

detection after IL-4 injection under the same conditions showed that *Mrc1* and *CD163* mRNA

expression in the choroid was enhanced 48 h after IL-4 injection (Fig. 3B). These results showed

that the optimal frequency of IL-4 injection is once every two days. Next, we examined whether

the promotion of M2 macrophage polarization by IL-4 injection on alternate days could inhibit

myopia progression in a LIM mouse model. All 3-weeks-old mice were subjected to LIM, in the

same manner as previously established (45), and administered with either PBS or 0.1 ug/100 ul

of IL-4 injection until 6 weeks of age (Fig. 3C). In the PBS injection group, eyes treated with −30

diopter (D) lenses showed refractive change (−11.52 ± 4.91 D vs. +12.45 ± 8.19 D, $P < 0.01$),

increased axial length (0.22 ± 0.01 mm vs. 0.18 ± 0.01 mm/g, $P < 0.05$), and decreased choroidal

thickness (−1.74 ± 0.66 mm vs. 3.91 ± 0.95 mm, $P < 0.001$) compared to eyes treated with 0D

lenses. In contrast, the IL-4 injection group showed smaller refractive change (−1.39 ± 2.91 D vs.

−11.52 ± 4.91 D, $P < 0.05$), smaller axial elongation (0.19 ± 0.01 mm vs. 0.22 ± 0.01 mm/g, $P <$

0.05), and thickening of the choroid (1.39 ± 0.19 mm vs. −1.74 ± 0.66 mm, $P < 0.001$) in eyes with

-30D lenses, compared to the PBS-injected group with −30D lenses (Fig. 3D). Meanwhile, to

demonstrate the influence of continuous IL-4 injection on choroidal blood perfusion, 3-weeks-old

mice were divided into three groups: control 0D group (both eyes were treated with 0D lenses),

control −30D group (binocular myopia induction), and IL-4 −30D group (binocular myopia

induction with IL-4 injection once every two days). OCTA was used to measure choroidal blood

perfusion at the beginning (3 weeks old) and end (6 weeks old) of myopic induction. The results

showed that choroidal blood perfusion was lower (8.59 ± 5.90 %area vs. −6.29 ± 8.73 %area, $P <$

0.01) after 3 weeks of myopia induction, compared to the control 0D group. Nevertheless, in the

IL4 −30D group, choroidal blood perfusion was improved (−6.29 ± 8.73 %area vs. 1.54 ± 9.34

%area, $P < 0.05$), compared to the control -30D group (Fig. 3E). To visualize the polarization of

M2 macrophages, after measuring choroidal blood perfusion at the end stage, choroid samples

were collected for flow cytometry, and data analysis revealed the corresponding growth in the

proportion of M2 macrophages (46.52 ± 0.05% vs. 35.71 ± 0.06%, $P < 0.05$) in the IL-4 injection

group with −30D lenses, compared to the PBS injection group with −30D lenses (Fig. 3F).

Collectively, these findings demonstrate that IL-4 injection can stimulate the polarization of M2

macrophages in the choroid and suppress the development of myopia by improving choroidal

blood perfusion in a LIM mouse model. To reinforce the conclusion that M2 macrophages can

inhibit myopia progression, we performed the same experiments with 0.1 μg /100 μl IL13 injection

instead of IL-4 injection, and the results also showed that M2 macrophages were polarized 48 h

after IL-13 injection (Supplementary Figs. 2A, B). The results of the LIM mouse model combined

with IL-13 injection were as expected; IL-13 also polarized M2 macrophages, suppressed myopia

progression, and improved choroidal blood perfusion in a murine model of LIM (Supplementary

Figs. 2C, D, E, F).

Berberine, an eye drop that polarizes M2 macrophages, inhibits myopia progression after consistent use

After realizing that M2 macrophage polarization may inhibit the development of myopia, we used

an M2 macrophage-activating drug as an eye drop to verify its effect on the development of

myopia. Some studies have shown that berberine promotes the polarization of M1 macrophages

to the M2 anti-inflammatory phenotype in mice (46, 47). To verify the role of berberine eye drops

in the development of myopia, we examined and compared changes in refraction, axial length,

and choroidal thickness in 3-weeks-old C57BL/6 J mice after myopia induction for 3 weeks and

the administration of either berberine or dimethyl sulfoxide (DMSO) eye drops. The control −30D

group (received −30D lenses for both eyes) and 2% DMSO −30D group (binocular myopia

induction with the application of 2% DMSO eye drops daily) showed a large refractive change (P

< 0.001), increased axial length elongation ($P < 0.05$), and decreased choroidal thickness ($P <$

0.001), compared to the control 0D group (received 0D lenses for both eyes). In addition, the

berberine −30D group (binocular myopia induction with daily 0.26 mg/ml eye drops) exhibited

smaller refractive changes (2.12 ± 1.63 D vs. −5.25 ± 2.89 D, $P < 0.001$), smaller axial elongation

(0.19 ± 0.02 mm vs. 0.22 ± 0.04 mm/g, $P < 0.05$), and a relatively thicker choroid (0.79 ± 0.64 mm

vs. −1.56 ± 0.76 mm, $P < 0.001$), compared to the 2% DMSO −30D group. These results indicate

that drugs used to activate the polarization of M2 macrophages may also have the potential to

inhibit the development of myopia (Supplementary Figs. 3A, B, C).

Effects of LPS and IL-4 injections on nicotinamide adenine dinucleotide phosphate (NADPH) oxidation activity

We inferred that the direction of macrophage polarization may influence the development or

inhibition of myopia through its pro-inflammatory or anti-inflammatory properties. However, the

other mechanisms underlying this association remain largely unknown. LPS-treated M1

macrophages are associated with high levels of reactive oxygen and nitrogen species (ROS and

RNS, respectively) and matrix metalloproteinase (MMP) activity, and ROS production is critical for

M1 macrophage activation and function (48, 49). Correspondingly, ROS overproduction may

impair blood perfusion to the retina, and circulatory disturbances during the development of

myopia can lead to oxidative stress (50, 51). To investigate whether M1 macrophage polarization

also induces myopia by promoting ROS and oxidative stress response, and whether M2

macrophage polarization suppressed myopia progression by inhibiting ROS release and oxidative

stress response, the mRNA expression of NADPH oxidase 2 (*Nox2*), which is the most important

source of ROS, and the expression of *Nox4, Mmp2*, and *Mmp9* in the choroid were observed

after either LPS or IL-4 injection (Fig. 4A). Quantitative analysis showed increased *Nox2, Nox4,*

Mmp2, and *Mmp9* mRNA expression in the LPS injection group compared to that in the PBS

injection group (Fig. 4B). In contrast, 48 h after IL-4 injection, the mice showed lower *Nox2,*

Mmp2, and *Mmp9* mRNA expression (Fig. 4C). Since 8-hydroxy-2-deoxyguanosine (8-OHdG) is

an effective marker for assessing oxidative DNA damage (52), we investigated the level of 8-

OHdG in the choroid after intraperitoneal injection of either LPS or IL-4 and found that the IL-4

injection group showed a reduction in choroidal 8-OHdG concentration compared to the control

group ($P < 0.01$). In contrast, the concentration of choroidal 8-OHdG was higher in the LPS

injection group, compared to the IL-4 group ($P < 0.05$) (Fig. 4D). Taken together, these data

suggest that polarized M1 macrophages may induce myopia by promoting oxidative stress,

whereas polarized M2 macrophages may inhibit myopia progression by suppressing the oxidative

stress response.

Discussion

In this paper, we have found that M1 macrophage accelerates myopia while M2 macrophage

suppressed myopia progression, this statement is supported by the data performed and

described. Since the choroid is now considered to be the most important tissue for the

development of myopia and maintenance of the choroidal vasculature is critical for the

suppression of myopia. Our findings on the role of macrophages are indeed a novel concept and

open up new avenues for further investigation.

Macrophages are tissue-resident or infiltrating immune cells that are present in all tissues. In

addition to their immune properties, macrophages play an important role in regulating normal

tissue development and homeostasis (14). Researchers have found that resident macrophages

are densely distributed in the choroid tissue, a connective tissue rich in pigments and blood

vessels that supply blood and nutrients to the outer retina (16, 53). However, the precise role of

choroidal-resident macrophages has rarely been directly studied. Recently, choroidal

macrophages have been suggested to perform an underappreciated trophic function by

preserving the vascular structure of the choroid (24). In the present study, we confirmed that

choroidal thinning occurred following depleted choroidal-resident macrophages and was

accompanied by axial length elongation and a significant refractive shift. This unexpected finding

supports the previously unknown concept that macrophages in the developing choroid are

associated with choroidal growth and myopia.

The ability of resident immune cells to maintain the vasculature changes depending on the

developmental state of the tissue (24). Microglia, as specialized macrophages, play an important

role in shaping the central nervous system (CNS) vasculature during development but are not

necessary for the maintenance of the mature uninjured CNS (17, 54). The choroidal thickness

steadily increases and reaches a steady state at 7–9 weeks of age; thereafter, it gradually

decreases with age (41). In our experiments, we used 8-weeks-old mice with a plateau in

choroidal thickness, depleted choroidal-resident macrophages, and found that the choroid

became thinner and myopia still occurred. These results demonstrate that choroidal

macrophages effectively support choroidal growth during development and maintain choroidal

stability after maturation.

Studies of the normal choroid in mice have revealed a network of F4/80$^+$ (CD160) and CD169$^+$

macrophages widely distributed around choroidal blood vessels (55). F4/80 is one of the most

specific cell surface markers of murine macrophages, and CD169$^+$ macrophages are a unique

subset of macrophages involved in multiple immune responses (56, 57). Activated macrophages

can be broadly divided into the M1 and M2 cell subsets, representing the two extremes of the

macrophage activation spectrum that can occur *in vivo* (58). M1 macrophage polarization occurs

in many ocular inflammatory diseases and is accompanied by the production of pro-inflammatory

cytokines; these inflammatory reactions induce local biochemical reactions, leading to local tissue

remodeling and disease progression. Numerous studies have shown that inflammatory diseases

of the eye can increase the risk of myopia. Myopic refractive errors are significantly increased in

patients with juvenile chronic arthritis (JCA)-related uveitis, and inflammatory cytokines IL6 and

TNFA were found to be overexpressed in the aqueous humor of uveitis patients (9, 59). Multifocal

choroiditis (MFC) was definitely associated with myopia, and statistical analysis of 41 MFC

patients showed that their average refractive error was -2.19 diopters (60). These findings

suggest that acute or chronic systemic inflammatory diseases increase the risk of myopia. In this

study, we demonstrated that LPS injection stimulates the expression of pro-inflammatory

cytokines in the choroid, reduces choroidal thickness, and induces myopia in a murine model.

This study suggests that the pro-inflammatory properties of M1 macrophages may contribute to

the pathogenesis of myopia.

In contrast, some anti-inflammatory substances have been shown to inhibit the development of

myopia when administered orally or as eye drops. Atropine eye drops are currently the most

effective therapy for myopia control and have been shown to be associated with effective

downregulation of eye inflammation (12, 61). Resveratrol eye drops reduced inflammation in a

form-deprivation myopia (FDM) animal model to prevent myopia development (62). In oral

preparations, the oral administration of lactoferrin (LF), which has anti-inflammatory properties,

inhibited myopia progression in a LIM mouse model (63). However, no studies have shown how

the polarization of M2 macrophages with anti-inflammatory properties directly affects the

development of myopia. Based on the highly plastic nature of macrophages, we administered IL-4

and IL-13 injections to induce the polarization of M2 macrophages in the choroid and found that

the expression of anti-inflammatory cytokines was activated in the choroid and inhibited the

progression of myopia. Taken together, these results indicate that direct stimulation of M1

macrophages to release inflammatory factors induces inflammatory reactions, leading to

accelerated myopia, which can be reversed by direct stimulation of M2 macrophages to release

anti-inflammatory factors.

We also found that with the activation of M2 macrophages, choroid blood perfusion, and choroid

thickness also increased correspondingly, this indicates M2 macrophages in the choroid play a

crucial role in choroidal thickness and maintenance of blood perfusion. This is consistent with its

property of maintaining the stability of the vascular structure. Under a physiological state, M2

macrophages contribute to endothelial barrier protection and facilitate vascular stability and

maturation (64, 65)**.** In a pathological condition, ameliorating macrophage inflammation or

polarizing it into the M2 phenotype can prevent the rupture of intracranial aneurysms (66). M2

macrophage polarization in the aneurysmal aortic wall heals the aortic wall and maintains

vascular integrity as the disease progresses (67). The ability of M2 macrophages to maintain

vascular stability provides strong evidence that it can maintain the stability and overall

morphology of the choroid vascular structure and thus against the development of myopia.

In addition to influencing myopia development through their anti-inflammatory or maintaining

vascular structural stability properties, M2 macrophages may also inhibit the progression of

myopia through their ability to inhibit oxidative stress. In contrast, the mechanism of myopia

induced by M1 macrophage polarization may be related to the activation of oxidative metabolism.

Polarized M1 macrophages produce reactive oxygen species (ROS) and drive inflammatory

responses, leading to the production of high levels of nitric oxide (NO) and NO synthase (iNOS);

excessive ROS can cause oxidative stress and damage biomolecules. Oxidative stress

upregulates the expression of matrix metalloproteinases-2 and -9 (MMP-2 and MMP-9) (68-71).

M2 macrophages have the opposite effects (72). Our study demonstrated that stimulating the

polarization of M1 macrophages in the choroid induces the expression of NADPH oxidases,

matrix metalloproteinase, and 8-OHdG while stimulating the polarization of M2 macrophages

results in the exact opposite effect. Moreover, there is evidence that oxidative stress plays a role

in the development of myopia-associated disorders at the cellular level (51). In pathological

myopia, retinal hypoxia and ischemia lead to the accumulation of ROS and insufficient ability of

the endogenous antioxidant system to clear ROS, which leads to an increase in hypoxia-inducible

factor-1 (HIF-1) levels and further promotes the development of myopia (51, 73). N-omega-nitro-

L-arginine methyl ester (L-NAME), an inhibitor of NOS, was injected intravitreally to prevent the

development of myopia in both form-deprivation myopia (FDM) and LIM animal models (74, 75).

In addition to its anti-inflammatory properties, LF can reverse high levels of MMP-2 activity in the

LIM model, thereby inhibiting myopia (63). Additionally, research has indicated that the MMP-9

gene locus may contribute to myopia (76). Combined with our results, the role of macrophages in

oxidative stress response may be a key factor affecting the development of myopia.

Currently, studies have shown that some natural compounds or supplements that modulate M1 to

M2 macrophages have therapeutic effects on specific diseases. Lupeol, as a multi-target agent,

ameliorates experimental inflammatory bowel disease by converting M1 to M2 macrophages (77).

Resveratrol through its anti-oxidant capacity and anti-inflammatory properties inhibits platelet

aggregation, thereby preventing atherosclerosis (78). Lactobacillus murinus alleviate intestinal

ischemia/reperfusion damage by encouraging M2 macrophages to secrete interleukin-10 (79). In

addition to all these, there are some compounds that treat ophthalmology-related diseases by

their properties of promoting the transformation of M1 into M2 macrophages. Quercetin may have

an impact on the treatment of ophthalmic diseases such as conjunctivitis, dry eye, and

retinopathy due to its anti-inflammatory and antioxidant properties (80). Forskolin is a medication

that can decrease intraocular pressure and be used as an anti-glaucoma agent (81). According to

the properties of various natural compounds, which are involved in regulating the phenotype

transformation of macrophages from M1 to M2, we targeted the agent named berberine, made it

into eye drops, and applied it in myopic-induced mice, we found that it has the ability to inhibit the

development of myopia. It indicates a possibility that control of macrophage phenotype may be a

new therapy applicable to myopia treatment.

In summary, our results have important implications for understanding the relationship between

macrophages and the progression of myopia. In addition to the notion that the two modes of

macrophage polarization can have specific effects on choroidal structure as well as the

development of myopia, our findings show that the effect of macrophages on myopia

development may be related to inflammation, vascular maintenance, and oxidative stress

response, and these possible co-existing mechanisms may play a key role in the development of

myopia. The above views provide new ideas for regulating the macrophage response to

interventions in myopia and provide a new method for drug screening for myopia treatment.

Materials and Methods

Mice

The Ethics Committee on Animal Research of the Keio University School of Medicine approved

all procedures, which followed the Association for Research in Vision and Ophthalmology

Statement for the Use of Animals in Ophthalmic and Vision Research, Keio University's

Institutional Guidelines on Animal Experimentation, and the Animal Research: Reporting of In

Vivo Experiments (ARRIVE) guidelines. In addition, this study implemented the principle of

random assignment. CLEA Japan, Inc. provided the wild-type male C57BL/6J mice. Those mice

were kept in standard transparent mouse cages (29×18×13 cm) with four or five per cage in

pathogen-free environments maintained at $23 \pm 3\,°C$, under approximately 50 lux of background

fluorescent lamp light (color temperature: 5000 K), with a 12-hour diurnal period and constant

access to normal chow (MF, Oriental Yeast Co., Ltd, Tokyo, Japan) and tap water. All the animals

were randomly allocated.

LIM mouse model

A murine LIM model was created as previously described (45). Briefly, mice were placed under

general anesthesia using a mixture of midazolam (Sandoz K.K., Tokyo, Japan), medetomidine

(Domitor®, Orion Corporation, Espoo, Finland), and butorphanol tartrate (MMB) (Meiji Seika

Pharma Co., Ltd., Tokyo, Japan). Eyeglass frames that fit the mouse's head's contours were

designed and printed using a three-dimensional printer. For myopia induction, a negative 30D

lens was constructed using polymethyl methacrylate (PMMA). The left and right eyes of the

spectacles were adjusted to fit the shape of the mouse skull frame and screwed onto a stick,

which was a joint enabling the left and right frame positions to be altered or removed for cleaning.

Subsequently, the stick was bonded to the mouse skull using a self-curing dental adhesive

system. Induction began once the mouse had fully recovered from anesthesia, and the lenses

were removed for cleaning at least twice a week.

The measurement of refraction, axial length, and choroidal thickness

Refraction was measured using an infrared photorefractor (Steinbeis Transfer Center, Stuttgart,

Baden-Württemberg, Germany) as previously reported (45). After measuring refraction, axial

length, and choroidal thickness were measured using an SD-OCT system (Envisu R4310, Leica

Microsystems, Wetzlar, Germany). For each mouse, measurements of refraction, axial length,

and choroidal thickness were performed before the start of LIM (0W) and at the end of LIM (3W).

Before the measurements, the mouse's eyes were treated with mydriasis eye drops containing

0.5% tropicamide and 0.5% phenylephrine (Santen Pharmaceutical Co., Ltd., Osaka, Japan) to

ensure mydriasis and cycloplegia. After pupil dilation, the mice were subjected to general

anesthesia using MMB. It is important to prevent the occurrence of corneal injury during

measurement. Axial length was measured as the vertical distance from the anterior corneal

surface to the retinal pigment epithelium layer near the optic nerve (45). Choroidal thickness was

determined by quantifying the circular area of the disc on the posterior surface of the choroid

using ImageJ software (82).

The measurement of choroidal blood perfusion

Choroidal blood perfusion was measured at the beginning (3-week-old) and end (6-week-old)

stages of myopia induction using an SS-OCT/OCTA device (XEPHILIO OCT-S1, CANON

Medical Systems, Tokyo, Japan). The measurement method has been described in a previous

report (83). Briefly, as with refraction measurements, the mice were dilated under general

anesthesia before measurement. Choroidal blood perfusion was measured using en face

angiography to identify the optic nerve as the central area. The choroidal blood perfusion signal

was obtained from B-scan images at the corresponding location. In the B-scan images, the area

covered by red noise points was the area without blood perfusion. ImageJ quantitative analysis

was used to calculate the non-blood perfusion area in the choroid and evaluate the percentage of

the area with blood perfusion in the choroid.

Flow cytometry

The mice were administered an overdose of MMB to induce profound anesthesia before being

euthanized by cervical dislocation. The eyes were immediately enucleated and the anterior

segment, vitreous, and retina were discarded. Choroid tissues were carefully scraped off the

sclera-choroid complex, collected in the tube (10 choroid tissues/tube), and then processed in

digestion buffer (0.75 mg/ml of collagenase A, from FUJIFILM, Tokyo, Japan) in complete

Dulbecco's Modified Eagle Medium (DMEM) solution for 45 min at 37 °C. The digested choroid

mixture was transferred onto the cell strainers and any undigested choroid pieces were mashed

using the end of a syringe plunger while adding 5 ml cold complete DMEM to allow the pass-

through of cells into new 50 ml tubes below the strainer. The aggregated choroidal cells were

dissociated into single-cell suspensions via differential centrifugation. The cells were washed by

adding 700 ul of flow cytometry staining (FACS) buffer (0.1 g of bovine serum albumin (BSA) and 80 ul of 0.5 mol/L ethylenediaminetetraacetic acid (EDTA) in 20 ml of PBS (pH 7.4) solution to the cell suspension. The pellet was resuspended with 10 ul Fc block solution (1:10 Fc block) (BD Biosciences, New Jersey, USA) for 10 min, stained with a mixture of fluorochrome-conjugated antibodies (1:200 FITC anti-mouse/human CD11b M1/70, BioLegend, California, USA), (1:200 APC/Cyanine7 anti-mouse F4/80, BioLegend, California, USA), (1:200 CD206, AbD Serotec, Kidlington, near Oxford, UK) for at least 45 min, and finally covered with 0.1 mg/ml Hoechst (DOJINDO, Amsterdam, Netherlands) for 10 min. All incubations were performed on ice and light was avoided. Data were acquired on a CytoFLEX S flow cytometer using CytExpert software (Beckman Coulter Life Sciences, Inc., Indiana, United States), and offline data analysis was performed using CyExpert software.

Clodronate liposome treatment

Clodronate liposome (Liposoma BV, Amsterdam, The Netherlands) was intraperitoneally injected at a dose of 0.10 ml/10 g (once every two days) to C57BL/6J mice from P21 to P28 or from P56 to P63. The control group was injected with an equal amount of PBS liposomes, as described above. Refraction, axial length, and choroidal thickness were measured before and after 4 four injections. The mice were then euthanized and choroid samples were collected.

LPS treatment

Lipopolysaccharides from *Escherichia coli* O111:B4 (Merck, Tokyo, Japan) were dissolved in PBS and injected intraperitoneally to induce systemic M1 macrophage polarization. The concentration of LPS was 10 mg/ml, and the injection dose was increased proportionally to the body weight of the mice, with 10 µg of LPS given per gram of mice. Since LPS injections caused a large reduction in mouse weight, we monitored and documented the mouse body weight before each injection. In addition, a corresponding volume of PBS was injected intraperitoneally as a vehicle control.

EPA administration

EPA-supplemented chow containing 5% of EPA (EPADEL®, Mochida, Pharmaceutical Co. Ltd.,

Tokyo, Japan) and normal chow were prepared as previously reported (43). Briefly, EPA ethyl

ester was blended with powdered ordinary chow, and tap water was added. The mixture was

shaped into tiny cylinders and dried for consumption. Feeding with EPA-mixed chow began and

ended with myopia induction.

IL-4/IL-13 treatment

To induce systemic M2 macrophage polarization, recombinant murine IL-4 (214-14) and IL-13

(210-13) (PeproTech, New Jersey, United States) were dissolved in PBS at the same

concentration of 0.1 μg/100 μl, and injected intraperitoneally into the mice once every two days.

The control group was intraperitoneally administered the same volume of PBS. The injections

were administered during the induction of myopia. To ensure that the mice grew normally, their

body weights were measured and recorded prior to each injection.

Berberine chloride eye drops

Berberine chloride hydrate (TCI, Tokyo, Japan) solution was freshly prepared in DMSO. When

determining the concentration of berberine chloride eye drops, we found that berberine chloride

hydrate could not be dissolved in water at room temperature. We attempted to use 100% DMSO

to dissolve it and found that the solubility of berberine chloride hydrate in 100% DMSO at room

temperature was approximately 13 mg/ml. At present, the concentration of berberine chloride eye

drops is 0.26 mg/ml. In addition, DMSO, a dissolution vector, has been reported to exert cytotoxic

effects at certain concentrations. Therefore, an untreated control group should be included along

with the DMSO vehicle control to check for solvent toxicity (84). To maintain the concentration of

berberine chloride hydrate at 0.26 mg/ml and reduce the concentration of DMSO as much as

possible, the final concentration of berberine chloride eye drops was 0.26 mg/ml, dissolved in 2%

DMSO. Moreover, a parallel control group of 2% DMSO was also set up in mice to observe its

effect on myopia.

Real-time Quantitative PCR (qPCR)

After the mice were euthanized, the eyeballs were enucleated immediately to collect retinal,

choroid, and scleral tissues, frozen in liquid nitrogen, and stored at -80 °C for later use. Mouse

total RNA was extracted from the retina, choroid, and scleral tissues using TRIzol (TRI reagent)

(MOR, Miami, FL, USA #TR118). Small RNAs were isolated using RWT (QIAGEN, Hilden,

Germany, #1067933) and RPE buffers (QIAGEN, Hilden, Germany, #1018013). The RNA

samples were dissolved in RNase-free water (TAKARA HOLDINGS Inc., Kyoto, Japan, 9012)

and measured with a spectrophotometer (NanoDrop; Thermo Fisher Scientific, Waltham, MA,

USA). According to the manufacturer's instructions, the extracted RNA was converted to cDNA

after RNA denaturation, DNase reaction, and reverse transcription. mRNA gene expression was

determined using SYBR Green RT-PCR on the cDNA template and PCR was performed using

StepOnePlus Real-Time PCR System (Applied Biosystems, Waltham, Massachusetts, USA). The

mRNA expression of each gene was evaluated using the comparative Ct (ΔΔCT) method and

normalized to glyceraldehyde-3-phosphate dehydrogenase (GAPDH) mRNA expression, which

was used as the reference gene. The following were the qPCR primer sequences:

mouse *Il6* forward: CTACCCCAATTTCCAATGCT

mouse *Il6* reverse: ACCACAGTGAGGAATGTCCA

mouse *Tnfa* forward: CTGTAGCCCACGTCGTAGC

mouse *Tnfa* reverse: TTGAGATCCATGCCGTTG

mouse Mrc1 forward: TCGAGACTGCTGCTGAGTCCA

mouse Mrc1 reverse: AGACAGGATTGTCGTTCAACCAAAG

mouse *Nox2* forward: ACTCCTTGGAGCACTGG

mouse *Nox2* reverse: GTTCCTGTCCAGTTGTCTTCG

mouse *Nox4* forward: TGAACTACAGTGAAGATTTCCTTGAAC

mouse *Nox4* reverse: GACACCCGTCAGACCAGGAA

mouse *Mmp2* forward: CAAGTTCCCCGGCGAT

mouse *Mmp2* reverse: TTCTGGTCAAGGTCAC

mouse *Mmp9* forward: GGACCCGAAGCGGACA

mouse *Mmp9* reverse: CGTCGTCGAAATGGGC

mouse *GAPDH* forward: AGGAGCGAGACCCCACTAAC

mouse *GAPDH* reverse: GATGACCCTTTTGGCTCCAC

8-OHdG ELISA

For assessing oxidative stress, the level of 8-Oxo-2'-deoxyguanosine (8-OHdG) in the choroid

tissues was measured using commercial Enzyme-Linked Immunosorbent Assay (ELISA) kits in

accordance with the manufacturer's instructions (Uscn Life Science; Wuhan, China). Because

this assay uses a competitive inhibition enzyme immunoassay approach, the concentration of 8-

OHdG in the sample and the assay signal strength were inversely correlated. The concentration

of 8-OHdG in each sample was calculated using standard curves generated from standard

proteins, which were calculated and analyzed using Boster's ELISA Online Calculator.

Statistical analyses

Our study used an independent *t*-test and one-way analysis of variance (ANOVA) for statistical

significance analyses of all data (Microsoft Excel 2003, USA). All results are presented as mean

standard deviation (SD), and results with *P* values < 0.05 were considered statistically significant.

Study Approval

The Institutional Animal Care and Use Committee at Keio University authorized all operations

(approval number: 16017). All methods and experimental protocols followed the National

Institutes of Health (NIH) guidelines for working with laboratory animals and the ARVO Animal

Statement for the Use of Animals in Ophthalmic and Vision Research.

Data Availability

All sources of data generated or analyzed during this study are included in this published article

and its supplementary information files.

Author Contributions

Conceptualization, J.H., S.I., T.K., and K.T.; Methodology, J.H., and S.I.; Formal Analysis, J.H.;

Investigation, J.H.; Data Curation, J.H.; Project administration, J.H., S.I. and T. K.; Writing –

Original Draft Preparation, J.H.; Writing – Review & Editing, S.I., K.M., J.H., H.T., K.N., T.K., and

K.T.; Supervision, K.N., T.K., and K.T. All authors made a substantial contribution in the revision

of the manuscript.

Acknowledgments

The authors thank Y. Katada, N. Ban, A. Nakai, N. Serizawa, C. Shoda, M. Ziyan, C. Junhan, and

A. Kawabata (Graduate School of Medicine, Keio University, Tokyo, Japan) for their technical

guidance and administrative support.

References

626   1.   Hawthorne FA, and Young TL. Genetic contributions to myopic refractive error: Insights
627   from human studies and supporting evidence from animal models. *Experimental eye*
628   *research.* 2013;114:141-9.
629   2.   Lopes MC, Andrew T, Carbonaro F, Spector TD, and Hammond CJ. Estimating heritability
630   and shared environmental effects for refractive error in twin and family studies.
631   *Investigative ophthalmology & visual science.* 2009;50(1):126-31.
632   3.   Morgan IG, Ohno-Matsui K, and Saw S-M. Myopia. *The Lancet.* 2012;379(9827):1739-48.
633   4.   Wojciechowski R. Nature and nurture: the complex genetics of myopia and refractive
634   error. *Clinical genetics.* 2011;79(4):301-20.
635   5.   Goldschmidt E, and Jacobsen N. Genetic and environmental effects on myopia
636   development and progression. *Eye.* 2014;28(2):126-33.
637   6.   Lim LS, Gazzard G, Low Y-L, Choo R, Tan DT, Tong L, et al. Dietary factors, myopia, and
638   axial dimensions in children. *Ophthalmology.* 2010;117(5):993-7. e4.

7.   Holden BA, Fricke TR, Wilson DA, Jong M, Naidoo KS, Sankaridurg P, et al. Global
prevalence of myopia and high myopia and temporal trends from 2000 through 2050.
Ophthalmology. 2016;123(5):1036-42.
8.   Yotsukura E, Torii H, Inokuchi M, Tokumura M, Uchino M, Nakamura K, et al. Current
prevalence of myopia and association of myopia with environmental factors among
schoolchildren in Japan. *JAMA ophthalmology.* 2019;137(11):1233-9.
9.   Fledelius H, Zak M, and Pedersen FK. Refraction in juvenile chronic arthritis: a long‐
term follow‐up study, with emphasis on myopia. *Acta Ophthalmologica Scandinavica.*
2001;79(3):237-9.
10.   Herbort CP, Papadia M, and Neri P. Myopia and inflammation. *Journal of ophthalmic &*
vision research. 2011;6(4):270.
11.   Kamath Y, Singh A, Bhat S, and Sripathi H. Acute onset myopia as a presenting feature of
systemic lupus erythematosus. *Journal of Postgraduate Medicine.* 2013;59(3):245.
12.   Lin H-J, Wei C-C, Chang C-Y, Chen T-H, Hsu Y-A, Hsieh Y-C, et al. Role of chronic
inflammation in myopia progression: clinical evidence and experimental validation.
EBioMedicine. 2016;10:269-81.
13.   Fujiwara N, and Kobayashi K. Macrophages in inflammation. *Current Drug Targets-*
Inflammation & Allergy. 2005;4(3):281-6.
14.   Wynn TA, Chawla A, and Pollard JW. Macrophage biology in development, homeostasis
and disease. *Nature.* 2013;496(7446):445-55.
15.   Epelman S, Lavine KJ, and Randolph GJ. Origin and functions of tissue macrophages.
Immunity. 2014;41(1):21-35.
16.   Chinnery HR, McMenamin PG, and Dando SJ. Macrophage physiology in the eye.
Pflügers Archiv - European Journal of Physiology. 2017;469(3):501-15.
17.   Wang X, Zhao L, Zhang J, Fariss RN, Ma W, Kretschmer F, et al. Requirement for
microglia for the maintenance of synaptic function and integrity in the mature retina.
Journal of Neuroscience. 2016;36(9):2827-42.
18.   Santos AM, Calvente R, Tassi M, Carrasco MC, Martín‐Oliva D, Marín‐Teva JL, et al.
Embryonic and postnatal development of microglial cells in the mouse retina. *Journal of*
Comparative Neurology. 2008;506(2):224-39.
19.   Rathnasamy G, Foulds WS, Ling E-A, and Kaur C. Retinal microglia–A key player in
healthy and diseased retina. *Progress in neurobiology.* 2019;173:18-40.
20.   Yu C, Roubeix C, Sennlaub F, and Saban DR. Microglia versus monocytes: distinct roles in
degenerative diseases of the retina. *Trends in neurosciences.* 2020;43(6):433-49.
21.   Forrester JV, McMenamin PG, Holthouse I, Lumsden L, and Liversidge J. Localization and
characterization of major histocompatibility complex class II-positive cells in the
posterior segment of the eye: implications for induction of autoimmune uveoretinitis.
Investigative ophthalmology & visual science. 1994;35(1):64-77.
22.   Kumar A, Zhao L, Fariss RN, McMenamin PG, and Wong WT. Vascular associations and
dynamic process motility in perivascular myeloid cells of the mouse choroid:
implications for function and senescent change. *Investigative ophthalmology & visual*
science. 2014;55(3):1787-96.
23.   Yang X, Zhao L, Campos MM, Abu-Asab M, Ortolan D, Hotaling N, et al. Choroid-resident
macrophages maintain local vasculature and RPE integrity and spontaneously
regenerate following depletion. *bioRxiv.* 2019:868430.

24. Yang X, Zhao L, Campos MM, Abu-Asab M, Ortolan D, Hotaling N, et al. CSF1R blockade
induces macrophage ablation and results in mouse choroidal vascular atrophy and RPE
disorganization. *Elife.* 2020;9:e55564.
25. Nishida Y, Fujiwara T, Imamura Y, Lima LH, Kurosaka D, and Spaide RF. Choroidal
thickness and visual acuity in highly myopic eyes. *Retina.* 2012;32(7):1229-36.
26. Liu Y, Wang L, Xu Y, Pang Z, and Mu G. The influence of the choroid on the onset and
development of myopia: from perspectives of choroidal thickness and blood flow. *Acta
Ophthalmologica.* 2021;99(7):730-8.
27. Flores-Moreno I, Lugo F, Duker JS, and Ruiz-Moreno JM. The relationship between axial
length and choroidal thickness in eyes with high myopia. *American journal of
ophthalmology.* 2013;155(2):314-9. e1.
28. Zhang S, Zhang G, Zhou X, Xu R, Wang S, Guan Z, et al. Changes in choroidal thickness
and choroidal blood perfusion in guinea pig myopia. *Investigative Ophthalmology &
Visual Science.* 2019;60(8):3074-83.
29. Huang X, Li Y, Fu M, and Xin H-B. *Macrophages*. Springer; 2018:119-26.
30. Ivashkiv LB. Epigenetic regulation of macrophage polarization and function. *Trends in
immunology.* 2013;34(5):216-23.
31. Mosser DM, and Edwards JP. Exploring the full spectrum of macrophage activation.
Nature reviews immunology. 2008;8(12):958-69.
32. Sica A, and Mantovani A. Macrophage plasticity and polarization: in vivo veritas. *The
Journal of clinical investigation.* 2012;122(3):787-95.
33. Wei C-C, Kung Y-J, Chen CS, Chang C-Y, Lin C-J, Tien P-T, et al. Allergic conjunctivitis-
induced retinal inflammation promotes myopia progression. *EBioMedicine.*
2018;28:274-86.
34. Zhu K, Yang C, Dai H, Li J, Liu W, Luo Y, et al. Crocin inhibits titanium particle-induced
inflammation and promotes osteogenesis by regulating macrophage polarization.
International Immunopharmacology. 2019;76:105865.
35. Talamonti E, Pauter AM, Asadi A, Fischer AW, Chiurchiù V, and Jacobsson A. Impairment
of systemic DHA synthesis affects macrophage plasticity and polarization: implications
for DHA supplementation during inflammation. *Cellular and Molecular Life Sciences.*
2017;74(15):2815-26.
36. Namgaladze D, and Brüne B. Macrophage fatty acid oxidation and its roles in
macrophage polarization and fatty acid-induced inflammation. *Biochimica et Biophysica
Acta (BBA)-Molecular and Cell Biology of Lipids.* 2016;1861(11):1796-807.
37. Hung L-F, Wallman J, and Smith EL. Vision-dependent changes in the choroidal thickness
of macaque monkeys. *Investigative Ophthalmology & Visual Science.* 2000;41(6):1259-
69.
38. NICKLA DL, WILDSOET C, and WALLMAN J. Visual influences on diurnal rhythms in ocular
length and choroidal thickness in chick eyes. *Experimental eye research.* 1998;66(2):163-
81.
39. Read SA, Alonso-Caneiro D, Vincent SJ, and Collins MJ. Longitudinal changes in choroidal
thickness and eye growth in childhood. *Investigative ophthalmology & visual science.*
2015;56(5):3103-12.
40. Weisser SB, van Rooijen N, and Sly LM. Depletion and reconstitution of macrophages in
mice. *JoVE (Journal of Visualized Experiments).* 2012(66):e4105.

41. Zhang Y, Jeong H, Mori K, Ikeda S-I, Shoda C, Miwa Y, et al. Vascular endothelial growth
factor from retinal pigment epithelium is essential in choriocapillaris and axial length
maintenance. *PNAS Nexus.* 2022;1(4):pgac166.
42. Mullen A, Loscher CE, and Roche HM. Anti-inflammatory effects of EPA and DHA are
dependent upon time and dose-response elements associated with LPS stimulation in
THP-1-derived macrophages. *The Journal of nutritional biochemistry.* 2010;21(5):444-50.
43. Mori K, Kuroha S, Hou J, Jeong H, Ogawa M, Ikeda Si, et al. Lipidomic analysis revealed
n‐3 polyunsaturated fatty acids suppressed choroidal thinning and myopia progression
in mice. *The FASEB Journal.* 2022;36(6):e22312.
44. Yao Y, Xu X-H, and Jin L. Macrophage polarization in physiological and pathological
pregnancy. *Frontiers in immunology.* 2019;10:792.
45. Jiang X, Kurihara T, Kunimi H, Miyauchi M, Ikeda S-i, Mori K, et al. A highly efficient
murine model of experimental myopia. *Scientific reports.* 2018;8(1):1-12.
46. Lin J, Cai Q, Liang B, Wu L, Zhuang Y, He Y, et al. Berberine, a traditional Chinese
medicine, reduces inflammation in adipose tissue, polarizes M2 macrophages, and
increases energy expenditure in mice fed a high-fat diet. *Medical Science Monitor:*
International Medical Journal of Experimental and Clinical Research. 2019;25:87.
47. WANG Q-Z, SHI J, LIU Q, YANG L-X, GUO L, YE C-Y, et al. Berberine promotes M1
proinflammatory phenotype to M2 anti-inflammatory phenotype polarization in
macrophage cell line RAW264. 7. *Basic & Clinical Medicine.* 2019;39(5):646.
48. Virág L, Jaén RI, Regdon Z, Boscá L, and Prieto P. Self-defense of macrophages against
oxidative injury: Fighting for their own survival. *Redox Biology.* 2019;26:101261.
49. Covarrubias A, Byles V, and Horng T. ROS sets the stage for macrophage differentiation.
Cell research. 2013;23(8):984-5.
50. Coviltir V, Burcel M, Cherecheanu AP, Ionescu C, Dascalescu D, Potop V, et al. Update on
myopia risk factors and microenvironmental changes. *Journal of ophthalmology.*
2019;2019.
51. Francisco B-M, Salvador M, and Amparo N. Oxidative stress in myopia. *Oxidative*
Medicine and Cellular Longevity. 2015;2015.
52. Halliwell B. Why and how should we measure oxidative DNA damage in nutritional
studies? How far have we come? *The American journal of clinical nutrition.*
2000;72(5):1082-7.
53. McMenamin PG, Saban DR, and Dando SJ. Immune cells in the retina and choroid: Two
different tissue environments that require different defenses and surveillance. *Progress*
in retinal and eye research. 2019;70:85-98.
54. Arnold T, and Betsholtz C. Erratum to: The importance of microglia in the development
of the vasculature in the central nervous system. *Vascular cell.* 2013;5(1):1-7.
55. McMenamin PG. Dendritic cells and macrophages in the uveal tract of the normal
mouse eye. *British Journal of Ophthalmology.* 1999;83(5):598-604.
56. Lin H-H, Faunce DE, Stacey M, Terajewicz A, Nakamura T, Zhang-Hoover J, et al. The
macrophage F4/80 receptor is required for the induction of antigen-specific efferent
regulatory T cells in peripheral tolerance. *The Journal of experimental medicine.*
2005;201(10):1615-25.
57. Liu Y, Xia Y, and Qiu C-H. Functions of CD169 positive macrophages in human diseases.
Biomedical Reports. 2021;14(2):1-.

58. Linehan E, and Fitzgerald D. Ageing and the immune system: focus on macrophages.
European Journal of Microbiology and Immunology. 2015;5(1):14-24.
59. Chen W, Zhao B, Jiang R, Zhang R, Wang Y, Wu H, et al. Cytokine expression profile in
aqueous humor and sera of patients with acute anterior uveitis. *Current Molecular*
Medicine. 2015;15(6):543-9.
60. Reddy CV, Brown Jr J, Folk JC, Kimura AE, Gupta S, and Walker J. Enlarged blind spots in
chonoretinal inflammatory disorders. *Ophthalmology.* 1996;103(4):606-17.
61. Wu P-C, Chuang M-N, Choi J, Chen H, Wu G, Ohno-Matsui K, et al. Update in myopia and
treatment strategy of atropine use in myopia control. *Eye.* 2019;33(1):3-13.
62. Hsu Y-A, Chen C-S, Wang Y-C, Lin E-S, Chang C-Y, Chen JJ-Y, et al. Anti-inflammatory
effects of resveratrol on human retinal pigment cells and a myopia animal model.
Current Issues in Molecular Biology. 2021;43(2):716-27.
63. Ikeda S-I, Kurihara T, Toda M, Jiang X, Torii H, and Tsubota K. Oral bovine milk lactoferrin
administration suppressed myopia development through matrix metalloproteinase 2 in
a mouse model. *Nutrients.* 2020;12(12):3744.
64. He H, Mack JJ, Güç E, Warren CM, Squadrito ML, Kilarski WW, et al. Perivascular
macrophages limit permeability. *Arteriosclerosis, thrombosis, and vascular biology.*
2016;36(11):2203-12.
65. Du Cheyne C, Tay H, and De Spiegelaere W. The complex TIE between macrophages and
angiogenesis. *Anatomia, Histologia, Embryologia*. 2020;49(5):585-96.
66. Muhammad S, Chaudhry SR, Dobreva G, Lawton MT, Niemelä M, and Hänggi D. Vascular
macrophages as therapeutic targets to treat intracranial aneurysms. *Frontiers in*
Immunology. 2021;12:630381.
67. Raffort J, Lareyre F, Clément M, Hassen-Khodja R, Chinetti G, and Mallat Z. Monocytes
and macrophages in abdominal aortic aneurysm. *Nature Reviews Cardiology.*
2017;14(8):457-71.
68. Deng Z, Shi F, Zhou Z, Sun F, Sun M-H, Sun Q, et al. M1 macrophage mediated increased
reactive oxygen species (ROS) influence wound healing via the MAPK signaling in vitro
and in vivo. *Toxicology and Applied Pharmacology.* 2019;366:83-95.
69. Castaneda OA, Lee S-C, Ho C-T, and Huang T-C. Macrophages in oxidative stress and
models to evaluate the antioxidant function of dietary natural compounds. *journal of*
food and drug analysis. 2017;25(1):111-8.
70. Valentin F, Bueb JL, Kieffer P, Tschirhart E, and Atkinson J. Oxidative stress activates
MMP‑2 in cultured human coronary smooth muscle cells. *Fundamental & clinical*
pharmacology. 2005;19(6):661-7.
71. Bittner A, Alcaíno H, Castro PF, Pérez O, Corbalán R, Troncoso R, et al. Matrix
metalloproteinase-9 activity is associated to oxidative stress in patients with acute
coronary syndrome. *International journal of cardiology.* 2010;143(1):98-100.
72. Pérez S, and Rius-Pérez S. Macrophage polarization and reprogramming in acute
inflammation: a redox perspective. *Antioxidants.* 2022;11(7):1394.
73. Bhatt L, Groeger G, McDermott K, and Cotter TG. Rod and cone photoreceptor cells
produce ROS in response to stress in a live retinal explant system. *Molecular vision.*
2010;16:283.
74. Fujikado T, Kawasaki Y, Fujii J, Taniguchi N, Okada M, Suzuki A, et al. The effect of nitric
oxide synthase inhibitor on form-deprivation myopia. *Current eye research.*
1997;16(10):992-6.

75. Fujikado T, Tsujikawa K, Tamura M, Hosohata J, Kawasaki Y, and Tano Y. Effect of a nitric
oxide synthase inhibitor on lens-induced myopia. *Ophthalmic Research.* 2001;33(2):75-
9.
76. Li Y, Zhang Y, Zhang P, Gao L, Ma Q, Li J, et al. Genetic susceptibility to high myopia in
Han Chinese population. *Open Life Sciences.* 2022;17(1):512-6.
77. Zhu Y, Li X, Chen J, Chen T, Shi Z, Lei M, et al. The pentacyclic triterpene Lupeol switches
M1 macrophages to M2 and ameliorates experimental inflammatory bowel disease.
International immunopharmacology. 2016;30:74-84.
78. Wang Z, Zou J, Huang Y, Cao K, Xu Y, and Wu JM. Effect of resveratrol on platelet
aggregation in vivo and in vitro. *Chinese medical journal.* 2002;115(03):377-80.
79. Hu J, Deng F, Zhao B, Lin Z, Sun Q, Yang X, et al. Lactobacillus murinus alleviate intestinal
ischemia/reperfusion injury through promoting the release of interleukin-10 from M2
macrophages via Toll-like receptor 2 signaling. *Microbiome.* 2022;10(1):1-21.
80. Zhao L, Wang H, and Du X. The therapeutic use of quercetin in ophthalmology: Recent
applications. *Biomedicine & Pharmacotherapy.* 2021;137:111371.
81. Wagh V, Patil P, Surana S, and Wagh K. Forskolin: upcoming antiglaucoma molecule.
Journal of postgraduate medicine. 2012;58(3):199.
82. Dysli C, Enzmann V, Sznitman R, and Zinkernagel MS. Quantitative analysis of mouse
retinal layers using automated segmentation of spectral domain optical coherence
tomography images. *Translational vision science & technology.* 2015;4(4):9-.
83. Hou J, Mori K, Ikeda S-i, Jeong H, Torii H, Negishi K, et al. Ginkgo biloba extracts improve
choroidal circulation leading to suppression of myopia in mice. *Scientific Reports.*
2023;13(1):3772.
84. Galvao J, Davis B, Tilley M, Normando E, Duchen MR, and Cordeiro MF. Unexpected
low‑dose toxicity of the universal solvent DMSO. *The FASEB Journal.* 2014;28(3):1317-
30.

Figures and Tables

Figure 1. Myopia is accompanied by sustained depletion of choroidal-resident
macrophages in mice at 3 and 8 weeks of age.

(**A**) Three-weeks-old wild-type C57BL/6JJc1 mice were injected with either 0.1 ml/10 g body
weight clodronate liposome or control liposome once every two days (n=12 per group).
Refraction, axial length, and choroidal thickness were measured at baseline and 8 days after 4
injections with an infrared photorefractor and SD-OCT system. **(B)** Compared to the control
group, the clodronate liposome (clo-lip) injection group showed a larger refractive change ($P <$
0.001), axial eye growth ($P < 0,01$), and thinner choroid ($P < 0.001$). Bars represent
mean +/− standard deviations. **(C)** The percentage of macrophages in the choroid was analyzed
using flow cytometry. Macrophages were identified as CD11b$^+$ and F4/80$^+$, appearing in the H3-
UR region. Compared with the control liposome injection group, the proportion of macrophages in
the clodronate liposome injection group was significantly reduced ($P < 0.05$). **(D)** Eight-weeks-old
wild-type C57BL/6JJc1 mice (4 per group unless otherwise indicated) were injected with 0.1
ml/10 g body weight clodronate liposome and control liposome once every two days. Refraction,
axial length, and choroidal thickness were measured at baseline and 8 days after 4 injections with
an infrared photorefractor and SD-OCT system. **(E)** Compared to the control group, the
clodronate liposome (clo-lip) injection group showed a remarkably larger refractive change ($P <$
0.01), axial length elongation ($P < 0.05$), and a thinner choroid ($P < 0.01$). Bars represent
mean +/− standard deviations. Differences between the groups were compared using a t-test. *P
< 0.05. **$P < 0.01$. ***$P < 0.001$.

Figure 2. The polarization of M1 macrophages in the choroid can induce myopia in a murine model.

(A) Three-weeks-old wild-type C57BL/6JJc1 mice were divided into two groups (4 per group unless otherwise indicated) and one group was intraperitoneally injected with 10 ul/g LPS and the other with PBS. The injection lasted for 2 weeks, and refraction, axial length, and choroidal thickness were measured weekly with an infrared photorefractor and SD-OCT system. Choroidal blood perfusion was measured with OCTA. In addition, choroid samples were collected for real-time PCR. **(B)** Compared to the PBS injection group, the LPS injection group showed a significantly larger refractive change ($P < 0.001$), significant axial eye growth ($P < 0,001$), and substantially thinner choroid ($P < 0.001$). Bars represent mean +/− standard deviations. **(C)** The choroidal blood perfusion decreased significantly after 2 weeks of LPS injection, compared to the PBS injection group. **(D)** mRNA levels of *Tnf-α* and *IL-6* significantly increased after 2 weeks of LPS injection. *P* values were obtained from a t-test. *$P < 0.05$. **$P < 0.01$. ***$P < 0.001$.

Figure 3. Inhibition of myopia progression by choroidal M2 macrophage polarization was demonstrated in a lens-induced myopia mouse model.

Three-week-old wild-type C57BL/6JJc1 mice were injected with 0.1 ug/100 ul IL-4, and choroid samples were collected at 0 h, 24 h, and 48 h after injection for flow cytometry and real-time PCR. **(A)** Macrophages in the H3-UR region were denoted as CD11b[+] and F4/80[+]. The proportion of macrophages increased significantly after IL-4 injection ($P < 0.001$). M2 macrophages in the Q2-UR were identified as CD206[+]. At 48 h after IL-4 injection, M2 macrophages were dramatically polarized ($P < 0.05$). **(B)** Real-time PCR results showed that the mRNA expression levels of *Mrc1* and *CD163* increased significantly 48 h after IL-4 injection. **(C)** Three-week-old wild-type C57BL/6JJc1 mice were divided into two groups and injected with either 0.1 ug/100 ul IL-4 or PBS on alternate days with myopia induction (4 per group unless otherwise indicated). Refraction, axial length, and choroidal thickness were measured at the initial (3-week-old) and final (6-week-old) stages of myopic induction using an infrared photorefractor and SD-OCT system. **(D)** Eyes treated with −30D lenses showed a significantly larger refractive change ($P < 0.01$), greater axial length elongation ($P < 0.05$), and increased choroidal thickness ($P < 0.001$) than those treated with 0D lenses in the PBS injection mice. Mice injected with 0.1 ug/100 ul IL-4 showed a significantly smaller refractive change ($P < 0.05$), smaller change in axial length($P < 0.05$), and positive change in choroidal thickness with a statistical significance ($P < 0.001$). Bars represent the mean +/− standard deviation. **(E)** To evaluate changes in choroidal blood perfusion, three-weeks-old mice were divided into three groups: both eyes with 0D lenses (control 0D group), binocular myopia induction (control -30D group), and binocular myopia induction with IL-4 injection once every two days group (IL-4 -30D group). Choroidal blood perfusion was assessed at the initial (3-week-old) and end (6-week-old) stages of myopic induction using OCTA.

Compared to the control 0D group, choroidal blood perfusion was significantly decreased in the
control -30D group. Accordingly, compared with the control -30D group, choroidal blood perfusion
was remarkably improved in the IL-4 -30D group. **(F)** Mice used to measure choroidal blood
perfusion were euthanized at 6 weeks of age, and the choroids were collected for flow cytometry.
The results showed that mice treated with -30D lenses and injected with IL-4 had a significant
proportion of M2 macrophages compared to those treated with -30D lenses and injected with
PBS. *P* values correspond to comparisons made using one-way ANOVA. *$P < 0.05$. **$P < 0.01$.
***$P < 0.001$.

Figure 4: Effect of macrophage polarization direction on oxidative stress response.

(A) Experimental plan for the assessment of *Nox2, Nox4, Mmp2,* and *Mmp9* mRNA expression in the choroid during M1 macrophage polarization and M2 macrophage polarization. Three-weeks-old wild-type C57BL/6JJc1 mice were injected with LPS to polarize M1 macrophages and with IL-4 to polarized M2 macrophages, while the control group was injected with PBS. Choroid samples were collected 24 h after LPS injection, 24 h and 48 h after IL-4 injection, and after PBS injection for real-time PCR and ELISA experiments. **(B)** Comparison of *Nox2, Nox4, Mmp2,* and *Mmp9* mRNA expression in the choroid between the control and M1 macrophage polarization group

showed a significant increase in the LPS-injected group. **(C)** Quantitative PCR analyses of the
choroid 48 h after IL-4 injection indicated a corresponding reduction in the expression
of *Nox2, Mmp2,* and *Mmp9* mRNA, corresponding to M2 macrophage polarization. **(D)** ELISA
analysis results showed that compared to the control group, the choroidal 8-OHdG concentration
was decreased in the IL-4 injection group ($P < 0.01$). On the contrary, the choroidal 8-OHdG
concentration was increased in the LPS injection group, compared to the IL-4 injection group ($P <$
0.05). *P* values indicate comparisons relative to the control, one-way ANOVA, or t-test. *$P < 0.05$.
**$P < 0.01$. ***$P < 0.001$.
952

Supplementary Figure 1: EPA administration inhibited the development of myopia caused by LPS injection.

Three-weeks-old wild-type C57BL/6JJc1 mice were divided into three groups: control group, LPS injection group, and LPS administered with EPA group (n=4). Changes in refraction, axial length, and choroidal thickness were measured after 3 weeks of feeding. **(A)** Compared to the large changes in refraction caused by LPS injection, the refractive shift was smaller after EPA administration ($P < 0.01$). **(B)** Compared to the increase in the ratio of axial length to body weight caused by LPS injection, EPA feeding reduced the ratio of axial length to body weight ($P < 0.001$). **(C)** Thinning of choroid occurred in the LPS injection group, whereas EPA feeding significantly improved choroidal thickness ($P < 0.05$). P values indicate comparisons relative to the control using one-way ANOVA. $*P < 0.05$. $**P < 0.01$. $***P < 0.001$.

Supplementary Figure 2: IL-13 injection suppressed myopia progression and increased choroidal blood perfusion in a murine model of lens-induced myopia.

(A) Choroid samples were collected after 48 h of IL-13 injection and PBS injection for flow cytometry analysis, which was designed to evaluate the optimal time for the polarization of M2 macrophages. **(B)** Flow cytometry analysis demonstrated that both macrophage ratio and M2 macrophage ratio were significantly increased 48 h after IL-13 injection, compared to the PBS injection group. **(C)** Three-week-old mice were divided into three groups: 0D lenses on both eyes (control 0D group), binocular myopia induction (control -30D group), and binocular myopia induction and IL-13 injection (IL-13 -30D group). IL-13 injection frequency was once every two days. Refraction, axial length, and choroidal thickness were measured at the initial (3-week-old) and end (6-week-old) stages of myopic induction with an infrared photorefractor and SD-OCT system. **(D)** The control -30D group showed a significantly larger refractive change ($P < 0.001$), greater axial length elongation ($P < 0.05$), and thinner choroidal ($P < 0.001$) compared to the control 0D group. The IL-13 -30D group indicated a significantly smaller refractive change ($P < 0.001$), lesser axial length change ($P < 0.05$), and positive change in choroidal thickness, with a statistical significance ($P < 0.001$). Bars represent mean +/− standard deviations. **(E)** Choroidal blood perfusion was measured using OCTA. Compared to the control 0D group, choroidal blood perfusion was significantly decreased in the control -30D group. Correspondingly, compared to the control -30D group, choroidal blood perfusion was significantly improved in the IL-13 -30D group. **(F)** The proportion of M2 macrophages in the choroid showed a significant increase in the IL-13 -30D group, compared to the control -30D group. P values indicate comparisons relative to control with one-way ANOVA or t-test. *$P < 0.05$. **$P < 0.01$. ***$P < 0.001$.

995 **Supplementary Figure 3: Continuous berberine eye drops can inhibit the progression of**
996 **myopia in mice.**
997
Three-weeks-old wild-type C57BL/6JJc1 mice were separated into control 0D group, control -30D
group, 2% DMSO -30D group, and berberine -30D group (n=4). The frequency of eye drops for
2% DMSO and 0.26 mg/ml berberine was once a day. After 3 weeks of eye drops, the changes in
refraction, axial length, and choroidal thickness were measured using SD-OCT. **(A)** Compared to
the control 0D group, a refractive shift was observed in the control -30D group and 2% DMSO
group ($P < 0.001$). Smaller refractive changes were observed in the berberine -30D group
compared to the 2% DMSO group ($P < 0.001$). **(B)** Compared to the growth of axial length in the
control -30D group and 2% DMSO group ($P < 0.05$), the berberine -30D group showed a smaller
axial length elongation ($P < 0.05$). **(C)** In the control -30D group and 2% DMSO -30D group, the
choroidal thickness was reduced compared to the control 0D group ($P < 0.001$). Contrarily,
choroid thickness was improved in the berberine -30D group compared to the 2% DMSO -30D
group ($P < 0.001$). *P* values indicate comparisons relative to the control using one-way ANOVA.
*$P < 0.05$. **$P < 0.01$. ***$P < 0.001$.
1011