The orientation of choroidal macrophage polarization significantly influences the development of myopia in murine models

Jing Hou[1,2#], Shin-ichi Ikeda[1,2#], Kiwako Mori[1,2], Heonuk Jeong[1,2], Hidemasa Torii[1,2], Kazuno Negishi[1], Kazuo Tsubota[1,3*], Toshihide Kurihara[1,2*]

[1]Department of Ophthalmology, Keio University School of Medicine, 35 Shinanomachi, Shinjuku-ku, 160-8582, Tokyo, Japan.

[2]Laboratory of Photobiology, Keio University School of Medicine, 35 Shinanomachi, Shinjuku-ku, 160-8582, Tokyo, Japan.

[3]Tsubota Laboratory, Inc., 304 Toshin Shinanomachi-ekimae Bldg., 34 Shinanomachi Shinjuku-ku, 160-0016, Tokyo, Japan.

[#] Those authors contributed equally to the work and request the double first authorship.

*Corresponding authors:

Toshihide Kurihara, MD, PhD

Principal Investigator of Laboratory of Photobiology and Associate Professor of the Department of Ophthalmology, Keio University School of Medicine; 35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan.

Tel: +81-3-5363-3204, Fax: +81-3- 5363-3274

E-mail: kurihara@z8.keio.jp

Kazuo Tsubota, MD, PhD

Professor Emeritus of the Department of Ophthalmology, Keio University School of Medicine, and

the CEO of Tsubota Laboratory, Inc.;304 Toshin Shinanomachi-Eki Mae Bldg., 34 Shinanomachi

Shinjuku-ku, Tokyo 160-0016, Japan.

Tel: +81-3-6384-2866

E-mail: tsubota@tsubota-lab.com

Competing Interest Statement

K.T. reports that he is the CEO of Tsubota Laboratory, Inc., Tokyo, Japan, a company that

develops products for the treatment of myopia. The other authors have declared that they have

no conflicts of interest. Patent-pending (J.H. S.I. T.K. K.T. 2022-72596).

Funding

This work is supported by AMED under Grant Number JP22gm1510007 to T. Kurihara.

Keywords: Macrophage; Myopia; Choroid; Vascular Maintenance; Anti-inflammatory.

Article Type: Research paper

Word count: 199 words for the abstract

Number of table(s) and figure(s): 4 figures, and 3 supplementary figures

Abstract

Myopia is a primary contributor to visual impairment and has emerged as a global public health

concern. Evidence indicates that one of the main structural features of myopia is the

corresponding decrease in choroidal thickness, and choroidal macrophages play an important

role in maintaining the choroidal thickness. Nevertheless, the effect of choroidal macrophages on

myopia remains unclear. Here, we discovered that the continuous intraperitoneal injection of

clodronate liposomes depleted choroidal macrophages and leads to myopia, which confirmed that

the presence of choroidal macrophages plays an important role in myopia development.

Subsequently, based on the phenotypic characteristics of macrophages, experiments were

designed to study the effects of different polarization directions of macrophages on myopia

development. We found that lipopolysaccharides (LPS) injection can induce the polarization of

choroidal M1 macrophages and result in myopia. Conversely, IL-4 or IL-13 injection causes

choroidal M2 macrophage polarization and suppresses the progression of myopia. Additionally,

we demonstrated that the opposite effects of M1 and M2 macrophages on myopia development

may be related to their impacts on inflammation and oxidative stress response. These findings

establish that choroidal macrophages are critically important in the development of myopia and

provide new strategies for the development of myopic therapies.

Main Text

Introduction

Myopia, commonly referred to as "short-sightedness," is the most prevalent vision-related

disease, which is caused by refractive error depending on the elongation of the eyeball along the

visual axis (1-4). Current research has shown that myopia can be influenced by a variety of

factors, including environmental impacts, dietary factors, and genetics; however, the

pathogenesis of myopia remains unclear (5-8). Numerous studies have suggested that

inflammation plays a role in the development of myopia: increased inflammation in the eye

accelerates the development of myopia, whereas lower inflammation inhibits it (9-12). It is worth

noting that macrophages are essential for the occurrence, maintenance, and resolution of

inflammation (13).

Macrophages are an essential component of the innate immune system that exists in all

mammalian tissues and exhibit great functional diversity. It not only plays a variety of functions

related to inflammation but also plays a crucial role in the development and maintenance of

homeostasis (14, 15). Healthy corneas contain a population of macrophages that maintain

corneal lymphatic homeostasis and promote lymphangiogenesis under inflammatory conditions

(16). The neural retina is populated by microglia, specialized resident macrophages that

contribute to the structural and functional integrity of neuronal synapses (16, 17). Under

homeostatic conditions, there are no microglia in the subretinal space or the choroid and are

restricted to the inner layers of the retina (18-20). Further, the choroid was found to contain a

dense network of resident tissue macrophages as early as the 1990s (21). At the level of

choroidal arteries and arterioles, dendritic resident macrophages were closely aligned along the

vessel wall. In the choroidal terminal choriocapillaris, they are not tightly packed on the vessel

wall, but are widely distributed in the connective tissue between blood vessels (22). These

findings provide a basis for exploring the relationship between choroidal vessels and

macrophages. Subsequently, researchers found that the ablation of a considerable majority of

choroidal macrophages was associated with broad and gradual atrophy of the choroidal

vasculature and thinning of the choroid (23, 24). This finding reveals that choroidal macrophages

play a hitherto underestimated role in the maintenance of the choroidal vascular structure. In the

field of myopia research, the stability of the choroidal vasculature is recognized as the most

important component for maintaining choroidal thickness, and choroidal thickness variation is a

structural characteristic of the development of myopia (25-28). Therefore, we hypothesized that

choroidal macrophages play a crucial role in the progression of myopia.

Macrophage subsets can be classified as classic M1 macrophages and alternative M2

macrophages, according to their pro- or anti-inflammatory properties in the polarized state (29).

M1 macrophages can be polarized by lipopolysaccharide (LPS), leading to a high expression of

inflammatory cytokines, such as tumor necrosis factor-α (TNFA) and interleukin-6 (IL6), as well

as reactive oxygen species. Correspondingly, M2 macrophage polarization induced by IL-4 and

IL-13 diminishes and alleviates inflammation following infection or damage, and inhibits the

production of reactive oxygen species (30-32). Studies have shown that inflammatory diseases

increase the risk of myopia, and inflammation in the eye directly contributes to the progression of

myopia (12, 33), all this underscores the importance of controlling inflammation in inhibiting the

development of myopia. In addition, it has been suggested that the existence of several

supplements which convert M1 into M2 macrophages, but this property has not been studied and

discussed in conjunction with myopia suppression (34-36). Therefore, we hypothesized that

different subsets of macrophages in the choroid have diametrically opposite effects on the

development of myopia according to their polarization direction and characteristics.

In this study, we investigated the consumption and polarization of choroidal macrophages in a

mouse model and combined a lens-induced myopia model (LIM) with the polarization of choroidal

macrophage to investigate the effect of different macrophage polarization directions on myopia

development. We found that choroidal macrophage ablation was associated with myopia

progression. We also discovered that the tendency for myopia development or inhibition is closely

related to the direction of macrophage polarization in the choroid and that macrophages interfere

with the development of myopia based on their effects on inflammation and oxidative stress.

Taken together, our findings suggest that macrophages play a key role in myopia and emphasize

the different characteristics of macrophages as a potential mechanism for developing or inhibiting

myopia. Regulating the phenotype of macrophages in the choroid may have therapeutic effects

on maintaining choroidal vasculature thereby slowing the progression of myopia.

Results

Sustained intraperitoneal injection of clodronate liposome depletes macrophages in the

choroid and can trigger myopia

It has been previously reported that resident macrophages in the choroid positively affect

choroidal thickness and maintain vascular status (24). In contrast, a series of animal models have

demonstrated that choroidal thinning is an inevitable change in the development of experimental

myopia and is a prominent feature in the development of myopia (26, 28, 37-39). To investigate

the effect of choroidal-resident macrophages on myopia in a mouse model, 3-weeks-old C57BL/6

J mice were randomly divided into 2 groups and were injected intraperitoneally with clodronate or

phosphate-buffered saline (PBS) liposomes. In the clodronate liposome group, after being

engulfed by macrophages, the clodronate liposomes were degraded by lysosomal

phospholipases, which released clodronate into the macrophages and induced apoptosis, thereby

promoting macrophage depletion (40). The injection volume of clodronate liposome was 0.1 ml

per 10 g of mice and the injection frequency was once every two days. Refraction, axial length,

and choroidal thickness were measured using an infrared photorefractor and a spectral domain-

optical coherence tomography (SD-OCT) system, before injection and after 8 days of injections,

respectively (Fig. 1A). Mice injected with clodronate liposome showed a higher refractive shift

(-7.30 ± 3.61 D vs. $+1.17 \pm 4.40$ D, $P < 0.001$), axial length elongation (0.08 ± 0.02 mm vs.

0.05 ± 0.02 mm, $P < 0.01$), and thinner choroid (−1.56 ± 0.89 mm vs. 1.09 ± 0.51 mm, $P < 0.001$)

compared to the PBS liposome injection group (Fig. 1B). Flow cytometry analysis of the

percentage of choroidal resident macrophages demonstrated a corresponding decrease after 8

days of clodronate liposome injection compared to the control group (2.583 ± 0.002% vs. 9.290

± 0.018%, $P < 0.05$) (Fig. 1C). These results indicated that clodronate liposome injection depletes

resident macrophages in the choroid, promotes choroidal thinning, and induces myopia in 3-

week-old mice. In previous studies, we observed that choroidal thickness did not increase and

tended to be stable from approximately 8 weeks of age in mice (41). However, whether the

depletion of choroidal macrophages in adult mice also contributes to myopia has not been

elucidated. To rule out the possibility that choroidal macrophage depletion can cause myopia

independent of choroidal thickness, 8-weeks-old adult mice were injected with clodronate

liposomes or PBS liposomes in the same manner, and changes in refraction, axial length, and

choroidal thickness were analyzed 8 days later (Fig. 1D). Intraperitoneal injection of clodronate

liposomes significantly reduced body weight while consuming resident macrophages, which

affects the comparison of changes in the axial length of the eye. To avoid the effects of body

weight changes on the axial length, we used the ratio of the axial length to body weight to

represent the changes in the axial length of the eye. In the clodronate liposome injection group,

refraction shift (−7.52 ± 2.57 D vs. +1.52 ± 4.91 D, $P < 0.01$), a higher axial length/body weight

(0.168 ± 0.015 mm/g vs. 0.155 ± 0.002 mm/g, $P < 0.05$), and thinner choroidal (−2.39 ± 1.33 mm

vs. −0.12 ± 1.48 mm, $P < 0.01$) were observed compared to the PBS liposome injection group

(Fig. 1E), suggesting that clodronate liposome injection causes choroidal resident macrophage

depletion and thinning of the choroid, which leads to myopia regardless of whether the choroid is

developing.

Continuous intraperitoneal injection of LPS stimulates choroidal M1 macrophage

polarization and induces myopia in a murine model

Macrophages are classified into two categories based on their activation state and functions: M1

type (classically activated macrophages) and M2 type (alternatively activated macrophages),

which exhibit pro-inflammatory and anti-inflammatory properties, respectively (31). As the

upregulation of allergic inflammation has been implicated in the progression of myopia in animal

models (33), we investigated whether M1 macrophage polarization can induce myopia in a

mouse model. Mice were divided into LPS (10 mg/ml) and PBS injection groups for daily

intraperitoneal injections. To determine whether myopia had developed, refraction, axial length,

and choroidal thickness were measured using SD-OCT before the injections, 1 week after, and 2

weeks after the injections (Fig. 2A). After 1 week of LPS injection, compared to PBS-injected

mice, LPS-injected mice showed refractive shift (−9.21 ± 5.91 D vs. +2.91 ± 3.91 D, $P < 0.001$),

rise in axial length/body weight (0.28 ± 0.01 mm/g vs. 0.22 ± 0.005 mm/g, $P < 0.001$), and thinning

of choroid (−1.11 ± 0.90 mm vs. 1.36 ± 0.70 mm, $P < 0.001$). The injections were continued until

the end of the 2 weeks, and LPS-injected mice maintained a significantly higher refractive shift

(−7.23 ± 4.36 D vs. +8.41 ± 7.01 D, $P < 0.001$), increase in the ratio of axial length to mouse body

weight (0.20 ± 0.005 mm/g vs. 0.17 ± 0.004 mm/g, $P < 0.01$), and thinning of choroid (−1.77 ± 1.02

mm vs. 2.05 ± 0.66 mm, $P < 0.001$) (Fig. 2B). Choroidal blood perfusion was measured by optical

coherence tomography angiography (OCTA), and the results showed that the choroidal blood

perfusion of mice injected with LPS for 2 weeks was reduced compared to that of mice injected

with PBS (37.36 ± 3.74 % area vs. 46.79 ± 4.74 % area, $P < 0.001$) (Fig. 2C). These data show

that the intraperitoneal injection of LPS can induce myopia in a mouse model. Choroid samples

were collected for real-time polymerase chain reaction (PCR) analysis to verify the polarization of

M1 macrophages in the choroid stimulated by intraperitoneal injection of LPS. The expression

levels of the pro-inflammatory cytokines *Tnf* and *Il6* were significantly upregulated in LPS-injected

choroid samples compared to those in PBS-injected choroid samples (Fig. 2D). These data

indicate that the intraperitoneal injection of LPS polarizes M1 macrophages in the choroid,

promotes the release of inflammatory cytokines, and causes myopia in a mouse model. As

reported in LPS-treated macrophages, pro-inflammatory cytokines (TNFA, and IL6) secretion was

significantly attenuated by eicosapentaenoic acid (EPA), which can inhibit myopia progression

(42, 43). Therefore, an experiment was designed to administer an EPA diet to LPS-induced

myopic mice, and it was found that myopia did not progress (Supplementary Figs. 1A, B, C).

Taken together, these data suggest that the LPS-induced phenotypic switch toward M1

macrophages is associated with the development of myopia.

The constant intraperitoneal injection of IL-4/IL-13 encouraged the polarization of

choroidal M2 macrophages and suppressed myopia progression in a murine model of LIM

We found that intraperitoneal injection of LPS enhanced the expression of pro-inflammatory

cytokines in the choroid and promoted the development of myopia; contrarily, choroidal M2

macrophage polarization inhibited the development of myopia. Previous studies have

demonstrated that IL-4 and IL-13, as anti-inflammatory interleukins, can directly activate M2

macrophages, which can be identified by the expression of M2 macrophage markers, including

mannose receptor 1 (Mrc1), also referred to as CD206 and CD163 (44). Above all, to investigate

the frequency of continuous IL-4 injection, we compared the proportion of M2 macrophages in the

choroid at 0 h, 24 h, and 48 h after being subjected to 0.1 µg/100 µl IL-4 intraperitoneal injection;

F4/80 and CD11b-positive cells were regarded as macrophages and CD206 was used as a

marker to identify M2 macrophages. Flow cytometry analysis showed a growth in the ratio of

macrophages (6.73 ± 0.22 % vs. 3.78 ± 0.41 %, $P < 0.001$) and M2 macrophages (43.57 ± 4.72 %

vs. 30.33 ± 8.35 %, $P < 0.05$) at 48 h after IL-4 intraperitoneal injection, compared to 0 h (Fig. 3A).

Real-time PCR detection after IL-4 injection under the same conditions showed that *Mrc1* and

CD163 mRNA expression in the choroid was enhanced 48 h after IL-4 injection (Fig. 3B). These

results showed that the optimal frequency of IL-4 injection is once every two days. Next, we

examined whether the promotion of M2 macrophage polarization by IL-4 injection on alternate

days could inhibit myopia progression in a LIM mouse model. All 3-weeks-old mice were

subjected to LIM, in the same manner as previously established (45), and administered with

either PBS or 0.1 ug/100 ul of IL-4 injection until 6 weeks of age (Fig. 3C). In the PBS injection

group, eyes treated with −30 diopter (D) lenses showed refractive change (−11.52 ± 4.91 D vs.

+12.45 ± 8.19 D, $P < 0.01$), increased axial length (0.22 ± 0.01 mm vs. 0.18 ± 0.01 mm/g, $P <$

0.05), and decreased choroidal thickness (−1.74 ± 0.66 mm vs. 3.91 ± 0.95 mm, $P < 0.001$)

compared to eyes treated with 0D lenses. In contrast, compared to the PBS- injected group with

−30D lenses, the IL-4 injection group showed smaller refractive change (−1.39 ± 2.91 D vs.

−11.52 ± 4.91 D, $P < 0.05$), smaller axial elongation (0.19 ± 0.01 mm vs. 0.22 ± 0.01 mm/g, $P <$

0.05), and thickening of the choroid (1.39 ± 0.19 mm vs. −1.74 ± 0.66 mm, $P < 0.001$) in eyes with

-30D lenses (Fig. 3D). Meanwhile, to demonstrate the influence of continuous IL-4 injection on

choroidal blood perfusion, 3-weeks-old mice were divided into three groups: control 0D group

(both eyes were treated with 0D lenses), control −30D group (binocular myopia induction), and IL-

4 −30D group (binocular myopia induction with IL-4 injection once every two days). OCTA was

used to measure choroidal blood perfusion at the beginning (3 weeks old) and end (6 weeks old)

of myopic induction. Compared to the control 0D group, choroidal blood perfusion was lower

(8.59 ± 5.90 %area vs. −6.29 ± 8.73 %area, $P < 0.01$) after 3 weeks of myopia induction.

Nevertheless, compared to the control -30D group, choroidal blood perfusion was improved

(−6.29 ± 8.73 %area vs. 1.54 ± 9.34 %area, $P < 0.05$) in the IL4 −30D group (Fig. 3E). To

visualize the polarization of M2 macrophages, after measuring choroidal blood perfusion at the

end stage, choroid samples were collected for flow cytometry, and data analysis revealed the

corresponding growth in the proportion of M2 macrophages (46.52 ± 0.05% vs. 35.71 ± 0.06%, P

< 0.05) in the IL-4 injection group with −30D lenses, compared to the PBS injection group with

−30D lenses (Fig. 3F). Collectively, these findings demonstrate that IL-4 injection can stimulate

the polarization of M2 macrophages in the choroid and suppress the development of myopia by

improving choroidal blood perfusion in a LIM mouse model. To reinforce the conclusion that M2

macrophages can inhibit myopia progression, we performed the same experiments with 0.1 μg

/100 μl IL13 injection instead of IL-4 injection, and the results also showed that M2 macrophages

were polarized 48 h after IL-13 injection (Supplementary Figs. 2A, B). The results of the LIM

mouse model combined with IL-13 injection were as expected; IL-13 also polarized M2

macrophages, suppressed myopia progression, and improved choroidal blood perfusion in a

murine model of LIM (Supplementary Figs. 2C, D, E, F).

Berberine, an eye drop that polarizes M2 macrophages, inhibits myopia progression after

consistent use

After realizing that M2 macrophage polarization may inhibit the development of myopia, we used

an M2 macrophage-activating drug as an eye drop to verify its effect on the development of

myopia. Some studies have shown that berberine promotes the polarization of M1 macrophages

to the M2 anti-inflammatory phenotype in mice (46, 47). To verify the role of berberine eye drops

in the development of myopia, we examined and compared changes in refraction, axial length,

and choroidal thickness in 3-weeks-old C57BL/6 J mice after myopia induction for 3 weeks and

the administration of either berberine or dimethyl sulfoxide (DMSO) eye drops. Compared to the

control 0D group (received 0D lenses for both eyes), the control −30D group (received −30D

lenses for both eyes) and 2% DMSO −30D group (binocular myopia induction with the application

of 2% DMSO eye drops daily) showed a large refractive change ($P < 0.001$), increased axial

length elongation ($P < 0.05$), and decreased choroidal thickness ($P < 0.001$). compared to the

2% DMSO −30D group, the berberine −30D group (binocular myopia induction with daily 0.26

mg/ml eye drops) exhibited smaller refractive changes (2.12 ± 1.63 D vs. -5.25 ± 2.89 D, $P <$

0.001), smaller axial elongation (0.19 ± 0.02 mm vs. 0.22 ± 0.04 mm/g, $P < 0.05$), and a relatively

thicker choroid (0.79 ± 0.64 mm vs. -1.56 ± 0.76 mm, $P < 0.001$). These results indicate that

drugs used to activate the polarization of M2 macrophages may also have the potential to inhibit

the development of myopia (Supplementary Figs. 3A, B, C).

Effects of LPS and IL-4 injections on nicotinamide adenine dinucleotide phosphate

(NADPH) oxidation activity

We inferred that the direction of macrophage polarization may influence the development or

inhibition of myopia through its pro-inflammatory or anti-inflammatory properties. However, the

other mechanisms underlying this association remain largely unknown. LPS-treated M1

macrophages are associated with high levels of reactive oxygen and nitrogen species (ROS and

RNS, respectively) and matrix metalloproteinase (MMP) activity, and ROS production is critical for

M1 macrophage activation and function (48, 49). Correspondingly, ROS overproduction may

impair blood perfusion to the retina, and circulatory disturbances during the development of

myopia can lead to oxidative stress (50, 51). To investigate whether M1 macrophage polarization

also induces myopia by promoting ROS and oxidative stress response, and whether M2

macrophage polarization suppressed myopia progression by inhibiting ROS release and oxidative

stress response, the mRNA expression of NADPH oxidase 2 (*Nox2*), which is the most important

source of ROS, and the expression of *Nox4, Mmp2*, and *Mmp9* in the choroid were observed

after either LPS or IL-4 injection (Fig. 4A). Quantitative analysis showed increased *Nox2, Nox4,*

Mmp2, and *Mmp9* mRNA expression in the LPS injection group compared to that in the PBS

injection group (Fig. 4B). In contrast, 48 h after IL-4 injection, the mice showed lower *Nox2,*

Mmp2, and *Mmp9* mRNA expression (Fig. 4C). Since 8-hydroxy-2-deoxyguanosine (8-OHdG) is

an effective marker for assessing oxidative DNA damage (52), we investigated the level of 8-

OHdG in the choroid after intraperitoneal injection of either LPS or IL-4 and found that the IL-4

injection group showed a reduction in choroidal 8-OHdG concentration compared to the control

group ($P < 0.01$). In contrast, the concentration of choroidal 8-OHdG was higher in the LPS

injection group, compared to the IL-4 group ($P < 0.05$) (Fig. 4D). Taken together, these data

suggest that polarized M1 macrophages may induce myopia by promoting oxidative stress,

whereas polarized M2 macrophages may inhibit myopia progression by suppressing the oxidative

stress response.

Discussion

In this paper, we have found that M1 macrophage accelerates myopia while M2 macrophage

suppressed myopia progression, this statement is supported by the data performed and

described. Since the choroid is now considered to be the most important factor for the

development of myopia, maintenance of the choroidal vascular is critical for the suppression of

myopia and our findings on the role of macrophages is indeed a novel concept, and open up new

avenues for further investigation.

Macrophages are tissue-resident or infiltrating immune cells that are present in all tissues. In

addition to their immune properties, macrophages play an important role in regulating normal

tissue development and homeostasis (14). Researchers have found that resident macrophages

are densely distributed in the choroid tissue, a connective tissue rich in pigments and blood

vessels that supply blood and nutrients to the outer retina (16, 53). However, the precise role of

choroidal-resident macrophages has rarely been directly studied. Recently, choroidal

macrophages have been suggested to perform an underappreciated trophic function by

preserving the vascular structure of the choroid (24). In the present study, we confirmed that

choroidal thinning occurred following depleted choroidal-resident macrophages and was

accompanied by axial length elongation and a significant refractive shift. This unexpected finding

supports the previously unknown concept that macrophages in the developing choroid are

associated with choroidal growth and myopia.

The ability of resident immune cells to maintain the vasculature changes depending on the

developmental state of the tissue (24). Microglia, as specialized macrophages, play an important

role in shaping the central nervous system (CNS) vasculature during development but are not

necessary for the maintenance of the mature uninjured CNS (17, 54). The choroidal thickness

steadily increases and reaches a steady state at 7–9 weeks of age; thereafter, it gradually

decreases with age (41). In our experiments, we used 8-weeks-old mice with a plateau in

choroidal thickness, depleted choroidal-resident macrophages, and found that choroidal thickness

was thinner and myopia still occurred. These results demonstrate that choroidal macrophages

effectively support choroidal growth during development and maintain choroidal stability after

maturation.

Studies of the normal choroid in mice have revealed a network of F4/80[+] (CD160) and CD169[+]

macrophages widely distributed around choroidal blood vessels (55). F4/80 is one of the most

specific cell surface markers of murine macrophages, and CD169[+] macrophages are a unique

subset of macrophages involved in multiple immune responses (56, 57). Activated macrophages

can be broadly divided into the M1 and M2 cell subsets, representing the two extremes of the

macrophage activation spectrum that can occur *in vivo* (58). M1 macrophage polarization occurs

in many ocular inflammatory diseases and is accompanied by the production of pro-inflammatory

cytokines; these inflammatory reactions induce local biochemical reactions, leading to local tissue

remodeling and disease progression. Numerous studies have shown that inflammatory diseases

of the eye can increase the risk of myopia. Hyperacute Vogt-Koyanagi-Harada (VKH) disease

often results in acute myopia due to ciliary body inflammatory exudation (59), Myopic refractive

errors are significantly increased in patients with juvenile chronic arthritis (JCA)-related uveitis,

and inflammatory cytokines IL6 and TNFA were found to be overexpressed in the aqueous humor

of uveitis patients (9, 60). These findings suggest that acute or chronic systemic inflammatory

diseases increase the risk of myopia. In this study, we demonstrated that LPS injection stimulates

the expression of pro-inflammatory cytokines in the choroid, reduces choroidal thickness, and

induces myopia in a murine model. This study suggests that the pro-inflammatory properties of

M1 macrophages may contribute to the pathogenesis of myopia.

In contrast, some anti-inflammatory substances have been shown to inhibit the development of

myopia when administered orally or as eye drops. Atropine eye drops are currently the most

effective therapy for myopia control and have been shown to be associated with effective

downregulation of eye inflammation (12, 61). Resveratrol eye drops reduced inflammation in a

monocular-form deprivation (MFD) animal model to prevent myopia development (62). In oral

preparations, the oral administration of lactoferrin (LF), which has anti-inflammatory properties,

inhibited myopia progression in a LIM mouse model (63). However, no studies have shown how

the polarization of M2 macrophages with anti-inflammatory properties directly affects the

development of myopia. Based on the highly plastic nature of macrophages, we administered IL-4

and IL-13 injections to induce the polarization of M2 macrophages in the choroid and found that

the expression of anti-inflammatory cytokines was activated in the choroid and inhibited the

progression of myopia. Taken together, these results indicate that direct stimulation of M1

macrophages to release inflammatory factors induces inflammatory reactions, leading to

accelerated myopia, which can be reversed by direct stimulation of M2 macrophages to release

anti-inflammatory factors.

We also found that with the activation of M2 macrophages, choroid blood perfusion, and choroid

thickness also increased correspondingly, this indicates M2 macrophages in the choroid play a

crucial role in choroidal thickness and maintenance of blood perfusion. This is consistent with its

property of maintaining the stability of the vascular structure. Under a physiological state, M2

macrophages contribute to endothelial barrier protection and facilitate vascular stability and

maturation (64, 65)**.** In a pathological condition, ameliorating macrophage inflammation or

polarizing it into the M2 phenotype can prevent the rupture of intracranial aneurysms (66). M2

macrophage polarization in the aneurysmal aortic wall heals the aortic wall and maintains

vascular integrity as the disease progresses (67). The ability of M2 macrophages to maintain

vascular stability provides strong evidence that it can maintain the stability and overall

morphology of the choroid vascular structure and thus against the development of myopia.

In addition to influencing myopia development through their anti-inflammatory or maintaining

vascular structural stability properties, M2 macrophages may also inhibit the progression of

myopia through their ability to inhibit oxidative stress. In contrast, the mechanism of myopia

induced by M1 macrophage polarization may be related to the activation of oxidative metabolism.

Polarized M1 macrophages produce reactive oxygen species (ROS) and drive inflammatory

responses, leading to the production of high levels of nitric oxide (NO) and NO synthase (iNOS);

excessive ROS can cause oxidative stress and damage biomolecules. Oxidative stress

upregulates the expression of matrix metalloproteinases-2 and -9 (MMP-2 and MMP-9) (68-71).

M2 macrophages have the opposite effects (72). Our study demonstrated that stimulating the

polarization of M1 macrophages in the choroid induces the expression of NADPH oxidases,

matrix metalloproteinase, and 8-OHdG while stimulating the polarization of M2 macrophages

results in the exact opposite effect. Moreover, there is evidence that oxidative stress plays a role

in the development of myopia-associated disorders at the cellular level (51). In pathological

myopia, retinal hypoxia and ischemia lead to the accumulation of ROS and insufficient ability of

the endogenous antioxidant system to clear ROS, which leads to an increase in hypoxia-inducible

factor-1 (HIF-1) levels and further promotes the development of myopia (51, 73). N-omega-nitro-

L-arginine methyl ester (L-NAME), an inhibitor of NOS, was injected intravitreally to prevent the

development of myopia in both form-deprivation myopia (FDM) and LIM animal models (74, 75).

In addition to its anti-inflammatory properties, LF can reverse high levels of MMP-2 activity in the

LIM model, thereby inhibiting myopia (63). Additionally, research has indicated that the MMP-9

gene locus may contribute to myopia (76). Combined with our results, the role of macrophages in

oxidative stress response may be a key factor affecting the development of myopia.

Currently, studies have shown that some natural compounds or supplements that modulate M1 to

M2 macrophages have therapeutic effects on specific diseases. Lupeol, as a multi-target agent,

ameliorates experimental inflammatory bowel disease by converting M1 to M2 macrophages (77).

Resveratrol through its anti-oxidant capacity and anti-inflammatory properties inhibits platelet

aggregation, thereby preventing atherosclerosis (78). Lactobacillus murinus alleviate intestinal

ischemia/reperfusion damage by encouraging M2 macrophages to secrete interleukin-10 (79). In

addition to all these, there are some compounds that treat ophthalmology-related diseases by

their properties of promoting the transformation of M1 into M2 macrophages. Quercetin may have

an impact on the treatment of ophthalmic diseases such as conjunctivitis, dry eye, and

retinopathy due to its anti-inflammatory and antioxidant properties (80). Forskolin is a medication

that can decrease intraocular pressure and be used as an anti-glaucoma agent (81). According to

the properties of various natural compounds, which are involved in regulating the phenotype

transformation of macrophages from M1 to M2, we targeted the agent named berberine, made it

into eye drops, and applied it in myopic-induced mice, we found that it has the ability to inhibit the

development of myopia. It indicates a possibility that control of macrophage phenotype may be a

new therapy applicable to myopia treatment.

In summary, our results have important implications for understanding the relationship between

macrophages and the progression of myopia. In addition to the notion that the two modes of

macrophage polarization can have specific effects on choroidal structure as well as the

development of myopia, our findings show that the effect of macrophages on myopia

development may be related to inflammation, vascular maintenance, and oxidative stress

response, and these possible co-existing mechanisms may play a key role in the development of

myopia. The above views provide new ideas for regulating the macrophage response to

interventions in myopia and provide a new method for drug screening for myopia treatment.

Materials and Methods

Mice

The Ethics Committee on Animal Research of the Keio University School of Medicine approved

all procedures, which followed the Association for Research in Vision and Ophthalmology

Statement for the Use of Animals in Ophthalmic and Vision Research, Keio University's

Institutional Guidelines on Animal Experimentation, and the Animal Research: Reporting of In

Vivo Experiments (ARRIVE) guidelines. In addition, this study implemented the principle of

random assignment. CLEA Japan, Inc. provided the wild-type male C57BL/6J mice. Those mice

were kept in standard transparent mouse cages (29×18×13 cm) with four or five per cage in

pathogen-free environments maintained at $23 \pm 3\,°C$, under approximately 50 lux of background

fluorescent lamp light (color temperature: 5000 K), with a 12-hour diurnal period and constant

access to normal chow (MF, Oriental Yeast Co., Ltd, Tokyo, Japan) and tap water. All the animals

were randomly allocated.

LIM mouse model

A murine LIM model was created as previously described (45). Briefly, mice were placed under general anesthesia using a mixture of midazolam (Sandoz K.K., Tokyo, Japan), medetomidine (Domitor®, Orion Corporation, Espoo, Finland), and butorphanol tartrate (MMB) (Meiji Seika Pharma Co., Ltd., Tokyo, Japan). Eyeglass frames that fit the mouse's head's contours were designed and printed using a three-dimensional printer. For myopia induction, a negative 30D lens was constructed using polymethyl methacrylate (PMMA). The left and right eyes of the spectacles were adjusted to fit the shape of the mouse skull frame and screwed onto a stick, which was a joint enabling the left and right frame positions to be altered or removed for cleaning. Subsequently, the stick was bonded to the mouse skull using a self-curing dental adhesive system. Induction began once the mouse had fully recovered from anesthesia, and the lenses were removed for cleaning at least twice a week.

The measurement of refraction, axial length, and choroidal thickness

Refraction was measured using an infrared photorefractor (Steinbeis Transfer Center, Stuttgart, Baden-Württemberg, Germany) as previously reported (45). After measuring refraction, axial length, and choroidal thickness were measured using an SD-OCT system (Envisu R4310, Leica Microsystems, Wetzlar, Germany). For each mouse, measurements of refraction, axial length, and choroidal thickness were performed before the start of LIM (0W) and at the end of LIM (3W). Before the measurements, the mouse's eyes were treated with mydriasis eye drops containing 0.5% tropicamide and 0.5% phenylephrine (Santen Pharmaceutical Co., Ltd., Osaka, Japan) to ensure mydriasis and cycloplegia. After pupil dilation, the mice were subjected to general anesthesia using MMB. It is important to prevent the occurrence of corneal injury during measurement. Axial length was measured as the vertical distance from the anterior corneal surface to the retinal pigment epithelium layer near the optic nerve (45). Choroidal thickness was determined by quantifying the circular area of the disc on the posterior surface of the choroid using ImageJ software (82).

The measurement of choroidal blood perfusion

The measurement of choroidal blood perfusion

Choroidal blood perfusion was measured at the beginning (3-week-old) and end (6-week-old)

stages of myopia induction using an SS-OCT/OCTA device (XEPHILIO OCT-S1, CANON

Medical Systems, Tokyo, Japan). The measurement method has been described in a previous

report (83). Briefly, as with refraction measurements, the mice were dilated under general

anesthesia before measurement. Choroidal blood perfusion was measured using en face

angiography to identify the optic nerve as the central area. The choroidal blood perfusion signal

was obtained from B-scan images at the corresponding location. In the B-scan images, the area

covered by red noise points was the area without blood perfusion. ImageJ quantitative analysis

was used to calculate the non-blood perfusion area in the choroid and evaluate the percentage of

the area with blood perfusion in the choroid.

Flow cytometry

The mice were administered an overdose of MMB to induce profound anesthesia before being

euthanized by cervical dislocation. The eyes were immediately enucleated and the anterior

segment, vitreous, and retina were discarded. Choroid tissues were carefully scraped off the

sclera-choroid complex, collected in the tube (10 choroid tissues/tube), and then processed in

digestion buffer (0.75 mg/ml of collagenase A, from FUJIFILM, Tokyo, Japan) in complete

Dulbecco's Modified Eagle Medium (DMEM) solution for 45 min at 37 °C. The digested choroid

mixture was transferred onto the cell strainers and any undigested choroid pieces were mashed

using the end of a syringe plunger while adding 5 ml cold complete DMEM to allow the pass-

through of cells into new 50 ml tubes below the strainer. The aggregated choroidal cells were

dissociated into single-cell suspensions via differential centrifugation. The cells were washed by

adding 700 ul of flow cytometry staining (FACS) buffer (0.1 g of bovine serum albumin (BSA) and

80 ul of 0.5 mol/L ethylenediaminetetraacetic acid (EDTA) in 20 ml of PBS (pH 7.4) solution to the

cell suspension. The pellet was resuspended with 10 ul Fc block solution (1:10 Fc block) (BD

Biosciences, New Jersey, USA) for 10 min, stained with a mixture of fluorochrome-conjugated

antibodies (1:200 FITC anti-mouse/human CD11b M1/70, BioLegend, California, USA), (1:200

APC/Cyanine7 anti-mouse F4/80, BioLegend, California, USA), (1:200 CD206, AbD Serotec,

Kidlington, near Oxford, UK) for at least 45 min, and finally covered with 0.1 mg/ml Hoechst

(DOJINDO, Amsterdam, Netherlands) for 10 min. All incubations were performed on ice and light

was avoided. Data were acquired on a CytoFLEX S flow cytometer using CytExpert software

(Beckman Coulter Life Sciences, Inc., Indiana, United States), and offline data analysis was

performed using CyExpert software.

Clodronate liposome treatment

Clodronate liposome (Liposoma BV, Amsterdam, The Netherlands) was intraperitoneally injected

at a dose of 0.10 ml/10 g (once every two days) to C57BL/6J mice from P21 to P28 or from P56

to P63. The control group was injected with an equal amount of PBS liposomes, as described

above. Refraction, axial length, and choroidal thickness were measured before and after 4 four

injections. The mice were then euthanized and choroid samples were collected.

LPS treatment

Lipopolysaccharides from *Escherichia coli* O111:B4 (Merck, Tokyo, Japan) were dissolved in

PBS and injected intraperitoneally to induce systemic M1 macrophage polarization. The

concentration of LPS was 10 mg/ml, and the injection dose was increased proportionally to the

body weight of the mice, with 10 µg of LPS given per gram of mice. Since LPS injections caused

a large reduction in mouse weight, we monitored and documented the mouse body weight before

each injection. In addition, a corresponding volume of PBS was injected intraperitoneally as a

vehicle control.

EPA administration

EPA-supplemented chow containing 5% of EPA (EPADEL®, Mochida, Pharmaceutical Co. Ltd.,

Tokyo, Japan) and normal chow were prepared as previously reported (43). Briefly, EPA ethyl

ester was blended with powdered ordinary chow, and tap water was added. The mixture was

shaped into tiny cylinders and dried for consumption. Feeding with EPA-mixed chow began and

ended with myopia induction.

IL-4/IL-13 treatment

To induce systemic M2 macrophage polarization, recombinant murine IL-4 (214-14) and IL-13

(210-13) (PeproTech, New Jersey, United States) were dissolved in PBS at the same

concentration of 0.1 µg/100 µl, and injected intraperitoneally into the mice once every two days.

The control group was intraperitoneally administered the same volume of PBS. The injections

were administered during the induction of myopia. To ensure that the mice grew normally, their

body weights were measured and recorded prior to each injection.

Berberine chloride eye drops

Berberine chloride hydrate (TCI, Tokyo, Japan) solution was freshly prepared in DMSO. When

determining the concentration of berberine chloride eye drops, we found that berberine chloride

hydrate could not be dissolved in water at room temperature. We attempted to use 100% DMSO

to dissolve it and found that the solubility of berberine chloride hydrate in 100% DMSO at room

temperature was approximately 13 mg/ml. At present, the concentration of berberine chloride eye

drops is 0.26 mg/ml. In addition, DMSO, a dissolution vector, has been reported to exert cytotoxic

effects at certain concentrations. Therefore, an untreated control group should be included along

with the DMSO vehicle control to check for solvent toxicity (84). To maintain the concentration of

berberine chloride hydrate at 0.26 mg/ml and reduce the concentration of DMSO as much as

possible, the final concentration of berberine chloride eye drops was 0.26 mg/ml, dissolved in 2%

DMSO. Moreover, a parallel control group of 2% DMSO was also set up in mice to observe its

effect on myopia.

Real-time Quantitative PCR (qPCR)

Real-time Quantitative PCR (qPCR)

After the mice were euthanized, the eyeballs were enucleated immediately to collect retinal,

choroid, and scleral tissues, frozen in liquid nitrogen, and stored at -80 °C for later use. Mouse

total RNA was extracted from the retina, choroid, and scleral tissues using TRIzol (TRI reagent)

(MOR, Miami, FL, USA #TR118). Small RNAs were isolated using RWT (QIAGEN, Hilden,

Germany, #1067933) and RPE buffers (QIAGEN, Hilden, Germany, #1018013). The RNA

samples were dissolved in RNase-free water (TAKARA HOLDINGS Inc., Kyoto, Japan, 9012)

and measured with a spectrophotometer (NanoDrop; Thermo Fisher Scientific, Waltham, MA,

USA). According to the manufacturer's instructions, the extracted RNA was converted to cDNA

after RNA denaturation, DNase reaction, and reverse transcription. mRNA gene expression was

determined using SYBR Green RT-PCR on the cDNA template and PCR was performed using

StepOnePlus Real-Time PCR System (Applied Biosystems, Waltham, Massachusetts, USA). The

mRNA expression of each gene was evaluated using the comparative Ct (ΔΔCT) method and

normalized to glyceraldehyde-3-phosphate dehydrogenase (GAPDH) mRNA expression, which

was used as the reference gene. The following were the qPCR primer sequences:

mouse *Il6* forward: CTACCCCAATTTCCAATGCT

mouse *Il6* reverse: ACCACAGTGAGGAATGTCCA

mouse *Tnfa* forward: CTGTAGCCCACGTCGTAGC

mouse *Tnfa* reverse: TTGAGATCCATGCCGTTG

mouse Mrc1 forward: TCGAGACTGCTGCTGAGTCCA

mouse Mrc1 reverse: AGACAGGATTGTCGTTCAACCAAAG

mouse *Nox2* forward: ACTCCTTGGAGCACTGG

mouse *Nox2* reverse: GTTCCTGTCCAGTTGTCTTCG

mouse *Nox4* forward: TGAACTACAGTGAAGATTTCCTTGAAC

mouse *Nox4* reverse: GACACCCGTCAGACCAGGAA

mouse *Mmp2* forward: CAAGTTCCCCGGCGAT

mouse *Mmp2* reverse: TTCTGGTCAAGGTCAC

mouse *Mmp9* forward: GGACCCGAAGCGGACA

mouse *Mmp9* reverse: CGTCGTCGAAATGGGC

mouse *GAPDH* forward: AGGAGCGAGACCCCACTAAC

mouse *GAPDH* reverse: GATGACCCTTTTGGCTCCAC

8-OHdG ELISA

For assessing oxidative stress, the level of 8-Oxo-2'-deoxyguanosine (8-OHdG) in the choroid

tissues was measured using commercial Enzyme-Linked Immunosorbent Assay (ELISA) kits in

accordance with the manufacturer's instructions (Uscn Life Science; Wuhan, China). Because

this assay uses a competitive inhibition enzyme immunoassay approach, the concentration of 8-

OHdG in the sample and the assay signal strength were inversely correlated. The concentration

of 8-OHdG in each sample was calculated using standard curves generated from standard

proteins, which were calculated and analyzed using Boster's ELISA Online Calculator.

Statistical analyses

Our study used an independent *t*-test and one-way analysis of variance (ANOVA) for statistical

significance analyses of all data (Microsoft Excel 2003, USA). All results are presented as mean

standard deviation (SD), and results with *P* values < 0.05 were considered statistically significant.

Study Approval

The Institutional Animal Care and Use Committee at Keio University authorized all operations

(approval number: 16017). All methods and experimental protocols followed the National

Institutes of Health (NIH) guidelines for working with laboratory animals and the ARVO Animal

Statement for the Use of Animals in Ophthalmic and Vision Research.

Data Availability

All sources of data generated or analyzed during this study are included in this published article

and its supplementary information files.

Author Contributions

Acknowledgments

618　The authors thank Y. Katada, N. Ban, A. Nakai, N. Serizawa, C. Shoda, M. Ziyan, C. Junhan, and

619　A. Kawabata (Graduate School of Medicine, Keio University, Tokyo, Japan) for their technical

620　guidance and administrative support.

References

1.　Hawthorne FA, and Young TL. Genetic contributions to myopic refractive error: Insights

from human studies and supporting evidence from animal models. *Experimental eye*

research. 2013;114:141-9.

2.　Lopes MC, Andrew T, Carbonaro F, Spector TD, and Hammond CJ. Estimating heritability

and shared environmental effects for refractive error in twin and family studies.

Investigative ophthalmology & visual science. 2009;50(1):126-31.

3.　Morgan IG, Ohno-Matsui K, and Saw S-M. Myopia. *The Lancet.* 2012;379(9827):1739-48.

4.　Wojciechowski R. Nature and nurture: the complex genetics of myopia and refractive

error. *Clinical genetics.* 2011;79(4):301-20.

5. Goldschmidt E, and Jacobsen N. Genetic and environmental effects on myopia

development and progression. *Eye.* 2014;28(2):126-33.

6. Lim LS, Gazzard G, Low Y-L, Choo R, Tan DT, Tong L, et al. Dietary factors, myopia, and

axial dimensions in children. *Ophthalmology.* 2010;117(5):993-7. e4.

7. Holden BA, Fricke TR, Wilson DA, Jong M, Naidoo KS, Sankaridurg P, et al. Global

prevalence of myopia and high myopia and temporal trends from 2000 through 2050.

Ophthalmology. 2016;123(5):1036-42.

8. Yotsukura E, Torii H, Inokuchi M, Tokumura M, Uchino M, Nakamura K, et al. Current

prevalence of myopia and association of myopia with environmental factors among

schoolchildren in Japan. *JAMA ophthalmology.* 2019;137(11):1233-9.

9. Fledelius H, Zak M, and Pedersen FK. Refraction in juvenile chronic arthritis: a long‐

term follow‐up study, with emphasis on myopia. *Acta Ophthalmologica Scandinavica.*

2001;79(3):237-9.

10. Herbort CP, Papadia M, and Neri P. Myopia and inflammation. *Journal of ophthalmic &*

vision research. 2011;6(4):270.

11. Kamath Y, Singh A, Bhat S, and Sripathi H. Acute onset myopia as a presenting feature of

systemic lupus erythematosus. *Journal of Postgraduate Medicine.* 2013;59(3):245.

12. Lin H-J, Wei C-C, Chang C-Y, Chen T-H, Hsu Y-A, Hsieh Y-C, et al. Role of chronic

inflammation in myopia progression: clinical evidence and experimental validation.

EBioMedicine. 2016;10:269-81.

13. Fujiwara N, and Kobayashi K. Macrophages in inflammation. *Current Drug Targets-*

Inflammation & Allergy. 2005;4(3):281-6.

14.   Wynn TA, Chawla A, and Pollard JW. Macrophage biology in development, homeostasis

655   and disease. *Nature.* 2013;496(7446):445-55.

656   15.   Epelman S, Lavine KJ, and Randolph GJ. Origin and functions of tissue macrophages.

657   *Immunity.* 2014;41(1):21-35.

658   16.   Chinnery HR, McMenamin PG, and Dando SJ. Macrophage physiology in the eye.

659   *Pflügers Archiv - European Journal of Physiology.* 2017;469(3):501-15.

660   17.   Wang X, Zhao L, Zhang J, Fariss RN, Ma W, Kretschmer F, et al. Requirement for

661   microglia for the maintenance of synaptic function and integrity in the mature retina.

662   *Journal of Neuroscience.* 2016;36(9):2827-42.

663   18.   Santos AM, Calvente R, Tassi M, Carrasco MC, Martín‐Oliva D, Marín‐Teva JL, et al.

664   Embryonic and postnatal development of microglial cells in the mouse retina. *Journal of*

665   *Comparative Neurology.* 2008;506(2):224-39.

666   19.   Rathnasamy G, Foulds WS, Ling E-A, and Kaur C. Retinal microglia–A key player in

667   healthy and diseased retina. *Progress in neurobiology.* 2019;173:18-40.

668   20.   Yu C, Roubeix C, Sennlaub F, and Saban DR. Microglia versus monocytes: distinct roles in

669   degenerative diseases of the retina. *Trends in neurosciences.* 2020;43(6):433-49.

670   21.   Forrester JV, McMenamin PG, Holthouse I, Lumsden L, and Liversidge J. Localization and

671   characterization of major histocompatibility complex class II-positive cells in the

672   posterior segment of the eye: implications for induction of autoimmune uveoretinitis.

673   *Investigative ophthalmology & visual science.* 1994;35(1):64-77.

674   22.   Kumar A, Zhao L, Fariss RN, McMenamin PG, and Wong WT. Vascular associations and

675   dynamic process motility in perivascular myeloid cells of the mouse choroid:

676  implications for function and senescent change. *Investigative ophthalmology & visual*

677  *science.* 2014;55(3):1787-96.

678 23. Yang X, Zhao L, Campos MM, Abu-Asab M, Ortolan D, Hotaling N, et al. Choroid-resident

679  macrophages maintain local vasculature and RPE integrity and spontaneously

680  regenerate following depletion. *bioRxiv.* 2019:868430.

681 24. Yang X, Zhao L, Campos MM, Abu-Asab M, Ortolan D, Hotaling N, et al. CSF1R blockade

682  induces macrophage ablation and results in mouse choroidal vascular atrophy and RPE

683  disorganization. *Elife.* 2020;9:e55564.

684 25. Nishida Y, Fujiwara T, Imamura Y, Lima LH, Kurosaka D, and Spaide RF. Choroidal

685  thickness and visual acuity in highly myopic eyes. *Retina.* 2012;32(7):1229-36.

686 26. Liu Y, Wang L, Xu Y, Pang Z, and Mu G. The influence of the choroid on the onset and

687  development of myopia: from perspectives of choroidal thickness and blood flow. *Acta*

688  *Ophthalmologica.* 2021;99(7):730-8.

689 27. Flores-Moreno I, Lugo F, Duker JS, and Ruiz-Moreno JM. The relationship between axial

690  length and choroidal thickness in eyes with high myopia. *American journal of*

691  *ophthalmology.* 2013;155(2):314-9. e1.

692 28. Zhang S, Zhang G, Zhou X, Xu R, Wang S, Guan Z, et al. Changes in choroidal thickness

693  and choroidal blood perfusion in guinea pig myopia. *Investigative Ophthalmology &*

694  *Visual Science.* 2019;60(8):3074-83.

695 29. Huang X, Li Y, Fu M, and Xin H-B. *Macrophages*. Springer; 2018:119-26.

696 30. Ivashkiv LB. Epigenetic regulation of macrophage polarization and function. *Trends in*

697  *immunology.* 2013;34(5):216-23.

31. Mosser DM, and Edwards JP. Exploring the full spectrum of macrophage activation.

Nature reviews immunology. 2008;8(12):958-69.

32. Sica A, and Mantovani A. Macrophage plasticity and polarization: in vivo veritas. *The*

Journal of clinical investigation. 2012;122(3):787-95.

33. Wei C-C, Kung Y-J, Chen CS, Chang C-Y, Lin C-J, Tien P-T, et al. Allergic conjunctivitis-

induced retinal inflammation promotes myopia progression. *EBioMedicine.*

2018;28:274-86.

34. Zhu K, Yang C, Dai H, Li J, Liu W, Luo Y, et al. Crocin inhibits titanium particle-induced

inflammation and promotes osteogenesis by regulating macrophage polarization.

International Immunopharmacology. 2019;76:105865.

35. Talamonti E, Pauter AM, Asadi A, Fischer AW, Chiurchiù V, and Jacobsson A. Impairment

of systemic DHA synthesis affects macrophage plasticity and polarization: implications

for DHA supplementation during inflammation. *Cellular and Molecular Life Sciences.*

2017;74(15):2815-26.

36. Namgaladze D, and Brüne B. Macrophage fatty acid oxidation and its roles in

macrophage polarization and fatty acid-induced inflammation. *Biochimica et Biophysica*

Acta (BBA)-Molecular and Cell Biology of Lipids. 2016;1861(11):1796-807.

37. Hung L-F, Wallman J, and Smith EL. Vision-dependent changes in the choroidal thickness

of macaque monkeys. *Investigative Ophthalmology & Visual Science.* 2000;41(6):1259-

69.

38. NICKLA DL, WILDSOET C, and WALLMAN J. Visual influences on diurnal rhythms in ocular

length and choroidal thickness in chick eyes. *Experimental eye research.* 1998;66(2):163-

81.

39. Read SA, Alonso-Caneiro D, Vincent SJ, and Collins MJ. Longitudinal changes in choroidal

thickness and eye growth in childhood. *Investigative ophthalmology & visual science.*

2015;56(5):3103-12.

40. Weisser SB, van Rooijen N, and Sly LM. Depletion and reconstitution of macrophages in

mice. *JoVE (Journal of Visualized Experiments).* 2012(66):e4105.

41. Zhang Y, Jeong H, Mori K, Ikeda S-I, Shoda C, Miwa Y, et al. Vascular endothelial growth

factor from retinal pigment epithelium is essential in choriocapillaris and axial length

maintenance. *PNAS Nexus.* 2022;1(4):pgac166.

42. Mullen A, Loscher CE, and Roche HM. Anti-inflammatory effects of EPA and DHA are

dependent upon time and dose-response elements associated with LPS stimulation in

THP-1-derived macrophages. *The Journal of nutritional biochemistry.* 2010;21(5):444-50.

43. Mori K, Kuroha S, Hou J, Jeong H, Ogawa M, Ikeda Si, et al. Lipidomic analysis revealed

n‐3 polyunsaturated fatty acids suppressed choroidal thinning and myopia progression

in mice. *The FASEB Journal.* 2022;36(6):e22312.

44. Yao Y, Xu X-H, and Jin L. Macrophage polarization in physiological and pathological

pregnancy. *Frontiers in immunology.* 2019;10:792.

45. Jiang X, Kurihara T, Kunimi H, Miyauchi M, Ikeda S-i, Mori K, et al. A highly efficient

murine model of experimental myopia. *Scientific reports.* 2018;8(1):1-12.

46. Lin J, Cai Q, Liang B, Wu L, Zhuang Y, He Y, et al. Berberine, a traditional Chinese

medicine, reduces inflammation in adipose tissue, polarizes M2 macrophages, and

increases energy expenditure in mice fed a high-fat diet. *Medical Science Monitor:*

International Medical Journal of Experimental and Clinical Research. 2019;25:87.

47. WANG Q-Z, SHI J, LIU Q, YANG L-X, GUO L, YE C-Y, et al. Berberine promotes M1

proinflammatory phenotype to M2 anti-inflammatory phenotype polarization in

macrophage cell line RAW264. 7. *Basic & Clinical Medicine.* 2019;39(5):646.

48. Virág L, Jaén RI, Regdon Z, Boscá L, and Prieto P. Self-defense of macrophages against

oxidative injury: Fighting for their own survival. *Redox Biology.* 2019;26:101261.

49. Covarrubias A, Byles V, and Horng T. ROS sets the stage for macrophage differentiation.

Cell research. 2013;23(8):984-5.

50. Coviltir V, Burcel M, Cherecheanu AP, Ionescu C, Dascalescu D, Potop V, et al. Update on

myopia risk factors and microenvironmental changes. *Journal of ophthalmology.*

2019;2019.

51. Francisco B-M, Salvador M, and Amparo N. Oxidative stress in myopia. *Oxidative*

Medicine and Cellular Longevity. 2015;2015.

52. Halliwell B. Why and how should we measure oxidative DNA damage in nutritional

studies? How far have we come? *The American journal of clinical nutrition.*

2000;72(5):1082-7.

53. McMenamin PG, Saban DR, and Dando SJ. Immune cells in the retina and choroid: Two

different tissue environments that require different defenses and surveillance. *Progress*

in retinal and eye research. 2019;70:85-98.

54. Arnold T, and Betsholtz C. Erratum to: The importance of microglia in the development

of the vasculature in the central nervous system. *Vascular cell.* 2013;5(1):1-7.

55. McMenamin PG. Dendritic cells and macrophages in the uveal tract of the normal

mouse eye. *British Journal of Ophthalmology.* 1999;83(5):598-604.

56. Lin H-H, Faunce DE, Stacey M, Terajewicz A, Nakamura T, Zhang-Hoover J, et al. The

macrophage F4/80 receptor is required for the induction of antigen-specific efferent

regulatory T cells in peripheral tolerance. *The Journal of experimental medicine.*

2005;201(10):1615-25.

57. Liu Y, Xia Y, and Qiu C-H. Functions of CD169 positive macrophages in human diseases.

Biomedical Reports. 2021;14(2):1-.

58. Linehan E, and Fitzgerald D. Ageing and the immune system: focus on macrophages.

European Journal of Microbiology and Immunology. 2015;5(1):14-24.

59. Mantovani A, Resta A, Herbort CP, Abu El Asrar A, Kawaguchi T, Mochizuki M, et al.

Work-up, diagnosis and management of acute Vogt-Koyanagi-Harada disease.

International Ophthalmology. 2007;27(2):105-15.

60. Chen W, Zhao B, Jiang R, Zhang R, Wang Y, Wu H, et al. Cytokine expression profile in

aqueous humor and sera of patients with acute anterior uveitis. *Current Molecular*

Medicine. 2015;15(6):543-9.

61. Wu P-C, Chuang M-N, Choi J, Chen H, Wu G, Ohno-Matsui K, et al. Update in myopia and

treatment strategy of atropine use in myopia control. *Eye.* 2019;33(1):3-13.

62. Hsu Y-A, Chen C-S, Wang Y-C, Lin E-S, Chang C-Y, Chen JJ-Y, et al. Anti-inflammatory

effects of resveratrol on human retinal pigment cells and a myopia animal model.

Current Issues in Molecular Biology. 2021;43(2):716-27.

63. Ikeda S-I, Kurihara T, Toda M, Jiang X, Torii H, and Tsubota K. Oral bovine milk lactoferrin

administration suppressed myopia development through matrix metalloproteinase 2 in

a mouse model. *Nutrients.* 2020;12(12):3744.

64.   He H, Mack JJ, Güç E, Warren CM, Squadrito ML, Kilarski WW, et al. Perivascular

macrophages limit permeability. *Arteriosclerosis, thrombosis, and vascular biology.*

2016;36(11):2203-12.

65.   Du Cheyne C, Tay H, and De Spiegelaere W. The complex TIE between macrophages and

angiogenesis. *Anatomia, Histologia, Embryologia.* 2020;49(5):585-96.

66.   Muhammad S, Chaudhry SR, Dobreva G, Lawton MT, Niemelä M, and Hänggi D. Vascular

macrophages as therapeutic targets to treat intracranial aneurysms. *Frontiers in*

Immunology. 2021;12:630381.

67.   Raffort J, Lareyre F, Clément M, Hassen-Khodja R, Chinetti G, and Mallat Z. Monocytes

and macrophages in abdominal aortic aneurysm. *Nature Reviews Cardiology.*

2017;14(8):457-71.

68.   Deng Z, Shi F, Zhou Z, Sun F, Sun M-H, Sun Q, et al. M1 macrophage mediated increased

reactive oxygen species (ROS) influence wound healing via the MAPK signaling in vitro

and in vivo. *Toxicology and Applied Pharmacology.* 2019;366:83-95.

69.   Castaneda OA, Lee S-C, Ho C-T, and Huang T-C. Macrophages in oxidative stress and

models to evaluate the antioxidant function of dietary natural compounds. *journal of*

food and drug analysis. 2017;25(1):111-8.

70.   Valentin F, Bueb JL, Kieffer P, Tschirhart E, and Atkinson J. Oxidative stress activates

MMP‐2 in cultured human coronary smooth muscle cells. *Fundamental & clinical*

pharmacology. 2005;19(6):661-7.

71.   Bittner A, Alcaíno H, Castro PF, Pérez O, Corbalán R, Troncoso R, et al. Matrix

metalloproteinase-9 activity is associated to oxidative stress in patients with acute

coronary syndrome. *International journal of cardiology.* 2010;143(1):98-100.

72.   Pérez S, and Rius-Pérez S. Macrophage polarization and reprogramming in acute

811   inflammation: a redox perspective. *Antioxidants.* 2022;11(7):1394.

812   73.   Bhatt L, Groeger G, McDermott K, and Cotter TG. Rod and cone photoreceptor cells

813   produce ROS in response to stress in a live retinal explant system. *Molecular vision.*

814   2010;16:283.

815   74.   Fujikado T, Kawasaki Y, Fujii J, Taniguchi N, Okada M, Suzuki A, et al. The effect of nitric

816   oxide synthase inhibitor on form-deprivation myopia. *Current eye research.*

817   1997;16(10):992-6.

818   75.   Fujikado T, Tsujikawa K, Tamura M, Hosohata J, Kawasaki Y, and Tano Y. Effect of a nitric

819   oxide synthase inhibitor on lens-induced myopia. *Ophthalmic Research.* 2001;33(2):75-

820   9.

821   76.   Li Y, Zhang Y, Zhang P, Gao L, Ma Q, Li J, et al. Genetic susceptibility to high myopia in

822   Han Chinese population. *Open Life Sciences.* 2022;17(1):512-6.

823   77.   Zhu Y, Li X, Chen J, Chen T, Shi Z, Lei M, et al. The pentacyclic triterpene Lupeol switches

824   M1 macrophages to M2 and ameliorates experimental inflammatory bowel disease.

825   *International immunopharmacology.* 2016;30:74-84.

826   78.   Wang Z, Zou J, Huang Y, Cao K, Xu Y, and Wu JM. Effect of resveratrol on platelet

827   aggregation in vivo and in vitro. *Chinese medical journal.* 2002;115(03):377-80.

828   79.   Hu J, Deng F, Zhao B, Lin Z, Sun Q, Yang X, et al. Lactobacillus murinus alleviate intestinal

829   ischemia/reperfusion injury through promoting the release of interleukin-10 from M2

830   macrophages via Toll-like receptor 2 signaling. *Microbiome.* 2022;10(1):1-21.

831   80.   Zhao L, Wang H, and Du X. The therapeutic use of quercetin in ophthalmology: Recent

832   applications. *Biomedicine & Pharmacotherapy.* 2021;137:111371.

833   81.   Wagh V, Patil P, Surana S, and Wagh K. Forskolin: upcoming antiglaucoma molecule.

834   *Journal of postgraduate medicine.* 2012;58(3):199.

835   82.   Dysli C, Enzmann V, Sznitman R, and Zinkernagel MS. Quantitative analysis of mouse

836   retinal layers using automated segmentation of spectral domain optical coherence

837   tomography images. *Translational vision science & technology.* 2015;4(4):9-.

838   83.   Hou J, Mori K, Ikeda S-i, Jeong H, Torii H, Negishi K, et al. Ginkgo biloba extracts improve

839   choroidal circulation leading to suppression of myopia in mice. *Scientific Reports.*

840   2023;13(1):3772.

841   84.   Galvao J, Davis B, Tilley M, Normando E, Duchen MR, and Cordeiro MF. Unexpected

842   low‐dose toxicity of the universal solvent DMSO. *The FASEB Journal.* 2014;28(3):1317-

843   30.

Figures and Tables

Figure 1. Myopia is accompanied by sustained depletion of choroidal-resident macrophages in mice at 3 and 8 weeks of age.

(**A**) Three-weeks-old wild-type C57BL/6JJc1 mice were injected with either 0.1 ml/10 g body weight clodronate liposome or control liposome once every two days (n=12 per group). Refraction, axial length, and choroidal thickness were measured at baseline and 8 days after 4 injections with an infrared photorefractor and SD-OCT system. **(B)** Compared to the control group, the clodronate liposome (clo-lip) injection group showed a larger refractive change ($P <$ 0.001), axial eye growth ($P <$ 0,01), and thinner choroid ($P <$ 0.001). Bars represent mean +/− standard deviations. **(C)** The percentage of macrophages in the choroid was analyzed using flow cytometry. Macrophages were identified as CD11b[+] and F4/80[+], appearing in the H3-UR region. Compared with the control liposome injection group, the proportion of macrophages in the clodronate liposome injection group was significantly reduced ($P <$ 0.05). **(D)** Eight-weeks-old

wild-type C57BL/6JJc1 mice (4 per group unless otherwise indicated) were injected with 0.1
ml/10 g body weight clodronate liposome and control liposome once every two days. Refraction,
axial length, and choroidal thickness were measured at baseline and 8 days after 4 injections with
an infrared photorefractor and SD-OCT system. **(E)** Compared to the control group, the
clodronate liposome (clo-lip) injection group showed a remarkably larger refractive change ($P <$
0.01), axial length elongation ($P < 0.05$), and a thinner choroid ($P < 0.01$). Bars represent
mean +/− standard deviations. Differences between the groups were compared using a t-test. *P
< 0.05. **$P < 0.01$. ***$P < 0.001$.
888

891   **Figure 2. The polarization of M1 macrophages in the choroid can induce myopia in a**
892   **murine model.**
893

(A) Three-weeks-old wild-type C57BL/6JJc1 mice were divided into two groups (4 per group
unless otherwise indicated) and one group was intraperitoneally injected with 10 ul/g LPS and the
other with PBS. The injection lasted for 2 weeks, and refraction, axial length, and choroidal
thickness were measured weekly with an infrared photorefractor and SD-OCT system. Choroidal
blood perfusion was measured with OCTA. In addition, choroid samples were collected for real-
time PCR. **(B)** Compared to the PBS injection group, the LPS injection group showed a
significantly larger refractive change ($P < 0.001$), significant axial eye growth ($P < 0,001$), and
substantially thinner choroid ($P < 0.001$). Bars represent mean +/− standard deviations. **(C)** The
choroidal blood perfusion decreased significantly after 2 weeks of LPS injection, compared to the

PBS injection group. **(D)** mRNA levels of *Tnf-α* and *IL-6* significantly increased after 2 weeks of
LPS injection. *P* values were obtained from a t-test. *$P < 0.05$. **$P < 0.01$. ***$P < 0.001$.

Figure 3. Inhibition of myopia progression by choroidal M2 macrophage polarization was
demonstrated in a lens-induced myopia mouse model.

Three-week-old wild-type C57BL/6JJc1 mice were injected with 0.1 ug/100 ul IL-4, and choroid
samples were collected at 0 h, 24 h, and 48 h after injection for flow cytometry and real-time
PCR. **(A)** Macrophages in the H3-UR region were denoted as CD11b$^+$ and F4/80$^+$. The proportion
of macrophages increased significantly after IL-4 injection ($P < 0.001$). M2 macrophages in the
Q2-UR and were identified as CD206$^+$. At 48 h after IL-4 injection, M2 macrophages were
dramatically polarized ($P < 0.05$). **(B)** Real-time PCR results showed that the mRNA expression
levels of *Mrc1* and *CD163* increased significantly 48 h after IL-4 injection. **(C)** Three-weeks-old
wild-type C57BL/6JJc1 mice were divided into two groups and injected with either 0.1 ug/100 ul
IL-4 or PBS on alternate days with myopia induction. Refraction, axial length, and choroidal
thickness were measured at the initial (3-week-old) and final (6-week-old) stages of myopic
induction using an infrared photorefractor and SD-OCT system. **(D)** Eyes treated with −30D
lenses showed a significantly larger refractive change ($P < 0.01$), greater axial length elongation
($P < 0.05$), and increased choroidal thickness ($P < 0.001$) than those treated with 0D lenses in the
PBS injection mice. Mice injected with 0.1 ug/100 ul IL-4 showed a significantly smaller refractive
change ($P < 0.05$), smaller change in axial length($P < 0.05$), and positive change in choroidal
thickness with a statistical significance ($P < 0.001$). Bars represent the mean +/− standard
deviation. **(E)** To evaluate changes in choroidal blood perfusion, three-weeks-old mice were
divided into three groups: both eyes with 0D lenses (control 0D group), binocular myopia
induction (control -30D group), and binocular myopia induction with IL-4 injection once every two
days group (IL-4 -30D group). Choroidal blood perfusion was assessed at the initial (3-week-old)
and end (6-week-old) stages of myopic induction using OCTA. Compared to the control 0D group,

choroidal blood perfusion was significantly decreased in the control -30D group. Accordingly,
compared with the control -30D group, choroidal blood perfusion was remarkably improved in the
IL-4 -30D group. **(F)** Mice used to measure choroidal blood perfusion were euthanized at 6 weeks
of age, and the choroids were collected for flow cytometry. The results showed that mice treated
with -30D lenses and injected with IL-4 had a significant proportion of M2 macrophages
compared to those treated with -30D lenses and injected with PBS. *P* values correspond to
comparisons made using one-way ANOVA. *$P < 0.05$. **$P < 0.01$. ***$P < 0.001$.

Figure 4: Effect of macrophage polarization direction on oxidative stress response.

(A) Experimental plan for the assessment of *Nox2, Nox4, Mmp2,* and *Mmp9* mRNA expression in the choroid during M1 macrophage polarization and M2 macrophage polarization. Three-weeks-old wild-type C57BL/6JJc1 mice were injected with LPS to polarize M1 macrophages and with IL-4 to polarized M2 macrophages, while the control group was injected with PBS. Choroid samples were collected 24 h after LPS injection, 24 h and 48 h after IL-4 injection, and after PBS injection for real-time PCR and ELISA experiments. **(B)** Comparison of *Nox2, Nox4, Mmp2,* and *Mmp9* mRNA expression in the choroid between the control and M1 macrophage polarization group

showed a significant increase in the LPS-injected group. **(C)** Quantitative PCR analyses of the
choroid 48 h after IL-4 injection indicated a corresponding reduction in the expression
of *Nox2, Mmp2,* and *Mmp9* mRNA, corresponding to M2 macrophage polarization. **(D)** ELISA
analysis results showed that compared to the control group, the choroidal 8-OHdG concentration
was decreased in the IL-4 injection group ($P < 0.01$). On the contrary, the choroidal 8-OHdG
concentration was increased in the LPS injection group, compared to the IL-4 injection group ($P < 0.05$). *P* values indicate comparisons relative to the control, one-way ANOVA, or t-test. *$P < 0.05$.
0.05). *P* values indicate comparisons relative to the control, one-way ANOVA, or t-test. *$P < 0.05$.
**$P < 0.01$. ***$P < 0.001$.
955

Supplementary Figure 1: EPA administration inhibited the development of myopia caused by LPS injection.

Three-weeks-old wild-type C57BL/6JJc1 mice were divided into three groups: control group, LPS injection group, and LPS administered with EPA group (n=4). Changes in refraction, axial length, and choroidal thickness were measured after 3 weeks of feeding. **(A)** Compared to the large changes in refraction caused by LPS injection, the refractive shift was smaller after EPA administration ($P < 0.01$). **(B)** Compared to the increase in the ratio of axial length to body weight caused by LPS injection, EPA feeding reduced the ratio of axial length to body weight ($P < 0.001$). **(C)** Thinning of choroid occurred in the LPS injection group, whereas EPA feeding significantly improved choroidal thickness ($P < 0.05$). P values indicate comparisons relative to the control using one-way ANOVA. *$P < 0.05$. **$P < 0.01$. ***$P < 0.001$.

Supplementary Figure 2: IL-13 injection suppressed myopia progression and increased choroidal blood perfusion in a murine model of lens-induced myopia.

(A) Choroid samples were collected after 48 h of IL-13 injection and PBS injection for flow cytometry analysis, which was designed to evaluate the optimal time for the polarization of M2 macrophages. **(B)** Flow cytometry analysis demonstrated that both macrophage ratio and M2 macrophage ratio were significantly increased 48 h after IL-13 injection, compared to the PBS injection group. **(C)** Three-week-old mice were divided into three groups: 0D lenses on both eyes (control 0D group), binocular myopia induction (control -30D group), and binocular myopia induction and IL-13 injection (IL-13 -30D group). IL-13 injection frequency was once every two days. Refraction, axial length, and choroidal thickness were measured at the initial (3-week-old) and end (6-week-old) stages of myopic induction with an infrared photorefractor and SD-OCT system. **(D)** The control -30D group showed a significantly larger refractive change ($P < 0.001$), greater axial length elongation ($P < 0.05$), and thinner choroidal ($P < 0.001$) compared to the control 0D group. The IL-13 -30D group indicated a significantly smaller refractive change ($P < 0.001$), lesser axial length change ($P < 0.05$), and positive change in choroidal thickness, with a statistical significance ($P < 0.001$). Bars represent mean +/− standard deviations. **(E)** Choroidal blood perfusion was measured using OCTA. Compared to the control 0D group, choroidal blood perfusion was significantly decreased in the control -30D group. Correspondingly, compared to the control -30D group, choroidal blood perfusion was significantly improved in the IL-13 -30D group. **(F)** The proportion of M2 macrophages in the choroid showed a significant increase in the IL-13 -30D group, compared to the control -30D group. P values indicate comparisons relative to control with one-way ANOVA or t-test. *$P < 0.05$. **$P < 0.01$. ***$P < 0.001$.

998 **Supplementary Figure 3: Continuous berberine eye drops can inhibit the progression of**
999 **myopia in mice.**
1000
Three-weeks-old wild-type C57BL/6JJc1 mice were separated into control 0D group, control -30D
group, 2% DMSO -30D group, and berberine -30D group (n=4). The frequency of eye drops for
2% DMSO and 0.26 mg/ml berberine was once a day. After 3 weeks of eye drops, the changes in
refraction, axial length, and choroidal thickness were measured using SD-OCT. **(A)** Compared to
the control 0D group, a refractive shift was observed in the control -30D group and 2% DMSO
group ($P < 0.001$). Smaller refractive changes were observed in the berberine -30D group
compared to the 2% DMSO group ($P < 0.001$). **(B)** Compared to the growth of axial length in the
control -30D group and 2% DMSO group ($P < 0.05$), the berberine -30D group showed a smaller
axial length elongation ($P < 0.05$). **(C)** In the control -30D group and 2% DMSO -30D group, the
choroidal thickness was reduced compared to the control 0D group ($P < 0.001$). Contrarily,
choroid thickness was improved in the berberine -30D group compared to the 2% DMSO -30D
group ($P < 0.001$). P values indicate comparisons relative to the control using one-way ANOVA.
*$P < 0.05$. **$P < 0.01$. ***$P < 0.001$.
1014