How many variables does WorldClim have, truly? Generative A.I. unravels the intrinsic

dimension of bioclimatic variables.

Russell Dinnage

Affiliations: 1) Department of Biological Sciences, University of Alberta, Edmonton, Alberta,

Canada; 2) Alberta Machine Learning Institute, Edmonton, Alberta, Canada; 3) Institute of

Environment, Department of Biological Science, Florida International University, Miami,

Florida, USA.

Corresponding author: Russell Dinnage, dinnage@ualberta.ca

Key words: Bioclimatic Variables, Dimension Reduction, Manifold Hypothesis, Variational

Autoencoder (VAE), Generative A.I., Species Distribution Models (SDM), Climate Data

Analysis, WorldClim Dataset, Ecological Data Interpretation

Abstract

The 19 standard bioclimatic variables available from the WorldClim dataset are some of the most

used data in ecology and organismal biology. It is well known that many of the variables are

correlated with each other, suggesting there are fewer than 19 independent dimensions of

information in them. But how much information is there? Here I explore the 19 WorldClim

bioclimatic variables from the perspective of the manifold hypothesis: that many high

dimensional datasets are actually confined to a lower dimensional manifold embedded in an

ambient space. Using a state-of-the-art generative probabilistic model (variational autoencoder)

to model the data on a non-linear manifold reveals that only 5 uncorrelated dimensions are

adequate to capture the full range of variation in the bioclimatic variables, with a clear data-

driven separation between informative and redundant dimensions that eliminates arbitrary

thresholds. I show that these 5 variables have meaningful structure and are sufficient to produce

species distribution models (SDMs) nearly as good and in some ways better than SDMs using

the original 19 bioclimatic variables. I have made the 5 synthetic variables available as a raster

dataset at 2.5 minute resolution in an R package that also includes functions to convert back and

forth between the 5 variables and the original 19.

Introduction

The WorldClim dataset, often referred to as bioclim, is a widely utilized resource providing

detailed global climate data suitable for a variety of ecological and biological applications (Fick

and Hijmans 2017). The dataset comprises 19 bioclimatic variables, each of which represents a

different aspect of annual trends, seasonality, and extreme or limiting environmental factors.

These variables are second-order summaries derived from just three first-order monthly

variables: minimum temperature, maximum temperature, and precipitation – via formulae

computing means, extremes, ranges, and seasonal aggregates, providing more biologically

meaningful summaries of environmental conditions than the raw monthly values alone. The set

of 19 variables were selected over the period 1984-1996 as part of the development of

BIOCLIM, the first species distribution modeling package (Booth et al. 2014), and this particular

selection has remained the standard set of bioclimatic variables used in ecological studies ever

since, despite lacking a formal analysis of their dimensionality.

The use of bioclim variables in species distribution modeling is well documented in the

literature. These models frequently employ bioclim data to establish the ecological niche of a

species, linking occurrences to the environmental conditions under which they are found (Elith

and Leathwick 2009). One reason the dataset is so widely used, besides its easy and wide

accessibility, is that equivalent variables have been estimated for past and future climates based

on paleoclimatic reconstructions and future climate model projections respectively. This makes it

straightforward to use species distribution models for prediction into the past and future,

especially to predict shifts in species distributions under climate change scenarios, providing

important information for conservation planning (Pearson et al. 2006). However, the reliability

of SDM predictions can be sensitive to the choice and number of input variables, and high-

dimensional correlated predictors can lead to spurious model performance (Fourcade et al. 2018).

However, a critical consideration when using bioclim variables is their inherent multicollinearity,

due to the shared underlying data used in their computation. Techniques such as Principal

Component Analysis (PCA), Variance Inflation Factor (VIF) analysis, and Regularization

methods have been employed to address this issue, with the aim of producing more reliable and

interpretable model outcomes (Dormann et al. 2013, Kriticos et al. 2014).

More generally, though each of the 19 bioclimatic variables measure an independently

meaningful and potentially useful aspect of climate, when put together they likely contain

considerable redundancy of information. Another way of putting this is that the 'true' or intrinsic

dimension of the data is less than the total number of dimensions in the data, which can lead to

problems for inference and prediction. For example, too many partially redundant variables can

lead to variance inflation, difficulty in assessing the importance of individual variables,

overfitting on noise, and decreased generalization ability in models (James et al. 2013). One way

to deal with too many variables using dimension reduction methods such as Principle

Components Analysis (PCA: (Pearson 1901)) to reduce the dimensionality of the dataset, a

frequently used strategy when using bioclim data. Using PCA on bioclim can reduce the first

95% of variance into 10 orthogonal variables, but a closer look at the PCA results suggests that a

linear dimension reduction method is inadequate. This is apparent from the fact that the

cumulative variance explained by each additional dimension forms a smooth relationship (Figure

1), whereas we expect a discontinuous 'jump' or 'elbow' in this plot in the case of a true linear

manifold (Cattell 1966). As I show below, a variational autoencoder provides a principled, data-

driven alternative: latent dimensions either contribute to reconstruction (low posterior variance)

or collapse to the prior (high variance), yielding an unambiguous separation without arbitrary

thresholds.

Manifold learning is an emerging concept rooted in the manifold hypothesis, which postulates

that high-dimensional real-world data lie on or near a lower-dimensional manifold embedded

within the high-dimensional space (Bengio et al. 2012). This hypothesis forms the basis for a

wide variety of advanced dimension reduction and data analysis techniques, which include the

new and exciting class of models known as deep generative learning. Classic PCA can also be

considered a simple method of manifold learning, but where the manifold is assumed to be linear

(e.g. a hyperplane). Here I explore the use of one deep generative model known as a variational

autoencoder (VAE: (Kingma and Welling 2013)) for analysing bioclim variables, a simple but

nevertheless useful example which I hope also demonstrates the exciting greater potential of

VAEs and other generative models for ecology.

Methods

Variational Autoencoders

Variational autoencoders (VAEs) are a class of generative models that employ deep learning

techniques for manifold learning (Kingma and Welling 2013). VAEs offer a probabilistic

approach for representing complex, high-dimensional data in a lower-dimensional latent space.

These models are built on the principles of autoencoders, which are neural networks designed to

learn a low dimensional representation of input data, through the use of a paired encoder and

decoder model, that respectively encode the input into a latent space, and decode the latent space

into the original input space (Hinton and Salakhutdinov 2006). VAEs introduce a layer of

probabilistic inference, framing the encoding and decoding process as a probabilistic graphical

model and utilizing Bayesian variational inference for learning and sampling (Kingma and

Welling 2013).

Notably, VAEs encourage a specific structure on the latent space by assuming a simple Bayesian

prior distribution for the latent space: an an independent Gaussian distribution for each latent

variable. This promotes a continuous and well-structured latent space that facilitates the learning

of high-dimensional manifold structures (Doersch 2016). Dai and Wipf 2019 and Zheng et al.

2023 recently showed that manifold dimensions can be identified by latent variables whose

posterior distributions have a low variance.

VAEs have begun to receive attention in molecular biology for a variety of uses such as

visualisation (Battey et al. 2020), estimating the probability density of sequence variants (Ziegler

et al. 2023), generating candidate functional proteins (Hawkins-Hooker et al. 2020), and

estimating reduced representations of complex metabolomics data useful for downstream

predictive modeling. (Gomari et al. 2022), but they have yet to widely used in ecology. I hope

this study, in addition to providing interesting new insights to a classical dataset used in ecology,

can act as a concise introduction to VAEs for ecologists, and to promote their wider use, as their

potential for ecology is substantial.

VAEs work by estimating two key functions, which are approximated using neural networks: an

encoder, and a decoder.

The encoder, also known as the recognition or inference model, takes the input data and provides

a probabilistic encoding in the latent space. Let's say we have a dataset $X = x_1, x_2, \ldots, x_n$ and we

wish to encode it into a latent variable z. The encoder would provide us with a distribution

$q_\phi(z|x)$ over the latent variable, given the input data.

The decoder, also known as the generative or the likelihood model, takes a point in the latent

space and generates a sample in the data space. It provides us with a distribution $p_\theta(x|z)$ over

the data given a latent variable.

The objective of VAE is to maximize the evidence lower bound (ELBO) on the log-likelihood of

the data:

$$\log p_\theta(x) \geq \mathbb{E}_{q_\phi(z|x)}[\log p_\theta(x|z)] - D_{KL}(q_\phi(z|x)||p_\theta(z))$$

The first term on the right-hand side represents the expected reconstruction error – it encourages

the VAE to reconstruct the input data accurately – and where the expectation is estimated using

Monte Carlo sampling. The second term represents the Kullback-Leibler (KL) divergence

between the approximate posterior $q_\phi(z|x)$ and the prior $p_\theta(z)$, which is typically chosen to be a

independent normal distribution. This term acts as a regularization, encouraging the distribution

of the latent variables to stay close to a unit Gaussian.

Training a VAE involves finding the parameters θ and ϕ that maximize the ELBO using

stochastic gradient ascent, enabled by the reparameterization gradient, often known as the

"reparameterization trick" (Kingma and Welling 2013). The reparameterization trick allows a

gradient calculation through the expectation in the ELBO, which is normally blocked by the non-

differentiability of a random sampling operation. It works by reparameterizing the independent

normal distribution $z \sim \text{Normal}(\mu, \sigma^2)$ to $z = \mu + \sigma \cdot \varepsilon$, where $\varepsilon \sim \text{Normal}(0, 1)$. ε is sampled from

a standard normal and then is injected into the model. Since the sampled stochastic noise is now

an external input to the computational graph, the gradient no longer needs to pass through the

sampling operation, allowing full backpropagation. After training a VAE, new data can be

generated by sampling from the prior $p_\theta(z)$, then passing these samples through the decoder.

A particularly useful aspect of VAEs for manifold learning is that maximizing the ELBO

encourages the model to converge on a minimal set of 'active' latent variables to explain the

data. A VAE learns an approximate Bayesian posterior distribution for each of the latent

variables. Variables that contribute strongly to the reconstruction of the data tend to have low

variance. On the other hand, any variables that are not important for reconstructing data will tend

to have their variances converge to 1, consistent with the prior distribution, and facilitated by the

flexibility of the decoder function, which can learn to ignore these high variance dimensions.

Therefore, the number of active dimensions can be deduced by the number of variables with low

variance posteriors (close to zero). Recent work has shown this to be a good estimate of the true

manifold dimension of simulated data (Dai and Wipf 2019; Zheng et al. 2023).

The encoder and decoder neural networks were generated using the in-development dagnn R

package (https://github.com/rdinnager/dagnn), which is designed to make specifying complex

neural network architecture simple by specifying a directed acyclic graph (DAG) using R

formula syntax.

Table 1. Glossary of Variational Autoencoder related terms.

Term	Description

Generative Models	A class of models that learn the full process generating a dataset, not just how to predict an outcome. Unlike models that only learn input-output relationships (e.g. a GLM predicting species presence from climate), generative models can also create (simulate) new realistic data points, because they capture the overall structure of the data.
Autoencoder (AE)	A neural network that learns to compress data into a small number of variables (encoding) and then reconstruct the original data from them (decoding). The compressed representation captures the most important patterns. A deterministic precursor to VAEs that lacks the probabilistic framework.
Variational Inference	A method for approximating complex probability distributions that would be too expensive to compute exactly. In VAEs, it is used to estimate the most likely values of the latent (hidden) variables given the observed data, analogous to how Bayesian methods estimate posterior distributions but using optimisation rather than sampling (e.g. MCMC).
Kullback-Leibler Divergence (KL)	A measure of how much one probability distribution differs from another. In VAEs, it measures how far the learned latent variable distributions deviate from the assumed prior (a standard normal distribution), acting as a penalty that keeps the latent space well-structured and prevents the model from encoding too much specific

	information about individual data points.
Regularization	Any technique that constrains a model during training to prevent it from fitting noise in the data (overfitting). In VAEs, the KL divergence term acts as a regularizer: it penalises latent distributions that deviate far from the prior, encouraging the model to use only as many latent dimensions as are truly needed to represent the data.
Evidence Lower Bound (ELBO)	The objective function maximised during VAE training. It balances two competing goals: (1) accurately reconstructing the input data (reconstruction loss) and (2) keeping the latent variable distributions simple and close to the prior (KL divergence). The name comes from the fact that maximising it approximates maximising the probability of the observed data (the "evidence") under the model.
Reparameterization Trick	A mathematical technique that enables gradient-based training of VAEs despite the random sampling step. Instead of sampling latent values z directly from a learned distribution, we compute z = mean + standard_deviation x noise, where the noise is drawn from a fixed standard normal distribution. This separates the randomness from the learnable parameters (mean and standard deviation), allowing the training algorithm to compute how to adjust them.
Manifold	A smooth, lower-dimensional surface embedded within a higher-

	dimensional space. For example, the surface of the Earth is a 2-dimensional manifold in 3-dimensional space. The manifold hypothesis proposes that complex high-dimensional data (such as 19 bioclim variables) actually lie on or near such a lower-dimensional surface, meaning fewer variables are needed to describe the data than the number of measured variables.
Latent Variables / Latent Space	Hidden (unobserved) variables that a model infers from the data. In a VAE, the latent space is the low-dimensional representation that the encoder produces and the decoder reconstructs from. Each point in the latent space corresponds to a particular combination of the original variables. "Active" latent variables are those the model actually uses; inactive ones are ignored by the decoder.
Posterior Distribution	In Bayesian statistics, the probability distribution of unknown quantities (e.g. latent variables) after accounting for the observed data. In a VAE, for each data point the encoder estimates a posterior distribution over the latent variables, characterised by a mean and variance for each latent dimension.

Model Training

I trained a VAE model coded in R with the torch package (Falbel and Luraschi 2022) on 19 global bioclimatic variables at 2.5 minute resolution, for a total of 12,532,612 data points, one for each global grid-cell. Full code to reproduce the model and all analyses in this study is available at: https://github.com/rdinnager/bioclim_intrinsic_dimension. I allowed for a maximum of 64 latent dimensions. The bioclim variables were standardised to zero mean and unit standard deviation before model input. Key hyperparameters included: 16 initial latent dimensions, a 3-layer encoder and decoder with skip connections, and the trainable precision parameter 'gamma' from Dai and Wipf (2019). The model was trained for 1000 epochs using the Adam optimiser with a one-cycle learning rate schedule (maximum learning rate 0.005). The VAE was trained using the procedure recommended in (Dai and Wipf 2019), which includes as trainable 'gamma' parameter. 'Gamma' in this case is the variance of the Gaussian distribution used to calculated the log likelihood of the VAE reconstruction. This gamma parameter tends to decrease during training as the reconstruction gets better, which has the effect of increasing the weighting of the reconstruction loss relative to the KL loss as the training goes on. Dai and Wipf (2019) showed this leads to a much better reconstruction whilst still maintaining a well-structured latent space.

Interpretation

To explore how to interpret the manifold variables extracted by the VAE, I looked at how the they were related to the original 19 variables, as well as how they were represented within the world's major biomes, using data from Dinerstein et al. (2017).

Comparing Species Distribution Models

I ran three sets of species distribution models (SDMs) using the *ENMTools* R package (Warren

et al. 2021): 1) SDMs using all 19 original bioclim variables; 2) SDMs using the first 10 PCA

axes of the 19 bioclim variables (computed as part of this study, explaining ~95% of total

variance; Figure 1); and 3) SDMs using the 5 manifold variables estimated in this study. This

was done using three different SDM methods: Random Forest, GLM, and the Bioclim model.

The three sets of models were run on occurrence data taken from (Elith et al. 2020), which is

available in the R package *disdat.* I used 5 out of the 6 datasets provided by Elith et al. (2020)

(the 6th proved too large to use easily in the parallelized pipeline developed for this study). The 5

datasets contained a total of 206 species from 5 different areas:

1) Australian Wet Tropics (Queensland, Australia): 20 bird species, and 20 plant species

2) Ontario, Canada: 30 bird species

3) New South Wales, Australia: 7 bat species, 10 bird species, 29 plant species, 8 small reptile

species

4) New Zealand: 52 vascular plant species

5) Switzerland: 30 tree species

For each species and model combination I ran 10 replicate SDMs using ENMTools, holding out

a random 10% of the occurrence points in each replicate. I then calculated the Reciever-Operator

Curve Area Under the Curve metric (AUC) of model fit on model predictions for the held out

10%. I also calculated the True Skill Statistic (TSS) as an alternative measure of model fit.

Additionally, I used the *ENMTools*' env.evaluate() function to calculate AUC and TSS in

195 environmental space. This uses latin hypercube sampling in environmental space to sample

196 background points which can be used in the same way that geographic background points are

197 used to estimate the ability of the model to predict presence or absence in environmental space.

198 To compare the bioclim manifold variables and the bioclim PCA variables I calculated a metric

199 ΔAUC for each replicate of each species and model, which is the different between the AUC

200 metric of the focal SDM (manifold or PCA) and the baseline model, which in this case is the

201 corresponding model run on all 19 original bioclim variables.

202 I report $\overline{\Delta\text{AUC}}$ in the main text, which is the average ΔAUC over all replicates, models and

203 species. In order to test whether this value was significantly different than zero, I fit a Linear

204 Mixed Model (LMM) using the *lme4* R package, including the variable type (manifold vs. PCA)

205 as a fixed effect and species, area (e.g. New Zealand vs. Canada), and model as random effects.

206 The reported $\overline{\Delta\text{AUC}}$ values are the estimated marginal means from this model (calculated using

207 the *emmeans* R package). The same was done for the True Skill Statistic.

208 Lastly, the whole above procedure was repeated but using a simple spatial block technique (in

209 which the study area is partitioned into geographic blocks and entire blocks are withheld for

210 testing, rather than random individual points) for generating the hold-out sample instead of a

211 simple random sample. For this I used the spatial block sampling option in ENMTools model

212 fitting function. This tests whether models can generalise to new geographic areas, a critical

213 requirement for SDMs used in conservation planning.

Results

The Instrinsic Dimension of WorldClim

After model training, examining the posterior distribution of the 64 latent dimensions revealed only 5 had low variance (Figure 2), suggesting an intrinsic dimensionality of 5. Using only these 5 variables to reconstruct the original data through the trained decoder, the reconstruction error is extremely low, with a mean squared error of reconstruction of 0.0036 (on the standardised variables, so in units of standard deviations).

Using SDM benchmark data from (Elith et al. 2020), and three different SDM methods (random forest, glm, and bioclim) from the ENMTools R package (Warren et al. 2021), I additionally tested whether these 5 'latent' variables are useful by comparing species distribution models fit with them with models fit on all 19 original variables, as well as the first 10 PCA dimensions as a comparison. For the 5 manifold dimensions and the PCA axes I calculated the model performance using $\overline{\varDelta\mathrm{AUC}}$, defined as the average difference in AUC between the focal model and the corresponding model using all 19 original variables, averaged over all species and methods tested. I find the 5 variables perform slightly less well (but not significantly) at predicting held-out test samples in geographic space (i.e. at spatially withheld occurrence locations) according to AUC ($\overline{\varDelta\mathrm{AUC}}$ = -0.037 [CI: -0.076, 0.002]; -0.02 for PCA [CI: -0.058, 0.019]; Figure S1) but perform considerably better at predicting them in environmental space (i.e. at background points sampled across the range of environmental conditions rather than geographic locations; Warren et al. 2018) ($\overline{\varDelta\mathrm{AUC}}$ = +0.11 [CI: 0.071, 0.15] , +0.047 for PCA [CI: 0.0096, 0.087]; Figure S1) (Warren et al. 2018). The same analysis using the true skill statistic (TSS) instead of AUC gave similar qualitative results (Figure S2). I suspected the

slightly worse performance in geographic space was because with fewer variables, models could

not overfit as strongly to approximate spatial structure in the occurrence data. To test this I

repeated the SDMs using a simple spatial block method for holding out test samples. This

confirmed my suspicion, with the reduction in AUC for spatially held-out test data being much

smaller and again, when evaluated in geographic space (i.e. predicting at spatially withheld

locations), the confidence interval for the mean delta-AUC included zero, indicating no

statistically significant difference from the full 19-variable models ($\overline{\Delta AUC}$ = -0.017 [CI: -0.043,

0.0081]; Figure S3), but still much improved in environmental space ($\overline{\Delta AUC}$ = +0.097 [CI:

0.071, 0.12]; Figure S3).

The 5 manifold dimensions learned by the VAE in this study have been provided in an R

package (https://github.com/rdinnager/biocman), which also provides function for converting

between the 5 manifold variables and the original 19 bioclimatic variable (e.g. functions for

encoding and decoding).

Discussion

Interpreting the Manifold Dimensions

Examining the 5 active variables shows that each has a unique global pattern (Figure 3) and link

to the original 19 variables (Figures S4-S8). Three of the variables admit a fairly straightforward

interpretation, BIOCMAN1, BIOCMAN2, and BIOCMAN4 – their values have a mostly

monotonic relationships with the original 19, and their global distribution maps show clear links

to important global climate factors, BIOCMAN1 being elevation, BIOCMAN2 being areas

supporting rainforest, including temperate rainforest, and BIOCMAN4 being latitude. The

remaining two are less straightforward, with some relationships with both the original 19

variables being unimodal or 'U-shaped'. On the other hand, BIOCMAN5 appears to clearly

differentiate areas with monsoonal climates with high values. BIOCMAN3 has the least clear

interpretation but globally low values seem to be associated with arid and semi-arid ecoregions

(Figure 1; Figure S6).

For the application of species distribution modeling and other ecological models where

interpretation of a model's coefficients might be of interest, the issue of challenging

interpretability can be mitigated using the decoder function, which allows a straightforward way

to convert the manifold variables back to the original data space. In this way they are no less

interpretable than using a transformation to make a variable better conform to model

assumptions (e.g. log or logistic transformations to reduce covariate skew). A simple procedure

to interpret model coefficients when using bioclim manifold variables as predictors is to output

predictions of the model across a range of manifold variable values of interest, then plot these

predictions against the original 19 variables after back-transforming the manifold variables using

the decoder. This then places the model prediction back into the context of full bioclimatic

variation. For instance, if an SDM shows a strong effect of BIOCMAN1, the decoder reveals this

corresponds primarily to variation in temperature-related variables (e.g. BIO1, Annual Mean

Temperature), allowing interpretation in terms of familiar bioclimatic variables. This approach is

analogous to examining PCA loadings, which map principal components back to original

variables. However, PCA loadings are single numbers, a direct consequence of the linear

assumption, making them easy to tabulate but potentially misleading when the true relationships

are non-linear. The VAE decoder relaxes this constraint, capturing non-linear mappings that may

require visualization rather than a single coefficient (see Supplementary Material: Figures S4–

S8), but in return provides a more faithful representation of the underlying data structure.

Though many ecologists probably suspected that bioclim had fewer intrinsic dimensions than 19,

including myself, it is perhaps surprising that the full panoply of bioclimatic variation on Earth

can be reconstructed with as few as 5 variables. It is also remarkable that these 5 variables have

close correspondence with known sources of climate variation such as latitude, elevation, and the

monsoon. This speaks to the potential power of generative A.I. methods in ecology. In this case

the physical and biological processes of the earth encode themselves in global climate variation,

which this VAE was then able to decode, thanks to its ability to learn flexible non-linear

functions of a simplified underlying space.

The existence of this low-dimensional manifold has a concrete explanation. The 19 bioclim

variables are derived from just three independent monthly variables – minimum temperature,

maximum temperature, and precipitation across 12 months (36 values in total) – through

formulae that compute means, extremes, ranges, and seasonal aggregates (Booth et al. 2014).

These derivation formulae impose deterministic algebraic relationships among the 19 variables,

confining them to a non-linear surface within the 19-dimensional ambient space. The 36

underlying monthly values are themselves highly correlated through seasonal cycles, latitude

effects, and temperature–precipitation coupling, so their effective dimensionality is considerably

less than 36. The bioclim formulae further compress this variation, consistent with the 5-

dimensional manifold the VAE recovers. Importantly, the manifold dimensions do not simply

invert the bioclim derivation formulae to recover monthly values; rather, they capture non-linear

combinations reflecting physical climate processes like elevation gradients and monsoon

dynamics. One might ask why not use the monthly variables directly – the answer is that the

bioclim summaries encode biologically meaningful aggregates (e.g. temperature of the warmest

quarter, precipitation seasonality) that are more directly relevant to species' physiological

tolerances, and the VAE's bidirectional encoder–decoder preserves this biological relevance

while removing redundancy.

A key methodological contribution of this work is therefore a principled, automatic approach to

dimensionality selection. Unlike PCA, where the number of components to retain requires

subjective judgment (e.g. variance thresholds, scree plot "elbows"), the VAE's regularization

structure produces a clear binary separation between active and inactive dimensions (Figure 2),

eliminating arbitrary choices entirely. This yielded 5 uncorrelated variables that perform just as

well as the original 19 in at least one downstream application (SDMs), and in some ways better.

If even just for this automatic dimensionality selection, VAE methods are currently under-used

in ecology.

A potential limitation of this study is that standard VAEs are non-identifiable: different training

runs may learn different rotations of the same latent subspace (Locatello et al. 2019). To address

this concern, I conducted an identifiability analysis by training 16 identifiable VAE (iVAE;

Khemakhem et al. 2020) models using latitude and longitude, encoded as sinusoidal angle

embeddings (see Supplementary Material), as auxiliary conditioning information, of which 9

converged to the same 5 active latent dimensions as the original VAE. Using the Mean

Correlation Coefficient (MCC) with Procrustes alignment to compare learned representations

across runs, I found that the manifold subspace is highly reproducible (mean Procrustes MCC =

0.84 across all pairwise comparisons). The original (unconditional) VAE showed strong

agreement with the iVAE runs (mean Procrustes MCC = 0.81) and clustered among them in a

hierarchical clustering analysis rather than as an outgroup, confirming that the 5-dimensional

manifold structure is a genuine feature of the bioclimatic data rather than an artifact of a

particular training run (see Supplementary Material: Identifiability of VAE Manifold Dimensions

for full details; Figure S14). Not all dimensions were equally reproducible: the best-matched

dimension pair across runs had a mean absolute correlation of 0.96, while the weakest had 0.65,

suggesting that some manifold dimensions capture more robust climatic gradients than others

(Figure S14D).

The VAE-based variable selection approach demonstrated here has potential applications beyond

bioclimatic data. The ability to automatically determine intrinsic dimensionality from data,

without requiring ad hoc thresholds, could prove valuable wherever ecologists work with large

suites of correlated environmental variables—for example, remote sensing spectral bands, soil

chemistry profiles, or oceanographic variables. Recent work has demonstrated that VAEs can

compress large climate reanalysis datasets by factors of several hundred while retaining essential

structure (Han et al. 2025), confirming that climate data generally occupies a low-dimensional

manifold. More broadly, deep learning approaches are increasingly being applied to species

distribution modeling (Kellenberger et al. 2026), and the manifold variables developed here can

serve as inputs to such models. The NicheFlow framework (Dinnage 2024) extends the present

work by using normalizing flows to build generative models of species' environmental niches

directly in manifold space, enabling both niche characterization and novel environmental

prediction.

The present analysis was conducted at 2.5 arc-minute resolution. Preliminary experiments at 5

arc-minute resolution also recovered 5 active manifold dimensions, suggesting that the intrinsic

dimensionality of bioclimatic variation is relatively robust to spatial resolution within this range.

A formal multi-resolution sensitivity analysis using the iVAE framework with warm-start

training across resolutions is an important avenue for future work. Previous studies have

generally found that changing spatial resolution affects the magnitude but not the qualitative

nature of species–environment relationships (Merkenschlager et al. 2023), consistent with our

observation that the manifold structure is preserved across the resolutions examined.

Figure Captions

Figure 1. Cumulative variance explained by 19 PCA dimensions of bioclim variables. There is no obvious jump in the amount of new variance explained from one PCA dimension to the next, which would indicate a possible linear manifold. Otherwise, convention suggests arbitrarily keeping the PCA dimensions that cumulatively sum up to >95% of variance explained.

Figure 2. The mean variances along each variational autoencoder latent dimension (there were 16 altogether). "Mean Variance" refers to the approximate posterior variance for each latent dimension, averaged across all ~12.5 million data points. All dimensions except for 5 had variance near or equal to 1, matching the prior distribution. The remaining 5 had variances close to zero. This is indicative that these are 'manifold' dimensions: they are necessary to reconstruct the full data. All other dimensions are just noise that the decoder network learns to ignore.

Figure 3. Global maps of the 5 VAE manifold variables. Most variables appear to be imperfect but reasonable approximations to known global climate factors. BIOCMAN1 looks like elevation. BIOCMAN2 is where rainforest exists (including temperate rainforest). BIOCMAN3 has the least clear pattern, but low values seem to be associated with arid and semi-arid places. BIOCMAN4 is close to a map of latitude, and BIOCMAN5 is high where there are monsoonal climates. Maps are presented in Danseiji V projection, which has low distortion in terrestrial areas at the cost of high distortion in the oceans (Kunimune 2020). High resolution versions with full scale bars can be found in Appendix S1: Figures S9–S13.

376 **Figure 1**

379   **Figure 2**

Figure 3

References

References

Battey, C. J., G. C. Coffing, and A. D. Kern. 2020. Visualizing population structure with

variational autoencoders. G3: Genes|Genomes|Genetics 11.

Bengio, Y., A. C. Courville, and P. Vincent. 2012. Representation Learning: A Review and New

Perspectives. IEEE Transactions on Pattern Analysis and Machine Intelligence 35:1798–1828.

Booth, T. H. 2022. Checking bioclimatic variables that combine temperature and precipitation

data before their use in species distribution models. Austral Ecology.

Booth, T. H., H. Nix, J. R. Busby, and M. F. Hutchinson. 2014. bioclim: the first species

distribution modelling package, its early applications and relevance to most current MaxEnt

studies. Diversity and Distributions 20.

Cattell, R. B. 1966. The Scree Test For The Number Of Factors.. Multivariate behavioral

research 1 2:245–76.

Dai, B., and D. P. Wipf. 2019. Diagnosing and Enhancing VAE Models. ArXiv abs/1903.05789.

Dinerstein, E., D. P. Olson, A. R. Joshi, C. Vynne, N. D. Burgess, E. D. Wikramanayake, N. R.

Hahn, S. Palminteri, P. Hedao, R. F. Noss, M. Hansen, H. Locke, E. C. Ellis, B. S. Jones, C. V.

Barber, R. Hayes, C. F. Kormos, V. G. Martin, E. Crist, W. Sechrest, L. Price, J. E. M. Baillie,

D. E. Weeden, K. F. Suckling, C. Davis, N. C. Sizer, R. Moore, D. Thau, T. Birch, P. V.

Potapov, S. Turubanova, A. Tyukavina, N. de Souza, L. Pintea, J. C. Brito, O. A.-ar-R.

Llewellyn, A. G. Miller, A. Patzelt, S. A. Ghazanfar, J. R. Timberlake, H. Klöser, Y.

Shennan□Farpón, R. Kindt, J.-P. B. Lillesø, P. van Breugel, L. Graudal, M. Voge, K. F. Al-

Shammari, and M. Saleem. 2017. An Ecoregion-Based Approach to Protecting Half the

Terrestrial Realm. Bioscience 67:534–545.

Doersch, C. 2016. Tutorial on Variational Autoencoders. ArXiv abs/1606.05908.

Dinnage, R. 2024. NicheFlow: Towards a foundation model for Species Distribution Modelling.

bioRxiv. https://doi.org/10.1101/2024.10.15.618541.

Dormann, C. F., J. Elith, S. Bacher, C. M. Buchmann, G. Carl, G. Carré, J. R. G. Márquez, B.

Gruber, B. Lafourcade, P. J. Leitão, T. Münkemüller, C. J. McClean, P. E. Osborne, B.

Reineking, B. Schröder, A. K. Skidmore, D. Zurell, and S. Lautenbach. 2013. Collinearity: a

review of methods to deal with it and a simulation study evaluating their performance.

Ecography 36:27–46.

Elith, J., C. H. Graham, R. Valavi, M. Abegg, C. Bruce, A. J. Ford, A. Guisan, R. J. Hijmans, F.

Huettmann, L. G. Lohmann, B. A. Loiselle, C. C. Moritz, J. M. C. Overton, A. T. Peterson, S. J.

Phillips, K. S. Richardson, S. E. Williams, S. K. Wiser, T. Wohlgemuth, and N. E. Zimmermann.

2020. Presence-only and Presence-absence Data for Comparing Species Distribution Modeling

Methods. Biodiversity Informatics 15:69–80.

Elith, J., and J. R. Leathwick. 2009. Species Distribution Models: Ecological Explanation and

Prediction Across Space and Time. Annual Review of Ecology, Evolution, and Systematics

40:677–697.

Falbel, D., and J. Luraschi. 2022. torch: Tensors and Neural Networks with 'GPU' Acceleration.

Fick, S. E., and R. J. Hijmans. 2017. WorldClim 2: new 1 km spatial resolution climate surfaces

for global land areas. International Journal of Climatology 37.

Fourcade, Y., A. G. Besnard, and J. Secondi. 2018. Paintings predict the distribution of species,

or the challenge of selecting environmental predictors and the modelling procedure. A reply to

Titley et al. Global Ecology and Biogeography 27:245–256.

Gomari, D. P., A. Schweickart, L. Cerchietti, E. Paietta, H. Fernandez, H. Al-Amin, K. Suhre,

and J. Krumsiek. 2022. Variational autoencoders learn transferrable representations of

metabolomics data. Communications Biology 5.

Han, T., Z. Chen, S. Guo, W. Xu, W. Ouyang, and L. Bai. 2025. Climate science data can be

compressed efficiently by dual-stage extreme compression with a variational auto-encoder

transformer. Communications Earth & Environment 6:955.

Hawkins-Hooker, A., F. Depardieu, S. Baur, G. Couairon, A. Chen, and D. Bikard. 2020.

Generating functional protein variants with variational autoencoders. PLoS Computational

Biology 17.

Hinton, G. E., and R. Salakhutdinov. 2006. Reducing the Dimensionality of Data with Neural

Networks. Science 313:504–507.

James, G., D. Witten, T. Hastie, and R. Tibshirani. 2013. An Introduction to Statistical Learning.

Springer, New York.

Kellenberger, B., K. Winner, and W. Jetz. 2026. The Performance and Potential of Deep

Learning for Predicting Species Distributions. Global Ecology and Biogeography 35:e70184.

Khemakhem, I., D. Kingma, R. Monti, and A. Hyvärinen. 2020. Variational autoencoders and

nonlinear ICA: A unifying framework. Proceedings of the 23rd International Conference on

Artificial Intelligence and Statistics (AISTATS), PMLR 108:2207–2217.

Kingma, D. P., and M. Welling. 2013. Auto-Encoding Variational Bayes. CoRR abs/1312.6114.

Kriticos, D. J., Jaroı́k Vojtěch, and N. Ota. 2014. Extending the suite of bioclim variables: a

proposed registry system and case study using principal components analysis. Methods in

Ecology and Evolution 5.

Kunimune, J. H. 2020. Minimum-error world map projections defined by polydimensional

meshes. International Journal of Cartography 7:78–99.

Locatello, F., S. Bauer, M. Lucic, G. Rätsch, S. Gelly, B. Schölkopf, and O. Bachem. 2019.

Challenging common assumptions in the unsupervised learning of disentangled representations.

Proceedings of the 36th International Conference on Machine Learning (ICML), PMLR

97:4114–4124.

Merkenschlager, C., F. Bangelesa, H. Paeth, and E. Hertig. 2023. Blessing and curse of

bioclimatic variables: A comparison of different calculation schemes and datasets for species

distribution modeling within the extended Mediterranean area. Ecology and Evolution

13:e10553.

Pearson, K. 1901. LIII. On lines and planes of closest fit to systems of points in space.

Philosophical Magazine Series 1 2:559–572.

Pearson, R. G., W. Thuiller, M. B. Araújo, E. Martínez☐Meyer, L. Brotóns, C. J. McClean, L.

Miles, P. Segurado, T. P. Dawson, and D. C. Lees. 2006. Model☐based uncertainty in species

range prediction. Journal of Biogeography 33.

Warren, D. L., L. J. Beaumont, R. Dinnage, and J. B. Baumgartner. 2018. New methods for

measuring ENM breadth and overlap in environmental space. Ecography.

Warren, D. L., N. J. Matzke, M. Cardillo, J. B. Baumgartner, L. J. Beaumont, M. Turelli, R. E.

Glor, N. A. Huron, M. V. P. Simões, T. L. Iglesias, J. C. Piquet, and R. Dinnage. 2021.

ENMTools 1.0: an R package for comparative ecological biogeography. Ecography.

Zheng, Y., T. He, Y. Qiu, and D. P. Wipf. 2023. Learning Manifold Dimensions with

Conditional Variational Autoencoders. ArXiv abs/2302.11756.

Ziegler, C., J. Martin, C. Sinner, and F. Morcos. 2023. Latent generative landscapes as maps of

functional diversity in protein sequence space. Nature Communications 14.