1 **Title**
2 Ubiquitin ligase and signalling hub MYCBP2 is required for efficient EPHB2 tyrosine kinase
3 receptor function
4

5 **Authors**
6 Chao Chang[1,2], Sara L. Banerjee[1,3], Sung Soon Park[1,2], Xiaolei Zhang[1], David Cotnoir-White[1,%],
7 Muriel Desbois[4,5], Brock Grill[4,6,7], and Artur Kania[1,2,3,8,#,*]
8
Affiliations
[1]Institut de recherches cliniques de Montréal (IRCM), Montréal, QC, H2W 1R7, Canada
[2]Integrated Program in Neuroscience, McGill University, Montréal, QC, H3A 2B4, Canada
[3]Division of Experimental Medicine, McGill University, Montréal, QC, H3A 2B2, Canada
[4]Center for Integrative Brain Research, Seattle Children's Research Institute, Seattle, WA 98101,
USA.
[5]School of Life Sciences, Keele University, Keele, Staffordshire ST5 5BG, UK
[6]Department of Pediatrics, University of Washington School of Medicine, Seattle, WA 98101,
USA
[7]Department of Pharmacology, University of Washington School of Medicine, Seattle, WA 98195,
USA.
[8]Department of Anatomy and Cell Biology, McGill University, Montréal, QC, H3A 0C7, Canada
21
22 [%]Current address: Modulari-T Biosciences, Montreal, QC, Canada H4B 1R6
23
24 [#]Lead contact
25 *Correspondence: artur.kania@ircm.qc.ca

26
27 **Running Title**
28 MYCBP2 functions in EPHB2 signalling.
29
30
31

Abstract

Eph receptor tyrosine kinases participate in a variety of normal and pathogenic processes during development and throughout adulthood. This versatility is likely facilitated by the ability of Eph receptors to signal through diverse cellular signalling pathways: primarily by controlling cytoskeletal dynamics, but also by regulating cellular growth, proliferation, and survival. Despite many proteins linked to these signalling pathways interacting with Eph receptors, the specific mechanisms behind such links and their coordination remain to be elucidated. In a proteomics screen for novel EPHB2 multi-effector proteins, we identified human MYC binding protein 2 (MYCBP2 or PAM or Phr1). MYCBP2 is a large signalling hub involved in diverse processes such as neuronal connectivity, synaptic growth, cell division, neuronal survival, and protein ubiquitination. Our biochemical experiments demonstrate that the formation of a complex containing EPHB2 and MYCBP2 is facilitated by FBXO45, a protein known to select substrates for MYCBP2 ubiquitin ligase activity. Formation of the MYCBP2-EPHB2 complex does not require EPHB2 tyrosine kinase activity and is destabilised by binding of ephrin-B ligands, suggesting that the MYCBP2-EPHB2 association is a prelude to EPHB2 signalling. Paradoxically, the loss of MYCBP2 results in increased ubiquitination of EPHB2 and a decrease of its protein levels suggesting that MYCBP2 stabilises EPHB2. Commensurate with this effect, our cellular experiments reveal that MYCBP2 is essential for efficient EPHB2 signalling responses in cell lines and primary neurons. Finally, our genetic studies in *C. elegans* provide *in vivo* evidence that the ephrin receptor VAB-1 displays genetic interactions with known MYCBP2 binding proteins. Together, our results align with the similarity of neurodevelopmental phenotypes caused by MYCBP2 and EPHB2 loss of function, and couple EPHB2 to a signaling effector that controls diverse cellular functions.

Introduction

Eph receptor tyrosine kinases and their membrane-tethered ligands, the ephrins, elicit short distance cell-cell signals that regulate many biological processes. Ephrin-Eph signalling primarily impacts the cytoskeleton with the immobilization of highly dynamic axonal growth cones being a classic example. Other processes that involve changes in transcription, growth, and survival such as angiogenesis, synaptic plasticity, stem cell fate, tumorigenesis and neurodegeneration also involve the Eph/ephrin system. Many proteins are postulated to couple Eph receptors to different intracellular effectors, but the molecular logic of this diversity remains fragmented (Kania and Klein, 2016;Bush, 2022).

EphB subfamily members preferentially bind transmembrane ephrin-Bs and although both molecules participate in bidirectional signalling, ephrin-B activation of EphB signalling cascades is more thoroughly studied (Gale et al., 1996;Mellitzer et al., 1999). To elicit robust Eph receptor forward signalling, ephrins multimerise in signalling clusters by intercalating with Ephs on a signal-recipient cell with array size correlating with signal amplitude (Kullander et al., 2001;Schaupp et al., 2014). Signalling initiation involves the activation of the receptor tyrosine kinase and phosphorylation of tyrosines proximal to the EphB transmembrane domain (Soskis et al., 2012;Binns et al., 2000). Eph-evoked cytoskeletal effects such as cell contraction and growth cone collapse result from changes in small GTPase activity modulated by EphB Guanine nucleotide exchange factors (GEFs) and GTPase activating proteins (GAPs) (Margolis et al., 2010;Shi et al., 2007). Eph signalling has also been linked to fundamental cellular pathways such as the Ras-MAPK pathway, mTOR-regulated protein synthesis, cell division and survival (Bush and Soriano, 2010;Nie et al., 2010;Fawal et al., 2018;Genander et al., 2009;Depaepe et al., 2005). Eph forward signalling eventually leads to their internalisation and either recycling or degradation via endosome/lysosome and ubiquitination/proteasome pathways (Zimmer et al., 2003;Okumura et al., 2017). While identification of a growing number of proteins interacting with Eph receptors has moved the field forward, we have yet to clarify the question of how Eph receptors activate various fundamental cellular processes, often within the same cell type.

Myc-binding protein 2 (MYCBP2), also known as Protein Associated with Myc (PAM) and Highwire, RPM-1, or Phr1 in different species, is a large signalling hub that regulates cytoskeletal dynamics, neuronal development, and axonal degeneration (Guo et al., 1998;Grill et al., 2016;Virdee, 2022). It has an atypical RING ubiquitin ligase activity that inhibits the p38/MAP and JNK kinase pathways thereby regulating cytoskeletal dynamics underlying axonal development and synaptic growth (Nakata et al., 2005;Collins et al., 2006;Pao et al., 2018;Wan et al., 2000;Lewcock et al., 2007;Borgen et al., 2017). MYCBP2 further regulates the Tuberin Sclerosis Complex linked to cell growth (Han et al., 2012), initiation of autophagy via ULK (Crawley et al., 2019) and NMNAT2-regulated neuronal survival and axonal degeneration (Babetto et al., 2013;Xiong et al., 2012). Biochemical mapping has shown that human MYCBP2

and its *C. elegans* ortholog RPM-1 rely upon the FBD1 domain to bind the F-box protein FBXO45 that acts as a ubiquitination substrate selector (Desbois et al., 2018;Sharma et al., 2014). Intriguingly, the neurodevelopmental phenotypes caused by MYCBP2 and EPHB2 loss of function are similar, raising the possibility that these two molecules could function in the same pathway (Henkemeyer et al., 1996;Lewcock et al., 2007;Dalva et al., 2000). Nonetheless, a biochemical or genetic interaction between MYCBP2 and EPHB2 has not been demonstrated in any system.

To shed light on how EphB receptors fulfil their multitude of functions, we used mass spectrometry (MS)-based proteomics to identify multi-effector proteins that bind EPHB2. One of our proteomic hits was MYCBP2, which we demonstrated forms a complex with EPHB2 using a combination of biochemical and cellular assays. Furthermore, we show that this interaction is required for efficient EPHB2 signalling in cell lines and primary neurons. Consistent with these findings, we observed *in vivo* genetic interactions in *C. elegans* between the Eph receptor, VAB-1, and known RPM-1 binding proteins. Our collective results indicate that the relationship between EPHB2 and MYCBP2 does not appear to involve the ubiquitin ligase activity of MYCBP2, and raise the possibility that MYCBP2 links EPHB2 to diverse fundamental cellular functions.

Results

Proteomics identifies MYCBP2 as a putative EPHB2-interacting protein

To understand the molecular logic underlying EPHB2 signalling diversity, we performed affinity purification coupled to MS (AP-MS) in order to identify EPHB2-interacting proteins, and prioritised those known to be signalling hubs. We used a stable HeLa cell line with tetracycline-inducible expression of BirA-linked EPHB2-FLAG, that we previously used to study EPHB2 signalling (Lahaie et al., 2019). To identify ephrin ligand-dependent EPHB2 protein complexes, we stimulated EPHB2-FLAG-overexpressing cells with pre-clustered Fc control or ephrinB2-Fc (eB2-Fc). We then harvested and lysed the cells, performed anti-FLAG immunoprecipitation, and used mass spectrometry to identify EPHB2 protein complexes (Fig. 1A).

To identify EPHB2-specific interactions and remove background contaminants, we performed Significance Analysis of INTeractome (SAINTexpress analysis (Teo et al., 2014)) using our EPHB2-related controls (Lahaie et al., 2019). To better visualize the changes in putative EPHB2 binding partners, we compared the average spectral counts of the identified proteins using ProHits-viz tool (Knight et al., 2017). Comparison of Fc and ephrin-B2-treated samples did not yield any significant differences. Since we applied the ligand and collected the samples on a time scale comparable to known EphB signalling dynamics, this limitation could be potentially due to autoactivation of Eph receptors when they are overexpressed, obscuring some ligand-dependent effects (Lackmann et al., 1998). However, the resulting scatter plot confirmed the presence of several known EPHB2 interactors, such as FYN and YES1 Src kinases and members of the Eph receptor family (Fig. 1B) (Banerjee et al., 2022). One of the most prominent, novel hits was the E3 ubiquitin ligase and signalling hub protein MYCBP2, and its binding partner FBXO45. FBXO45 was previously identified as a putative EPHB2 interacting protein in large-scale, cell-based interactome studies (Huttlin et al., 2021;Salokas et al., 2022).

Biochemical validation of MYCBP2 binding to EPHB2

To confirm that MYCBP2 can indeed form a molecular complex with EPHB2, we tested whether endogenous MYCBP2 co-immunoprecipitates (co-IP) with FLAG-tagged EPHB2 in HEK 293T cells. Based on differences in EPHB2 and EPHA3 interactomes, we reasoned that EPHA3 may serve as a negative control for the EPHB2-MYCBP2 association (Huttlin et al., 2021). We found that MYCBP2 coprecipitated with affinity-purified EPHB2-FLAG, but not EPHA3 (Fig. 1C). This confirmed MYCBP2 binding to EPHB2 and suggested MYCPB2 displays EPH receptor subtype specificity. Importantly, the EPHB2-MYCBP2 interaction was reduced by 24.5% and 53% following ephrin-B1 and ephrin-B2 treatment respectively in HeLa EPHB2 cells, suggesting the involvement of MYCBP2 in ephrin-B:EPHB2 signalling (Fig. 1D, E, eB1-Fc, p=0.1365; eB2-Fc,

158   p=0.0002). To test this interaction *in vivo*, we performed co-IP using dissociated rat cortical
159   neurons, which further confirmed MCYBP2 association with EPHB2 (Fig. 1F).
160
We next asked whether the kinase activity of EPHB2 is required for the formation of the
EPHB2-MYCBP2 complex. To test this, we expressed GFP-tagged wild type (WT) EPHB2 or
EPHB2 with a kinase-dead mutation (KD) in HeLa cells. We found that MYCBP2 showed
comparable coprecipitation with WT and KD EPHB2 (Fig. 1G). Thus, EPHB2 kinase activity is
not required for the formation of MYCBP2 - EPHB2 complexes, suggesting that MYCBP2
association with EPHB2 may be a prelude to ephrin-B-evoked EPHB2 signalling.
167
168
169   **FBXO45 enhances the association between MYCBP2 and EPHB2**
170
Our proteomics screen for EPHB2 interactors also identified FBXO45, the F-box protein that
forms a ubiquitin ligase complex with MYCBP2 (Sharma et al., 2014;Saiga et al., 2009). Thus,
we initially reasoned that FBXO45 might perform a similar role in the formation of the MYCBP2-
EPHB2 complex. We first tested whether FBXO45 binds to EPHB2 by co-expressing MYC-
tagged FBXO45 with EPHB2-FLAG or EPHA3-FLAG in HEK293T cells. Co-IP revealed that
FBXO45 can associate with EPHB2 but not EPHA3, suggesting that EPHB2, FBXO45 and
MYCBP2 could form a ternary complex (Fig. 2A). To test this idea, we co-expressed EPHB2-
FLAG and GFP-MYCBP2 in the presence or absence of MYC-FBXO45 and examined co-IP
efficiency between EPHB2 and MYCBP2. Interestingly, FBXO45 enhanced the interaction
between EPHB2 and MYCBP2 (Fig. 2B, C, p=0.0068). Together, these data suggest that EPHB2
can form a complex with both MYCBP2 and FBXO45, and that FBXO45 increases the efficiency
of MYCBP2-EPHB2 interaction.
183
184
185   **Biochemical mapping of EPHB2-MYCBP2 interaction**
186
To identify the MYCBP2 protein domain(s) required for the formation of the ternary complex with
EPHB2 and FBXO45, we co-expressed EPHB2-FLAG and three GFP-MYCBP2 fragments in
HEK293T cells. Co-IP revealed that the central region of MYCBP2 was sufficient for binding with
EPHB2 (Fig. 2D, E). In addition, co-expression of MYC-FBXO45 demonstrated that the
association of the central domain of MYCBP2 with EPHB2 is enhanced by FBOX45 and is
consistent with the presence of an FBXO45 binding site within this MYCBP2 fragment (Fig. 2E)
(Sharma et al., 2014).
194
195   To identify the domains of EPHB2 required for the formation of the tripartite complex, we
196   took advantage of the observation that EPHA3 does not readily form a complex with MYCBP2
197   or FBXO45. Thus, we performed domain swapping experiments between EPHB2 and EPHA3

reasoning that placing EPHA3-specific sequences in EPHB2 would inhibit the formation of the tripartite complex. We constructed a series of FLAG-EPHB2/EPHA3 chimeras and determined whether they bound MYCBP2 in the presence of FBXO45 (Fig. 2F). Co-IP revealed that the ability of a particular chimera to associate with MYCBP2 was correlated with its association with FBXO45, in line with the MYCBP2-FBXO45 complex interacting with EPHB2. Surprisingly, we found that EPHB2-EPHA3 chimeras with an EPHA3 identity of intracellular juxtamembrane, kinase, SAM or PDZ binding domains retained the ability to associate with FBXO45 and MYCBP2, suggesting that the formation of the tripartite complex is driven by the extracellular domain and/or the transmembrane domain of EPHB2 (Fig. 2G). However, these results are also consistent with the possibility that the EPHB2 identity of the extracellular fragments could alter the conformation of the intracellular domains of EPHA3 identity, allowing the interaction with FBXO45-MYCBP2 to occur. To exclude this possibility, we created EPHB2 mutants lacking the intracellular or extracellular domains and tested their ability to complex with MYCBP2 and FBXO45 by co-IP (Fig. 2H). In these experiments only the deletion mutant lacking the intracellular domain retained its ability to form the tripartite complex. Collectively, these results argue that the combination of extracellular and transmembrane domains of EPHB2 are necessary and sufficient for formation of the MYCBP2–FBXO45–EPHB2 complex (Fig. 2I).

MYCBP2 is required for EPHB2-mediated cellular responses

Given the decrease in MYCBP2–EPHB2 association evoked by ephrin-B treatment (Fig. 1D, E), we next sought to determine whether MYCBP2 fulfils a specific function in ephrin-B:EPHB2 forward signalling. Thus, we infected EPHB2-FLAG HeLa cells using lentivirus containing CRISPR sgRNA targeting the *MYCBP2* exon 6, and pooled *MYCBP2*[CRISPR] cells after puromycin selection (Desbois et al., 2018). Using EPHB2-FLAG HeLa cells carrying a stably integrated empty expression vector (CTRL[CRISPR]) as controls, we found that *MYCBP2*[CRISPR] led to a reduction in endogenous MYCBP2 protein levels (Fig. 3A).

To study the role of MYCBP2 in EphB signalling, we took advantage of an experimental paradigm in which exposure to ephrin-B2 evokes cytoskeletal contraction of HeLa cells expressing EPHB2 (Lahaie et al., 2019). Thus, following induction of EPHB2 expression, *MYCBP2*[CRISPR] and CTRL[CRISPR] cells were stimulated with pre-clustered Fc control or ephrin-B2 for 15 minutes and scored as collapsed or uncollapsed. While the proportion of collapsed cells for the two cell lines treated with Fc was similar (CTRL[CRISPR], 5.4%; *MYCBP2*[CRISPR], 5.7%; p=0.9841), ephrin-B2 treatment resulted in the collapse of 19.0% of CTRL[CRISPR] cells but only 12.1% of *MYCBP2*[CRISPR] cells (n=9 coverslips; Fig. 3B, C; p=0.0017). Moreover, when compared to the Fc conditions, *MYCBP2*[CRISPR] cells exhibited a less drastic change in collapse rate upon ephrin-B2 treatment (Fig. 3B, C; eB2 vs Fc: CTRL[CRISPR], p<0.0001; *MYCBP2*[CRISPR], p=0.0115). In addition, time-lapse imaging of *MYCBP2*[CRISPR] and CTRL[CRISPR] cells transiently transfected

238 with an EPHB2-GFP expression plasmid revealed a similar attenuation of ephrin-B2-induced
239 cellular contraction (Fig. 3D, E; p=0.0268). These data argue that MYCBP2 regulates a short-
240 term cellular response evoked by ephrin-B2:EPHB2 signalling.
241
To study longer-term cellular responses evoked by ephrin-B2:EPHB2 signalling, we
turned to a stripe assay in which the preference of CTRL[CRISPR] and *MYCBP2*[CRISPR] cells for
immobilized ephrin-B2 or Fc was measured. To do this, cells of either line were deposited over
alternating stripes of ephrin-B2 or Fc, and stripe preference was scored for individual cells after
overnight incubation. While only 33.2% of CTRL[CRISPR] cells resided on ephrin-B2 stripes, this
proportion was significantly increased to 47.4% for *MYCBP2*[CRISPR] cells, suggesting the loss of
MYCBP2 function led to a decreased repulsion from ephrin-B2 stripes (Fig. 3F, G; n=5 and 7
carpets respectively, p=0.0109). When cells were plated on Fc:Fc stripes, CTRL[CRISPR] and
MYCBP2[CRISPR] cells exhibited no preference over cy3-conjugated Fc stripes (49.18% vs 51.88%,
p=0.5386, images not shown). Native HeLa cells only respond to ephrin-B2 once they are made
to express EphB2. Thus, our data suggest that MYCBP2 is required for EPHB2-mediated cellular
responses in HeLa cells.
254
255
256 **Loss of MYCBP2 decreases cellular levels of EPHB2 protein**
257
The association of EPHB2 with MYCBP2 and its substrate recognition protein FBXO45 suggests
that the MYCBP2 ubiquitin ligase complex could target EPHB2 for degradation, a mechanism
frequently deployed to terminate transmembrane receptor signalling (Foot et al., 2017). However,
the results of our cellular assays contradicted this model, and rather suggested that loss of
MYCBP2 function decreased EPHB2 signalling. To further evaluate these two scenarios, we first
compared EPHB2 protein levels in HeLa *MYCBP2*[CRISPR] cells and CTRL[CRISPR] cells by using
tetracycline to induce the EPHB2 overexpression. We found that MYCBP2 loss reduced EPHB2
protein levels (Fig. 4A). To further confirm this, instead of inducing EPHB2 overexpression with
tetracycline, we transfected EPHB2-FLAG plasmid into both CTRL[CRISPR] and *MYCBP2*[CRISPR]
cells and examined EPHB2-FLAG levels after two days. As shown in Fig. 4B, C, levels of
EPHB2-FLAG were significantly lowered by 26.1% in HeLa *MYCBP2*[CRISPR] cells compared to
CTRL[CRISPR] cells (p=0.0046). We further investigated whether MYCBP2 affects EPHB2 protein
turnover when cycloheximide is added to prevent new protein synthesis. EPHB2-FLAG
expression was induced by tetracycline for 12 hours followed by protein synthesis inhibition with
cycloheximide. We found that EPHB2 half-life was reduced in HeLa cells lacking MYCBP2
compared to control (Fig. 4D, E; 8h treatment, p=0.0474).
274
275   Ephrin ligand treatment eventually results in Eph receptor degradation, a process
276 associated with signalling termination. We therefore asked whether ligand-mediated EPHB2
277 receptor degradation depends on MYCBP2. We induced EPHB2 expression in CTRL[CRISPR] and

MYCBP2[CRISPR] cells, and exposed them to ephrin-B2 for a different length of time. Western blotting revealed that 4-8 hours stimulation with ephrin-B2 reduced EPHB2 levels in CTRL[CRISPR] cells, but this effect was more drastic in *MYCBP2*[CRISPR] cells (Fig. 4F, G). Taken together, our results are not consistent with MYCBP2 ubiquitinating EPHB2 and causing its degradation. Unexpectedly, our data indicate that MYCBP2 stabilizes EPHB2 in HeLa cells under both naïve and ligand-challenged conditions.

Loss of MYCBP2 enhances ligand-induced EPHB2 receptor ubiquitination

Ubiquitination of receptor tyrosine kinases, including Eph receptors, can herald their degradation via the proteasome and thus termination of signalling (Haglund and Dikic, 2012;Sabet et al., 2015). This model is not supported by our results, which suggest that MYCBP2 is required for EPHB2 protein maintenance. Nonetheless, we investigated whether EPHB2 receptor ubiquitination is altered in HeLa cells depleted of MYCBP2. HeLa CTRL[CRISPR] and *MYCBP2*[CRISPR] cells were transfected with HA-tagged ubiquitin, EPHB2 expression was induced with tetracycline, and cells were treated with ephrin-B2 for 30min. We observed that EPHB2 receptor ubiquitination was not significantly increased in CTRL[CRISPR] cells after this short-term ligand treatment (Fig. 4H, I CTRL[CRISPR], p=0.1349). In contrast, EPHB2 ubiquitination was significantly increased in *MYCBP2*[CRISPR] cells (Fig. 4H, I *MYCBP2*[CRISPR], p=0.0195). This effect argues against the concept that EPHB2 is a MYCBP2 ubiquitination substrate, and suggests that in the absence of MYCBP2 degradation of the EPHB2 receptor is enhanced due to increased ubiquitination.

Loss of MYCBP2 promotes EPHB2 degradation through the lysosomal pathway

EPHB2 can be degraded by either a proteasomal or lysosomal pathway depending on the cellular context (Cissé et al., 2011;Litterst et al., 2007;Fasen et al., 2008). Thus, we wanted to shed light on how EPHB2 is degraded and understand why EPHB2 degradation is enhanced by MYCBP2 loss of function. To do so, we induced EPHB2 expression in CTRL[CRISPR] and *MYCBP2*[CRISPR] cells and applied the S26 proteasome inhibitor MG132, or the lysosomal inhibitors BafilomycinA1 or Chloroquine. We found that MG132 did not have significant effects on EPHB2 levels in both cell types (Fig. 4J, K). However, we found that BafilomycinaA1 (BafA1) significantly increased EPHB2 protein levels in both HeLa CTRL[CRISPR] cells and *MYCBP2*[CRISPR] cells by 19% and 40%, respectively (Fig. 4J, K). We also observed a trend towards increased EPHB2 levels with Chloroquine (coQ) treatment in CTRL[CRISPR] (14%) and MYCBP2[CRISPR] (35%), further suggesting a role for lysosomal degradation (Fig. 4J, K). These observations suggest that the loss of MYCBP2 promotes EPHB2 receptor degradation through the lysosomal pathway, which expands our understanding of how lysosomal degradation of EPHB2 is regulated.

Loss of MYCBP2 does not decrease EPHB1, EPHB3, or EPHA3 receptor levels

The above experiments raise the question of whether MYCBP2 is a general regulator of Eph receptor stability. Since EPHA3 does not form a complex with MYCBP2, and EphA receptor levels are controlled by the proteasomal pathway (Sharfe et al., 2003;Walker-Daniels et al., 2002), we hypothesized that MYCBP2 might regulate the levels of the entire EphB receptor class. To test this, we co-transfected plasmids encoding GFP-tagged EPHB1 and HA-tagged EPHB3 into HeLa CTRL^{CRISPR} and *MYCBP2*^{CRISPR} cells. Compared to CTRL^{CRISPR} cells, EPHB1 and EPHB3 levels were not significantly reduced in *MYCBP2*^{CRISPR} cells (Fig. 4L, M, EPHB1, p=0.0588; EPHB3, p=0.4253). Likewise, FLAG-EPHA3 levels were similar in CTRL^{CRISPR} and *MYCBP2*^{CRISPR} cells (Fig. 4N, O, p=0.5369). Taken together, these data suggest that MYCBP2 specifically affects EPHB2 receptor stability.

Loss of MYCBP2 attenuates the magnitude of EPHB2 cellular signalling

EPHB2 receptor activation evokes signal transduction events such as tyrosine phosphorylation of EPHB2, activation of EPHB2 tyrosine kinase function and phosphorylation of the ERK1/2 downstream effector (Poliakov et al., 2008). We therefore measured EPHB2 and ERK1/2 phosphorylation in CTRL^{CRISPR} and *MYCBP2*^{CRISPR} cells for up to 8 hours after ephrin-B2 application (Fig. 5A). P-EPHB2 (pY20) and p-ERK1/2 signals were normalised to GAPDH and ERK1/2, respectively. In HeLa CTRL^{CRISPR} cells, pTyr-EPHB2 response reached a plateau after 1-2 h treatment and remained up 8 hours post-stimulation (Fig. 5A-C). On the contrary, ephrin-B2-evoked phosphorylation of EPHB2 was reduced in *MYCBP2*^{CRISPR} cells with quantitative results showing a significant reduction by 8 hours of treatment (Fig. 5A, C; 8h p=0.0331). We were also able to detect significantly lower p-ERK1/2 levels in *MYCBP2*^{CRISPR} cells relative to CTRL^{CRISPR} cells, although activation of ERK1/2 by ephrin-B2 was variable (Fig. 5D. 4h, p=0.0494; 8h, p=0.0078; n=6). We again noted significantly enhanced EPHB2 degradation in *MYCBP2*^{CRISPR} cells (Fig. 5E; 8h, p=0.0437). This decrease was also observed when CTRL^{CRISPR} and *MYCBP2*^{CRISPR} cells were treated with ephrin-B1 (data not shown).

Next, we asked whether the decrease in ligand-evoked EPHB1/2 and ERK1/2 activation in *MYCBP2*^{CRISPR} cells reflects a requirement for MYCBP2 in the EPHB2 signalling cascade per se, or whether it is explained by decreased EPHB2 protein levels caused by MYCBP2 loss. We thus normalized p-EPHB2 and p-ERK1/2 signal to the levels of EPHB2 protein at all time points, which revealed that kinetics and magnitude of EPHB2 and ERK1/2 phosphorylation are similar between CTRL^{CRISPR} and *MYCBP2*^{CRISPR} cells (Fig. 5F, G). Although these data argue against a direct role of MYCBP2 in the early events of EPHB2 signalling, they nevertheless indicate that MYCBP2 loss results in marked attenuation of EPHB2 activation and its downstream pERK1/2 signalling in line with decreased cellular responses to ephrin-B2.

Exogenous Fbxo45 binding domain of MYCBP2 disrupts the EPHB2-MYCBP2 interaction

Previous studies showed that FBXO45 binds to the FBD1 domain of MYCBP2, and exogenous FBD1 overexpression can disrupt the FBXO45-MYCBP2 association (Sharma et al., 2014). We thus tested whether FBD1 overexpression can interfere with the formation of the EPHB2-MYCBP2 complex (Fig. 6A). Indeed, the expression of GFP-FBD1 wild-type (WT) in HEK cells expressing EPHB2-FLAG and MYC-FBXO45 reduced binding between EPHB2 and MYCBP2 (Fig. 6B). This effect was not observed with a GFP-FBD1 mutant (mut) fragment that harbours three point mutations that inhibit binding to FBXO45. Reduction of the EPHB2-MYCBP2 interaction was also observed in cells that were not overexpressing FBXO45 (Fig. 6C). Thus, exogenous FBD1 can specifically disrupt the EPHB2-MYCBP2 association. This suggests that FBXO45 binding to MYCBP2 may facilitate for formation of the MYCBP2-EPHB2 complex.

FBD1 expression impairs EPHB2-mediated neuronal responses

To determine whether the EPHB2-MYCBP2 interaction is required for EPHB2 function, we disrupted binding via FBD1 ectopic expression in cell lines or primary neurons, and studied cellular responses to ephrin-B treatment. We introduced GFP-FBD1 mut or GFP-FBD1 WT into HeLa cells with inducible EPHB2 expression and cultured these cells on ephrin-B2 and Fc stripes. Following overnight culture, only 19.2% of cells expressing GFP-FBD1 mut resided on ephrin-B2 stripes. In contrast, 28.6% of cells expressing GFP-FBD1 WT were found on ephrin-B2 stripes indicating that FBD1 expression can dampen ephrin-B2:EPHB2 mediated cell repulsion (Fig. 6D, E; p=0.0107). In contrast, cells expressing GFP-FBD1 mut or GFP-FBD1 WT displayed no preference for either one of Fc:Fc control stripes (49.63% vs 49.85%, p=0.9560, images not shown).

Since EPHB2 is expressed in the embryonic chicken spinal cord, we performed *in ovo* electroporation to introduce plasmids encoding GFP-FBD1 mut or GFP-FBD1 WT into the spinal cords of 2.5-day old embryonic chickens (Hamburger-Hamilton; (HH) Stage 15-17; (Hamburger and Hamilton, 1951;Luria et al., 2008)). Two days later spinal cords were dissected, divided into explants, cultured overnight on alternating stripes containing ephrin-B2 or Fc, and axonal GFP signal present over ephrin-B2 versus Fc stripes was determined. When explants expressed GFP-FBD1 mut (negative control), we observed 26.5% of axonal GFP signal resided on ephrin-B2 stripes (Fig. 7A, B). In contrast, we found a significant increase of 32.2% of neurites expressing FBD1 WT on ephrin-B2 stripes (Fig. 7A, B; p=0.0410). Thus, FBD1 expression impairs long-term repulsive responses to ephrinB2-EPHB2 signalling in spinal explants.

To study short-term neuronal responses to ephrin-Bs, we turned to mouse hippocampal neurons and a growth cone collapse assay (Srivastava et al., 2013). Here, we electroporated

GFP-FBD1 mut or GFP-FBD1 WT expression plasmids into dissociated hippocampal neurons and treated them with pre-clustered Fc, ephrin-B1 or ephrin-B2 for one hour (Fig. 7C). Neurons expressing GFP-FBD1 mut showed 33.9% growth cone collapse with Fc treatment, while Ephrin-B1 and Ephrin-B2 treatment elicited significant increases to 57.9% and 53.3% collapse, respectively (Fig. 7C, D). Importantly, both ephrin-B ligands significantly increased the frequency of growth cone collapse with respect to Fc treatment when FBD1 mut was expressed (Fig. 7D; p<0.0001, and p=0.0008). In contrast, this effect was either significantly reduced or less evident in neurons expressing GFP-FBD1 WT: with only 48.1% and 57.5% growth cones being collapsed by ephrin-B1 and ephrin-B2, respectively, compared to 41.7% being collapsed by Fc treatment (Fig. 7D; p=0.4357 and p=0.0065, respectively). Together, these data indicate that impairing MYCBP2 function via FBD1 expression disrupts ephrin:B-EPHB2 signalling in axonal guidance.

C. elegans ephrin receptor VAB-1 interacts genetically with known RPM-1/MYCBP2 binding proteins

Next, we sought to test genetic interactions between an ephrin receptor and the MYCBP2 signalling network using an *in vivo* animal model. To do so, we turned to *C. elegans* which has a sole Eph family receptor called VAB-1(George et al., 1998) , and a single MYCBP2 ortholog called RPM-1(Grill et al., 2016). Previous studies have shown that RPM-1/MYCBP2 is required to terminate axon outgrowth with failed termination defects in *rpm-1* mutants arising due to impaired growth cone collapse(Schaefer et al., 2000;Borgen et al., 2017). Prior studies have also demonstrated that RPM-1 functions via numerous downstream pathways, and that RPM-1 binding proteins display genetic enhancer interactions (Grill et al., 2007;Grill et al., 2012;Tulgren et al., 2014;Baker et al., 2014). Given our cell-based biochemical results showing that MYCBP2 binds EphB2, we opted to evaluate two known RPM-1 binding proteins for genetic interactions with the VAB-1 Ephrin receptor. 1) The Rab GEF GLO-4 is an RPM-1 binding protein that is orthologous to mammalian SERGEF. GLO-4 functions via the GLO-1/RAB32 small GTPase and is not involved in RPM-1 ubiquitin ligase activity (Grill et al., 2007;Sharma et al., 2014)(Fig 8A). 2) We also tested the *C. elegans* FBXO45 ortholog, FSN-1, which is the F-box protein of the RPM-1 ubiquitin ligase complex (Liao et al., 2004;Sharma et al., 2014;Baker and Grill, 2017;Desbois et al., 2022) (Fig 8A).

To evaluate genetic interactions between the VAB-1 Eph receptor and RPM-1/MYCBP2 binding proteins, we used the PLM mechanosensory neurons of *C. elegans*. There are two PLM neurons with one on either side of the body. Each PLM neuron extends a primary axon that grows anteriorly and terminates at a precise location before the cell body of another mechanosensory neuron called ALM (Fig. 8B). Consistent with previous studies (Mohamed and Chin-Sang, 2006), *vab-1/EphR* loss-of-function (lf) mutants presented moderate failed termination defects where the PLM axon overextends beyond the ALM cell body (Fig. 8B).

Quantitation indicated that this phenotype occurred at a low, but significant frequency (Fig. 8C).
We then examined the genetic relationship between *vab-1* and the RPM-1 binding protein *glo-*
4/SERGEF. In *vab-1; glo-4* double mutants, we observed both moderate overextension defects
and more severe failed termination defects where the axon hooked ventrally (Fig. 8B). The
frequency of both types of axon termination defects were significantly enhanced in *vab-1; glo-4*
double mutants (Fig. 8B, C). Overextension defects were significantly rescued by transgenic
expression of VAB-1 in *vab-1; glo-4* double mutants (Fig. 8C). We also tested the relationship
between the F-box protein *fsn-1/FBXO45* and *vab-1*, and observed enhanced defects in *vab-1;*
fsn-1 double mutants (Fig. 8D). Finally, we evaluated *vab-1*; *rpm-1* double mutants and did not
observe a significant change in phenotypic severity or penetrance compared to *rpm-1* single
mutants (Fig. 8E).
449
Collectively, these genetic experiments support two conclusions. 1) Enhanced defects in
vab-1; glo-4 and *vab-1; fsn-1* double mutants indicate that the VAB-1 Ephrin receptor functions
in parallel to known RPM-1 binding proteins to facilitate axon termination. These results are
consistent with the VAB-1 Ephrin receptor functioning within the RPM-1 signalling network. 2)
Lack of suppression in *vab-1; fsn-1* and *vab-1; rpm-1* double mutants indicates that VAB-1 is not
inhibited or degraded by the RPM-1/FSN-1 ubiquitin ligase complex. Thus, our findings in *C.*
elegans are consistent with our biochemical studies in mammalian cells that demonstrate
MYCBP2 stabilizes EPHB2 rather than degrading it. We note that because *C. elegans* contains
only a single Ephrin receptor, our findings with *C. elegans* do not necessarily pertain specifically
to EPHB2. Nonetheless, these results add further evidence indicating there are substantial
genetic interactions between known MYCBP2 binding proteins and Eph receptors.
461

Discussion

Our MS-based proteomics efforts to identify EPHB2 interacting proteins yielded MYCBP2, a signalling hub and ubiquitin ligase that is functionally linked to many of the cellular processes also mediated by EPHB2, including cellular growth, proliferation, synapse formation, and axon development. Our experiments argue against EPHB2 being a MYCBP2 ubiquitination substrate. Instead, we envisage a model where MYCBP2 controls EPHB2 signalling indirectly by preventing its lysosomal degradation and maintaining EPHB2 protein levels sufficiently high to mediate efficient cellular and axonal growth cone repulsion from ephrin-B ligands. The interaction between MYCBP2 and EPHB2 may allow the coupling EPHB2 to fundamental cellular processes that control growth, proliferation, and survival. Here, we discuss the molecular logic of MYCBP2 and EPHB2 association in the context of Eph receptor signalling, its potential implications for neural development and diversification of EphB2 signalling.

Functional significance of MYCBP2-EPHB2 complex formation

Our biochemical experiments validate the formation of a MYCBP2-EPHB2 complex and suggest that this association is decreased following ligand application. Because PHR proteins like MYCBP2 are large signalling hubs, MYCBP2 association with EPHB2 might sterically hinder the formation of EPHB2 multimers and clusters necessary for signalling. Thus, ligand-induced dissociation of the MYCBP2-EPHB2 complex could be a prelude to signalling. Our findings indicate that MYCBP2 association with EPHB2 is enhanced by FBXO45, a subunit of the MYCBP2/FBXO45 complex that mediates ubiquitination substrate binding. Previous work showed that MYCBP2 functions to polyubiquitinate specific protein substrates, targeting them for degradation and inhibition (Crawley et al., 2019;Han et al., 2012;Nakata et al., 2005;Desbois et al., 2022). We considered the possibility that the ubiquitin ligase activity of MYCBP2 is important for the termination of EPHB2 signalling, but several lines of evidence argue against this idea: (1) the application of ephrin-B2 ligand results in the dissociation of the MYCBP2-EPHB2 complex (Walker-Daniels et al.), (2) loss of MYCBP2 function results in decreased cellular levels of EPHB2 protein, and (3) impairing MYCBP2 increases ligand-stimulated EPHB2 ubiquitination. In addition, our quantification of EPHB2 signalling suggested that decreased EPHB2 and ERK phosphorylation seen in MYCBP2-deficient cells can be accounted for by the decrease in EPHB2 receptor levels. Thus, a more plausible model that is consistent with our results is that MYCBP2 association with EPHB2 protects EPHB2 from turnover by lysosome-mediated degradation and prevents EPHB2 ubiquitination by the action of unidentified ubiquitin ligases. Interestingly, studies in *C. elegans* have shown that RPM-1/MYCBP2 regulates lysosome biogenesis via the GLO-4/GLO-1 pathway (Grill et al., 2007). While this established a link between MYCBP2 signalling and the endo-lysosomal degradation system, our results now

501 indicate that MYCBP2 can also influence turnover of EPHB2 via lysosomal degradation.
502 Consistent with these findings, we observed that the Eph receptor VAB-1 functions in parallel
503 genetic pathways with the known RPM-1 binding protein GLO-4/SERGEF that regulates
504 lysosome biogenesis.
505
The "protective" effect of MYCBP2 vis-à-vis EPHB2 ubiquitination and lysosomal
degradation might be secondary to effects on other aspects of EPHB2 signalling that we have
not explored experimentally. For example, since MYCBP2 is a signalling hub with multiple
substrates and binding proteins, it could bring EphB2 into close proximity to components of the
MYCBP2 signalling network integrating cell-cell communication via ephrin:Eph signals with
MYCBP2 intracellular signalling to influence fundamental cellular processes. One example of
this is suggested by a recent study demonstrating a link between EPHB2 and mTOR-mediated
cell growth signalling pathways (Gagne et al., 2021). One potential mechanism could involve
Ephrin binding-induced dissociation of the EPHB2-MYCBP2 interaction, allowing MYCBP2
ubiquitination of the TSC complex that regulates mTOR function (Han et al., 2012).
516
The formation of a MYCBP2-EPHB2 complex that includes other signalling receptors
could also explain the curious result that the extracellular domain of EPHB2 is critical for
MYCBP2 association. The finding would be consistent with the extracellular domain of EPHB2
interacting with the extracellular domains of other receptors, whose intracellular domains are
linked more directly with MYCBP2 and FBXO45. An alternative explanation may be a non-
classical extracellular MYCBP2-EPHB2 interaction similar to the one proposed for the
extracellular domain of N-Cadherin and FBXO45 (Na et al., 2020).
524
525
526 **MYCBP2 and EPHB2 functions in the developing nervous system**
527
Our proteomic, biochemical, and genetic experiments indicate that MYCBP2 and EPHB2
function in the same pathway. This is also supported by the striking similarity of MYCBP2 and
EPHB2 loss of function phenotypes in the mouse nervous system. In the context of developing
neuronal connections, *EphB2* and *Mycbp2/Phr1* mouse mutants display similar axon guidance
phenotypes such as abnormal limb nerve trajectories by motor axons, defective growth cone
crossing of the midline at the level of the optic chiasm and decreased connectivity between the
two cortical hemispheres through the corpus callosum (Lewcock et al., 2007;Luria et al.,
2008;Williams et al., 2003;Henkemeyer et al., 1996;D'Souza et al., 2005). Both proteins also
function in synaptic development and their loss of function leads to decreased numbers of
synapses and altered synapse morphology (Wan et al., 2000;Bloom et al., 2007;Dalva et al.,
2000;Zhen et al., 2000). Furthermore, outside the nervous system, both are involved in a variety
of cancers with recent evidence linking them to esophageal adenocarcinoma and c-MYC-

dependent control of cell proliferation (Venkitachalam et al., 2022;Han et al., 2012;Genander et
al., 2009). Our proteomic, biochemical, and cellular experiments together with genetic interaction
studies in *C. elegans* now provide a new framework in which to consider phenotypic and disease
links between EPHB2 and MYCBP2.

Importantly, the biomedical relevance of our findings are heightened by a recent study
that identified genetic variants in MYCBP2, which cause a neurodevelopmental disorder termed
MYCBP2-related Developmental delay with Corpus callosum Defects (MDCD) (Al-Abdi et al.,
2022). MDCD features defective neuronal connectivity including a hypoplastic or absent corpus
callosum, neurobehavioral deficits including intellectual disability and epilepsy, and abnormal
craniofacial development. This constellation of comorbidities in MDCD closely resembles some
of the phenotypes observed in mice with deficient EPHB2 signalling. Given our finding that
MYCBP2 loss reduces EPHB2 levels and influences Eph receptor effects on axon development,
the question of whether EPHB2 expression levels are normal in MYCBP2 patients remains
pertinent.

In conclusion, our study has revealed numerous biochemical and genetic links between
MYCBP2 and EPHB2. Our findings indicate that the MYCBP2/FBXO45 complex protects
EPHB2 from degradation, and these are functionally integrated signalling players with an
evolutionarily conserved role in axonal development. Future studies will be needed to address
how the EPHB2-MYCBP2 interaction affects nervous system development in mammals *in vivo*
and to identify further regulators of EPHB2 degradation. Additionally, another idea worthy of
closer examination in the future is the possibility that MYCBP2 signalling could provide routes
through which EPHB2-initiated signals access numerous fundamental cellular functions.

Acknowledgments

We thank Dominic Fillion for assistance with microscopy, Denis Flaubert and the IRCM Proteomics Core for proteomics analysis, Gu Hua for HA-Ubiquitin plasmid, Sylvie Lahaie, Louis-Philippe Croteau and Farin B. Bourojeni for discussions, and Meirong Liang for technical support. This work was supported by project grants from the Canadian Institutes of Health Research (PJT-162225, MOP-77556, PJT-153053, and PJT-159839) to AK and a grant from the National Institutes of Health (R01 NS072129) to BG. CC received IRCM Jacques-Gauthier Scholarship. S. L. B. is a Fonds de recherche du Québec – Santé (FRQS) scholar. SSP is funded by IRCM Jean-Coutu Scholarship.

Author contribution

All authors: Conceptualisation, Writing – Editing and Review.

CC: Methodology, Investigation, Data Curation, Writing – Original Draft, Visualisation.

SLB: Methodology, Investigation, Data Curation, Visualisation, Supervision, Writing – Original Draft.

SSP: Methodology, Investigation, Data Curation, Visualisation.

XZ: Methodology, Investigation.

DCW: Methodology, Investigation.

MD: Methodology, Investigation, Resources, Data Curation, Visualisation, Supervision, Writing – Original Draft.

BG: Supervision, Funding Acquisition, Writing – Original Draft.

AK: Writing – Original Draft, Supervision, Project Administration, Funding Acquisition.

Conflict of interest

The authors declare no competing financial interests.

Materials and Methods

Vertebrate animals

All animal experiments were carried out in accordance with the Canadian Council on Animal Care guidelines and approved by the IRCM Animal Care Committee. Fertilized chicken eggs (FERME GMS, Saint-Liboire, QC, Canada) were incubated at 38-39°C and staged according to Hamburger and Hamilton (HH) (Hamburger and Hamilton, 1951). C57BL/6 mice were used for hippocampal neuron collapse assay. Timed mating vaginal plug was designated as E0.5.

Cell culture

HeLa and HEK293T cells were maintained in DMEM (Thermo Fisher Scientific, #11965092) supplemented with 10% Fetal Bovine Serum (FBS; Wisent Bioproducts, #080-150) and 1% Penicillin/Streptomycin (Wisent Bioproducts, #450-200-EL) at 37°C with 5% CO_2. Tetracycline inducible HeLa EPHB2-FLAG cells were generated by transfecting Flp-In T-REx HeLa cells with EPHB2-BirA*-FLAG expression plasmid using Lipofectamine 3000 (Invitrogen, #L3000015) followed by hygromycin selection (200µg/ml). Stable but not clonal CTRL[CTRISPR] or MYCBP2[CRISPR] HeLa EPHB2-FLAG cell lines were generated by infecting cells with packaged lentivirus, followed by puromycin selection (1µg/ml). Lentivirus particles were packaged using MYCBP2 sgRNA CRISPR plasmid designed to target the *MYCBP2* exon6 (pLentiCRISPR2-sgMYCBP2) and pLentiCRISPR empty vector was used as Ctrl CRISPR. EPHB2-FLAG overexpression was induced using 1µg/ml tetracycline simultaneously with cell starvation in DMEM supplemented with 0.5%FBS, 1% penicillin/streptomycin for 12-20h. Prior to cell stimulation, Fc control (Millipore, #401104), ephrinB1-Fc (R&D, #473-EB) or ephrinB2-Fc (R&D, #496-EB) were pre-clustered using goat anti-human Fc IgG (Sigma, #I2136) in 4:1 ratio for 30min.

Affinity purification - mass spectrometry

HeLa EPHB2-FLAG cells cultured in DMEM supplemented with 0.5% FBS and 1µg/ml tetracycline in 15cm cell culture plates for 20h, were treated with pre-clustered Fc control or ephrinB2-Fc for 15min. After treatment, cells were washed twice with PBS and lysed using a lysis buffer (50mM Tris, pH7.4; 150mM NaCl; 1% NP-40) supplemented with protease (Roche, #11836153001) and phosphatase inhibitors (Roche, #04906837001). The lysates were collected in 1.5ml Eppendorf tubes and centrifuged at 13,000rpm for 15min at 4°C. Supernatants were transferred to new tubes with prewashed anti-FLAG agarose beads (Sigma, #A2220) and incubated on a rotator overnight at 4°C. The following day, beads were washed four times using 50mM Ammonium Bicarbonate. The on-bead proteins were diluted in 2M Urea/50mM ammonium bicarbonate and on-bead trypsin digestion was performed overnight at 37°C. The samples were then reduced with 13 mM dithiothreitol at 37°C and, after cooling for 10min, alkylated with 23 mM iodoacetamide at room temperature for 20min in the dark. The supernatants were acidified with trifluoroacetic acid and cleaned from residual detergents and

reagents with MCX cartridges (Waters Oasis MCX 96-well Elution Plate) following the
manufacturer's instructions. After elution in 10% ammonium hydroxide /90% methanol (v/v),
samples were dried with a Speed-vac, reconstituted under agitation for 15min in 12μL of
2%ACN-1%FA and loaded into a 75μm i.d. × 150 mm Self-Pack C18 column installed in the
Easy-nLC II system (Proxeon Biosystems). Peptides were eluted with a two-slope gradient at a
flowrate of 250nL/min. Solvent B first increased from 2 to 35% in 100min and then from 35 to
80% B in 10min. The HPLC system was coupled to Orbitrap Fusion mass spectrometer (Thermo
Scientific) through a Nanospray Flex Ion Source. Nanospray and S-lens voltages were set to
1.3-1.7 kV and 60 V, respectively. Capillary temperature was set to 225°C. Full scan MS survey
spectra (m/z 360-1560) in profile mode were acquired in the Orbitrap with a resolution of 120,000
with a target value at 3e5. The 25 most intense peptide ions were fragmented in the HCD
collision cell and analyzed in the linear ion trap with a target value at 2e4 and a normalized
collision energy at 29 V. Target ions selected for fragmentation were dynamically excluded for
15 sec after two MS2 events.
647
Mass spectrometry data analysis
The peak list files were generated with Proteome Discoverer (version 2.3) using the following
parameters: minimum mass set to 500 Da, maximum mass set to 6000 Da, no grouping of
MS/MS spectra, precursor charge set to auto, and minimum number of fragment ions set to 5.
Protein database searching was performed with Mascot 2.6 (Matrix Science) against the UniProt
Human protein database. The mass tolerances for precursor and fragment ions were set to 10
ppm and 0.6 Da, respectively. Trypsin was used as the enzyme allowing for up to 1 missed
cleavage. Cysteine carbamidomethylation was specified as a fixed modification, and methionine
oxidation as variable modification. Data interpretation was performed using Scaffold (version 4.8)
and further statistical analysis was performed through ProHits integrated with SAINT
(Significance Analysis of INTeractome) (Liu et al., 2010).
659
Cell lysis, co-immunoprecipitation and Western blotting
Cells were washed with PBS, lysed with RIPA buffer (50mM Tris, pH 7.4; 150mM NaCl; 1% NP-
40; 0.1% SDS) supplemented with protease and phosphatase inhibitors. For co-IP experiments,
cells were lysed with co-IP buffer (50mM Tris, pH 7.4; 150mM NaCl; 0.1% NP-40) with protease
and phosphatase inhibitors. Cell lysates were centrifuged at 13,000 rpm for 20min at 4°C, then
the supernatants were collected, and total protein concentrations were quantified using BCA kit
(Thermo Scientific, #23225). For FLAG co-IP, 500-1000μg of total protein was incubated with
20-40 μl of prewashed anti-FLAG agarose beads (Sigma, #A2220) for 3h at 4°C. After incubation,
the beads were centrifuged at 2,600rpm for 1min at 4℃ and washed three times with the co-IP
buffer. The beads were resuspended in 2xSDS-PAGE loading buffer (5x loading buffer: Tris,
150mM, pH 6.8; SDS, 10%; Glyercol, 30%; b-Mercaptoethanol, 5%; Bromophenol Blue, 0.02%).
For western blotting, proteins were separated on 6-10% SDS-PAGE gels and transferred to
methanol pre-activated PVDF membranes (Millipore, #IPVH00010). For MYCBP2 blots, gels

were wet transferred overnight at 4°C using 33V. Membranes were incubated in blocking buffer
(TBST: 20mM Tris, pH 7.6; 150mM NaCl; 0.1% Tween 20; 5% skim milk) for 1h at room
temperature, followed by primary antibody incubation (1-2h at room temperature or overnight at
4°C) and corresponding secondary antibody incubation (1h at room temperature). Primary
antibodies were: rabbit polyclonal anti-MYCBP2 (Abcam, #ab86078; RRID:AB_1925230), rabbit
anti-pERK1/2 (Thr202/Tyr204; Cell Signaling Technology, #9101; RRID:AB_331646), rabbit
anti-ERK1/2 (Cell Signaling Technology, #9102; RRID:AB_330744), goat polyclonal anti-EPHB2
(R&D Systems, #AF467; RRID:AB_355375), mouse monoclonal anti-Actin (Sigma-Aldrich,
#A5441; RRID:AB_476744), mouse monoclonal anti-pTyr (PY20; Santa Cruz Biotechnology,
#sc-508; RRID:AB_628122), mouse monoclonal anti-GAPDH (Santa Cruz Biotechnology, #sc-
47724; RRID:AB_627678), mouse monoclonal anti-FLAG-HRP (Sigma-Aldrich, #A8592;
RRID:AB_439702), rabbit monoclonal anti-HA (Cell Signaling Technology, #3724;
RRID:AB_1549585), mouse monoclonal anti-MYC (Santa Cruz Biotechnology, #sc-40;
RRID:AB_627268), rabbit polyclonal anti-GFP (Thermo Fisher Scientific, #A-11122;
RRID:AB_221569). Secondary antibodies were: Donkey anti-Goat HRP (Jackson, 705-035-003),
Donkey anti-Mouse HRP (Jackson, 715-035-151), Donkey anti-Rabbit HRP (Jackson, 711-035-
152). After three washes with TBST, membranes were incubated with ECL reagent (Cytiva,
RPN2106) for 1min and chemiluminescence signal was acquired using film or Bio-Rad
ChemiDoc Imaging machine. Band intensity was quantified using ImageJ or Bio-Rad Image Lab
software.
693
HeLa cell collapse assay
HeLa-EPHB2 CTRL[CRISPR] and MYCBP2[CRISPR] cells were seeded on glass coverslips (Electron
Microscopy Sciences, #7223101) in 24 well plates at a density of 20 000 cells/well. After 24
hours, cell media was changed for DMEM supplemented with 0.5% FBS, 1% P/S and 1μg/ml
tetracycline to starve the cells and induce EPHB2 expression for 16-20 hours. Cells were then
stimulated with 1.5μg/ml of pre-clustered Fc control or ephrin-B2-Fc for 15min. Cells were fixed
with 3.2% paraformaldehyde (Lewcock et al.), 6% sucrose in PBS for 12-15min. Nuclei were
stained with DAPI and F-actin with Phalloidin Alexa Fluor 568 conjugate (Thermo Fisher,
#A12380). Images were acquired using Zeiss LSM710 confocal microscope and 20x objective.
Fully rounded cells are scored as collapsed cells.
704
For time lapse imaging experiments, CTRL[CRISPR] or *MYCBP2*[CRISPR] cells were plated on
Poly-D-Lysine coated glass bottom 35mm dishes (MATTEK, #P35GC-1.5-10-C) at a density of
300,000 cells/dish. The next day, cells were transfected with 1.5μg of EPHB2-GFP plasmid for
4-5 h, using lipofectamine 3000 in opti-MEM (ThermoFisher, #31985070), and then media was
changed for DMEM supplemented with 0.5% FBS. The following day, the images were acquired
under Zeiss Spinning Disk Microscope using a 20x objective. During the imaging, the cells were
maintained at 37°C with 5% CO_2. Pre-clustered ephrinB2-Fc was added to a final concentration

of 2μg/mL directly into the dishes at the beginning of each experiment. The images were
acquired every minute for 1h.
714

HeLa cell stripe assay

Alternative ephrin-B2-Fc or Fc stripes were prepared using silicon matrices with a micro-well
system (Poliak et al., 2015). HeLa EPHB2 CTRL[CRISPR] and MYCBP2[CRISPR] cells, or HeLa EPHB2
cells transiently transfected with GFP-FBD1 wild-type or GFP-FBD1 mutant (GRR/AAA: G2404A,
R2406A, R2408A), were cultured with tetracycline for 20h, trypsinized and plated on stripes (~10
000 cells per carpet); see (Desbois et al., 2018) for specific sequences. The next day, cells were
fixed with paraformaldehyde (3.2% PFA, 6% sucrose in PBS), and stained with DAPI and
Phalloidin Alexa Fluor 488; or DAPI, Rabbit anti-GFP (Thermo Fisher Scientific, # A-11122;
RRID:AB_221569) and Phalloidin Alexa Fluor 647 (Abcam, #ab176759). Images were acquired
using Zeiss LSM710 confocal microscope and 20x objective (three vision fields for each carpet).
Cells on or off the ephrin-B2 stripes were decided by the nuclei localization of the cells.
726

Ubiquitination assay

HeLa EphB2 CTRL[CRISPR] and MYCBP2[CRISPR] cells were seeded in 6-well plates at a density of
0.5 million cells per well. Next day, cells were transfected with 1.2μg HA-Ubiquitin (gift of Gu
Hua) using lipofectamine 3000 (2μl/well) for 4-5h in opti-MEM (ThermoFisher, #31985070), then
media was changed to DMEM with 10%FBS. The following day, EPHB2 expression was induced
using 1μg/ml tetracycline in DMEM with 0.5% FBS for 12h followed by 2μg/ml Fc or eB2-Fc
treatment for 30min. After IP using anti-FLAG beads, precipitates were eluted with 2xSDS
loading buffer, resolved using 8% SDS gel and transferred onto PVDF membranes. Membranes
were blocked in 5% milk following an incubation with anti-HA antibody (Cell Signaling
Technology, #3724) to detect EPHB2 ubiquitination levels. The membrane was then stripped
using mild stripping buffer (1L: 15g glycine, 1g SDS, 10mL Tween 20, pH 2.2) and probed with
anti-FLAG antibody to reveal EPHB2 levels.
739

Lysosome and proteasome inhibition

741 HeLa EPHB2 CTRL[CRISPR] and MYCBP2[CRISPR] cells were seed in 6-well plates at a density of 0.5
742 million cells per well. Next day, EPHB2 overexpression was induced using 1μg/ml tetracycline
743 in DMEM with 10% FBS for 16h, followed by 26S proteasome inhibitor (MG132, 50 μM, Sigma,
744 #474790) or lysosome inhibitor treatment (BafilomycinA1, 0.2 μM, Sigma, #B1793; Chloroquine,
745 50 μM, Tocris, #4109) for 6hr.
746

Chick *in ovo* electroporation

Fertilized eggs (FERME GMS, Saint-Liboire, QC) were incubated in an incubator (Lyon
Technologies, model PRFWD) at 39°C with a humidity level of around 40%-60% according to
standard protocols. At HH stage.15-17, chick embryo spinal neural tubes were electroporated
with expression constructs (TSS20 Ovodyne electroporator at 30V, 5 pulses, 50ms wide,

1000ms interval). Following electroporation, eggs were sealed with double layer of parafilm
(Pechiney Plastic Packaging Company) and incubated till HH stage 24-26.
754

Chick spinal explants stripe assay
Alternative eprhinB2-Fc/Fc stripes were prepared using silicon matrices with a micro-well system
and pre-coated with laminin. At HH stage 24-26, chick embryos were harvested, and the lumbar
part neural tubes were dissected with tungsten needles (World Precision Instruments) in MN
medium (20ml motor neuron medium: 19.2ml Neurobasal medium, 400µl B27 supplement, 200µl
50mM L-glutamic acid, 200µl 100xP/S antibiotics,73mg L-glutamine). The lumbar neural tube
was then cut into around 20 explants and the explants were plated on stripes (a 1cmx1cm square
covers the whole stripe area). After overnight incubation, the explants were fixed with 4% PFA
for 12min at 37°C, washed once with PBS, and incubated with blocking buffer, primary antibodies,
and secondary antibodies. After three times PBS washes, the explants on stripes were mounted
and explants with extensions were imaged using LSM710 (at least three explants per carpet).
The percentage of branches on ephrinB2 stripes was calculated using the total length of
branches localized on ephrinB2 stripes divided by the total length of the branches of the explant.
768

Dissociated mouse hippocampal neuron culture and electroporation
Primary hippocampal neurons were cultured from wild-type C57BL/6 mice at embryonic day 16–
18 (E16-18). The hippocampi were dissected out and collected in 4.5ml dissection buffer
(calcium- and magnesium-free Hank's BSS: 500 ml distilled water (Gibco, #15230162), 56.8 ml
10xHBSS (Gibco, #14185052), 5.68ml 1M HEPES (Gibco, #15630080), 2.84ml HyClone
(Thermo Scientific, #SV30010)). Hippocampi were added with 0.5ml 2.5%Tyrpsin and incubated
at 37°C for 13-15min. After 5 times of thorough wash with dissection buffer, hippocampal
neurons were dissociated in 0.8ml DMEM (Gibco, #11965118) with 10% FBS (Wisent
Bioproducts, 080-150) by pipetting 10 times up and down. Then cell numbers were counted and
desired number of neurons were directed for electroporation. After spin down at 2,000 rpm for
2min, dissociated hippocampal neurons ($1x10^6$/condition) were resuspended with 100µl
homemade nucleofection solution, mixed with 5ug of DNA, and transferred into the aluminum
cuvettes (AMAXA/Lonza). Electroporation was achieved by Nucleofector I (AMAXA/Lonza)
using program O-05 (Mouse CNS neurons). 1ml Plating Medium (PM; 500ml MEM (Sigma,
#M4655), 17.5ml 20% Glucose (Sigma, #G8270), 5.8ml 100mM pyruvate (Sigma, #P2256),
58ml heat-inactivated horse serum (Thermo Scientific, #26050088)) was added to the cuvettes
immediately, and desired number of neurons were plated on 1mg/ml Poly-L-Lysine (Sigma,
#P2636) coated coverslips in 12-well plate. At 1 day *in vitro* (DIV1), medium was replaced with
Neuron growth and maintenance medium (NBG; 500ml Neurobasal medium (Gibco,
#21103049), 10ml SM1 neuronal supplement (Stemcell, #05711), and 1.25ml GlutaMAX-I
(Gibco, #35050)).
790

Hippocampal neuron growth cone collapse assay

Electroporated hippocampal neurons were cultured on glass coverslips (18 mm, 100 thousand
neurons/well in a 12-well plate). At DIV2, the neurons were treated with pre-clustered ephrinB1-
Fc, ephrinB2-Fc or Fc control in NBG medium at a final concentration of 2µg/ml for 60min at
37°C. After treatment, neurons were fixed with paraformaldehyde (3.2% PFA,6% sucrose in PBS)
for 12min, followed by two PBS washes, and blocked with Blocking Buffer (PBS, 0.15% TritonX-
100, 2% FBS) for 60min at room temperature. Neurons were then incubated with rabbit anti-
GFP antibody (1:5000, Thermo Fisher Scientific, #A11122; RRID:AB_221569) in Blocking Buffer
for 90min at room temperature. Followed by three PBS washes, neurons were incubated with
DAPI, donkey anti-rabbit IgG Alexa Fluor 488 conjugate (1:1000, Jacksonimmuno, #711545152)
and Phalloidin Alexa Fluor 568 conjugate (1:300, Invitrogen, #A12380) in Blocking Buffer for
60min. Neurons on coverslips were then washed and mounted on microscope slides
(Fisherbrand, #1255015). Collapsed growth cones were scored using followed criteria.
804
Collapsed hippocampal neuron growth cone quantification
Neuron selection: only neurons with moderate GFP staining (neurons with strong GFP signalling
were excluded) and only neurons with more than three branches were scored. Branch selection:
branches shorter than the diameter of the neuron cell body were excluded; branches
intermingling with others were excluded. Collapsed growth cone: growth cones of a fan shape
are scored as full, growth cones with a width smaller than that of the branches are scored as
collapsed, and growth cones' size in between are scored as "hard to tell". The collapse rate was
calculated using collapse growth cone numbers divided by the total growth cone numbers.
813
814 **Microscopy and imaging**
815 HeLa cell collapse assay images were acquired using Leica DM6 or Zeiss LSM710 confocal
816 microscopy. Stripe assay and growth cone collapse assay images were acquired with a Zeiss
817 LSM710 or LSM700 confocal microscope.
818
Quantification and statistical analysis
All cell counts, collapsed/uncollapsed visualization and explant neurite length measurements
were performed with ImageJ 2.9.0 (Schindelin et al., 2012). All numbers are illustrated in figure
legends. In western blotting, each n represents one independent experiment; in neuron growth
cone collapse assay, each n represents one independent experiment with neurons pooled from
multiple embryos. All data statistical analyses were performed using GraphPad Prism 9.1.1. Test
methods and p values were described in figure legends, with p value 0.05 as a significance
threshold.
827
828 **Data Availability**

829   The mass spectrometry proteomics data have been deposited to the ProteomeXchange
830   Consortium via the PRIDE partner repository (Perez-Riverol et al., 2019) with the dataset
831   identifier PXD041786 and 10.6019/PXD041786.
832

C. elegans genetics and strains

834   *C. elegans* N2 isolate was used for all experiments. Animals were maintained using standard
835   procedures. The following mutant alleles were used: *vab-1*(*dx31*) II, *fsn-1*(*gk429*) III, *rpm-1*(*ju44*)
836   V, *glo-4*(*ok623*) V. The integrated transgene used to evaluate axon termination was *muIs32*
837   [P$_{mec-7}$GFP] II. For genetic analysis the animals were grown at 23 ℃.
838

Transgenic extrachromosomal arrays were generated using standard microinjection
procedures for *C. elegans*. *vab-1* minigene (pCZ47) was injected at either 25 ng/μL or 50 ng/μL,
and co-injection markers used for transgene selection were either neomycin resistance (pBG-
264) or P*ttx-3*::RFP (pBG-41). pBluescript (pBG-49) was used to reach a final concentration of
100 ng/μl in all injection mixes. pCZ47 was a gift from Andrew Chisholm (Addgene plasmid #
128414 ; RRID:Addgene_128414)

C. elegans axon termination analysis and imaging

Axon termination defects were defined as PLM axons that extended beyond the normal
termination point adjacent to the ALM cell body. Two different failed termination phenotypes
were scored: axon overextension (moderate phenotype) where the PLM axon grew beyond ALM
cell body, axons that overextend and form a ventral hook (severe phenotype). To quantify axon
termination defects, 20-30 young adult animals were anesthetized (10 μM levamisole in M9
buffer) on a 2% agar pad on glass slides and visualized with a Leica DM5000 B (CTR5000)
epifluorescent microscope (40x oil-immersion objective). For image acquisition, young adult
animals were mounted on a 3% agarose pad and a Zeiss LSM 710 (40x oil-immersion objective)
was used to generate z stacks.
855

For statistical analysis of axon termination defects, comparisons were done using a
Student's *t*-test with Bonferroni correction on GraphPad Prism software. Error bars represent
standard error of the mean (SEM). Significance was defined as $p < 0.05$ after Bonferroni
correction. Bar graphs represent averages from 8 to 10 counts (20–30 animals/count) obtained
from eight or more independent experiments for each genotype. For transgenic rescue
experiments, data shown in Figure 8B was obtained from two, independently derived transgenic
extrachromosomal arrays.
863
864

Figures and figure legends

Figure 1. MS-proteomics and biochemistry in HeLa cells identifies MYCBP2 as EPHB2 binding protein.

A) Schematic of EPHB2 affinity purification coupled to mass spectrometry (AP-MS) workflow. B) Scatter plot of AP-MS data showing known and putative EPHB2 binding proteins, including MYCBP2. Y axis and X axes represent the average spectral counts of the identified protein hits in the EPHB2 protein complexes from cells stimulated with Fc control or ephrin-B2 (eB2-Fc), respectively. C) In HEK 293T cells, endogenous MYCBP2 is pulled down by transiently overexpressed EPHB2-FLAG but not by EPHA3-FLAG. D) In EPHB2-FLAG stable HeLa cell line, ephrin-B stimulation reduces the interaction between MYCBP2 and EPHB2. E) Quantification of MYCBP2-EPHB2 association intensity after Fc, ephrin-B1 (eB2-Fc) or ephrin-B2 (eB2-Fc) treatment (eB1-Fc, p=0.1365; eB2-Fc, p=0.0002; one-sample t-test). Error bars represent standard deviation (SD). F) Representative image of MYCBP2 pull down with anti-

878   EPHB2 or IgG control antibodies from rat cortical neurons. Asterisk indicates MYCBP2. G)

879   Representative images from western blot analysis of endogenous MYCBP2 following IP of GFP-EPHB2

880   wild-type (WT) or its kinase dead (KD) counterpart.

Figure 2. Mapping binding regions for EPHB2-MYCBP2 reveals role of FBXO45 in this interaction.

A) Co-IP of EPHB2-FLAG with MYC-FBXO45 using transfected HEK293 cells. EPHB2 co-precipitates FBXO45, but EPHA3 does not. Asterisk indicates MYC-FBXO45. B) In HEK 293T cells, FBXO45 overexpression enhances EPHB2-MYCBP2 binding. Asterisk indicates MYC-FBXO45. C) Quantification of the association intensity of MYCBP2 and EPHB2 upon FBXO45 overexpression (EPHB2, $p=0.0068$; EPHB2 vs EPHA3, $p=0.0005$; one sample t-test). Error bars represent SD. D) Schematic representation of MYCBP2 N-terminal, Central and C-terminal fragments. E) Co-IP of EPHB2-FLAG with GFP-MYCBP2 fragments in HEK 293T cells. EPHB2 coprecipitates with MYCBP2 central fragment. Asterisks indicate GFP-MYCBP2 fragments. F) Schematic of chimeric domain swapping of EPHB2 (orange) and EPHA3 (grey). G) Co-IP of MYC-FBXO45 and endogenous MYCBP2 with EPHB2/EPHA3 domain swapped chimeras. H) Schematic representation of EPHB2 ΔECD (extracellular domain, aa deletions of 19-530) and ΔICD (intracellular domain, aa deletions of 590-986) truncations. I) Co-IP of endogenous MYCBP2 with EPHA3, EPHB2 and EPHB2 truncation mutants. ECD, extracellular domain; TM, transmembrane; JM, juxtamembrane; SAM, Sterile alpha motif; PBM, PDZ (PSD-95, Dlg1, Zo-1) binding motif.

Figure 3. MYCBP2 CRISPR HeLa cells exhibit reduced ephrin-B2 evoked cell retraction and ephrin-B2 stripe avoidance.

A) MYCBP2 is reduced in HeLa *MYCBP2*CRISPR cells generated by sgRNA targeting MYCBP2 exon 6. B) Representative images of cell collapse assays using CTRLCRISPR or *MYCBP2*CRISPR HeLa cells that were stimulated with ephrin-B2. Red arrows indicate rounded/collapsed cells. C) Quantification of collapsed cells. Statistical significance between CTRLCRISPR and *MYCBP2*CRISPR cells was determined using two-way ANOVA followed by Sidak's multiple comparison test (CTRLCRISPR vs *MYCBP2* CRISPR: Fc, p=0.9841; eB2-Fc, p=0.0017. Fc vs eB2-Fc: CTRLCRISPR, p<0.0001; *MYCBP2* CRISPR, p=0.0115). D) Representative time-lapse sequences of CTRLCRISPR and *MYCBP2*CRISPR HeLa cells after ephrin-B2 treatment. E) Quantification of cell area reduction after 60 min exposure to ephrin-B2 (p=0.0268, two-tailed unpaired t

908   test). F) Ephrin-B2 stripe assays using CTRL[CRISPR] or *MYCBP2*[CRISPR] HeLa cells. Cells are visualized with
909   Phalloidin 488 staining and nuclei are stained with DAPI (black, Fc stripes; red, ephrin-B2 stripes). G)
910   Quantification of cells present on ephrin-B2 stripes (%). Statistical significance was determined using
911   two-tailed unpaired t-test (p=0.0109). Error bars represent SD.
912

Figure 4. MYCBP2 loss-of-function increases EPHB2 protein turnover in HeLa cells.

A) Induced EPHB2-FLAG expression is reduced in *MYCBP2*^{CRISPR} HeLa cells. B) Western blotting for transfected EPHB2-FLAG in CTRL^{CRISPR} and *MYCBP2*^{CRISPR} cells. C) Quantification of transfected EPHB2-FLAG levels (p=0.0046, one-sample t-test). D) Representative western blot of EPHB2-FLAG in CTRL^{CRISPR} and *MYCBP2*^{CRISPR} cells treated with DMSO or cycloheximide for 4h and 8h. E) Quantification of EPHB2-FLAG turnover with cycloheximide (CTRL^{CRISPR} vs. *MYCBP2*^{CRISPR} at 8h eB2-Fc stimulation, p= 0.0474, two-way ANOVA followed by Tukey's multiple comparison test). F) Western blot showing EPHB2-FLAG degradation when cells are challenged with ephrin-B2 (1µg/ml) for different periods of time. G) Quantification of ephrin-B2-evoked EPHB2 degradation (ns, not significant; two-way ANOVA followed by Tukey's multiple comparison test). H) Western blot of EPHB2 ubiquitination in CTRL^{CRISPR} and *MYCBP2*^{CRISPR} cells. I) Quantification of ubiquitinated EPHB2. CTRL^{CRISPR} cells stimulated with Fc vs. eB2-Fc, p= 0.1349 (One sample t-test); *MYCBP2*^{CRISPR} cells stimulated with Fc vs. EB2-Fc, p= 0.0195 (Unpaired two-tailed t-test). J) After tetracycline induction of EPHB2-FLAG expression for 16hrs, CTRL^{CRISPR} and *MYCBP2*^{CRISPR} HeLa cells were treated with DMSO (1:500) or inhibitors of the proteasome (MG132 50µM) or lysosome (BafA1 0.2µM; CoQ 50µM) for 6 hours, and EPHB2 levels were analysed by western blotting. K) Quantification of EPHB2 levels following treatment with proteasome or lysosome inhibitors. Statistical significance for the comparison between CTRL^{CRISPR} cells treated with DMSO or inhibitors was determined by one-sample t-test (MG132, p=0.0598; BafA1, p=0.0200; CoQ, p=0.3632), whereas statistical significance for the comparison between *MYCBP2*^{CRISPR} cells treated with DMSO and individual inhibitors was determined by two-tailed paired t-test (MG132, p=0.8893; BafA1, p=0.0361; CoQ, p=0.0835). L) GFP-EPHB1 and HA-EPHB3 transfected into CTRL^{CRISPR} and *MYCBP2*^{CRISPR} HeLa cells and detected by Western blot. M) Quantification of GFP-EPHB1 and HA-EPHB3 levels (EPHB1, p=0.0588; EPHB3, p=0.4253; one-sample t-test). N) FLAG-EPHA3 transfected into CTRL^{CRISPR} and *MYCBP2*^{CRISPR} HeLa cells and detected by WB. O) Quantification of FLAG-EPHA3 (p=0.5369, one-sample t-test). Error bars represent SD.

Figure 5. MYCBP2 depletion impairs EPHB2 phosphorylation and ERK1/2 activation in HeLa cells.
A) Representative western blot for pERK1/2 and pTyr-EPHB2 detected in CTRL[CRISPR] and *MYCBP2*[CRISPR]
cells treated with ephrin-B2 (eB2-Fc) for different periods (n=6). Membranes were striped and reblotted
with anti-ERK1/2, anti-EPHB2, anti-GAPDH and anti-MYCBP2 antibodies as controls. B) Quantification
of EPHB2 tyrosine phosphorylation in CTRL[CRISPR] cells evoked by ephrin-B2 treatment (15 min, p=0.1363;

30 min, p=0.2056; 1h, p=0.0342; 2h, p=0.0234; 4h, p=0.0068; 8h, p= 0.0231; one-sample t-test). C)
Quantification of EPHB2 tyrosine phosphorylation in CTRL^{CRISPR} and *MYCBP2*^{CRISPR} HeLa cells
(unstimulated, p=0.5589, one-sample t-test; stimulated for 15 min, p=0.7463; 30 min, p=0.5520; 1h,
p=0.1920; 2h, p=0.2009; 4h, p=0.1550; 8h, p=0.0331; two-tailed unpaired t-test). D) Quantification of
pERK1/2 in CTRL^{CRISPR} and *MYCBP2*^{CRISPR} HeLa cells (0 min, p=0.0168, one-sample t-test; 15 min,
p=0.6695; 30 min, p= 0.6649; 1h, p= 0.1776; 2h, p= 0.1479; 4h, p= 0.0494; 8h, p= 0.0078; two-tailed
unpaired t-test). E) Quantification of EPHB2 in CTRL^{CRISPR} and *MYCBP2*^{CRISPR} HeLa (0 min, p=0.4604,
one-sample t-test; 15 min, p=0.2222; 30 min, p=0.1376; 1h, p=0.0651; 2h, p=0.0736; 4h, p=0.2451; 8h,
p=0.0437, two-tailed unpaired t-test). F) Quantification of ephrin-B2-evoked EPHB2 tyrosine
phosphorylation levels relative to total EPHB2 protein levels (0 min, p=0.7058, one-sample t-test; 15 min,
p=0.2464; 30 min, p=0.7835; 1h, p=0.9164; 2h, p=0.7196; 4h, p=0.8625; 8h, p=0.5750, two-tailed
unpaired t-test). G) Quantification of ephrin-B2-evoked pERK1/2 relative to EPHB2 total protein levels (0
min, p=0.3308, one-sample t-test; 15 min, p=0.3856; 30 min, p=0.0624; 1h, p=0.2683; 2h, p=0.1284; 4h,
p=0.7998; 8h, p=0.7790, two-tailed unpaired t-test. Error bars represent SD.
960

Figure 6. Exogenous FBD1 fragment of MYCBP2 disrupts EPHB2-MYCBP2 binding and impairs EPHB2 function in HeLa cells.

A) Schematic illustrating competition of exogenous MYCBP2-FBD1 fragment that disrupts MYCBP2-FBXO45 binding and leads to MYCBP2 reduction in EPHB2 complexes. B) Exogenous FBD1 WT overexpression leads to reduced EPHB2-MYCBP2 binding in HEK293 cells despite co-expression of FBXO45. C) FBD1 overexpression also disrupts EPHB2-MYCBP2 binding in the absence of FBXO45 overexpression. D) Representative images of ephrin-B2 stripe assays using HeLa cells expressing GFP-FBD1 mut or GFP-FBD1 WT. E) Quantification of cells present on eB2 stripes (p=0.0107, two-tailed unpaired t-test). Error bars represent SD.

Figure 7. Exogenous FBD1 overexpression impairs EPH receptor functions in chick spinal cord explants and mouse hippocampal neurons.

A) Representative images of ephrin-B2 stripe assays with chick embryonic spinal cord explants overexpressing GFP-FBD1 mut (negative control) or GFP-FBD1 WT. Images with inverted GFP signal in dark pixels on Fc (white) / eB2 (pink) stripes are placed beside the original images. B) Quantification of GFP-positive branches present on ephrin-B2 stripes (GFP-FBD1 mut vs. GFP-FBD1 WT, p=0.0410, two-tailed unpaired t-test). C) Representative images of DIV2 mouse hippocampal neurons overexpressing GFP-FBD1 mut or GFP-FBD1 WT and challenged with Fc control, ephrin-B1 (eB1-Fc) or ephrin-B2 (eB2-Fc). D) Quantification of growth cone collapse rate for hippocampal neurites. GFP-FBD1 mut: Fc vs. eB1-Fc, p=<0.0001; Fc vs. eB2-Fc, p=0.0008; GFP-FBD1 WT: Fc vs. eB1-Fc p=0.4357; Fc vs eB2-Fc p=0.0065. Two-way ANOVA followed by Tukey's multiple comparisons test. Error bars represent SD.

Figure 8. *C. elegans VAB-1 ephrin receptor interacts genetically with two known RPM-1/MYCBP2 binding proteins FSN-1/FBXO45 and GLO-4/SERGEF.*

A) Schematic shows the known RPM-1/MYCBP2 binding proteins GLO-4/SERGEF and FSN-1/FBXO45. GLO-4 functions independent of RPM-1 ubiquitin ligase. FSN-1 is the F-box protein that forms a ubiquitin ligase complex with RPM-1. Adapted from Grill et al., 2016. B) Schematic of axon morphology and axon termination site for PLM mechanosensory neurons. Shown are representative images of failed axon termination defects observed for PLM neurons of indicated genotypes. Examples of moderate overextension defects (arrowhead) observed in *vab-1/EphR* and *glo-4/SERGEF* single mutants. Example of severe hook defects (arrow) observed in *vab-1; glo-4* double mutants. C) Quantitation of axon termination defects for indicated genotypes. *vab-1; glo-4* double mutants show enhanced frequency of both severe hook (black) and moderate overextension (grey) failed termination defects. Overextension defects are significantly reduced by transgenic expression of VAB-1. D) Quantitation of axon termination defects for indicated genotypes. *vab-1; fsn-1* double mutants show enhanced termination defects. E)

998   Quantitation of axon termination defect for indicated genotypes. Axon termination defects are not
999   suppressed in *vab-1; rpm-1* double mutants compared to *rpm-1* single mutants. n is defined as a single
1000   count of 20-30 animals. Means are shown from 8 to 10 counts (20-30 animals per count) for each
1001   genotype, and error bars represent SEM. Significance determined using Student's *t*-test with Bonferroni
1002   correction. ** $p < 0.01$; *** $p < 0.001$; n.s, not significant. Scale bar is 20 μm.
1003
1004
1005

References

AL-ABDI, L., DESBOIS, M., RUSNAC, D. V., SULAIMAN, R. A., ROSENFELD, J. A., LALANI, S., MURDOCK, D. R., BURRAGE, L. C., UNDIAGNOSED DISEASES, N., BILLIE AU, P. Y., TOWNER, S., WILSON, W. G., WONG, L., BRUNET, T., STROBL-WILDEMANN, G., BURTON, J. E., HOGANSON, G., MCWALTER, K., BEGTRUP, A., ZARATE, Y. A., CHRISTENSEN, E. L., OPPERMAN, K. J., GILES, A. C., HELABY, R., KANIA, A., ZHENG, N., GRILL, B. & ALKURAYA, F. S. 2022. Loss-of-function variants in MYCBP2 cause neurobehavioural phenotypes and corpus callosum defects. *Brain*, awac364.

BABETTO, E., BEIROWSKI, B., RUSSLER, E. V., MILBRANDT, J. & DIANTONIO, A. 2013. The Phr1 ubiquitin ligase promotes injury-induced axon self-destruction. *Cell Reports,* 3**,** 1422-1429.

BAKER, S. T. & GRILL, B. 2017. Defining Minimal Binding Regions in Regulator of Presynaptic Morphology 1 (RPM-1) Using Caenorhabditis elegans Neurons Reveals Differential Signaling Complexes. *Journal of Biological Chemistry,* 292**,** 2519-2530.

BAKER, S. T., OPPERMAN, K. J., TULGREN, E. D., TURGEON, S. M., BIENVENUT, W. & GRILL, B. 2014. RPM-1 uses both ubiquitin ligase and phosphatase-based mechanisms to regulate DLK-1 during neuronal development. *PLoS Genet,* 10**,** e1004297.

BANERJEE, S. L., LESSARD, F., CHARTIER, F. J. M., JACQUET, K., OSORNIO-HERNANDEZ, A. I., TEYSSIER, V., GHANI, K., LAVOIE, N., LAVOIE, J. N., CARUSO, M., LAPRISE, P., ELOWE, S., LAMBERT, J. P. & BISSON, N. 2022. EPH receptor tyrosine kinases phosphorylate the PAR-3 scaffold protein to modulate downstream signaling networks. *Cell Rep,* 40**,** 111031.

BINNS, K. L., TAYLOR, P. P., SICHERI, F., PAWSON, T. & HOLLAND, S. J. 2000. Phosphorylation of tyrosine residues in the kinase domain and juxtamembrane region regulates the biological and catalytic activities of Eph receptors. *Mol Cell Biol,* 20**,** 4791-805.

BLOOM, A. J., MILLER, B. R., SANES, J. R. & DIANTONIO, A. 2007. The requirement for Phr1 in CNS axon tract formation reveals the corticostriatal boundary as a choice point for cortical axons. *Genes Dev,* 21**,** 2593-606.

BORGEN, M. A., WANG, D. & GRILL, B. 2017. RPM-1 regulates axon termination by affecting growth cone collapse and microtubule stability. *Development,* 144**,** 4658-4672.

BUSH, J. O. 2022. Cellular and molecular mechanisms of EPH/EPHRIN signaling in evolution and development. *Curr Top Dev Biol,* 149**,** 153-201.

BUSH, J. O. & SORIANO, P. 2010. Ephrin-B1 forward signaling regulates craniofacial morphogenesis by controlling cell proliferation across Eph-ephrin boundaries. *Genes Dev,* 24**,** 2068-80.

CISSÉ, M., HALABISKY, B., HARRIS, J., DEVIDZE, N., DUBAL, D. B., SUN, B., ORR, A., LOTZ, G., KIM, D. H., HAMTO, P., HO, K., YU, G.-Q. & MUCKE, L. 2011. Reversing EphB2 depletion rescues cognitive functions in Alzheimer model. *Nature,* 469**,** 47-52.

COLLINS, C. A., WAIRKAR, Y. P., JOHNSON, S. L. & DIANTONIO, A. 2006. Highwire restrains synaptic growth by attenuating a MAP kinase signal. *Neuron,* 51**,** 57-69.

CRAWLEY, O., OPPERMAN, K. J., DESBOIS, M., ADRADOS, I., BORGEN, M. A., GILES, A. C., DUCKETT, D. R. & GRILL, B. 2019. Autophagy is inhibited by ubiquitin ligase activity in the nervous system. *Nat Commun,* 10**,** 5017.

D'SOUZA, J., HENDRICKS, M., LE GUYADER, S., SUBBURAJU, S., GRUNEWALD, B., SCHOLICH, K. & JESUTHASAN, S. 2005. Formation of the retinotectal projection requires Esrom, an ortholog of PAM (protein associated with Myc). *Development,* 132**,** 247-56.

DALVA, M. B., TAKASU, M. A., LIN, M. Z., SHAMAH, S. M., HU, L., GALE, N. W. & GREENBERG, M. E. 2000. EphB receptors interact with NMDA receptors and regulate excitatory synapse formation. *Cell,* 103**,** 945-56.

DEPAEPE, V., SUAREZ-GONZALEZ, N., DUFOUR, A., PASSANTE, L., GORSKI, J. A., JONES, K. R., LEDENT, C. & VANDERHAEGHEN, P. 2005. Ephrin signalling controls brain size by regulating apoptosis of neural progenitors. *Nature,* 435**,** 1244-50.

DESBOIS, M., CRAWLEY, O., EVANS, P. R., BAKER, S. T., MASUHO, I., YASUDA, R. & GRILL, B. 2018. PAM forms an atypical SCF ubiquitin ligase complex that ubiquitinates and degrades NMNAT2. *J Biol Chem,* 293**,** 13897-13909.

DESBOIS, M., OPPERMAN, K. J., AMEZQUITA, J., GAGLIO, G., CRAWLEY, O. & GRILL, B. 2022. Ubiquitin ligase activity inhibits Cdk5 to control axon termination. *PLoS Genet,* 18**,** e1010152.

FASEN, K., CERRETTI, D. P. & HUYNH-DO, U. 2008. Ligand binding induces Cbl-dependent EphB1 receptor degradation through the lysosomal pathway. *Traffic,* 9**,** 251-66.

FAWAL, M.-A., JUNGAS, T., KISCHEL, A., AUDOUARD, C., IACOVONI, J. S. & DAVY, A. 2018. Cross Talk between One-Carbon Metabolism, Eph Signaling, and Histone Methylation Promotes Neural Stem Cell Differentiation. *Cell Reports,* 23**,** 2864-2873.e7.

FOOT, N., HENSHALL, T. & KUMAR, S. 2017. Ubiquitination and the Regulation of Membrane Proteins. *Physiol Rev,* 97**,** 253-281.

GAGNE, L. M., MORIN, N., LAVOIE, N., BISSON, N., LAMBERT, J. P., MALLETTE, F. A. & HUOT, M. E. 2021. Tyrosine phosphorylation of DEPTOR functions as a molecular switch to activate mTOR signaling. *J Biol Chem,* 297**,** 101291.

GALE, N. W., HOLLAND, S. J., VALENZUELA, D. M., FLENNIKEN, A., PAN, L., RYAN, T. E., HENKEMEYER, M., STREBHARDT, K., HIRAI, H., WILKINSON, D. G., PAWSON, T., DAVIS, S. & YANCOPOULOS, G. D. 1996. Eph receptors and ligands comprise two major specificity subclasses and are reciprocally compartmentalized during embryogenesis. *Neuron,* 17**,** 9-19.

GENANDER, M., HALFORD, M. M., XU, N.-J., ERIKSSON, M., YU, Z., QIU, Z., MARTLING, A., GREICIUS, G., THAKAR, S., CATCHPOLE, T., CHUMLEY, M. J., ZDUNEK, S., WANG, C., HOLM, T., GOFF, S. P., PETTERSSON, S., PESTELL, R. G., HENKEMEYER, M. & FRISÉN, J. 2009. Dissociation of EphB2 signaling pathways mediating progenitor cell proliferation and tumor suppression. *Cell,* 139**,** 679-692.

GEORGE, S. E., SIMOKAT, K., HARDIN, J. & CHISHOLM, A. D. 1998. The VAB-1 Eph receptor tyrosine kinase functions in neural and epithelial morphogenesis in C. elegans. *Cell,* 92**,** 633-43.

GRILL, B., BIENVENUT, W. V., BROWN, H. M., ACKLEY, B. D., QUADRONI, M. & JIN, Y. 2007. C. elegans RPM-1 regulates axon termination and synaptogenesis through the Rab GEF GLO-4 and the Rab GTPase GLO-1. *Neuron,* 55**,** 587-601.

GRILL, B., CHEN, L., TULGREN, E. D., BAKER, S. T., BIENVENUT, W., ANDERSON, M., QUADRONI, M., JIN, Y. & GARNER, C. C. 2012. RAE-1, a novel PHR binding protein, is required for axon termination and synapse formation in Caenorhabditis elegans. *J Neurosci,* 32**,** 2628-36.

GRILL, B., MURPHEY, R. K. & BORGEN, M. A. 2016. The PHR proteins: intracellular signaling hubs in neuronal development and axon degeneration. *Neural development,* 11**,** 8-18.

GUO, Q., XIE, J., DANG, C. V., LIU, E. T. & BISHOP, J. M. 1998. Identification of a large Myc-binding protein that contains RCC1-like repeats. *Proc Natl Acad Sci U S A,* 95**,** 9172-7.

HAGLUND, K. & DIKIC, I. 2012. The role of ubiquitylation in receptor endocytosis and endosomal sorting. *J Cell Sci,* 125**,** 265-75.

HAMBURGER, V. & HAMILTON, H. L. 1951. A series of normal stages in the development of the chick embryo. *J Morphol,* 88**,** 49-92.

HAN, S., KIM, S., BAHL, S., LI, L., BURANDE, C. F., SMITH, N., JAMES, M., BEAUCHAMP, R. L., BHIDE, P., DIANTONIO, A. & RAMESH, V. 2012. The E3 ubiquitin ligase protein associated with Myc (Pam) regulates mammalian/mechanistic target of rapamycin complex 1 (mTORC1) signaling in vivo through N- and C-terminal domains. *J Biol Chem,* 287**,** 30063-72.

HENKEMEYER, M., ORIOLI, D., HENDERSON, J. T., SAXTON, T. M., RODER, J., PAWSON, T. & KLEIN, R. 1996. Nuk controls pathfinding of commissural axons in the mammalian central nervous system. *Cell,* 86**,** 35-46.

HUTTLIN, E. L., BRUCKNER, R. J., NAVARRETE-PEREA, J., CANNON, J. R., BALTIER, K., GEBREAB, F., GYGI, M. P., THORNOCK, A., ZARRAGA, G., TAM, S., SZPYT, J., GASSAWAY, B. M., PANOV, A., PARZEN, H., FU, S., GOLBAZI, A., MAENPAA, E., STRICKER, K., GUHA THAKURTA, S., ZHANG, T., RAD, R., PAN, J., NUSINOW, D. P., PAULO, J. A., SCHWEPPE, D. K., VAITES, L. P., HARPER, J. W. & GYGI, S. P. 2021. Dual proteome-scale networks reveal cell-specific remodeling of the human interactome. *Cell,* 184**,** 3022-3040 e28.

KANIA, A. & KLEIN, R. 2016. Mechanisms of ephrin-Eph signalling in development, physiology and disease.
Nature reviews Molecular cell biology, 17**,** 240-256.

KNIGHT, J. D. R., CHOI, H., GUPTA, G. D., PELLETIER, L., RAUGHT, B., NESVIZHSKII, A. I. & GINGRAS, A. C.
2017. ProHits-viz: a suite of web tools for visualizing interaction proteomics data. *Nat Methods,*
14**,** 645-646.

KULLANDER, K., MATHER, N. K., DIELLA, F., DOTTORI, M., BOYD, A. W. & KLEIN, R. 2001. Kinase-
dependent and kinase-independent functions of EphA4 receptors in major axon tract formation
in vivo. *Neuron,* 29**,** 73-84.

LACKMANN, M., OATES, A. C., DOTTORI, M., SMITH, F. M., DO, C., POWER, M., KRAVETS, L. & BOYD, A.
W. 1998. Distinct subdomains of the EphA3 receptor mediate ligand binding and receptor
dimerization. *J Biol Chem,* 273**,** 20228-37.

LAHAIE, S., MORALES, D., BAGCI, H., HAMOUD, N., CASTONGUAY, C.-E., KAZAN, J. M., DESROCHERS, G.,
KLAR, A., GINGRAS, A.-C., PAUSE, A., CÔTÉ, J.-F. & KANIA, A. 2019. The endosomal sorting adaptor
HD-PTP is required for ephrin-B:EphB signalling in cellular collapse and spinal motor axon
guidance. *Scientific reports,* 9**,** 11945.

LEWCOCK, J. W., GENOUD, N., LETTIERI, K. & PFAFF, S. L. 2007. The Ubiquitin Ligase Phr1 Regulates Axon
Outgrowth through Modulation of Microtubule Dynamics. *Neuron,* 56**,** 604-620.

LIAO, E. H., HUNG, W., ABRAMS, B. & ZHEN, M. 2004. An SCF-like ubiquitin ligase complex that controls
presynaptic differentiation. *Nature,* 430**,** 345-350.

LITTERST, C., GEORGAKOPOULOS, A., SHIOI, J., GHERSI, E., WISNIEWSKI, T., WANG, R., LUDWIG, A. &
ROBAKIS, N. K. 2007. Ligand binding and calcium influx induce distinct ectodomain/gamma-
secretase-processing pathways of EphB2 receptor. *J Biol Chem,* 282**,** 16155-63.

LIU, G., ZHANG, J., LARSEN, B., STARK, C., BREITKREUTZ, A., LIN, Z. Y., BREITKREUTZ, B. J., DING, Y.,
COLWILL, K., PASCULESCU, A., PAWSON, T., WRANA, J. L., NESVIZHSKII, A. I., RAUGHT, B., TYERS,
M. & GINGRAS, A. C. 2010. ProHits: integrated software for mass spectrometry-based interaction
proteomics. *Nat Biotechnol,* 28**,** 1015-7.

LURIA, V., KRAWCHUK, D., JESSELL, T. M., LAUFER, E. & KANIA, A. 2008. Specification of motor axon
trajectory by ephrin-B:EphB signaling: symmetrical control of axonal patterning in the developing
limb. *Neuron,* 60**,** 1039-53.

MARGOLIS, S. S., SALOGIANNIS, J., LIPTON, D. M., MANDEL-BREHM, C., WILLS, Z. P., MARDINLY, A. R.,
HU, L., GREER, P. L., BIKOFF, J. B., HO, H. Y., SOSKIS, M. J., SAHIN, M. & GREENBERG, M. E. 2010.
EphB-mediated degradation of the RhoA GEF Ephexin5 relieves a developmental brake on
excitatory synapse formation. *Cell,* 143**,** 442-55.

MELLITZER, G., XU, Q. & WILKINSON, D. G. 1999. Eph receptors and ephrins restrict cell intermingling and communication. *Nature,* 400**,** 77-81.

MOHAMED, A. M. & CHIN-SANG, I. D. 2006. Characterization of loss-of-function and gain-of-function Eph receptor tyrosine kinase signaling in C. elegans axon targeting and cell migration. *Dev Biol,* 290**,** 164-76.

NA, Y., CALVO-JIMENEZ, E., KON, E., CAO, H., JOSSIN, Y. & COOPER, J. A. 2020. Fbxo45 Binds SPRY Motifs in the Extracellular Domain of N-Cadherin and Regulates Neuron Migration during Brain Development. *Mol Cell Biol,* 40.

NAKATA, K., ABRAMS, B., GRILL, B., GONCHAROV, A., HUANG, X., CHISHOLM, A. D. & JIN, Y. 2005. Regulation of a DLK-1 and p38 MAP kinase pathway by the ubiquitin ligase RPM-1 is required for presynaptic development. *Cell,* 120**,** 407-20.

NIE, D., DI NARDO, A., HAN, J. M., BAHARANYI, H., KRAMVIS, I., HUYNH, T., DABORA, S., CODELUPPI, S., PANDOLFI, P. P., PASQUALE, E. B. & SAHIN, M. 2010. Tsc2-Rheb signaling regulates EphA-mediated axon guidance. *Nature Neuroscience,* 13**,** 163-172.

OKUMURA, F., JOO-OKUMURA, A., OBARA, K., PETERSEN, A., NISHIKIMI, A., FUKUI, Y., NAKATSUKASA, K. & KAMURA, T. 2017. Ubiquitin ligase SPSB4 diminishes cell repulsive responses mediated by EphB2. *Molecular biology of the cell,* 28**,** 3532-3541.

PAO, K. C., WOOD, N. T., KNEBEL, A., RAFIE, K., STANLEY, M., MABBITT, P. D., SUNDARAMOORTHY, R., HOFMANN, K., VAN AALTEN, D. M. F. & VIRDEE, S. 2018. Activity-based E3 ligase profiling uncovers an E3 ligase with esterification activity. *Nature,* 556**,** 381-385.

PEREZ-RIVEROL, Y., CSORDAS, A., BAI, J., BERNAL-LLINARES, M., HEWAPATHIRANA, S., KUNDU, D. J., INUGANTI, A., GRISS, J., MAYER, G., EISENACHER, M., PEREZ, E., USZKOREIT, J., PFEUFFER, J., SACHSENBERG, T., YILMAZ, S., TIWARY, S., COX, J., AUDAIN, E., WALZER, M., JARNUCZAK, A. F., TERNENT, T., BRAZMA, A. & VIZCAINO, J. A. 2019. The PRIDE database and related tools and resources in 2019: improving support for quantification data. *Nucleic Acids Res,* 47**,** D442-D450.

POLIAK, S., MORALES, D., CROTEAU, L.-P., KRAWCHUK, D., PALMESINO, E., MORTON, S., JEAN-FRANÇOIS, C., CHARRON, F., DALVA, M. B., ACKERMAN, S. L., KAO, T.-J. & KANIA, A. 2015. Synergistic integration of Netrin and ephrin axon guidance signals by spinal motor neurons. *eLife,* 4**,** e10841.

POLIAKOV, A., COTRINA, M. L., PASINI, A. & WILKINSON, D. G. 2008. Regulation of EphB2 activation and cell repulsion by feedback control of the MAPK pathway. *The Journal of cell biology,* 183**,** 933-947.

SABET, O., STOCKERT, R., XOURI, G., BRUGGEMANN, Y., STANOEV, A. & BASTIAENS, P. I. H. 2015. Ubiquitination switches EphA2 vesicular traffic from a continuous safeguard to a finite signalling mode. *Nat Commun,* 6**,** 8047.

SAIGA, T., FUKUDA, T., MATSUMOTO, M., TADA, H., OKANO, H. J., OKANO, H. & NAKAYAMA, K. I. 2009.
Fbxo45 forms a novel ubiquitin ligase complex and is required for neuronal development.
Molecular and Cellular Biology, 29**,** 3529-3543.

SALOKAS, K., LIU, X., OHMAN, T., CHOWDHURY, I., GAWRIYSKI, L., KESKITALO, S. & VARJOSALO, M. 2022.
Physical and functional interactome atlas of human receptor tyrosine kinases. *EMBO Rep,* 23**,**
e54041.

SCHAEFER, A. M., HADWIGER, G. D. & NONET, M. L. 2000. rpm-1, a conserved neuronal gene that
regulates targeting and synaptogenesis in C. elegans. *Neuron,* 26**,** 345-56.

SCHAUPP, A., SABET, O., DUDANOVA, I., PONSERRE, M., BASTIAENS, P. & KLEIN, R. 2014. The composition
of EphB2 clusters determines the strength in the cellular repulsion response. *J Cell Biol,* 204**,** 409-
22.

SCHINDELIN, J., ARGANDA-CARRERAS, I., FRISE, E., KAYNIG, V., LONGAIR, M., PIETZSCH, T., PREIBISCH, S.,
RUEDEN, C., SAALFELD, S., SCHMID, B., TINEVEZ, J. Y., WHITE, D. J., HARTENSTEIN, V., ELICEIRI, K.,
TOMANCAK, P. & CARDONA, A. 2012. Fiji: an open-source platform for biological-image analysis.
Nat Methods, 9**,** 676-82.

SHARFE, N., FREYWALD, A., TORO, A. & ROIFMAN, C. M. 2003. Ephrin-A1 induces c-Cbl phosphorylation
and EphA receptor down-regulation in T cells. *J Immunol,* 170**,** 6024-32.

SHARMA, J., BAKER, S. T., TURGEON, S. M., GURNEY, A. M., OPPERMAN, K. J. & GRILL, B. 2014.
Identification of a peptide inhibitor of the RPM-1 . FSN-1 ubiquitin ligase complex. *J Biol Chem,*
289**,** 34654-66.

SHI, L., FU, W. Y., HUNG, K. W., PORCHETTA, C., HALL, C., FU, A. K. & IP, N. Y. 2007. Alpha2-chimaerin
interacts with EphA4 and regulates EphA4-dependent growth cone collapse. *Proc Natl Acad Sci*
U S A, 104**,** 16347-52.

SOSKIS, M. J., HO, H. Y., BLOODGOOD, B. L., ROBICHAUX, M. A., MALIK, A. N., ATAMAN, B., RUBIN, A. A.,
ZIEG, J., ZHANG, C., SHOKAT, K. M., SHARMA, N., COWAN, C. W. & GREENBERG, M. E. 2012. A
chemical genetic approach reveals distinct EphB signaling mechanisms during brain development.
Nat Neurosci, 15**,** 1645-54.

SRIVASTAVA, N., ROBICHAUX, M. A., CHENAUX, G., HENKEMEYER, M. & COWAN, C. W. 2013. EphB2
receptor forward signaling controls cortical growth cone collapse via Nck and Pak. *Mol Cell*
Neurosci, 52**,** 106-16.

TEO, G., LIU, G., ZHANG, J., NESVIZHSKII, A. I., GINGRAS, A. C. & CHOI, H. 2014. SAINTexpress:
improvements and additional features in Significance Analysis of INTeractome software. *J*
Proteomics, 100**,** 37-43.

TULGREN, E. D., TURGEON, S. M., OPPERMAN, K. J. & GRILL, B. 2014. The Nesprin family member ANC-1
regulates synapse formation and axon termination by functioning in a pathway with RPM-1 and
beta-Catenin. *PLoS Genet,* 10**,** e1004481.

VENKITACHALAM, S., BABU, D., RAVILLAH, D., KATABATHULA, R. M., JOSEPH, P., SINGH, S.,
UDHAYAKUMAR, B., MIAO, Y., MARTINEZ-URIBE, O., HOGUE, J. A., KRESAK, A. M., DAWSON, D.,
LAFRAMBOISE, T., WILLIS, J. E., CHAK, A., GARMAN, K. S., BLUM, A. E., VARADAN, V. & GUDA, K.
2022. The Ephrin B2 Receptor Tyrosine Kinase Is a Regulator of Proto-oncogene MYC and
Molecular Programs Central to Barrett's Neoplasia. *Gastroenterology,* 163**,** 1228-1241.

VIRDEE, S. 2022. An atypical ubiquitin ligase at the heart of neural development and programmed axon
degeneration. *Neural Regen Res,* 17**,** 2347-2350.

WALKER-DANIELS, J., RIESE, D. J., 2ND & KINCH, M. S. 2002. c-Cbl-dependent EphA2 protein degradation
is induced by ligand binding. *Mol Cancer Res,* 1**,** 79-87.

WAN, H. I., DIANTONIO, A., FETTER, R. D., BERGSTROM, K., STRAUSS, R. & GOODMAN, C. S. 2000.
Highwire regulates synaptic growth in Drosophila. *Neuron,* 26**,** 313-29.

WILLIAMS, S. E., MANN, F., ERSKINE, L., SAKURAI, T., WEI, S., ROSSI, D. J., GALE, N. W., HOLT, C. E., MASON,
C. A. & HENKEMEYER, M. 2003. Ephrin-B2 and EphB1 mediate retinal axon divergence at the optic
chiasm. *Neuron,* 39**,** 919-35.

XIONG, X., HAO, Y., SUN, K., LI, J., LI, X., MISHRA, B., SOPPINA, P., WU, C., HUME, R. I. & COLLINS, C. A.
2012. The Highwire ubiquitin ligase promotes axonal degeneration by tuning levels of Nmnat
protein. *PLoS Biol,* 10**,** e1001440.

ZHEN, M., HUANG, X., BAMBER, B. & JIN, Y. 2000. Regulation of presynaptic terminal organization by C.
elegans RPM-1, a putative guanine nucleotide exchanger with a RING-H2 finger domain. *Neuron,*
26**,** 331-43.

ZIMMER, M., PALMER, A., KÖHLER, J. & KLEIN, R. 2003. EphB-ephrinB bi-directional endocytosis
terminates adhesion allowing contact mediated repulsion. *Nature Cell Biology,* 5**,** 869-878.
1233
1234