1   A nucleoid-associated protein is involved in the emergence of antibiotic resistance by promoting

2   the frequent exchange of the replicative DNA polymerase in *M. smegmatis*

3   Wei L. Ng [a] and E. Hesper Rego [1][a]

Affiliations

6   [a] Department of Microbial Pathogenesis, Yale University School of Medicine, New Haven, CT

7   06519

8   [1] To whom correspondence should be addressed: hesper.rego@yale.edu

SUMMARY

Antibiotic resistance in *M. tuberculosis* exclusively originates from chromosomal mutations,

either during normal DNA replication or under stress, when the expression of error-prone DNA

polymerases increases to repair damaged DNA. To bypass DNA lesions and catalyze error-

prone DNA synthesis, translesion polymerases must be able to access the DNA, temporarily

replacing the high-fidelity replicative polymerase. The mechanisms that govern polymerase

exchange are not well understood, especially in mycobacteria. Here, using a suite of

quantitative fluorescence imaging techniques, we discover that, as in other bacterial species, in

M. smegmatis, the replicative polymerase, DnaE1, exchanges at a timescale much faster than

that of DNA replication. Interestingly, this fast exchange rate depends on an actinobacteria-

specific nucleoid-associated protein (NAP), Lsr2. In cells missing *lsr2*, DnaE1 exchanges less

frequently, and the chromosome is replicated more faithfully. Additionally, in conditions that

damage DNA, cells lacking *lsr2* load the complex needed to bypass DNA lesions less effectively

and, consistently, replicate with higher fidelity but exhibit growth defects. Together, our results

show that Lsr2 promotes dynamic flexibility of the mycobacterial replisome, which is critical for

robust cell growth and lesion repair in conditions that damage DNA.

Importance. Unlike many other pathogens, *M. tuberculosis* has limited ability for horizontal

gene transfer, a major mechanism for developing antibiotic resistance. Thus, the mechanisms

that facilitate chromosomal mutagenesis are of particular importance in mycobacteria. Here, we

show that Lsr2, a nucleoid-associated protein, has a novel role in DNA replication and

mutagenesis in the model mycobacterium *M. smegmatis*. We find that Lsr2 promotes the fast

exchange rate of the replicative DNA polymerase, DnaE1, at the replication fork and is

important for the effective loading of the DnaE2-ImuA'-ImuB translesion complex. Without *lsr2*,

M. smegmatis replicates its chromosome more faithfully and acquires resistance to rifampin at a

lower rate, but at the cost of impaired survival to DNA damaging agents. Together, our work

establishes Lsr2 as a potential factor in the emergence of mycobacterial antibiotic resistance.

INTRODUCTION

DNA replication and repair are fundamental to bacterial evolution. The function of DNA

replication and repair pathways are key determinants of the balance between genetic stability

and mutability. To establish this balance, bacteria encode and regulate numerous DNA

polymerases with specialized functions in different conditions (1). Importantly, the emergence

of antibiotic resistance in the major human pathogen *M. tuberculosis* is mainly driven by

chromosomal mutagenesis, as *M. tuberculosis* has no ability for horizontal gene transfer (2, 3).

Thus, the mechanisms regulating polymerase expression and activity are of particular

importance for mycobacterial evolution.

In mycobacteria, the fidelity of chromosome replication is established by the major

replicative polymerase DnaE1, which possesses endogenous proofreading activity (4). In

addition to DnaE1, mycobacteria encode a suite of other polymerases with specialized repair

functions that are only expressed under certain conditions, such as oxidative or DNA-damaging

stress (5-7). For example, the mycobacterial specialist polymerases DinB1, DinB2, DinB3, and

DnaE2 are induced under different conditions and have varying repair activities and mutational

signatures (5-10). Of particular importance is the error-prone polymerase DnaE2. It and its

accessory proteins ImuA' and ImuB – collectively referred to as the 'mutasome' – are highly

expressed in conditions that damage DNA and are the primary drivers of stress-induced

mutagenesis *in vitro* and potentially *in vivo* (5, 6, 11). In general, however, the mechanisms that

facilitate switching between polymerases at the site of repair are not well understood, especially

in mycobacteria (12).

It is known that in non-DNA damaging conditions, in *E. coli*, components within the large

multi-protein complex that replicates DNA (the replisome) exchange rapidly, likely when

encountering DNA-protein obstacles. More specifically, fluorescence recovery after

photobleaching (FRAP) and single molecule studies have revealed that, despite being highly

processive *in vitro*, replisome components are highly dynamic within the cell (13). For example,

at room temperature, the alpha subunit of Pol III in *E. coli* exchanges with the cytoplasmic pool

at a time scale of approximately 4 seconds, suggesting that subunits are constantly being

replaced during the 150 minutes it takes to replicate the chromosome at this temperature (13).

These data underlie the notion that the replisome is dynamic and flexible, allowing new subunits

the opportunity to exchange frequently and suggest that protein obstacles influence the

exchange rate of replisome components.

Mycobacteria differ from model bacteria like *E. coli* in several ways. Most notable is their

slow growth and long doubling times. Pathogenic mycobacteria, like *M. tuberculosis* and *M.*

leprae, are notorious for their long doubling times, which can vary from 18 hours to several

weeks. Even so-called fast growers like *M. smegmatis* double on the hour-long timescale

compared to the minute-long timescale of commonly studied organisms like *E. coli*. Intriguingly,

the duration of chromosomal replication at least partially scales with growth rate: *M. smegmatis*

exhibits a longer duration of chromosome replication than *E. coli* (~100-150 minutes vs ~41

minutes, respectively) despite the ability of DnaE1 to polymerize DNA at a faster rate than *E.*

coli Pol IIIα *in vitro* (4, 14, 15).

These observations led us to wonder if components within the replisome exchange at a

different timescale in mycobacteria than in faster-growing bacteria. Using various fluorescence

imaging techniques, we find that, as in *E. coli*, the replicative polymerase is exchanged rapidly

in *M. smegmatis*. Surprisingly, this fast exchange rate is, in part, mediated by an actinobacteria-

specific DNA-binding protein Lsr2, as DnaE1 is exchanged more slowly in cells missing *lsr2*.

Interestingly, when the DNA is damaged, ImuB, a key scaffolding protein that recruits the error-

prone polymerase into an active complex, is exchanged more rapidly, suggesting ineffective

loading of the mutasome. Consistently, loss of *lsr2* results in growth defects in conditions that

damage DNA, and reduced mutagenesis. Overall, our data deepen our understanding of DNA

metabolism in mycobacteria and provide another unexpected layer to the mechanisms

mycobacteria use to mutate and acquire drug resistance.

RESULTS

The mycobacterial replisome is highly dynamic

To investigate the dynamics of DnaE1, we created a strain expressing DnaE1-mScarlet

from its native promoter. A similar fusion encodes functional DnaE1 (16). To verify that DnaE1-

mScarlet reports on the localization of the replisome, we performed time-lapse microscopy and

analyzed fluorescence distributions as a function of both cell length and cell cycle (**Fig. 1A, B**).

Figure 1. DnaE1 localization over time in *M. smegmatis*. (A) A representative cell expressing DnaE1-mScarlet from its native promoter is imaged over time with both phase and fluorescence microscopy. Fluorescent spots appear at mid-cell, disappear, and then re-appear at roughly one- and three-quarters positions at the end of one cell cycle/beginning of another. **(B)** To quantify fluorescence probability distributions over time and space, we constructed kymographs by averaging individual distributions that were normalized to cell cycle time and length (N=43).

Consistent with the reported placement of the replisome in mycobacteria, DnaE1 localizes at the

approximately mid-cell before disappearing, presumably after DNA replication has been

completed (16). Towards the end of the cell cycle, DnaE1 reappears at one- and three-quarter

positions to initiate replication in newly synthesized sister chromosomes (**Fig. 1A, B**). These

data show that DnaE1-mScarlet is reporting on the localization of the replisome in *M.*

smegmatis.

To test the exchange rate of DnaE1, we took the same approach as Beattie, TR *et al*

and used FRAP to measure the dynamics of recovery of DnaE1 in cells (13). Fast recovery of

fluorophores into a bleached spot indicates fast dissociation of bleached molecules. Conversely,

slow, or incomplete recovery of non-bleached fluorophores indicates slow dissociation of

bleached molecules, and a stably associated complex. To account for differences in the degree

of bleaching per cell, we normalized the fluorescence intensity after bleaching to zero, and the

final intensity to the maximum recovery we expected based on the percentage of fluorophores

bleached. We fit the data to a diffusion exchange reaction and calculated both the timescale of

recovery, reflecting the dissociation time of bound subunits, and the maximum recovery,

reflecting the mobile fraction of proteins. Imaging was done in the same conditions as above, in

which we measured the duration of chromosomal replication to be approximately 100 minutes

Figure 2. The replicative polymerase is highly dynamic, which is in part mediated by Lsr2. **(A)** Representative time sequence of FRAP of DnaE1-mScarlet in WT (top) and Δ*lsr2* (bottom) imaged at 2 and 4-second intervals, respectively. Blue arrow indicates the location of the focus targeted for bleaching. **(B)** Fluorescence recovery curves for DnaE1-mScarlet (N=21 for WT, 12 for Δ*lsr2*). Solid lines show a one-phase exponential association fit to the data. Individual points show the mean±SEM of fluorescence intensity after bleach correction. Dashed line represents the estimated maximum possible fluorescence recovery. **(C)** DnaE1-mScarlet bound times (mean±SD) obtained from one-phase exponential association fits in panel B for both WT and Δ*lsr2* for DnaE1. (****), $p<0.0001$ by unpaired t-test. **(D)** Frequency of spontaneous rifampin-resistant mutants per 10^8 cfu. LOD refers to the limit of detection. *lsr2* complements were expressed from the native *lsr2* promoter on a phage integrative plasmid. p-values were obtained by one-way ANOVA. (*), $p<0.0021$ (**), $p<0.0002$.

based on the time between the appearance and disappearance of DnaE1 (**Fig. 1B, S2B**). We

found that, as in *E. coli*, DnaE1 exchanged quickly and nearly completely with a timescale of

approximately 8 +/- 3 seconds (mean±SD) (**Fig. 2A-C**). These data are consistent with the

model that has emerged in *E. coli,* which suggests that the replisome constantly encounters

obstacles that promote the exchange of individual subunits with the cytoplasmic pool.

Lsr2 promotes the frequent exchange of DnaE1

In bacteria, nucleoid-associated proteins (NAPS) are important for bacterial

chromosome organization and influence replication, transcription, and cell cycle regulation

either directly or indirectly. Lsr2 is one of the most well-studied NAPs in mycobacteria. Lsr2 is

conserved across mycobacteria and related actinomycetes (17-21). Like H-NS in *E. coli*, Lsr2

primarily binds AT-rich regions of DNA and functions as a global transcriptional repressor (18).

Notably, *in vitro* studies show that Lsr2 binding interferes with transcription, inhibits

topoisomerase I-dependent supercoil relaxation, and protects DNA from DNase I digestion and

H_2O_2-mediated degradation (17). Loss of *lsr2* is associated with a wide range of different

phenotypes in mycobacteria, including altered cell cycle timing and growth defects upon

entering/exiting hypoxic conditions and when exposed to oxidative stress and various antibiotics

(17, 22-25). Lsr2 is also involved in resistance against phage infection, protection against

reactive oxygen intermediates during macrophage infection, and cells missing *lsr2* show

reduced virulence in various hosts, including mice (20, 22, 23, 26).

ChIP-seq data show an abundance of Lsr2 binding sites across the chromosome, and

consistent with this, Lsr2 decorates the entire nucleoid in discrete and highly dynamic puncta

(24). Interestingly, the major Lsr2 foci are localized near, but not co-localized with, the

replisome. We verified this localization pattern in our hands by creating strains that express

Lsr2-mScarlet and DnaE1-mGFPmut3. Consistent with prior reports, Lsr2 and DnaE1 exhibit

distinct localizations, and although Lsr2 localizes nearby DnaE1 at certain times in the cell

cycle, Lsr2 does not co-localize with DnaE1 despite Lsr2's abundance throughout nucleoid (**Fig.**

S1).

Given the importance of Lsr2 to mycobacteria physiology and its localization, we

wondered if deletion of *lsr2* would influence replisome exchange dynamics. To that end, we

generated an *lsr2* deletion strain expressing DnaE1-mScarlet. Consistent with prior findings

using a DnaN reporter, we find that DnaE1 fluorescence disappears from the mid-cell region

faster in cells missing *lsr2*, suggesting that DNA replication is faster by approximately 20% (**Fig.**

S2A, B) (25). Next, as before, we measured the rate of DnaE1 recovery in Δ*lsr2* cells using

FRAP (**Fig. 2A**). We observed a much longer recovery time (38 +/- 10 seconds) compared to

wild type cells (**Fig. 2B,C**). These data suggest that DnaE1 remains associated with the

replisome for longer in cells missing *lsr2*. Alternatively, a longer recovery time in the *lsr2* mutant

could be due to reduced DnaE1 access to the replisome (e.g*.,* due to increased access by other

polymerases to the replisome). However, reduced access by DnaE1, an essential polymerase in

M. smegmatis, would likely result in a growth rate difference in the mutant, which we and others

do not observe (**Fig. S2C**) (25). Likewise, reduced access by DnaE1 would also increase basal

mutation rates, as this polymerase is responsible for faithful chromosome replication in standard

laboratory medium (4). Instead, we observe that that Δ*lsr2* cells mutate at decreased frequency

as measured by spontaneous resistance to rifampin (**Fig. 2D**), further supporting the notion of

increased stability of DnaE1 in Δ*lsr2.* Taken together, these data show that Lsr2 promotes the

rapid exchange of DnaE1. In its absence, DnaE1 is more stably associated, and cells replicate

their chromosome more quickly and more faithfully.

Lrs2 is important for the acquisition of mutations by promoting the stable formation of

the mutasome

We next considered if Lsr2 also affects polymerase exchange in conditions that elevate

the expression of the error-prone polymerase DnaE2 and its associated complex. To test this,

we created strains carrying fluorescent proteins fusions to ImuA', ImuB, and DnaE2 integrated

either at a phage site (*imuA'* and *imuB* in Δ*imuAB*) or the chromosomal locus (*dnae2*). When

exposed to UV light, these strains acquired resistance to rifampin at a similar rate as wild type,

showing that the fusion proteins retain the ability to repair damaged DNA (**Fig. S3**) (27).

ImuB recruits ImuA' and DnaE2 into an active complex by binding to the β-clamp subunit

of the replisome. ImuA' and DnaE2 bind ImuB but not the β-clamp, and therefore require ImuB

to associate with the replisome and catalyze mutations (8, 11, 27, 28). Despite being co-

transcribed, ImuA'-mScarlet and mScarlet-ImuB had dramatically different fluorescence

intensities: ImuA'-mScarlet was barely detectable above background fluorescence levels, while

mScarlet-ImuB was brightly fluorescent (**Fig. S4A,B**). Thus, cellular levels of ImuA' may be

much lower than ImuB, implying that additional post-transcriptional mechanisms exist (e.g.

translation regulation or proteolysis) to limit ImuA' abundance. Alternatively, mScarlet may be

cleaved off the ImuA'-mScarlet translational fusion and targeted for degradation. However,

consistent with the low abundance of ImuA', we also observed very low fluorescence of DnaE2-

mGFPmut3 (**Fig. S4C**). Others have reported similar results using N-terminal fusions (27).

These data suggest that as in *E. coli,* the translesion polymerase abundance may be tightly

regulated post-transcriptionally (29). Further, these data suggest that one limiting factor in

forming an active mutasome complex may be the low cellular concentrations of ImuA' and/or

DnaE2.

Figure 3. Lsr2 mediates stable loading of ImuB. Cells expressing mScalet-ImuB were treated with 1.56ng/ml mitomycin C for 4 hours before forming FRAP. **(A)** Fluorescence recovery curves for mScarlet-ImuB (N=13 for WT, 10 for Δ*lsr2*). Solid lines show a one-phase exponential association fit to the data. Individual points show the mean±SEM of fluorescence intensity after bleach correction. Dashed line represents the estimated maximum possible fluorescence recovery. **(B)** Bound times (mean±SD) obtained from one-phase exponential association fits from panel A. p<0.0001 by unpaired t-test.

Due to the dim fluorescence of ImuA' and DnaE2, it was difficult to draw conclusions

about their localization or dynamics in cells with or without *lsr2*. Thus, we decided to determine

the exchange rate of ImuB. We treated cells with a low concentration of mitomycin (MMC) to

induce expression of ImuB and, as before, performed FRAP. In wild type cells, ImuB displayed

a much longer timescale of exchange (74 +/- 14 seconds) compared to DnaE1, suggesting

ImuB remains associated with the β-clamp for a longer time (**Fig. 3A,B**). The long bound time of

ImuB is consistent with the need to recruit low-abundance proteins ImuA' and/or DnaE2. In cells

missing *lsr2*, ImuB exchanged more quickly (46 +/- 6 seconds) than in wild type, at a rate similar

to DnaE1-mScarlet (**Fig. 2C**). Thus, as ImuB and DnaE1 both bind the β -clamp, these data

suggest that the mutasome complex is less stably associated in cells missing *lsr2*, possibly due

to the increased stability of DnaE1.

To further test this hypothesis, we reasoned that Δ*lsr2* would phenocopy cells defective

in mutasome expression or stability. *M. smegmatis* and *M. tuberculosis* missing ImuA' and ImuB

are more sensitive to MMC treatment and are less likely to mutate when exposed to DNA-

damaging agents like ultraviolet light (**Fig. S3A**) (11, 27). Thus, we hypothesized that cells

missing *lsr2* would be altered in these key phenotypes. We first confirmed by RNA-seq that the

FIGURE 4. Lsr2 mediates survival to mitomycin and UV-induced mutagenesis. (A) Serial dilutions of WT, Δ*lsr2*, and complemented strains on agar plates with or without mitomycin C (1.56 ng/ml). Results shown are representative spots of biological triplicates (**Fig. S5**). **(D)** Frequency of rifampin-resistant mutants per 10^8 cfu with UV-induced DNA damage (20 mJ/cm^2). LOD refers to the limit of detection. *lsr2* complements were expressed from the native *lsr2* promoter on a phage integrative plasmid. p-values were obtained by one-way ANOVA (****), p<0.0001.

expression of mutasome components or other genes related to the DNA-damage response was

unaltered in Δ*lsr2* (**Supplementary Tables 1-9**). Then, to test our model, we grew cells on agar

plates with different concentrations of MMC. Deletion of *lsr2* results in a severe growth defect in

the presence of MMC, which could be complemented by the expression of *lsr2* from *M.*

smegmatis or *M. tuberculosis* (**Fig. 4A, Fig. S5**). Next, we assayed the ability of Δ*lsr2* to

acquire resistance to rifampicin when exposed to UV. We chose a dose of UV that resulted in

no survival difference between the two genotypes (**Fig. S6**). Consistent with a role in DNA

damage response, we found that cells deleted for *lsr2* are approximately 4-5 times less likely to

acquire resistance to rifampicin when exposed to UV (**Fig. 4B**). Together, our data show that

Lsr2 is involved in the exchange of polymerase at the replication fork in both standard culture

medium and in conditions that damage DNA and up-regulate repair pathways.

DISCUSSION

Lsr2 acts as a transcriptional repressor for many genes on the *M. smegmatis* and *M.*

tuberculosis chromosomes (17, 22-24). Additionally, multiple lines of evidence suggest that Lsr2

has other roles aside from its function as a transcription factor (22, 25, 26). Our results implicate

Lsr2 as a potential factor in mycobacterial mutagenesis and antibiotic resistance (**Fig. 6**).

While Lsr2 binds to and/or alters the transcription of many genes in both *M. smegmatis*

and *M. tuberculosis,* very few phenotypes associated with the deletion mutant have been

connected to altered expression of target genes (17, 22, 24). One of the few exceptions is the

change in colony morphology, which can be attributed to the increased expression of a putative

polyketide synthase (MSMEG_4727) in *M. smegmatis* (24, 30). Why, then, is Lsr2 associated

Loss of Lsr2:

Altered DnaE1 and
mutasome dynamics
(this study)

↓mutation frequency
(this study)

↓chromosome replication duration
Kolodziej (*et al*, 2021)

defect in phage DNA replication
Dulberger (et al, 2023)

↓growth under DNA damaging stress
(this study)

DnaE2/ImuA'/ImuB
mutasome

DnaE1

Lsr2

mutation, other polymerases (DinB1, DinB2, DinB3), DnaN

FIGURE 6. Model of Lsr2 regulation of DNA replication and mutagenesis. Our data suggest that loss of *lsr2* results in a decrease in mutagenesis due to a novel role for Lsr2 in modulating replisome dynamics. Specifically, our data are consistent with a model in which Lsr2 acts as a roadblock on the DNA to promote the exchange of the major replicative polymerase DnaE1 with others in the cytoplasmic pool. In conditions that do not induce DNA damage, these polymerases may include DinB1, which contributes to spontaneous mutagenesis (12). In conditions that damage DNA, the mutasome, which is comprised of DnaE2 and its accessory proteins (ImuA' and ImuB), is induced and needs to exchange with the replicative polymerase to repair lesions (13, 18, 19). Deletion of Lsr2 reduces the flexibility of the replisome needed for sufficient exchange, thereby resulting in cells that are more reliant on DnaE1, a higher fidelity polymerase that is unable to bypass lesions. Thus, cells missing *lsr2* are impaired in their ability to grow in conditions that damage DNA in their ability to acquire mutations.

with so many bacterial behaviors, including bacterial growth, the cell cycle, robust infection,

survival to hydrogen peroxide and antibiotics, survival in low oxygen, and resistance to phage?

Our results indicate that one of Lsr2's functions in the cell is to promote the flexibility of

the mycobacterial replisome, enabling specialist polymerases like the DnaE2 translesion

complex (ImuB/DnaE2/ImuA') to access the replication fork, leading to efficient, but error-prone,

DNA replication and cell growth in conditions that damage DNA. Specifically, our findings

support a model whereby in non-DNA damaging conditions, Lsr2 serves as an obstacle that

promotes the frequent exchange of DnaE1, allowing other polymerases to exchange into the

replisome (**Fig. 6**). This model is consistent with *Δlsr2* having a slightly reduced basal mutation

rate, as Lsr2 could be allowing exchange of polymerases besides DnaE1 at a low frequency.

Likewise, in DNA damaging conditions, out data suggest a model in which Lsr2-mediated

exchange of DnaE1 promotes loading of ImuB and thus DnaE2 and ImuA'. DnaE1 is a more

faithful enzyme but is unable to bypass lesions; thus, cells missing *lsr2* grow more slowly under

conditions that damage DNA but replicate with higher fidelity.

Numerous studies in eukaryotes and other bacteria show that the rate of replication fork

progression is actively regulated, in some cases by machinery that slow down DNA replication,

to ensure coordination amongst DNA replication, DNA repair, and transcription (31-34). We

hypothesize that Lsr2 may function to regulate replication fork progression in mycobacteria, and

that this function may provide a unifying explanation for many of the phenotypes associated with

loss of *lsr2,* including slower DNA replication in the host and other environmental conditions

likely to assault DNA. Likewise, in conditions that do not significantly damage DNA, like those

found in laboratory growth media, loss of *lsr2* leads to faster and higher fidelity chromosomal

replication, possibly explaining the emergence of spontaneous *lsr2* mutants in several

laboratories (30, 35, 36). Additionally, loss of *lsr2* is associated with increased resistance

against infection by certain phages, and fewer and smaller zones of phage DNA replication (26).

We speculate that efficient turnover of the bacterial host DNA replication machinery mediated by

Lsr2 could be required for robust replication of phage DNA and productive infection.

Intriguingly, Lsr2 is phosphorylated by PknB, a plasma membrane-bound eukaryotic

serine/threonine kinase involved in cell growth and division (37). Future research is needed but

we speculate that such regulation could be used to coordinate cell wall synthesis and DNA

synthesis during environmental changes or even during a typical cell cycle.

Lsr2 may be a promising drug target for preventing the emergence of antibiotic

resistance in mycobacteria. Our work adds to the growing body of evidence Lsr2 is at the nexus

of several evolutionary tradeoffs, including fast chromosome replication, mutagenesis,

replisome flexibility, and phage resistance. Indeed, loss of Lsr2 results in severe growth defects

in *M. tuberculosis* and *M. abscessus in vivo* (20, 23). Thus, we speculate that targeting Lsr2

therapeutically may help treat drug-susceptible infections, while preventing the emergence of

drug resistance.

261
MATERIALS AND METHODS

Strains and growth conditions

All mycobacterial strains are derivatives of *Mycobacterium smegmatis* MC2 155. Cells were

grown in Middlebrook 7H9 broth supplemented with 0.05% (v/v) Tween80, ADC, and 0.2% (v/v)

glycerol for liquid culture. LB agar was used as solid media. Liquid cultures were grown at 37 °C

on a rotator. Agar plates were incubated at 37 °C. If applicable, antibiotics were added at the

following final concentrations: hygromycin B (50 ug/mL), kanamycin (25ug/mL), nourseothricin

(20 ug/ml), zeocin (20 ug/mL), and gentamicin (5 ug/ml).

Plasmid DNA for transformation into *M. smegmatis* was prepared from *E. coli* (DH5a,

XL1). If applicable, antibiotics were added at the following final concentrations: hygromycin B

(100 ug/mL), kanamycin (50 ug/mL), and nourseothricin (40 ug/ml). Plasmid constructs were

generated using Gibson cloning and phage-based integrative vectors. Fluorescent protein

fusions were generated using mGFPmut3 and mScarlet-I variants.

Genetic modifications to the native locus were generated as previously described using

the mycobacterial Che9c phage RecET recombination system (38). To construct a knockout, we

designed a dsDNA substrate encoding a zeocin resistance cassette flanked by loxP sites and

~500bp of homology on either end of the locus of interest. The dsDNA substrate was then

electroporated into *M. smegmatis* strains carrying a plasmid encoding isovaleronitrile-inducible

expression of RecET. Colonies were selected by zeocin resistance and screened by PCR using

primers annealing to regions flanking dsDNA insertion. In some cases, the zeocin resistance

cassette was removed by Cre-Lox recombination. Strains and primers are listed in Tables S12

and S13, respectively.

Fluorescence microscopy

All microscopy experiments were performed on a Nikon TI-E inverted wide-field microscope

equipped with the Nikon Perfect Focus System, an ORCA-Flash4.0 Digital CMOS Camera

(Hamamatsu) and an environmental control chamber maintained at 37 °C.

Time-lapse and kymograph analysis: Time-lapse imaging was done using a 60X Oil 1.45NA

Plan Apochromat phase-contrast objective. Bacterial samples were loaded and grown in a B04A

microfluidic plate (CellAsic ONIX, B04A-03-5PK). Phase and fluorescence images were

acquired every 30 minutes using NIS Elements software. Fluorescence images were taken

sequentially using a 470 nm excitation laser and 515/30 emission filter for msfGFP and

mGFPmut3, and a 555 nm excitation laser and a 595/40 emission filter for mScarletI.

Kymograph analysis was performed using a custom pipeline for segmenting and tracking

cells, and extracting fluorescence profiles. Cell segmentation was performed on the phase

image using a U-Net network trained for segmenting mycobacterial cells (39). Semi-automated

cell tracking and extraction of fluorescence profiles from background-subtracted images was

done using a custom MATLAB program and open-source image analysis software Fiji (40).

Average kymographs were generated by 2D interpolation of at least twenty individual

kymographs.

FRAP: FRAP experiments were done using a 60X oil DIC objective. Bacterial cells were imaged

under 7H9 agar pads (1% agar, 0.2% (v/v) glycerol) in 50 mm glass-bottomed dishes.

Fluorescent spots were manually selected for localized bleaching with a focused 405-nm laser.

Two pre-bleach fluorescence images were acquired: one to identify and center a target focus,

and a second for quantification of total pre-bleach fluorescence. Fluorescence recovery was

monitored by acquisition at pre-defined intervals (2 to 10 sec) following photobleaching using an

exposure time of 200 ms.

Cell outlines for determination of total pre-bleach fluorescence were manually drawn

using the brightfield image. Fluorescence images were background subtracted and corrected for

photobleaching. To quantify the fluorescence of bleached foci over time, a circular ROI was

manually drawn around the bleached foci and intensity profiles over time were obtained using

Fiji (40). Fluorescence recovery curves were generated by first normalizing fluorescence

intensity profiles to pre-bleach focus fluorescence. The post-bleach value was then subtracted

from all time points. Maximum recovery was estimated by calculating the total amount of

fluorescence in the cell after photobleaching outside the focus. Finally, all time points were

divided by the estimated maximum recovery value, such that a value of 0 and 1 correspond to

no recovery and maximum recovery, respectively.

RNA-seq of WT _M. smegmatis_ and Δ_lsr2_

WT _M. smegmatis_ and Δ_lsr2_ were grown in triplicate at 37°C to OD600 ~0.7-1.0. For mitomycin

treatment, 1 ml of cells were added to 4 ml of 7H9 containing 100 ng/ml mitomycin C for a final

concentration of 80 ng/ml mitomycin C. Approximately 2-4 ml of cells were harvested at each

timepoint and incubated with 2 volumes of RNAprotect (Qiagen) at room temperature for 5

minutes prior to centrifugation for 5 minutes at 4000xg 4°C. The supernatant was discarded,

and the cells were then immediately used for RNA extraction or stored at -80°C.

RNA extraction was performed using the Qiagen RNeasy Kit (74104). Cells were

resuspended in 500 ul of RLT buffer (Qiagen) with 2-mercaptoethanol and homogenized by

bead beating for 3 minute pulses at maximum speed, three times, with 5 minutes of rest on ice

in-between. The lysate was then centrifuged at maximum speed for 1 minute, and the

supernatant was transferred to a new tube. The remainder of the RNA extraction was performed

using the Qiagen RNeasy Kit according to manufacturer instructions. RNA quality was verified

using a bioanalyzer (Agilent Technologies TapeStation 2200). DNase treatment and ribosomal

RNA depletion were performed at the Microbial Genome Sequencing Center before library

preparation and sequencing.

Mutation frequency analysis

10 ml of bacterial cells were grown to OD600 ~0.7-1.0. 5 ml of cells were UV irradiated (20

mJ/cm2 UV; Stratalinker UV crosslinker), while the remaining 5 ml of cells were used for the

untreated control. Treated and untreated samples were then added to an equal volume of fresh

7H9 media and outgrown for 3 hours at 37°C. 100 ul liquid culture was used for CFU

determination, and the remainder of the culture was plated onto Rif-containing LB agar plates

(200 ug/ml) for mutant detection. All mutation frequency measurements were performed using 8

biological replicates. Rif-resistant colonies were enumerated after 5-7 days of incubation at

37°C. CFU plates were read after about 2-3 days of growth on non-selective media. Mutation

frequencies were calculated by the number of Rif-resistant colonies per ml of culture by the

CFU/ml.

Spot assay

Bacterial cells were grown to OD600 ~0.7-1.0, normalized to the same OD, and then 10-fold

serially diluted. 5 ul of culture were spotted on LB agar containing varying concentrations of

antibiotic. Plates were imaged after incubation at 37°C for 2-3 days. All assays were done in

biological triplicate.

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the lead contact, Dr. Hesper Rego (hesper.rego@yale.edu).

<u>Materials availability</u>

All unique/stable reagents generated in this study are available from the lead contact.

<u>Data availability</u>

RNA-seq sequencing data are deposited to the Short Read Archive (SRA) under project

number: PRJNA1060683

ACKNOWLEDGMENTS

We thank members of the Rego lab for helpful suggestions, and Dr. Eduardo Groisman for

discussions. This work was supported by NIH R01 AI148255 and the Searle and Pew

Foundations to EHR.

AUTHOR CONTRIBUTIONS

WLN and EHR conceived experiments and analyzed data. WLN performed experiments. WLN

and EHR wrote the manuscript. EHR secured funding for the work.

TABLES S1-6. All quantified RNA-seq read counts for WT and Δ*lsr2*, in triplicate, treated with
mitomycin-C at 0, 1, 3, 6, or 24 hours.
*see attached

381
382 **TABLES S7-11.** Differentially expressed genes in WT and Δ*lsr2* treated with mitomycin-C 0, 1,
383 3, 6, or 24 hours
384 *see attached

386   **TABLE S12.** Strain list

STRAIN	GENOTYPE	SOURCE	Additional information
WN788	mc2155Δlsr2::zeoR	This work	*lsr2* deletion strain
WN906	mc2155Δlsr2::zeoR L5::plsr2-mScarletI-FLAG Giles::pimuAB-msfGFP-myc Tweety::pNative-MSMEG6092	This work	*lsr2* deletion strain, complemented with untagged *M. smegmatis* Lsr2
WN874	mc2155Δlsr2::zeoR Giles::pimuAB-msfGFP-myc Tw::pdnaE2-mScarletI-FLAG L5::pNative-Rv3597c	This work	*lsr2* deletion strain, complemented with untagged *M. tuberculosis* Lsr2
WN1384	mc2155 Giles::pNative-dnae1-mScarletI-FLAG	This work	C-terminal DnaE1 fusion in a WT background
WN1385	mc2155Δlsr2::zeoR Giles::pNative-dnae1-mScarletI-FLAG	This work	C-terminal DnaE1 fusion in a Δ*lsr2* background
WN1390	mc2155Δlsr2::zeoR L5::pNative-MSMEG6092-mScarletI-FLAG Giles::pTetO-DnaE1-mGFPmut3-myc	This work	Dual-labeled C-terminal Lsr2 and C-terminal DnaE1 fusions in a Δ*lsr2* background
WN573	mc2155ΔimuAB::loxP	This work	*imuAB* deletion strain
WN1278	mc2155ΔimuAB::loxP L5::pJR962-LexA sgRNA Giles::pNative-imuA_FLAG-mScarletI-imuB-1550DS	This work	Translational fusion of ImuB
WN1280	mc2155Δimuab::loxP Δlsr2::zeoR L5::pJR962-LexA sgRNA Giles::pNative-imuA_FLAG-mScarletI-imuB-1550DS	This work	Translational fusion of ImuB in Δ*lsr2* background
WN1318	mc2155Δimuab::loxP L5::pJR962-LexA sgRNA Giles::pNative-imuA-mScarletI-FLAG_imuB-1550DS	This work	Translational fusion of ImuA'
WN273	mc2155Δdnae2::zeoR	This work	*dnaE2* deletion strain
WN171	mc2155 rpoB::mApple-hygR dnaE2::mGFPmut3-zeoR	This work	Translational fusion of DnaE2

389   **TABLE S11.** Primer list

390   *see attached
391

References

1. Fuchs RP, Fujii S. 2013. Translesion DNA synthesis and mutagenesis in prokaryotes. Cold Spring Harb Perspect Biol 5:a012682.
2. Galagan JE. 2014. Genomic insights into tuberculosis. Nature Reviews Genetics 15:307-320.
3. Boritsch EC, Khanna V, Pawlik A, Honoré N, Navas VH, Ma L, Bouchier C, Seemann T, Supply P, Stinear TP, Brosch R. 2016. Key experimental evidence of chromosomal DNA transfer among selected tuberculosis-causing mycobacteria. Proceedings of the National Academy of Sciences 113:9876-9881.
4. Rock JM, Lang UF, Chase MR, Ford CB, Gerrick ER, Gawande R, Coscolla M, Gagneux S, Fortune SM, Lamers MH. 2015. DNA replication fidelity in Mycobacterium tuberculosis is mediated by an ancestral prokaryotic proofreader. Nat Genet 47:677-81.
5. Dupuy P, Ghosh S, Adefisayo O, Buglino J, Shuman S, Glickman MS. 2022. Distinctive roles of translesion polymerases DinB1 and DnaE2 in diversification of the mycobacterial genome through substitution and frameshift mutagenesis. Nature Communications 13:4493.
6. Boshoff HI, Reed MB, Barry CE, 3rd, Mizrahi V. 2003. DnaE2 polymerase contributes to in vivo survival and the emergence of drug resistance in Mycobacterium tuberculosis. Cell 113:183-93.
7. Dupuy P, Ghosh S, Fay A, Adefisayo O, Gupta R, Shuman S, Glickman MS. 2023. Roles for mycobacterial DinB2 in frameshift and substitution mutagenesis. Elife 12.
8. Kana BD, Abrahams GL, Sung N, Warner DF, Gordhan BG, Machowski EE, Tsenova L, Sacchettini JC, Stoker NG, Kaplan G, Mizrahi V. 2010. Role of the DinB homologs Rv1537 and Rv3056 in Mycobacterium tuberculosis. J Bacteriol 192:2220-7.
9. Ordonez H, Shuman S. 2014. Mycobacterium smegmatis DinB2 misincorporates deoxyribonucleotides and ribonucleotides during templated synthesis and lesion bypass. Nucleic Acids Res 42:12722-34.
10. Ordonez H, Uson ML, Shuman S. 2014. Characterization of three mycobacterial DinB (DNA polymerase IV) paralogs highlights DinB2 as naturally adept at ribonucleotide incorporation. Nucleic Acids Res 42:11056-70.
11. Warner DF, Ndwandwe DE, Abrahams GL, Kana BD, Machowski EE, Venclovas Č, Mizrahi V. 2010. Essential roles for *imuA*′- and *imuB*-encoded accessory factors in DnaE2-dependent mutagenesis in *Mycobacterium tuberculosis*. Proceedings of the National Academy of Sciences 107:13093-13098.
12. Joseph AM, Badrinarayanan A. 2020. Visualizing mutagenic repair: novel insights into bacterial translesion synthesis. FEMS Microbiol Rev 44:572-582.
13. Beattie TR, Kapadia N, Nicolas E, Uphoff S, Wollman AJ, Leake MC, Reyes-Lamothe R. 2017. Frequent exchange of the DNA polymerase during bacterial chromosome replication. Elife 6.
14. Logsdon MM, Ho PY, Papavinasasundaram K, Richardson K, Cokol M, Sassetti CM, Amir A, Aldridge BB. 2017. A Parallel Adder Coordinates Mycobacterial Cell-Cycle Progression and Cell-Size Homeostasis in the Context of Asymmetric Growth and Organization. Curr Biol 27:3367-3374.e7.
15. Helmstetter CE, Cooper S. 1968. DNA synthesis during the division cycle of rapidly growing Escherichia coli Br. Journal of Molecular Biology 31:507-518.

16. Trojanowski D, Kołodziej M, Hołówka J, Müller R, Zakrzewska-Czerwińska J. 2019.
Watching DNA Replication Inhibitors in Action: Exploiting Time-Lapse Microfluidic
Microscopy as a Tool for Target-Drug Interaction Studies in Mycobacterium. Antimicrob
Agents Chemother 63.
17. Colangeli R, Helb D, Vilchèze C, Hazbón MH, Lee CG, Safi H, Sayers B, Sardone I,
Jones MB, Fleischmann RD, Peterson SN, Jacobs WR, Jr., Alland D. 2007.
Transcriptional regulation of multi-drug tolerance and antibiotic-induced responses by
the histone-like protein Lsr2 in M. tuberculosis. PLoS Pathog 3:e87.
18. Gordon BR, Imperial R, Wang L, Navarre WW, Liu J. 2008. Lsr2 of Mycobacterium
represents a novel class of H-NS-like proteins. J Bacteriol 190:7052-9.
19. Gehrke EJ, Zhang X, Pimentel-Elardo SM, Johnson AR, Rees CA, Jones SE, Hindra,
Gehrke SS, Turvey S, Boursalie S, Hill JE, Carlson EE, Nodwell JR, Elliot MA. 2019.
Silencing cryptic specialized metabolism in Streptomyces by the nucleoid-associated
protein Lsr2. eLife 8:e47691.
20. Le Moigne V, Bernut A, Cortès M, Viljoen A, Dupont C, Pawlik A, Gaillard JL, Misguich
F, Crémazy F, Kremer L, Herrmann JL. 2019. Lsr2 Is an Important Determinant of
Intracellular Growth and Virulence in Mycobacterium abscessus. Front Microbiol 10:905.
21. Wiechert J, Filipchyk A, Hünnefeld M, Gätgens C, Brehm J, Heermann R, Frunzke J.
2020. Deciphering the Rules Underlying Xenogeneic Silencing and Counter-Silencing of
Lsr2-like Proteins Using CgpS of Corynebacterium glutamicum as a Model. mBio 11.
22. Colangeli R, Haq A, Arcus VL, Summers E, Magliozzo RS, McBride A, Mitra AK,
Radjainia M, Khajo A, Jacobs WR, Salgame P, Alland D. 2009. The multifunctional
histone-like protein Lsr2 protects mycobacteria against reactive oxygen intermediates.
Proceedings of the National Academy of Sciences 106:4414-4418.
23. Bartek IL, Woolhiser LK, Baughn AD, Basaraba RJ, Jacobs WR, Jr., Lenaerts AJ,
Voskuil MI. 2014. Mycobacterium tuberculosis Lsr2 is a global transcriptional regulator
required for adaptation to changing oxygen levels and virulence. mBio 5:e01106-14.
24. Kołodziej M, Łebkowski T, Płociński P, Hołówka J, Paściak M, Wojtaś B, Bury K,
Konieczny I, Dziadek J, Zakrzewska-Czerwińska J. 2021. Lsr2 and Its Novel Paralogue
Mediate the Adjustment of Mycobacterium smegmatis to Unfavorable Environmental
Conditions. mSphere 6.
25. Kołodziej M, Trojanowski D, Bury K, Hołówka J, Matysik W, Kąkolewska H, Feddersen
H, Giacomelli G, Konieczny I, Bramkamp M, Zakrzewska-Czerwińska J. 2021. Lsr2, a
nucleoid-associated protein influencing mycobacterial cell cycle. Scientific Reports
11:2910.
26. Dulberger CL, Guerrero-Bustamante CA, Owen SV, Wilson S, Wuo MG, Garlena RA,
Serpa LA, Russell DA, Zhu J, Braunecker BJ, Squyres GR, Baym M, Kiessling LL,
Garner EC, Rubin EJ, Hatfull GF. 2023. Mycobacterial nucleoid-associated protein Lsr2
is required for productive mycobacteriophage infection. Nature Microbiology 8:695-710.
27. Gessner S, Martin ZA-M, Reiche MA, Santos JA, Dinkele R, Ramudzuli A, Dhar N, de
Wet TJ, Anoosheh S, Lang DM, Aaron J, Chew T-L, Herrmann J, Müller R, McKinney
JD, Woodgate R, Mizrahi V, Venclovas Č, Lamers MH, Warner DF. 2023. Investigating
the composition and recruitment of the mycobacterial ImuA′–ImuB–DnaE2 mutasome.
eLife 12:e75628.
28. Santos JA, Timinskas K, Lamers MH, Venclovas Č, Warner DF, Gessner SJ. 2023.
RecA-NT homology motif in ImuB is essential for mycobacterial ImuA'-ImuB protein
interaction and mutasome function. bioRxiv doi:10.1101/2023.03.28.534377.
29. Goodman MF, McDonald JP, Jaszczur MM, Woodgate R. 2016. Insights into the
complex levels of regulation imposed on Escherichia coli DNA polymerase V. DNA
Repair (Amst) 44:42-50.

30. Báez-Ramírez E, Querales L, Aranaga CA, López G, Guerrero E, Kremer L, Carrère-
Kremer S, Viljoen A, Daffé M, Laval F, Cole ST, Benjak A, Alzari P, André-Leroux G,
Jacobs WR, Vilcheze C, Takiff HE. 2021. Elimination of PknL and MSMEG_4242 in
Mycobacterium smegmatis alters the character of the outer cell envelope and selects for
mutations in Lsr2. The Cell Surface 7:100060.
31. Tan KW, Pham TM, Furukohri A, Maki H, Akiyama MT. 2015. Recombinase and
translesion DNA polymerase decrease the speed of replication fork progression during
the DNA damage response in Escherichia coli cells. Nucleic Acids Res 43:1714-25.
32. Munden A, Rong Z, Sun A, Gangula R, Mallal S, Nordman JT. 2018. Rif1 inhibits
replication fork progression and controls DNA copy number in Drosophila. eLife
7:e39140.
33. Kobayashi T, Horiuchi T. 1996. A yeast gene product, Fob1 protein, required for both
replication fork blocking and recombinational hotspot activities. Genes Cells 1:465-74.
34. Tsirkas I, Dovrat D, Thangaraj M, Brouwer I, Cohen A, Paleiov Z, Meijler Michael M,
Lenstra T, Aharoni A. 2022. Transcription-replication coordination revealed in single live
cells. Nucleic Acids Research 50:2143-2156.
35. Arora K, Whiteford DC, Lau-Bonilla D, Davitt CM, Dahl JL. 2008. Inactivation of lsr2
results in a hypermotile phenotype in Mycobacterium smegmatis. J Bacteriol 190:4291-
300.
36. Kocíncová D, Singh AK, Beretti JL, Ren H, Euphrasie D, Liu J, Daffé M, Etienne G,
Reyrat JM. 2008. Spontaneous transposition of IS1096 or ISMsm3 leads to
glycopeptidolipid overproduction and affects surface properties in Mycobacterium
smegmatis. Tuberculosis (Edinb) 88:390-8.
37. Alqaseer K, Turapov O, Barthe P, Jagatia H, De Visch A, Roumestand C, Wegrzyn M,
Bartek IL, Voskuil MI, O'Hare HM, Ajuh P, Bottrill AR, Witney AA, Cohen-Gonsaud M,
Waddell SJ, Mukamolova GV. 2019. Protein kinase B controls Mycobacterium
tuberculosis growth via phosphorylation of the transcriptional regulator Lsr2 at threonine
112. Mol Microbiol 112:1847-1862.
38. Murphy KC, Papavinasasundaram K, Sassetti CM. 2015. Mycobacterial recombineering.
Methods Mol Biol 1285:177-99.
39. Falk T, Mai D, Bensch R, Çiçek Ö, Abdulkadir A, Marrakchi Y, Böhm A, Deubner J,
Jäckel Z, Seiwald K, Dovzhenko A, Tietz O, Dal Bosco C, Walsh S, Saltukoglu D, Tay
TL, Prinz M, Palme K, Simons M, Diester I, Brox T, Ronneberger O. 2019. U-Net: deep
learning for cell counting, detection, and morphometry. Nature Methods 16:67-70.
40. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K,
Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image
analysis. Nature Methods 9:676-682.

A

530   **Figure S1. DnaE1 and Lsr2 do not co-localize. (A)** Microscopy images of cells expressing

531   DnaE1-mGFPmut3 and Lsr2-mScarlet in Δ*lsr2*. **(B)** Average kymographs for each fusion (N=28

532   cells).

536
537
Figure S2. (A) Kymograph of DnaE1-mScarlet expressed from the native promoter on a phage
integrative plasmid in WT (left, N=43) or Δ*lsr2* (right, N=59). WT is the same as main text
Figure 1B for comparison. **(B)** From the cells shown in panel A, the fluorescence at mid-cell
from was normalized to time 0 and plotted over time. **(C)** Time between birth and division as
determine by visible invaginations on the phase contrast images was measured for both WT
and Δ*lsr2* cells.

Figure S3. ImuA' and DnaE2 translational fusions maintain functionality in UV-induced

mutagenesis. Frequency of rifampin-resistant mutants per 10^8 cfu without or with UV-induced

DNA damage (20 mJ/cm^2 UV). **(A)** Results shown are for WT, *imuA'B* deletion, or *imuA'B*

deletion complemented with a fluorescent translational fusion of ImuA' and ImuB. **(B)** Results

shown are for WT, *dnaE2* deletion, and *dnaE2* deletion complemented with a fluorescent

translational fusion of DnaE2. Complements are expressed from the native promoter on a

phage integrative plasmid. LOD refers to the limit of detection. p-values were obtained by one-

way ANOVA. 0.1234 (ns), 0.0332 (*), 0.0021 (**), 0.0002 (***), <0.0001 (****).

Figure S4. Microscopy images of mutasome translational fusions. Images shown are of
(A) *imuA'B* deletion mutant complemented with a fluorescent translational fusion of ImuA' and
ImuB, **(B)** *imuA'B* deletion mutant complemented with ImuA and a fluorescent translational
fusion of ImuB, and **(C)** *dnaE2* deletion mutant complemented with a fluorescent translational
fusion of DnaE2. Complements are expressed from the native promoter on a phage integrative
plasmid. DNA damage was induced by mitomycin C treatment (80 ng/ml) for 4-6 hours. Images
were acquired using 1 second **(A, B)** or 500 ms **(C)** exposure times.

Figure S5. Loss of *lsr2* leads to growth defects on mitomycin C. Serial dilutions of WT,

Δ*lsr2*, and complemented strains on agar plates with or without mitomycin C (1.56 ng/ml) in

biological triplicate. *M. smegmatis* and *M. tuberculosis lsr2* complements were expressed from

the native *lsr2* promoter on a phage integrative plasmid.

574
575 **Figure S6. Loss of *lsr2* is not associated with a decrease in survival following UV**

576 **treatment.** WT, Δ*lsr2*, and complement, were treated with 20 mJ/cm^2 UV and plated for cfu.

577 Percent survival is the ratio of cfu/ml for UV-treated cells divided by the cfu/ml for untreated cells

578 multiplied by 100. p-values were obtained by one-way ANOVA. 0.1234 (ns), 0.0332 (*), 0.0021

579 (**), 0.0002 (***), <0.0001 (****).

582