Mycobiome analyses of critically ill COVID-19 patients suggests antifungal prophylaxis may be protective against *A. fumigatus*-associated mortality

Danielle Weaver[1], Sara Gago[1], Matteo Bassetti[2,3], Daniele Roberto Giacobbe[2,3], Juergen Prattes[4], Martin Hoenigl[4,5,6], Florian Reizine[7], Hélène Guegan[8,9], Jean-Pierre Gangneux[8,9], Michael John Bromley[1] & Paul Bowyer[1]

Affiliations:

1. University of Manchester, Manchester, United Kingdom
2. Department of Health Sciences (DISSAL), University of Genoa, Genoa, Italy
3. Infectious Diseases Unit, IRCCS San Martino Polyclinic Hospital, Genoa, Italy
4. Division of Infectious Diseases, Medical University of Graz, Graz, Austria
5. Biotech Med, Graz
6. Translational Medical Mycology Research Unit, ECMM Excellence Center for Medical Mycology, Medical University of Graz, Graz, Austria
7. Centre Hospitalier Universitaire de Rennes Medical Intensive Care Unit, Rennes, France
8. Univ Rennes, CHU Rennes, Inserm, EHESP, IRSET (Institut de recherche en santé, environnement et travail) – UMR_S 1085, F-35000 Rennes, France
9. Centre Hospitalier Universitaire de Rennes, Laboratoire de Parasitologie-Mycologie, Centre National de Référence Mycoses et Antifongiques-Laboratoire Associé Asp-C, European Excellence Center for Medical Mycology (ECMM), Rennes, France

Corresponding author:

- Paul Bowyer.
- Core Technology Facility, University of Manchester, M13 9WU.
- paul.bowyer@manchester.ac.uk
- 0161 275 5411

Author's contributions

- Conception and design: DW, SG, MB, DRG, JP, MH, JPG, MJB and PB.
- Processing of specimens and generation of data: DW, SG, MB, JP, JPG, FR and HG.
- Analysis and interpretation of data: DW, SG, MH, MJB and PB
- Final approval of the manuscript: DW, SG, MB, DRG, JP, MH, FR, HG, JPG, MJB and PB.

Funding sources

- SG was co-funded by the NIHR Manchester Biomedical Research Centre, The Fungal Infection Trust, Manchester Academy of Health Sciences and The Dowager Countess Eleanor Peel Trust.
- DW was funded by the NIHR Manchester Biomedical Research Centre.

Word count: 2,856

Abstract

Rationale: Covid-associated pulmonary aspergillosis (CAPA) is a life-threatening complication

in patients with severe COVID-19. Previously, acute respiratory distress syndrome in patients

with COVID-19 has been associated with lung fungal dysbiosis, evidenced by reduced

microbial diversity and *Candida* colonisation. Increased fungal burden in the lungs of critically

ill COVID-19 patients is linked to prolonged mechanical ventilation and increased mortality.

However, specific mycobiome signatures associated with severe COVID-19 in the context of

survival and antifungal drug prophylaxis have not yet been determined and such knowledge

could have an important impact on treatment.

Objectives: To understand the composition of the respiratory mycobiome in critically ill

COVID-19 patients with and without CAPA, the impact of antifungal use and its role in patient

outcome.

Methods: We performed a multi-national study of 39 COVID-19 patients in intensive care units

(ICU) with and without CAPA. Respiratory mycobiome was profiled using ITS1 sequencing and

Aspergillus fumigatus burden was further validated using qPCR. Fungal communities were

investigated using alpha diversity, beta diversity, taxa prevalence and taxa abundances.

Measurements and Main Results: Respiratory mycobiomes were dominated by *Candida* and

Aspergillus. There was no significant association with corticosteroid use or CAPA diagnosis and

respiratory fungal communities. Increased *A. fumigatus* burden was associated with mortality.

The use of antifungals at ICU admission was associated with an absence of *A. fumigatus.*

Conclusions: Our findings suggest that systemic antifungal treatment at ICU admission may

be protective against *A. fumigatus*-associated mortality in CAPA.

Introduction

COVID-19 is a pulmonary disease caused by severe acute respiratory syndrome coronavirus 2. There have been over 500 million confirmed cases of COVID-19 since December 2019 with mortality > 6 million[1]. Around 5% of patients with COVID-19 require admission into the intensive care unit (ICU)[2,3] and, 50% of those patients need mechanical ventilation[4], thus increasing the risk of hospital-acquired pneumonia[5]. The role of the pulmonary microbiome and its associations with disease outcomes in patients with COVID-19 has been explored since the beginning of the pandemic[6–8]. However, our knowledge on the role of fungi in the pathophysiology of COVID-19 is limited. Specifically, the association between respiratory mycobiome composition and patient outcome, and the interplay of antifungal use, is yet to be investigated.

Mycobiome sequencing of the upper respiratory tract (nasopharyngeal swabs) suggests COVID-19 infection significantly reduces fungal diversity, with a higher abundance of *Alternaria* and *Cladosporium* spp.*,* and a lower abundance of other taxa such as *Candida* and *Aspergillus* in those infected[9]. In the lower respiratory tract (tracheal aspirates), bacterial and fungal microbiome analyses of patients with severe COVID-19 have shown changes over time that might be linked to antimicrobial pressure[10]. A variety of respiratory mycobiome clusters were identified, including those dominated by *Candida* and *Cladosporium.* Using 18S qPCR in bronchoalveolar lavage (BAL) samples it has been reported that critically ill COVID-19 patients with high fungal burdens are less likely to be liberated from mechanical ventilation[11]. However, the taxa responsible for this outcome remains unclear. Lastly, mycobiome sequencing of BAL found COVID-19 patients with acute respiratory distress syndrome (ARDS)

to be associated with reduced fungal diversity and an increase in *Candida* colonisation[12]. In

patients without *Candida* colonisation, an increased abundance of an unclassified *Ascomycota*

species was identified. However, little is known of the fungal species present in the airways of

patients with fungal disease, in particular *Aspergillus*-related pulmonary infections, with the

exception of fungal allergic diseases[13].

Covid-associated pulmonary aspergillosis (CAPA) is an important complication of COVID-19.

CAPA has mainly been described in critically ill patients. Multicentre cohort studies of CAPA

conducted in the ICU setting report incidence rates varying between 10-15% per centre[14–16].

Nevertheless, mortality rates in patients with CAPA were double that observed in critically ill

COVID-19 patients without CAPA[17,18]. Airway epithelial cell damage due to viral replication and

COVID-19 associated downregulation of interferon γ signalling pathway, aberrant immune

responses due to ARDS, corticosteroids, azithromycin, or the use of immunomodulators have

been linked with susceptibility to CAPA[19–22]. Here we performed a multi-national mycobiome

analysis of 39 respiratory samples from critically ill COVID-19 patients with and without CAPA.

Methods

Study design, participants, and sample collection

This study was based on a multinational retrospective study on the prevalence of invasive

pulmonary aspergillosis in critically ill COVID-19 patients in intensive care units (ICUs) during

2020[15]. Inclusion criteria consisted of: PCR confirmed COVID-19 infection, bronchoscopy or

tracheal aspiration performed during routine clinical investigations, and chest imaging

available seven days before or after respiratory samples were collected. Patients less than 18

years of age were excluded. Respiratory samples not passing quality control (described in PCR

& Sequencing section) were also excluded. Respiratory specimens obtained at ICU admission

or during ICU stay were collected, aliquoted (at least 1ml) and stored at -80 °C. Criteria for

defining aspergillosis were according to previous guidelines[23] with the following

modifications: COVID-19 requiring ICU admission was included as an additional host factor,

tracheal aspirates were equated to bronchoalveolar lavage (BAL) fluid for microbiological

tests, and serum and BAL GM was added as entry criterion. For a summary of full patient

demographics, see Table E1.

IRB approval was obtained at each participating center: Medical University of Graz EC #32-296

ex 19/20. University of Genoa Liguria Region Ethics Committee registry number 163/2020.

Sample processing

113   BAL DNA was extracted using a cetyltrimethylammonium bromide (CTAB) method[24]. Full

114   details on sample processing are provided in the online supplement. Briefly, for mycobiome

115   analysis, the ITS1 region was amplified using Nextera XT compatible versions of ITS1[25] and

116   ITS2degen (a degenerate version of ITS2 primer) (see Table 1). The presence of *A. fumigatus*

117   in respiratory samples was validated using a TaqMan probe assay targeting the ITS1 region[26].

Data analysis

Paired end reads were subject to quality trimming at Q30 and a minimum length filter of 75

nucleotides using bbduk[27] (BBMap v38.22). Primer sequences were removed using Cutadapt[28]

(v1.18). Reads were mapped to UNITE database using bowtie2[29] (v2.3.5.1). Count data was

further processed in R (v4.1.3) using the following packages: phyloseq[30] v1.38.0, vegan[31] v2.5-

7, DESeq2[32] v1.34.0, stringr[33] v1.4.0, ggplot2[34] v3.3.5 and tidyr[35] v1.2.0.

To minimise contamination in the mycobiome data, samples were subject to a multi-staged

quality control process using negative extraction control (NEC) samples, along with qPCR and

sequencing data. Firstly, samples from a DNA extraction batch with *A. fumigatus* detected (Ct

<40) in the corresponding NEC by qPCR were excluded. Secondly, samples were excluded if

they produced less than 500 fungal reads (read count filter) and/or a lower number of fungal

reads than the corresponding NEC sample (NEC count filter). Of 99 initial samples, 22 were

removed due to contamination of an extraction batch, 35 samples did not pass the read count

filter and 3 did not pass the NEC count filter.

Abundances were standardised to the median sequencing depth. Extremely low abundance

taxa were removed by only retaining those occurring > 0.2% in any sample. DESeq2 was used

to identify significantly differentially abundant taxa (adjusted *p* value < 0.05 and basemean >

500). Differences in diversity (Shannon, Chao1 and observed OTUs) were assessed using

pairwise Wilcoxon rank sum tests. PERMANOVA test was used to assess differences in Bray-

Curtis ordination.

Results

Patient cohort

The respiratory mycobiome of 91 critically ill COVID-19 patients in the intensive care unit (ICU)

was analysed using internal transcribed spacer 1 (ITS1) amplicon sequencing of BAL. Samples

from 39 patients harboured significant fungal communities which passed quality control (See

Data analysis section in methods). Table 1 describes demographic and clinical characteristics

of the 39 patients maintained in the mycobiome analysis, stratified by CAPA diagnosis. Six

patients had CAPA (15%). Patients were from Genoa, Graz and Rennes (64%, 26% and 10%,

respectively). Mean age of those with and without CAPA was 61 and 64, respectively. Most

patients received systemic corticosteroids (74% overall, 83% of those with CAPA and 73% of

those without CAPA). No patients diagnosed with CAPA received systemic antifungals at the

time of ICU admission. Mortality at end of follow-up in patients with CAPA was 33% (2/6) while

mortality in patients without CAPA was also 33% (11/33).

Table 1. Clinical and demographic characteristics of patients with and without probable

CAPA diagnosis.

Variable	Patients with CAPA (N=6)	Patients without CAPA (N=33)
Age		
Mean (SD)	61 (3.4)	64 (7.9)
valid (missing)	6 (0)	32 (1)
Sex		
male	100% (6)	79% (26)
female	0% (0)	18% (6)
missing	0% (0)	3% (1)
Ethnicity		
Caucasian	100% (6)	91% (30)
other		6.1% (2)
missing	0% (0)	3% (1)
BMI > 30		
yes	50% (3)	18% (6)
no	50% (3)	79% (26)
missing	0% (0)	3% (1)
Smoking		
yes	33% (2)	12% (4)
no	67% (4)	85% (28)
missing	0% (0)	3% (1)
Institution		
Graz	33% (2)	24% (8)
Genoa	17% (1)	73% (24)
Rennes	50% (3)	3% (1)
Hematology oncology		
yes	17% (1)	9.1% (3)
no	83% (5)	88% (29)
missing	0% (0)	3% (1)
Solid organ transplant		
yes	17% (1)	3% (1)
no	83% (5)	94% (31)
missing	0% (0)	3% (1)
Cardiovascular disease		
yes	67% (4)	45% (15)

no	33% (2)	52% (17)
missing	0% (0)	3% (1)
Pulmonary disease		
yes	33% (2)	24% (8)
no	67% (4)	73% (24)
missing	0% (0)	3% (1)
Diabetes Mellitus		
yes	17% (1)	9.1% (3)
no	83% (5)	88% (29)
missing	0% (0)	3% (1)
Corticosteroids		
yes	83% (5)	73% (24)
no	17% (1)	24% (8)
missing	0% (0)	3% (1)
Tocilizumab		
yes	17% (1)	6.1% (2)
no	83% (5)	91% (30)
missing	0% (0)	3% (1)
Azithromycin		
yes	33% (2)	33% (11)
no	67% (4)	61% (20)
missing	0% (0)	6.1% (2)
Systemic antifungals at ICU admission		
yes	0% (0)	15% (5)
no	100% (6)	6.1% (2)
missing	0% (0)	79% (26)
Life Support		
mechanical	100% (6)	70% (23)
ECMO		3% (1)
noninvasive		6.1% (2)
mechanical & noninvasive		12% (4)
none		6.1% (2)
missing	0% (0)	3% (1)
Duration ICU (days)		
Mean (SD)	27 (11)	32 (28)
valid (missing)	6 (0)	32 (1)
Palliative (Day 28 or 32)		
yes	67% (4)	48% (16)
no	33% (2)	48% (16)
missing	0% (0)	3% (1)
Survival (at end of follow up)		
yes	33% (2)	33% (11)
no	67% (4)	64% (21)
missing	0% (0)	3% (1)

Days from ICU admission to CAPA		
Mean (SD)	6.2 (3.5)	-
BAL GM (ODI > 1)		
positive	83% (5)	9.1% (3)
negative	17% (1)	82% (27)
missing	0% (0)	9.1% (3)
BAL PCR		
positive	50% (3)	0% (0)
negative	0% (0)	58% (19)
missing	50% (3)	42% (14)
BAL culture		
positive	50% (3)	0% (0)
negative	50% (3)	97% (32)
missing	0% (0)	3% (1)
BAL LFD		
positive	17% (1)	6.1% (2)
negative	0% (0)	9.1% (3)
missing	83% (5)	85% (28)
Tracheal Aspirate GM		
positive	0% (0)	3% (1)
negative	0% (0)	3% (1)
missing	100% (6)	94% (31)
Tracheal Aspirate PCR		
positive	33% (2)	0% (0)
negative	17% (1)	21% (7)
missing	50% (3)	79% (26)
Tracheal Aspirate Culture		
positive	17% (1)	0% (0)
negative	33% (2)	3% (1)
missing	50% (3)	97% (32)
Serum GM (> 0. 5)		
positive	33% (2)	0% (0)
negative	50% (3)	42% (14)
missing	17% (1)	58% (19)
Bronchial Aspirate culture		
positive	17% (1)	0% (0)
negative	33% (2)	21% (7)
missing	50% (3)	79% (26)

CAPA; COVID-19 associated pulmonary aspergillosis. BMI; body mass index. ICU; intensive care unit. ECMO;

extracorporeal membrane oxygenation. BAL; bronchoalveolar lavage. GM; galactomannan. LFD; lateral flow

device.

***Candida* and *Aspergillus spp.* dominate respiratory mycobiome communities in critically ill**

COVID-19 patients

Median read counts per sample for the 39 samples that passed our quality metric was 57,913

(range 6,970 – 342,640). Total taxa identified was 58, with a median of 7 taxa per sample

(range 2 – 28 taxa). At the genus level, there were 36 taxa in total and a median of 5 taxa per

sample (range 1 – 22). There was no significant clustering between samples batches (Fig. E1),

suggesting that sample processing had no impact on mycobiome communities. Mycobiomes

were predominantly consisted of *Candida* and *Aspergillus* species (Fig. 1A). *Candida albicans*,

Aspergillus fumigatus and *Candida parapsilosis* were the most abundant species (Fig. E2A-B).

Fig. 1. *Aspergillus* **and** *Candida spp.* **dominate the respiratory mycobiome in critically ill COVID-19 patients.** (A) *Aspergillus* and *Candida* were the main genus observed in the lungs from COVID-19 patients included in the study (n = 39). Median *A. fumigatus* abundance was higher in patients with coronavirus associated pulmonary aspergillosis (CAPA) compared to those without CAPA, measured as (B) Raw sequence reads generated by ITS1 sequencing (Median ~16,700 vs. 35) and (C) *A. fumigatus* specific qPCR (Median 0.33 vs. 0 genome equivalents) As one outlier (Grubbs test, *P* < 0.0001) had a genome equivalents value > 10, a Y axis break is included for ease of visualising majority of data points. (D) Alpha diversity measures (Observed OTUs, Chao1 and Shannon) trended towards higher diversity in CAPA patients (E) Corticosteroid treatment caused no apparent effect on alpha diversity as measured by Observed OTUs and Chao1. Shannon diversity trended towards lower diversity in patients receiving corticosteroids. (F) Median *A. fumigatus* abundance (sequence reads) was lower in patients receiving corticosteroids (Median 8 reads compared to 5,800 in those without corticosteroids). Hypothesis testing was applied using Wilcoxon Rank Sum tests. CS: Corticosteroids. Boxplot data represent median and interquartile range.

CAPA patients have higher median *A. fumigatus* abundance and trend towards lower mycobiome diversity whereas corticosteroid use has a less apparent effect on the mycobiome

In our study, higher median *A. fumigatus* levels were observed in CAPA when assessed by read count (Median ~16,700 vs. 35) and (C) *A. fumigatus* specific qPCR (Median 0.3 vs. 0 genome equivalents) (Fig. 1B-C), but there was overlap between the patient groups and, possibly due to low sample size for CAPA patients, statistical significance was not reached. *A. fumigatus* burden in patients without CAPA was varied, as some patients had little to no *A. fumigatus* and others contained particularly high burden. Furthermore, species differential abundance analysis did not find significant differences between *A. fumigatus* levels based on CAPA status. Instead, *Cladosporium delicatulum, Mycosphaerella tassiana* and *Filobasidium magnum* were found to be at higher abundance in CAPA patients (Fig. E4). In addition, probable CAPA patients had higher median α diversity (Observed OTUs, Chao1 and Shannon), however, this difference was not significant, and no significant effect on β diversity (Bray-Curtis ordination) was observed (Fig. 1D, Fig. E3).

Corticosteroid use resulted in lower median α diversity (Shannon only), however, this difference was not significant (Fig. 1E). Median levels of *A. fumigatus* were lower in the corticosteroid treated group as assessed by sequencing (8 vs. ~5,800 reads). and qPCR (0 vs. 0.1 genome equivalents) (Fig.1F, Fig. E3). However, patients on corticosteroids were highly heterogenous in terms of *A. fumigatus* abundance, some harboured little to no *A. fumigatus* and others displayed considerably high amounts. Species differential abundance analysis found no significant differences between corticosteroid usage. In addition, corticosteroid use had no significant impact on β diversity (Bray-Curtis ordination) (Fig. E3).

Increased *A. fumigatus* burden is associated with mortality

The mycobiome of individuals surviving CAPA (at the end of follow-up) showed a trend towards higher median α diversity (Fig. 2A). Alongside an increased number of OTUs, grouped mean abundances indicated a lower level of *Aspergillus* was present upon survival (Fig. 2B). Differences in species prevalence suggested this difference was due to a reduction in *A. fumigatus* in particular (Fig. 2C). Furthermore, mycobiome differential abundance analysis using DESeq2 found *A. fumigatus* and *Candida albicans* to be significantly less abundant upon survival, with log fold change values of -4.3 and -4.6, respectively (*padj* < 0.05, baseMean > 500) (Fig. 2D). Although *A. fumigatus* counts and genome equivalents (as determined by qPCR) trended towards lower levels upon survival, these changes were not significant by Wilcoxon Rank sum tests (Fig. 2E-F). Notably, *A. fumigatus* qPCR burden in those that did not survive appeared to be bimodal; 8 samples had a burden above a threshold of 0.1 haploid genome equivalents (HGE), 3 of which were patients with probable CAPA. Most patients which survived had HGE =< 0.1, excluding one outlier which had extremely high burden (> 10, higher than any other sample). In addition, BAL galactomannan index values were not significantly different between the patient groups, with again an outlier in the survival group having a very high BAL GM index (Fig. 2G).

Fig. 2. A higher *A. fumigatus* burden is associated with mortality in critically ill COVID-19 patients.
(A) Alpha diversity measures (Observed OTUs, Chao1 and Shannon) trended towards higher diversity in critically ill COVID-19 patients which survived. (B) At the genus level, BAL ITS1 mycobiomes from patients which survived (n =13) indicated a lower proportion of *Aspergillus*, and an apparent increase in the number of observed taxa overall. (C) Increased *A. fumigatus* burdens in lungs of COVID-19 patients which did not survive (blue, 0.5 prevalence) compared to those that did survive (grey, 0.3 prevalence). Taxa were counted if present at over 5% in a sample, and only taxa found in at least 10% of samples of either group are shown. (D) Analysis using DESeq2 identified *A. fumigatus* and *C. albicans* to be at significantly lower abundance in patients which survived. S: Survival; NS: No survival. Data represent median and interquartile range. (E) Median *A. fumigatus* abundance (raw sequence reads) was not significantly different between the patients. (F) *A. fumigatus* burden measured by qPCR in patients which did not survive appeared biomodal, with some patients exhibiting a high burden (>0.1 genome equivalents). (G) BAL galactomannan levels between patient groups were not significantly different. Hypothesis testing in A and D-F was applied using Wilcoxon Rank Sum tests.

Systemic antifungal use at intensive care unit admission reduces *A. fumigatus* burden in

critically ill COVID-19 patients and is associated with COVID-19 survival

Use of systemic antifungals at ICU admission in critically ill COVID-19 patients resulted in significantly reduced α diversity when analysing raw mycobiomes (prior to filtering for very rare taxa). Upon removal of very rare taxa, there was a trend towards reduced median α diversity upon systemic antifungal use at ICU admission (Fig. 3A). At the genus level, there appeared to be a strong reduction in *Aspergillus* counts that correlated with treatment (Fig. 3B). At the species level, the most prevalent taxa in patients without systemic antifungal treatment was *A. fumigatus*, whereas this species was not detectable in patients receiving treatment (Fig. 3C). Furthermore, differential abundance analysis of mycobiome data found a significant reduction of *A. fumigatus* in patients which received systemic antifungal treatment (LFC -6.3, *padj* 0.04) (Fig. 3D). This analysis also found *Candida albicans, Candida parapsilosis* and *Candida tropicalis* had significantly higher abundance in COVID-19 patients receiving antifungal treatment at ICU admission (LFC 5.3, 9.4 and 14.5, respectively). Although no significant difference was found between *A. fumigatus* counts or qPCR burden in groups with or without systemic antifungals, three patients not receiving antifungals did have high *A. fumigatus* levels and were also positive for CAPA (Fig. 3E-F). BAL galactomannan index was not statistically different between patients with or without systemic antifungal use, however, all individuals receiving treatment were GM negative (ODI < 1) whereas only half of the patients without treatment were GM negative (Fig.3G).

Fig. 3. ***A. fumigatus*** **is associated with the absence of systemic antifungals** at **intensive care unit**

admission in critically ill COVID-19 patients. (A) Alpha diversity measures (Observed OTUs, Chao1 and

Shannon) trended higher diversity in COVID-19 patients without systemic antifungals at ICU admission

(sAF). (B) At the genus level, BAL ITS1 mycobiomes from COVID-19 patients which received sAF (n =8)

display an absence of *Aspergillus.* (C) Species prevalence in each group indicated a complete lack of *A. fumigatus* in patients received sAF (grey) compared to those that did not (blue). Taxa were counted if present at over 5% in a sample, and only taxa found in at least 10% of samples of either group are shown. (D) Analysis using DESeq2 identified *A. fumigatus* to be at significantly lower abundance in the mycobiome of patients who received antifungals, compared to patients who did not. In contrast, three *Candida* species were found to be significantly more abundant in the patients receiving antifungals. (E) Median *A. fumigatus* abundance (raw sequence reads) was not significantly different between the patients. (F) *A. fumigatus* burden measured by qPCR in COVID-19 patients which did not receive sAF appeared bimodal, with three patients exhibiting a high burden (~1 genome equivalents). Notably these patients were all diagnosed with COVID19-associated pulmonary aspergillosis (CAPA)(red). (G) All COVID-19 patients receiving antifungals where BAL galactomannan negative (1 or lower). Half of patients not receiving antifungals were galactomannan positive. Hypothesis testing in A and E-G was applied using Wilcoxon Rank Sum tests. sAF: Systemic antifungals.

Our findings suggest an association between *A. fumigatus* abundance and mortality in critically ill COVID-19 patients, and that systemic antifungals at ICU significantly reduces *A. fumigatus* levels. Therefore, we combined these factors to assess the association between *A. fumigatus* and survival outcomes depending on the presence or absence of systemic antifungal treatment. Upon combining these factors, it was apparent that a higher level of *Aspergillus* was associated with mortality only in COVID-19 patients who had not received systemic antifungal treatment (Fig. 4A). Presence of *A. fumigatus* in only those patients that did not survive or receive systemic antifungal treatment was confirmed by qPCR data (Fig. 4B). Therefore, these findings suggest systemic antifungal treatment at ICU admission may have been protective against *A. fumigatus*-associated mortality in patients with severe COVID-19.

Fig. 4. _A. fumigatus_ is associated with mortality in patients with COVID-19 who have not received systemic antifungals at intensive care unit admission. (A) When combining the use of systemic antifungals at ICU and survival outcomes, _Aspergillus_ was only found in the BAL ITS1 mycobiomes from patients which did not receive sAF or survive (n=6). (B) _A. fumigatus_ levels (measured by qPCR) did not differ significantly between groups (Wilcoxon Rank Sum test); however, _A. fumigatus_ burden was observed only in the patient group which did not survive or receive systemic antifungals at ICU admission. sAF: Systemic antifungals. S: Survival. NS: No survival.

Discussion

This multinational study found that higher _Aspergillus fumigatus_ burdens in critically ill

COVID-19 patients are associated with increased mortality. In addition, the association of _A._

fumigatus with mortality was found only in patients who did not receive antifungal

treatment at ICU admission, suggesting prophylactic use of antifungals in severe COVID-19

patients may reduce _A. fumigatus_-associated mortality in this cohort.

The respiratory mycobiomes of critically ill COVID-19 patients described here were

dominated by _Candida_ and _Aspergillus._ Previous studies using similar patient groups have

also reported fungal communities in the lung to be dominated by _Candida_[10,12]. Furthermore,

one study identified a significant increase in unidentified Ascomycota spp. in patients

without _Candida_ colonisation[12]. Due to the reported incidence of CAPA in severe COVID-19,

the authors hypothesised that _Aspergillus_ could be present in these patients and, although

mycobiome samples were mostly negative for *Aspergillus*, the presence of *Aspergillus* was

confirmed by PCR in follow up BAL samples in over 20% of patients.

A recent report suggests higher fungal burden in the lung microbiota of patients with

proven/probable CAPA[11]. However, a considerable overlap in fungal burdens between those

with and without CAPA was noted. Our study found CAPA patients had higher median levels

of *A. fumigatus* and trended towards higher fungal diversity. However, these findings did not

meet statistical significance, which may have been driven by the low sample size (n = 6 in

CAPA group). Some patients within the non-CAPA group harboured considerable levels of *A.*

fumigatus. It is known that high *Aspergillus* burdens can be found in the lungs of healthy

individuals[36]. These observations suggest that as long as a sufficient level of Aspergillus is

present in the lung, other factors such as disease susceptibility or strain virulence in the

context of CAPA may be more important than burden in the outcome of infection. This study

was limited to one sample time point, and it would be interesting to assess how *Aspergillus*

burden changes during the course of CAPA or COVID-19 infection.

It has been suggested that corticosteroid treatment increases lung fungal burden

(particularly increased abundance of *A. fumigatus*)[13] and lowers Shannon diversity[37] in

asthmatic patients. In contrast, no significant differences were found between mycobiome

diversity or taxa abundance in respiratory fungal communities of COPD patients with or

without inhaled corticosteroid treatment[38]. Another recent study found that alterations in

the airway mycobiome in COPD were not significantly affected by oral or inhaled

corticosteroid use[39]. Large cohort studies suggest systemic corticosteroids are a risk factor

for CAPA[16,19], however, there are no specific reports of the influence corticosteroid use has

on lung fungal communities in COVID-19. In our study, patients receiving corticosteroids

displayed lower median levels of *A. fumigatus* and lower fungal diversity (by Shannon metric

only) compared to those not receiving corticosteroids. However, these differences did not

meet statistical significance. As most (~74%) patients received corticosteroids in this cohort,

this may have contributed to low power in these statistical comparisons, warranting further

data on this topic.

A high fungal burden has previously been associated with a lower likelihood of release from

mechanical ventilation and increased mortality risk in severe COVID-19 patients[11]. However,

as this study utilised pan-fungal qPCR to identify burden, there was no indication of the

specific fungal taxa responsible for this association. Our findings suggest that higher levels of

A. fumigatus in particular are associated with increased mortality in severe COVID-19. There

are no previous reports on the impact of antifungal use on the respiratory mycobiome in

COVID-19 patients. In this study, the use of antifungals at ICU admission appeared protective

against *A. fumigatus*-associated mortality.

Our study has investigated the composition of respiratory fungal communities in critically ill

COVID-19 patients with and without CAPA. *Candida* and *Aspergillus* were predominant in the

respiratory communities. CAPA diagnosis was associated with higher median *A. fumigatus*

level and fungal diversity, and a higher prevalence of *A. fumigatus* was associated with

mortality and a lack of systemic antifungals at ICU admission. Our data provides evidence

suggesting the use of prophylactic antifungals in seriously ill COVID-19 patients may prevent

A. fumigatus-associated mortality. A limitation of this study is the small number of patients

included, particularly the low number of CAPA cases. In addition, incomplete clinical data

with respect to antifungal use reduced the sample sizes for this comparison, which may have

resulted in low power for these statistical tests. Therefore, a further study of a larger cohort

is warranted to further understand the association between prophylactic antifungal use, the

presence of *A. fumigatus* in the respiratory mycobiome and patient outcome in COVID-19

critical care.

Conflicts of Interest

MH received research funding from Gilead, Astellas, MSD, Euroimmune, IMMY, Scynexis,

Pulmocide, F2G and Pfizer, outside the submitted work.

MJB is a former employee and has previously received research funding from F2G outside

the submitted work.

In the past 5 years SG has received speaker fees from Gilead Sciences and research grant

support from Pfizer outside of the submitted work.

Outside the submitted work, DRG reports investigator-initiated grants from Pfizer, Shionogi,

and Gilead Italia and speaker and/or advisor fees from Pfizer and Tillotts Pharma.

Outside the submitted work, MB reports research grants and/or personal fees for

advisor/consultant and/or speaker/chairman from Bayer, BioMérieux, Cidara, Cipla, Gilead,

Menarini, MSD, Pfizer, and Shionogi. In the past 5 years JPG has received speaker fees from

Gilead Sciences, MundiPharma, Pfizer, and Shionogi outside of the submitted work. In the

past 5 years HG has received speaker fees from Gilead Sciences. JP has received speakers'

fees from Gilead Sciences, Pfizer, Swedish Orphan Biovitrum, Associated of Cape Cod outside

of the submitted work, served at advisor boards Gilead Sciences and Pfizer, and holds stocks

of NovoNordisk and AbbVie Inc. All remaining authors declare no competing interests.

Acknowledgements

This work was part of the ECMM CAPA initiative under the umbrella of the ECMM NGS Working group. This work was funded by the NIHR Manchester Biomedical Research Centre (NIHR203308). The views expressed are those of the author(s) and not necessarily those of the NIHR of the Department of Health and Social Care.

References

1. World Health Organisation. WHO Coronavirus Dashboard. https://covid19.who.int/.

2. Wiersinga, W. J., Rhodes, A., Cheng, A. C., Peacock, S. J. & Prescott, H. C. Pathophysiology, Transmission, Diagnosis, and Treatment of Coronavirus Disease 2019 (COVID-19): A Review. *JAMA* **324**, 782–793 (2020).

3. Osuchowski, M. F. *et al.* The COVID-19 puzzle: deciphering pathophysiology and phenotypes of a new disease entity. *Lancet. Respir. Med.* **9**, 622–642 (2021).

4. Dongelmans, D. A. *et al.* Characteristics and outcome of COVID-19 patients admitted to the ICU: a nationwide cohort study on the comparison between the first and the consecutive upsurges of the second wave of the COVID-19 pandemic in the Netherlands. *Ann. Intensive Care* **12**, (2022).

5. Grasselli, G. *et al.* Hospital-Acquired Infections in Critically Ill Patients With COVID-19. *Chest* **160**, 454–465 (2021).

6. Merenstein, C. *et al.* Signatures of COVID-19 Severity and Immune Response in the Respiratory Tract Microbiome. *MBio* **12**, (2021).

362   7.   Lloréns-Rico, V. *et al.* Clinical practices underlie COVID-19 patient respiratory microbiome

363   composition and its interactions with the host. *Nat. Commun. 2021 121* **12**, 1–12 (2021).

364   8.   Ren, L. *et al.* Dynamics of the Upper Respiratory Tract Microbiota and Its Association with

365   Mortality in COVID-19. *Am. J. Respir. Crit. Care Med.* **204**, 1379–1390 (2021).

366   9.   Gupta, A. *et al.* Mycobiome profiling of nasopharyngeal region of SARS-CoV-2 infected

367   individuals. *Microbes Infect.* 105059 (2022) doi:10.1016/J.MICINF.2022.105059.

368   10.   Ruiz-Rodriguez, A. *et al.* Bacterial and fungal communities in tracheal aspirates of intubated

369   COVID-19 patients: a pilot study. *Sci. Reports 2022 121* **12**, 1–10 (2022).

370   11.   Kullberg, R. F. J. *et al.* Lung Microbiota of Critically Ill COVID-19 Patients are Associated with

371   Non-Resolving Acute Respiratory Distress Syndrome. *Am. J. Respir. Crit. Care Med.* 1–74

372   (2022) doi:10.1164/rccm.202202-0274OC.

373   12.   Viciani, E. *et al.* Critically ill patients with COVID-19 show lung fungal dysbiosis with reduced

374   microbial diversity in patients colonized with Candida spp. *Int. J. Infect. Dis.* **117**, 233–240

375   (2022).

376   13.   Fraczek, M. G. *et al.* Corticosteroid treatment is associated with increased filamentous fungal

377   burden in allergic fungal disease. *J. Allergy Clin. Immunol.* 1–8 (2017)

378   doi:10.1016/j.jaci.2017.09.039.

379   14.   Janssen, N. A. F. *et al.* Multinational Observational Cohort Study of COVID-19–Associated

380   Pulmonary Aspergillosis. *Emerg. Infect. Dis.* **27**, 2892 (2021).

381   15.   Prattes, J. *et al.* Risk factors and outcome of pulmonary aspergillosis in critically ill coronavirus

382   disease 2019 patients—a multinational observational study by the European Confederation of

383   Medical Mycology. *Clin. Microbiol. Infect.* **28**, 580–587 (2022).

16.   Gangneux, J. P. *et al.* Fungal infections in mechanically ventilated patients with COVID-19

during the first wave: the French multicentre MYCOVID study. *Lancet Respir. Med.* **10**, 180–

190 (2022).

17. Bartoletti, M. *et al.* Epidemiology of Invasive Pulmonary Aspergillosis Among Intubated

Patients With COVID-19: A Prospective Study. *Clin. Infect. Dis.* **73**, e3606–e3614 (2021).

18. Salmanton-García, J. *et al.* COVID-19–Associated Pulmonary Aspergillosis, March–August 2020

- Volume 27, Number 4—April 2021 - Emerging Infectious Diseases journal - CDC. *Emerg.*

Infect. Dis. **27**, 1077–1086 (2021).

19. Leistner, R. *et al.* Corticosteroids as risk factor for COVID-19-associated pulmonary

aspergillosis in intensive care patients. *Crit. Care* **26**, 1–11 (2022).

20. Koehler, P. *et al.* Defining and managing COVID-19-associated pulmonary aspergillosis: the

2020 ECMM/ISHAM consensus criteria for research and clinical guidance. *Lancet Infect. Dis.*

21, e149–e162 (2021).

21. Feys, S. *et al.* Lung epithelial and myeloid innate immunity in influenza-associated or COVID-

19-associated pulmonary aspergillosis: an observational study. *Lancet Respir. Med.* **10**, 1147–

1159 (2022).

22. Hoenigl, M. *et al.* COVID-19-associated fungal infections. *Nat. Microbiol. 2022 78* **7**, 1127–

1140 (2022).

23. Blot, S. I. *et al.* A clinical algorithm to diagnose invasive pulmonary aspergillosis in critically ill

patients. *Am. J. Respir. Crit. Care Med.* **186**, 56–64 (2012).

24. Fraczek, M. G. *et al.* The cdr1B efflux transporter is associated with non-cyp51a-mediated

itraconazole resistance in Aspergillus fumigatus. *J. Antimicrob. Chemother.* **68**, 1486–1496

(2013).

25. White, T. J., Bruns, T., Lee, S. & Taylor, J. Amplification and direct sequencing of fungal

ribosomal RNA genes for phylogenetics. *PCR Protoc.* 315–322 (1990) doi:10.1016/B978-0-12-

372180-8.50042-1.

26.   Walsh, T. J. *et al.* Molecular Detection and Species-Specific Identification of Medically

Important Aspergillus Species by Real-Time PCR in Experimental Invasive Pulmonary

Aspergillosis. *J. Clin. Microbiol.* **49**, 4150 (2011).

27.   Bushnell, B. BBMap. sourceforge.net/projects/bbmap/ (2018).

28.   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal **17**, 10 (2011).

29.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**,

357–359 (2012).

30.   McMurdie, P. J. & Holmes, S. phyloseq: An R Package for Reproducible Interactive Analysis

and Graphics of Microbiome Census Data. *PLoS One* **8**, e61217 (2013).

31.   Dixon, P. VEGAN, a package of R functions for community ecology. *J. Veg. Sci.* **14**, 927–930

(2003).

32.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol.* **15**, 1–21 (2014).

33.   Wickham, H. stringr: Simple, Consistent Wrappers for Common String Operations. R package

version 1.4.0. https://CRAN.R-project.or at (2019).

34.   Wickham, H. ggplot2: Elegant Graphics for Data Analysis. at (2019).

35.   Wickham, H. & Henry, L. tidyr: Tidy Messy Data. R package version 1.1.0. https://CRAN.R-

project.or at (2020).

36.   Nguyen, L. D. N., Viscogliosi, E. & Delhaes, L. The lung mycobiome: an emerging field of the

human respiratory microbiome. *Front. Microbiol.* **6**, 89 (2015).

37. Huang, C. *et al.* Fungal and bacterial microbiome dysbiosis and imbalance of trans-kingdom

network in asthma. *Clin. Transl. Allergy* **10**, 1–13 (2020).

38. Martinsen, E. M. H. *et al.* The pulmonary mycobiome-A study of subjects with and without

chronic obstructive pulmonary disease. *PLoS One* **16**, 1–16 (2021).

39. Tiew, P. Y. *et al.* A high-risk airway mycobiome is associated with frequent exacerbation and

mortality in COPD. *Eur. Respir. J.* **57**, (2021).

40. Wickham, H. stringr: Simple, Consistent Wrappers for Common String Operations. at

https://cran.r-project.org/package=stringr (2019).

41. Wickham, H. ggplot2: Elegant Graphics for Data Analysis. *Springer-Verlag New York* (2016).

Supplementary data

443 Raw sequence data has been deposited at the NCBI sequence read archive (SRA) under accession

444 number PRJNA905224. Code used for analysis is available at

445 https://github.com/Danweaver1/COVID_respiratory_mycobiome.

Fig. E1. Mycobiome Bray-Curtis ordination by sample batch

450 **Fig. E2. ITS1 respiratory mycobiomes at the species level.** (A) Heatmap of fungal species identified in all samples. To remove extremely rare taxa, only those present at > 0.2% in one sample were retained. (B) *A. fumigatus* and *C. albicans* are the most prevalent species. Taxa were counted if present at over 5% in a sample, and only taxa found in at least 10% of samples of either group are shown.

Fig. E3. CAPA diagnosis and corticosteroid use had no significant impact on fungal burden, *Aspergillus fumigatus* levels or beta diversity.

Fig. E4. Significantly differentially abundant taxa identified by DESeq2 analysis.

457   **Table E1. Full clinical patient demographics**

Characteristics	N=39
Age	
Mean (SD)	64 (7.4)
valid (missing)	38 (1)
Sex	
male	82% (32)
female	15% (6)
missing	2.6% (1)
Ethnicity	
Caucasian	92% (36)
other	5.1% (2)
missing	2.6% (1)
BMI > 30	
yes	23% (9)
no	74% (29)
missing	2.6% (1)
Smoking	
yes	15% (6)
no	82% (32)
missing	2.6% (1)
Institution	
Graz	26% (10)
Genoa	62% (24)
Rennes	13% (5)
Hematology oncology	
yes	10% (4)
no	87% (34)

missing	2.6% (1)
Solid organ transplant	
yes	5.1% (2)
no	92% (36)
missing	2.6% (1)
Cardiovascular disease	
yes	49% (19)
no	49% (19)
missing	2.6% (1)
Pulmonary disease	
yes	26% (10)
no	72% (28)
missing	2.6% (1)
Diabetes Mellitus	
yes	10% (4)
no	87% (34)
missing	2.6% (1)
Corticosteroids	
yes	74% (29)
no	23% (9)
missing	2.6% (1)
Tocilizumab	
yes	7.7% (3)
no	90% (35)
missing	2.6% (1)
Azithromycin	
yes	33% (13)
no	62% (24)
missing	5.1% (2)

Systemic antifungals at ICU admission	
yes	13% (5)
no	21% (8)
missing	67% (26)
Life Support	
mechanical	74% (29)
ECMO	2.6% (1)
noninvasive	5.1% (2)
mechanical & noninvasive	10% (4)
none	5.1% (2)
missing	2.6% (1)
Duration ICU (days)	
Mean (SD)	31 (26)
valid (missing)	38 (1)
Palliative (Day 28 or 32)	
yes	51% (20)
no	46% (18)
missing	2.6% (1)
Survival (at end of follow up)	
yes	33% (13)
no	64% (25)
missing	2.6% (1)
Antifungal Treatment Outcome	
yes	5.1% (2)
no	5.1% (2)
partial	0% (0)
missing	90% (35)
Primary Treatment	
yes	5.1% (2)

no	0% (0)
missing	95% (37)
Antifungals initiated for CAPA	
yes	18% (7)
no	0% (0)
missing	82% (32)
Antifungal initiated	
Posa Isa	5.2% (2)
Isav	0% (0)
IsavCasp	2.6% (1)
LAmb	0% (0)
Voriconazole	10% (4)
missing	82% (32)
BAL GM (ODI > 1)	
positive	21% (8)
negative	72% (28)
missing	7.7% (3)
BAL LFD	
positive	7.7% (3)
negative	7.7% (3)
missing	85% (33)
BAL PCR	
positive	7.7% (3)
negative	49% (19)
missing	44% (17)
BAL culture	
positive	7.7% (3)
negative	90% (35)
missing	2.6% (1)

Bronchial Aspirate culture	
positive	2.6% (1)
negative	23% (9)
missing	74% (29)
Tracheal Aspirate culture	
Positive	2.6% (1)
negative	7.7% (3)
missing	90% (35)
Tracheal Aspirate GM	
positive	2.6% (1)
negative	2.6% (1)
missing	95% (37)
Tracheal Aspirate PCR	
positive	5.1% (2)
negative	21% (8)
missing	74% (29)
Serum GM (> 0.5)	
positive	5.1% (2)
negative	44% (17)
missing	51% (20)

462 **Table E2. Full results for significantly differentially abundant taxa determined by DESeq2**

463 **analysis when assessing CAPA diagnosis, survival, and systemic antifungal use at ICU**

464 **admission.**

Variable	Taxa	base-Mean	log2Fold-Change	lfcSE	stat	p	Adjusted p
CAPA diagnosis	Cladosporium delicatulum	4389.2	13.71	4.00	3.42	0.000619	0.003719
	Filobasidium magnum	1165.8	12.93	3.13	4.13	3.59E-05	0.000368
	Fungi sp	1017.7	9.18	2.62	3.51	0.000456	0.003719
	Mycosphaerella tassiana	312608.4	20.62	4.01	5.15	2.65E-07	3.62E-06
Survival (at end of follow up)	Aspergillus fumigatus	8266.3	-4.55	1.30	-3.50	0.000462	0.017097
	Candida albicans	83190.1	-4.48	1.42	-3.15	0.001611	0.029809
Systemic anti-fungals at ICU admission	Aspergillus fumigatus	943.1	-6.33	2.43	-2.6036	0.009225	0.042435
	Candida albicans	66157.1	5.32	2.13	2.499698	0.01243	0.047648
	Candida parapsilosis	8112.9	9.35	2.68	3.486701	0.000489	0.003749
	Candida tropicalis	51931.1	14.52	2.99	4.85916	1.18E-06	1.36E-05

Supplementary methods

DNA extraction

BAL DNA was extracted using a cetyltrimethylammonium bromide (CTAB) method[24]. Negative

extraction controls (NECs) consisting of molecular grade Tris-EDTA (TE) buffer (Promega) were

included with each sample batch. First, 500 µl samples were incubated with 0.1%

dithiothreitol (Thermo Scientific) at 37 °C for 45 minutes. To lyse fungal cells, CTAB DNA

Extraction Buffer (Generon Ltd) was added and a short mechanical disruption (2 x 20 seconds

FastPrep-24) using glass beads (425-600µM, Sigma) was performed, followed by 3 cycles of

alternated heating to 65 °C and gentle vortexing for 10 mins each. Cell lysates were treated

with 4 µl RNase A (100 mg/ml, Sigma) for 30 mins prior to DNA isolation using 1:1 ratio of

Phenol-Chloroform (25:24:1, Merck) and 1:1 Chloroform:isoamyl alcohol (24:1, VWR). DNA

was precipitated at -20 °C overnight using 2 volumes of ice-cold 100% ethanol and 0.1 volumes

of 3M sodium acetate pH 4.8. Pelleted DNA was washed in 100% ethanol and twice in 70%

ethanol. Final DNA was suspended in 50 µl TE buffer.

qPCR

The presence of *A. fumigatus* in respiratory samples was validated using a TaqMan probe

assay targeting the ITS1 region[26]. Assays were performed on an Applied Biosystems 7500 fast

system using TaqMan Fast Advanced Master Mix (Thermo Fisher Scientific). Standard curves

were a nine sample 10-fold dilution series beginning at 100 ng *A. fumigatus* genomes per

reaction.

PCR & Sequencing

The ITS1 region was amplified using Nextera XT compatible versions of ITS1[25] and ITS2degen

(a degenerate version of ITS2 primer) (see Table 1). PCR setup used Phusion Green Hot Start

II HF PCR master mix to create 25 µl reactions with 300 nM forward primer, 1.2 µM reverse

primer and 2 µl DNA. Parameters included annealing temperature of 55 °C for 40 cycles. PCRs

included positive (*Aspergillus niger* genomic DNA), negative (molecular grade water) and

negative extraction controls (NECs) from DNA extractions. Controls were processed fully and

sequenced alongside the BAL samples. For sequencing library preparation, samples were

processed as per Illumina fungal metagenomic demonstrated protocol (Starting at 'Clean up'

section) using Nextera XT index kit v2 (Illumina). However, this protocol was modified to

include sample pooling prior to the second clean up. Libraries were sequenced 2x150 on an

Illumina iSeq100, with a final library loading concentration of 50 pM.

Data analysis

Paired end reads were subject to quality trimming at Q30 and a minimum length filter of 75

nucleotides using bbduk[27] (BBMap v38.22). Primer sequences were removed using Cutadapt[28]

(v1.18). Reads were mapped to UNITE database using bowtie2[29] (v2.3.5.1). Count data was

further processed in R (v4.1.3) using the following packages: phyloseq[30] v1.38.0, vegan[31] v2.5-

7, DESeq2[32] v1.34.0, stringr[40] v1.4.0, ggplot2[41] v3.3.5 and tidyr[35] v1.2.0.

To minimise contamination in the mycobiome data, samples were subject to a multi-staged

quality control process using NEC samples, along with qPCR and sequencing data. Firstly,

samples from a DNA extraction batch with *A. fumigatus* detected (Ct <40) in the

corresponding NEC by qPCR were excluded. Secondly, samples were excluded if they

produced less than 500 fungal reads (read count filter) and/or a lower number of fungal

reads than the corresponding NEC sample (NEC count filter). Of 99 initial samples, 22 were

removed due to contamination of an extraction batch, 35 samples did not pass the read

count filter and 3 did not pass the NEC count filter.

Abundances were standardised to the median sequencing depth. Extremely low abundance

taxa were removed by only retaining those occurring > 0.2% in any sample. DESeq2 was used

to identify significantly differentially abundant taxa (adjusted p value < 0.05 and basemean >

500). Differences in diversity (Shannon, Chao1 and observed OTUs) were assessed using

pairwise Wilcoxon rank sum tests. PERMANOVA test was used to assess differences in Bray-

Curtis ordination.