The respiratory enzyme complex Rnf is vital for metabolic adaptation and virulence in
Fusobacterium nucleatum

Timmie A. Britton[a], Chenggang Wu[b], Yi-Wei Chen[c], Dana Franklin[a], Yimin Chen[c], Martha I.
Camacho[c], Truc T. Luong[c], Asis Das[d], and Hung Ton-That[a,c,e,1]

[a]*Molecular Biology Institute, University of California, Los Angeles, California, USA;* [b]*Department*
of Microbiology & Molecular Genetics, University of Texas McGovern Medical School, Houston,
Texas, USA; [c]*Division of Oral & Systemic Health Sciences,, School of Dentistry, University of*
California, Los Angeles, California, USA; [d]*Department of Medicine, Neag Comprehensive Cancer*
Center, University of Connecticut Health Center, Farmington, CT, USA; [e]*Department of*
Microbiology, Immunology & Molecular Genetics, University of California, Los Angeles, Los
Angeles, CA, USA.

[1]To whom correspondence may be addressed; email htonthat@dentistry.ucla.edu

Running Title: Rnf complex in bacterial virulence

Keywords: *Fusobacterium nucleatum*, Rnf complex, metabolism, hydrogen sulfide,
coaggregation, preterm birth, virulence

ABSTRACT

A prominent oral commensal and opportunistic pathogen, *Fusobacterium nucleatum* can traverse to extra-oral sites such as placenta and colon, promoting adverse pregnancy outcomes and colorectal cancer, respectively. How this anaerobe sustains many metabolically changing environments enabling its virulence potential remains unclear. Informed by our genome-wide transposon mutagenesis, we report here that the highly conserved Rnf complex, encoded by the *rnfCDGEAB* gene cluster, is key to fusobacterial metabolic adaptation and virulence. Genetic disruption of the Rnf complex via non-polar, in-frame deletion of *rnfC* (Δ*rnfC*) abrogates polymicrobial interaction (or coaggregation) associated with adhesin RadD and biofilm formation. The defect in coaggregation is not due to reduced cell surface of RadD, but rather an increased level of extracellular lysine, which binds RadD and inhibits coaggregation. Indeed, removal of extracellular lysine via washing Δ*rnfC* cells restores coaggregation, while addition of lysine inhibits this process. These phenotypes mirror that of a mutant (Δ*kamA*) that fails to metabolize extracellular lysine. Strikingly, the Δ*rnfC* mutant is defective in ATP production, cell growth, cell morphology, and expression of the enzyme MegL that produces hydrogen sulfide from cysteine. Targeted metabolic profiling demonstrated that catabolism of many amino acids, including histidine and lysine, is altered in Δ*rnfC* cells, thereby reducing production of ATP and metabolites including H_2S and butyrate. Most importantly, we show that the Δ*rnfC* mutant is severely attenuated in a mouse model of preterm birth. The indispensable function of Rnf complex in fusobacterial pathogenesis via modulation of bacterial metabolism makes it an attractive target for developing therapeutic intervention.

IMPORTANCE

Although viewed as an oral commensal, the Gram-negative *F. nucleatum* is an opportunistic pathogen that can spread to extra-oral sites such as placenta and colon, promoting adverse pregnancy outcomes and colorectal cancer, respectively. How this anaerobe sustains various metabolically changing environments enabling its virulence potential remains unclear. We demonstrate here that the highly conserved Rnf complex is key to fusobacterial metabolic adaptation and virulence. Genetic disruption of this Rnf complex causes global defects in polymicrobial interaction, biofilm formation, cell growth and morphology, H_2S production, and ATP synthesis. Targeted metabolomic profiling demonstrates that the loss of this respiratory enzyme significantly diminishes catabolism of numerous amino acids, which negatively impacts fusobacterial virulence as tested in a preterm birth model in mice.

.

INTRODUCTION

A prominent member of the human oral microbiota that is known to harbor over 700 bacterial and fungal species (1, 2), the Gram-negative anaerobe *Fusobacterium nucleatum* plays an integral role in oral biofilm development and dental plaque formation, by virtue of its inherent capacity to adhere to diverse microbial species, notably the *Actinomyces* spp., *Streptococcus* spp., *Porphyromonas gingivalis*, and *Candida albicans* (3-12). Beyond commensalism, *F. nucleatum* is also an opportunistic pathogen that can induce preterm birth, promote tumor growth and metastatic progression of breast cancer cells, proliferation and migration of pancreatic cancer cells, and colorectal cancer (CRC), and noted for its ability to spread from the oral cavity and colonize many extra-oral sites, including placental, breast, pancreatic, and colorectal tissues (13-17). To date, a number of adhesins have been shown to mediate host-pathogen interactions. These include Fap2 and FadA, with the former aiding fusobacterial binding to tumor- and placenta-expressed Gal-GalNAc (18, 19), while the latter promoting placental colonization and CRC progression (17, 20-22). The other known adhesin RadD is a major fusobacterial virulence factor that not only mediates polymicrobial interaction (or coaggregation), a process that is inhibited by arginine and lysine (23, 24), but it is also critical for adverse pregnancy outcomes in a mouse model of preterm birth (24).

Recently, a genome-wide Tn5 transposon mutagenesis screen for identifying additional coaggregation factors revealed that the genetic disruption of a lysine metabolic pathway (LMP), e.g. deletion of *kamA* and *kamD* genes, blocks coaggregation through the excess accumulation of extracellular lysine, which binds and inhibits RadD (24). The subsequent discovery that the *kamA* deletion mutant is significantly attenuated in inducing preterm birth in mice (24) led to the realization that amino acid metabolism might play a key role in fusobacterial virulence. Consistent with this, targeted genetic analyses showed that a mutant, $\Delta megL$, defective in cysteine/methionine metabolism leading to decreased hydrogen sulfide (H_2S) production, is also

attenuated in virulence in the mouse model of preterm birth (25). Yet another set of genes

whose disruption by Tn5 insertion mutagenesis obliterated both coaggregation and biofilm

development by fusobacteria encode a putative respiratory enzyme known as the _Rhodobacter_

nitrogen-fixation (Rnf) complex (26).

Originally identified in _Rhodobacter capsulatus_ (27), the Rnf complex is a highly

conserved and evolutionarily ancient membrane-bound ferredoxin:NAD+ oxidoreductase that is

thought to couple reversible electron transfer from reduced ferredoxin (fd_{red}) to NAD^+ with the

establishment of an ion-motive force (IMF), enabling substrate import and/or ATP biosynthesis

(27-29). Found in over 150 bacterial genomes and 2 archaeal genomes, with high occurrence in

anaerobes (28), the Rnf complex has been demonstrated to be a multifunctional respiratory

enzyme that acts as a versatile metabolic exchange center for N_2-fixation (27), carbon dioxide-

fixation (30), metabolism of low-energy substrates, such as ethanol and lactate (31-33), gene

regulation to some extent (34, 35), and gut colonization in mice (36). Importantly, whether the

Rnf complex contributes to bacterial virulence remains to be explored. Here, we show that _F._

nucleatum encodes a functional _rnf_ locus, _rnfC-rnfD-rnfG-rnfE-rnfA-rnfB_, and that genetic

disruption of the Rnf complex, via deletion of _rnfC_ or _rnfD_, causes severe defects in many

virulence traits of _F. nucleatum_, including coaggregation, biofilm formation, H_2S production, and

ATP production, in addition to altering cell morphology and growth. Intriguingly, while the defect

in coaggregation is mechanistically linked to a failure of lysine catabolism, leading to an

increased level of extracellular lysine that inhibits RadD-mediated coaggregation, the defects in

other traits are attributed to global reduction of amino acid metabolism as determined by

targeted metabolomic analysis. Most significantly, the _rnfC_ mutant is severely impaired in

inducing preterm birth in mice. This study establishes that the Rnf complex is a central

component in _F. nucleatum_ metabolism that broadly impacts bacterial virulence.

RESULTS

Transposon mutagenesis reveals the involvement of the *F. nucleatum* Rnf complex in

biofilm formation, coaggregation, and H$_2$S-production. In our previous genome-wide

screens that identified *F. nucleatum* mutants defective in biofilm formation and coaggregation,

multiple subunit-encoding genes for the predicted Rnf complex were frequently targeted by Tn5

transposon insertion (24, 26) (Fig. 1A). To establish the potential multifunctional role of the Rnf

complex, we further characterized these mutants in standard biofilm and coaggregation assays

(24, 26). To cultivate monospecies biofilms, normalized fusobacterial cultures of different strains

were seeded into sterile multi-well plates, anaerobically cultured at 37°C for 48 h, and the

resultant biofilms were then stained with crystal violet. Compared to the wild-type strain (ATCC

23726), each of the four tested mutant strains were defective in biofilm formation (Fig. 1B). To

examine whether these mutants are also deficient in coaggregation, normalized, unwashed

fusobacterial cells were mixed and incubated with *S. oralis* or *A. oris* in a standard assay,

followed by imaging (see Methods). The tested Tn5 mutants also failed to co-aggregate with

these two oral co-colonizers at a level similar to that of a *radD* mutant used as a negative

control (Fig. 1C).

Unexpectedly, we noticed that all of the *rnf* mutants lacked the rotten-egg odor

characteristically produced by wild-type fusobacteria, suggestive of severe deficiency in

hydrogen sulfide production by these mutants. Indeed, in a bismuth sulfide assay, in which

bismuth trichloride reacts with H$_2$S to yield precipitation of brown bismuth sulfide, all *rnf*::Tn5

mutants showed a significantly reduced level of hydrogen sulfide relative to the wild-type strain

(Fig. 1D-1E). Overall, these results point to a multifaceted role of the Rnf complex in many

cellular processes in *F. nucleatum*.

Genetic deletion of *rnfC* disrupts RadD-mediated coaggregation via blockage of lysine

catabolism. A Tn5 insertion can potentially cause polar effects on transcription of downstream

genes. Although this is unlikely the case here (Fig. S1A), we decided to generate non-polar, in-

frame deletions of *rnf* genes utilizing our published protocols (24, 26, 37). Since all Tn5 mutants

displayed similar phenotypes, we chose *rnfC* for further characterizations, as RnfC is predicted

to be a peripheral membrane protein, enabling generation of polyclonal antibodies against a

recombinant RnfC protein (Fig. S1B). To examine how the Rnf complex mediates polymicrobial

interaction, this *rnfC* mutant (Δ*rnfC*) and other strains were subjected to the coaggregation

assay, in which unwashed fusobacterial cells were mixed with *Streptococcus gordonii*, like the

coaggregation assays with *S. oralis* and *A. oris* as mentioned above (see Fig. 1C). As expected,

deletion of *rnfC* abrogated fusobacterial coaggregation with *S. gordonii*, as compared to the

parent strain, and this defect was rescued by ectopic expression of RnfC from a plasmid (Fig.

2A; the unwashed panel).

Because RadD is essential as the adhesin for fusobacterial coaggregation (23, 24), we

examined if deletion of *rnfC* causes a defect in surface expression of RadD by immuno-

fluorescence microscopy, wherein normalized fusobacterial cells of different strains were

stained with antibodies against RadD (α-RadD), followed by counter-staining with Alexa 488-

conjugated IgG. Microscopic examination presented in Fig. S2A revealed no significant defects

of RadD-surface expression in the Δ*rnfC* as compared to the parent strain. Consistent with this

observation, western blotting of whole cell lysates showed similar levels of RadD antigen

production in all strains – parent, Δ*rnfC*, and its complement (Fig. S2B). Thus, the

coaggregation defect of Δ*rnfC* is not due to reduced RadD expression.

Unexpectedly, when we utilized Δ*rnfC* cells free of the culture medium before mixing with

S. gordonii, the mutant fusobacteria were able to adhere to oral streptococci at a level

comparable to the parent strain (Fig. 2A; compare panels washed vs unwashed). This behavior

of sensitivity to the culture medium mirrored that of a deletion mutant of *kamA*, which is

defective in lysine catabolism and hence accumulates excess lysine in the culture medium (24).

Given that lysine inhibits RadD-dependent coaggregation (23, 24), the results prompted us to

examine the relative extracellular levels of lysine in these strains by using liquid

chromatography-mass spectrometry as previously reported (24). Strikingly, compared to the

parent strain, the extracellular level of lysine for the ΔrnfC mutant remained high and was quite

comparable to that of the ΔkamA mutant (Fig. 2B). Thus, this accumulation of extracellular

lysine is likely the inhibitor that blocks coaggregation by ΔrnfC mutant bacteria. To obtain

support for this hypothesis, we re-performed the coaggregation assay with the same set of

strains in the absence or presence of added 15 mM of lysine in the coaggregation mix.

Regardless of whether fusobacterial cells were washed or not, the addition of lysine blocked

fusobacterial coaggregation with *S. gordonii* as expected (Fig. 2C).

Since deletion of *kamA* leads to accumulation of extracellular lysine (Fig. 2B) (24), we

examined the transcript level of *kamA* in the ΔrnfC mutant by qRT-PCR. Remarkably,

expression of *kamA* was significantly reduced in the absence of *rnfC*, as compared to the parent

strain, and this defect was rescued by ectopic expression of *rnfC* from a plasmid (Fig. S3). In

addition, expression of *kamD*, another lysine metabolic gene in the *kamA* locus (24), mirrored

that of *kamA* in these strains (Fig. S3). Altogether, the results indicate that the defect of RadD-

mediated coaggregation by genetic disruption of *rnfC* is due to increased extracellular levels of

lysine, likely due to the reduced expression of lysine metabolic genes by an as yet unexplored

mechanism.

Genetic deletion of *rnf*C disrupts biofilm formation, growth, cell morphology, and

hydrogen sulfide production. Next, to determine how biofilm formation is dependent on the

Rnf complex, we first analyzed the non-polar, in-frame *rnfC* deletion mutant in the

aforementioned biofilm assay using crystal violet staining (as described in Fig. 1B above). As

expected, this mutant was also unable to form monospecies biofilms compared to the parent

strain, and ectopic expression of RnfC on a plasmid was sufficient to restore biofilm formation

(Fig. 3A). Since the Rnf complex plays a key role in energy conservation through several

bacterial metabolic pathways (27, 38, 39), we reasoned that the defects in biofilm formation by

rnf mutants could be due to alterations with aspect to fusobacterial physiology. To investigate

this further, we tested whether the Rnf complex is important in ATP biosynthesis by subjecting

normalized fusobacterial cells to a microbial cell viability assay (BacTiter-Glo^{TM}), which relies on

the mono-oxygenation of luciferin catalyzed by luciferase in the presence of ATP. Indeed,

compared to the parent strain, the Δ*rnfC* mutant drastically reduced ATP production, and

complementation with an RnfC-expressing plasmid rescued the defect (Fig. 3B). Consistent with

this observation, the Δ*rnfC* mutant was also severely defective in growth displaying premature

growth cessation after the culture reached a suboptimal density (Fig. 3C); the mutant also

displayed a morphological abnormality with short/stubby cells as examined by electron

microscopy (EM) (Fig. 3D-3E).

Potentially, the physiological deficits reported above might be due to reduced

metabolism of amino acids, esp. cysteine as it is one of eight important amino acids for

fusobacteria (40). To test this hypothesis, we employed the aforementioned bismuth sulfide

assay to measure hydrogen sulfide since it is a product of cysteine metabolism in *F. nucleatum*

(25, 41-44). Consistent with the results of our Tn5 mutants, the Δ*rnfC* mutant showed a

significant defect in H_2S production, and overexpression of RnfC from a plasmid substantially

enhanced this process (Fig. 3F). Because methionine-gamma lyase MegL was previously

shown to be responsible for the bulk of H_2S production from cysteine metabolism in *F.*

nucleatum (25), we went on to determine the overall expression level of MegL in cells by

immunoblotting of whole cell lysates with antibodies against MegL (α-MegL). Remarkably,

compared with the parent and rescued strains, the Δ*rnfC* mutant expressed a significantly

reduced level of MegL (Fig. 3G). This reduction of MegL enzyme corresponded to a reduced

level of *megL* mRNA as determined by qRT-PCR (Fig. S3). Notably, the expression of *cysK1*

and *cysK2*, which encode two additional H_2S-producing enzymes (24), was also reduced in the

absence of *rnfC* (Fig. S3).

To ascertain that the physiological defects reported above are a reflection of the loss of

Rnf function, we repeated these experiments with a non-polar, in-frame *rnfD* deletion mutant

(Δ*rnfD*). Like the Δ*rnfC* mutant, the Δ*rnfD* mutant showed significant defects in coaggregation,

biofilm formation, cell growth, and H_2S production, and these defects were rescued by ectopic

expression of *rnfD* (Fig. S4). Altogether, these findings support our hypothesis that the biofilm

formation defect caused by genetic disruption of the Rnf complex is due to gross deficiencies in

ATP biosynthesis and amino acid metabolism.

Loss of RnfC complex disrupt global *F. nucleatum* metabolism. To gain further insight into

the function of the *F. nucleatum* Rnf complex in relation to the aforementioned phenotypes

associated with genetic disruption of the Rnf complex, we performed metabolomic analysis of

parent and Δ*rnfC* cells using ultra-performance liquid chromatography coupled to mass

spectrometry (UPLC/MS). Among over 80 metabolites detected in these samples, 10

metabolites (Gln, Tyr, Ile, etc) were elevated, whereas 7 metabolites, including purine

ribonucleoside precursors, were significantly depleted in the Δ*rnfC* mutant, as compared to the

parent strain (Fig. 4A).

To obtain a better view of the specific metabolic pathways altered upon loss of *rnfC*, we

performed a quantitative pathway enrichment and pathway topology analysis of the parent and

Δ*rnfC* cell using the free, web-based software, MetaboAnalyst 5.0. In keeping with our initial

findings (Fig. 4A), several pathways involved in amino acid and purine metabolism were

significantly enriched in the Δ*rnfC* mutant compared to the parent strain (Fig. 4B-4C). Pathway

analysis of the "Lys degradation", "His degradation", and "Ala/Asp/Gln metabolism" nodes in the

topology analysis revealed that many amino acids are directly degraded into glutamate, which is

subsequently fermented into the short-chain fatty acids, butyrate, and acetate (Fig. 4D).

Significantly, the Δ*rnfC* mutant had elevated levels of urocanate, glutamine, N-acetyl-L-

glutamate, lysine, and the lysine-degradation pathway intermediate, L-2-aminoadipate, relative

to the parent strain (Fig. 4E-I), indicating possible disruptions in glutamate fermentation.

Consistent with this result, we observed a reduced level of butyrate in this mutant as determined

by high-performance liquid chromatography coupled to mass-spectrometry (HPLC/MS) (Fig.

4J). Additionally, as demonstrated by pathway analysis of the "Cys/Met metabolism" node from

the topology analysis, hydrogen sulfide production is derived primarily from methionine

degradation into L-homocysteine and subsequent L-cysteine degradation into pyruvate (Fig.

S5A), both catalyzed by MegL (25). Correspondingly, elevated levels of methionine and slightly

lower levels of the pathway intermediate, S-adenosyl-L-homocysteine, were observed in the

Δ*rnfC* mutant, relative to the parent strain (Fig. S5B), suggesting the Rnf complex may also be

important for methionine/cysteine metabolism. In sum, the results support a broad role of the

Rnf complex in *F. nucleatum* metabolism.

The Rnf complex is required for *Fusobacterium nucleatum* virulence. Because genetic

disruption of the Rnf complex causes significant defects in many virulence traits of *F. nucleatum*

(Fig. 1-3), we sought to determine whether this complex contributes to *F. nucleatum*

pathogenesis. Utilizing a mouse model of pre-term birth as previously reported (24, 37), we

infected via tail vein injection groups of five CF-1 pregnant mice at day 16/17 of gestation with

roughly 5.0 x 10^7 CFU of either the parent or Δ*rnfC* strain, and pup delivery was monitored (Fig.

5A). Strikingly, the Δ*rnfC* mutant was severely attenuated in virulence, with nearly 50% of pups

born alive by the end-point, while the parent strain with almost no pub survival observed within

the same timeframe (Fig. 5B). Clearly, the Rnf complex plays a significant role in metabolism

that is central to *F. nucleatum* pathogenesis.

DISCUSSION

The dual life of *F. nucleatum* as a commensal in the human oral cavity and an opportunistic

pathogen in extra-oral sites raises an intriguing question as to how *F. nucleatum* is able to

maintain its plasticity in metabolically changing environments. Here we have demonstrated that

the highly conserved multicomponent Rnf complex plays a central role in mediating metabolism

of various amino acids in *F. nucleatum* (Fig. 6), which impacts many physiological traits critical

for fusobacterial virulence. Specifically, various *rnf* mutants emerged in unbiased forward

genetic screens in which we sought to uncover genetic factors that govern two different aspects

of *F. nucleatum* pathobiology – its ability to form biofilms and to co-aggregate with certain

partner co-colonizers of the oral cavity. We then showed that that targeted disruption of the Rnf

complex, via deletion of *rnfC* or *rnfD*, not only causes severe defects in coaggregation, and

biofilm formation but also hydrogen sulfide production, cell morphology and growth, ATP

production, and induction of preterm birth (Fig. 1-5 and Fig. S4).

The ability of fusobacteria to co-aggregate with streptococci and other partners such as

A. oris depends on a well characterized adhesin RadD present on the fusobacterial cell surface.

We have shown that the loss of Rnf complex function does not alter RadD expression or its

display at the bacterial surface (Fig. S2). Rather, the defect in RadD-mediated interbacterial

coaggregation appears to be a consequence of a defect in lysine metabolism mediated by the

LMP, which in turn raises the level of extracellular lysine that can bind and inhibit RadD to block

coaggregation (Fig. 2). Intriguingly, our results revealed that this reduced lysine metabolism

parallels a decreased expression of the LMP genes, i.e. *kamA* and *kamD*, (Fig. S3). Our

metabolomic profiling further revealed that the defect in amino acid metabolism is not limited to

lysine alone but rather the metabolism of many additional amino acids (viz. cysteine, histidine,

and glutamate) is also affected (Fig. 4 and Fig. S5). Critically, cysteine metabolism that

generates hydrogen sulfide in *F. nucleatum* is mainly catalyzed by MegL, a conserved L-

methionine γ-lyase (25). Logically, we expected that the defective cysteine metabolism in the

Δ*rnfC* mutant would lead to reduced hydrogen sulfide production by the mutant. What we did not

anticipate, however, is that the level of hydrogen sulfide production in the Δ*rnfC* mutant is

significantly lower than that of the *megL* mutant (Fig. 3F), indicating the Rnf complex plays a

broader role in amino acid metabolism.

Indeed, pathway and gene expression analyses demonstrated that multiple pathways

leading to hydrogen sulfide production are significantly affected in the absence of *rnfC* (Fig. S5)

and that expression of other cysteine metabolic genes, e.g. *cysK1* and *cysK2*, is markedly

reduced (Fig. S3). Intriguingly, given CysK1 catalyzes the conversion of cysteine to hydrogen

sulfide and L-lanthionine, an essential amino acid required for formation of fusobacterial

peptidoglycan (45, 46), the reduced expression of *cysK1* by deletion of *rnfC* can be attributed to

aberrant cell growth, short and stumpy cell size, coupled with reduced ATP production (Fig. 3).

How the absence of *rnfC* reduces expression of LMP genes and *cysK1*/*cysK2* genes remains a

significant puzzle. It is noteworthy that the expression of LMP genes and *megL* is controlled by

the two-component transduction systems (TCSs), CarRS and ModRS, respectively (24, 25); one

intriguing possibility, which remains to be tested in future studies, is that metabolic blockage by

genetic disruption of the Rnf complex causes accumulation of certain metabolites and

intermediates that may trigger gene expression responses from these TCSs and perhaps other

transcriptional regulators.

How is the Rnf complex central to *F. nucleatum* metabolism and virulence? We propose

that the *F. nucleatum* Rnf complex promotes metabolism of many amino acids, via the oxidation

of reduced ferredoxin to the reduction of NAD+ that generates an electrochemical ion gradient

across the cytoplasmic membrane, leading to the formation of several key metabolites and ATP

(Fig. 6). As such, genetic disruption of the Rnf complex causes pleiotropic defects that negatively

impact the pathophysiology of *F. nucleatum*. Given the wide conservation of the Rnf complex in

many pathogens and its absence in eukaryotes, this multi-subunit respiratory enzyme may serve

as an important target for broad anti-infective therapeutic strategies.

MATERIALS AND METHODS

Bacterial strains, plasmids, and media. The bacterial strains and plasmids used in this study

are listed in *Supplemental Information* (*SI*) Table S1. *F. nucleatum* cells were grown in tryptic soy

broth (TSB) supplemented with 1% Bacto™ peptone and 0.25% fresh cysteine (TSPC) or on

TSPC agar plates in an anaerobic chamber (10% CO_2, 10% H_2, and 80% N_2). Heart infusion broth

(HIB) or agar (HIA) was used to culture *A. oris* and supplemented with 0.5% glucose to grow *S.*

oralis and *S. gordonii*. *E. coli* strains were grown in Luria broth (LB). All bacterial strains, except

F. nucleatum, were cultured in a 5% CO_2 incubator. When required, chloramphenicol or

thiamphenicol were added into the medium at a concentration of 15 µg/ml and 5 µg/ml,

respectively. All reagents were purchased from Sigma-Aldrich unless noted otherwise.

Plasmid construction. **(i)** To generate pRnfC, the primer set com-rnfC-F/R (*SI* Table S2) was

used to amplify the *rnfC* coding sequence and its promoter region from chromosomal DNA of *F.*

nucleatum ATCC 23726, while appending the restriction sites KpnI/NdeI to the amplicon. The

PCR product was digested with KpnI and NdeI restriction enzymes and cloned into pCWU6 (*SI*

Table S2) precut with the same enzymes. **(ii)** To generate pRnfD, a segment encompassing the

rnfC promoter region and the *rnfD* coding region was PCR-amplified from chromosomal DNA of

F. nucleatum Δ*rnfC*, using the primer set com-rnfD-F/R (SI Table S2), while appending NdeI and

XhoI restriction sites to the amplicon. The PCR product was digested with NdeI and XhoI

restriction enzymes and cloned into pCWU6 precut with the same enzymes. **(iii)** To generate

pMCSG7-RnfC, the primer set, LIC-RnfC-F/R, was used to PCR-amplify the *rnfC* coding

sequence from chromosomal DNA of *F. nucleatum* ATCC 23726, while appending adapter

sequences for subsequent ligation-independent cloning (LIC) into pMCSG7 as previously

reported (47). All generated vectors were subjected to DNA sequence to confirm cloned

sequences.

Gene deletion in *F. nucleatum*. Generation of non-polar, in-frame deletion mutants, $\Delta rnfC$ and

$\Delta rnfD$, was performed according to our published protocol (24, 26, 37), with primers for generation

of deletion constructs listed in *SI* Table S2.

qRT-PCR. Overnight-grown (~ 17 h) fusobacterial cultures were normalized to OD_{600} of ~2.0, and

cells were harvested by centrifugation for RNA extraction using RNeasy Mini Kits (Qiagen)

according to the manufacturer's instructions, as described previously (25). Purified RNA, free from

DNA by treatment with DNase I (Qiagen), was used for cDNA synthesis using iScript^{TM} RT

supermix (Bio-Rad) based on the manufacturers protocol. Real-time PCR reactions were

prepared using the SYBR® Green PCR Master Mix with the appropriate primers (Table S2), and

analysis was performed using the CFX96 Real-Time System (Bio-Rad). The $\Delta\Delta C_T$ method was

used to calculate changes in gene expression between samples. Briefly, $\Delta\Delta C_T = \Delta C_{T1} - \Delta C_{T2}$,

where $\Delta C_T = C_T$ (target) - C_T (housekeeping gene). Fold changes were calculated as $\log_{10}(2^{\Delta\Delta Ct})$.

The *16S* rRNA gene was used as reference. Reactions without reverse transcriptase were used

as control to assess genomic DNA contamination.

Biofilm assay. Stationary-phase cells of fusobacterial strains normalized to an OD_{600} of ~0.6 in

fresh TSPC were seeded into flat-bottom, multi-well plates (Greiner Bio-One) and grown

anaerobically for 48 h at 37°C. Obtained biofilms were washed gently in phosphate-buffered saline

(PBS) and dried before being stained with 0.2 ml of 1% crystal violet (w/v) solution for 10 min at

room temperature. Biofilms were gently washed with sterile water before drying and imaging. For

quantification, dried biofilms were de-stained with acetic acid (30% v/v) for 10 min at room

temperature, followed by measurement at OD_{590}. Results were obtained from at least three

independent experiments performed in triplicate. Statistical analysis was performed by GraphPad

Prism 9.0.

Bacterial coaggregation assays. Fusobacterial interaction with bacterial partner strains was

assessed using a previously published coaggregation assay (24), with some modifications.

Briefly, overnight cultures of *F. nucleatum* and partner strains were harvested by centrifugation

and washed twice in coaggregation buffer [0.02 M tris-buffered saline, 150 mM NaCl, 0.1 mM

$CaCl_2$]. Fusobacterial cells were normalized to an OD_{600} of ~0.4 in coaggregation buffer without

or with 15 mM L-lysine, when necessary, whereas partner strains were normalized to an OD_{600} of

2.0 in coaggregation buffer. 0.25 ml aliquots of fusobacterial and partner strains were mixed in

multi-well plates (GenClone), and coaggregation was imaged. For the experiments using

unwashed fusobacterial cells, a similar procedure was used, except that fusobacterial cultures

were directly used without washing. Results were obtained from at least three independent

experiments performed in triplicate.

H_2S detection. Detection of H_2S was performed according to a published protocol with slight

modification (25). Briefly, overnight cultures of fusobacterial cells were normalized to an OD_{600} of

~0.6 in TSPC, and 0.1 ml of normalized cell suspension was mixed with 0.1 ml of bismuth solution

[0.4 M triethanolamine pH 8.0, 10 mM bismuth chloride, 20 mM pyridoxal 5-phosphate

monohydride, 20 mM EDTA, 40 mM L-cysteine] and incubated at 37°C anaerobic for 1 h, at which

point images were taken and H_2S was quantified by OD_{405}. The results were presented as average

of at least three independent experiments performed in triplicate.

Measurement of lysine levels by liquid chromatography-mass spectrometry. The relative

lysine levels from the culture supernatant of various fusobacterial strains grown to a stationary

phase were determined at the Metabolomics Core at Baylor College of Medicine, according to our

previous publication (24). Significance analysis was performed by GraphPad Prism.

ATP quantification assay. Mid-log phase fusobacterial cells (OD_{600} of ~ 0.5) were harvested by

centrifugation and combined in equal volumes (0.1 ml) with the BacTiter-Glo Reagent (Promega)

in multi-well plates (Cellstar). The plates were incubated at room temperature in dark for ~2 min

prior to measurement of relative luminescent units (RLU's) using a plate reader (Tecan). ATP

concentrations were determined from an ATP standard curve with ATP concentrations ranging

from 1 to 1000 nM. The results, presented as $pmol/10^6$ CFU, were obtained from at least three

independent experiments performed in triplicate.

Western blotting. Expression of fusobacterial proteins was analyzed by immunoblotting with

antibodies against RadD (α-RadD; 1:5000), MegL (α-MegL; 1:5000), FtsX (α-FtsX; 1:2000) and

RnfC (α-RnfC; 1:3000). Generation of the first 3 antibodies was described elsewhere (24-26). For

α-RnfC, the procedure was followed according to a previously published protocol (26). Briefly, *E.*

coli BL21 (DE3) harboring pMCSG7-RnfC was used to purify recombinant protein RnfC by affinity

chromatography. The purified protein was used for antibody production (Cocalico Biologicals,

Inc.). For immunoblotting, overnight fusobacterial cultures were normalized to an OD_{600} of 1.0,

and 1-ml aliquots of bacterial cultures were taken for trichloroacetic acid (TCA) precipitation and

acetone-wash, as previously reported (24). Protein samples obtained were suspended in SDS-

containing sample buffer, separated by SDS-PAGE using 4-15% Tris-Glycine gradient gels (RadD

and MegL) or a 12% Tris-Glycine gel (RnfC), and immunoblotted by specific antibodies.

Electron microscopy. Electron microscopy was performed according to previously published

protocols (24, 26). Briefly, overnight cultures of fusobacteria were harvested by centrifugation and

re-suspended in PBS supplemented with 0.1 M NaCl. A drop of bacterial suspension was added

to carbon-coated grids, stained with 1% uranyl acetate, and washed prior to imaging with an

electron microscope (JEOL 1200).

Immunofluorescence microscopy. Overnight-grown fusobacterial cells were harvested by centrifugation and washed twice in PBS before normalizing to an OD_{600} of 0.5 in PBS. 0.2-ml aliquots of cell suspension were used to coat circular glass coverslips placed in a 24-well plate for 20 min at room temperature. Cells were fixed with 2.5% formaldehyde (in PBS) for 20 min, washed with PBS, and blocked with 3% w/v bovine serum albumin (BSA) for 1 h. Cells were incubated with α-RadD (1:200) for 1 h and then AlexaFluor 488 goat anti-rabbit IgG for another hour, followed by washing in PBS three times in the dark. Cells were analyzed by a fluorescence microscope (Keyence BZ-X800).

Targeted metabolic analysis. Parent and Δ*rnfC* mutant cells were harvested by centrifugation from mid-log phase cultures in triplicate and normalized to an OD_{600} of 0.5 before drying in a speed vacuum concentrator (Thermo Scientific). Dried pellet samples were subjected to metabolomics analysis at the UC Riverside Metabolomics Core Facility as previously described with some modifications (48). In brief, 10 mg of dry pellet was extracted with 100 µl/mg of monophasic extraction solvent (30:30:20:20/acetonitrile, methanol, isopropanol and water) by sonication on ice for 5 min, followed by 90-min vortexing at 4°C. Cell-free supernatants were collected by centrifugation (3000 g) at 4°C for 30 min, and 1-ml aliquots were analyzed by a TQ-XS triple quadrupole mass spectrometer (Waters) coupled to an I-class UPLC system (Waters), with separations performed using a ZIC-pHILIC column (2.1 x 150 mm, 5 µM) (EMD Millipore). The mobile phases were (A) water supplemented with 15 mM ammonium bicarbonate titrated to a pH of 9.6 with ammonium hydroxide and (B) acetonitrile. The flow rate was 200 µL/min and the columns were held at 50°C. The injection volume was 2 µL. The gradient was as follows: 0 min, 90% B; 1.5 min, 90% B; 16 min, 20% B; 18 min, 20% B; 20 min, 90% B; 28 min, 90% B. The MS was operated in multiple reaction monitoring mode (MRM). Source and desolvation temperatures were 150° C and 600° C, respectively. Desolvation gas was set to 1100 L/h and cone gas to 150 L/h. Collision gas was set to 0.15 mL/min. All gases were nitrogen except the collision gas of

argon. Capillary voltage was 1 kV in positive ion mode and 2 kV in negative ion mode. A quality

control (QC) sample, generated by pooling equal aliquots of each sample, was analyzed

periodically to monitor system stability and performance. Samples were analyzed in random

order. Statistical analysis was performed relative to the parent strain and determined by a

Student's *t* test, using GraphPad Prism 9.0.

Raw mass-spectral data for all 81 identified metabolites with corresponding KEGG identifiers

above for *F. nucleatum* ATCC 23726 were used to perform quantitative enrichment and pathway

analysis using MetaboAnalyst 5.0 (49, 50). Statistical analysis was performed relative to the

parent strain and determined by the Global test of MetaboAnalyst 5.0.

Mouse model of preterm birth. The virulence potential of Δ*rnfC* was evaluated using a

published mouse model of preterm birth (25, 37). Briefly, groups of five CF-1 (Charles River

Laboratories) pregnant mice were infected with ~ 5 x 10^7 CFU of the parent or Δ*rnfC* strain at day

16 or 17 of gestation via tail vein injection. Pup survival was recorded for the next 7 days.

Statistical analysis was determined via the Mantel-Cox text, using GraphPad Prism 9.0, and

specified in corresponding figure legends.

All animal procedures were approved by the UCLA Animal Research Committee.

ACKNOWLEDGMENTS

We thank Amancio De Souza (Metabolomics Core Facility, UCR) for technical assistance and

our lab members for their discussion and critical review of the manuscript. Research reported in

this publication was supported by the National Institute of Dental & Craniofacial Research

(NIDCR) of the National Institutes of Health under Award Numbers DE030895 (to C.W.) and

DE026758 (to H.T.-T). T.A.B. was supported by the UCLA Dentist-Scientist and Oral Health-

Researcher Training Program, NIDCR Grant T90DE030860. D.F. was supported by the Ruth L.

Kirschstein National Research Service Award (T32AI007323). The content is solely the

responsibility of the authors and does not necessarily represent the official views of the National

Institutes of Health.

AUTHOR CONTRIBUTIONS

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

1.   Dewhirst FE, Chen T, Izard J, Paster BJ, Tanner AC, Yu WH, Lakshmanan A, Wade

WG. 2010. The human oral microbiome. J Bacteriol 192:5002-17.

2.   Caselli E, Fabbri C, D'Accolti M, Soffritti I, Bassi C, Mazzacane S, Franchi M. 2020.

Defining the oral microbiome by whole-genome sequencing and resistome analysis: the

complexity of the healthy picture. BMC Microbiol 20:120.

3.   Kolenbrander PE, Andersen RN, Moore LV. 1989. Coaggregation of *Fusobacterium*

nucleatum, *Selenomonas flueggei*, *Selenomonas infelix*, *Selenomonas noxia*, and

Selenomonas sputigena with strains from 11 genera of oral bacteria. Infect Immun

57:3194-203.

4.   Lancy P, Jr., Dirienzo JM, Appelbaum B, Rosan B, Holt SC. 1983. Corncob formation

between *Fusobacterium nucleatum* and *Streptococcus sanguis*. Infect Immun 40:303-9.

5.   Coppenhagen-Glazer S, Sol A, Abed J, Naor R, Zhang X, Han YW, Bachrach G. 2015.

Fap2 of *Fusobacterium nucleatum* is a galactose-inhibitable adhesin involved in

coaggregation, cell adhesion, and preterm birth. Infect Immun 83:1104-13.

6.   Mutha NVR, Mohammed WK, Krasnogor N, Tan GYA, Choo SW, Jakubovics NS. 2018.

Transcriptional responses of *Streptococcus gordonii* and *Fusobacterium nucleatum* to

coaggregation. Mol Oral Microbiol 33:450-464.

7.   Wu T, Cen L, Kaplan C, Zhou X, Lux R, Shi W, He X. 2015. Cellular Components

Mediating Coadherence of *Candida albicans* and *Fusobacterium nucleatum*. J Dent Res

94:1432-8.

8.   Rickard AH, Gilbert P, High NJ, Kolenbrander PE, Handley PS. 2003. Bacterial

coaggregation: an integral process in the development of multi-species biofilms. Trends

Microbiol 11:94-100.

9.   Kolenbrander PE, Andersen RN. 1989. Inhibition of coaggregation between

Fusobacterium nucleatum and *Porphyromonas* (*Bacteroides*) *gingivalis* by lactose and

related sugars. Infect Immun 57:3204-9.

10.   Karched M, Bhardwaj RG, Asikainen SE. 2015. Coaggregation and biofilm growth of

Granulicatella spp. with *Fusobacterium nucleatum* and *Aggregatibacter*

actinomycetemcomitans. BMC Microbiol 15:114.

11.   Polak D, Shapira L, Weiss EI, Houri-Haddad Y. 2012. The role of coaggregation

between *Porphyromonas gingivalis* and *Fusobacterium nucleatum* on the host response

to mixed infection. J Clin Periodontol 39:617-25.

12.   Yamaguchi-Kuroda Y, Kikuchi Y, Kokubu E, Ishihara K. 2023. *Porphyromonas gingivalis*

diffusible signaling molecules enhance *Fusobacterium nucleatum* biofilm formation via

gene expression modulation. J Oral Microbiol 15:2165001.

13. Han YW, Shen T, Chung P, Buhimschi IA, Buhimschi CS. 2009. Uncultivated bacteria as

etiologic agents of intra-amniotic inflammation leading to preterm birth. J Clin Microbiol

47:38-47.

14. Payne MS, Bayatibojakhi S. 2014. Exploring preterm birth as a polymicrobial disease: an

overview of the uterine microbiome. Front Immunol 5:595.

15. Parhi L, Alon-Maimon T, Sol A, Nejman D, Shhadeh A, Fainsod-Levi T, Yajuk O,

Isaacson B, Abed J, Maalouf N, Nissan A, Sandbank J, Yehuda-Shnaidman E, Ponath

F, Vogel J, Mandelboim O, Granot Z, Straussman R, Bachrach G. 2020. Breast cancer

colonization by *Fusobacterium nucleatum* accelerates tumor growth and metastatic

progression. Nat Commun 11:3259.

16. Udayasuryan B, Ahmad RN, Nguyen TTD, Umana A, Monet Roberts L, Sobol P, Jones

SD, Munson JM, Slade DJ, Verbridge SS. 2022. *Fusobacterium nucleatum* induces

proliferation and migration in pancreatic cancer cells through host autocrine and

paracrine signaling. Sci Signal 15:eabn4948.

17. Rubinstein MR, Wang X, Liu W, Hao Y, Cai G, Han YW. 2013. Fusobacterium

nucleatum promotes colorectal carcinogenesis by modulating E-cadherin/beta-catenin

signaling via its FadA adhesin. Cell Host Microbe 14:195-206.

18. Abed J, Emgard JE, Zamir G, Faroja M, Almogy G, Grenov A, Sol A, Naor R, Pikarsky E,

Atlan KA, Mellul A, Chaushu S, Manson AL, Earl AM, Ou N, Brennan CA, Garrett WS,

Bachrach G. 2016. Fap2 Mediates *Fusobacterium nucleatum* Colorectal

Adenocarcinoma Enrichment by Binding to Tumor-Expressed Gal-GalNAc. Cell Host

Microbe 20:215-25.

19. Parhi L, Abed J, Shhadeh A, Alon-Maimon T, Udi S, Ben-Arye SL, Tam J, Parnas O,

Padler-Karavani V, Goldman-Wohl D, Yagel S, Mandelboim O, Bachrach G. 2022.

Placental colonization by *Fusobacterium nucleatum* is mediated by binding of the Fap2

lectin to placentally displayed Gal-GalNAc. Cell Rep 38:110537.

20. Rubinstein MR, Baik JE, Lagana SM, Han RP, Raab WJ, Sahoo D, Dalerba P, Wang

TC, Han YW. 2019. *Fusobacterium nucleatum* promotes colorectal cancer by inducing

Wnt/beta-catenin modulator Annexin A1. EMBO Rep doi:10.15252/embr.201847638.

21. Xu M, Yamada M, Li M, Liu H, Chen SG, Han YW. 2007. FadA from *Fusobacterium*

nucleatum utilizes both secreted and nonsecreted forms for functional oligomerization for

attachment and invasion of host cells. J Biol Chem 282:25000-9.

22. Ikegami A, Chung P, Han YW. 2009. Complementation of the *fadA* mutation in

Fusobacterium nucleatum demonstrates that the surface-exposed adhesin promotes

cellular invasion and placental colonization. Infect Immun 77:3075-9.

23. Kaplan CW, Lux R, Haake SK, Shi W. 2009. The *Fusobacterium nucleatum* outer

membrane protein RadD is an arginine-inhibitable adhesin required for inter-species

adherence and the structured architecture of multispecies biofilm. Mol Microbiol 71:35-

47.

24. Wu C, Chen YW, Scheible M, Chang C, Wittchen M, Lee JH, Luong TT, Tiner BL, Tauch

A, Das A, Ton-That H. 2021. Genetic and molecular determinants of polymicrobial

interactions in *Fusobacterium nucleatum*. Proc Natl Acad Sci U S A 118.

25. Chen YW, Camacho MI, Chen Y, Bhat AH, Chang C, Peluso EA, Wu C, Das A, Ton-

That H. 2022. Genetic Determinants of Hydrogen Sulfide Biosynthesis in *Fusobacterium*

nucleatum Are Required for Bacterial Fitness, Antibiotic Sensitivity, and Virulence. mBio

13:e0193622.

26. Wu C, Al Mamun AAM, Luong TT, Hu B, Gu J, Lee JH, D'Amore M, Das A, Ton-That H.

2018. Forward Genetic Dissection of Biofilm Development by *Fusobacterium nucleatum*:

Novel Functions of Cell Division Proteins FtsX and EnvC. mBio 9.

27. Schmehl M, Jahn A, Meyer zu Vilsendorf A, Hennecke S, Masepohl B, Schuppler M,

Marxer M, Oelze J, Klipp W. 1993. Identification of a new class of nitrogen fixation genes

in Rhodobacter capsulatus: a putative membrane complex involved in electron transport

to nitrogenase. Mol Gen Genet 241:602-15.

28. Reyes-Prieto A, Barquera B, Juarez O. 2014. Origin and evolution of the sodium -

pumping NADH: ubiquinone oxidoreductase. PLoS One 9:e96696.

29. Biegel E, Muller V. 2010. Bacterial Na+-translocating ferredoxin:NAD+ oxidoreductase.

Proc Natl Acad Sci U S A 107:18138-42.

30. Biegel E, Schmidt S, Muller V. 2009. Genetic, immunological and biochemical evidence

for a Rnf complex in the acetogen *Acetobacterium woodii*. Environ Microbiol 11:1438-43.

31. Neumann-Schaal M, Jahn D, Schmidt-Hohagen K. 2019. Metabolism the Difficile Way:

The Key to the Success of the Pathogen *Clostridioides difficile*. Front Microbiol 10:219.

32. Weghoff MC, Bertsch J, Muller V. 2015. A novel mode of lactate metabolism in strictly

anaerobic bacteria. Environ Microbiol 17:670-7.

33. Seedorf H, Fricke WF, Veith B, Bruggemann H, Liesegang H, Strittmatter A, Miethke M,

Buckel W, Hinderberger J, Li F, Hagemeier C, Thauer RK, Gottschalk G. 2008. The

genome of *Clostridium kluyveri*, a strict anaerobe with unique metabolic features. Proc

Natl Acad Sci U S A 105:2128-33.

34. Hreha TN, Mezic KG, Herce HD, Duffy EB, Bourges A, Pryshchep S, Juarez O,

Barquera B. 2015. Complete topology of the RNF complex from *Vibrio cholerae*.

Biochemistry 54:2443-55.

35. Koo MS, Lee JH, Rah SY, Yeo WS, Lee JW, Lee KL, Koh YS, Kang SO, Roe JH. 2003.

A reducing system of the superoxide sensor SoxR in *Escherichia coli*. EMBO J 22:2614-

22.

36. Liu Y, Chen H, Van Treuren W, Hou BH, Higginbottom SK, Dodd D. 2022. *Clostridium*

sporogenes uses reductive Stickland metabolism in the gut to generate ATP and

produce circulating metabolites. Nat Microbiol 7:695-706.

37.   Peluso EA, Scheible M, Ton-That H, Wu C. 2020. Genetic Manipulation and Virulence

Assessment of *Fusobacterium nucleatum*. Curr Protoc Microbiol 57:e104.

38.   Biegel E, Schmidt S, Gonzalez JM, Muller V. 2011. Biochemistry, evolution and

physiological function of the Rnf complex, a novel ion-motive electron transport complex

in prokaryotes. Cell Mol Life Sci 68:613-34.

39.   Tremblay PL, Zhang T, Dar SA, Leang C, Lovley DR. 2012. The Rnf complex of

Clostridium ljungdahlii is a proton-translocating ferredoxin:NAD+ oxidoreductase

essential for autotrophic growth. mBio 4:e00406-12.

40.   Dzink JL, Socransky SS. 1990. Amino acid utilization by *Fusobacterium nucleatum*

grown in a chemically defined medium. Oral Microbiol Immunol 5:172-4.

41.   Fukamachi H, Nakano Y, Yoshimura M, Koga T. 2002. Cloning and characterization of

the L-cysteine desulfhydrase gene of *Fusobacterium nucleatum*. FEMS Microbiol Lett

215:75-80.

42.   Yoshida Y, Suwabe K, Nagano K, Kezuka Y, Kato H, Yoshimura F. 2011. Identification

and enzymic analysis of a novel protein associated with production of hydrogen sulfide

and L-serine from L-cysteine in *Fusobacterium nucleatum* subsp. *nucleatum* ATCC

25586. Microbiology 157:2164-71.

43.   Yoshida Y, Ito S, Kamo M, Kezuka Y, Tamura H, Kunimatsu K, Kato H. 2010. Production

of hydrogen sulfide by two enzymes associated with biosynthesis of homocysteine and

lanthionine in *Fusobacterium nucleatum* subsp. *nucleatum* ATCC 25586. Microbiology

(Reading) 156:2260-2269.

44.   Suwabe K, Yoshida Y, Nagano K, Yoshimura F. 2011. Identification of an L-methionine

gamma-lyase involved in the production of hydrogen sulfide from L-cysteine in

Fusobacterium nucleatum subsp. *nucleatum* ATCC 25586. Microbiology 157:2992-3000.

45.   Vasstrand EN, Hofstad T, Endresen C, Jensen HB. 1979. Demonstration of lanthionine

594   as a natural constituent of the peptidoglycan of *Fusobacterium nucleatum*. Infect Immun

595   25:775-80.

596   46.   Fredriksen A, Vasstrand EN, Jensen HB. 1991. Peptidoglycan precursor from

597   *Fusobacterium nucleatum* contains lanthionine. J Bacteriol 173:900-2.

598   47.   Siegel SD, Amer BR, Wu C, Sawaya MR, Gosschalk JE, Clubb RT, Ton-That H. 2019.

599   Structure and Mechanism of LcpA, a Phosphotransferase That Mediates Glycosylation

600   of a Gram-Positive Bacterial Cell Wall-Anchored Protein. mBio 10.

601   48.   Vliet SMF, Dasgupta S, Sparks NRL, Kirkwood JS, Vollaro A, Hur M, Zur Nieden NI,

602   Volz DC. 2019. Maternal-to-zygotic transition as a potential target for niclosamide during

603   early embryogenesis. Toxicol Appl Pharmacol 380:114699.

604   49.   Xia J, Psychogios N, Young N, Wishart DS. 2009. MetaboAnalyst: a web server for

605   metabolomic data analysis and interpretation. Nucleic Acids Res 37:W652-60.

606   50.   Pang Z, Zhou G, Ewald J, Chang L, Hacariz O, Basu N, Xia J. 2022. Using

607   MetaboAnalyst 5.0 for LC-HRMS spectra processing, multi-omics integration and

608   covariate adjustment of global metabolomics data. Nat Protoc 17:1735-1761.

 FIGURE LEGENDS

Figure 1: Transposon mutagenesis reveals the involvement of an Rnf complex in biofilm

formation, coaggregation, and H₂S production. (A) Shown is the *rnf* gene locus in *F.*

nucleatum, with arrows indicating the Tn5 insertion sites of previously identified Tn5 mutants

(24, 26). **(B)** Mono-species biofilms of the parent (WT) and Tn5 *rnf* mutant strains were

cultivated in multi-well plates, stained with 1% crystal violet, and imaged. **(C)** Indicated

fusobacterial strains were examined for their adherence to *S. oralis* and *A. oris* by a

coaggregation assay. A *radD*::Tn5 mutant was used as a negative control. **(D-E)** Production of

H₂S from indicated strains was determined by the bismuth sulfide method (D) and quantified by

absorbance measurement at 405 nm (E). Cell-free samples were used as a negative control. All

results were obtained from three independent experiments performed in triplicate.

Figure 2: A non-polar, in-frame *rnfC* deletion mutant is defective in RadD-mediated

coaggregation and lysine uptake. (A) The parent, *rnfC* deletion mutant, and *rnfC*-

complemented strains were tested for their ability to co-aggregate with *S. gordonii.* Fusobacterial

cells were washed or unwashed before being mixed in equal volumes with washed *S. gordonii*

and imaged. Mutants devoid of *radD* or *kamA*, a lysine metabolic pathway gene (24), were used

as references. **(B)** The relative level of lysine in the cell-free culture medium of indicated strains

grown to stationary phase was determined by liquid chromatography-mass spectrometry, with the

level of media without bacteria set to 100. **(C)** The same coaggregation experiment as (A) was

performed with or without addition of 15 mM lysine. All results were obtained from three

independent experiments performed in triplicate. Significance was calculated by a student's t-test;

*P < 0.05; **P < 0.01.

Figure 3: Deletion of *rnfC* causes pleiotropic defects. (A) 48-h grown biofilms of indicated

strains were stained with 1% crystal violet and quantified by optical density measurement at 590

nm (OD$_{590}$). **(B)** ATP production of indicated strains was assessed by a luciferase assay using a

BacTiter-Glo™ kit (Promega). Mid-log phase cells were mixed with equal volumes of the BacTiter-

Glo™ reagent and incubated at room temperature for 2 min before measuring luminescence on

a plate reader. Normalized cell suspensions were serially diluted and plated for bacterial

enumeration and subsequent normalization of relative luminescent signal. **(C)** Bacterial growth of

indicated strains was determined by optical density at 600 nm at timed intervals. **(D-E)** Mid-log

phase cells of parent and Δ*rnfC* mutant strains were immobilized on carbon-coated nickel grids

and stained with 1% uranyl acetate prior to imaging by electron microscopy; scale bars: 1 μm. **(F)**

H$_2$S-production of indicated strains was determined by the bismuth sulfide assay. **(G)** Protein

samples obtained from the whole-cell lysates of normalized cultures of indicated strains were

subjected to immunoblotting with antibodies against MegL (α-MegL). A Coomassie Blue stained

band (*) was used as a loading control. All results were obtained from three independent

experiments performed in triplicate. Significance was calculated by a student's t-test; *P < 0.05;

***P < 0.001; ****P < 0.0001.

Figure 4: Deletion of *rnfC* disrupts amino acid metabolism. (A) The parent and Δ*rnfC* cells

grown to mid-log phase were subjected to metabolomics analysis using liquid chromatography-

mass spectrometry (LC-MS). Shown is a volcano plot of 81 differentially expressed metabolites

(DEMs) between the parent and Δ*rnfC* mutant strains. Metabolites significantly depleted or

enriched in the Δ*rnfC* strain, relative to the parent, are marked in red or green, respectively. **(B)**

Shown is a graphical overview of quantitative enrichment analysis of DEMs between the parent

and Δ*rnfC* strains generated using MetaboAnalyst 5.0. The top 24 pathway-associated metabolic

sets in Δ*rnfC* compared to the parent strain were sorted based on their fold enrichment and P-

values. **(C)** Pathway analysis was performed using MetaboAnalst 5.0, which combines pathway

enrichment and topology analysis. A range of P-values is shown from red to yellow. **(D)** Shown

are amino acid metabolic nodes in *F. nucleatum*, based on the significance level in (C), generating

acetate and butyrate. **(E-I)** The relative concentrations of DEMs in the amino acid nodes shown

in D between the parent and Δ*rnfC* mutant strains are presented. P-values were calculated using

the Global test. **(J)** The relative level of butyrate (mM) in the overnight cultures of the parent and

Δ*rnfC* strains was determined by liquid chromatography-mass spectrometry (LC-MS).

Figure 5: The *rnfC* mutant is attenuated in virulence. (A-B) Groups of pregnant CF-1 mice

were infected with ~5.0 x 10^7 CFU/mL of the parent or Δ*rnfC* strain via tail-vein injection at day 16

or 17 of gestation. Pup survival was recorded over time. Statistical analysis was determined by

Mantel-Cox testing.

Figure 6: A working model of Rnf-mediated energy conservation through amino acid

metabolism in *F. nucleatum*. See text for details.

A

rnfB | rnfA | rnfE | rnfG | rnfD | rnfC

B

WT | rnfA::Tn5 | rnfB::Tn5 | rnfC::Tn5 | rnfD::Tn5   biofilm

C

S. oralis

A. oris

WT | rnfA::Tn5 | rnfB::Tn5 | rnfC::Tn5 | radD::Tn5

D

rnfC::Tn5 | rnfA::Tn5 | rnfB::Tn5 | radD::Tn5 | rnfD::Tn5 | control

WT

E

OD$_{405}$

2.0
1.5
1.0
0.5
0

WT | Tn5::rnfA | Tn5::rnfB | Tn5::rnfC | Tn5::rnfD

*** *** *** ***

Figure 1: Britton et al.

A

unwashed

washed

parent Δ*rnfC* Δ*rnfC*/ Δ*kamA* Δ*radD*
                 pRnfC

B

Relative lysine level

cell-free parent Δ*rnfC* Δ*rnfC*/ Δ*kamA*
medium   pRnfC

** **

C

unwashed cells

-

washed cells

-

Lys parent Δ*rnfC* Δ*rnfC*/ Δ*kamA* Δ*radD*
(mM)   pRnfC

Figure 2: Britton et al.

Figure 3: Britton et al.

Figure 4: Britton et al.

A

mating ~5x10⁷ CFU birth

day 0   16/17   20+

B

P < 0.0001

ΔrnfC

parent

Pup survival (%)

Time (h)

Figure 5: Britton et al.

Figure 6: Britton et al.